

Fourier transform, null variety, and Laplacian's eigenvalues

Article

Accepted Version

Benguria, R., Levitin, M. and Parnovski, L. (2009) Fourier transform, null variety, and Laplacian's eigenvalues. *Journal of Functional Analysis*, 257 (7). pp. 2088-2123. ISSN 0022-1236 doi: <https://doi.org/10.1016/j.jfa.2009.06.022> Available at <http://centaur.reading.ac.uk/16152/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1016/j.jfa.2009.06.022>

To link to this article DOI: <http://dx.doi.org/10.1016/j.jfa.2009.06.022>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Fourier transform, null variety, and Laplacian's eigenvalues*

Rafael Benguria

Facultad de Física

P. Universidad Católica de Chile

Casilla 306, Santiago 22

Chile

rbenguri@fis.puc.cl

Michael Levitin

Cardiff School of Mathematics

Cardiff University, and WIMCS

Senghennydd Road, Cardiff CF24 4AG

United Kingdom

Levitin@cardiff.ac.uk

Leonid Parnovski

Department of Mathematics

University College London

Gower Street, London WC1E 6BT

United Kingdom

leonid@math.ucl.ac.uk

version 7

June 16, 2009

Abstract

We consider a quantity $\kappa(\Omega)$ — the distance to the origin from the null variety of the Fourier transform of the characteristic function of Ω . We conjecture, firstly, that $\kappa(\Omega)$ is maximized, among all convex balanced domains $\Omega \subset \mathbb{R}^d$ of a fixed volume, by a ball, and also that $\kappa(\Omega)$ is bounded above by the square root of the second Dirichlet eigenvalue of Ω . We prove some weaker versions of these conjectures in dimension two, as well as their validity for domains asymptotically close to a disk, and also discuss further links between $\kappa(\Omega)$ and the eigenvalues of the Laplacians.

Keywords: Laplacian, Dirichlet eigenvalues, Neumann eigenvalues, eigenvalue estimates, Fourier transform, characteristic function, Pompeiu problem, Schiffer's conjecture, convex sets

2000 Mathematics Subject Classification: 42B10, 35P15, 52A40

*The research has been supported by the Royal Society International collaborative grant with Chile. The work of RB has also been supported by CONICYT/PBCT (Chile) Proyecto Anillo de Investigación en Ciencia y Tecnología ACT30/2006. The work of LP has been also supported by a Leverhulme Trust grant.

1 Introduction

Let Ω be a bounded open domain in \mathbb{R}^d with boundary $\partial\Omega$, let $\mathbf{x} = (x_1, \dots, x_d)$ be a vector of Cartesian coordinates in \mathbb{R}^d , and let

$$\chi_\Omega(\mathbf{x}) = \begin{cases} 1, & \text{if } \mathbf{x} \in \Omega, \\ 0 & \text{if } \mathbf{x} \notin \Omega \end{cases}$$

denote the characteristic function of Ω .

The complex Fourier transform of $\chi_\Omega(\mathbf{x})$,

$$\widehat{\chi}_\Omega(\boldsymbol{\xi}) = \mathcal{F}[\chi_\Omega](\boldsymbol{\xi}) := \int_{\Omega} e^{i\boldsymbol{\xi} \cdot \mathbf{x}} d\mathbf{x}$$

or, more importantly, its *complex null variety*, or *null set*,

$$\mathcal{N}_{\mathbb{C}}(\Omega) := \{\boldsymbol{\xi} \in \mathbb{C}^d : \widehat{\chi}_\Omega(\boldsymbol{\xi}) = 0\}$$

has been studied extensively. Particular attention has been attracted by the role it plays in numerous attempts to prove the famous *Pompeiu problem* and *Schiffer's conjecture*. We can refer for example to [Agr, Avi, Ber, BroKah, BroSchTay, GarSeg, Kob1, Kob2]; this list is by no means complete.

Although our paper is not directly related to these still open questions, we recall them as part of the motivation for further study of the null variety.

Let $\mathcal{M}(d)$ be a group of rigid motions of \mathbb{R}^d , and Ω be a bounded simply connected domain with piecewise smooth boundary. The Pompeiu problem is to prove that the existence of a non-zero continuous function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ such that $\int_{\mathbf{m}(\Omega)} f(\mathbf{x}) d\mathbf{x} = 0$ for all $\mathbf{m} \in \mathcal{M}(d)$ implies that Ω is a ball.

Schiffer's conjecture is that the existence of an eigenfunction v (corresponding to a non-zero eigenvalue μ) of a Neumann Laplacian on a (simply connected) domain Ω such that $v \equiv \text{const}$ along the boundary $\partial\Omega$ (or, in other words, the existence of a non-constant solution v to the over-determined problem $-\Delta v = \mu v$, $\partial v / \partial n|_{\partial\Omega} = 0$, $v|_{\partial\Omega} = 1$) implies that Ω is a ball.

It is known that the positive answer to the Pompeiu problem is equivalent to Schiffer's conjecture. Moreover, a domain Ω would be a counterexample to both if there exists $r > 0$ such that $\mathcal{N}_{\mathbb{C}}(\Omega)$ contains the complex sphere $\{\boldsymbol{\xi} \in \mathbb{C}^d : \sum_{j=1}^d \xi_j^2 = r^2\}$. One of the common tools in attacking the conjectures has been an asymptotic analysis of the null variety far from the origin in an attempt to prove that such counterexample cannot exist.

In many cases, the study of the null variety in the papers cited above has been restricted to the case of a *convex* domain Ω . Additionally, it is convenient to assume that Ω is *balanced* (i.e., centrally symmetric with respect to the origin), and to deal instead with the *real null variety*

$$\mathcal{N}(\Omega) := \mathcal{N}_{\mathbb{C}}(\Omega) \cap \mathbb{R}^d = \{\boldsymbol{\xi} \in \mathbb{R}^d : \widehat{\chi}_\Omega(\boldsymbol{\xi}) = 0\} = \{\boldsymbol{\xi} \in \mathbb{R}^d : \int_{\Omega} \cos(\boldsymbol{\xi} \cdot \mathbf{x}) d\mathbf{x} = 0\}.$$

We assume that Ω is convex and balanced in most parts of this paper.

The purpose of this paper is to study the behaviour of the null variety *near* the origin, and its relation with the classical spectral theory. Namely, we define the numbers

$$\kappa_{\mathbb{C}}(\Omega) := \text{dist}(\mathcal{N}_{\mathbb{C}}(\Omega), \mathbf{0}) = \min\{|\xi| : \xi \in \mathcal{N}_{\mathbb{C}}(\Omega)\}$$

and

$$\kappa(\Omega) := \text{dist}(\mathcal{N}(\Omega), \mathbf{0}) = \min\{|\xi| : \xi \in \mathcal{N}(\Omega)\}$$

(if there are no real zeros, we set $\kappa(\Omega) = \infty$). Throughout most of the paper, we will be dealing with the real zeros of Fourier transform and the quantity $\kappa(\Omega)$, so, unless specified otherwise, we always assume that the argument of the Fourier transform $\mathcal{F}[\chi_{\Omega}]$ is real.

On the basis of some partial cases presented below in Section 3, we conjecture that, firstly, $\kappa(\Omega)$ is maximized, among all convex balanced domains of the same volume as Ω , by a ball (see Conjecture 2.2), and, secondly, that for all convex balanced domains $\kappa(\Omega)$ is bounded above by the square root of the second Dirichlet eigenvalue of Ω (see Conjecture 2.3). Note that it is very easy to see that $\kappa(\Omega)$ is always (i.e. without the convexity and central symmetry conditions) bounded below by the square root of the second Neumann eigenvalue of Ω , see Lemma 3.3.

Unfortunately we are unable to prove Conjectures 2.2 and 2.3 as stated. Even in the planar case $d = 2$, when the geometry of convex domains is easier to deal with, we are only able to establish some weaker versions of these conjectures, see Theorems 2.4 and 2.5. However, even these weaker results shed some extra light on the links between $\kappa(\Omega)$ and Dirichlet and Neumann eigenvalues, and in particular show some surprising links with Friedlander's inequalities between the eigenvalues of these two problems, see Remark 3.9 and Remark 4.9. Additionally, we can also establish the validity of Conjectures 2.2 and 2.3 for small star-shaped perturbations of a disk, see Theorem 2.7.

We also indicate that our results and conjectures can not be extended to wider classes of domains, in particular when the convexity condition is dropped, see Theorems 2.8, 2.9, and Corollary 2.10.

The rest of this paper is organized as follows. Section 2 contains the statements of our Conjectures and main Theorems. Some particular cases making the conjectures plausible are collated in Section 3. Some preliminary estimates (which in particular imply the validity of Conjecture 2.2 for relatively "long and thin" planar convex balanced domains) are presented and proved in Section 4. Extra notation and facts from convex geometry are in Section 5. Section 6 contains the proof of Theorem 2.4; some auxiliary technical Lemmas used in the proofs are collected in a separate Section 7. The perturbation-type results are proved in Section 8, and the counterexamples for non-convex domains are proved in Section 9.

We finish this Section by introducing some additional notation used throughout the paper. We write $\text{vol}_d(\cdot)$ for a d -dimensional Lebesgue measure of a set. Given a unit vector $\mathbf{e} \in S^{d-1}$, we write $x_{\mathbf{e}} = \mathbf{x} \cdot \mathbf{e}$ and $\mathbf{x}'_{\mathbf{e}} = \mathbf{x} - x_{\mathbf{e}}\mathbf{e}$. We write a real vector $\boldsymbol{\xi} \in \mathbb{R}^d$ in spherical coordinates as $\boldsymbol{\xi} = (\rho, \boldsymbol{\omega})$, with $\rho = |\boldsymbol{\xi}|$ and $\boldsymbol{\omega} = \boldsymbol{\xi}/\rho \in S^{d-1}$. $B_d(R) = \{\mathbf{x} \in \mathbb{R}^d : |\mathbf{x}| < R\}$ denotes a ball of radius R centred at $\mathbf{0}$, and a shorthand for a unit ball will be $B_d = B_d(1)$. Ω^* stands for a ball in \mathbb{R}^d centred at $\mathbf{0}$ and of the same volume as Ω .

Additionally, for a direction $\mathbf{e} \in S^{d-1}$, we define $\kappa_j(\mathbf{e}) = \kappa_j(\mathbf{e}; \Omega)$ as the j -th positive real ρ -zero of $\widehat{\chi}_{\Omega}(\rho\mathbf{e})$ (counting multiplicities); note that $\mathcal{N}(\Omega) = \bigcup_{j=1}^{\infty} \mathcal{N}_j(\Omega)$, where

$$\mathcal{N}_j(\Omega) := \bigcup_{\mathbf{e} \in S^{d-1}} \kappa_j(\mathbf{e}; \Omega)\mathbf{e}.$$

Finally, $J_a(r)$ are the usual Bessel functions of order a , and $j_{a,k}$ are their positive zeros numbered in increasing order. The eigenvalues of the Dirichlet Laplacian on Ω are denoted by $\lambda_k(\Omega)$, $k = 1, \dots$, and of the Neumann Laplacian by $\mu_j(\Omega)$, $j = 1, \dots$ ($\mu_1 = 0$).

Acknowledgments

We would like to express our gratitude to I. Polterovich for stimulating discussions, to F. Nazarov and N. Sidorova for helping with the proof of Theorem 2.9, and to N. Filonov for letting us use his Lemma 3.4. We are also grateful to the referee for useful suggestions.

2 Conjectures and statements

Definition 2.1. Ω is *balanced* if it is invariant with respect to the mapping $\mathbf{x} \mapsto -\mathbf{x}$.

Conjecture 2.2. *If Ω is convex and balanced, then*

$$\kappa(\Omega) \leq \kappa(\Omega^*), \quad (2.1)$$

with the equality iff Ω is a ball.

Conjecture 2.3. *If Ω is convex and balanced, then*

$$\kappa(\Omega) \leq \sqrt{\lambda_2(\Omega)}, \quad (2.2)$$

with the equality iff Ω is a ball.

In the next section we consider several explicit examples for which we demonstrate the validity of these conjectures.

Although we believe these Conjectures to be true, we are unable to prove them without some additional assumptions. We can however establish somewhat weaker forms in the two-dimensional case as stated in the next two theorems. Also, we can prove (2.1) subject to some additional conditions on Ω , see Corollaries 4.3 and 4.4, and Remark 4.5.

Theorem 2.4. *If $d = 2$, and Ω is convex and balanced, then*

$$\kappa(\Omega) \leq C\kappa(\Omega^*), \quad (2.3)$$

with

$$C = \tilde{C} := \frac{2j_{0,1}}{j_{1,1}} \approx 1.2552. \quad (2.4)$$

Theorem 2.5. *If $d = 2$, and Ω is convex and balanced, then*

$$\kappa(\Omega) \leq 2\sqrt{\lambda_1(\Omega)}. \quad (2.5)$$

Remark 2.6. Note that Theorem 2.5 immediately follows from Theorem 2.4 by the Faber-Krahn inequality,

$$\lambda_1(\Omega) \geq \lambda_1(\Omega^*) = \frac{\pi j_{0,1}^2}{\text{vol}_2(\Omega)},$$

and rescaling properties of Lemma 3.2. Note also that (2.5) is clearly weaker than (2.2) in the two-dimensional case, since, by the Payne-Pólya-Weinberger inequality [PayPólWei], in two dimensions

$$\lambda_2(\Omega) < 3\lambda_1(\Omega),$$

or by the even stronger Ashbaugh-Benguria inequality [AshBen],

$$\lambda_2(\Omega) \leq \left(\frac{j_{1,1}}{j_{0,1}}\right)^2 \lambda_1(\Omega) \approx 2.539\lambda_1(\Omega).$$

Finally, in the one-dimensional case, a convex balanced domain is an interval $(-a, a) = B_1(a)$ for some $a > 0$, and

$$\kappa(B_1(a)) = \sqrt{\lambda_2(B_1(a))} = \frac{\pi}{a},$$

so that (2.1) and (2.2) hold with equality.

We can also establish the validity of (2.1) and (2.2) for balanced star-shaped (but not necessarily convex) domains which are close to a disk. Namely, let $F : \mathbb{S}^1 \rightarrow \mathbb{R}$ be a C^2 function on the unit circle; we additionally assume that F is periodic with period π :

$$F(\theta + \pi) = F(\theta). \quad (2.6)$$

For $\epsilon \geq 0$, define a domain in polar coordinates (r, θ) as

$$\Omega_{\epsilon F} := \{(r, \theta) : 0 \leq r \leq 1 + \epsilon F(\theta)\}. \quad (2.7)$$

Condition 2.6 implies that $\Omega_{\epsilon F}$ is balanced.

Assume additionally that F is *area preserving*, that is

$$\int_0^{2\pi} F(\theta) \, d\theta = 0, \quad (2.8)$$

and so

$$\text{vol}_2(\Omega_{\epsilon F}) = \pi + O(\epsilon^2).$$

As we shall see from the re-scaling properties summarized in Lemma 3.2, condition (2.8) can be assumed without any loss of generality.

The unperturbed domain (when $\epsilon = 0$), Ω_{0F} , is just a unit planar disk B_2 .

We have

Theorem 2.7. *Let us fix a non-zero function F as above satisfying (2.6) and (2.8). Then the one-sided derivatives satisfy*

$$\left. \frac{d\kappa(\Omega_{\epsilon F})}{d\epsilon} \right|_{\epsilon=0+} < 0, \quad (2.9)$$

and

$$\left. \frac{d\kappa(\Omega_{\epsilon F})}{d\epsilon} \right|_{\epsilon=0+} < \left. \frac{d\sqrt{\lambda_2(\Omega_{\epsilon F})}}{d\epsilon} \right|_{\epsilon=0+}. \quad (2.10)$$

Consequently, for sufficiently small $\epsilon > 0$ (depending on F), Conjectures 2.2 and 2.3 with $\Omega = \Omega_{\epsilon F}$ hold.

On the other hand, there exist arbitrarily small star-shaped non-convex perturbations of the disk for which at least (2.1) does not hold. Namely, we have

Theorem 2.8. *For each positive $\tilde{\delta}$, there exists a balanced star-shaped domain Ω with $\text{vol}_2(\Omega) = \pi$ and such that $B(0, 1 - \tilde{\delta}) \subset \Omega \subset B(0, 1 + \tilde{\delta})$, for which $\kappa(\Omega) > j_{1,1}$.*

Continuing formulating negative results, we have the following

Theorem 2.9. *There is no C such that (2.3) holds uniformly for all (not necessarily connected) balanced one-dimensional domains Ω .*

From this, we immediately have

Corollary 2.10. *There is no C such that (2.3) holds uniformly for all balanced connected two-dimensional domains Ω .*

Theorem 2.8 and Corollary 2.10 show that convexity plays a crucial role in Theorem 2.4 and Conjecture 2.2

3 Motivation and elementary domains

We start with two trivial results, which are immediate by the change of variables, and which in particular show that our conjectures are scale invariant. Let \mathcal{R}_α denotes a mapping $(x_1, x_2, \dots, x_d) \mapsto (\alpha x_1, x_2, \dots, x_d)$, $\alpha > 0$.

Lemma 3.1. *For any $\Omega \subset \mathbb{R}^d$,*

$$\mathcal{N}(\mathcal{R}_\alpha \Omega) = \mathcal{R}_{1/\alpha} \mathcal{N}(\Omega).$$

Lemma 3.2. *Let Ω' be the image of $\Omega \subset \mathbb{R}^d$ under a homothety with coefficient $\alpha > 0$. Then*

$$\kappa(\Omega') = \frac{1}{\alpha} \kappa(\Omega), \quad \lambda_j(\Omega') = \frac{1}{\alpha^2} \lambda_j(\Omega).$$

The following result illustrates that there exists a relation between the null variety and eigenvalues of the Neumann Laplacian, which makes Conjecture 2.3 even more intriguing.

Lemma 3.3. *For any $\Omega \subset \mathbb{R}^d$,*

$$\kappa(\Omega) \geq \kappa_{\mathbb{C}}(\Omega) \geq \sqrt{\mu_2(\Omega)}. \quad (3.1)$$

Proof. Let $\xi_0 \in \mathcal{N}_{\mathbb{C}}(\Omega)$, and so $\int_{\Omega} e^{i\xi_0 \cdot x} dx = 0$. This means that $\langle e^{i\xi_0 \cdot x}, 1 \rangle_{L_2(\Omega)} = 0$, so that $\phi := e^{i\xi_0 \cdot x}$ is a test function for $\mu_2(\Omega)$ (obviously, $\phi \in H^1(\Omega)$). But, by direct computation,

$$\frac{\|\nabla \phi\|_{L_2(\Omega)}^2}{\|\phi\|_{L_2(\Omega)}^2} = |\xi_0|^2.$$

Thus, $|\xi_0|^2 \geq \mu_2(\Omega)$ for any $\xi_0 \in \mathcal{N}_{\mathbb{C}}(\Omega)$, whence the result. \square

In fact, as was shown to us by N. Filonov [Fil2], one can improve this result to obtain

Lemma 3.4. *For any $\Omega \subset \mathbb{R}^d$,*

$$\kappa(\Omega) \geq 2\sqrt{\mu_2(\Omega)}. \quad (3.2)$$

Proof. By the variational principle,

$$\mu_2(\Omega) \leq \sup_{\phi \in \mathcal{L}_2} \frac{\|\nabla \phi\|_{L_2(\Omega)}^2}{\|\phi\|_{L_2(\Omega)}^2}$$

for any linear subspace $\mathcal{L}_2 \subset H^1(\Omega)$ such that $\dim \mathcal{L}_2 = 2$. Choose $\xi_0 \in \mathcal{N}(\Omega)$, and set $\mathcal{L}_2 = \text{span}(e^{i\xi_0 \cdot x/2}, e^{-i\xi_0 \cdot x/2})$. The elements of \mathcal{L} are linearly independent, and the result immediately follows by direct computation. \square

Example 3.5 (A ball in \mathbb{R}^d). For a unit ball B_d and real ξ , we have:

$$\widehat{\chi}_{B_d}(\xi) = (2\pi)^{d/2} \frac{J_{d/2}(|\xi|)}{|\xi|^{d/2}}, \quad (3.3)$$

and so

$$\kappa(B_d) = j_{d/2,1}. \quad (3.4)$$

On the other hand,

$$\lambda_2(B_d) = \lambda_3(B_d) = \cdots = \lambda_{1+d}(B_d) = j_{d/2,1}^2 = (\kappa(B_d))^2.$$

For illustration, we give a proof of (3.3) in dimension $d = 2$. We choose the direction of ξ as the x_1 -axis, and write, in polar coordinates, $\mathbf{x} = (r \cos \theta, r \sin \theta)$.

Thus,

$$\widehat{\chi}_{B_d}(\xi) = \int_0^1 \int_0^{2\pi} e^{i|\xi|r \cos \theta} r \, dr \, d\theta.$$

Then we use formula [AbrSte, formula 9.1.18], i.e.,

$$J_0(z) = \frac{1}{2\pi} \int_0^{2\pi} \cos(z \cos \theta) \, d\theta,$$

to express the previous integral as

$$\widehat{\chi}_{B_d}(\xi) = 2\pi \int_0^1 J_0(|\xi|r) r \, dr = \frac{2\pi}{|\xi|^2} \int_0^{|\xi|} J_0(s) s \, ds.$$

Finally, we use the raising and lowering relations for Bessel functions embodied in [AbrSte, formula 9.1.27] (third formula with $\nu = 1$), i.e.,

$$J_1'(r) = J_0(r) - \frac{1}{r} J_1(r),$$

which can be expressed in the more convenient form,

$$r J_0(r) = (r J_1(r))'.$$

Thus, we get,

$$\widehat{\chi}_{B_d}(\xi) = \frac{2\pi}{|\xi|^2} \int_0^{|\xi|} (s J_1(s))' \, ds = \frac{2\pi}{|\xi|} J_1(|\xi|),$$

which is the desired equality (3.3) in two dimensions. The corresponding formula in any dimension is equally simple to establish.

Example 3.6 (A cuboid in \mathbb{R}^d). Consider, for $d \geq 2$, a cuboid P with edge lengths $a_1 \geq a_2 \geq \dots \geq a_d > 0$. We have:

$$\lambda_2(P) = \pi^2 \left(4a_1^{-2} + \sum_{j=2}^d (a_j)^{-2} \right). \quad (3.5)$$

On the other hand, if $\xi \in \mathcal{N}(P)$, we have $\prod_{j=1}^d \sin(\xi_j a_j / 2) = 0$, and so $|\xi|$ is minimized by the vector $(2\pi/a_1, 0, \dots, 0)$, giving

$$\kappa(P) = \frac{2\pi}{a_1} < \sqrt{\lambda_2(P)}. \quad (3.6)$$

Proving (2.1) for P requires a bit more effort. We have

$$P^* = B_d(R) \quad \text{with } R = \frac{(a_1 \cdot a_2 \cdots a_d \cdot \Gamma(1 + d/2))^{1/d}}{\sqrt{\pi}},$$

and, after some transformations, the required inequality is reduced to

$$j_{d/2,1} \geq 2\sqrt{\pi} \left(\Gamma \left(1 + \frac{d}{2} \right) \right)^{1/d}.$$

This, in turn, is proved using a combination of Stirling's formula, Lorch's lower bound $j_{\nu,1} \geq \sqrt{(\nu+1)(\nu+5)}$ [Lor], and numerical checks for low d .

Example 3.7 (A right-angled triangle in \mathbb{R}^2). Let $T = T_{1,a}$ be a right-angled triangle with sides 1, $a > 1$, and $\sqrt{1+a^2}$. One can check, after some computations, that

$$\kappa(T_{1,a}) = 2\pi\sqrt{1+a^{-2}}.$$

We remark that both inequalities (2.1) and (2.2) with $\Omega = T$ hold for values of a sufficiently close to one, but fail for large a or small a . This can be checked either by direct computation (in case of (2.1)) or by domain monotonicity (in case of (2.2)), by comparing $\lambda_2(T)$ with either $\lambda_2(T_{a,a}) = 10\pi^2/a^2$ (for small a) or with the second eigenvalue of the rectangle with sides $3/4$ and $a/4$ (for large a).

Note that T is not balanced and we do not conjecture that (2.1) and (2.2) hold in general for such domains. It may be plausible that $\kappa_{\mathbb{C}}(\Omega) \leq \kappa(\Omega^*)$ and $\kappa_{\mathbb{C}}(\Omega) \leq \sqrt{\lambda_2(\Omega)}$ for general convex domains, however the study of complex null varieties is outside the scope of this paper.

Example 3.8 (Numerics). We have also verified Conjectures 2.2 and 2.3 numerically. We have conducted (jointly with Brian Krushave, an undergraduate student at Heriot-Watt University, whose research was funded by a Nuffield Foundation undergraduate bursary) a large number of calculations for different multiparametric families of balanced convex domains in the two-dimensional case. A typical example would be a family of rectangles with different circular

or elliptic segments added along their sides, in order to produce some stadium-like domains.

The zeros of Fourier transform were found by analytic or numerical integration and minimization, and the eigenvalues of the Dirichlet Laplacian by the finite element method.

Remark 3.9 (Estimates of the spectrum). We would like to show how to use estimates of $\kappa(\Omega)$ in spectral inequalities between the eigenvalues $\lambda_n = \lambda_n(\Omega)$ of the Dirichlet Laplacian on Ω and the eigenvalues $\mu_n = \mu_n(\Omega)$ of the Neumann Laplacian on the same domain. It is known that for general domains we have

$$\mu_{n+1} < \lambda_n \quad (3.7)$$

for each n , and, moreover, for convex domains in \mathbb{R}^d we have

$$\mu_{n+d} < \lambda_n \quad (3.8)$$

[LevWei]. It was conjectured that (3.8) holds for all domains; this conjecture remains open, and we remark that a ‘counterexample’ given in the paper by Levine and Weinberger is erroneous.

Estimate (3.7) was proved by Friedlander [Fri] for domains with smooth boundaries; later, an elegant proof for arbitrary domains was obtained by Filonov [Fil1]. Filonov’s proof goes like this. Let n be fixed. Denote by ϕ_j the Dirichlet eigenfunctions of Ω . By the min-max principle, in order to prove $\mu_{n+1} \leq \lambda_n$, it is enough to find a subspace \mathcal{L} of $H^1(\Omega)$ such that $\dim \mathcal{L} = n + 1$ and for each $\phi \in \mathcal{L} \setminus \{0\}$ we have

$$\int_{\Omega} |\nabla \phi|^2 \, dx \leq \lambda_n \int_{\Omega} \phi^2 \, dx. \quad (3.9)$$

Put $\mathcal{L} = \text{span}(\phi_1(\mathbf{x}), \dots, \phi_n(\mathbf{x}), e^{i\xi \cdot \mathbf{x}})$, where ξ is any real vector satisfying $|\xi|^2 = \lambda_n$. Obviously, $\dim \mathcal{L} = n + 1$. Suppose now that $\phi \in \mathcal{L}$. This means that

$$\phi = \sum_{j=1}^n a_j \phi_j + b e^{i\xi \cdot \mathbf{x}}.$$

Then the left-hand side of (3.9) is

$$\sum_{j=1}^n |a_j|^2 \lambda_j + |b|^2 \lambda_n \text{vol}_d(\Omega) + 2 \sum_{j=1}^n \text{Re} \left(a_j b \lambda_n \int_{\Omega} \phi_j e^{i\xi \cdot \mathbf{x}} \, dx \right)$$

(in the last sum, we have integrated by parts using the fact that ϕ_j satisfies Dirichlet boundary conditions on $\partial\Omega$ and that $|\xi|^2 = \lambda_n$). The right-hand side of (3.9) is

$$\lambda_n \left(\sum_{j=1}^n |a_j|^2 + |b|^2 \text{vol}_d(\Omega) + 2 \sum_{j=1}^n \text{Re} \left(a_j b \int_{\Omega} \phi_j e^{i\xi \cdot \mathbf{x}} \, dx \right) \right).$$

Comparing the last two expressions leads to (3.9).

Now suppose we want to improve this result and to show (3.8) that for some class of (not necessarily convex) domains $\mu_{n+2} \leq \lambda_n$. The natural approach to try is to add one more exponential to L , namely to put

$$\mathcal{L} = \text{span}(\phi_1(\mathbf{x}), \dots, \phi_n(\mathbf{x}), e^{i\xi_1 \cdot \mathbf{x}}, e^{i\xi_2 \cdot \mathbf{x}}),$$

$$|\xi_j|^2 = \lambda_n, \quad j = 1, 2. \quad (3.10)$$

Then, in order for (3.9) to hold, we must get rid of the cross-term with two exponentials, i.e. we must assume that

$$\int_{\Omega} e^{i(\xi_1 - \xi_2) \cdot \mathbf{x}} d\mathbf{x} = 0.$$

In the notation introduced above, this means that

$$\xi_1 - \xi_2 \in \mathcal{N}(\Omega). \quad (3.11)$$

Obviously, we can choose vectors ξ_1, ξ_2 satisfying both (3.10) and (3.11) iff $\kappa(\Omega) \leq 2\sqrt{\lambda_n(\Omega)}$. Thus, if we could show that for some, not necessarily convex, d -dimensional domain Ω , the estimate (2.5) holds, then the inequality $\mu_{n+2}(\Omega) \leq \lambda_n(\Omega)$ will hold for each n . Similarly, for any Ω , if we know a number n_0 such that $\kappa(\Omega) \leq 2\sqrt{\lambda_{n_0}(\Omega)}$, then the inequality $\mu_{n+2}(\Omega) \leq \lambda_n(\Omega)$ is guaranteed to hold for $n \geq n_0$.

4 Some estimates of $\kappa(\Omega)$ for convex balanced domains

Throughout this section Ω is convex and balanced, and Ω dependence is frequently dropped; also we always work with *real* zeros of the Fourier transform.

Our aim here is to prove the following

Theorem 4.1. *Suppose that $d = 2$ and $D(\Omega)$ is the diameter of Ω . Then*

$$\kappa(\Omega) \leq \frac{4\pi}{D(\Omega)}. \quad (4.1)$$

Remark 4.2. After this paper was written, we have discovered that Theorem 4.1 had been previously proved in [Zas].

Note that Theorem 4.1 immediately implies

Corollary 4.3. *Conjecture 2.2 holds for convex, balanced domains $\Omega \subset \mathbb{R}^2$ such that the diameter $D(\Omega)$ satisfies*

$$\frac{\sqrt{\pi}D(\Omega)}{2\sqrt{\text{vol}_2(\Omega)}} \geq \frac{2\pi}{j_{1,1}}. \quad (4.2)$$

The scaling in (4.2) is chosen in such a way that its left-hand side equals one for a disk.

Let $r_-(\Omega)$ be the *inradius* of a convex balanced domain Ω . Then it is easy to see that there exists a rectangle with sides $2r_-(\Omega)$ and $D(\Omega)$ which contains Ω . Thus $2r_-(\Omega)D(\Omega) \geq \text{vol}_2(\Omega)$, which together with Corollary 4.3 immediately implies

Corollary 4.4. *Conjecture 2.2 holds for convex, balanced domains $\Omega \subset \mathbb{R}^2$ such that inradius $r_-(\Omega)$ satisfies*

$$\frac{\sqrt{\pi}r_-(\Omega)}{\sqrt{\text{vol}_2(\Omega)}} \leq \frac{j_{1,1}}{8}. \quad (4.3)$$

Remark 4.5. In the same spirit, one can also establish the validity of Conjecture 2.3 subject to additional geometric constraints: if a domain is sufficiently “long” (i.e. the left-hand side of (4.2) is sufficiently large or the left-hand side of (4.3) is sufficiently small), then (2.2) holds. However such an estimate would be non-explicit, as there is no explicit isoperimetric bound on the second Dirichlet eigenvalue for convex domains, see [Hen].

Before proving Theorem 4.1, we need to introduce some auxiliary notation, and establish some technical facts.

Fix $\mathbf{e} \in S^{d-1}$, and define the function $\nu_{\mathbf{e}} : \mathbb{R} \rightarrow \mathbb{R}$ by

$$\nu_{\mathbf{e}}(t) = \text{vol}_{d-1}(\{\mathbf{x} : x_{\mathbf{e}} = t\} \cap \Omega)$$

It is easy to see that $\nu_{\mathbf{e}}$ is an even function and has a compact support $\text{supp } \nu_{\mathbf{e}} = [-w(\mathbf{e}), w(\mathbf{e})]$, where w is the support function of Ω , i.e. $w(\mathbf{e})$ is a half-breadth of Ω in direction \mathbf{e} . If Ω is convex and $d = 2$, then $\nu_{\mathbf{e}}$ is a concave function on $[-w(\mathbf{e}), w(\mathbf{e})]$ (this is not true if $d \geq 3$, e.g. when Ω is a cube and \mathbf{e} is a diagonal but in general Brunn–Minkowski inequality implies that $(\nu_{\mathbf{e}}(\rho))^{1/(d-1)}$ is concave on $[-w(\mathbf{e}), w(\mathbf{e})]$). Thus, $\nu_{\mathbf{e}}(t) \leq \nu_{\mathbf{e}}(0)$ and the function $\nu_{\mathbf{e}}$ is non-increasing on $[0, w(\mathbf{e})]$.

As we are working with real zeros of the Fourier transform, we can instead work with

$$\widehat{\chi}_{\mathbf{e}}(\rho) := \widehat{\chi}(\rho\mathbf{e}) = \int_{\Omega} \cos(\rho\mathbf{e} \cdot \mathbf{x}) \, d\mathbf{x} = 2 \int_0^{w(\mathbf{e})} \cos(t\rho) \nu_{\mathbf{e}}(t) \, dt.$$

Lemma 4.6. *Let $Z : [0, z] \rightarrow \mathbb{R}$ be non-increasing and concave. Then*

$$\int_{2\pi k}^{2\pi(k+1)} Z(t) \cos(t) dt \leq 0, \quad (4.4)$$

$$\int_{2\pi(k+1/2)}^{2\pi(k+3/2)} Z(t) \cos(t) dt \geq 0, \quad (4.5)$$

$$\int_{2\pi k}^{2\pi(k+1/2)} Z(t) \cos(t) dt \geq 0, \quad (4.6)$$

$$\int_{2\pi(k+1/2)}^{2\pi(k+1)} Z(t) \cos(t) dt \leq 0 \quad (4.7)$$

for $k \in \mathbb{N}$ (assuming that all intervals of integration are inside $[0, z]$).

Proof. Let $L(t)$ be a linear function such that $L(2\pi(k+1/4)) = Z(2\pi(k+1/4))$ and $L(2\pi(k+3/4)) = Z(2\pi(k+3/4))$. Then, by concavity of $Z(t)$, we have $Z(t) \geq L(t)$ for $t \in [2\pi(k+1/4), 2\pi(k+3/4)]$ (note that $\cos(t) \leq 0$ for these values of t) and also $Z(t) \leq L(t)$ for $t \in [2\pi k, 2\pi(k+1/4)] \cup [2\pi(k+3/4), 2\pi(k+1)]$ (note that $\cos(t) \geq 0$ for these values of t). Therefore,

$$\int_{2\pi k}^{2\pi(k+1)} Z(t) \cos(t) dt \leq \int_{2\pi k}^{2\pi(k+1)} L(t) \cos(t) dt = 0,$$

the last equality easily checked by a direct computation. This proves (4.4), and (4.5) is being dealt with similarly.

Further,

$$\int_{2\pi k}^{2\pi(k+1/2)} Z(t) \cos(t) dt = \int_{2\pi k}^{2\pi(k+1/4)} (Z(t) - Z(2\pi(k+1/2) - t)) \cos(t) dt \geq 0,$$

since the integrand is non-negative. Inequality (4.7) is similar. \square

Remark 4.7. Note that (4.4) and (4.5) require only concavity of a function Z , whereas (4.6) and (4.7) require only its monotonicity.

Recall that $\kappa_j(\mathbf{e})$ denotes the j -th ρ -root (counted in increasing order with account of multiplicities) of $\widehat{\chi}_{\mathbf{e}}(\rho)$.

Lemma 4.8. *Let $d = 2$, then*

$$\kappa_j(\mathbf{e}) \leq \frac{\pi(j+1)}{w(\mathbf{e})}.$$

Proof. We have $\widehat{\chi}_{\mathbf{e}}(0) > 0$. Set $\rho_j = \frac{j\pi}{w(\mathbf{e})}$. Let us show that

$$\widehat{\chi}_{\mathbf{e}}(\rho_j) = \widehat{\chi}_{\mathbf{e}}\left(\frac{j\pi}{w(\mathbf{e})}\right)$$

is non-negative when j is odd, and is non-positive when j is even.

Assume $j = 2k$. Then

$$\begin{aligned}\widehat{\chi}_{\mathbf{e}}(\rho_j) &= 2 \int_0^w \cos\left(\frac{2\pi kt}{w}\right) \nu_{\mathbf{e}}(t) dt = \frac{w}{\pi k} \int_0^{2\pi k} \cos(\tau) \nu_{\mathbf{e}}\left(\frac{\tau w}{2\pi k}\right) d\tau \\ &= \frac{w}{\pi k} \sum_{\ell=0}^{k-1} \int_{2\pi\ell}^{2\pi(\ell+1)} \cos(\tau) \nu_{\mathbf{e}}\left(\frac{\tau w}{2\pi k}\right) d\tau,\end{aligned}$$

which is non-positive by (4.4).

Assume now that $j = 2k + 1$. Then

$$\begin{aligned}\widehat{\chi}_{\mathbf{e}}(\rho_j) &= 2 \int_0^w \cos\left(\frac{2\pi(k+1)t}{w}\right) \nu_{\mathbf{e}}(t) dt = \frac{w}{\pi k} \int_0^{2\pi(k+1)} \cos(\tau) \nu_{\mathbf{e}}\left(\frac{\tau w}{2\pi(k+1)}\right) d\tau \\ &= \frac{w}{\pi k} \int_0^{\pi} \cos(\tau) \nu_{\mathbf{e}}\left(\frac{\tau w}{2\pi(k+1)}\right) d\tau \\ &\quad + \frac{w}{\pi k} \sum_{\ell=0}^{k-1} \int_{2\pi(\ell+1/2)}^{2\pi(\ell+3/2)} \cos(\tau) \nu_{\mathbf{e}}\left(\frac{\tau w}{2\pi(k+1)}\right) d\tau,\end{aligned}$$

which is non-negative by (4.5) and (4.6).

The result now follows from the Intermediate Value theorem (if, for example, $\widehat{\chi}_{\mathbf{e}}(\rho)$ is positive except at the points ρ_{2k} where it is zero, then each point ρ_{2k} is a zero of multiplicity (at least) two, so we still have $\kappa_j(\mathbf{e}) \leq \frac{\pi(j+1)}{w(\mathbf{e})}$) \square

Lemma 4.8 immediately leads to the main result of this section.

Proof of Theorem 4.1. By Lemma 4.8,

$$\kappa(\Omega) = \inf_{\mathbf{e} \in S^1} \kappa_1(\mathbf{e}) \leq \inf_{\mathbf{e} \in S^1} \frac{2\pi}{w(\mathbf{e})} = \frac{2\pi}{\sup_{\mathbf{e} \in S^1} w(\mathbf{e})} = \frac{4\pi}{D(\Omega)}.$$

\square

Remark 4.9. It was proved in [Zas] that the function $\kappa_1(\mathbf{e})$ is continuous. Using this fact, one can establish further relationship between this function and Neumann eigenvalues, similar to Lemma 3.3. For example, we have:

$$\max_{\mathbf{e} \in S^1} \kappa_1(\mathbf{e}) \geq \sqrt{\mu_3(\Omega)}. \quad (4.8)$$

Indeed, recall that $\mathcal{N}_1 = \mathcal{N}_1(\Omega) = \bigcup_{\mathbf{e} \in S^1} \kappa_1(\mathbf{e})\mathbf{e}$. Obviously, $\mathcal{N}_1(\Omega) \subset \mathcal{N}(\Omega)$.

Assuming the continuity of $\kappa_1(\mathbf{e})$, we see that \mathcal{N}_1 is a continuous closed curve having the origin inside it. Let \mathbf{e}_0 be arbitrary unit vector so that $\mathbf{p}_0 := \kappa_1(\mathbf{e}_0)\mathbf{e}_0 \in \mathcal{N}_1$. Then the closed curve $\mathbf{p}_0 + \mathcal{N}_1$ obviously contains both the points inside \mathcal{N}_1 (the origin) and outside \mathcal{N}_1 (for example, the point $2\mathbf{p}_0$). Therefore, the intersection $(\mathbf{p}_0 + \mathcal{N}_1) \cap \mathcal{N}_1$ is non-empty, say $\mathbf{p}_1 \in (\mathbf{p}_0 + \mathcal{N}_1) \cap \mathcal{N}_1$. Then three points \mathbf{p}_0 , \mathbf{p}_1 , and $\mathbf{p}_0 - \mathbf{p}_1$ all belong to \mathcal{N}_1 . Now we can argue as in the proof of Lemma 3.3, with $e^{i\mathbf{p}_0 \cdot \mathbf{x}}$, $e^{i\mathbf{p}_1 \cdot \mathbf{x}}$ and 1 being three mutually orthogonal test-functions. This shows that $\sqrt{\mu_3(\Omega)} \leq \max(|\mathbf{p}_1|, |\mathbf{p}_2|) \leq \max_{\mathbf{e} \in S^1} \kappa_1(\mathbf{e})$.

Remark 4.10. Using the results of this section and the fact that $D(\Omega) \geq 2\sqrt{\text{vol}_2(\Omega)/\pi}$, we obtain

$$\kappa(\Omega) \leq \frac{4\pi}{D(\Omega)} \leq \frac{2\pi^{3/2}}{\sqrt{\text{vol}_2(\Omega)}} = \frac{2\pi}{j_{1,1}} \kappa(\Omega^*),$$

thus proving (2.3) with a numerical constant

$$C = \tilde{C}_1 = \frac{2\pi}{j_{1,1}} \approx 1.6398.$$

Remark 4.11. It should be noted that there is no analog of Theorem 4.1 in dimensions higher than two, i.e. one cannot estimate $\kappa(\Omega)$ in terms of the diameter $D(\Omega)$. Indeed, let $S = \{(x_1, x_2) \in \mathbb{R}^2 : |x_1| + |x_2| < 1\}$, and let T be the three-dimensional body of revolution obtained by rotating S around the x_1 -axis. Also, choose $\alpha > 0$, and set $T_\alpha = \{\mathbf{x} \in \mathbb{R}^3 : (x_1, \alpha x_2, \alpha x_3) \in T\}$. Then, as $\alpha \rightarrow \infty$, the distances to origin of all zeros of $\widehat{\chi_{T_\alpha}}$, which are not proportional to $\mathbf{e}_1 = (1, 0, 0)$, tend to ∞ by Lemma 3.1. On the other hand,

$$\widehat{\chi_{T_\alpha}}(\xi \mathbf{e}_1) = \alpha^{-2} \widehat{\chi_T}(\xi \mathbf{e}_1) = 2\pi\alpha^{-2} \int_0^1 (1-x)^2 \cos(x\xi) dx = \frac{2\pi\alpha^{-2}}{\xi^3} (\xi - \sin \xi) > 0$$

for all $\xi \in \mathbb{R}$. Thus, $\kappa(T_\alpha) \rightarrow \infty$ as $\alpha \rightarrow \infty$, while $D(T_\alpha) = 2$. A similar example works in any higher dimension.

5 Geometric notation for planar balanced star-shaped domains

We set, for a balanced star-shaped domain $\Omega \subset \mathbb{R}^2$, and $r \geq 0$,

$$\eta(r; \Omega) := \text{vol}_1(\Omega \cap \{|\mathbf{x}| = r\})$$

and

$$\alpha(r; \Omega) := \frac{1}{\text{vol}_2(\Omega)} \int_0^r \eta(\rho; \Omega) d\rho = \frac{\text{vol}_2(\Omega \cap B_2(r))}{\text{vol}_2(\Omega)}$$

(the normalizing factor $1/\text{vol}_2(\Omega)$ will simplify the computations later on).

Let us also define the numbers

$$r_- = r_-(\Omega) = \min_{\mathbf{e} \in S^1} w(\mathbf{e}), \quad r_+ := \max_{\mathbf{e} \in S^1} w(\mathbf{e}).$$

Obviously, r_- is the inradius of Ω and $2r_+$ is its diameter.

Some properties of the functions η and α and the numbers r_{\pm} are obvious:

- Both $\eta(r)$ and $\alpha(r)$ are non-negative; additionally, $\alpha(r)$ is non-decreasing;
- $\eta(r) \equiv 2\pi r$ and $\alpha(r) \equiv \pi r^2 / \text{vol}_2(\Omega)$ for $r \leq r_-$; moreover, $r_- = \sup\{r : \eta(r) = 2\pi r\} = \sup\{r : \alpha(r) = \pi r^2 / \text{vol}_2(\Omega)\}$;
- $\eta(r) \equiv 0$ and $\alpha(r) \equiv \text{const} = 1$ for $r \geq r_+$; moreover, $r_+ = D/2 = \inf\{r : \eta(r) = 0\} = \inf\{r : \alpha(r) = 1\}$ and $\text{supp } \eta = [0, D/2]$.

An additional important property is valid for convex domains.

Lemma 5.1. *Let $\Omega \subset \mathbb{R}^2$ be a balanced convex domain. Then for $r \in [r_-(\Omega), r_+(\Omega)]$, the function $\eta(r)$ is decreasing and the function $\alpha(r)$ is concave.*

Proof. Let us prove that the function η is decreasing in the given interval. Indeed, suppose $r_- < r_1 < r_2$. Since $\eta(r_1) < 2\pi r_1$, we have:

$$(\Omega \cap \{|\mathbf{x}| = r_1\}) \neq \{|\mathbf{x}| = r_1\}.$$

Thus, the set $G := \Omega \cap \{|\mathbf{x}| = r\}$ consists of several (possibly, infinitely many, but at least two) circular arcs, say G_1, \dots, G_n, \dots . Note that G is obviously symmetric with respect to the origin, so if G_j is one of the arcs of G , then the symmetric arc, \tilde{G}_j is also a part of G . Let S_j be the strip based on G_j and \tilde{G}_j (i.e. S_j is the smallest centrally symmetric strip containing G_j and \tilde{G}_j , see Figure 1).

Then a little thought shows that the convexity of Ω implies

$$(\Omega \cap \{|\mathbf{x}| \geq r_1\}) \subset (\cup_j S_j).$$

Thus,

$$\eta(r_2) \leq \text{vol}_1((\cup_j S_j) \cap \{|\mathbf{x}| = r_2\}).$$

However, for each j we have:

$$\text{vol}_1(S_j \cap \{|\mathbf{x}| = r_2\}) < 2 \text{vol}_1 G_j$$

(see figure Figure 1). Summing this over j , we obtain $\eta(r_1) > \eta(r_2)$.

The concavity of α follows immediately from its definition as an integral of η . \square

Remark 5.2. In a similar manner, one can define the analogues of functions η and α in a higher-dimensional setting. Unfortunately, in general, the function η is no longer decreasing on the interval $[r_-, r_+]$; the simplest counterexample is a strip $\Omega = \{\mathbf{x} = (x_1, x_2, x_3) \in \mathbb{R}^3, |x_1| < 1\}$.




Figure 1: Arcs G_j , \tilde{G}_j and strip S_j .

6 Proof of Theorem 2.4

Set

$$\tau := 2j_{0,1}.$$

Without loss of generality we assume that

$$\text{vol}_2(\Omega) = \pi\tau^2 = 4\pi j_{0,1}^2, \tag{6.1}$$

and so $\Omega^* = B(\tau)$. Thus,

$$\kappa(\Omega^*) = \frac{j_{1,1}}{\tau} = \frac{j_{1,1}}{2j_{0,1}} = \frac{1}{\tilde{C}}$$

with \tilde{C} as in (2.4), and in order to prove Theorem 2.4, we need to prove

$$\kappa(\Omega) \leq 1. \tag{6.2}$$

We prove (6.2), and therefore Theorem 2.4 by a sequence of Lemmas. Some of them are rather technical, and for convenience the proofs of these Lemmas are collected in the next section.

First, Theorem 4.1 implies that if $D(\Omega) \geq 4\pi$, then the statement is proved.

Correspondingly, if the half-breadth of Ω in some direction \mathbf{e} , $w(\mathbf{e}) < j_{0,1}^2/2$, then, by Theorem 4.1, the statement is proved since $2r_-D \geq \text{Vol}_2(\Omega)$ as in Corollary 4.4. Thus without loss of generality we can assume that

$$r_+ = D/2 < 2\pi \tag{6.3}$$

and

$$r_- > \frac{j_{0,1}^2}{2}. \tag{6.4}$$

The following averaging result is one of the central points of the proof.

Lemma 6.1. *Suppose that*

$$\int_{\Omega} J_0(|\mathbf{x}|) \, d\mathbf{x} \leq 0. \quad (6.5)$$

Then (6.2) holds.

Proof of Lemma 6.1. To prove (6.2), it is enough to show that there exists $\mathbf{e} \in S^1$ such that

$$\int_{\Omega} \cos(x_{\mathbf{e}}) \, d\mathbf{x} \leq 0.$$

Suppose this inequality is wrong for all $\mathbf{e} \in S^1$. Then

$$\int_{S^1} \int_{\Omega} \cos(x_{\mathbf{e}}) \, d\mathbf{x} \, d\mathbf{e} > 0. \quad (6.6)$$

Changing the order of integration and acting as in in the proof of (3.3), we get

$$\int_{\Omega} J_0(|\mathbf{x}|) \, d\mathbf{x} > 0.$$

Now the Lemma follows by contradiction. \square

We now show that the condition of Lemma 6.1 in fact follows from some integral inequality being satisfied by a class of functions. Namely, consider a class \mathcal{A} of continuous functions $\alpha : [0, \infty) \rightarrow \mathbb{R}$ with the following properties:

- (a) $\alpha(r)$ is non-negative and non-decreasing;
- (b) $\alpha(r) = r^2/(4j_{0,1}^2)$ for $0 \leq r \leq r_-$;
- (c) $\alpha(r) = 1$ for $r \geq r_+$;
- (d) $\alpha(r)$ is concave for $r_- \leq r \leq r_+$;
- (e) $j_{0,1}^2/2 < r_- \leq 2j_{0,1} \leq r_+ < 2\pi$.

Lemma 6.2. *If*

$$\sup_{\alpha \in \mathcal{A}} \int_0^{j_{0,3}} \alpha(r) J_1(r) \, dr \leq 0 \quad (6.7)$$

holds, then (6.2) holds for all planar convex balanced domains Ω normalized by (6.1).

Proof of Lemma 6.2. We continue the calculations in the proof of Lemma 6.1. Using the geometric notation introduced in the previous Section, we have:

$$\int_{\Omega} J_0(|\mathbf{x}|) \, d\mathbf{x} = \int_0^{\infty} \eta(r) J_0(r) \, dr = \text{vol}_2(\Omega) \int_0^{\infty} \alpha(r) J_1(r) \, dr$$

(the last identity is proved by integration by parts using $J_0' = -J_1$).

By Lemma 6.1, we need to show that

$$\int_0^{\infty} \alpha(r) J_1(r) \, dr \leq 0.$$

If for some $k \in \mathbb{N}$, $\alpha(j_{0,k}) = 1$, then $\alpha(r) = 1$ for $r \geq j_{0,k}$, and so, after integration by parts,

$$\int_{j_{0,k}}^{\infty} \alpha(r) J_1(r) \, dr = \int_{j_{0,k}}^{\infty} J_1(r) \, dr = J_0(j_{0,k}) = 0.$$

We need to choose which k to take. In our case $\alpha(r) = 1$ whenever $r \geq D/2$, so by (6.3) we need to choose k such that $j_{0,k} > 2\pi$ and we can take $k = 3$, see the Table 1 below.

Thus, we need to show that

$$I := \int_0^{j_{0,3}} \alpha(r) J_1(r) \, dr \leq 0.$$

The conditions (a)–(d) are just the re-statement of the properties of the function α summarized at the start of the previous Section with account of normalization (6.1); condition (e) re-states (6.4), (6.3), and also the obvious inequalities $\pi r_-^2 \leq \text{vol}_2(\Omega) \leq \pi r_+^2$. \square

It is useful here to plot the function $J_1(r)$ and other quantities appearing above.

For future use, we also give two tables of approximate decimal values of various constants appearing here and below. The first one lists the values appearing along the horizontal axis in various graphs, and the second one lists the values along the vertical axis. In both Tables the values are sorted out in increasing order.

The key points of the proof are the estimates of the function $\alpha(r)$ which are collected in the following sequence of Lemmas.

We start by denoting $y_{1,1} := \alpha(j_{1,1})$, and we also introduce a new constant

$$y_{\min} := 1 - \frac{(2\pi - j_{1,1})(64 - \tau^2)}{8(16\pi - \tau^2)}. \quad (6.8)$$




Figure 2: Graph of $J_1(r)$.

$2\pi - \sqrt{4\pi^2 - \tau^2}$	2.240206980
$\tau^2/8 = j_{0,1}^2/2$	2.891592982
$j_{1,1}$	3.831705970
$\tau = 2j_{0,1}$	4.809651116
2π	6.283185308
$j_{1,2}$	7.015586670
$j_{0,3}$	8.653727913
$2\pi + \sqrt{4\pi^2 - \tau^2}$	10.326163640

Table 1: Decimal values of constants appearing along the horizontal axis.

L (see (6.14))	-0.0852948043
$y_{\min}L + M$ (see (6.16))	-0.0072444612
M (see (6.15))	0.0386824043
$c(j_{1,1}^2/\tau^2) = \frac{\tau^2 - j_{1,1}^2}{\tau^2(2\pi - j_{1,1})}$ (see (6.9))	0.3403496255
y_{\min} (see (6.8))	0.5384485717
$j_{1,1}^2/\tau^2$	0.6346834915

Table 2: Decimal values of constants appearing along the vertical axis.

Lemma 6.3. For the functions $\alpha(r)$ satisfying the conditions (a)–(e) above,

$$y_{1,1} \geq y_{\min}.$$

The proof of this Lemma is in the next section.

Now, given the function $\alpha(r)$ and using the value of $y_{1,1} = \alpha(j_{1,1})$ as a parameter, we construct two new functions. One of them is a linear function $v(r) = c(y_{1,1})r + d(y_{1,1})$, where the coefficients c and d are chosen to be

$$c = c(y_{1,1}) = \frac{1 - y_{1,1}}{2\pi - j_{1,1}}; \quad d = d(y_{1,1}) := 1 - 2\pi c = \frac{2\pi y_{1,1} - j_{1,1}}{2\pi - j_{1,1}}. \quad (6.9)$$

The graph of $v(r)$ is a straight line joining the points $A_1 = (j_{1,1}, \alpha(j_{1,1})) = (j_{1,1}, y_{1,1})$ and $A_+ = (2\pi, \alpha(2\pi)) = (2\pi, 1)$.

The other function is a piecewise-continuous one given by

$$\alpha_{\text{approx}}(r) := \begin{cases} r^2/\tau^2 & \text{for } r \in [0, \tau^2/8]; \\ y_{1,1} & \text{for } r \in (\tau^2/8, j_{1,1}]; \\ v(r) & \text{for } r \in [j_{1,1}, 2\pi]; \\ 1 & \text{for } r \in [2\pi, j_{0,3}]. \end{cases} \quad (6.10)$$




Figure 3: A typical graph of $\alpha(r)$ (solid line) and $\alpha_{\text{approx}}(r)$ (dashed line). The points have coordinates $A_- = (r_-, \alpha(r_-)) = (r_-, r_-^2/\tau^2)$, $A_1 = (j_{1,1}, \alpha(j_{1,1})) = (j_{1,1}, y_{1,1})$, and $A_+ = (2\pi, \alpha(2\pi)) = (2\pi, 1)$.

Obviously, $\alpha(r) \equiv \alpha_{\text{approx}}(r) \equiv 1$ for $r \geq 2\pi$.

Lemma 6.4. *Let $\alpha(r)$ satisfy conditions (a)–(e). Then*

$$\alpha(r) \leq \alpha_{\text{approx}}(r) \quad \text{for } r \in [0, j_{1,1}]. \quad (6.11)$$

and

$$\alpha(r) \geq \alpha_{\text{approx}}(r) \quad \text{for } r \in [j_{1,1}, 2\pi]. \quad (6.12)$$

The proof of this Lemma is in the next section.

Lemma 6.4 immediately implies, with account of the fact that $J_1(r)$ changes sign from plus to minus at $r = j_{1,1}$, the following

Corollary 6.5.

$$\int_0^{j_{0,3}} \alpha(r) J_1(r) dr \leq \int_0^{j_{0,3}} \alpha_{\text{approx}}(r) J_1(r) dr. \quad (6.13)$$

The integral in the right-hand side of (6.13) can be explicitly calculated as a function of the parameter $y_{1,1}$, although the expressions are quite complicated. We introduce two constants,

$$\begin{aligned} L := & J_0\left(\frac{\tau^2}{8}\right) \\ & - \frac{1}{2\pi - j_{1,1}} \left(\pi^2 J_1(2\pi) \mathbf{H}_0(2\pi) - \pi^2 J_0(2\pi) \mathbf{H}_1(2\pi) + \frac{\pi j_{1,1}}{2} J_0(j_{1,1}) \mathbf{H}_1(j_{1,1}) \right. \\ & \left. + j_{1,1} J_0(j_{1,1}) + 2\pi J_0(2\pi) \right) \end{aligned} \quad (6.14)$$

and

$$\begin{aligned} M := & \frac{1}{8} J_2\left(\frac{\tau^2}{8}\right) \\ & + \frac{1}{2\pi - j_{1,1}} \left(\pi^2 J_1(2\pi) \mathbf{H}_0(2\pi) - \pi^2 J_0(2\pi) \mathbf{H}_1(2\pi) + \frac{\pi j_{1,1}}{2} J_0(j_{1,1}) \mathbf{H}_1(j_{1,1}) \right. \\ & \left. - j_{1,1} J_0(j_{1,1}) + 2\pi J_0(2\pi) \right); \end{aligned} \quad (6.15)$$

in the above formulae \mathbf{H} denote the Struve functions [AbrSte, Chapter 12]. The numerical values of L and M can be found in the table above.

Lemma 6.6.

$$\int_0^{j_{0,3}} \alpha_{\text{approx}}(r) J_1(r) dr = Ly_{1,1} + M.$$

With account of Lemmas 6.2, 6.3, and 6.6, and Corollary 6.5 we immediately have

$$\int_0^{j_{0,3}} \alpha(r) J_1(r) dr \leq Ly_{\min} + M \approx -0.007244446126 < 0, \quad (6.16)$$

which finishes the proof of the Theorem.

7 Proofs of Lemmas

Proof of Lemma 6.3. There are two possibilities. If $r_- \geq j_{1,1}$, then $\alpha(j_{1,1}) = \frac{j_{1,1}^2}{\tau^2}$ by condition (b), and the claim of the Lemma is true. We thus need to consider a case when $r_- \leq j_{1,1}$.

Let us introduce a linear function $y(r) := a(r_-)r + b(r_-)$, where the constants a and b depend upon r_- as a parameter and are chosen to be

$$a = a(r_-) = \frac{\tau^2 - r_-^2}{\tau^2(2\pi - r_-)}; \quad b = b(r_-) := 1 - 2\pi a = \frac{r_-(2\pi r_- - \tau^2)}{\tau^2(2\pi - r_-)}. \quad (7.1)$$

The graph of $y(r)$ is a straight line joining the points $A_- = (r_-, \alpha(r_-)) = (r_-, r_-^2/\tau^2)$ and $A_+ = (2\pi, \alpha(2\pi)) = (2\pi, 1)$.

The function $\alpha(r)$ is concave on the interval $[r_-, 2\pi]$ by conditions (c) and (d), and its graph passes through the points A_- and A_+ . Thus, this graph lies above the straight line joining A_- and A_+ , and therefore

$$\alpha(r) \geq y(r) \quad \text{for } r \in [r_-, 2\pi]. \quad (7.2)$$

As $j_{1,1} \in [r_-, 2\pi]$, (7.2) implies

$$\alpha(j_{1,1}) \geq a(r_-)j_{1,1} + b(r_-) = 1 - (2\pi - j_{1,1})a(r_-),$$

and as in our case r_- can take values only in the interval $[\tau^2/8, j_{1,1}]$, we have

$$\alpha(j_{1,1}) \geq 1 - (2\pi - j_{1,1}) \max_{r_- \in [\tau^2/8, j_{1,1}]} a(r_-)$$

We have

$$\frac{da(r_-)}{dr_-} = \frac{r_-^2 - 4\pi r_- + \tau^2}{\tau^2(2\pi - r_-)^2}.$$

The roots of the numerator in the right-hand side are $2\pi \pm \sqrt{4\pi^2 - \tau^2}$, and as seen from the table above the derivative is negative for $r_- \in [\tau^2/8, j_{1,1}]$. Thus

$$y_{1,1} \geq 1 - (2\pi - j_{1,1})a(\tau^2/8). \quad (7.3)$$

It is an easy manipulation to check that the right-hand side of (7.3) equals y_{\min} . \square

Proof of Lemma 6.4. As $\alpha_{\text{approx}}(r) = \alpha(r)$ for r in the interval $[0, \tau^2/8]$, and $\alpha_{\text{approx}}(r) = y_{1,1} = \alpha(j_{1,1})$ for r in the interval $[\tau^2/8, j_{1,1}]$, inequality (6.11) follows immediately from the monotonicity condition (a).

In order to prove (6.12), we again need to consider two cases. First, if $r_- < j_{1,1}$, then $\alpha(r)$ is concave for $r \in [j_{1,1}, 2\pi]$, and its graph between the points A_1 and A_+ lies above the straight line joining this points. Thus, it remains to consider the case $r_- \geq j_{1,1}$ (and so $y_{1,1} = j_{1,1}^2/\tau^2$).

We now show that in this case

$$\frac{r^2}{\tau^2} \geq v(r) = c(j_{1,1}^2/\tau^2)r + d(j_{1,1}^2/\tau^2) \quad \text{for } r \geq j_{1,1}. \quad (7.4)$$

Indeed, consider

$$u(r) := \frac{r^2}{\tau^2} - c(j_{1,1}^2/\tau^2)r - d(j_{1,1}^2/\tau^2).$$

We have

$$u(j_{1,1}) = 0,$$

and also for $r \geq j_{1,1}$,

$$\frac{du}{dr}(r) = 2\frac{r}{\tau^2} - c(j_{1,1}^2/\tau^2) \geq 2\frac{j_{1,1}}{\tau^2} - \frac{\tau^2 - j_{1,1}}{\tau^2(2\pi - j_{1,1})} > 0$$

(see table above for numerical values), which proves (7.4).

Thus, $\alpha(r_-) \geq \alpha_{\text{approx}}(r_-)$. As, by concavity, the graph of $\alpha(r)$ between the points A_- and A_+ lies above the straight line joining these points, and the graph of $\alpha_{\text{approx}}(r)$ is a straight line joining the point $(r_-, \alpha_{\text{approx}}(r_-))$ (which is located below A_-) with A_+ , inequality (6.12) follows. \square

Proof of Lemma 6.6. The result follows from straightforward integration of (6.10) using the standard relations

$$\begin{aligned} \int J_1(x)dx &= -J_0(x) && \text{[AbrSte, formula 11.1.6];} \\ \int xJ_1(x)dx &= -\int xJ_0'(x)dx = -xJ_0(x) + \int J_0(x)dx \\ &= \frac{\pi x}{2} (J_1(x)\mathbf{H}_0(x) - J_0(x)\mathbf{H}_1(x)) && \text{[AbrSte, formula 11.1.7];} \\ \int x^2J_1(x)dx &= x^2J_2(x) && \text{[AbrSte, formula 11.3.20].} \end{aligned}$$

\square

8 Perturbation-type results

In this Section, we prove Theorem 2.7. In order to do this, we need to compute the one-sided derivatives of $\sqrt{\lambda_2(\Omega_{\epsilon F})}$ and $\kappa(\Omega_{\epsilon F})$ with respect to the parameter ϵ describing the deformations of the disk. The first derivative is easily computable from the following classical result (see e.g., [Hen, Rel]) which we state without proof:

Theorem 8.1 (Derivative of a multiple Dirichlet eigenvalue). *Let $\Omega_0 \subset \mathbb{R}^d$ be a bounded domain with C^2 boundary. Assume that $\lambda_k(\Omega_0) = \dots = \lambda_{k+p-1}(\Omega_0)$ is a multiple Dirichlet eigenvalue of order $p \geq 2$. Let us denote by $u_{k_1}, u_{k_2}, \dots, u_{k_p}$ an orthonormal family of eigenfunctions associated to λ_k . Let $\mathbf{S}(t) : \mathbb{R}^d \rightarrow \mathbb{R}^d$, $t \in [0, t_0)$, be a continuously differentiable with respect to t family of mappings such that $\mathbf{S}(0)$ is an identity, and let $\Omega_t = \mathbf{S}(t)(\Omega_0)$. Then, the function $t \rightarrow \lambda_k(\Omega_t)$ has a (directional) derivative at $t = +0$ which is one of the eigenvalues of the $p \times p$ matrix $M = [m_{i,j}]$ defined by*

$$m_{i,j} = - \int_{\partial\Omega} \left(\frac{\partial u_{k_i}}{\partial n} \frac{\partial u_{k_j}}{\partial n} \right) \mathbf{S}'(0)(\sigma) \cdot \mathbf{n}(\sigma) \, d\sigma, \quad i, j = 1, \dots, p, \quad (8.1)$$

where $\mathbf{n}(\sigma)$ is an exterior normal to $\partial\Omega$ at the point $\sigma \in \partial\Omega$.

Theorem 8.1 implies,

Lemma 8.2. *Let $\Omega_{\epsilon F}$ be as in Theorem 2.7. Then*

$$\left. \frac{d\sqrt{\lambda_2(\Omega_{\epsilon F})}}{d\epsilon} \right|_{\epsilon=+0} = -\frac{j_{1,1}}{2\pi} \left| \int_0^{2\pi} F(\theta) e^{2i\theta} \, d\theta \right|. \quad (8.2)$$

Proof of Lemma 8.2. In our case, $\Omega_0 = \Omega_{0F}$ is a disk of radius 1, $\lambda_2(\Omega_0)$ is doubly degenerate, so the matrix M is of dimension 2, and we can choose the orthonormal eigenfunctions $u_2(r, \theta) = N J_1(j_{1,1}r) \cos \theta$, $u_3(r, \theta) = N J_1(j_{1,1}r) \sin \theta$. The constant N is introduced in order for the eigenfunctions u_2 and u_3 to have L_2 -norm one in the unit disk. Using standard properties of Bessel functions (in particular [AbrSte, formulas 11.45 and 9.1.30]) one gets,

$$N = \sqrt{\frac{2}{\pi} \frac{1}{|J_0(j_{1,1})|}}.$$

Using the lowering property of Bessel functions,

$$\frac{1}{z} \frac{d}{dz} (z J_1(z)) = J_0,$$

([AbrSte, formula 9.1.30]) the value of N just obtained, and the fact that $J_1(j_{1,1}) = 0$ and $J_0(j_{1,1}) < 0$, in the expression for u_2 and u_3 we obtain,

$$\frac{\partial u_2}{\partial n} = -j_{1,1} \sqrt{\frac{2}{\pi}} \cos \theta,$$

and

$$\frac{\partial u_3}{\partial n} = -j_{1,1} \sqrt{\frac{2}{\pi}} \sin \theta,$$

at the boundary of the disk (i.e., at $r = 1$).

Taking these remarks into account, the elements of the matrix M in our case are given by

$$m_{1,1} = \frac{2j_{1,1}^2}{\pi} \int_0^\pi F(\theta) \cos^2 \theta \, d\theta,$$

$$m_{1,2} = m_{2,1} = \frac{2j_{1,1}^2}{\pi} \int_0^\pi F(\theta) \cos \theta \sin \theta \, d\theta,$$

and

$$m_{2,2} = \frac{2j_{1,1}^2}{\pi} \int_0^\pi F(\theta) \sin^2 \theta \, d\theta.$$

For area preserving deformations of the ball, i.e. for functions F satisfying (2.8), we can write the matrix M in the simple form,

$$M = \begin{pmatrix} \operatorname{Re} a & \operatorname{Im} a \\ \operatorname{Im} a & -\operatorname{Re} a \end{pmatrix}$$

where

$$a = \frac{j_{1,1}^2}{\pi} \int_0^{2\pi} e^{2i\theta} F(\theta) \, d\theta.$$

It is simple to compute the two eigenvalues of M in this case, and they are given by

$$\pm |a|.$$

Hence, using Theorem 8.1 we obtain the (directional) derivative of the second eigenvalue of the perturbed domain by using a smaller of these two eigenvalues:

$$\left. \frac{d\lambda_2(\Omega_{\epsilon F})}{dt} \right|_{t=0} = -\frac{j_{1,1}^2}{\pi} \left| \int_0^{2\pi} F(\theta) e^{2i\theta} \, d\theta \right|. \quad (8.3)$$

From (8.3), taking into account that $\lambda_2 = j_{1,1}^2$, we finally get (8.2). \square

Now, we will compute the derivative of $\kappa(\Omega_{\epsilon F})$ at $\epsilon = 0$, for area preserving deformations of the disk. For brevity, we shall use the notation $f_\epsilon(\boldsymbol{\xi}) := \widehat{\chi_{\Omega_{\epsilon F}}}(\boldsymbol{\xi})$ for the Fourier transform of the characteristic function of $\Omega_{\epsilon F}$ and $\mathcal{N}_\epsilon = \mathcal{N}(\Omega_{\epsilon F}) = \{\boldsymbol{\xi} \in \mathbb{R}^d : f_\epsilon(\boldsymbol{\xi}) = 0\}$ for its null variety.

We know that \mathcal{N}_0 contains a circle of radius $j_{1,1}$, and we seek to characterize the elements of \mathcal{N}_ϵ . Pick an element $\xi_0 \in \mathcal{N}_0$. For definiteness, we choose coordinates in such a way that

$$\xi_0 = (1, 0)j_{1,1}, \quad (8.4)$$

and we write an element of \mathcal{N}_ϵ as

$$\xi = \xi_0 + \epsilon \xi_1.$$

It is precisely ξ_1 which we would like to determine by requiring $f_\epsilon(\xi) = 0$ to hold up to first order in ϵ . Using polar coordinates, we write

$$\xi_1 = (\cos \omega, \sin \omega)\rho_1.$$

With the above notation, we have

$$f_\epsilon(\xi) = \int_0^{2\pi} \int_0^{1+\epsilon F(\theta)} e^{ij_{1,1}r \cos \theta} e^{i\epsilon \rho_1 r \cos(\theta-\omega)} r \, dr \, d\theta. \quad (8.5)$$

In the sequel, we use the fact that $f_0(j_{1,1}) = 0$, i.e.,

$$\int_0^{2\pi} \int_0^1 e^{ij_{1,1}r \cos \theta} r \, dr \, d\theta = 0, \quad (8.6)$$

and split the integral in the variable r in (8.5) as an integral from $r = 0$ to $r = 1$ plus an integral from $r = 1$ to $r = 1 + \epsilon F(\theta)$. After some algebraic computations we obtain

$$f_\epsilon(\xi) = \epsilon N(\rho_1, \omega) + O(\epsilon^2), \quad (8.7)$$

where

$$\begin{aligned} N(\rho_1, \omega) = i\rho_1 \int_0^{2\pi} \left(\int_0^1 e^{ij_{1,1}r \cos \theta} r^2 \cos(\theta - \omega) \, dr \right) \\ + \int_0^{2\pi} F(\theta) e^{ij_{1,1} \cos \theta} \, d\theta. \end{aligned} \quad (8.8)$$

Using the fact that the perturbed domain is *balanced*, i.e., that $F(\theta) = F(\theta+\pi)$, we get

$$\int_0^{2\pi} F(\theta) e^{ij_{1,1} \cos \theta} \, d\theta = \int_0^{2\pi} F(\theta) \cos(j_{1,1} \cos \theta) \, d\theta.$$

Since we also have $\int_0^{2\pi} e^{ij_{1,1}r \cos \theta} \sin \theta \, d\theta = 0$, we arrive at

$$N(\rho_1, \omega) = i\rho_1 \int_0^{2\pi} \left(\int_0^1 e^{ij_{1,1}r \cos \theta} \cos \theta \cos \omega r^2 \, dr \right) + \int_0^{2\pi} F(\theta) \cos(j_{1,1} \cos \theta) \, d\theta. \quad (8.9)$$

Integrating the first integral in (8.9) by parts in r gives

$$N(\rho_1, \omega) = \frac{\rho_1}{j_{1,1}} \cos \omega \left(\int_0^{2\pi} e^{ij_{1,1} \cos \theta} \, d\theta - \int_0^1 \int_0^{2\pi} 2r e^{ij_{1,1}r \cos \theta} \, dr \, d\theta \right) + \int_0^{2\pi} F(\theta) \cos(j_{1,1} \cos \theta) \, d\theta. \quad (8.10)$$

Now, we use the integral representation

$$J_0(x) = \frac{1}{2\pi} \int_0^{2\pi} e^{ix \cos \theta} \, d\theta$$

to simplify the first two terms in (8.10). We finally get

$$N(\rho_1, \omega) = 2\pi \frac{\rho_1}{j_{1,1}} \cos \omega J_0(j_{1,1}) + \int_0^{2\pi} F(\theta) \cos(j_{1,1} \cos \theta) \, d\theta. \quad (8.11)$$

Here we have used the fact that $\int_0^1 r J_0(j_{1,1}r) \, dr = J_1(j_{1,1})/j_{1,1}^2 = 0$, since $j_{1,1}$ is a zero of J_1 . The vector $\xi_1 = \rho_1(\cos \omega, \sin \omega)$ is determined by the condition

$$N(\rho_1, \omega) = 0.$$

Therefore, (8.11) implies

$$\rho_1 \cos \omega = -\frac{j_{1,1}}{2\pi J_0(j_{1,1})} \int_0^{2\pi} F(\theta) \cos(j_{1,1} \cos \theta) \, d\theta. \quad (8.12)$$

In the case when ξ_0 is not given by (8.4), but by

$$\xi_0 = (\cos \alpha, \sin \alpha) j_{1,1},$$

we have

$$\rho_1 \cos \omega = -\frac{j_{1,1}}{2\pi J_0(j_{1,1})} \int_0^{2\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) \, d\theta. \quad (8.13)$$

In order to compute $\kappa(\Omega_{\epsilon F})$ to first order in ϵ all we have to compute is $|\xi_0 + \epsilon \xi_1|$ to first order in ϵ , which in turn is given by

$$j_{1,1} + \epsilon \rho_1 \cos \omega + O(\epsilon^2).$$

Using (8.13), we obtain

$$\kappa(\Omega_{\epsilon F}) = \min_{\alpha} \left[j_{1,1} \left(1 - \frac{\epsilon}{2\pi J_0(j_{1,1})} \int_0^{2\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta \right) + O(\epsilon^2) \right]. \quad (8.14)$$

From this result, we immediately have,

Lemma 8.3. *Let F be a C^2 function on a unit circle satisfying periodicity condition (2.6). Then*

$$\left. \frac{d\kappa(\Omega_{\epsilon F})}{d\epsilon} \right|_{\epsilon=0} = \min_{\alpha} \left[-\frac{j_{1,1}}{2\pi J_0(j_{1,1})} \int_0^{2\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta \right]. \quad (8.15)$$

Remark 8.4. Note that Lemma 8.3 does not assume the area preservation condition (2.8).

In order to finish the proof of Theorem 2.7 for perturbations around the circle, we need to prove that the right-hand side of (8.15) is always less or equal than the right-hand side of (8.2), i.e.,

$$\min_{\alpha} \int_0^{\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta \leq -A \left| \int_0^{\pi} F(\theta) e^{2i\theta} d\theta \right|, \quad (8.16)$$

where $A = -J_0(j_{1,1}) \approx 0.408\dots$, assuming additionally that F satisfies the area preservation condition (2.8).

For future reference we denote the left-hand side and the right-hand side of (8.16) by L_F and R_F , respectively.

Remark 8.5. Note also that if the average of F is 0 and, additionally, F has zero two-modes, i.e., $\int_0^{\pi} F(\theta) e^{2i\theta} d\theta = 0$, then (8.16) is valid. In fact, in this case, the right-hand side R_F vanishes whereas the left-hand side is given by

$$L_F = \min_{\alpha} \int_0^{\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta,$$

so we have

$$L_F \leq \int_0^{\pi} F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta, \quad (8.17)$$

for every α , and averaging over α we get,

$$\begin{aligned} L_F &\leq \frac{1}{\pi} \int_0^\pi \left(\int_0^\pi F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta \right) d\alpha, \\ &= \frac{1}{\pi} \int_0^\pi d\theta \cos(j_{1,1} \cos \theta) \int_0^\pi F(\theta + \alpha) d\alpha = 0, \end{aligned}$$

and we are done.

Before we conclude, we need to analyze the case of equality in (2.10), i.e., we need to show that equality is only attained in (2.10) if the domain is a ball, or, in other words, if $F(\theta) \equiv 0$. In order to have equality in (2.10), we need equality in (8.17), which in turn implies,

$$\int_0^\pi F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta = 0, \quad (8.18)$$

for all α . Since $F(\theta)$ has zero average, and moreover $F(\theta + \pi) = F(\theta)$ (which is required so that the perturbed domain is balanced), the Fourier series of $F(\theta)$ can be written as

$$F(\theta) = \sum_{k=-\infty}^{\infty} c_k e^{ik\theta} = \sum_{m \neq 0} c_{2m} e^{2im\theta}, \quad (8.19)$$

with $c_0 = 0$ (because F has zero average) and $c_{2k+1} = 0$, for all k (because the domain is balanced). Replacing (8.19) in (8.18) we get,

$$\sum_{m \neq 0} c_{2m} \int_0^\pi e^{2im(\theta+\alpha)} \cos(j_{1,1} \cos \theta) d\theta = 0, \quad (8.20)$$

Using the integral representation for $J_n(z)$, i.e.,

$$J_n(z) = \frac{1}{\pi} \int_0^\pi \cos(z \sin \theta - n\theta) d\theta,$$

after some computation we can write (8.20) as

$$\sum_{m \neq 0} c_{2m} (-1)^m J_{2m}(j_{1,1}) e^{2im\alpha} = 0, \quad (8.21)$$

all $0 \leq \alpha \leq 2\pi$. Since the $\exp(2im\alpha)$ form an orthogonal set of functions, we finally get,

$$c_{2m} (-1)^m J_{2m}(j_{1,1}) = 0,$$

all m . Since $j_{m,1} > j_{1,1}$ for all $m > 1$, $J_{2m}(j_{1,1}) \neq 0$, thus, $c_{2m} = 0$, all m , hence, $F(\theta) \equiv 0$ as it was to be shown. A similar argument can be used to show that equality is attained only for the ball in the general case.

Before we go into the proof of (8.16) for a general F satisfying both the periodicity condition (giving a balanced domain) and the zero average condition (area preserving domain perturbation), we need the following result.

Lemma 8.6. *Assuming F averages up to zero, it is always possible to rotate F in such a way that the following two conditions are fulfilled simultaneously:*

$$\int_0^\pi F(\theta + \phi) \sin(2\theta) d\theta = 0, \quad (8.22)$$

and

$$\int_0^\pi F(\theta + \phi) \cos(2\theta) d\theta \geq 0. \quad (8.23)$$

Here $F(\theta + \phi)$ is F rotated by an angle ϕ .

Proof of Lemma 8.6. Consider the function

$$T(\phi) := \int_0^\pi F(\theta + \phi) \sin(2\theta) d\theta.$$

Since F averages to zero, $\int_0^\pi T(\phi) d\phi = 0$, so there exists a point $\phi_1 \in [0, \pi]$, such that $T(\phi_1) = 0$, and (8.22) holds.

Now, consider

$$Q(\phi) := \int_0^\pi F(\theta + \phi) \cos(2\theta) d\theta.$$

Clearly, $T(\phi_1) = T(\phi_1 + \pi/2) = 0$. On the other hand, $Q(\phi_1) = -Q(\phi_1 + \pi/2)$. So, either $Q(\phi_1) \geq 0$, or $Q(\phi_1 + \pi/2) \geq 0$, and we have obtained (8.23) by choosing $\phi = \phi_1$ or $\phi = \phi_1 + \pi/2$. \square

After proving this Lemma we are ready to prove (8.16). Consider F with zero average and such that $\int_0^\pi F(\theta) \cos(2\theta) d\theta \geq 0$ and $\int_0^\pi F(\theta) \sin(2\theta) d\theta = 0$. In this case, the right-hand side of (8.16) is given by

$$R_F = -A \int_0^\pi F(\theta) \cos(2\theta) d\theta. \quad (8.24)$$

On the other hand, the left-hand side L_F satisfies (8.17) for each α . Now, multiply (8.17) by

$$\cos^2 \alpha / \int_0^\pi \cos^2 \alpha d\alpha \equiv (2/\pi) \cos^2 \alpha$$

and integrate in α from 0 to π (notice that $(2/\pi) \cos^2 \alpha \geq 0$). We thus have,

$$L_F \leq \frac{2}{\pi} \int_0^\pi \left(\int_0^\pi F(\theta + \alpha) \cos(j_{1,1} \cos \theta) d\theta \right) \cos^2 \alpha d\alpha. \quad (8.25)$$

Now, split $\cos^2 \alpha = (e^{2i\alpha} + e^{-2i\alpha} + 2)/4$ in (8.25). If we do the integral over α first, using the fact that the average of F vanishes, we get

$$\begin{aligned} \int_0^\pi F(\theta + \alpha) \cos^2 \alpha d\alpha &= \frac{1}{4} \int_0^\pi e^{2i\alpha} F(\theta + \alpha) d\alpha + \frac{1}{4} \int_0^\pi e^{-2i\alpha} F(\theta + \alpha) d\alpha \\ &= \frac{1}{4} e^{-2i\theta} \int_0^\pi e^{2i\beta} F(\beta) d\beta + \frac{1}{4} e^{2i\theta} \int_0^\pi e^{-2i\beta} F(\beta) d\beta. \end{aligned}$$

By Lemma 8.6, and the choice of orientation of F , we have

$$\int_0^\pi e^{2i\beta} F(\beta) d\beta = \int_0^\pi e^{-2i\beta} F(\beta) d\beta = \int_0^\pi \cos(2\beta) F(\beta) d\beta =: P,$$

and so

$$\int_0^\pi F(\theta + \alpha) \cos^2 \alpha d\alpha = \frac{1}{2} \cos(2\theta) P.$$

Then,

$$L_F \leq \frac{2}{\pi} \frac{1}{2} \int_0^\pi \cos(2\theta) \cos(j_{1,1} \cos \theta) d\theta P = P J_0(j_{1,1}). \quad (8.26)$$

Here we have used the fact that $\int_0^\pi \cos(2\theta) \cos(j_{1,1} \cos \theta) d\theta = \pi J_0(j_{1,1})$ (this follows by taking real part in [AbrSte, formula 9.1.21], with $n = 2$, and the fact that $J_2(j_{1,1}) = -J_0(j_{1,1})$ [AbrSte, 9.1.27]). Hence,

$$L \leq J_0(j_{1,1}) P = -A P = R_F.$$

This proves (8.16) and therefore Theorem 2.7. \square

9 Non-convex domains: counterexamples

We start by proving Theorem 2.8.

First, we introduce some notation. For a domain Ω we put

$$\zeta(r) = \zeta_\Omega(r) := \frac{\eta(r)}{2\pi r}.$$

Then the function ζ satisfies the following properties: $\zeta(r) \leq 1$; if Ω is star-shaped, ζ is non-increasing, $\text{supp } \zeta = \text{supp } \eta = [0, D(\Omega)/2]$, and $\text{vol}_2(\Omega) = 2\pi \int_0^{D/2} r\zeta(r)dr$. The strategy of the proof is the following: first, we construct a function $\tilde{\zeta}$ which is non-increasing, $\tilde{\zeta}(r) = 1$ for $0 \leq r \leq 1 - \tilde{\delta}$, $\text{supp } \tilde{\zeta} \subset [0, 1 + \tilde{\delta}]$, $\int_0^{1+\tilde{\delta}} r\tilde{\zeta}(r)dr = \pi$ and, finally,

$$\int_0^{\infty} r\tilde{\zeta}(r)J_0(\gamma r)dr > 0 \quad (9.1)$$

for all $\gamma \leq j_{1,1}$. Then we construct a domain Ω such that $\tilde{\zeta} = \zeta_{\Omega}$ and $\int_{\Omega} \cos(\gamma x_{\mathbf{e}})d\mathbf{x}$ is close to the left-hand side of (9.1) for all \mathbf{e} , $|\mathbf{e}| = 1$ and all $\gamma \leq j_{1,1}$. This Ω will be a required domain.

Let $\zeta_0(r) = \begin{cases} 1, & 0 < r \leq 1 \\ 0, & r > 1 \end{cases}$ be the ζ -function for the ball of radius one.

Suppose that $\tilde{\delta}$ is fixed. Let δ be a small positive parameter, and put

$$\xi_{\delta}(r) = \begin{cases} -1/2, & 1 - \delta < r < 1, \\ a, & 1 < r < 1 + \tilde{\delta}, \\ 0, & \text{otherwise.} \end{cases}$$

Here, we choose $a = a(\delta)$ from the condition

$$\int_0^{\infty} r\xi_{\delta}(r)dr = 0, \quad (9.2)$$

which is equivalent to

$$a = \frac{\int_{1-\delta}^1 r dr}{2 \int_1^{1+\tilde{\delta}} r dr}. \quad (9.3)$$

Obviously, $a \rightarrow 0$ as $\delta \rightarrow 0$. Note also that for small δ we have

$$\int_0^{\infty} r\xi_{\delta}(r)J_0(j_{1,1}r)dr > 0. \quad (9.4)$$

Indeed, we obviously have

$$\frac{d \int_0^1 r\xi_{\delta}(r)J_0(j_{1,1}r)dr}{d\delta} = -\frac{1}{2} \frac{d \int_{1-\delta}^1 rJ_0(j_{1,1}r)dr}{d\delta} = -\frac{J_0(j_{1,1})}{2},$$

so

$$\int_0^1 r\xi_{\delta}(r)J_0(j_{1,1}r)dr \sim \frac{-J_0(j_{1,1})\delta}{2}. \quad (9.5)$$

Similarly, using (9.3) we obtain

$$\begin{aligned} b &:= \frac{d \int_1^\infty r \xi_\delta(r) J_0(j_{1,1}r) dr}{d\delta} = \frac{d(a \int_1^{1+\tilde{\delta}} r J_0(j_{1,1}r) dr)}{d\delta} \\ &= \frac{\int_1^{1+\tilde{\delta}} r J_0(j_{1,1}r) dr}{2 \int_1^{1+\tilde{\delta}} r dr} \frac{d \int_{1-\delta}^1 r dr}{d\delta} = \frac{\int_1^{1+\tilde{\delta}} r J_0(j_{1,1}r) dr}{2 \int_1^{1+\tilde{\delta}} r dr}, \end{aligned} \quad (9.6)$$

so

$$\int_1^\infty r \xi_\delta(r) J_0(j_{1,1}r) dr \sim b\delta \quad (9.7)$$

as $\delta \rightarrow 0$. Since $j_{1,1}$ is a local minimum of J_0 , we have $b > \frac{J_0(j_{1,1})}{2}$. Now formulas (9.5) and (9.7) imply (9.4). We now fix a small $\delta < \tilde{\delta}$ for which (9.4) holds and put $\tilde{\zeta}(r) = \zeta_0(r) + \xi_\delta(r)$. Then, since

$$\int_0^\infty r \zeta_0(r) J_0(j_{1,1}r) dr = 0, \quad (9.8)$$

we have

$$\int_0^\infty r \tilde{\zeta}(r) J_0(j_{1,1}r) dr > 0. \quad (9.9)$$

It is easy to show that in fact for all positive $\gamma \leq j_{1,1}$ we have

$$\int_0^\infty r \tilde{\zeta}(r) J_0(\gamma r) dr > 0. \quad (9.10)$$

Indeed, the function

$$l(\gamma) := \int_0^\infty r \tilde{\zeta}(r) J_0(\gamma r) dr$$

decreases for $\gamma < j_{1,1}$, since its derivative

$$l'(\gamma) = - \int_0^\infty r^2 \tilde{\zeta}(r) J_1(\gamma r) dr$$

is negative as $J_1(r)$ is positive for $r \in [0, j_{1,1}]$. Note also that (9.2) implies

$$\int_0^\infty r \tilde{\zeta}(r) dr = 1. \quad (9.11)$$

Now let us construct a sequence of domains Ω_n which satisfy the following properties:

- (i) the domain Ω_n is invariant under the rotation on $\frac{2\pi}{n}$ around the origin;
- (ii) in the sector $-\frac{\pi}{n} \leq \phi \leq \frac{\pi}{n}$ in polar coordinates (r, ϕ) the domain Ω_n is given by $\{(r, \phi), |\phi| < \frac{\pi \tilde{\zeta}(r)}{n}\}$.

Thus, for large n the domain Ω_n has many thin spikes, see Figure 4 for the picture of such a domain.




Figure 4: Domain Ω_n .

It is obvious that these properties determine the domain Ω_n uniquely and, moreover, that $\zeta_{\Omega_n} = \tilde{\zeta}$ for all n . Note also that for all positive $\gamma \leq j_{1,1}$ and all unit vectors \mathbf{e} we have

$$\int_{\Omega_n} \cos(\gamma x_{\mathbf{e}}) d\mathbf{x} \rightarrow \int_0^{\infty} r \tilde{\zeta}(r) J_0(\gamma r) dr \tag{9.12}$$

as $n \rightarrow \infty$ uniformly over γ and \mathbf{e} . Therefore, (9.10) implies that for sufficiently large n

$$\int_{\Omega_n} \cos(\gamma x_{\mathbf{e}}) d\mathbf{x} > 0 \tag{9.13}$$

for all positive $\gamma \leq j_{1,1}$ and all unit vectors \mathbf{e} . Thus, for this domain Ω_n we have $\kappa(\Omega_n) > j_{1,1}$, finishing the proof of Theorem 2.8.

It remains to prove Theorem 2.9 and Corollary 2.10. Suppose that we have proved Theorem 2.9, and thus constructed a sequence I_n of one-dimensional balanced domains for which $\kappa(I_n) \text{vol}_1(I_n) \rightarrow \infty$ as $n \rightarrow \infty$. Consider

$$A_n := (I_n)^2 = \{\mathbf{x} = (x_1, x_2), x_1, x_2 \in I_n\}. \tag{9.14}$$

Then $\text{vol}_2(A_n) = (\text{vol}_1(I_n))^2$, and $\kappa(A_n) = \kappa(I_n)$, and so $\kappa(A_n) \sqrt{\text{vol}_2(A_n)} \rightarrow \infty$ as $n \rightarrow \infty$. Now it remains to connect the disjoint rectangles in A_n by narrow

corridors to construct connected domains \tilde{A}_n with $\kappa(\tilde{A}_n)\sqrt{\text{vol}_2(\tilde{A}_n)} \rightarrow \infty$ as $n \rightarrow \infty$, proving Corollary 2.10.

Let us prove Theorem 2.9. We formulate the following

Lemma 9.1. *For each positive C there exist a natural number n and real numbers w_1, \dots, w_n such that $w_1 \geq 1$, $w_{j+1} \geq w_j + 1$, and the function $f(\xi) := \sum_{j=1}^n \cos(w_j \xi)$ is positive for $\xi \in [-C/n, C/n]$.*

Proof. The proof is due to F. Nazarov [Naz]. Put, for real t , $g(t) := (1 - |t|)_+^2$. Then the Fourier transform $\hat{g}(s)$ of g is positive for real s . Denote, for real x ,

$$G(x) := 1 + 2 \sum_{k=1}^n \left(1 - \frac{k}{n}\right)^2 \cos(kx) = \sum_{k \in \mathbb{Z}} g\left(\frac{k}{n}\right) e^{ikx}. \quad (9.15)$$

Then the Poisson summation formula implies that

$$G(x) = n \sum_{m \in \mathbb{Z}} \hat{g}(n(x + 2\pi m)) \geq n\hat{g}(nx), \quad (9.16)$$

and so $G(x) \geq cn$ whenever $|x| \leq C/n$. Now put

$$F(x) := \sum_{k=1}^n a_k^n \cos(kx), \quad (9.17)$$

where a_k^n is a collection of independent random variables such that $a_k^n = 1$ with probability $(1 - \frac{k}{n})^2$; otherwise $a_k^n = 0$. Then the standard probabilistic arguments based on the large deviation principle imply that for each fixed point x the probability of the event

$$|F(x) - (G(x) - 1)/2| \geq n^{3/4} \quad (9.18)$$

is $O(e^{-n^{1/4}})$. In particular, putting $x = 0$ in (9.18), we see that the number of coefficients a_k^n which are equal to one, is at least $n/10$ with probability $1 - O(n^3 e^{-n^{1/4}})$. Put $x_j := \frac{j}{n^3}$, $j = 0, \dots, n^3$. Then the probability of the event that for all $j = 0, \dots, n^3$ we have

$$|F(x_j) - (G(x_j) - 1)/2| \leq n^{3/4} \quad (9.19)$$

is at least $1 - O(n^3 e^{-n^{1/4}})$ and thus is positive for sufficiently large n . Since the derivative of both F and G is $O(n^2)$, this means that the probability that (9.18) is satisfied for all $x \in [0, 1]$ is positive when n is large. Therefore, for each large n there is at least one F such that (9.18) is satisfied for all $x \in [0, 1]$. Thus chosen F satisfies $F(x) \geq cn/2$ for $|x| \leq C/n$. \square

To finish the proof of Theorem 2.9, we now take, for a given $C > 0$, the numbers n and w_j , $j = 1, \dots, n$, from Lemma 9.1, and define $I_n := \{x \in \mathbb{R}, ||x| - w_j| \leq 1/2 \text{ for some } j\}$. Then $\text{vol}_1(I_n) \leq 2n$, and

$$\widehat{\chi_{I_n}}(\xi) = \frac{4 \sin(\xi/2)}{\xi} \sum_{j=1}^n \cos(w_j \xi).$$

Therefore by Lemma 9.1 for any constant C we have $\kappa(I_n) \geq C/n$ for sufficiently large n .

References

- [AbrSte] Abramowitz, M., and Stegun, I. A., **Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables**, Dover, N. Y., 1964.
- [Agr] Agranovsky, M. L., *On the stability of the spectrum in the Pompeiu problem*, J. Math. Anal. Appl. **178**, no. 1, 269–279 (1993).
- [AshBen] Ashbaugh, M. S., and Benguria, R. D., *Proof of the Payne-Pólya-Weinberger conjecture*, Bull. Amer. Math. Soc. (N.S.) **25**, no. 1, 19–29 (1991).
- [Avi] Aviles, P., *Symmetry theorems related to Pompeiu's problem*, Amer. J. Math. **108**, no. 5, 1023–1036 (1986).
- [Ber] Berenstein, C. A., *An inverse spectral theorem and its relation to the Pompeiu problem*, J. Analyse Math. **37**, 128–144 (1980).
- [BroKah] Brown, L., and Kahane, J.-P., *A note on the Pompeiu problem for convex domains*, Math. Ann. **259**, no. 1, 107–110 (1982).
- [BroSchTay] Brown, L., Schreiber, B. M., and Taylor, B. A., *Spectral synthesis and the Pompeiu problem*, Ann. Inst. Fourier (Grenoble) **23**, no. 3, 125–154 (1973).
- [Fil1] Filonov, N., *On an inequality for the eigenvalues of the Dirichlet and Neumann problems for the Laplace operator*, Algebra i Analiz **16**, no. 2, 172–176 (2004) (Russian); translation in St. Petersburg Math. J. **16**, no. 2, 413–416 (2005).
- [Fil2] Filonov, N., *Private communication* (2008).
- [Fri] Friedlander, L., *Some inequalities between Dirichlet and Neumann eigenvalues*, Arch. Rational Mech. Anal. **116**, no. 2, 153–160 (1991).

- [GarSeg] Garofalo, N., and Segàla, F., *Asymptotic expansions for a class of Fourier integrals and applications to the Pompeiu problem*, J. Analyse Math. **56**, 1–28 (1991).
- [Hen] Henrot, A., **Extremum Problems for Eigenvalues of Elliptic Operators**, Frontiers in Mathematics, Birkhäuser, Basel, 2006.
- [Kob1] Kobayashi, T., *Bounded domains and the zero sets of Fourier transforms*, **75 years of Radon transform (Vienna, 1992)**, 223–239, Conf. Proc. Lecture Notes Math. Phys., IV, Int. Press, Cambridge, MA, 1994.
- [Kob2] Kobayashi, T., *Perturbation of domains in the Pompeiu problem*, Comm. Anal. Geom. **1**, no. 3-4, 515–541 (1993).
- [LevWei] Levine, H. A., and Weinberger, H. F., *Inequalities between Dirichlet and Neumann eigenvalues*, Arch. Rational Mech. Anal. **94**, 193–208 (1986).
- [Lor] Lorch, L., *Some inequalities for the first positive zeros of Bessel functions*, SIAM J. Math. Anal. **24**, 814–823 (1993).
- [Naz] Nazarov, F., *Private communication* (2008).
- [PayPólWei] Payne, L. E., Pólya, G., and Weinberger, H. F., *Sur le quotient de deux fréquences propres consécutives*, C. R. Acad. Sci. Paris **241**, 917–919 (1955); and *On the ratio of consecutive eigenvalues*, J. Math. and Phys. **35**, 289–298 (1956).
- [Rel] Rellich, F., **Perturbation theory of eigenvalue problems**, Gordon and Breach, NY, 1969.
- [Zas] Zastavnyi, V.P., *Zero set of the Fourier transform of measures and the summation of double Fourier series of methods of Bernshtein–Ragozinskii type*, Ukr. Math. J. **36**, 459–464 (1984).