

A universal primer for isolation of fragments of a gene encoding phytoene desaturase for use in virus-induced gene silencing (VIGS) studies

Article

Accepted Version

Yin-Chih, L., Imsabai, W., Reid, M. S., Cai-Zhong, J. and Wagstaff, C. ORCID: <https://orcid.org/0000-0001-9400-8641> (2010) A universal primer for isolation of fragments of a gene encoding phytoene desaturase for use in virus-induced gene silencing (VIGS) studies. *Acta Horticulturae*, 877. pp. 1697-1704. ISSN 0567-7572 Available at <https://centaur.reading.ac.uk/18147/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: http://www.actahort.org/books/877/877_233.htm

Publisher: International Society for Horticultural Science

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A Universal Primer for Isolation of Fragments of a Gene Encoding Phytoene Desaturase for Use in Virus-Induced Gene Silencing (VIGS) Studies

L. Yin-Chih, W. Imsabai and M.S. Reid
Department of Plant Sciences
University of California
One Shields Avenue
Davis, CA 95616
USA

C. Wagstaff
Food Biosciences Department
University of Reading, Whiteknights
PO Box 217, Reading
Berkshire, RG6 6AH
United Kingdom

J. Cai-Zhong
Crops Pathology and Genetic Research Unit
Agricultural Research Service
US Department of Agriculture
Davis, CA 95616
USA

Keywords: *PDS* (phytoene desaturase), VIGS (virus-induced gene silencing), TRV (tobacco rattle virus), PCR

Abstract

We have been using Virus-Induced Gene Silencing (VIGS) to test the function of genes that are candidates for involvement in floral senescence. Although VIGS is a powerful tool for assaying the effects of gene silencing in plants, relatively few taxa have been studied using this approach, and most that have are in the *Solanaceae*. We typically use silencing of phytoene desaturase (PDS) in preliminary tests of the feasibility of using VIGS. Silencing this gene, whose product is involved in carotene biosynthesis, results in a characteristic photobleaching phenotype in the leaves. We have found that efficient silencing requires the use of fragments that are more than 90% homologous to the target gene. To simplify testing the effectiveness of VIGS in a range of species, we designed a set of universal primers to a region of the PDS gene that is highly conserved among species, and that therefore allows an investigator to isolate a fragment of the homologous PDS gene from the species of interest. We report the sequences of these primers and the results of VIGS experiments in horticultural species from the *Asteraceae*, *Leguminosae*, *Balsaminaceae* and *Solanaceae*.

INTRODUCTION

Virus-induced gene silencing (VIGS) is a method of transiently interrupting gene function through RNA interference. The VIGS process is a natural plant defense mechanism and is initiated experimentally by *Agrobacterium*-mediated introduction of viral genomes containing a fragment of a targeted plant gene. When the virus replicates and spreads in the infected plants, the plant cells recognize the foreign double-stranded chimeric RNAs synthesized during viral replication and cut them into small oligonucleotides (siRNA). The siRNA molecules that serve as guides for an RNA-induced silencing complex (RISC) that recognizes homologous gene transcripts (Baulcombe, 1999; Chicas and Macino, 2001) targeting them for hydrolysis and thereby reduces expression of the target gene.

VIGS provides a rapid method for loss of gene function assay, avoiding the need for time-consuming transformation and regeneration processes. In addition, VIGS provides the opportunity to overcome functional redundancy by suppressing all or most members of a gene family (Burch-Smith et al., 2004). Vectors based on tobacco rattle virus (TRV) with a wide host range have been shown to function effectively in *Nicotiana benthamiana* (Liu et al., 2002a), tomato (Liu et al., 2002b), petunia (Chen et al., 2004),

poppy (Wege et al., 2007) and *Aquilegia* (Gould and Kramer, 2007) for phytoene desaturase (*PDS*) silencing. Infected plants show characteristic photo-bleaching symptoms resulting from the decreased production of photoprotective carotenoid proteins and the subsequent breakdown of chlorophyll pigments. This phenotypic marker can be a potential reporter for examining the effect of silencing other target genes. In these previous studies, researchers reported that a highly homologous gene fragment was required to obtain high efficiency silencing. However, the sequence of *PDS* is not known for many species. In this paper, we describe a universal primer for isolation of *PDS* fragment from different plant species and report its successful use in a range of taxa.

MATERIALS AND METHODS

Plant Material

Seedlings of a range of species including petunia (*Petunia × hybrida*) ‘Primetime Blue’; tomato (*Lycopersicon esculentum*) ‘New Yorker’; Yesterday, today and tomorrow (*Brunfelsia glandiflora*); daisy ‘Alaska’; gerbera (*Gerbera × hybrida*); chrysanthemum (*Dendranthema grandiflorum*) ‘Veracruz’; lettuce (*Lactuca sativa*) ‘Red Sails’; impatiens (*Impatiens wallerana*) ‘Orange Bling’; Phalaenopsis orchid (*Phalaenopsis amabilis*); rose (*Rosa × hybrida*) ‘Magnum’; four o’clock (*Mirabilis jalapa*); and bean (*Phaseolus vulgaris*) ‘Kentucky Blue’ and ‘Golden Wax’ were grown under 16h light/8h dark cycles at approximately 20-25°C.

Plasmid Construct

PDS constructs: A 520 bp fragment of the *PDS* gene was amplified from leaf cDNA of each species using primers 5’-GGTGGAAAG (g/a) T (a/g/t) GCTGC (a/t/c) TGG-3’ and 5’-GCCAT (t/c) TT (g/a/t) GAACCATG (t/c) TT (t/c) TCCTG-3’. The resulting product was cloned into pGEM-T easy (Promega) for amplification, sequencing and cloning. The *PDS* cDNA fragment was excised using *EcoRI* digestion from the plasmid, then subcloned into pTRV2 to generate a pTRV2 *PDS* construct specific to each species.

Agrobacterium-Mediated Infection

The constructs, pTRV1 (TRV RNA1 construct), and pTRV2 (RNA2 (empty vector control) or RNA2+*PDS* fragment) constructs were transformed into *Agrobacterium* strain GV3101 by electroporation. The bacteria were cultured overnight at 28°C in LB medium with 40 mg/L Kanamycin and 20 mg/L Gentamycin. The *Agrobacterium* cells were then harvested and resuspended in inoculation buffer (10 mM MgCl₂, 10 mM MES, 200 μM acetosyringone) to an O.D. of 1~2 and left at room temperature for at least 4h. The bacteria containing pTRV1 and the bacteria containing pTRV2 or pTRV2 *PDS* were mixed together in a 1:1 ratio. The leaves of seedlings were inoculated with the mixed bacteria culture using a 1 ml disposable syringe without a needle (for daisy and tomato) or by vacuum infiltration (for lettuce and impatiens). Infected plants were incubated in growth chambers under 16h light/8h dark cycles at the temperature of 20-22°C.

RT-PCR Analysis

Total RNA was extracted from lettuce leaves using TRIzol Reagent (Invitrogen) and treated with RNase-free DNase (Ambion) to remove any contaminating genomic DNA. The first strand cDNA was synthesized using 2 μg total RNA, oligo d(T) primer, random hexamer, and superscript reverse transcriptase (Invitrogen). This cDNA was normalized by real-time quantitative PCR using 18S rRNA primer (5’-CATGGCCGTTCTT AGTTGGTGGAG-3’ and 5’-AAGAAGCTGGCCGCGAAGGGAT AC-3’).

The primers for amplifying *PDS* transcripts by semi-quantitative PCR were designed outside the region targeted for gene silencing to avoid amplification from the pTRV *PDS* construct. These primers were 5’-CTTTGCAAGCCAATTGTTGA-3’ and 5’-TTGTAGGGGTTCGACATGGTT-3’.

RESULTS AND DISCUSSION

Design of Primers to Amplify *PDS* Gene Fragments from Different Plant Species

A BLAST search for *PDS* sequences from a wide range of taxa permitted us to design a set of degenerate primers that would amplify a *PDS* fragment from known species, and therefore, most probably from species where the gene's sequence is unknown. An approximately 520 bp fragment of the *PDS* gene was amplified by the degenerate primers from all tested species (Fig. 1). A comparison of the *PDS* fragment sequences from 9 different plant species reveals a similarity of higher than 80% but in most cases less than the 90% homology required for efficient VIGS (Fig. 2). It seems likely that most other species will include the highly conserved region of the *PDS* gene amplified by this pair of degenerate primers.

Host Range of TRV-Induced *PDS* Gene Silencing

Fragments isolated from the target species using our universal primers were very effective when used in VIGS silencing of *PDS*. Developing leaves of pTRV2-*PDS* infected plants showed the strong photobleaching phenotype that indicated movement of the virus and its effectiveness in silencing the native *PDS* gene in bean, impatiens, daisy, and tomato (Fig. 3). No photo-bleached phenotype was found in the H₂O or pTRV2 empty vector controls (data not shown).

PDS gene transcripts were clearly evident in RNA extracted from non-silenced leaves and vector control, but barely detected in that from photo-bleaching leaves of lettuce by RT-PCR (Fig. 4). The result suggested that the photo-bleaching phenotype was caused by the down-regulation of the *PDS* gene.

In previous VIGS studies of a wide range of different ornamentals (Chen et al., 2003), only *N. benthamiana* and *P. hybrida* (both from the *Solanaceae*) showed the *PDS* phenotype. We now suspect that this negative result reflected a lack of homology between the *PDS* of *N. benthamiana* (the source of the *PDS* fragment used in the study) and the *PDS* sequences in the other target species.

Efficiency of VIGS in Different Species

Although we were able to silence *PDS* in all of the species that we tested, efficiency (% plants showing phenotype), and extent of silencing varied among taxa. Increasing the concentration of the *Agrobacterium* inoculum (to an OD₆₀₀ of 2.0) increased the frequency of silencing in impatiens plants (Fig. 5). In most tested species, however, the silencing symptom was local, confined to one or two leaves (Fig. 3). To obtain systemic symptoms in such species, it may be necessary to develop techniques to improve viral movement, and/or inhibit the plant's antiviral mechanism.

ACKNOWLEDGEMENTS

We gratefully acknowledge the support of the American Floral Endowment, the USDA Floriculture Initiative, and the US-Israel Binational Agricultural Research and Development (BARD, IS-3815-05) and a BBSRC LINK grant in collaboration with Vitacress Salads Ltd. (CW).

Literature Cited

- Baulcombe, D.C. 1999. Fast forward genetics based on virus-induced gene silencing. *Current Opinion in Plant Biology* 2:109-123.
- Burch-Smith, T.M., Anderson, J.C., Martin, G.B. and Dinesh-Kumar, S.P. 2004. Applications and advantages of virus-induced gene silencing for gene function studies in plants. *Plant Journal* 39(5):734-746.
- Chen, J.C., Jiang, C.Z., Gookin, T.E., Hunter, D.A., Clark, D.G. and Reid, M.S. 2004. Chalcone synthase as a reporter in virus-induced gene silencing studies of flower senescence. *Plant Mol. Biol.* 55:521-530.
- Chen, J.C., Johnson, F., Clark, D.G., Gookin, T. and Reid, M.S. 2003. Potential

- application of virus-induced gene silencing (VIGS) in flower senescence studies. *Acta Hort.* 669:147-152.
- Chicas, A. and Macino, G. 2001. Characteristics of post-transcriptional gene silencing. *EMBO Reports* 2:992-996.
- Gould, B. and Kramer, E.M. 2007. Virus-induced gene silencing as a tool for functional analyses in the emerging model plant *Auilegia* (columbine, *Ranunculaceae*). *Plant Methods* 3:6.
- Liu, Y., Schiff, M., Marathe, R. and Dinesh-Kumar, S.P. 2002a. Tobacco Rar1, EDS1 and NPR1/NIM1 like genes are required for N-mediated resistance to tobacco mosaic virus. *Plant J.* 30:415-429.
- Liu, Y., Schiff, M. and Dinesh-Kumar, S.P. 2002b. Virus-induced gene silencing in tomato. *Plant J.* 31:777-786.
- Wege, S., Scholz, A., Gleissberg, S. and Becke, A. 2007. Highly efficient virus-induced gene silencing (VIGS) in California poppy (*Eschscholzia californica*): An evaluation of VIGS as a strategy to obtain functional data from non-model plants. *Annals of Botany* 100:641-649.

Figures




Fig. 1. *PDS* gene fragments were amplified by the universal primers from different species. 1: Petunia; 2: Tomato; 3: Brunfelsia; 4: Daisy; 5: Gerbera; 6: Chrysanthemum; 7: Lettuce; 8: Impatiens; 9: Orchid; 10: Rose; 11: Mirabilis; 12: Pole bean; 13: Bush bean. The size of the *PDS* fragment is approximately 520 bp (arrow).

Rose	GGTGGAAAG-ATTGCTGCTTGGAAAGATAAAGATGGAGACTGGTATGAGACAGGCGCTACA
Mirabilis	GGTGGAAAG-ATTGCTGCTTGGAAAGATGAAGATGGAGATTGGTATGAACTGGACTACA
Impatiens	GGTGGAAAG-GTTGCTGCTTGGAAAGATGAGGATGGAGACTGGTATGAAACAGGCTCTACA
Bean	GGTGGAAAG-GTTGCTGCTTGGAAAGATGAGGATGGAGACTGGTATGAAACAGGCTCTACA
Mum	GGTGGAAAG-ATTGCTGCTTGGAAAGACGAGGATGGAGATTGGTACGAGACGGGTTTACA
Daisy	GGTGGAAAG-ATTGCTGCTTGGAAAGACGAGGATGGAGATTGGTACGAGACGGGTTTACA
Gerbera	GGTGGAAAG-ATTGCTGCTTGGAAAGATGATGATGGAGATTGGTACGAGACAGGTTTACA
Tomato	GGTGGAAAG-GTAGCTGCATGGAAAGATGATGATGGAGATTGGTACGAGACTGGTTTGCA
Brunfelsia	GGTGGAAAG-ATTGCTGCTTGGAAAGATGACGATGGAGATTGGTATGAGACTGGTTTGCA ***** * ****
Rose	TATATTTTTTGGGGCTTATCCAAATATTCAAACTTGTTTGGAGAGCTTGGTATCAATGA
Mirabilis	CATATTTCTTGGGGCTTACCCAAATATTCAAACTTGTTTGGAGAGCTTGGTATTAATGA
Impatiens	CATATTTCTTGGGGCTTACCCGAATGTGCAGAACCTTTTGGAGAACTTGGCATCAATGA
Bean	CATCTTCTTGGGGCTTACCTAATGTGCAGAACCTTATTGGAGAACTTGGCCTTAATGA
Mum	CATATTTTTTGGAGCTTACCCAAATGTACAGAACCTGTTTGGAGAGCTAGGCATTAACGA
Daisy	CATATTTTTTGGAGCTTATCCAAATGTACAGAACCTGTTTGGAGAGCTAGGCATTAACGA
Gerbera	CATATTTCTTGGGGCTTACCCAAATATACAGAACCTTTTGGAGAACTAGGCATTAATGA
Tomato	TATATTTCTTGGGGCTTACCCAAATATACAGAACCTGTTTGGAGAACTAGGCATTAACGA
Brunfelsia	CATATTTCTTGGGGCTTACCCAAATATACAGAACCTGTTTGGAGAACTAGGCATTAATGA * * * * *
Rose	TCGATTGCAGTGAAGGAACACTCTATGATATTGCAATGCCAAACAGCCAGGAGATT
Mirabilis	TCGGTTGCAATGGAAGGAGCACTCCATGATTTTCGCAATGCCAAAGTAAGCCGGTGAATT
Impatiens	CAGGCTCCAAATGGAAGGAACATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Bean	TCGGTTACAAATGGAAGGAGCATTTCATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Mum	TAGATTCGATTGGAGGAGCATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Daisy	TAGATTGCAGTGAAGGAGCATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Gerbera	AAGATTGCAGTGAAGGAGCATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Tomato	TCGATTGCAGTGAAGGAGCATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Brunfelsia	TCGATTGCAGTGAAGGAGCATTCTATGATTTTTCGCAATGCCAAATAGCCTGGAGAAAT * * * * *
Rose	CAGCCGCTTTGATTTCCCTGAAGTCTGCCAGCACCTTAAATGGAATTTGGGCGATATT
Mirabilis	CAGCCGCTTTGATTTTCCCTGAAGTCTGCCAGCACCTTAAATGGAATTTGGGCGATATT
Impatiens	CAGCCGATTTGACTTCCCGATGTTTTCGCAATGCCAAATAGCCTGGAGAAAT
Bean	TAGTCGATTTGATTTCCCGAAATCCTTCGCTCCCATTAATGGAATCTGGGCGATATT
Mum	TAGTCGCTTTGACTTCCCGATGTTTTCGCTCCCATTAATGGAATCTGGGCGATATT
Daisy	TAGTCGCTTTGACTTCCCGATGTTTTCGCTCCCATTAATGGAATCTGGGCGATATT
Gerbera	TAGTCGCTTTGACTTCCCGATGTTTTCGCTCCCATTAATGGAATCTGGGCGATATT
Tomato	CAGCCGCTTTGATTTTCCCGAAGCTTACCCGCTCCTTTAAATGGAATTTAGCGATCTT
Brunfelsia	CAGCCGCTTTGATTTTCCCGAAGCTTACCCGCTCCTTTAAATGGAATTTAGCGATCTT * * * * *
Rose	AAAGAACAATGAGATGCTGACATGGCCAGAAAAAGTGAAGTTTGCCTATCGGACTTGTGCC
Mirabilis	AAGGAACAATGAAATGCTAATCTGGCCGAGAAAAAGTGAAGTTTGCCTATAGGCTCTTGCC
Impatiens	ACCGAATAATGAAATGCTTACATGGCCGAGAAAAAGTGAAGTTTGCCTATAGGCTCTTGCC
Bean	GAGGAACAATGAGATGCTGACATGGCCAGAGAAAAATCAAAATTTGCAATTTGGGCTTCTGCC
Mum	GAGGAACAATGAAATGCTGACATGGCCAGAGAAAAATCAAAATTTGCAATTTGGGCTTCTGCC
Daisy	GAGGAACAATGAAATGCTGACATGGCCAGAGAAAAATCAAAATTTGCAATTTGGGCTTCTGCC
Gerbera	GAGGAACAATGAAATGCTGACATGGCCAGAGAAAAATCAAAATTTGCAATTTGGGCTTCTGCC
Tomato	AAAGAATAACGAAATGCTTACATGGCCAGAGAAAAAGTGAAGTTTGCCTATCGGACTTGTGCC
Brunfelsia	AAAGAACAATGAAATGCTTACATGGCCAGAGAAAAAGTGAAGTTTGCCTATCGGACTTGTGCC * * * * *
Rose	AGCAATTTCTTGGTGGACAGGCTTATGTTGAAGCTCAGGATGGCTTGACTGTAAAGGAGTG
Mirabilis	TGCTATTGTGCGTGGACAAGATTATGTTGGTGGCTCAAGATGGTTTGAAGTGTCCAAAGAGTG
Impatiens	AGCTATGAAATGCTTGGACAAGATTATGTTGGTGGCTCAAGATGGTTTGAAGTGTCCAAAGAGTG
Bean	AGCTATGCTTTGCTGACAGGCTATGTTGAGGCTCAAGATGGTCTTCTGCTTAAGAGATG
Mum	TGCAATGTTAGGTGGGCGAGGCTATGTTGAGGCTCAAGATGGGCTAAGTGTTCAGACTG
Daisy	TGCAATGTTAGGTGGGCGAGGCTATGTTGAGGCTCAAGATGGGCTAAGTGTTCAGACTG
Gerbera	AGCAATGTTAGGAGGACAGGCTTATGTTGAGGCTCAAGATGGTTTGAAGTGTTCAGACTG
Tomato	AGCAATGCTTGGAGGGCAATCTTATGTTGAAGCTCAAGATGGGATAAGTGTTCAGACTG
Brunfelsia	TGCAATGCTTGGAGGGCAATCTTATGTTGAAGCTCAAGACGGGATAAGGCTTAAGGACTG * * * * *
Rose	GATGACAAAACAGGGGATACCTGATCGAGTAACTACTGAGGTGTTTATGCCATGTCAAA
Mirabilis	GATGAGGAAGCAAGGTGTACCTGATCGAGTAACTACTGAGGTGTTTATGCCATGTCAAA
Impatiens	GATGCAAAAACAGGATATACCGAGAGCTGTAACCTACTGAGGTGTTTATGCCATGTCAAA
Bean	GATGATAAAACAGGGTGTACCTGAACGAGTAACTGACGAGGTGTTTATGCAATGTCAAA
Mum	GATGAGACAGCGAGGCATACCGGATCGAGTCAACCCGAGGTGTTTATGCCATGTCAAA
Daisy	GATGAGACAGCGAGGCATACCGGATCGAGTCAACCCGAGGTGTTTATGCCATGTCAAA
Gerbera	GATGAGAAAGCAAGGCATACCGATCGGGTACTACTGAGGTGTTTATGCCATGTCAAA
Tomato	GATGAGAAAGCAAGGTGTGCCGAGAGGTGACAGATGAGGTGTTTATGCTATGTCAAA
Brunfelsia	GATGAGAAAGCAAGGTGTGCCGATAGGTTGACAGATGAGGTGTTTATGCTATGTCAAA * * * * *
Rose	GGCCCTTAACCTTATTAATCCTGATGAGCTCTCAATGCAATGCATATTGATTGCTTTGAA
Mirabilis	GGCATTGAACCTTATAAATCCGATGAGCTCTCAATGCAATGCATATTGATTGCTTTGAA
Impatiens	GGCTTTAAACTTCATTAAACCTGACGAGCTTTCAATGCAATGTATTTGATTGCTCTGAA
Bean	GGCATTAACCTTCATCAATCCTGATGAACCTTCAATGCAATGTATTTGATTGCTTTAAA
Mum	GGCATTAACCTTCATAAATCCTGATGAACCTTCAATGCAATGTATTTCTCATAGCTCTAAA
Daisy	GGCATTAACCTTCATAAATCCTGATGAACCTTCAATGCAATGTATTTCTCATAGCTCTAAA
Gerbera	GGCGTTAAACTTCATAAATCCAGATGAACCTTCTATGCAATGTATTTCTCATGCTTTGAA
Tomato	GGCACTCAACTTTATAAACCTGACGAACCTTCAATGCAATGCATTTTATGCGATTGAA
Brunfelsia	AGCACTTAACCTTCATAAACCTGACGAGCTTCAATGCAATGCATTTGATTGCTTTGAA * * * * *
Rose	TCGATTCTTTCAGGAAAAACATGGTTCAAAAATGGCA
Mirabilis	TCGATTCTTTCAGGAAAAACATGGTTCAAAAATGGCA
Impatiens	TCGGTTTCTTTCAGGAAAAACATGGTTCAAAAGTGGCA
Bean	CCGATTCTTTCAGGAAAAACATGGTTCTAAAATGGCA
Mum	CCGGTTTCTTTCAGGAAAAACATGGTTCAAAAGTGGCA
Daisy	CCGGTTTCTTTCAGGAAAAACATGGTTCTAAAATGGCA
Gerbera	CAGGTTTCTTTCAGGAAAAACATGGTTCAAAAATGGCA
Tomato	CAGGTTTCTTTCAGGAAAAACATGGTTCAAAAATGGCC
Brunfelsia	CAGATTCTTTCAGGAAAAACATGGTTCAAAAATGGCA * * * * *

Fig. 2. Multiple sequence alignment of the *PDS* fragments isolated from the 9 target plant species.




Fig. 3. Photobleaching phenotype resulting from VIGS silencing of *PDS* in different plant species using fragments isolated with the universal primers. A: Henderson's Baby Bush Lima Bean, B: Daisy 'Alaska', C: Impatiens 'Orange Bling', D: Tomato 'New Yorker'.




Fig. 4. Abundance of *PDS* transcripts in 'Red Sails' lettuce plants after VIGS silencing. P-silenced tissue, NP-non-silenced tissue, E-empty vector control.



Fig. 5. Effect of *Agrobacterium* concentration on silencing PDS in impatiens plants.

