

# *Geographical limits of the Southeastern distribution of Aedes aegypti (Diptera, Culicidae) in Argentina*

Article

Accepted Version

Diaz-Nieto, L. M., Macia, A., Perotti, A. and Beron, C. M. (2013) Geographical limits of the Southeastern distribution of *Aedes aegypti* (Diptera, Culicidae) in Argentina. PLoS Neglected Tropical Diseases, 7 (1). e1963. ISSN 1935-2735 doi: <https://doi.org/10.1371/journal.pntd.0001963> Available at <http://centaur.reading.ac.uk/29437/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1371/journal.pntd.0001963>

Publisher: Public Library of Science

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1       **Geographical limits of the Southeastern distribution of *Aedes aegypti* (Diptera,**  
2                                   **Culicidae) in Argentina**

3  
4       **Leonardo M. Díaz-Nieto<sup>1</sup>, Arnaldo Maciá<sup>2</sup>, M. Alejandra Perotti<sup>3</sup>, Corina M.**  
5       **Berón<sup>1\*</sup>**

6  
7       **1** Centro de Estudios de Biodiversidad y Biotecnología (CEBB-CIB-FIBA-Mar del  
8       Plata) – CONICET, Mar del Plata, Argentina, **2** División Entomología, Facultad de  
9       Ciencias Naturales y Museo de La Plata (UNLP), La Plata, Argentina, **3** School of  
10      Biological Sciences, University of Reading, Reading, UK

11     \* E-mail: cberon@fiba.org.ar

12  
13  
14     **Background**

15           *Aedes aegypti* (Linnaeus) is a human-biting mosquito and the primary vector of  
16      human dengue and yellow fever viruses; it is also considered the principal vector of  
17      Chikungunya virus in Asia [1,2]. In particular, dengue and dengue hemorrhagic fever  
18      constitute an important burden to humankind in terms of morbidity and mortality. About  
19      3.6 billion people in the tropics, mainly in Asia, the Western Pacific region, the  
20      Caribbean, as well as Central and South America, live under risk of infection with one  
21      or more of the four dengue virus serotypes (DEN-1 to DEN-4), and recent reports  
22      estimate over 230 million infections, over 2 million cases of the severe form of the  
23      disease and 21,000 deaths [3].

24           It is believed that *A. aegypti* originally migrated from West Africa to the New  
25      World beginning with the 15th century aboard slave ships, and after that yellow fever

26 appeared in the new world. Presumably the yellow fever virus was introduced by  
27 travellers on these ships, especially African slaves. The adaptation of this insect to  
28 survive in human environments was crucial for colonization and development in water  
29 storage containers in the holds of sailing ships [4]. At present, *A. aegypti* lives in close  
30 proximity to people, in urban areas, breeding in all types of domestic and peridomestic  
31 collections of fresh water, including flower vases, water drums, tins, broken coconut  
32 shells, old tyres and gutters. A major range of expansion of *Aedes* mosquitoes into these  
33 urban areas is also attributable to the adaptation of the genera *Aedes* to breed in water-  
34 holding automobile tyres [5].

35 *A. aegypti* is a tropical and subtropical species spanning a geographical  
36 distribution from 35°N to 35°S. Its lower thermal threshold corresponds to 10°C  
37 isotherms during the winter, and although it has been found up to 45°N, its presence in  
38 colder regions is due to its ability to colonise new areas during the warm season [6]. In  
39 South America, the historic direction of dispersal of *Aedes* mosquitoes has been towards  
40 higher latitudes and from tropical to sub-tropical areas, in particular in the Southern  
41 Cone. We propose, that the south eastern movement of *A. aegypti* might be related to  
42 human migrations from rural areas to towns lacking in a proper housing policy and  
43 essential services like water, and sewage disposal systems  
44 ([http://www.migraciones.gov.ar/pdf\\_varios/estadisticas/Patria\\_Grande.pdf](http://www.migraciones.gov.ar/pdf_varios/estadisticas/Patria_Grande.pdf)) [7].

45 Between the 1950s, 1960s and most of the 1970s, in Central and South America  
46 epidemic dengue was rare because *A. aegypti* had been eliminated from most of the  
47 countries. The eradication program organized by the Pan American Health Organization  
48 (PAHO) was discontinued in the early 1970s, and consequently the mosquito was  
49 reintroduced in countries from which it had been eradicated [6,8]. In Argentina, the  
50 earliest records of *A. aegypti* go back to the 1900s and are concurrent with the dengue-

51 like epidemic of 1916, which affected the coastal areas of the Uruguay River (31°44'S,  
52 60°31'W) [9]. However, in 1986 re-infestation took place in the northern border with  
53 Paraguay, deriving in its spread over wide areas of the country. Nowadays, the current  
54 geographical distribution of *A. aegypti* in Argentina is wider than during its eradication  
55 in 1967 [10,11]. Recently it has been demonstrated that the three *A. aegypti* main  
56 haplogroups identified in Argentina would represent different colonization events,  
57 probably from neighboring countries: Bolivia, Paraguay, and Brazil (Fig. 1A and B) [7].  
58 Particularly, in Buenos Aires province, the most densely populated area of the country,  
59 the records of high abundances of well-established populations of *A. aegypti* were taken  
60 in La Plata (capital of the province) and in Buenos Aires (capital city of the country),  
61 both located on the East coast, and the southernmost findings in Chascomús, from 132  
62 Km from Buenos Aires city (35°33'S, 58°00'W, Fig 1) [10-15].

63 On the other hand, cases of dengue have increased in the last few years in  
64 Argentina. From January to June 2012, 2,043 patients with symptoms were reported,  
65 and 194 were confirmed with serotypes DEN-1, DEN-2 or DEN-3  
66 ([http://www.msal.gov.ar/dengue/images/stories/partes\\_dengue/parte74.pdf](http://www.msal.gov.ar/dengue/images/stories/partes_dengue/parte74.pdf)). In 2011  
67 PAHO emitted an epidemiological alert due to the introduction of DEN-4 serotype in  
68 the Americas (<http://new.paho.org>), being Brazil, Paraguay and Bolivia countries of  
69 high risk of dengue infection, with 57,267 possible cases and 5 deaths (Brazil); 10,827  
70 suspected cases, 30 victims (Paraguay) and 3,233 notified cases with 28 deaths  
71 (Bolivia) (Fig 1B)  
72 ([http://www.msal.gov.ar/dengue/images/stories/partes\\_dengue/parte74.pdf](http://www.msal.gov.ar/dengue/images/stories/partes_dengue/parte74.pdf)).

73 In the USA, the dispersal of *Aedes albopictus* Skuse offered an opportunity to  
74 understand the synanthropic behavior of *Aedes* mosquitoes. The mosquito was  
75 introduced in 1985 in the continental territory through shipments of used tyres

76 containing eggs originated in Asia [16]. In subsequent years, the pattern of spread of  
77 this container-dwelling species followed the main interstate highways [17], quickly  
78 reaching and colonizing several new areas of the USA in a few years. We wondered  
79 whether *A. aegypti* would present a similar behavior, and is making use of human  
80 transportation [18]. For this, we investigated the occurrence of the mosquito in major  
81 roads connecting densely populated cities with the Southeast of Argentina (Table 1).

82         One of the most important highways in Argentina is the Provincial Route N°2,  
83 connecting Buenos Aires and La Plata cities with Mar del Plata city and the most visited  
84 beaches of the country, principally in summer time, representing about two million  
85 people commuting between those places (Fig 1C and Table 1)  
86 (<http://www.indec.mecon.ar>). Route N°2 crosses the most prominent wetland areas of  
87 the Pampas, and its construction has definitely reshaped the landscape, making available  
88 new manmade wetlands, which offer shelter to an increasing diversity of flora and  
89 fauna, including mosquitoes [19]. On this artery there are some small towns that offer  
90 several travel services such as tyre-repair stations or “gomerías”, which store used  
91 automobile and truck tyres for long periods of time, thus these tyres accumulate  
92 rainwater (Fig 2 and 3A). Moreover, along this highway a lot of vehicles transport  
93 goods from the north of the country to the coastal area without any sanitary control to  
94 prevent insects exchange from one region to the other. The latest scientific  
95 southernmost record of *A. aegypti* carried on in Buenos Aires province, was obtained in  
96 Chascomús a town located on Route N°2 [11]. Route N°2 takes the bulk of the traffic  
97 and people in south-eastern direction. On the other hand, Route N°11, connecting  
98 Buenos Aires and La Plata cities with the Atlantic coast, is a short motorway parallel to  
99 the coastline and Route N°226 runs south-west and is mostly used by freight transport  
100 (Fig 1C).

101           **Present distribution of *A. aegypti* in the most populated areas of the Buenos**  
102 **Aires province**

103           In order to understand the status of the southern distribution of *A. aegypti*, we  
104 sampled mosquito larvae and pupae during the rain period, in January and March 2011,  
105 and only in March 2012 because rainfall levels were very low in January (Fig. 2). The  
106 sampling stations were located in towns situated along Route N°2 and the other two  
107 major arteries that connect Buenos Aires with the South. The sampling stations were  
108 cemeteries, that are far from the towns and are shortly visited and “gomerías” located in  
109 densely populated areas of each town, both at the edge of the roads (Fig. 3 and Table 2).  
110 Larval specimens were collected and reared until fourth instar or adult stage to facilitate  
111 identification using specific keys [13,20]. Voucher specimens, prepared from all  
112 localities, were submitted to the local museum, Museo de Ciencias Naturales “Lorenzo  
113 Scaglia” (Mar del Plata, Argentina).

114           Larvae of *A. aegypti* were found in March 2011 and 2012 in Chascomús,  
115 agreeing with and confirming previous records [11,13]. Here we report the finding of *A.*  
116 *aegypti* in the towns of Lezama, Castelli and Dolores for the first time. A population of  
117 mosquitoes was found in Lezama in March 2011, 39.2 Km southeast of Chascomús,  
118 being both localities separated by farmland and uniquely connected by Route N°2. As a  
119 high number of larvae of all stages and pupae were found in multiple containers in this  
120 locality, we feel confident that Lezama holds a natural, well-established population. In  
121 March 2012, we found a higher number of larvae of all stages and pupae in the same  
122 type of containers for a second time in Lezama; and for the first time in Castelli (27.7  
123 Km south from Lezama) and Dolores (59.5 Km south from Lezama), stating Dolores  
124 the southernmost limit of the species’ range within Argentina, now 98.7 Km south from

125 Chascomús (Fig 1C). In Routes N°11 and 226 *A. aegypti* was not found in any of the  
126 water containers examined.

127 In the south of Argentina *A. aegypti* is very likely to be moving by passive  
128 dispersal using the major highway connecting the North with the Southeast of the  
129 country. It is noteworthy that this same behavior has been studied and documented in a  
130 closely related species, *A. albopictus* in the USA. Previous observations on this  
131 mosquito in North America are consistent with the hypothesis of mosquito migration  
132 facilitated by anthropic action, presumably by transportation of scrapped tyres through  
133 the interstate highway system [17]. In *A. aegypti*, egg resistance in absence of water, a  
134 feature shared with *A. albopictus*, can lead to a similar way of transferring to new places  
135 in order to breed. Therefore, passive dispersal of *Aedes* species using frequented  
136 freeways should be considered at the time of designing new monitoring programs.

137 According to Shepherd *et al.* [21] dengue virus transmission follows two general  
138 patterns: epidemic dengue and hyperendemic dengue. Epidemic dengue transmission  
139 occurs when dengue virus is introduced into a region as an isolated event that involves a  
140 single viral strain. If the number of vectors and susceptible hosts are sufficient,  
141 explosive transmission can occur with an infection incidence of 25-50%. Hyperendemic  
142 dengue transmission is characterized by the circulation of multiple viral serotypes in an  
143 area with susceptible hosts and competent vector (with or without seasonal variation)  
144 and appears to be a major risk for dengue hemorrhagic fever. Travelers to these areas  
145 are more likely to be infected than travelers going to areas that experience only  
146 epidemic transmission.

147 In South America, particularly in Buenos Aires Province, it is known that the  
148 provincial Health Ministry has a program of surveillance of *A. aegypti*, which involves  
149 mosquito larvae and eggs monitoring and their control. However, this surveillance does



150 not follow a regular pattern, being erratic in terms of time and each council or  
151 municipality decides to carry it on or not. In addition, to obtain official data from  
152 concrete actions is sometimes unlikely to find.

153 The new biogeographical record of Central and Southern Argentina, reported in  
154 this article, is an important fact of the constant expansion of *A. aegypti* into new  
155 southernmost areas. Together with the presence of the different dengue serotypes  
156 indicate that the situation is far more dangerous than previously thought. Urgent and  
157 responsible actions must be taken to control the Dengue vector and its further expansion  
158 into new areas.

159

## 160 **Acknowledgments**

161 We especially thank Prof. Ana Tassi for critical revision of the manuscript.  
162 LMDN is a PhD student at the National University of Mar del Plata, and recipient of the  
163 ANPCyT doctoral scholarship (Argentina). AM is Research Assistant of Comisión de  
164 Investigaciones Científicas (CIC), Argentina. CMB is a Research Fellow. This work is  
165 part of LMDN's PhD thesis.

166

## 167 **References**

- 168 1. Becker N, Petrić D, Zgomba M, Boase C, Dahl C, et al. (2003) Mosquitoes and  
169 their control. New York. Kluwer Academic/ Plenum Publishers 498 p.
- 170 2. Pialoux G, Gaüzère BA, Jauréguiberry S, Strobel M (2007) Chikungunya, an  
171 epidemic arbovirosis. Lancet Infect Dis 7:319-327.
- 172 3. Gubler DJ (2012) The economic burden of dengue. Am J Trop Med Hyg  
173 86:743-744.

- 174 4. Lounibos LP (2002) Invasions by insect vectors of human disease. *Annu Rev*  
175 *Entomol* 47: 233-266.
- 176 5. Berry WJ, Craig GB (1984) Bionomics of *Aedes atropalpus* breeding in scrap  
177 tires in northern Indiana. *Mosq News* 44: 476–484.
- 178 6. Pan-American Health Organization (1994) Dengue and dengue hemorrhagic  
179 fever in the Americas: Guidelines for Prevention and Control. Washington:  
180 PAHO Scientific Publication N° 548. 98 pp.
- 181 7. Albrieu Llinás G, Gardenal CN (2012) Phylogeography of *Aedes aegypti* in  
182 Argentina: long-distance colonization and rapid restoration of fragmented relicts  
183 after a continental control campaign. *Vector Borne Zoon Dis* 12: 254–261.
- 184 8. Gubler DJ (1998) Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev.*  
185 11:480-496.
- 186 9. Gaudino NM (1916) Dengue. *Rev San Mil* 15: 617-627.
- 187 10. Curto SI, Boffi R, Carbajo AE, Plastina R, Schweigmann N (2002)  
188 Reinfestación del territorio argentino por *Aedes aegypti*. Distribución geográfica  
189 (1994-1999). In: Salomón OD. Actualizaciones en Artropodología sanitaria  
190 Argentina. Buenos Aires: Fundación Mundo Sano. pp. 127-137.
- 191 11. Rossi GC, Lestani EA, D’Oria JM (2006) Nuevos registros y distribución de  
192 mosquitos de la Argentina (Diptera: Culicidae). *Rev Soc Entomol Argent* 65:  
193 51-56.
- 194 12. Maciá A (2006) Differences in performance of *Aedes aegypti* larvae raised at  
195 different densities in tires and ovitraps under field conditions in Argentina. *J*  
196 *Vector Ecol* 31: 371-377.

- 197 13. Rossi GC, Mariluis JC, Schnack JA, Spinelli GR (2002) Dípteros vectores  
198 (Culicidae y Calliphoridae) de la provincia de Buenos Aires. La Plata:  
199 COBIOBO N° 4. PROBIOTA N° 3. 45 p.
- 200 14. Schweigmann N, Orellano P, Kuruc J, Vera TM, Vezzani D, et. al. (2002)  
201 Distribución y abundancia de *Aedes aegypti* (Diptera: Culicidae) en la ciudad de  
202 Buenos Aires. In: Salomón OD. Actualizaciones en Artropodología sanitaria  
203 Argentina. Buenos Aires: Fundación Mundo Sano. pp. 155-160.
- 204 15. Vezzani D, Carbajo AE (2008) *Aedes aegypti*, *Aedes albopictus*, and dengue in  
205 Argentina: current knowledge and future directions. Mem Inst Oswaldo Cruz  
206 103: 66-74.
- 207 16. Sprenger D, Wuithiranyagool T (1986) The discovery and distribution of *Aedes*  
208 *albopictus* in Harris County, Texas. J Am Mosq Control Assoc 2: 217-9.
- 209 17. Moore CG, Mitchell CJ (1997) *Aedes albopictus* in the United States: Ten-Year  
210 Presence and Public Health Implications. Emerg Infec Dis 3: 329-34.
- 211 18. Hemme RR, Thomas CL, Chadee DD, Severson DW (2010) Influence of urban  
212 landscapes on population dynamics in a short-distance migrant mosquito:  
213 evidence for the dengue vector *Aedes aegypti*. PLoS Negl Trop Dis 4: e634.
- 214 19. Schnack JA, de Francesco FO, Colado UR, Novoa ML, Schnack EJ (2000)  
215 Humedales antrópicos: su contribución para la conservación de la biodiversidad  
216 en los dominios subtropical y pampásico de la Argentina. Ecología Austral 10:  
217 63-80.
- 218 20. Darsie RF, Mitchell CJ (1985) The mosquitoes of Argentina. In: Lewis T.  
219 Nielsen. Mosquito systematics. Utah: American Mosquito control Association.  
220 pp. 153-253.

221 21. Shepherd SM, Hinfey PB, Shoff WH (2009) Dengue. Available from:  
222 <http://emedicine.medscape.com/article/215840-overview>.

223

## 224 **Legends to figures**

225

226 **Figure 1. *Aedes aegypti* and dengue fever in South America. (A)** Historical  
227 distribution of *A. aegypti* in Argentina, indicating: 1916, first dengue-like  
228 epidemic; 1986 re-infestation places and biogeographical records between  
229 1991 and 1999, **(B)** Current geographic distribution of *A. aegypti* and regions  
230 with risk of transmission of dengue in South America, **(C)** Studied area, showing  
231 highways between Buenos Aires and Mar del Plata cities, sampling points and  
232 distances between them. (A and B) adapted from Curto *et al.*, Vezzani and  
233 Carbajo, [10,15] and <http://www.healthmap.org/dengue/index.php>.

234

235 **Figure 2. Weather conditions of the studied area, from July 2010 to June**  
236 **2012.** On the left mean temperature in °C (T), on the right % of relative humidity  
237 (RH) and total precipitation in mm (PP). <http://www.tutiempo.net/clima>. Arrows  
238 indicate sampling times.

239

240 **Figure 3. Sampling places in Buenos Aires province: (A)** Tyre-repair  
241 stations showing tyres with accumulated rainwater, **(B)** flowerpots at  
242 cemeteries.

243

244  
245  
246  
247

**Table 1.** Information of cities connected by Route N° 2 in Buenos Aires Province  
(<http://www.censo2010.indec.gov.ar/>).

<b>City</b>	<b>Area km<sup>2</sup></b>	<b>Population size</b>	<b>Number of households</b>
Buenos Aires	2,681	12,801,365	3,147,638
Chascomús	3,452	38,477	18,277
Lezama	1,102	4,111	nd <sup>c</sup>
Castelli	2,063	8,206	3,448
Dolores	1,973	26,601	10,687
General Guido	2,814	2,814	1,508
Maipú	2,641	10,172	4,375
Mar del Plata	1,461	618,989	308,570
MdP, Summer time <sup>a</sup>	1,461	2,000,000	nd <sup>c</sup>

248  
249  
250  
251  
252

<sup>a</sup>MdP, Mar del Plata.

<sup>c</sup>No data

253  
254

**Table 2.** Sampling stations and species collected in cities along Route N° 2, in the Southeast of Argentina.




City	2011			2012								
	Flowerpots <sup>a</sup>	<i>Culex</i> sp.	<i>A. aegypti</i>	Tyre-repair stations <sup>b</sup>	<i>Culex</i> sp.	<i>A. aegypti</i>	Flowerpots <sup>a</sup>	<i>Culex</i> sp.	<i>A. aegypti</i>	Tyre-repair stations <sup>b</sup>	<i>Culex</i> sp.	<i>A. aegypti</i>
<b>Chascomús</b>	239 (12/0)	+	-	1 (1/1)	+	+	300 (8/2)	+	+	2 (2/2)	+	+
<b>Lezama</b>	200 (0/0)	-	-	3 (3/1)	+	+	200 (5/0)	+	-	3 (3/3)	+	+
<b>Castelli</b>	480 (0/0)	-	-	3 (3/0)	+	-	200 (3/2)	+	+	3 (3/3)	+	+
<b>Dolores</b>	730 (29/0)	+	-	2 (2/0)	+	-	400 (12/1)	+	+	2 (2/2)	+	+
<b>Gral. Guido</b>	280 (1/0)	+	-	2 (2/0)	+	-	300 (7/0)	+	-	3 (3/0)	+	-
<b>Maipú</b>	440 (5/0)	+	-	2 (2/0)	+	-	nd <sup>c</sup>	nd <sup>c</sup>	nd <sup>c</sup>	2 (1/0)	+	-
<b>Mar del Plata</b>	3,600 (~45/0)	+	-	10 (8/0)	+	-	3,600 (~45/0)	+	-	10 (8/0)	+	-


255  
256  
257  
258  
259  
260

<sup>a</sup> Number of flowerpots sampled, in brackets positive ones for *Culex sp* and for *A. aegypti* respectively.

<sup>b</sup> The number of *A. aegypti* was 500 larvae or more in each tyre-repair station, in brackets positive ones for *Culex sp* and for *A. aegypti* respectively.

<sup>c</sup> No data





262

263





A



B