

*A dramatic isotope effect in the reaction of
ClSiH with trimethylsilane-1-d:
experimental evidence for intermediate
complexes in silylene Si-H(D) insertion
reactions*

Article

Accepted Version

Becerra, R., Boganov, S. E., Krylova, I. V., Promyslov, V. M. and Walsh, R. (2011) A dramatic isotope effect in the reaction of ClSiH with trimethylsilane-1-d: experimental evidence for intermediate complexes in silylene Si-H(D) insertion reactions. *Organometallics*, 30 (16). pp. 4225-4227. ISSN 1520-6041 doi: <https://doi.org/10.1021/om200411x> Available at <http://centaur.reading.ac.uk/32063/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1021/om200411x>

Publisher: American Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A Dramatic Isotope Effect in the Reaction of ClSiH with Trimethylsilane-1-d: Experimental Evidence for Intermediate Complexes in Silylene Si-H(D) Insertion Reactions

Rosa Becerra^{*a}, Sergey E. Boganov^b, Irina V. Krylova^b, Vladimir M. Promyslov^b and Robin Walsh^{*c}


^a *Instituto de Química-Física "Rocasolano", C.S.I.C., C/ Serrano 119, 28006 Madrid, Spain,*

^b *N.D. Zelinsky Institute of Organic Chemistry, Russian Academy of Sciences, Leninsky Prospekt 47, 119991 Moscow, Russian Federation and* ^c *Department of Chemistry, University of Reading, Whiteknights, P.O. Box 224, Reading RG6 6AD, UK.*

ABSTRACT: A kinetic isotope effect (k_D/k_H) of 7.4 has been found for the reaction of chlorosilylene with trimethylsilane (Me_3SiD vs Me_3SiH). Such a value can be accounted for by theoretical modelling, but only if an internal rearrangement of the initially formed complex is included in the mechanism. This provides the first concrete evidence for such complexes.


The insertion reaction of silylenes into Si-H bonds is one of the fundamental and most widely studied reactions of silylenes.¹ The mechanism of this process is thought to involve weakly bound intermediate complexes.² The evidence for this comes largely from the interpretation of gas-phase kinetic studies, allied to theoretical (ab initio) calculations. The main arguments in favour of such complexes are the existence of low energy minima on the calculated potential energy surfaces³ for such reactions and the experimental finding of negative activation energies^{3,4}. Despite these arguments the experimental evidence may still be regarded as inconclusive. The complexes themselves have not been seen directly nor has any process unique to the complex itself been identified. We report here an experiment which provides unambiguous evidence of such a process.

The basis of our choice of this experiment was the prediction from ab initio calculations⁵ that, in the reaction of ClSiH with Me_3SiH , not only does the reaction proceed via a complex, but the complex has a degenerate rearrangement process as shown in scheme 1 below, indicated by the exchange of H and H*, via a low energy transition state in which both H-atoms are coordinated to each Si-atom.




Scheme 1

If correct this indicates that the introduction of a D-label would provide a means of lifting the degeneracy. Although the process cannot be observed directly (the complex has no known spectroscopic signature), its consequences should be detectable, since formation of the complex from its reactants is reversible, viz



This fact is a necessary consequence of the low A factor and negative activation energy found in our earlier kinetic study of this reaction.⁵ Thus an investigation of the reaction of $\text{ClSiH} + \text{Me}_3\text{SiD}$ should lead to H/D exchange and reveal the occurrence of the process




We report here studies of the kinetics of the gas phase reaction between ClSiH and Me_3SiD using laser flash photolysis. Details of the technique are given in our earlier review article.² ClSiH was made by excimer laser photolysis of 1-chlorosilacyclopent-3-ene⁶ at 193 nm. ClSiH was monitored by means of a CW Argon Ion laser tuned to its known absorption at 457.9 nm.⁶ Me_3SiD was prepared by standard means.⁷ Experiments were carried out in a spectroil quartz reaction vessel and signal decays (exponential) were collected and stored in a transient recorder. These yielded values for the pseudo first order rate constant, k_{obs} . A set of experiments was undertaken using different pressures of Me_3SiD . The results are shown in the Figure. The good linearity of the plot indicates that the reaction obeys second order kinetics. Also shown is the plot for $\text{ClSiH} + \text{Me}_3\text{SiH}$ obtained previously.⁵ The gradients of these plots yield values for the rate constants of $(1.395 \pm 0.052) \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Me_3SiD) and $(1.876 \pm 0.065) \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ (Me_3SiH), thus giving a kinetic isotope effect value of $k_{\text{D}}/k_{\text{H}} = 7.4 \pm 0.4$. Such a value is far in excess of the expectation of a value of less than one⁸⁻¹⁰ for a direct insertion process (ie leading to $\text{Me}_3\text{SiSiClHD}$ or

Me₃SiSiClH₂ in a single step). Only a mechanism change between the reactions of ClSiH with Me₃SiH and Me₃SiD can account for such a large value. Unfortunately we are not able to observe directly the formation of ClSiD with our detection system¹¹, but it is clear that a substantial additional pathway is needed to account for the disappearance of ClSiH in the reaction with Me₃SiD. One possibility can be eliminated, viz.



where Me₃SiSiClHD* represents the vibrationally excited disilane product. If this process were to occur it would indicate that collisional stabilisation of the disilane product was incomplete and the measured rate constant would be pressure dependent. This has been checked experimentally and found not to be the case (as indeed it was also not the case for ClSiH + Me₃SiH studied previously⁵). Only the rearrangement process involving the intermediate complex, predicted by calculation^{5,12,13}, would appear to offer an explanation. The full reaction mechanism is shown in scheme 2 below.



Scheme 2

In order to provide more concrete support for this mechanism we have undertaken theoretical calculations to investigate the isotope effect. This involved ab initio calculations (G3MP2B3 level) reported previously⁵ followed by RRKM calculations¹⁴ of the rate constants for rearrangement of the intermediate complex (k_3) relative to the rate constants for reaction to product (k_2) and to redissociation (k_{-1}), based on the structures and energies of transition states involved. The value of the isotope effect was obtained as the ratio of rate constants for ClSiH consumption with Me₃SiH and with Me₃SiD. Of course the major difference in the latter case is the inclusion of the scrambling process which ceases to be degenerate with the inclusion of the D-label. The vibrational assignments of the species involved in this calculation as well as their energies were taken from the ab initio calculations

for both the H- or D- labelled species. Table 1 summarises the energy surface details and Table 2 the RRKM results.

Since the intermediate complex and rearrangement transition states are all more stable than the reactants a chemical activation version of RRKM theory was used in these calculations¹⁴. Table 2 shows that the isotope effect is dependent on the magnitude of the *A* factor (and transition state structure) for redissociation of the complex, a quantity not available from theory. Values were selected based on an initial assumption that the association of ClSiH with Me₃SiH (or Me₃SiD) was collision controlled. This gives the maximum possible value¹⁵ for log (*A*₋₁/s⁻¹) of 16.08 corresponding to *k*_D/*k*_H = 9.27. The table shows that *k*_D/*k*_H decreases as log (*A*₋₁/s⁻¹) decreases. For the latter a value of 15.90, corresponding to *k*_D/*k*_H = 7.21 gives a good fit to experiment. This corresponds to ca 66% collision efficiency in the initial, complex-forming reaction step. Since silylenes, in general, are very efficient at insertion into Si-H bonds² (when, unlike in this case, the initial association step is rate controlling) this is a reasonable result. In order to check further, the isotope effect was calculated in the absence of the scrambling process (step (3)), viz when the only pathway open to the complex (apart from redissociation, step (-1)) was rearrangement to form disilane product (step (2)). The value obtained for *k*_D/*k*_H, in these circumstances, was 0.73, in line with expectation for a normal insertion process⁸⁻¹⁰, but clearly very different from experiment.

In summary, the large isotope effect found in this work, independent of pressure, can only be explained by a rearrangement process of an intermediate complex. This is the first experimental evidence for such a process and necessarily validates the existence of the complex in the Si-H insertion reaction. We have been able to reproduce the magnitude of this isotope effect by theoretical modelling. Further experiments are underway to extend these results to higher temperatures.

AUTHOR INFORMATION

Corresponding authors:

E-mails: r.walsh@reading.ac.uk, r.becerra@iqfr.csic.es

ACKNOWLEDGEMENT

This work was supported by the award of an RSC Journals grant to R.B. R.B. also thanks the Ministerio de Educacion y Ciencia(Spain) for support under project CTQ2006-10512.

REFERENCES and NOTES

1. See for example: Gaspar, P. P.; West, R. Silylenes. In *The Chemistry of Organosilicon Compounds*; Z. Rappoport and Y. Apeloig, Eds.; Wiley: Chichester, 1998, Vol. 2, Chapter 43, p 2463.
2. Becerra, R.; Walsh, R. *Phys. Chem. Chem. Phys.* **2007**, *9*, 2817 and references cited therein.
3. Becerra, R.; Frey, H. M.; Mason, B. P.; Walsh, R.; Gordon, M. S. *J. Chem. Soc., Faraday Trans.* **1995**, *91*, 2723
4. Baggott, J.E.; Blitz, M.A.; Frey, H.M.; Walsh, R. *J. Am. Chem. Soc.* **1990**, *112*, 8337.
5. Becerra, R.; Boganov, S. E.; Egorov, M. P.; Faustov, V.I.; Krylova, I.V.; Nefedov, O. M.; Promyslov, V.M.; Walsh, R. *J. Phys. Chem. A* **2009**, *113*, 5512.
6. Becerra, R.; Boganov, S. E.; Egorov, M. P.; Krylova, I.V.; Nefedov, O. M.; Walsh, R. *Chem. Phys. Lett.* **2005**, *413*, 194.
7. Skell, P. S.; Owen, P. W. *J. Am. Chem. Soc.* **1972**, *94*, 1578.
8. Values in the range 0.3 to 0.9 for primary isotope effects are found for a variety of mechanistic processes⁹, and an actual value of 0.9 obtained in the reaction of $\text{SiMe}_2 + \text{Et}_3\text{SiH(D)}$ ¹⁰.
9. Melander, L. Isotope effects on reaction rates. The Ronald Press, New York, 1960.
10. Conlin, R. T.; Netto-Ferreira, J. C.; Zeng, S.; Scaiano, J. C. *Organometallics* **1990**, *9*, 1332.
11. Unlike ClSiH , ClSiD has no vibronic band coincident with any line of our Ar Ion Laser monitor.
12. Swihart and Carr¹³ were the first to predict the possibility of such processes.
13. Swihart, M.T.; Carr, R.W. *J. Phys. Chem. A* **1997**, *101*, 7434.
14. Holbrook, K.A.; Pilling, M.J.; Robertson, S.H. *Unimolecular Reactions*, 2nd ed.; Wiley: Chichester, 1996.
15. Using microscopy reversibility and the calculated value of $\Delta S^\circ = -136.4 \text{ J K}^{-1} \text{ mol}^{-1}$ for forming the complex.




Figure. Second-order plot for reaction of ClSiH with Me₃SiD in SF₆ (5 Torr) at 297 K. Dashed line shows fit to data for ClSiH + Me₃SiH (ref. 5).

Table 1. Ab initio calculation of energies of species involved in the reaction of ClSiH + Me₃SiH(D) at the G3MP2B3 level

Species	$E_0(H)/\text{kJ mol}^{-1}$	$E_0(D)/\text{kJ mol}^{-1}$
ClSiH + Me ₃ SiH(D)	0	0
Complex (C1)	-36.90	-36.64
TS2 ^a	-9.30	-8.50
TS3 ^b	-21.90	-21.27

^aTransition state for step (2)

^bTransition state for step (3)

Table 2. RRKM calculations of the isotope effect based on the ab initio energy surface and different structures for redissociation of the intermediate complex

Quantity	Value		
$\log(A_{-1}/\text{s}^{-1})^a$	16.08 ^b	15.90 ^b	15.74 ^b
k_D/k_H	9.27	7.21	5.45

^aA factor for redissociation of intermediate complex, step(-1)

^bValues obtained after judicious modification of vibrational wavenumbers of transitional modes in TS(-1)

Table of contents graphic

