

*In Vitro Fermentation of NUTRIOSE®
FB06, a wheat dextrin soluble fibre, in a
continuous culture human colonic model
system*

Article

Published Version

Hobden, M. R., Martin-Morales, A., Guerin-Deremaux, L., Wils, D., Costabile, A., Walton, G. E., Rowland, I., Kennedy, O. B. and Gibson, G. R. (2013) In Vitro Fermentation of NUTRIOSE® FB06, a wheat dextrin soluble fibre, in a continuous culture human colonic model system. PLoS ONE, 8 (10). e77128. doi:
<https://doi.org/10.1371/journal.pone.0077128> Available at
<http://centaur.reading.ac.uk/34961/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1371%2Fjournal.pone.0077128>

To link to this article DOI: <http://dx.doi.org/10.1371/journal.pone.0077128>

Publisher: Public Library of Science

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

In Vitro Fermentation of NUTRIOSE® FB06, a Wheat Dextrin Soluble Fibre, in a Continuous Culture Human Colonic Model System

Mark R. Hobden^{1*}, Agustin Martin-Morales¹, Laetitia Guérin-Deremaux², Daniel Wils², Adele Costabile¹, Gemma E. Walton¹, Ian Rowland¹, Orla B. Kennedy¹, Glenn R. Gibson¹

1 Department of Food and Nutritional Sciences, The University of Reading, Reading, United Kingdom, **2** Roquette, Lestrem, France

Abstract

Wheat dextrin soluble fibre may have metabolic and health benefits, potentially acting via mechanisms governed by the selective modulation of the human gut microbiota. Our aim was to examine the impact of wheat dextrin on the composition and metabolic activity of the gut microbiota. We used a validated *in vitro* three-stage continuous culture human colonic model (gut model) system comprised of vessels simulating anatomical regions of the human colon. To mimic human ingestion, 7 g of wheat dextrin (NUTRIOSE® FB06) was administered to three gut models, twice daily at 10.00 and 15.00, for a total of 18 days. Samples were collected and analysed for microbial composition and organic acid concentrations by 16S rRNA-based fluorescence *in situ* hybridisation and gas chromatography approaches, respectively. Wheat dextrin mediated a significant increase in total bacteria in vessels simulating the transverse and distal colon, and a significant increase in key butyrate-producing bacteria *Clostridium* cluster XIVa and *Roseburia* genus in all vessels of the gut model. The production of principal short-chain fatty acids, acetate, propionate and butyrate, which have been purported to have protective, trophic and metabolic host benefits, were increased. Specifically, wheat dextrin fermentation had a significant butyrogenic effect in all vessels of the gut model and significantly increased production of acetate (vessels 2 and 3) and propionate (vessel 3), simulating the transverse and distal regions of the human colon, respectively. In conclusion, wheat dextrin NUTRIOSE® FB06 is selectively fermented *in vitro* by *Clostridium* cluster XIVa and *Roseburia* genus and beneficially alters the metabolic profile of the human gut microbiota.

Citation: Hobden MR, Martin-Morales A, Guérin-Deremaux L, Wils D, Costabile A, et al. (2013) *In Vitro* Fermentation of NUTRIOSE® FB06, a Wheat Dextrin Soluble Fibre, in a Continuous Culture Human Colonic Model System. PLoS ONE 8(10): e77128. doi:10.1371/journal.pone.0077128

Editor: Stefan Bereswill, Charité-University Medicine Berlin, Germany

Received: August 9, 2013; **Accepted:** September 5, 2013; **Published:** October 24, 2013

Copyright: © 2013 Hobden et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The study was funded by Roquette, France. The funders were involved in study design, decision to publish and preparation of the manuscript, however they had no role in running the experiments, data collection or analysis.

Competing interests: Roquette provided financial support for this study. Laetitia Guérin-Deremaux and Daniel Wils are employees of Roquette. NUTRIOSE® is a commercially available food product. This does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

* E-mail: m.r.hobden@reading.ac.uk

Introduction

The increasing worldwide prevalence of obese and overweight individuals is a major public health concern [1]. Obesity, and more specifically the accretion of excess adipose tissue, is associated with elevated chronic systemic low-grade inflammation and increased risk of metabolic diseases, such as type 2 diabetes and cardiovascular disease (CVD) [2]. The pathogenesis of these conditions is attributable to a complex interaction between genetic, metabolic, environmental and behavioural factors, however the specific contribution of each of these determinants is not fully understood [3]. The composition and metabolic activity of microbial inhabitants of the human gut has recently been acknowledged as an

environmental factor that may influence the development of obesity and associated metabolic diseases [4,5].

The human gut microbiota is a diverse ecosystem comprising of up to 100 trillion archaeal and bacterial cells. Any of between 1,000 - 1,150 bacterial species may reside in the gut, with most individuals harbouring at least 160 different species. Although over 90% of gut bacteria belong to Bacteroidetes and Firmicutes, the specific composition of the gut microbiota, at the phylum, genus and species level, is highly individual and affected by various factors, including adiposity [6]. Obesity and diet-induced weight gain have been associated with a modified microbial composition, with some studies [7-10], but not all [11-13], providing evidence for an altered Firmicutes-Bacteroidetes ratio. Reduced numbers of *Eubacterium rectale-Clostridium coccoides* cluster,

Bifidobacterium spp., *Lactobacillus* spp. and *Roseburia* spp., have been observed in mice subjected to an obesogenic diet [14-17]. Furthermore, a recent rodent model study by Liou et al., has provided the first empirical evidence that changes in the gut microbiota may contribute towards reduced host weight and adiposity [18].

The gut microbiota has a major influence on host metabolism, with microbe-host interactions connecting with organs, including the gut, liver, adipose tissue, muscle and brain. Accordingly, dietary modulation of the composition and activity of the gut microbiota, with food ingredients such as prebiotics, has been highlighted as a potential target for obesity and metabolic diseases [5]. Prebiotics are defined as 'selectively fermented dietary ingredients that result in specific changes in the composition and/or activity of the gastrointestinal microbiota, thus conferring benefit(s) upon host health' [19]. Woods and Gorbach included in this definition an increase in beneficial bacteria and/or a decrease in harmful types, a reduction in intestinal pH, production of SCFA and changes in bacterial enzyme concentrations [20]. The fermentation of a food ingredient by the gut microbiota is dependent on its physicochemical structure [4]. Thus far, most attention has focused on the prebiotic potential of soluble fibres, in particular non-digestible oligosaccharides such as inulin-type fructooligosaccharides (FOS) and transgalactooligosaccharides (TOS), with the daily intake of 2.75 - 20 g shown to positively alter gut microbial composition after a short feeding period [21]. Other soluble fibres, including resistant dextrins (wheat or starch), glucans, gums and pectins are also increasingly recognised as having prebiotic potential. The intake of wheat dextrin (WD) and FOS has been shown to have satiogenic and weight management benefits, possibly attributable to elevated synthesis of anorexigenic gut hormones (peptide YY (PYY) and glucagon-like peptide 1 (GLP-1)) and decreased synthesis of the orexigenic gut hormone, ghrelin [22-24]. Furthermore, TOS has recently been found to beneficially impact on metabolic markers of immune function, systemic inflammation and blood glucose regulation [25]. The mechanisms responsible for these effects remain to be elucidated, however up regulation of short-chain fatty acids (SCFA), acetate, propionate and butyrate, which are key metabolic end-products of bacterial fermentation, may play pivotal roles [26].

NUTRIOSE® (NUTRIOSE® FB06, Roquette, France) is a non-viscous WD with a total fibre content of ~ 85% and a mono- and disaccharide content of ≤ 0.5% [27]. NUTRIOSE® has a structure of linear and branched glucosidic linkages that make it resistant to hydrolysis in the small intestine and consequently available for bacterial fermentation in the human large gut [28]. NUTRIOSE® induces a low glycaemic response and is well tolerated by the human digestive system, even at high doses [29,30]. Emerging evidence indicates that NUTRIOSE® has prebiotic potential, however most of the studies have been done in animal models, which have a different microbiota from that of humans, or have investigated a limited number of bacterial groups in humans. Nevertheless these studies have demonstrated that the intake of NUTRIOSE® may modulate gut microbial ecology, with

evidence for increased faecal counts of *Bacteroides* and *Lactobacillus* spp., reduced *Clostridium perfringens*, increased total SCFA production, elevated α - and β - glucosidase activity and decreased faecal pH [29-32]. Furthermore, NUTRIOSE® has been shown to have promising effects on energy metabolism, appetite regulation and weight management [33-36]. In a recent human intervention study by Guerin-Deremaux et al., the daily intake of either 14 g, 18 g, or 24 g of NUTRIOSE®, over a period of nine weeks, was found to increase perceived feelings of satiety and subsequently lead to a reduction in energy intake, bodyweight and percentage body fat in a group of overweight males [34]. Preliminary evidence also suggests that NUTRIOSE® may improve indices of lipid and glucose homeostasis, however additional work is warranted to verify these findings [37].

The aim of the present study was to examine more fully the impact of WD fermentation on gut microbial ecology and metabolic end products of microbial fermentation in distinct anatomical regions of the human colon using an *in vitro* three-stage continuous culture human colonic model (gut model) system. This system has been used extensively at our institution and provides a controlled and steady-state environment in which to study the composition and metabolic activities of the gut microbiota in relation to external perturbations, such as with the administration of food ingredients [38]. The intake of 14 g/day NUTRIOSE® was the lowest dose to significantly increase satiety, reduce energy intake and improve body composition in previous investigations by Guerin-Deremaux and colleagues [34,36]. Accordingly, this dosage was chosen for the present study and was administered in two equal doses each day to mimic human ingestion. The impact of WD fermentation on the proliferation of a selection of the main bacterial groups of the gut microbiota, including groups with purported benefits or detriments to health, was assessed using 16S rRNA-based fluorescence in situ hybridisation (FISH) and organic acid production analysed by gas chromatography (GC).

Materials and Methods

Three-stage continuous culture colonic model (gut model) system

Three gut models, each comprising of a cascade of three glass fermenters connected in series that simulate the different physical and nutritional characteristics of the proximal (V1), transverse (V2) and distal colon (V3), were implemented under conditions previously detailed by Macfarlane et al. [38]. Gut models were inoculated with faecal samples from healthy human donors, all of whom had no previous history of gastrointestinal disorders and had not received antibiotics or pre/probiotics for at least three months prior to the study. All faecal donors had the experimental procedure explained to them and were given the opportunity to ask questions. All donors then provided verbal informed consent for the use of their faeces in the study. The University of Reading research ethics committee exempted this study from review because no donors were involved in any intervention and waived the need for written consent due to the fact the samples received were

Table 1. Probes used for bacterial enumeration by FISH.

Probe name	Sequence (5' to 3')	Target species	Hybridization- Washing Temperature (°C)	Reference
Ato291	GGTCGGTCTCTCAACCC	<i>Atopobium</i> , <i>Colinsella</i> , <i>Olsenella</i> and <i>Eggerthella</i> spp.; <i>Cryptobacterium curtum</i> ; <i>Mycoplasma equigenitalium</i> and <i>Mycoplasma elephantis</i>	50 - 50	[54]
Bac303	CCAATGTGGGGACCTT	Most <i>Bacteroides sensu stricto</i> and <i>Prevotella</i> spp.; all <i>Parabacteroides</i> ; <i>Barnesiella viscericola</i> and <i>Odoribacter splanchnicus</i>	46 - 48	[55]
Bif164	CATCCGGCATTACCACCC	Most <i>Bifidobacterium</i> spp. and <i>Parascardovia denticolens</i>	50 - 50	[56]
Chis150	TTATGCGGTATTAATCTYCCTTT	Most members of <i>Clostridium</i> cluster I; all members of <i>Clostridium</i> cluster II; <i>Clostridium tyrobutyricum</i> ; <i>Adhaeribacter aquaticus</i> and <i>Flexibacter canadensis</i> (family <i>Flexibacteriaceae</i>); [<i>Eubacterium</i>] <i>combesii</i> (family <i>Propionibacteriaceae</i>)	50 - 50	[57]
Erec482	GCTTCTTAGTCARGTACCG	Most members of <i>Clostridium</i> cluster XIVa; <i>Syntrophococcus sucromutans</i> , [<i>Bacteroides</i>] <i>galacturonicus</i> and [<i>Bacteroides</i>] <i>xylanolyticus</i> , <i>Lachnospira pectinschiza</i> and <i>Clostridium saccharolyticum</i>	50 - 50	[57]
Lab158	GGTATTAGCAYCTGTTTCCA	Most <i>Lactobacillus</i> , <i>Leuconostoc</i> and <i>Weissella</i> spp.; <i>Lactococcus lactis</i> ; all <i>Vagococcus</i> , <i>Enterococcus</i> , <i>Melisococcus</i> , <i>Tetragenococcus</i> , <i>Catelicoccus</i> , <i>Pediococcus</i> and <i>Paralactobacillus</i> spp.	50 - 50	[58]
Rrec584	TCAGACTTGCCGYACCGC	<i>Roseburia - Eubacterium rectale</i> (a component of cluster XIVa)	50 - 50	[59]
Eco1531	CACCGTAGTGCCCTCGTCATC	<i>Escherichia coli</i>	37 - 37	[59]
EUB338	GCTGCCTCCCGTAGGAGT	Total bacteria	46 - 48	[60]
EUB338II	GCAGCCACCCGTAGGTGT	Total bacteria	46 - 48	[60]
EUB338III	GCTGCCACCCGTAGGTGT	Total bacteria	46 - 48	[60]

doi: 10.1371/journal.pone.0077128.t001

not collected by means of intervention. Equilibrium of the system, steady state 1 (SS1), was reached after eight full turnovers at 15, 16 and 17 days. Thereafter, the test product, NUTRIOSE®, was administered into V1 in 7 g doses at 10.00 and 15.00 each day until the second steady state (SS2) had been reached at 33, 34, and 35 days. Samples (1 mL) were collected from all vessels of the colonic system and centrifuged at 13,000 x g for 10 min to remove all particulate matter. A shortened GC method was used to determine the stabilisation of SCFA concentrations over three consecutive days and confirm steady state (SS1 and SS2) [39]. Further samples were stored at -18 °C for future analysis.

In vitro enumeration of bacterial populations by FISH

Enumeration of bacterial populations was performed by FISH using synthetic oligonucleotide probes designed to target specific diagnostic regions of 16S rRNA and labelled with the fluorescent Cy3 dye (Sigma Aldrich Ltd., Poole, Dorset, UK), as previously described [40]. As detailed in Table 1, probes were used for the determination of total bacteria, bifidobacteria, lactobacilli/enterococci, *Bacteroides*, *Clostridium perfringens*/ *histolyticum* subgroup, *Clostridium* cluster XIVa, *Clostridium* cluster I and II, *Roseburia* genus and *Atopobium - Coriobacterium*.

Organic acid determination by GC

Samples were extracted and derivatised as previously described by Richardson et al. [41]. Briefly, aliquots of 1 mL of supernatant were transferred into glass tubes, followed by the addition of 50 µL of internal standard (100 mM; 2-ethylbutyric

acid), 500 µL of concentrated hydrochloric acid (HCl) and 2 mL of diethyl ether (Sigma Aldrich Ltd., Poole, Dorset, UK). Samples were then vortexed for 1 min before centrifugation at 3,000 x g for 10 min. The top ether layer was transferred from each tube into clean glass tubes. A second extraction step was then completed using a further 1mL of diethyl ether. The diethyl layer was again collected and pooled with the layer from the first extraction. Aliquots of 400µL of this pooled extract were transferred into glass vials, alongside 50 µL of *N*-methyl-*N*-*t*-butyldimethylsilyltrifluoroacetamide (Cheshire Sciences, Chester, UK). The samples were then incubated at 80°C in a water bath for 20min and left at room temperature for 48 h to allow for the complete derivatisation of lactic acid.

The derivatized samples were run on a 5890 SERIES II Gas Chromatograph (Hewlett Packard, UK) with flame ionization detector, using an Rtx-1 10m×0.18mm column coated with a 0.20µm coating (Crossbond 100% dimethyl polysiloxane; Restek, Buckinghamshire, UK). Injector and detector temperature were set at 275°C and the column temperature was programmed from 63°C for 3 min to 190°C at 10°C/min⁻¹ and held at 190 °C for 3 min. Helium was used as the carrier gas (flow rate 1.2 mL/min; head pressure 90 MPa). External standards contained (mM): sodium formate, 10; acetic acid, 30; propionic acid, 20; iso-butyric acid, 5; n-butyric acid, 20; iso-valeric acid, 5; n-valeric acid, 5; sodium lactate, 10; sodium succinate, 20. Chemstation B.03.01 (Agilent Technologies, Cheshire, UK) was used for calibration and calculation of the internal response factor for quantification of peak areas within samples.




Figure 1. Bacterial populations. Bacterial populations in the three different vessels (V1, V2 and V3) of the gut models before (SS1) and after (SS2) WD treatment. Data presented as means of the three gut models \pm standard deviations. SS1 and SS2 are calculated as mean values over three consecutive days. ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, significantly different from SS1. doi: 10.1371/journal.pone.0077128.g001

Statistical analysis

All statistical tests were performed on GraphPad Prism 6.0 (GraphPad Software, La Jolla, CA, USA). One-way AVOVA tests were used to compare SS1 and SS2 data for bacterial counts and organic acid concentrations. Where significant differences were identified, *post hoc* analysis was performed using Tukey multiple comparison tests. Statistical significance was accepted at $P < 0.05$ for all analyses.

Results

The impact of WD on the human gut microbiota

Average bacterial counts, as enumerated by FISH, are displayed in Figure 1 and expressed as log CFU/ml \pm standard deviations. Following administration of WD, total bacterial populations significantly increased by 0.37 log₁₀ in V2 simulating the transverse colon ($P < 0.0001$) and 0.30 log₁₀ in V3 simulating the distal colon ($P < 0.001$). No significant differences for total bacteria were found in V1, simulating the proximal colon. Concentrations of *Clostridium coccooides* – *Eubacterium rectale* significantly increased by 0.71 log₁₀ in V1 ($P < 0.0001$), 0.67 log₁₀ in V2 ($P < 0.0001$) and 0.37 log₁₀ in V3 ($P < 0.01$). Furthermore, concentrations of *Roseburia* - *E. rectale* significantly increased by 1.07 log₁₀ in V1 ($P < 0.0001$), 1.14 log₁₀ in V2 ($P < 0.0001$) and 1.33 log₁₀ in V3 ($P < 0.0001$). No significant differences were found for the other bacterial groups analysed, including *Bifidobacterium* spp., *Lactobacillus* - *Enterococcus*., *Bacteroides* - *Prevotella*, *Atopobium* - *Coriobacterium* and *Escherichia coli*. There were trends for increases in *Bifidobacterium* spp. and *Escherichia coli* in all vessels, however these did not reach significance. Counts for

the *Clostridium histolyticum* group were below the detection level (data not shown).

Organic acid production

Relative concentrations of organic acids, as determined by GC, are reported in Table 2 and expressed as mM \pm standard deviations. Fermentation of WD mediated a significant 19.45 mM increase in butyrate in V1 ($P < 0.0001$), 30.38 mM increase in V2 ($P < 0.001$) and 27.01 mM increase in V3 ($P < 0.01$) of the gut models. In V1, simulating the proximal colon, WD administration resulted in a significant 9.82 mM reduction in acetate ($P < 0.01$) and 14.71 mM reduction in propionate ($P < 0.05$) concentrations. In V2, simulating the transverse colon, acetate concentrations significantly increased by 16.11 mM ($P < 0.01$), whereas propionate concentrations did not change. In V3, simulating the distal colon, acetate concentrations significantly increased by 29.16 mM ($P < 0.001$) and propionate concentrations significantly increased by 14.64 mM ($P < 0.01$). Concentrations of formate, iso-valeric, valeric and lactate were below the detection limit.

Discussion

The aim of the present study was to extend knowledge of the impact of WD on the microbial ecology of the human gut. A validated gut model system was used to establish the effects of *in vitro* fermentation of WD on the proliferation of a range of known bacterial groups. These included the purported health-promoting *Bifidobacterium* and *Lactobacillus* genera, but also other bacteria increasingly recognised for having important roles in the human colon, including *Bacteroides* and *Roseburia*

Table 2. SCFA concentrations.

	Acetate		Propionate		Butyrate	
	SS1	SS2	SS1	SS2	SS1	SS2
Vessel 1	31.92 ± 6.12	22.10 ± 4.28 **	33.82 ± 7.65	19.12 ± 8.01 *	30.75 ± 3.10	50.20 ± 2.47 ****
Vessel 2	45.77 ± 3.56	61.88 ± 7.65 **	41.02 ± 5.53	40.87 ± 6.70	34.78 ± 6.11	65.16 ± 8.72 ***
Vessel 3	50.82 ± 8.37	79.97 ± 10.42 ***	43.15 ± 6.81	57.80 ± 8.24 **	36.98 ± 6.01	63.99 ± 9.43 **

Mean SCFA concentrations reported as mM ± standard deviations for the three vessels of the gut models before (SS1) and after (SS2) WD treatment. SS1 and SS2 are calculated as mean values over three consecutive days. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, significantly different from SS1.

doi: 10.1371/journal.pone.0077128.t002

genus. In addition, metabolic end products of microbial fermentation, including SCFAs, acetate, propionate and butyrate, were analysed to elucidate potential mediators by which this novel food ingredient may exert metabolic and health benefits to the human host through modulation of the gut microbiota.

The provision of WD, at a dose of 14 g/day, resulted in marked increases in total bacterial populations in vessels simulating the transverse and distal colon. This increased proliferation of total bacteria is concurrent with findings from previous studies in rodents and humans [29-32]. To our knowledge, we have demonstrated for the first time that WD significantly increases the proliferation of Gram-positive *Clostridium* Cluster XIVa bacteria and *Roseburia* genus, as categorised by increased counts for *Clostridium coccooides* – *Eubacterium rectale* and *Roseburia* - *E. rectale* groups, respectively. These bacterial groups are major components of the human gut microbiota and have important roles in the fermentation and putrefaction of food-derived substances, resulting in the production of SCFAs and intestinal gases [42]. Furthermore, *Clostridium* Cluster XIVa and *Roseburia* genus are key butyrate producers, with ~ 80% of the butyrate-producing isolates originating in human faeces belonging to these groups [43]. Experimental work in rodents suggests that obesity is inversely correlated to numbers of *Clostridium* Cluster XIVa and *Roseburia* genus. Interestingly, these studies found that the provision of dietary fibres, chitin–glucan and wheat arabinoxylan, to mice subjected to an obesogenic diet ameliorated the reduction in these two bacterial groups [14,17]. Owing to the findings of the present study, it is feasible that the intake of WD may also have restorative effects on gut microbial populations of *Clostridium* Cluster XIVa and *Roseburia* genus in obese and overweight individuals.

Whilst there was a trend for an increase in *Bifidobacterium* spp. across the gut model, neither populations of this genus nor *Lactobacillus* - *Enterococcus*., significantly increased in response to the provision of WD. This, in combination with previous findings, suggests that the physicochemical structure of WD may not confer selectivity to bifidobacteria, which has been shown to prefer oligosaccharide structures, including

TOS and various forms of galacto-oligosaccharides [44]. Pasman et al. observed increases in lactobacilli in response to WD, however this was following the intake of a higher dose of 45 g/day [31], with a recent study by Lefranc-Millot et al. [30] failing to observe such an effect with lower dosages. Lefranc-Millot et al. did however find that the intake of 10 g/day WD increased numbers of *Bacteroides*, a genus of bacteria that may benefit the human host by preventing potential pathogens from colonising the gut. Furthermore, the administration of 8 g/day and 15 g/day reduced numbers of *Clostridium perfringens*, an opportunistic human pathogen [30]. In the current study *Clostridium perfringens* levels were below the detection threshold in all three human donors and subsequently we were unable to determine the impact of WD on this bacterial group. Whilst there was a trend for an increase in *Bacteroides* in the transverse and distal regions of the gut model, this did not reach significance.

The observed increases in butyrate production across the three vessels of the gut model, together with increases in *Clostridium* cluster XIVa and *Roseburia* genus provide evidence for a butyrogenic effect of WD on gut microbial ecology. Moreover, the increasing magnitude of butyrate production through the gut model from V1-V3 strengthens support for a butyrogenic effect of WD in all regions of the human colon and eliminates the possibility that increased concentrations in the latter regions of the gut model were solely due to butyrate transiting through from a previous vessel in the system. Our findings support previous evidence from a rodent model that found that 14 days of WD supplementation increased faecal concentrations of butyrate, acetate and propionate [32]. Increased production of SCFA butyrate in the human large gut may have numerous health benefits. Firstly, butyrate is the preferred energy-providing substrate for colonocytes [45]. Secondly, butyrate has been shown to have a beneficial trophic effect on the gut epithelium [46]. Lastly, butyrate has been demonstrated to modulate colonic cell proliferation / apoptosis and may have protective properties against colon cancer and ulcerative colitis [47]. As the incidence of colon cancer and other colonic conditions, such as ulcerative colitis, is highest in the distal regions of the colon [48], the elevated production of butyrate and SCFAs, propionate and acetate, in V3 of the gut model is of considerable interest as regional differences in SCFA concentrations has implications for disease risk [49]. Furthermore, increased production of SCFAs indicates a preferential increase in carbohydrate fermentation rather than protein fermentation, which mainly occurs in the distal regions of the human colon due to reduced nutrient availability [50]. Rather than producing beneficial SCFAs and other organic acids, the breakdown of protein by proteolytic bacteria results in the formation of ammonia, phenols, indoles, thiols, amines and sulphides, which are potentially detrimental to health [51].

There is mounting evidence that prebiotics impact on appetite regulation and gut hormone release however the underlying mechanisms are not yet clear. Evidence from animal studies has recently highlighted the potential mechanistic importance of SCFA production by the gut microbiota. Findings from rodent models have demonstrated

that butyrate and propionate, but not acetate, stimulate the secretion of anorexigenic hormones, GLP-1, PYY and gastric inhibitory polypeptide (GIP). These effects may be expressed through the activation of G-protein coupled receptors free-fatty acid receptor 2 and free-fatty acid receptor 3 by butyrate and propionate, however mechanisms independent of these receptors may also be involved [52]. Gut hormones are fundamental in the regulation of appetite but also in the control of glucose metabolism, with GIP involved in insulin dependent glucose uptake into muscle tissue and GLP-1 involved in insulin secretion from the pancreas [53]. In the present study, the WD mediated increases in butyrate and propionate production has identified possible metabolic mediators for any effects of WD fermentation on appetite regulation and energy metabolism.

In conclusion, WD had a beneficial impact on gut microbial ecology, not only in V1 where administered, but persisting through all vessels of gut model system. Whilst not exerting a classical prebiotic effect, such as increased *Bifidobacterium*

spp. or *Lactobacillus* spp., WD was selectively fermented by *Clostridium* Cluster XIVa and *Roseburia* genus, which are increasingly recognised as important health promoting components of the gut microbiota. Furthermore, observed increases in SCFA production, in particular butyrate and propionate, provide possible metabolic mediators for any effects of WD on host appetite regulation and energy metabolism. Future human intervention studies are required to establish the concomitant effects of WD on gut microbial ecology, appetite regulation and systemic markers of metabolism in order to elucidate underlying mechanisms of action.

Author Contributions

Conceived and designed the experiments: LGD OBK GRG IR. Performed the experiments: AMM MRH GEW. Analyzed the data: MRH AMM AC LGD OBK GRG. Wrote the manuscript: MRH AMM AC GEW LGD DW IR OBK GRG.

References

- Wang YC, McPherson K, Marsh T, Gortmaker SL, Brown M (2011) Health and economic burden of the projected obesity trends in the USA and the UK. *Lancet* 378: 815-825. doi:10.1016/S0140-6736(11)60814-3. PubMed: 21872750.
- Haffner SM (2006) Relationship of Metabolic Risk Factors and Development of Cardiovascular Disease and Diabetes. *Obesity* (Silver Spring) 14: 121S-127S. doi:10.1038/oby.2006.291. PubMed: 16931493.
- Norris JM, Rich SS (2012) Genetics of Glucose Homeostasis Implications for Insulin Resistance and Metabolic Syndrome. *Arterioscler Thromb Vasc Biol* 32: 2091-2096. doi:10.1161/ATVBAHA.112.255463. PubMed: 22895670.
- Tremaroli V, Bäckhed F (2012) Functional interactions between the gut microbiota and host metabolism. *Nature* 489: 242-249. doi:10.1038/nature11552. PubMed: 22972297.
- Nicholson JK, Holmes E, Kinross J, Burcelin R, Gibson G et al. (2012) Host-gut microbiota metabolic interactions. *Science* 336: 1262-1267. doi:10.1126/science.1223813. PubMed: 22674330.
- Qin J, Li R, Raes J, Arumugam M, Burgdorf KS et al. (2010) A human gut microbial gene catalogue established by metagenomic sequencing. *Nature* 464: 59-65. doi:10.1038/nature08821. PubMed: 20203603.
- Ley RE, Bäckhed F, Turnbaugh P, Lozupone CA, Knight RD et al. (2005) Obesity alters gut microbial ecology. *Proc Natl Acad Sci U S A* 102: 11070-11075. doi:10.1073/pnas.0504978102. PubMed: 16033867.
- Ley RE, Turnbaugh PJ, Klein S, Gordon JI (2006) Microbial ecology: human gut microbes associated with obesity. *Nature* 444: 1022-1023. doi:10.1038/4441022a. PubMed: 17183309.
- Turnbaugh PJ, Bäckhed F, Fulton L, Gordon JI (2008) Diet-induced obesity is linked to marked but reversible alterations in the mouse distal gut microbiome. *Cell Host Microbe* 3: 213-223. doi:10.1016/j.chom.2008.02.015. PubMed: 18407065.
- Furet JP, Kong LC, Tap J, Poitou C, Basdevant A et al. (2010) Differential Adaptation of Human Gut Microbiota to Bariatric Surgery-Induced Weight Loss Links With Metabolic and Low-Grade Inflammation Markers. *Diabetes* 59: 3049-3057. doi:10.2337/db10-0253. PubMed: 20876719.
- Duncan SH, Lohley GE, Holtrop G, Ince J, Johnstone AM et al. (2008) Human colonic microbiota associated with diet, obesity and weight loss. *Int J Obes* 32: 1720-1724. doi:10.1038/ijo.2008.155. PubMed: 18779823.
- Schwartz A, Taras D, Schäfer K, Beijer S, Bos NA et al. (2009) Microbiota and SCFA in lean and overweight healthy subjects. *Obesity* 18: 190-195. PubMed: 19498350.
- Zhang H, DiBaise JK, Zuccolo A, Kudrna D, Braidotti M et al. (2009) Human gut microbiota in obesity and after gastric bypass. *Proc Natl Acad Sci USA* 106: 2365-2370. doi:10.1073/pnas.0812600106. PubMed: 19164560.
- Neyrinck AM, Possemiers S, Verstraete W, De Backer F, Cani PD et al. (2012) Dietary modulation of clostridial cluster XIVa gut bacteria (< i> Roseburia</i> spp.) by chitin-glucan fiber improves host metabolic alterations induced by high-fat diet in mice. *The Journal of nutritional biochemistry* 23: 51-59.
- Cani PD, Neyrinck AM, Fava F, Knauf C, Burcelin RG et al. (2007) Selective increases of bifidobacteria in gut microflora improve high-fat-diet-induced diabetes in mice through a mechanism associated with endotoxaemia. *Diabetologia* 50: 2374-2383. doi:10.1007/s00125-007-0791-0. PubMed: 17823788.
- Cani PD, Bibiloni R, Knauf C, Waget A, Neyrinck AM et al. (2008) Changes in gut microbiota control metabolic endotoxemia-induced inflammation in high-fat diet-induced obesity and diabetes in mice. *Diabetes* 57: 1470-1481. doi:10.2337/db07-1403. PubMed: 18305141.
- Neyrinck AM, Possemiers S, Druart C, Van de Wiele T, De Backer F et al. (2011) Prebiotic Effects of Wheat Arabinoxylan Related to the Increase in Bifidobacteria, Roseburia and Bacteroides/Prevotella in Diet-Induced Obese Mice. *PLOS ONE* 6: e20944. doi:10.1371/journal.pone.0020944. PubMed: 21695273.
- Liou AP, Paziuk M, Luevano J-M, Machineni S, Turnbaugh PJ et al. (2013) Conserved Shifts in the Gut Microbiota Due to Gastric Bypass Reduce Host Weight and Adiposity. *Sci Transl Med* 5: 178ra141. PubMed: 23536013.
- Gibson GR, Scott KP, Rastall RA, Tuohy KM, Hotchkiss A et al. (2010) Dietary prebiotics: current status and new definition. *Foods Science Technol Bulletin Funct Foods* 7: 1-19. doi:10.1616/1476-2137.15880.
- Woods MN, Gorbach SL (2001) Influences of fibre on the ecology of the intestinal flora. *CRC Handbook of Dietary Fibre in Human Nutrition*: 257-270.
- Walton G, Swann J, Gibson G (2013) Prebiotics. In: E RosenbergE DeLongS LoryE StackebrandtF Thompson. *The Prokaryotes*. Berlin Heidelberg: Springer Verlag, pp. 25-43.
- Nazare JA, Sauvinet V, Normand S, Guérin-Deremaux L, Gabert L et al. (2011) Impact of a resistant dextrin with a prolonged oxidation pattern on day-long ghrelin profile. *J Am Coll Nutr* 30: 63-72. doi:10.1080/07315724.2011.10719945. PubMed: 21697540.
- Cani PD, Lecourt E, Dewulf EM, Sohet FM, Pachikian BD et al. (2009) Gut microbiota fermentation of prebiotics increases satiety and incretin gut peptide production with consequences for appetite sensation and glucose response after a meal. *Am J Clin Nutr* 90: 1236-1243. doi:10.3945/ajcn.2009.28095. PubMed: 19776140.
- Parnell JA, Reimer RA (2009) Weight loss during oligofructose supplementation is associated with decreased ghrelin and increased peptide YY in overweight and obese adults. *Am J Clin Nutr* 89: 1751-1759. doi:10.3945/ajcn.2009.27465. PubMed: 19386741.
- Vulevic J, Juric A, Tzortzis G, Gibson GR (2013) A mixture of transgalactooligosaccharides reduces markers of metabolic syndrome and modulates the fecal microbiota and immune function of overweight

- adults. *J Nutr* 143: 324-331. doi:10.3945/jn.112.166132. PubMed: 23303873.
26. Sleeth ML, Thompson EL, Ford HE, Zac-Varghese SE, Frost G (2010) Free fatty acid receptor 2 and nutrient sensing: a proposed role for fibre, fermentable carbohydrates and short-chain fatty acids in appetite regulation. *Nutr Res Rev* 23: 135-145. doi:10.1017/S0954422410000089. PubMed: 20482937.
 27. Fu B, Wang J, Roturier JM, Tang Z, Li H et al. (2008) Determination of Total Dietary Fiber in Selected Foods Containing Resistant Maltodextrin by a Simplified Enzymatic-Gravimetric Method and Liquid Chromatography: Interlaboratory Study in China. *J AOAC Int* 91: 614-621. PubMed: 18567308.
 28. Lefranc-Millot C, Wils D, Roturier JM, Le Bihan C, Saniez-Degrave MH et al. (2009) NUTRIOSE® Soluble Fiber. *Fibers Ingredients Foods Applications Health Benefits*: 19-40.
 29. van den Heuvel EG, Wils D, Pasman WJ, Bakker M, Saniez MH et al. (2004) Short-term digestive tolerance of different doses of NUTRIOSE FB, a food dextrin, in adult men. *Eur J Clin Nutr* 58: 1046-1055. doi: 10.1038/sj.ejcn.1601930. PubMed: 15220947.
 30. Lefranc-Millot C, Guérin-Deremaux L, Wils D, Neut C, Miller LE et al. (2012) Impact of a resistant dextrin on intestinal ecology: how altering the digestive ecosystem with NUTRIOSE(R), a soluble fibre with prebiotic properties, may be beneficial for health. *J Int Med Res* 40: 211-224. doi:10.1177/147323001204000122. PubMed: 22429361.
 31. Pasman W, Wils D, Saniez MH, Kardinaal A (2006) Long-term gastrointestinal tolerance of NUTRIOSE FB in healthy men. *Eur J Clin Nutr* 60: 1024-1034. doi:10.1038/sj.ejcn.1602418. PubMed: 16482066.
 32. Guerin-Deremaux L, Ringard F, Desailly F, Wils D (2010) Effects of a soluble dietary fibre NUTRIOSE® on colonic fermentation and excretion rates in rats. *Nutrition research and practice* 4: 470-476
 33. Guerin-Deremaux L, Li S, Pochat M, Wils D, Mubasher M et al. (2011) Effects of NUTRIOSE(R) dietary fiber supplementation on body weight, body composition, energy intake, and hunger in overweight men. *Int J Food Sci Nutr* 62: 628-635. doi:10.3109/09637486.2011.569492. PubMed: 21591985.
 34. Guérin-Deremaux L, Pochat M, Reifer C, Wils D, Cho S et al. (2011) The soluble fiber NUTRIOSE induces a dose-dependent beneficial impact on satiety over time in humans. *Nutr Res* 31: 665-672. doi: 10.1016/j.nutres.2011.09.004. PubMed: 22024490.
 35. Li S, Guerin-Deremaux L, Pochat M, Wils D, Reifer C et al. (2010) NUTRIOSE dietary fiber supplementation improves insulin resistance and determinants of metabolic syndrome in overweight men: a double-blind, randomized, placebo-controlled study. *Appl Physiol Nutr Metab* 35: 773-782. doi:10.1139/H10-074. PubMed: 21164548.
 36. Guérin-Deremaux L, Pochat M, Reifer C, Wils D, Cho S et al. (2013) Dose-response impact of a soluble fiber, NUTRIOSE®, on energy intake, body weight and body fat in humans. *Global Epidemic. Obesity (Silver Spring)* 1.
 37. Li SLS, Guerin-Deremaux LG-DL, Pochat MPM, Wils DWD, Reifer CRC et al. (2010) NUTRIOSE dietary fiber supplementation improves insulin resistance and determinants of metabolic syndrome in overweight men: a double-blind, randomized, placebo-controlled study. *Appl Physiol Nutr, Metab* 35: 773-782. doi:10.1139/H10-074. PubMed: 21164548.
 38. Macfarlane GT, Macfarlane S, Gibson GR (1998) Validation of a Three-Stage Compound Continuous Culture System for Investigating the Effect of Retention Time on the Ecology and Metabolism of Bacteria in the Human Colon. *Microb Ecol* 35: 180-187. doi:10.1007/s002489900072. PubMed: 9541554.
 39. Pereira DI, Gibson GR (2002) Cholesterol assimilation by lactic acid bacteria and bifidobacteria isolated from the human gut. *Appl Environ Microbiol* 68: 4689-4693. doi:10.1128/AEM.68.9.4689-4693.2002. PubMed: 12200334.
 40. Martín-Peláez S, Gibson GR, Martín-Orúe SM, Klinder A, Rastall RA et al. (2008) In vitro fermentation of carbohydrates by porcine faecal inocula and their influence on *Salmonella Typhimurium* growth in batch culture systems. *FEMS Microbiol Ecol* 66: 608-619. doi:10.1111/j.1574-6941.2008.00610.x. PubMed: 19049655.
 41. Richardson AJ, Calder AG, Stewart CS, Smith A (1989) Simultaneous determination of volatile and non-volatile acidic fermentation products of anaerobes by capillary gas chromatography. *Lett Appl Microbiol* 9: 5-8. doi:10.1111/j.1472-765X.1989.tb00278.x.
 42. Konstantinov S, Smidt H, Akkermans A (2005) *Handbook on Clostridia: Ecology and activity of clostridia in the intestine of mammals*. Germany: University of Ulm. CRC Press. pp. 986-997.
 43. Barcenilla A, Pryde SE, Martin JC, Duncan SH, Stewart CS et al. (2000) Phylogenetic relationships of butyrate-producing bacteria from the human gut. *Appl Environ Microbiol* 66: 1654-1661. doi:10.1128/AEM.66.4.1654-1661.2000. PubMed: 10742256.
 44. Macfarlane GT, Steed H, Macfarlane S (2008) Bacterial metabolism and health-related effects of galacto-oligosaccharides and other prebiotics. *J Appl Microbiol* 104: 305-344. PubMed: 18215222.
 45. Sakata T (1987) Stimulatory effect of short-chain fatty acids on epithelial cell proliferation in the rat intestine: a possible explanation for trophic effects of fermentable fibre, gut microbes and luminal trophic factors. *Br J Nutr* 58: 95-103. doi:10.1079/BJN19870073. PubMed: 3620440.
 46. Sakata T (1984) Short chain fatty acids as the luminal trophic factor. *Can J Anim Sci* 64: 189-190. doi:10.4141/cjas84-218.
 47. Rowland IR (2009) The role of the gastrointestinal microbiota in colorectal cancer. *Curr Pharm Des* 15: 1524-1527. doi: 10.2174/138161209788168191. PubMed: 19442169.
 48. Correa P, Haenszel W (1978) The epidemiology of large bowel cancer. *Adv Cancer Res* 26: 1-141. doi:10.1016/S0065-230X(08)60086-X. PubMed: 343522.
 49. Cats A, Devries E, Mulder N, Kleibeuker J (1996) Regional differences of physiological functions and cancer susceptibility in the human large intestine. *Int J Oncol* 9: 1055-1069. PubMed: 21541613.
 50. Macfarlane G, Gibson G, Beatty E, Cummings J (1992) Estimation of short-chain fatty acid production from protein by human intestinal bacteria based on branched-chain fatty acid measurements. *FEMS Microbiol Lett* 101: 81-88. doi:10.1016/0168-6496(92)90049-Y.
 51. Cummings JH, Macfarlane GT (1991) The control and consequences of bacterial fermentation in the human colon. *J Appl Microbiol* 70: 443-459. doi:10.1111/j.1365-2672.1991.tb02739.x. PubMed: 1938669.
 52. Lin HV, Frassetto A, Kowalik EJ Jr, Nawrocki AR, Lu MM et al. (2012) Butyrate and Propionate Protect against Diet-Induced Obesity and Regulate Gut Hormones via Free Fatty Acid Receptor 3-Independent Mechanisms. *PLOS ONE* 7: e35240. doi:10.1371/journal.pone.0035240. PubMed: 22506074.
 53. Murphy KG, Bloom SR (2006) Gut hormones and the regulation of energy homeostasis. *Nature* 444: 854-859. doi:10.1038/nature05484. PubMed: 17167473.
 54. Harmsen HJM, Wildeboer-Veloo ACM, Grijpstra J, Knol J, Degener JE et al. (2000) Development of 16S rRNA-Based Probes for the Coriobacterium Group and the Atopobium Cluster and Their Application for Enumeration of Coriobacteriaceae in Human Feces from Volunteers of Different Age Groups. *Appl Environ Microbiol* 66: 4523-4527. doi:10.1128/AEM.66.10.4523-4527.2000. PubMed: 11010909.
 55. Manz W, Amann R, Ludwig W, Vancanneyt M, Schleifer KH (1996) Application of a suite of 16S rRNA-specific oligonucleotide probes designed to investigate bacteria of the phylum Cytophaga-Flavobacter-Bacteroides in the natural environment. *Microbiology* 142: 1097-1106. doi:10.1099/13500872-142-5-1097. PubMed: 8704951.
 56. Laitinen R, Malinen E, Palva A (2002) PCR-ELISA: I: Application to simultaneous analysis of mixed bacterial samples composed of intestinal species. *Syst Appl Microbiol* 25: 241-248. doi: 10.1078/0723-2020-00118. PubMed: 12353879.
 57. Franks AH, Harmsen HJM, Raangs GC, Jansen GJ, Schut F et al. (1998) Variations of bacterial populations in human feces measured by fluorescent in situ hybridization with group-specific 16S rRNA-targeted oligonucleotide probes. *Appl Environ Microbiol* 64: 3336-3345. PubMed: 9726880.
 58. Harmsen Hermie JM, Schut Frits, Gjalit W, Welling (1999) A 16S rRNA-targeted Probe for Detection of Lactobacilli and Enterococci in Faecal Samples by Fluorescent In Situ Hybridization. *Microb Ecol Health Dis* 11: 3-12. doi:10.1080/089106099435862.
 59. Walker AW, Duncan SH, McWilliam Leitch EC, Child MW, Flint HJ (2005) pH and peptide supply can radically alter bacterial populations and short-chain fatty acid ratios within microbial communities from the human colon. *Appl Environ Microbiol* 71: 3692-3700. doi:10.1128/AEM.71.7.3692-3700.2005. PubMed: 16000778.
 60. Daims H, Brühl A, Amann R, Schleifer KH, Wagner M (1999) The domain-specific probe EUB338 is insufficient for the detection of all Bacteria: development and evaluation of a more comprehensive probe set. *Syst Appl Microbiol* 22: 434-444. doi:10.1016/S0723-2020(99)80053-8. PubMed: 10553296.