

Adjustments in the forcing-feedback framework for understanding climate change

Article

Accepted Version

Sherwood, S. C., Bony, S., Boucher, O., Bretherton, C., Forster, P. M., Gregory, J. M. ORCID: https://orcid.org/0000-0003-1296-8644 and Stevens, B. (2015) Adjustments in the forcing-feedback framework for understanding climate change. Bulletin of the American Meteorological Society, 96 (2). pp. 217-228. ISSN 1520-0477 doi: 10.1175/BAMS-D-13-00167.1 Available at https://centaur.reading.ac.uk/38337/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

To link to this article DOI: http://dx.doi.org/10.1175/BAMS-D-13-00167.1

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the <u>End User Agreement</u>.

www.reading.ac.uk/centaur



CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Bulletin of the American Meteorological Society

Adjustments in the forcing-feedback framework for understanding climate change --Manuscript Draft--

Manuscript Number:	BAMS-D-13-00167
Full Title:	Adjustments in the forcing-feedback framework for understanding climate change
Article Type:	Article
Corresponding Author:	Steven Sherwood UNSW Kensington, NSW AUSTRALIA
Corresponding Author's Institution:	UNSW
First Author:	Steven Sherwood
Order of Authors:	Steven Sherwood
	Sandrine Bony
	Olivier Boucher
	Chris Bretherton
	Piers Forster
	Jonathan Gregory
	Bjorn Stevens
Manuscript Classifications:	4.088: Climate change; 4.100: Climate sensitivity; 4.208: Energy budget/balance; 4.228: Feedback; 8.020: Climate models
Abstract:	The traditional forcing-feedback framework has provided an indispensable basis for discussing global climate changes. However, as analyses of model behavior has become more detailed, shortcomings and ambiguity in the framework have become more evident and physical effects unaccounted for by the traditional framework have become interesting. In particular, the new concept of adjustments, which are responses to forcings that are not mediated by the global mean temperature, has emerged. This concept, related to the older ones of climate efficacy and stratospheric adjustment, is a more physical way of capturing unique responses to specific forcings. We present a pedagogical review of the adjustment concept, why it is important, and how it can be used. The concept is particularly useful for aerosols, where it helps to organize what has become a complex array of forcing mechanisms. It also helps clarify issues around cloud and hydrological response, transient vs. equilibrium climate change, and geoengineering.
Author Comments:	One figure (3) is excerpted from a figure in a paper by Zelinka et al. published last year in J. Climate. Another figure (4) is taken from the latest IPCC report, but we have added one small additional feature to it which communicates an important point not addressed by that report. Please advise if either of these might raise a copyright issue.
Suggested Reviewers:	TImothy Andrews timothy.andrews@metoffice.uk.gov
	Robert Colman R.Colman@bom.gov.au
	Youichi Kamae kamae@aori.u-tokyo.ac.jp
	Drew Shindell drew.t.shindell@nasa.gov



Never Stand Still

Science

Climate Change Research Centre

26 June 2014

To: Editors, BAMS

Re: BAMS-D-13-00167

We hereby submit the first revision of the article "Adjustments in the forcing-feedback framework for understanding climate change". The text length and figure number are unchanged from the previous draft. We attach a separate "response to reviewers" file.

Many thanks for considering this submission.

Sincerely,

and that

Steven Sherwood

1	Adjustments in the forcing-feedback framework for understanding
2	climate change
3	Steven C. Sherwood, *
	Climate Change Research Centre and ARC Centre of Excellence for Climate System Science,
	University of New South Wales, Sydney, NSW Australia
4	SANDRINE BONY AND OLIVIER BOUCHER
	Laboratoire de Météorologie Dynamique/IPSL, Université P. et M. Curie, Paris, France
5	CHRIS BRETHERTON
	Dept. of Meteorology, University of Washington, Washington, USA
6	Piers M. Forster
	School of Earth and Environment, University of Leeds, Leeds, UK
7	Jonathan M. Gregory
	NCAS-Climate, University of Reading, Reading, UK
	Met Office Hadley Centre, Exeter, UK
8	BJORN STEVENS
	Max Planck Institute for Meteorology, Hamburg, Germany

^{*} Corresponding author address: S. Sherwood, Climate Change Research Centre, University of New South

ABSTRACT

The traditional forcing-feedback framework has provided an indispensable basis for dis-10 cussing global climate changes. However, as analysis of model behavior has become more 11 detailed, shortcomings and ambiguities in the framework have become more evident and 12 physical effects unaccounted for by the traditional framework have become interesting. In 13 particular, the new concept of adjustments, which are responses to forcings that are not 14 mediated by the global mean temperature, has emerged. This concept, related to the older 15 ones of climate efficacy and stratospheric adjustment, is a more physical way of capturing 16 unique responses to specific forcings. We present a pedagogical review of the adjustment 17 concept, why it is important, and how it can be used. The concept is particularly useful for 18 aerosols, where it helps to organize what has become a complex array of forcing mechanisms. 19 It also helps clarify issues around cloud and hydrological response, transient vs. equilibrium 20 climate change, and geoengineering. 21

9

Wales, Sydney, Australia.

E-mail: S.Sherwood@unsw.edu.au

25

The traditional and now ubiquitous framework for understanding global climate change 26 involves an external forcing, a response whereby the climate system opposes the forcing in 27 order to regain equilibrium, and feedbacks which amplify or damp the response. The concept 28 is most often applied to the global mean surface temperature \overline{T} , where the external forcing 29 is a radiative perturbation (effective power input) $d\overline{F}$, and $d\overline{T}$ is the change in \overline{T} produced 30 by $d\overline{F}$ at the new equilibrium¹. This new equilibrium is achieved when the system response 31 has caused a change $d\overline{R}$ in net rate of energy loss by the planet that balances the effect of 32 the imposed perturbation, i.e., so that $\overline{N} = d\overline{F} - d\overline{R} = 0$ where \overline{N} is the net power into 33 the planet from space. Feedback arises because there are various quantities X_i (atmospheric 34 water vapor or sea ice cover for example) which depend on \overline{T} and alter the planetary energy 35 budget. The new equilibrium encompasses all of their effects as well: 36

$$\mathrm{d}\overline{R} = \mathrm{d}\overline{T} \left(\frac{\partial\overline{R}}{\partial\overline{T}} + \sum_{i} \frac{\partial\overline{R}}{\partial X_{i}} \frac{\mathrm{d}X_{i}}{\mathrm{d}\overline{T}} \right) = \mathrm{d}\overline{F}$$
(1)

The ratio $(d\overline{R}/d\overline{T})^{-1}$ is called the "climate sensitivity parameter" (the "equilibrium climate sensitivity" being usually defined as $d\overline{T}$ for a forcing equivalent to a doubling of CO₂). The term $(\partial \overline{R}/\partial \overline{T})$ is the "Planck response," or change that \overline{R} would undergo if the climate system behaved as a black body with no feedbacks. The black-body system is stable to radiative perturbations because $(\partial \overline{R}/\partial \overline{T}) > 0$. This traditional approach is illustrated in

¹Following the custom in climate literature, we use "equilibrium" in the loose sense of a system that is in a statistically steady state of energy balance. This is not a strict equilibrium since the Earth is constantly generating and exporting entropy.

⁴² Fig. 1b.

There are ambiguities across disciplines in what is meant by "feedback," a concept well 43 known in electronics and control theory (Bates 2007). We will follow the usual custom of 44 detailed climate feedback analysis studies (see Hansen et al. 1984; Schlesinger and Mitchell 45 1987; Soden and Held 2006) by referring to the amplifying role of a system property X_i , 46 quantified by $\alpha_i \equiv -(\partial \overline{R}/\partial X_i)(\mathrm{d}X_i/\mathrm{d}\overline{T})$ as in (1), as a feedback. A feedback is "positive" 47 if it amplifies the change in \overline{T} ; in this case $\alpha_i > 0$. A system including feedbacks is stable 48 if the sum of the α_i is smaller than the Planck response². The forcing-feedback paradigm 49 has helped establish, for example, the dominant role of water vapor in amplifying global 50 temperature change and the role of clouds in accounting for its uncertainty (Cess 1990). 51

Many potential feedbacks within the Earth system can be conceived, involving system 52 components having a wide range of characteristic response times (Dickinson and Schaudt 53 1998; Jarvis and Li 2011). Clouds and water vapor can respond to climate changes in days 54 to weeks, whereas the deep ocean or ice sheets may require centuries or millennia. Feedbacks 55 that are not fast enough to fully keep pace with responses of interest, due to the involvement 56 of a slowly-varying component such as the deep ocean or ice sheets, may appear to have time-57 varying strengths (e.g. Senior and Mitchell 2000); those may be better treated as exogenous 58 forcings in transient calculations. The "equilibrium" (sometimes called "Charney") climate 59 sensitivity (ECS) has become the standard measure of the climate sensitivity of the Earth 60 system relevant to the anthropogenic warming problem. It was adopted from early slab-61 ocean model experiments that were run to a less-complete equilibrium in which ice sheets, 62 vegetation and atmospheric composition were all specified. The paradigm can however be 63 extended to include, for example, changes to natural sources of CO_2 as "carbon cycle" 64

²Alternatively, the Planck response can also be considered as a strong negative feedback that is more direct and simpler than the others. This avoids giving it a special status and thus makes (1) more symmetrical (Gregory et al. 2009) in the sense that the system is stable if the sum of all the feedbacks, including the Planck response, is negative. However, use of the word "feedback" to describe the Planck response is confusing because there is nothing being "fed back upon."

feedbacks (e.g. Gregory et al. 2009; Arneth et al. 2010; Raes et al. 2010). Additional feedbacks
not considered in the ECS will enter on longer (e.g., geologic) time scales.

In applying (1) to the global climate one normally assumes that all partial derivatives represent constants of the climate system, but this implies at least two bold assumptions. The first is that responses vary *linearly* with perturbation amplitudes (or equivalently, are not state dependent). Formally this must hold for sufficiently small perturbations, but possibly not for multiple doublings of CO₂ as some feedbacks may become stronger or weaker in significantly different climates (Crucifix 2006; Caballero and Huber 2013; Colman and McAvaney 2009).

The second assumption is that all responses are uniquely determined by the scalar $d\overline{T}$ 74 regardless of how the temperature change is brought about, a situation that may be called 75 fungibility. Complete fungibility requires either that temperature changes always occur with 76 the same spatial and seasonal pattern, or that different patterns produce the same $d\overline{R}/d\overline{T}$ 77 where $d\overline{R}$ is the change in global- and annual-mean radiation balance. However this will 78 not be the case since different forcings will generally produce different warming patterns. 79 Moreover, during transient warming regardless of how it is forced, some parts of the ocean 80 may warm more slowly than others, temporarily producing anomalous warming patterns. In 81 the absence of feedbacks these pattern differences should not strongly affect the paradigm, 82 since the Planck non-linearity is sufficiently weak (Bates 2012). Many quantities X that 83 affect the global radiation budget, however, are sensitive to spatial or seasonal variations in 84 temperature. This sensitivity lies at the root of difficulties that have emerged over the years 85 with the traditional framework (Hansen et al. 1997). 86

WHY ADJUSTMENTS? The radiative forcing concept is, in effect, a "common currency" that we may use to compare various types of perturbation: emissions of CO₂ or other pollutants, changes in land use, solar activity, etc. The concept is useful only to the extent that it accurately predicts the magnitude of the response without having to worry about any other details of the perturbation—that is, insofar as feedbacks are independent of the ₉₂ perturbation.

While one might imagine that the instantaneous impact of a perturbation on the top-93 of-atmosphere (TOA) radiation balance would be a good measure of its radiative forcing, 94 early studies quickly recognized that this measure was not optimal. The temperature of the 95 stratosphere, in particular, was not closely tied to that of the surface. For example it warms 96 under a positive solar forcing, yet cools under a positive greenhouse gas forcing (Fels et al. 97 1980) therefore requiring the surface and troposphere to warm more to balance the same 98 instantaneous TOA net flux perturbation (Hansen et al. 1997). This problem was resolved 99 by allowing for a "stratospheric adjustment" prior to calculating the radiative forcing, which 100 has been the standard approach at least since the first IPCC report (IPCC 1990). 101

Recent work reveals that heterogeneous responses also occur within the troposphere and 102 can produce similar, but more subtle problems. Some of the most important mechanisms 103 by which this can occur are illustrated in Fig. 2. For example, increasing the concentration 104 of CO_2 in the atmosphere affects longwave radiative fluxes and slightly warms the mid-105 and lower troposphere, even with no surface temperature change (see Fig. 2b). In models 106 this subtle change in stratification and relative humidity reduces middle and low-altitude 107 cloud cover (Fig. 3), further altering the TOA net flux even before any global warming or 108 cloud feedbacks take place (Andrews and Forster 2008; Gregory and Webb 2008; Colman 109 and McAvaney 2011; Kamae and Watanabe 2012a; Wyant et al. 2012). This change in cloud 110 cover is quite different to that which occurs subsequently due to the increase in \overline{T} (compare 111 Fig. 3a and Fig. 3b). Likewise, any forcing that is horizontally inhomogeneous (for example 112 changes in tropospheric ozone, or aerosols, discussed below) or that significantly affects the 113 tropospheric radiative cooling will drive changes in atmospheric circulation that may alter 114 the planetary albedo by changing patterns of cloud cover. Conceptually these complications 115 are no different from the one long recognized for the stratosphere. 116

¹¹⁷ We refer to changes that occur directly due to the forcing, without mediation by the ¹¹⁸ global-mean temperature, as "adjustments" and the accordingly modified top-of-atmosphere radiative imbalance as the "effective" radiative forcing, following Boucher et al. (2013). Their
role is illustrated in Fig. 1a.

Most adjustments are rapid, but there is no fundamental time scale that separates rapid 121 adjustments from feedback responses. The time scales of the two can, in principle, over-122 lap significantly. Some adjustments can for example occur through state variables X that 123 respond very slowly (e.g., vegetation cover or soil humidity responses to CO₂-induced stom-124 atal closure (Doutriaux-Boucher et al. 2009) or aerosol-induced diffuse sunlight (Mercado 125 et al. 2009)—see Fig. 2d—or responses involving stratospheric composition and chemistry). 126 Meanwhile some feedbacks, such as the water-vapor feedback, can be triggered by warming 127 of the land and atmosphere (Colman and McAvaney 2011) that occurs within days or weeks 128 of an applied forcing. Indeed the original stratospheric adjustment requires several months 129 to complete and has to be calculated by a special model run with the troposphere and surface 130 held fixed. However, the largest tropospheric adjustments are likely due to changes in clouds 131 driven by changes in tropospheric radiative fluxes, and appear to occur within days (Dong 132 et al. 2009). 133

ADJUSTING OUR VIEW OF AEROSOLS It turns out that the climate community 134 has been grappling with tropospheric adjustments for years, but without calling them by 135 this name: they play a dominant role in the climatic impact of aerosols. One example 136 is the "semi-direct effect" of aerosols, triggered by the uneven distribution of tropospheric 137 radiative heating by the aerosol. This can subtly alter atmospheric stability which will affect 138 convection (Fig. 2b), and because it is horizontally heterogeneous, it can drive circulations 139 (Fig. 2a) that alter both the global cloud radiative effect and patterns of temperature and 140 rainfall. This response should be regarded as a rapid adjustment to aerosol perturbations to 141 the radiation field, since it occurs even in the absence of a change in T. 142

Likewise, the cloud-mediated (or "indirect") impact of aerosols, which serve as cloud condensation nuclei (CCN), on climate involves rapid adjustments. This impact begins with an increase in the number of nucleated droplets which, in the absence of any changes to the ¹⁴⁶ water content or circulation of air within the cloud, would produce a cloud with higher albedo ¹⁴⁷ (often called the "Twomey effect"). Model studies indicate however that the knock-on effects ¹⁴⁸ that occur via changes in the flow field or the microphysical evolution of clouds can lead to ¹⁴⁹ final changes in albedo that differ substantially from this initial droplet number effect. A ¹⁵⁰ number of such knock-on effects have been articulated in the literature including "lifetime ¹⁵¹ effect," liquid water path effect, etc. Most of these are based on idealized conceptual models, ¹⁵² and their applicability to real clouds remains controversial (Boucher et al. 2013).

The initial and the various knock-on effects are often conceptualized as each having 153 distinct physical significance, but in more realistic simulations it is typically not possible to 154 distinguish them individually, and the assumptions under which they were deduced often 155 do not hold. Only their combined effect can be properly diagnosed. We argue that the 156 subsequent change in \overline{T} is a response to this net radiative effect of aerosol—the aerosol 157 effective radiative forcing or ERF, which includes the initial droplet-number effect and all 158 adjustments. This concept is not new, and has been referred to in the literature before as a 159 quasi-forcing (Rotstayn and Penner 2001) or a radiative flux perturbation (Lohmann et al. 160 2010). 161

In both the CO₂ and aerosol cases, adjustments are more uncertain than instantaneous forcings because they involve cloud and other dynamical responses that models may not calculate reliably. Regardless of this, models suggest these effects can be large, so they cannot be ignored.

WAYS OF DEFINING AND CALCULATING ADJUSTMENTS The ERF concept is motivated mainly by the desire to improve fungibility within the forcing-response framework, that is, minimize the quantitative differences of $d\overline{T}$ to various types of forcing $d\overline{F}$. Ideally we would choose an adjustment framework that optimises this, aiming for the ERF to be the forcing experienced by the system when $d\overline{T} = 0$. There is however no unambiguous way to specify this, because regionally heterogeneous surface temperature changes occur immediately after a forcing is applied.

Two common approaches are available for quantifying the adjustment, with different 173 advantages and disadvantages. The first or "regression method" (sometimes called Gregory 174 method) is to regress the net TOA flux perturbation \overline{N} onto $d\overline{T}$ in a transient warming 175 simulation, yielding a plot, (see Fig. 4) in which the $d\overline{T} = 0$ intercept is the ERF (Gregory 176 et al. 2004). The second or "fixed-SST" method (sometimes called the Hansen method) 177 diagnoses ERF, $d\overline{F}$ and \overline{N} from a simulation including the forcing agent but with sea-surface 178 temperatures and sea ice prescribed to their unperturbed climatology (Cess and Potter 1988; 179 Hansen et al. 2005, see also Fig. 1c). 180

The regression method implicitly defines adjustments as those changes which occur rel-181 atively soon (within a few years), including those mediated by regional variations in SST 182 change. The latter are excluded by the fixed-SST approach, which does on the other hand 183 include all other forcing-related adjustments no matter how long they take to occur (provided 184 the simulation is long enough). The regression method can be thrown off by time-varying 185 feedbacks, in which case \overline{N} versus $d\overline{T}$ will not be a straight line. However this method sat-186 isfies the principle of cleanly separating adjustments from global mean temperature change 187 $d\overline{T}$, whereas the fixed-SST method permits land temperature change which contributes to 188 $\mathrm{d}\overline{T}$ and affects the air-sea temperature difference over oceans (see Kamae and Watanabe 189 2012b; Shine et al. 2003; Vial et al. 2013). This enhances the global Planck response and 190 triggers some warming-related changes such as an increase in global atmospheric water vapor 191 (Colman and McAvaney 2011), the effects of which should be subtracted out if one wishes 192 to isolate true adjustments from changes that result from feedbacks. 193

These two methods are shown for a typical CMIP model in Fig. 4. A third method that has been used in the literature for precipitation responses (examined further below) is to assume that the change during some limited time period (e.g., one year) following an abrupt forcing, compared to the climatology before, is due to adjustments. However, the $d\overline{T}$ during this period is substantial, making it difficult to quantitatively compare with the other approaches.

The regression-based ERF estimate from a single simulation is inherently noisier than the 200 fixed-SST one and is best suited for global-mean rather than regional responses. However, it 201 can be made more precise by averaging across an ensemble of at least 5-10 shorter coupled 202 simulation pairs of 10-20 years in which the step change in CO_2 from the control is made 203 at different times to average over natural climate variability (e.g. Watanabe et al. 2012, 204 and see next section). The fixed-SST ERF estimate is naturally more robust to internal 205 climate variability because it takes advantage of the long averaging time, and the fact that 206 the interannual ocean variability is either absent or identical in the perturbed and control 207 simulations. 208

To the extent that the temperature-mediated response of the climate system is linear and invariant to the warming pattern, these methods should give almost identical results when the latter is corrected for the change in $d\overline{T}$ at fixed SST. As seen in Fig. 4, this is approximately true for global mean quantities, but there are noteworthy differences.

In all CMIP5 models for which the needed output has been published, the fixed-SST 213 $4 \times CO_2$ ERF exceeds the regression-based one, usually by a statistically significant margin. 214 This point has been obscured because the literature has reported the former without the 215 aforementioned $\sim 0.5 \text{ W m}^{-2}$ feedback correction for land warming. Applying this correction 216 to the seven CMIP5 models in Table 1 of Andrews et al. (2012) reporting both estimates, the 217 fixed-SST ERF exceeds the regression ERF by about 15% (0.2-1.6 W m⁻² at $4 \times CO_2$, with 218 a mean of 1 W m⁻²). The HadGEM2-ES model exhibits a particularly large discrepancy 219 due to a somewhat nonlinear response of \overline{N} to the warming $d\overline{T}$ during the first year or 220 two. Andrews et al. (2012) traced this response to an increase over time in cloud shortwave 221 radiative feedback over oceans. Since this increase seems to occur in many models, it merits 222 further study. 223

Watanabe et al. (2012) showed in one model how an ENSO-like SST anomaly can set up in the first year or two after CO_2 increase due to weakening of the Walker circulation (see also Bony et al. 2013); this is an example of an adjustment not captured by the fixedSST framework. In parts of the oceans with relatively shallow mixed layers the SST can respond more rapidly than in others, leading to the emergence of fast changes in SST patterns while the global $d\overline{T}$ is still small (Armour et al. 2013). These changes influence cloud and circulation patterns (see next Section), amplify the atmospheric adjustments and can be aliased onto changes in global-mean cloud radiative effects in some models.

GCM-based estimates of the radiative forcing of anthropogenic aerosol on climate have often been based on comparison of fixed-SST simulations with pre-industrial and present-day aerosol emissions. These estimates are universally uncorrected for the associated change in global surface air temperature due to land temperature change; in principle an approximately 1 W m^{-2} correction per K of $d\overline{T}$ should be added to them to be fully consistent with the ERF paradigm. However, at least for one model checked (CAM5), the surface air temperature change is less than 0.1 K, so this correction is negligible for most purposes.

In summary, ERF is a construct designed to fit the global radiative response of a model 239 as a linear function of $d\overline{T}$ over timescales of decades to a century. From this perspective, 240 the regression ERF is preferable to the fixed-SST one since it is based on precisely the 241 linear fit which is used for global feedback analysis, but this fit is imperfect, especially if 242 applied to regional responses to $d\overline{T}$ rather than global-mean ones. The difference in results 243 between the two methods can be interpreted as an indicator of short-term deviations from 244 linearity in the relation of \overline{N} to $d\overline{T}$. Such deviations seem to arise from the knock-on effects 245 of inhomogeneous surface warming. The attribution of this to adjustment or feedback is 246 inherently ambiguous, should depend on the circumstances and goals of the analysis, and 247 will be different between the two methods considered here. 248

PRECIPITATION Rapid adjustments to \overline{N} caused by CO₂ and aerosol are difficult if not impossible to detect in observations. We can however look for these physical effects in quantities other than the TOA radiative flux. Notably, we can consider the direct impact of a CO₂ change on precipitation, in the absence of any global-mean (or ocean-mean) \overline{T} change. We should note however that because precipitation patterns are sensitive to small changes in the temperature pattern, we would expect regional precipitation changes to be relatively forcing-dependent even in the absence of adjustments—for example, a forcing that causes warming asymmetrically distributed between the hemispheres shifts tropical rain maxima toward the hemisphere of greater warming(Seo et al. 2014). Thus rapid adjustments alone may not explain all forcing-dependence of precipitation responses.

Possible adjustments of precipitation to aerosol perturbations (both radiative and cloud-259 microphysical) are now well recognized in principle, but poorly understood, hence controver-260 sial. For instance, by absorbing solar radiation, increased black carbon aerosols will cause 261 a slight decrease in global-mean precipitation for the same surface temperature (Andrews 262 et al. 2010). However, regional precipitation changes may be more important, and can occur 263 far away from the aerosol that drives them (Wang 2013). Models suggest that, due to their 264 heterogeneous heating of the atmosphere and surface, aerosol-radiation interactions can af-265 fect monsoons (Ramanathan et al. 2005; Lau et al. 2006), shift the inter-tropical convergence 266 zone (Rotstayn and Lohmann 2002), and displace atmospheric jets poleward (Allen et al. 267 2012). Some of these studies have argued that these effects can be detected in observed 268 rainfall trends. CCN-mediated effects on precipitation also have attracted great attention 269 but are even more controversial (e.g. Tuttle and Carbone 2011; Tao et al. 2012). 270

Less recognized are the direct effects of solar or greenhouse-gas perturbations on precip-271 itation. CO_2 warms and stabilizes the lower troposphere, slowing the global hydrological 272 cycle for a given \overline{T} (Allen and Ingram 2002; Andrews et al. 2010) and slowing and causing 273 a redistribution of the tropical overturning circulation (Andrews et al. 2010; Wyant et al. 274 2012; Bony et al. 2013). The shifts in tropical rainfall associated with this effect make up 275 a substantial part of the total circulation-driven rainfall change in climates simulated by 276 the end of the 21st century (Bony et al. 2013). The change in global-mean rainfall is also 277 nontrivial compared to that from warming. These effects on rainfall are somewhat more 278 pronounced than those on TOA radiative balance, where adjustments appear to account for 279 no more than 20% of global-mean $d\overline{T}$ in a multi-model average (though also contributing to 280

forcing uncertainty, Forster et al. 2013; Vial et al. 2013). Much of the precipitation adjustment to CO_2 occurs very rapidly, within a week (Fig. 5). Thus, precipitation adjustments to CO_2 may stand a better chance of eventually being detectable in observations than would the TOA radiation adjustments.

Determining the spatial distribution of the precipitation adjustment is challenging be-285 cause precipitation is highly variable on interannual and longer time scales, and sensitive to 286 gradients in tropical sea surface temperature. Unforced anomalies that happen to occur after 287 a step increase in forcing will be confounded with adjustments, necessitating an ensemble 288 average to obtain the latter accurately from abrupt-forcing scenarios (Fig. 6). Moreover, 289 atmospheric responses to forcings can quickly drive changes to the surface oceans, espe-290 cially near the equator, that can strongly amplify or otherwise alter regional precipitation 291 responses (compare panels a,b of Fig. 6; see also Chadwick et al. 2014). Such knock-on 292 responses (which also affect top-of-atmosphere radiation) should be regarded as part of the 293 adjustment to the extent that they involve SST gradients rather than the global mean \overline{T} , 294 although again there is no unambiguous separation. 295

CONCLUSION In response to changing concentrations of CO_2 or other forcings, the 296 climate system changes in ways that are independent of any global-mean surface temperature 297 change, but which subsequently influence the global-mean radiation budget and hence surface 298 temperature. These adjustments also appear to affect other climate quantities significantly, 299 in particular precipitation. They are physically significant, depend on the forcing agent, and 300 need to be accounted for when computing the radiative forcing of the agent. Many of them 301 develop on a time scale of days (Cao et al. 2012; Kamae and Watanabe 2012a; Bony et al. 302 2013). Accounting more appropriately for adjustments offers new opportunities to better 303 understand, predict, and evaluate impacts of different perturbations. 304

The fact that adjustments scale with the amplitude of the forcing rather than that of the global warming response means that even if global-mean temperature were for some reason very insensitive to forcing—due for example to some hypothetical strong negative feedback from clouds—the adjustments would remain unaffected. It also implies that part
 of the climate response to forcing is independent of the long-standing uncertainty in climate
 sensitivity.

While the forcing-feedback paradigm has always been recognised as imperfect, such discrepancies have previously been attributed to variations in "efficacy" (Hansen et al. 1984), which did not clarify their nature. Decomposing the climate response into a forcing-specific adjustment and a \overline{T} -mediated response that is more forcing-independent provides a clearer way of understanding climate changes, especially transient ones.

The adjustment concept needs to be fully integrated into energy budget studies (e.g. 316 Otto et al. 2013). Estimates of radiative forcings and climate sensitivity ought to be defined 317 consistently when observations are used to constrain estimates of the radiative forcings, 318 climate sensitivity or both quantities. The traditional climate sensitivity is actually the 319 product of two quantities, the radiative forcing of a doubling of CO_2 and a climate sensitivity 320 parameter in units of K $(W m^{-2})^{-1}$, where it has been assumed that the former is known 321 exactly. In fact it is not, especially when adjustments are considered as part of the forcing 322 (see Webb et al. 2013; Stevens and Schwartz 2012). Because adjustments make the forcing 323 uncertain, future studies should distinguish between the traditional climate sensitivity, which 324 depends on adjustments, and the climate sensitivity parameter, which does not. 325

This decomposition may clarify some past reports of feedbacks appearing to be state-326 dependent, forcing-dependent or time-dependent, although not all such complexities are 327 likely be resolved and some variations in efficacy will remain. Studies already show that 328 transient climate changes at arbitrary times while the system is out of equilibrium, can be 329 approximately recovered by adding the rapid adjustment to CO_2 at that time to the (lagging) 330 temperature-mediated response (Andrews et al. 2010). To a considerable extent this also 331 works for multi-model mean precipitation responses (Bony et al. 2013). This leads to a 332 considerable simplification and will be useful to those exploring climate change using simple 333 models that are only a function of global-mean radiative forcings (e.g., Huntingford and 334

Cox 2000), especially if such models explore scenarios (e.g., overshooting of carbon targets, ramping up and down of greenhouse gas forcings) that stray from those for which they have been fitted to the behavior of a more detailed climate model.

This approach also helps us to understand and anticipate the limitations of potential geo-338 engineering strategies. Idealized climate simulations of solar radiation management (Bala 339 et al. 2008; Schmidt et al. 2012) show that when the warming due to CO_2 increase is coun-340 teracted by a decrease in the solar constant, the warming-induced impacts are reversed to 341 a large extent but adjustment responses may linger. In the case of precipitation, the rapid 342 adjustment to the higher CO_2 amount is not counteracted by a commensurate adjustment 343 in response to lower solar radiation, leaving a net decrease in global-mean precipitation as 344 well as larger residual changes at the regional level. 345

Indeed changes to solar radiation, volcanic aerosol, and orbital properties are each likely to lead to distinct adjustments to global rainfall and regional climate patterns in addition to their common impact on global-mean temperature. Recognizing and understanding the adjustments may be crucial in helping to make sense of both present and past climate changes. For instance accounting for adjustments helps interpret differences in the precipitation response to natural vs anthropogenic forcing (Liu et al. 2013).

Rapid adjustments involve rapid processes, and may present an opportunity to use model 352 evaluations on very short time scales to constrain processes that are also important for 353 longer-term climate change. Such systematic weather-forecast type of verification has been 354 suggested as a possible way to improve the representation of model processes (Brown et al. 355 2012), but may also aid our understanding of the multiple adjustments associated with 356 aerosol-cloud interactions (Boucher 2012) or the physical processes that control the direct 357 effect of CO_2 on circulation and precipitation (Bony et al. 2013). Since rapid adjustments 358 closely track the forcing variations in time, they may contribute substantially to initial tran-359 sient climate changes even if making up a relatively small part of the long-term equilibrium 360 response. This may offer an opportunity to better detect and disentangle the relative role of 361

different forcings on climate, including on regional responses, and thus to facilitate detection and attribution studies. There are thus two broad reasons for future studies to distinguish more carefully between forcing-specific adjustments and temperature-driven feedback responses: to clarify our understanding of past climate changes, and to exploit what the relationships among various adjustment and feedback responses may be able to tell us about the climate system and how it will respond in the future.

368 Acknowledgments.

We thank N. Huneeus for generous help leading to Figure 5. CB acknowledges support from NOAA grant NA13OAR4310104. PF was supported by a Royal Society Wolfson Merit Award and EPSRCEP/I014721/1(http://gow.epsrc.ac.uk/NGBOViewGrant.aspx

372 ?GrantRef=EP/I014721/1>. Additional support (for BS and SB) was provided through
373 funding from the European Union Seventh Framework Programme (FP7/2007-2013) under
374 grant agreement number 244067

REFERENCES

- Allen, M. R. and W. J. Ingram, 2002: Constraints on future changes in climate and the hydrologic cycle. *Nature*, **419**, 224–232.
- Allen, R. J., S. C. Sherwood, J. R. Norris, and C. S. Zender, 2012: Recent Northern Hemisphere tropical expansion primarily driven by black carbon and tropospheric ozone. *Nature*,
 485, 350–U93.
- Andrews, T. and P. M. Forster, 2008: CO₂ forcing induces semi-direct effects with consequences for climate feedback interpretations. *Geophys. Res. Lett.*, **35** (4), L04 802.
- Andrews, T., P. M. Forster, O. Boucher, N. Bellouin, and A. Jones, 2010: Precipitation, radiative forcing and global temperature change. *Geophys. Res. Lett.*, **37**, L14701.
- Andrews, T., J. M. Gregory, M. J. Webb, and K. E. Taylor, 2012: Forcing, feedbacks and
 climate sensitivity in CMIP5 coupled atmosphere-ocean climate models. *Geophys. Res. Lett.*, **39**, L09712.
- Armour, K. C., C. M. Bitz, and G. H. Roe, 2013: Time-varying climate sensitivity from
 regional feedbacks. J. Climate, 26, 4518–4534.
- Arneth, A., et al., 2010: Terrestrial biogeochemical feedbacks in the climate system. Nat.
 Geosci., 3, 525–532.
- Bala, G., P. B. Duffy, and K. E. Taylor, 2008: Impact of geoengineering schemes on the
 global hydrological cycle. *Proc. Nat. Acad. Sci. US.*, 105, 7664–7669.
- Bates, J. R., 2007: Some considerations of the concept of climate feedback. Q. J. R. Meteorol.
 Soc., 133, 545–560.

376

- Bates, J. R., 2012: Climate stability and sensitivity in some simple conceptual models. *Clim. Dyn.*, 38, 455–473.
- Bony, S., G. Bellon, D. Klocke, S. Sherwood, S. Fermepin, and S. Denvil, 2013: Robust direct
 effect of carbon dioxide on tropical circulation and regional precipitation. *Nat. Geosci.*, 6,
 447–451.
- ⁴⁰² Boucher, O., 2012: The aerosol effect: An essential pursuit. *Nature*, **490**, 40–41.
- Boucher, O., et al., 2013: Clouds and aerosols. Climate Change 2013: The Physical Science
 Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, T. F. Stocker, D. Qin, G.-K. Plattner, S. K. A.
 M. Tignor, J. Boschung, A. Nauels, Y. Xia, V. Bex, and P. M. Midgley, Eds., Cambridge
- ⁴⁰⁷ University Press, Cambridge, United Kingdom and New York, NY, USA.
- Brown, A., S. Milton, M. Cullen, B. Golding, J. Mitchell, and A. Shelly, 2012: Unified
 modeling and prediction of weather and climate: A 25-year journey. *Bull. Amer. Meteorol.*Soc., 93, 1865–1877.
- Caballero, R. and M. Huber, 2013: State-dependent climate sensitivity in past warm climates
 and its implications for future climate projections. *Proc. Nat. Acad. Sci. US.*, **110**, 14162–
 14167.
- Cao, L., G. Bala, and K. Caldeira, 2012: Climate response to changes in atmospheric carbon
 dioxide and solar irradiance on the time scale of days to weeks. *Env. Res. Lett.*, 7, 034 015.
- ⁴¹⁶ Cess, R. D. and G. L. Potter, 1988: A methodology for understanding and intercomparing
 ⁴¹⁷ atmospheric climate feedback processes in general-circulation models. J. Geophys. Res.,
 ⁴¹⁸ 93, 8305–8314.
- ⁴¹⁹ Cess, R. D. *et. al.*, 1990: Intercomparison and interpretation of climate feedback processes
 ⁴²⁰ in 19 atmospheric general circulation models. *J. Geophys. Res.*, **95**, 16,601–16,615.

- ⁴²¹ Chadwick, R., P. Good, T. Andrews, and G. Martin, 2014: Surface warming patterns drive
 ⁴²² tropical rainfall pattern responses to CO₂ forcing on all timescales. *Geophys. Res. Lett.*,
 ⁴²³ 41, 610–615.
- Colman, R. and B. McAvaney, 2009: Climate feedbacks under a very broad range of forcing. *Geophys. Res. Lett.*, 36 (1), L01702.
- Colman, R. A. and B. J. McAvaney, 2011: On tropospheric adjustment to forcing and climate
 feedbacks. *Climate Dynamics*, 36 (9-10), 1649–1658.
- 428 Crucifix, M., 2006: Does the last glacial maximum constrain climate sensitivity? *Geophys.*429 *Res. Lett.*, **33**, L18 701.
- ⁴³⁰ Dickinson, R. E. and K. J. Schaudt, 1998: Analysis of timescales of response of a simple
 ⁴³¹ climate model. J. Climate, 11, 97–106.
- ⁴³² Dong, B. W., J. M. Gregory, and R. T. Sutton, 2009: Understanding land-sea warming con⁴³³ trast in response to increasing greenhouse gases. part i: Transient adjustment. J. Climate,
 ⁴³⁴ 22, 3079–3097.
- ⁴³⁵ Doutriaux-Boucher, M., M. J. Webb, J. M. Gregory, and O. Boucher, 2009: Carbon dioxide
 ⁴³⁶ induced stomatal closure increases radiative forcing via a rapid reduction in low cloud.
 ⁴³⁷ *Geophys. Res. Lett.*, **36**, L02 703.
- Fels, S. B., J. D. Mahlman, M. D. Schwarzkopf, and R. W. Sinclair, 1980: Stratospheric sensitivity to perturbations in ozone and carbon-dioxide—radiative and dynamical response.
 J. Atmos. Sci., 37, 2265–2297.
- Forster, P. M., T. Andrews, P. Good, J. M. Gregory, L. S. Jackson, and M. Zelinka, 2013:
 Evaluating adjusted forcing and model spread for historical and future scenarios in the
 CMIP5 generation of climate models. J. Geophys. Res., doi:10.1002/jgrd.50174.

- Gregory, J. and M. Webb, 2008: Tropospheric adjustment induces a cloud component in CO₂ forcing. J. Climate, **21**, 58–71.
- Gregory, J. M., C. D. Jones, P. Cadule, and P. Friedlingstein, 2009: Quantifying carbon
 cycle feedbacks. J. Climate, 22, 5232–5250.
- Gregory, J. M., et al., 2004: A new method for diagnosing radiative forcing and climate
 sensitivity. *Geophys. Res. Lett.*, **31**, L03 205.
- Hansen, J., M. Sato, and R. Ruedy, 1997: Radiative forcing and climate response. J. Geophys. *Res.*, 102, 6831–6864.
- 452 Hansen, J., et al., 1984: Climate sensitivity: Analysis of feedback mechanisms. *Climate*
- Processes and Climate Sensitivity, J. E. Hansen and T. Takahashi, Eds., American Geo physical Union, AGU Geophysical Monograph, Vol. 5, 130–163.
- Hansen, J., et al., 2005: Efficacy of climate forcings. J. Geophys. Res., 110, D18104.
- ⁴⁵⁶ Huntingford, C. and P. M. Cox, 2000: An analogue model to derive additional climate change
 ⁴⁵⁷ scenarios from existing GCM simulations. *Clim. Dyn.*, **16**, 575–586.
- ⁴⁵⁸ IPCC, 1990: Climate Change: the IPCC Scientific Assessment. Cambridge University Press,
- ⁴⁵⁹ report by the Intergovernmental Panel on Climate Change.
- Jarvis, A. and S. Li, 2011: The contribution of timescales to the temperature response of climate models. *Clim. Dyn.*, **36**, 523–531.
- Kamae, Y. and M. Watanabe, 2012a: On the robustness of tropospheric adjustment in
 CMIP5 models. *Geophys. Res. Lett.*, **39**, L23808.
- Kamae, Y. and M. Watanabe, 2012b: Tropospheric adjustment to increasing CO₂: its
 timescale and the role of land-sea contrast. *Clim Dyn*, 41 (11-12), 3007–3024.

- Lau, K. M., M. K. Kim, and K. M. Kim, 2006: Asian summer monsoon anomalies induced
 by aerosol direct forcing: the role of the Tibetan Plateau. *Clim. Dyn.*, 26, 855–864.
- Liu, J., B. Wang, M. A. Cane, S. Y. Yim, and J. Y. Lee, 2013: Divergent global precipitation changes induced by natural versus anthropogenic forcing. *Nature*, **493**, 656–659.
- 470 Lohmann, U., et al., 2010: Total aerosol effect: radiative forcing or radiative flux perturba471 tion? Atm. Chem. Phys., 10, 3235–3246.
- Mercado, L. M., N. Bellouin, S. Sitch, O. Boucher, C. Huntingford, M. Wild, and P. M. Cox,
 2009: Impact of changes in diffuse radiation on the global land carbon sink. *Nature*, 458,
 1014–U87.
- 475 Otto, A., et al., 2013: Energy budget constraints on climate response. Nat. Geosci., 6,
 476 415–416.
- ⁴⁷⁷ Raes, F., H. Liao, W. T. Chen, and J. H. Seinfeld, 2010: Atmospheric chemistry-climate
 ⁴⁷⁸ feedbacks. J. Geophys. Res., 115, D12 121.
- Ramanathan, V., et al., 2005: Atmospheric brown clouds: Impacts on South Asian climate
 and hydrological cycle. *Proc. Nat. Acad. Sci. USA*, **102**, 5326–5333.
- ⁴⁸¹ Rotstayn, L. D. and U. Lohmann, 2002: Tropical rainfall trends and the indirect aerosol
 ⁴⁸² effect. J. Climate, 15, 2103–2116.
- Rotstayn, L. D. and J. E. Penner, 2001: Indirect aerosol forcing, quasi forcing, and climate
 response. J. Climate, 14, 2960–2975.
- Schlesinger, M. E. and J. F. B. Mitchell, 1987: Climate model simulations of the equilibrium
 climatic response to increased carbon-dioxide. *Rev. Geophyss*, 25, 760–798.
- 487 Schmidt, H., et al., 2012: Solar irradiance reduction to counteract radiative forcing from a
- quadrupling of CO₂: climate responses simulated by four Earth system models. *Earth Sys.* Dyn., 3, 63-78.

- Senior, C. A. and J. F. B. Mitchell, 2000: The time-dependence of climate sensitivity. *Geo- phys. Res. Lett.*, 27, 2685–2688.
- Seo, J., S. M. Kang, and D. M. W. Frierson, 2014: Sensitivity of intertropical convergence
 zone movement to the latitudinal position of thermal forcing. *JClim*, 27, 3035–3042.
- Shine, K. P., J. Cook, E. J. Highwood, and M. M. Joshi, 2003: An alternative to radiative
 forcing for estimating the relative importance of climate change mechanisms. *Geophys. Res. Lett.*, **30**, 2047.
- ⁴⁹⁷ Soden, B. J. and I. M. Held, 2006: An assessment of climate feedbacks in coupled ocean⁴⁹⁸ atmosphere models. J. Climate, 19, 3354–3360.
- Stevens, B. and S. E. Schwartz, 2012: Observing and modeling earth's energy flows. Surveys
 Geophys., 33, 779–816.
- Tao, W. K., J. P. Chen, Z. Q. Li, C. Wang, and C. D. Zhang, 2012: Impact of aerosols on convective clouds and precipitation. *Rev. Geophys.*, **50**, RG2001.
- Tuttle, J. D. and R. E. Carbone, 2011: Inferences of weekly cycles in summertime rainfall. *J. Geophys. Res.*, 116, D20213.
- Vial, J., J.-L. Dufresne, and S. Bony, 2013: On the interpretation of inter-model spread in
 CMIP5 climate sensitivity estimates. *Clim. Dyn.*, 41, 3339–3362.
- ⁵⁰⁷ Wang, C., 2013: Impact of anthropogenic absorbing aerosols on clouds and precipitation: A
 ⁵⁰⁸ review of recent progresses. Atm. Res., **122**, 237–249.
- Watanabe, M., H. Shiogama, M. Yoshimori, T. Ogura, T. Yokohata, H. Okamoto, S. Emori,
 and M. Kimoto, 2012: Fast and slow timescales in the tropical low-cloud response to
 increasing CO₂ in two climate models. *Clim. Dyn.*, **39**, 1627–1641.
- ⁵¹² Webb, M. J., F. H. Lambert, and J. M. Gregory, 2013: Origins of differences in climate ⁵¹³ sensitivity, forcing and feedback in climate models. *Clim. Dyn.*, **40**, 677–707.

- ⁵¹⁴ Wyant, M. C., C. S. Bretherton, P. N. Blossey, and M. Khairoutdinov, 2012: Fast cloud
 ⁵¹⁵ adjustment to increasing CO₂ in a superparameterized climate model. J. Adv. Model.
 ⁵¹⁶ Earth Systems, 4, M05 001.
- ⁵¹⁷ Zelinka, M., S. Klein, K. Taylor, T. Andrews, M. Webb, J. Gregory, and P. Forster, 2013:
- ⁵¹⁸ Contributions of different cloud types to feedbacks and rapid adjustments in CMIP5. J.
- ⁵¹⁹ *Climate*, **26**, 50075027.

520 List of Figures

1

521

522

523

524

525

526

527

528

529

ods of diagnosing them. (a) Full system, with shortwave albedo effects in upper part and longwave in lower part. Traditionally-defined forcing occurs via green arrow (in the case of solar forcing) or red arrows (other forcings) from perturbation to the top-of-atmosphere energy imbalance \overline{N} . Adjustments also occur via red arrows. Feedbacks occur via blue arrows, with the Planck response shown by the direct arrow from $d\overline{T}$ to \overline{N} . Feedbacks and adjustments can be diagnosed simultaneously by the regression method. (b) Traditional view of Planck system with no adjustments (nor feedbacks). (c-d)

Diagram showing forcing-feedback concepts for global temperature and meth-

Reduced atmosphere-only system with fixed SST. Adjustments can be diagnosed by observing change in \overline{N} after applying a perturbation with SST fixed (c); feedbacks can be diagnosed by observing changes in \overline{N} after changing the SST with no (other) perturbation (d).

25

26

2Several examples of forcing adjustment mechanisms. (a) Solar, aerosol and 534 greenhouse gas perturbations each can cause horizontal variations (red/blue 535 regions) in net radiative heating of the atmosphere, which can drive circula-536 tions that alter cloud cover regionally and possibly change the global-mean ra-537 diative effect of clouds, modifying the conventional radiative forcings of these 538 perturbations. (b) These perturbations can also cause vertical variations in 539 the heating rate, altering atmospheric stratification, and affecting convection 540 and local cloud development. (c) Perturbations may affect land and ocean 541 surfaces differently, further affecting cloud cover. (d) CO_2 and aerosol pertur-542 bations can increase the growth of plants (affecting land albedo), or increase 543 their water-use efficiency, affecting fluxes of water vapor (yellow arrow) and 544 ultimately cloud cover. 545

3 Response of zonal-mean cloud cover over oceans to (a) a quadrupling of CO_2 546 (with warming effects removed), and (b) warming (with CO_2 effects removed), 547 averaged over several GCMs, determined by regression method. Because 548 cloudiness is generally reduced in both cases, these responses produce in-549 creased effective radiative forcing and positive cloud feedback, respectively, 550 although the details of the cloud changes vary among models and can be seen 551 here to differ significantly between the adjustment and feedback responses. 552 Reproduced from Zelinka et al. (2013). 553

27

28

Stratospherically-adjusted RF and ERF estimates by regression and fixed-4 554 SST methods using instantaneous $4 \times CO_2$ experiments, for a typical CMIP5 555 climate model. N is the net radiative flux imbalance at the top of atmosphere 556 and $d\overline{T}$ the global mean surface temperature change. The green cross gives 557 the stratospherically-adjusted radiative forcing of CO_2 quadrupling, 7.1 W 558 m^{-2} in this model. This is estimated as the instantaneous net downward 559 radiative flux change at the tropopause when CO_2 is quadrupled. The black 560 diamonds are annual means of the differences between the first 150 years of 561 step- $4 \times CO_2$ and control simulations. The blue line is the regression fit to 562 these points. Its $d\overline{T} = 0$ intercept (blue cross) is the Gregory estimate of 563 ERF (6.5 W m⁻² in this model). The red cross is a 30 year mean difference 564 of a pair of fixed sea surface temperature runs, one with standard CO_2 and 565 one with quadrupled CO_2 . To make it consistent with our basic definition, it 566 needs to be adjusted to $d\overline{T} = 0$ by adding 0.5 W m⁻² thereby removing the 567 feedback contribution (dashed red line), giving a fixed-SST ERF estimate of 568 7.0 W m^{-2} for this model. Adapted from Fig. 7.2 of Boucher et al. (2013). 569

570	5	Rapid development of the CO_2 -induced precipitation response (units mm
571		$\mathrm{day}^{-1})$ as simulated by the ECMWF-IFS (Integrated Forecast System) model
572		for October 2011 upon instantaneous quadrupling of CO_2 . As CO2 increases,
573		the reduction of the atmospheric radiative cooling warms the troposphere rel-
574		ative to the ocean surface, and warms the surface relative to the atmosphere
575		over land. This adjustment response, which takes place within a few days, re-
576		duces precipitation over ocean but enhances it over most land areas. Adapted
577		from Bony et al. (2013).
578	6	Adjustment of precipitation to a quadrupling of CO_2 estimated two ways,
579		using a single climate model (IPSL-CM5A). (a) Difference between first year
580		after quadrupling and control climatology, averaged over an ensemble of 12
581		realizations (this result, which is similar to that obtained by regression method

29

30

582on the ensemble mean time series, shows strong regional influences in the583Tropics from SST changes). (b) Change with SST held fixed everywhere,

averaged over 12 years. Note scale has double the range of that in Fig. 5.



FIG. 1. Diagram showing forcing-feedback concepts for global temperature and methods of diagnosing them. (a) Full system, with shortwave albedo effects in upper part and longwave in lower part. Traditionally-defined forcing occurs via green arrow (in the case of solar forcing) or red arrows (other forcings) from perturbation to the top-of-atmosphere energy imbalance \overline{N} . Adjustments also occur via red arrows. Feedbacks occur via blue arrows, with the Planck response shown by the direct arrow from $d\overline{T}$ to \overline{N} . Feedbacks and adjustments can be diagnosed simultaneously by the regression method. (b) Traditional view of Planck system with no adjustments (nor feedbacks). (c-d) Reduced atmosphere-only system with fixed SST. Adjustments can be diagnosed by observing change in \overline{N} after applying a perturbation with SST fixed (c); feedbacks can be diagnosed by observing changes in \overline{N} after changing the SST with no (other) perturbation (d).



FIG. 2. Several examples of forcing adjustment mechanisms. (a) Solar, aerosol and greenhouse gas perturbations each can cause horizontal variations (red/blue regions) in net radiative heating of the atmosphere, which can drive circulations that alter cloud cover regionally and possibly change the global-mean radiative effect of clouds, modifying the conventional radiative forcings of these perturbations. (b) These perturbations can also cause vertical variations in the heating rate, altering atmospheric stratification, and affecting convection and local cloud development. (c) Perturbations may affect land and ocean surfaces differently, further affecting cloud cover. (d) CO_2 and aerosol perturbations can increase the growth of plants (affecting land albedo), or increase their water-use efficiency, affecting fluxes of water vapor (yellow arrow) and ultimately cloud cover.



FIG. 3. Response of zonal-mean cloud cover over oceans to (a) a quadrupling of CO_2 (with warming effects removed), and (b) warming (with CO_2 effects removed), averaged over several GCMs, determined by regression method. Because cloudiness is generally reduced in both cases, these responses produce increased effective radiative forcing and positive cloud feedback, respectively, although the details of the cloud changes vary among models and can be seen here to differ significantly between the adjustment and feedback responses. Reproduced from Zelinka et al. (2013).



FIG. 4. Stratospherically-adjusted RF and ERF estimates by regression and fixed-SST methods using instantaneous $4 \times CO_2$ experiments, for a typical CMIP5 climate model. N is the net radiative flux imbalance at the top of atmosphere and $d\overline{T}$ the global mean surface temperature change. The green cross gives the stratospherically-adjusted radiative forcing of CO₂ quadrupling, 7.1 W m⁻² in this model. This is estimated as the instantaneous net downward radiative flux change at the tropopause when CO₂ is quadrupled. The black diamonds are annual means of the differences between the first 150 years of step- $4 \times CO_2$ and control simulations. The blue line is the regression fit to these points. Its $d\overline{T} = 0$ intercept (blue cross) is the Gregory estimate of ERF (6.5 W m⁻² in this model). The red cross is a 30 year mean difference of a pair of fixed sea surface temperature runs, one with standard CO₂ and one with quadrupled CO₂. To make it consistent with our basic definition, it needs to be adjusted to $d\overline{T} = 0$ by adding 0.5 W m⁻² thereby removing the feedback contribution (dashed red line), giving a fixed-SST ERF estimate of 7.0 W m⁻² for this model. Adapted from Fig. 7.2 of Boucher et al. (2013).



FIG. 5. Rapid development of the CO_2 -induced precipitation response (units mm day⁻¹) as simulated by the ECMWF-IFS (Integrated Forecast System) model for October 2011 upon instantaneous quadrupling of CO_2 . As CO2 increases, the reduction of the atmospheric radiative cooling warms the troposphere relative to the ocean surface, and warms the surface relative to the atmosphere over land. This adjustment response, which takes place within a few days, reduces precipitation over ocean but enhances it over most land areas. Adapted from Bony et al. (2013).



FIG. 6. Adjustment of precipitation to a quadrupling of CO_2 estimated two ways, using a single climate model (IPSL-CM5A). (a) Difference between first year after quadrupling and control climatology, averaged over an ensemble of 12 realizations (this result, which is similar to that obtained by regression method on the ensemble mean time series, shows strong regional influences in the Tropics from SST changes). (b) Change with SST held fixed everywhere, averaged over 12 years. Note scale has double the range of that in Fig. 5.

Response to comments of Reviewer #1.

We have addressed all minor editorial issues raised; below are the more substantive comments with our responses.

Below are some comments on specific wording and other details. In addition, I hope that the authors go over the paper again and identify the main points, express them more clearly, and summarize them in the conclusions.

We have tried to do this in the revision, thanks for the suggestion.

First of all, the title is misleading. It could easily be interpreted to mean "adjusting the framework". I suggest changing it to : "The role of (rapid) adjustments in the forcing-feedback framework"

We have chosen to keep the current title, which we do not think is misleading and was already the result of much discussion among the authors.

19-21 The last sentence includes concepts that are only marginally part of the paper. As such, I am not sure they merit being included in the abstract.

We have considered this but do not think they are marginal, and hope they come out a bit more strongly in the revision.

14 "which are direct responses to forcings that are not mediated by surface temperature changes" - global mean temperature is in the feedback definition, and used to calculate feedbacks, but arguably, feedbacks in the real climate system are mediated by temperature changes on a veriey of spatial scales.

We have tried to clarify this point in the revision. Clearly temperature changes heterogeneously in the real world; the question is whether changes specifically associated with that heterogeneity (and orthogonal to the mean) count as feedback responses or as direct consequences of the forcing. This Is somewhat ambiguous but there are principles that apply, as articulated in the revision.

35 surface snow cover - Why not mention sea ice? It's a stronger, more straightforward, feedback mechanism because the albedo change is larger.

Agreed, done

37 ".. called the "climate sensitivity". "Climate sensitivity parameter" is not a term used widely in the literature. I believe that in particular for the purpose of making the information contained in this paper accessible to a broad audience, it is best not to introduce terminology that is not standard.

We believe clarity is needed on this. The terminology is standard and our use is consistent with e.g. the IPCC AR5.

41-42 Rewrite this sentence. It sounds out of context, especially since the "new approaches" have not been specifically mentioned yet. In the preceding sentence, you could reference Figure 1b. Write a sentence or two to describe and reference the other parts of the Figure as well.

We now postpone reference to "new approaches" and provide pointers to this figure later in the text (middle of "Why Adjustments?" and beginning of "Ways of Defining and Calculating Adjustments").

123 Again, consider replacing global-mean temperature with surface temperature.

No, it is crucial that this be the global-mean temperature for the framework to be logically consistent. This relates to an earlier point that we hope we have clarified better.

131 You provide some specific examples for slow adjustment processes, an example or two of fast feedback processes relevant here would be helpful.

Done.

165 The subtitle "How should rapid adjustments be defined and diagnosed?" leads the reader to expect specific guidelines for calculating these quantities. However, the section does not provide this information, but rather a discussion of methods to perform these calculations and their shortcomings. Consider changing either the title or the contents of this section.

We have changed the subtitle.

171 What is the definition of a "transient simulation" you are using here? I assume that you are referring to a simulation where a forcing was applied

instantaneously in the beginning of the run, and the simulation traces the climate's response over time. However, "transient" is often also used when considering transient forcings that evolve with time. Be specific. It is my understanding, that the Gregory method is best applied to equilibrium, rather than transient (in the second sense of the term) model simulations. But I may be wrong.

No, it is not essential that the forcing be applied instantaneously here, only that the simulation be transient in the usual sense: one where the system is out of equilibrium. We have ensured that all uses of the word "transient" consistently mean this and that if we specifically mean abrupt forcing application we say so.

238-253 This may be personal preference, but I would refrain from invoking the Gaia hypothesis, even if it is just by referencing Daisy world. Aside from being controversial, it is not a straightforward enough analogy to be useful to a broad audience.

We have eliminated this.

The last sentences of the paper should reiterate your main points. The current ending is rather weak. Try and end on a strong note.

Good advice, we have tried to do this.

Figure 3 - I understand this is taken from a different paper. It would still be useful, if the raw data is available to the authors, to plot the cloud changes using the same scale. This will highlight not only the spatial differences, but also the differences in magnitude. The Figure suggests the cloud changes are of roughly the same magnitude, but different distribution, unless one carefully looks at the scale.

The two panels have different units so it is not possible to put them on the same scale. The more important effect in most models when appropriately scaled is the temperature-induced one, which does appear bolder as drafted by Zelinka et al., so in this respect the figure is in no way misleading.

Figure 4 - Please remove the negative values of N from this Figure. The y-Axis can just as easily go from 0 to 8 without any loss of information. Removing the dashed 0 line however, will make the figure look cleaner. Also make sure the tick marks on the x-Axis are spaced the same as for the y-Ax.

Thanks, done.

Responses to Reviewer #2

I still wonder, though, if this message could have been formulated simpler. Maybe because I consider myself a more visual person, I like conceptional drawings (like Figure 1). But I struggled at times with complicated content of the text, especially when sentences where seemingly unnecessarily elongated.

We have tried to simplify the language in those places that may have caused problems for the reviewer.

minor comments

Figure 1

Nice conceptional illustration. Sill N is not explained (apparently the net input to the system from the text). I wonder if the direct arrow from the perturbation is needed in a) and c) as N is modified by pl.albedo and IR greenhouse? Should not also the link from N to dT in a) be left out?

No the link is needed to represent solar forcing. This is now explained in the caption, as is N.

Figure 2

Please add "uneven distributed" after a), processes for c) and d) are not that easily visually understood as a) or b) cane we improve?

Done w/respect to wording. No changes made to figure.

Figure 3

Both a) and b) have 'warming effects removed". If this correct, what is the case difference?

No one is warming-removed and one CO2-removed. Wording altered to reduce the chance of confusion here.

The (rapid) adjustment vs feedback is an interesting approach. I am not sure, however, that it makes it easier for aerosol. An adjustment component for aerosol is more difficult to quantify then a direct (radiative) effect (when separating oldfashioned in aerosol direct and the sum of all aerosol indirect effects). In order to include the semi-direct effect complicated modeling is required, which in part depends on less than perfect cloud parameterizations in modeling.

The framework is already being used for aerosols, all we have done is to point out the parallels with CO2 and proposed a clarification of the framework. The semi-direct effect does indeed require complicated modeling, but is automatically part of a GCM experiment with aerosols so it cannot be avoided, and is no more complicated than the adjustments to other forcings which also involve clouds.

I also do not think that the CCN/IN impact is not also affected by the semi-direct effect so any split will be artificially (as also the authors admit that timescales of adjustments and feedbacks can be 'comparable')

We have not split them, semi-direct effects associated with CCN are all part of the adjustment. Not sure what change the reviewer wants here.

Agreed, similar to the direct effect also the Twomey effect is one of many effect, which has been picked, because the ability to observe and easily simulate. I agree that "only the combined effect" matters ... but what if such simulated effects are derived with models having deficiencies in cloud-representation and convection? (shouldn't we stick to something that can be understood?)

This, and the above comments indicate a philosophical difference. Such effects are already being examined, and models indicate they are sometimes of significant magnitude. We therefore need a framework for understanding them. The fact that they are uncertain is no reason to ignore them, when they are already present in comprehensive models and (implicitly) in observations.

Gregory's method is based on global averages. Can we apply such an averaging linear concept for uneven distributed responses? (... well the next page talks about this).

It is now stated more explicitly that the regression method may not be optimal for regional responses but can be used with appropriate ensemble averaging.

Can we be sure that adding absorbing aerosol decreases global precipitation? Any reference?

Reference now given.

Why is response in Figure 6 (twice the scale but similar colors) so much larger than in Figure 5?

First it is two different models; second, one is showing the response after only a few days while the other is the total response.

The last sentence (which I like) is a nice summary of what this contribution intends (... although the given examples for aerosol and precipitation come across as complex, so that better detection and attribution will remain an uphill battle).

Thanks we have moved this toward the end.

Copyright Form Click here to download Copyright Form: BAMS all copyright forms.pdf