

# *Climate change and water in the UK: past changes and future prospects*

Article

Published Version

Creative Commons: Attribution 3.0 (CC-BY)

Open Access

Watts, G., Battarbee, R. W., Bloomfield, J. P., Crossman, J., Daccache, A., Durance, I., Elliott, J. A., Garner, G., Hannaford, J., Hannah, D. M., Hess, T., Jackson, C. R., Kay, A. L., Kernan, M., Knox, J., Mackay, J., Monteith, D. T., Ormerod, S. J., Rance, J., Stuart, M. E., Wade, A. J., Wade, S. D., Weatherhead, K., Whitehead, P. G. and Wilby, R. L. (2015) Climate change and water in the UK: past changes and future prospects. *Progress in Physical Geography*, 39 (1). pp. 6-28. ISSN 1477-0296 doi: <https://doi.org/10.1177/0309133314542957> Available at <http://centaur.reading.ac.uk/39254/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1177/0309133314542957>

Publisher: Sage

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online



# Climate change and water in the UK – past changes and future prospects

Progress in Physical Geography  
2015, Vol. 39(1) 6–28  
© The Author(s) 2015  
Reprints and permission:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/0309133314542957  
ppg.sagepub.com



## **Glenn Watts**

Environment Agency, UK

## **Richard W. Battarbee**

University College London, UK

## **John P. Bloomfield**

British Geological Survey, UK

## **Jill Crossman**

University of Oxford, UK

## **Andre Daccache**

Cranfield University, UK

## **Isabelle Durance**

Cardiff University, UK

## **J. Alex Elliott**

Centre for Ecology and Hydrology, UK

## **Grace Garner**

University of Birmingham, UK

## **Jamie Hannaford**

Centre for Ecology and Hydrology, UK

## **David M. Hannah**

University of Birmingham, UK

## **Tim Hess**

Cranfield University, UK

## **Christopher R. Jackson**

British Geological Survey, UK

## **Alison L. Kay**

Centre for Ecology and Hydrology, UK

## **Martin Kernan**

University College London, UK

## **Jerry Knox**

Cranfield University, UK

## **Jonathan Mackay**

British Geological Survey, UK

## **Don T. Monteith**

Centre for Ecology and Hydrology, UK

## **Steve J. Ormerod**

Cardiff University, UK

## **Jemima Rance**

HR Wallingford, UK

## **Marianne E. Stuart**

British Geological Survey, UK

## **Andrew J. Wade**

University of Reading, UK

## **Steven D. Wade**

Met Office, UK

## **Keith Weatherhead**

Cranfield University, UK

## **Paul G. Whitehead**

University of Oxford, UK

## **Robert L. Wilby**

Loughborough University, UK

---

### **Corresponding author:**

Glenn Watts, Environment Agency, Horizon House, Deanery Road, Bristol BS1 5AH, UK.

Email: glenn.watts@environment-agency.gov.uk

## Abstract

Climate change is expected to modify rainfall, temperature and catchment hydrological responses across the world, and adapting to these water-related changes is a pressing challenge. This paper reviews the impact of anthropogenic climate change on water in the UK and looks at projections of future change. The natural variability of the UK climate makes change hard to detect; only historical increases in air temperature can be attributed to anthropogenic climate forcing, but over the last 50 years more winter rainfall has been falling in intense events. Future changes in rainfall and evapotranspiration could lead to changed flow regimes and impacts on water quality, aquatic ecosystems and water availability. Summer flows may decrease on average, but floods may become larger and more frequent. River and lake water quality may decline as a result of higher water temperatures, lower river flows and increased algal blooms in summer, and because of higher flows in the winter. In communicating this important work, researchers should pay particular attention to explaining confidence and uncertainty clearly. Much of the relevant research is either global or highly localized: decision-makers would benefit from more studies that address water and climate change at a spatial and temporal scale appropriate for the decisions they make.

## Keywords

adaptation, climate change, climate change impacts, decision-making, freshwater ecosystems, hydrological change, water environment, water quality change

## 1 Introduction

Observed change in the climate over recent decades has been linked with changes in the global hydrological cycle, including increased atmospheric water content and changing precipitation patterns (Allan, 2011; Bates et al., 2008). It has been suggested that many of the most severe global impacts of anthropogenic climate change may be mediated by water (Stern, 2006) and that rivers may be among the ecosystems most sensitive to climate change (Kernan et al., 2010; Millennium Ecosystem Assessment, 2005; Ormerod, 2009). It is also anticipated that further warming will intensify the hydrological cycle, leading globally to an increased risk of floods and droughts (Bates et al., 2008; Giorgi et al., 2011; Rockström et al., 2009) and increased exposure to water resources stress, even under the most stringent emissions mitigation scenarios (Arnell et al., 2011, 2013). According to the UK's 2012 Climate Change Risk Assessment (CCRA), three of the six potential risks requiring early action are water-related: flood and coastal erosion risk

management; specific aspects of natural ecosystems including the management of soils, water and biodiversity; and management of water resources (CCRA, 2012: 8). The Intergovernmental Panel on Climate Change (IPCC) is 'virtually certain' that global precipitation will increase with increased global temperature, and has 'high confidence' that the contrast between wet and dry regions and wet and dry seasons will increase over most of the world (IPCC, 2013b), with Europe at high risk from flooding and water shortages over the rest of the 21st century without further adaptation effort (IPCC, 2014).

Regionally, the impact of anthropogenic climate change on the past and future water cycle is less clear. Projections of climate change at a regional level show considerable variability, with disagreement even about the sign of change in variables such as precipitation in some regions and seasons (Bates et al., 2008). This is a particular problem for the UK, where weather patterns are strongly influenced by the North Atlantic Oscillation, storm tracks and blocking (Murphy et al., 2009), and where Global Climate Model (GCM) output needs to be

downscaled to reflect local and regional scale weather and climate processes (Fowler et al., 2007a), making the direct use of global or continental scale studies problematic.

It is at the regional level that most steps towards adaptation must be taken. To develop and test adaptation responses, policy-makers and decision-makers require a good understanding of the scale and scope of possible change and the uncertainties associated with these regional projections. Lack of information, or perhaps just a perception of a lack of information, seems to act as a barrier to adaptation action (Moser and Ekstrom, 2010). Even when decision-makers consciously choose approaches that favour flexible or low-regrets solutions, such as scenario-neutral approaches (Wilby and Dessai, 2010) or robust decision-making (Lempert et al., 2006), they need a plausible range of context-relevant futures against which they can test solutions (Adger et al., 2009).

This paper aims to synthesize the available research on the impact of anthropogenic climate change on the water environment in the UK in a way that will help decision-makers understand the scale and possible direction of change. We assume that decision-makers, for example in water companies, regulators and government, are best placed to evaluate the impact of climate change on their own area of work, but that to do this they need clear information about the scale and rate of change. As part of this synthesis, we identify areas where further research could reduce uncertainties or improve understanding in a way that would help decision-making. While we focus on the UK, we expect that the construction of this synthesis and the identification of gaps will be relevant to others planning research to aid decision-making.

This paper is organized as follows. Section II provides a summary of the evidence of changes to the UK water cycle and water environment, focusing mainly on the second half of the 20th century and the first decade of the 21st century. Section III looks at projected changes

through the rest of the 21st century. In Section IV we consider the implications for decision-making, identify research gaps and draw conclusions. To avoid any confusion, whenever in the paper we mention seasons, these are defined climatologically for the UK: winter is December, January and February; spring is March, April and May; summer is June, July and August; and autumn is September, October and November.

## **II Historical changes to the water cycle and water environment in the UK**

### *I Precipitation and evapotranspiration*

The England and Wales Precipitation (EWP) series includes monthly totals from January 1766 (Alexander and Jones, 2001), and has been shown to be suitable for use in a wide range of studies across the UK (Croxtton et al., 2006). Annual average rainfall has not changed significantly through this series, but there is an increasing trend in winter rainfall, although with little change over the last 50 years (Jenkins et al., 2008). More winter rain is falling in intense events (Burt and Ferranti, 2011; Jenkins et al., 2008; Jones et al., 2012; Osborn and Hulme, 2002), with the changes being most significant in long-duration events (5–10 days) (Fowler and Kilsby, 2003). Long summer events also show increased rainfall intensity (Jones et al., 2012). There is also some evidence that the within-year clustering of extreme rainfall events has increased in recent years (Jones et al., 2012). There is insufficient evidence to suggest a link between anthropogenic climate change and these changes in precipitation; it has been suggested that at the UK scale such a link may not be apparent until the 2050s in most regions (Fowler and Wilby, 2010).

Evapotranspiration includes both direct evaporation to the atmosphere from soil and water

surfaces, and transpiration from trees and other plants. Potential evapotranspiration (PE) is an estimate of the maximum volume of water that could pass to the atmosphere if there is no limit to supply. Actual evapotranspiration (AE) is constrained by water availability (usually soil moisture), and cannot be greater than PE. Evapotranspiration is a function of the energy balance, humidity and wind speed (Allen et al., 1998; Monteith, 1965; Penman, 1948).

There are few studies of historical changes in evapotranspiration in the UK (Kay et al., 2013a). Burt and Shahgedanova (1998) calculate evaporation from 1815 to 1996 for a site in Oxford, finding increases in PE but decreases in AE. Kay et al. (2013a) find some signs of increasing PE since the 1960s across the UK. Temperature trends can be used to infer changes in PE as the two tend to be correlated (see, for example, Dai, 2011, and Sheffield et al., 2012, for a discussion of the impact of different PE formulations on drought indices), though it is not clear that the relationship will remain constant in a changing climate. Karoly and Stott (2006) conclude that it is likely that there has been a human contribution to warming in the Central England Temperature (CET) series. CET has increased by about a degree Celsius since 1980 (Jenkins et al., 2008). On this basis, it might be reasonable to hypothesize that UK PE has also increased over the same period, but there is no formal study to confirm this.

## 2 River flows and groundwater levels

The UK has an exceptionally dense river gauging and groundwater level network, essential for the detection of changes and trends in river flow and groundwater level (Hannah et al., 2011). However, there are few records from before the 1960s, catchment change and water use affects many of the sites, the uplands are under-represented, and poor-quality and missing data is a problem for the estimation of low

and high extremes of both river flow and groundwater level.

Groundwater levels are highly variable in both space and time, and there have been no studies of historical changes in groundwater levels across the UK. No links have yet been made between UK groundwater levels and anthropogenic climate change (Jackson et al., 2014).

There are more large-scale studies of changes in UK river flow, mainly concentrating on the 'benchmark network' of sites where the net impact of human disturbance on flow regimes has been relatively minor (Bradford and Marsh, 2003). Hannaford and Buys (2012) find a high degree of spatial variability in seasonal flow trends from 1969 to 2008. Winter flows have increased in upland, western areas and autumn flows have increased across much of the UK. There is a weak trend towards decreasing spring flows since 1960, particularly in lowland England, and no clear pattern in summer.

A considerable effort has been devoted to detecting changes in flooding. Increases in high flow magnitude and duration have been found from the 1960s to the 2000s, especially in the north and west (Hannaford, 2014). However, there is little compelling evidence for any long-term increase in flooding, but pronounced interdecadal variability, with notable 'flood rich' and 'flood poor' periods in long hydrological records (Hannaford, 2014; Wilby and Quinn, 2013). In contrast, less work has addressed changes in low flows or drought: studies have generally been inconclusive, with marked spatial variations and high sensitivity to the period of study (Hannaford, 2014). As with floods, there is pronounced interdecadal variability in long-term records of low flow or drought, and Marsh et al. (2007) demonstrate that some of the droughts of the 19th century were longer and more severe than those of the 20th century.

These changes in flow have not been attributed to anthropogenic climate change (Hannaford, 2014). The only explicit flow attribution published is for the unusual flood event of

autumn 2000 (Kay et al., 2011; Pall et al., 2011) which is thought to have been more likely as a result of greenhouse gas forcing, though even here the two different studies offer different assessments of the increased probability attributed to the forcing.

### **3 River, lake and groundwater temperature, water quality and freshwater ecosystems**

Along with flow, water temperature is one of the most important influences on ecosystem state in surface waters (Caissie, 2006; Webb et al., 2008). Water temperature exerts a direct control on the metabolisms and lifecycles of aquatic organisms and affects water quality, with most chemical and bacteriological processes operating faster at higher temperatures, leading, for example, to increased occurrence of algal blooms (Whitehead et al., 2009). River water temperature is controlled by energy and hydrological fluxes at the air-water and water-riverbed interfaces (Hannah et al., 2008). River water temperature varies seasonally and diurnally in response to the dominant solar forcing, with day-to-day thermal variability in response to weather conditions and flow change. Lake surface water temperature is closely related to air temperature, with the nature of the relationship changed by lake area, volume, exposure to convective cooling by winds and, in summer, thermal stratification (George et al., 2007).

For the UK, there are very few studies of long-term changes in water temperature, though there is some sign of increasing water temperatures since the 1970s or 1980s in rivers and lakes (Durance and Ormerod, 2007; Foley et al., 2012; Hannah and Garner, 2014; Langan et al., 2001; Orr et al., 2014; Webb and Walling, 1992). There is very little information about changes in groundwater temperature in the UK.

River water quality is influenced by many factors including water temperature, hydrological regime, nutrient status, point source and

diffuse discharges, mobilization of toxic substances and acidification potential (Whitehead et al., 2009), making understanding and attributing change difficult. Pollution from point source discharges and toxic substances has decreased in recent years, mainly as a result of tighter regulation and the introduction of tertiary treatment at sewage treatment works (Jarvie et al., 2006; Neal et al., 2010), but there is a long history of increasing nutrient levels in UK catchments (Whitehead et al., 2009), mainly related to changes in land-use and fertilizer application (Bennion et al., 2012; Whitehead, 1990). Upland catchments have begun to recover from acidification since the 1980s, as a result of reductions in anthropogenic atmospheric sulphur emissions (Battarbee et al., 2012; Monteith et al., 2012; Wright et al., 2005). Dissolved organic carbon (DOC) concentrations have doubled across the UK since the 1980s (Evans et al., 2001, 2005; Freeman et al., 2001; Monteith et al., 2000, 2001; Worrall et al., 2003, 2004), most likely in response to reductions in acid deposition (Monteith et al., 2007). Changes in UK river water quality have not been linked to anthropogenic climate change, with changes in the lowlands in particular being driven by changes in land-use and point sources (Howden et al., 2010).

The most significant impact on UK groundwater quality in the second half of the 20th century was the intensification of agriculture and the consequent impact on groundwater quality from diffuse pollution, in particular from nitrate (Shand et al., 2007). Such changes are thought to be greater than any direct effects from a change in climate (Stuart et al., 2011) and there have been no studies linking historical groundwater quality changes to change in climate.

Freshwaters are considered to be among the most sensitive of all ecosystems to the effects of climate change (Durance and Ormerod, 2007, 2009), partly because of the magnitude of possible changes in water temperature and

flow, but also because most river animals are ectotherms (their body temperatures are controlled by their surroundings). However, establishing long-term climate effects on freshwater organisms and ecological processes is difficult. Detecting change requires long, systematic records. There are also many confounding factors that affect biodiversity and ecosystem form and function; these include flow variation, improving or declining water quality, habitat degradation and invasive species. Even where climate change is the underlying cause of local extinction, this may not occur as a direct response to climate forcing but to other related factors that have responded to climate forcing, such as loss of host or pollinator species or changes in pathogens or competitors (Cahill et al., 2013). Clews et al. (2010) suggest that warmer, drier summers may explain reductions in salmon and brown trout populations in the Welsh Wye catchment between 1985 and 2004, and a decline in Arctic Charr *Salvelinus alpinus* and increase in invasive Roach *Rutilus rutilus* in Lake Windermere, Cumbria, has been related to increased water temperature (Jeppesen et al., 2012). Increased spates may destroy habitats and extended low flow conditions can cause both scouring and siltation (Meyer et al., 1999; Wright et al., 2004). Altered environmental conditions may result in the loss of taxa allowing non-native species to enter freshwater ecosystems (Verdonschot et al., 2010).

#### 4 Summary of impacts

Anthropogenic influences on global climate are clear (IPCC, 2013a), but the change in climate and its impacts in the UK are so far less apparent (Table 1). Air temperature increases have been attributed formally to anthropogenic climate change, and water temperature changes may be linked to the same processes that have changed air temperature, though water temperature is only partly dependent on air temperature. Changes in rainfall patterns have not been linked

to anthropogenic climate change, and the impact of climate change on river flows and groundwater levels has not yet been detected. This suggests that adaptation to climate change will need to start before formal attribution of changes; for example, Fowler and Wilby (2010) estimate that it will be at least a decade before climate change induced changes in winter extreme rainfall can be detected in south-west England, with later detection for other seasons and regions of the UK.

#### 5 Possible changes to the UK water cycle and water environment through the 21st century

In evaluating possible changes, it is essential to consider uncertainty in climate projections. Uncertainties in climate prediction come from three main sources: internal variability of the climate system, model uncertainty, and emissions uncertainty (Hawkins and Sutton, 2009). For many regions, including western Europe, natural climate variability is the biggest source of uncertainty for up to 30 years ahead, with greater uncertainty in precipitation projections than temperature projections (Hawkins and Sutton, 2011). Model uncertainty – the way that different GCMs simulate changes in climate for a given radiative forcing – then becomes the most important source of uncertainty (Hawkins and Sutton, 2011).

The UK has a rich and growing academic literature on the future impact of climate change, much of which covers some aspect of water or the water environment. UK-scale assessments of climate-change impacts have been conducted for river flows (Christiensen et al., 2012; Prudhomme et al., 2012), droughts (Blenkinsop and Fowler, 2007; Burke et al., 2010; Rahiz and New, 2013; Vidal and Wade, 2009), extreme rainfall (Fowler and Ekström, 2009; Fowler et al., 2007b) and floods (Kay et al., 2013b, 2013c).



## 6 Precipitation and evapotranspiration

The previous generation of UK climate projections, known as UKCIP02 (Jenkins et al., 2002), presented a relatively simple picture of increasingly warmer, wetter winters and hotter, drier summers through the 21st century, suggesting, for example, that by the 2080s virtually every summer may be hotter and drier than 2001. The UK Climate Projections 2009 (UKCP09; Murphy et al., 2009), on which many recent UK climate impact studies are based, generate a probabilistic ensemble of climate projections that combine climate variability and model structural uncertainty for three separate emissions scenarios (Murphy et al., 2008). The three emissions scenarios, called High, Medium and Low in UKCP09, correspond to the IPCC's A1FI, A1B and B1 scenarios (IPCC, 2000). UKCP09 allows an explicit understanding of the possible range of climate change, though the range remains constrained by the processes represented in the climate models used in the assessment.

UKCP09 shows little change in annual average rainfall by the 2080s, with the 10th to 90th percentile range from a 16% reduction to a 14% increase for the medium emissions scenario. Seasonal precipitation changes may be greater. In winter, UKCP09 projects the biggest increases to be along the west coast, with a median change of +33% (10th to 90th percentile range +9 to +70%). In contrast, small decreases are seen in Scotland in winter (10th to 90th percentile range -11 to +7%). In summer, UKCP09's biggest median change of about -40% is in southern England (10th to 90th percentile range -65 to -6%), with little change in northern Scotland (10th to 90th percentile range -8 to +10%). It will be noted that, by definition, there is a 20% chance of values being outside the ranges quoted. Fowler and Ekström (2009) examine changes in extreme rainfall using a range of Regional Climate Models (RCMs), finding increases in winter,

spring and autumn extreme precipitation by the 2080s, with increases ranging from 5 to 30% depending on region and season. Summer changes are less clear, and changes in short-duration (sub-daily) rainfall events remain unclear, as current climate models are unreliable at these scales (Fowler et al., 2007b).

UKCP09, in common with most RCMs and GCMs, does not offer direct estimates of future evapotranspiration. It is possible to use output from climate models to calculate PE using one of the many formulae that exist, but this can be problematical. The more physically based methods are considered most accurate (e.g. Allen et al., 1998) but GCMs do not project all climate parameters with equal reliability, raising the question of whether it is better to use physically based methods with uncertain data or empirical methods with more certain data (Kingston et al., 2009). It is clear that different methods lead to different projections of potential evapotranspiration (Kay et al., 2013a), at scales from local (Kay and Davies, 2008) to global (Kingston et al., 2009).

For the UK, gridded Penman-Monteith estimates generally show increases in PE by the 2050s (Prudhomme et al., 2012) and the 2080s (Kay and Davies, 2008). Patterns of increase vary seasonally: Kay and Davies (2008) find that some months can show small decreases in PE using the Penman-Monteith formulation, but annual changes range from +6 to +56%. Temperature-based PE estimates for the UK show a similar range (e.g. Kay and Jones, 2012) but tend towards increases throughout the year (e.g. Christensen et al., 2012; Kay and Davies, 2008).

## 7 River flows and groundwater levels

Much attention has been paid to possible changes in flow as a result of climate change. Most studies concentrate either on long-term changes in average flow or on changes in flood flow. A

**Table 1.** Observed changes to water and the water environment in the UK.

Area	Observed change	Confidence	Sources of information
Air temperature	Increase of about 1°C in Central England Temperature (CET) since 1980.	High confidence of increase; medium confidence of link to climate change.	Jenkins et al. (2008); Karoly and Stott (2006)
Precipitation	Little change in annual average rainfall over last 300 years.	High confidence of little change.	Jenkins et al. (2008)
	More winter rainfall falling in intense events over the last 30 years.	Medium confidence that more winter rainfall is falling in intense events. Insufficient evidence to link to climate change.	Burt and Ferranti (2011); Fowler and Kilsby (2003); Jenkins et al. (2008); Jones et al. (2012); Osborn and Hulme (2002)
	Increased clustering of within-year extreme rainfall events.	Medium confidence of increased clustering. Insufficient evidence to link to climate change.	Jones et al. (2012)
Evapotranspiration	No systematic studies of change; possible increases since 1960s across the UK.	Low confidence in increases. Insufficient evidence to link to climate change.	Kay et al. (2013a)
River flows	Increase in winter runoff and high flows over last 40 years.	Medium confidence in increases; insufficient evidence to link to climate change.	Hannaford (2014); Hannaford and Buys (2012); Werrity (2002)
	No apparent trend in summer flows, low flows or drought.	Low confidence in changes. Insufficient evidence to link to climate change.	Hannaford (2014); Hannaford and Buys (2012)
Groundwater levels and recharge	No systematic studies of change.	Insufficient evidence to link to climate change.	Jackson et al. (2014)
River water temperature	Increase in river water temperature over recent decades.	Medium confidence of increase; low confidence that this is a response to climate change.	Durance and Ormerod (2007); Hannah and Garner (2014); Orr et al. (2014); Stuart et al. (2011); Webb and Walling (1992); Webb et al. (2008)
Groundwater temperature	Little is known about changes in groundwater temperature.	Insufficient evidence to understand changes.	
River water quality	Improved river water quality over last 30 years, especially in the lowlands.	High confidence in improved quality in the lowlands; low confidence in any link to climate change.	Howden et al. (2010); Whitehead et al. (2009); Wright et al. (2005)

*(continued)*

**Table 1.** (continued)

Area	Observed change	Confidence	Sources of information
Groundwater quality	Increased diffuse pollution, especially from nitrates, over the second half of the 20th century.	High confidence in increased pollution; high confidence that this is not linked to climate change.	Bloomfield et al. (2006); Shand et al. (2007); Stuart et al. (2011)
Freshwater ecosystems	Some evidence of reductions in populations of invertebrates and fish that depend on cold conditions.	Medium confidence in changes; low confidence that these changes are linked to climate change.	Clews et al. (2010); Durance and Ormerod (2007)

Note: confidence statements are subjective and based on the quantity of information, its spatial coverage, and the level of agreement between different studies.

common approach is to scale historic rainfall and potential evapotranspiration time series, referred to as a perturbation or delta change approach (Fowler et al., 2007a; Prudhomme et al., 2002). This has the effect of maintaining current sequencing of wet and dry events but changing their magnitude. Identification of the scaling factors for a river catchment or other area of interest requires downscaling from a GCM, for which a variety of different methods exists, often giving different hydrological results (Fowler et al., 2007a). Combined with the uncertainty in hydrological modelling (e.g. New et al., 2007), this means that projections of changes in UK flow cover a wide range, though as the range of uncertainty is poorly sampled this is probably not the full range of possible outcomes (Arnell et al., 2013; Hall, 2008).

Two recent studies look at the impact of climate change on monthly and seasonal river flows across the UK. Christensen et al. (2012) use the UKCP09 climate ensemble to model flows in the 2020s at 70 locations across the UK, choosing sites where the catchment is thought to be broadly undisturbed by artificial influences (Bradford and Marsh, 2003). The ensemble approach allows Christensen et al. (2012) to draw conclusions about the range of possible changes in flow. The median

projection is a reduction in spring and summer flows, a mixed pattern in autumn, and small increases in winter flows across the UK. The greatest reductions in flow are in August, with a median projection of up to 30% reduction compared to the 1961–90 baseline. For most seasons the 25th to 75th percentile range spans a range from lower to higher flows; the exception is for the summer where most sites have a reduction in flow even at the 75th percentile. This distribution leads Christensen et al. (2012) to some confidence in a reduction in summer flows across the UK by the 2020s.

Prudhomme et al. (2012) use a different approach to examine changes in flow across Britain by the 2050s. Their model is a generalized rainfall-runoff model parameterized on catchment characteristics (Young, 2006) which means that it is possible to model natural flows at reach level across the country. Such a model needs spatially coherent climate projections; Prudhomme et al. (2012) use change factors derived from the 11-member perturbed physics ensemble of RCMs that form part of the basis of UKCP09, based on a single emissions scenario (UKCP09's Medium scenario). These provide realizations of possible change to which it is not possible to assign probabilities, but which give some sense of spatial

variability in change that cannot be gathered from the probabilistic approach of Christensen et al. (2012). Prudhomme et al. (2012) find a mixed picture of changes in winter flow in England and Wales, with changes of  $-20$  to  $+40\%$ . In Scotland the changes in winter are smaller, in the range of  $\pm 20\%$ . In spring most of the ensemble has reduced flows across the UK, with reductions of up to  $40\%$ , but three of the 11 RCMs show increases in flow of up to  $60\%$  in central England. In summer, there is a more consistent picture of reduced flows with reductions of up to  $80\%$  especially in the north and west. However, even in summer in some ensemble members there are small increases in flow in some areas, most notably northeast Scotland in one scenario, and southeast England in another. Autumn changes, like spring, are mixed, with a range of  $-80\%$  to  $+60\%$ , with decreases appearing most frequently in southern England. Annual average flows change little across the century.

The results of Christensen et al. (2012) and Prudhomme et al. (2012) are broadly consistent, given the different time horizons, but this is perhaps not surprising as both are derived from the same RCM results. Overall, summer flows are more likely to reduce through the century across Britain, but Prudhomme et al. (2012) demonstrate that increases remain possible.

Changes in drought frequency and severity are perhaps the most important question for water supply, but neither Christensen et al. (2012) nor Prudhomme et al. (2012) are able to consider such changes, because the change factor approach can only scale historic weather sequences, while GCMs and therefore RCMs are not good at representing the processes (such as blocking weather patterns) that lead to the persistence of extended dry weather across northern Europe (Murphy et al., 2009). Long droughts lasting two years or more are particularly important for water supply in the UK (Marsh et al., 2007; Watts et al., 2012).

Blenkinsop and Fowler (2007) consider precipitation deficit droughts in six different RCMs for the 2080s. Short summer droughts are projected to increase over most of the UK, though there is uncertainty about the sign of change in Scotland and northern England. The longest droughts are projected to become shorter and less severe in most of the RCMs. Blenkinsop and Fowler (2007) caution that climate models may not be able to simulate persistent low-rainfall events, making it difficult to draw conclusions about long droughts. In an apparently contradictory finding, Vidal and Wade (2009) use a slightly different definition of precipitation deficit drought, and find an increase in long droughts in southeast England by the end of the 21st century. Vidal and Wade (2009) agree with Blenkinsop and Fowler (2007) that disagreement between different climate models means that uncertainty is large, and that water supply planners need to consider a range of possible future droughts. Burke et al. (2010) look at droughts from three to 18 months' duration and find an overall increase in droughts of all duration through the 21st century, though with a wide spread that spans decreases as well as increases in drought frequency. As an example, by the end of the 21st century a drought like 1976 (Doornkamp et al., 1980) could occur at anything from the current frequency (perhaps 1 in 100) to 1 in 10 years (Burke et al., 2010). Rahiz and New (2013) consider the spatial coherence of future droughts, showing that by the end of the century droughts may be more coherent, with a higher probability of droughts occurring in different areas at the same time.

Changes in flood magnitude and frequency are as interesting and important as changes in drought occurrence, and equally difficult to assess. There are two main approaches (Wilby et al., 2008): inference from projections of extreme precipitation, and downscaling from GCMs followed by hydrological simulation. Fowler et al. (2007b) use a multimodel ensemble to look at changes in extreme rainfall, dividing

the UK into nine regions based on rainfall coherence. Changes in the ensemble median indicate an increase of 10–20% on 1961–90 extreme rainfall values by 2071, with more confidence about changes in 10-day values than in 1- or 2-day extremes, though the authors report considerable uncertainty in all of these changes. Fowler and Wilby (2010) confirm these projected increases in extreme rainfall across the UK in autumn and winter. Kay et al. (2013b, 2013c) use the UKCP09 probabilistic projections to estimate the range of changes in 20-year return period flood peaks for river basin regions across England, Wales and Scotland. Results showed the largest potential increases in northwest Scotland and southeast England, with the latter showing the greatest range of uncertainty, related to the balance of catchment types present as well as the range of climate change projections (Kay et al., 2013d). Bell et al. (2012) use a distributed hydrological model to examine changes in flood in the Thames catchment, southeast England, and report that for the 2080s the average estimated change in modelled 20-year return period flood peaks is +36% with a range of –11% to +68%. An emerging approach to determining the response of floods to climate change is to link synoptic features, such as atmospheric rivers, to flood occurrence directly and then determine how these features change in a warming climate (Lavers et al., 2011). For example, projections of future atmospheric rivers suggest that by the end of the century there could be a 50–100% increase in the frequency of the largest winter floods in upland Britain (Lavers et al., 2013).

Changes in rainfall and evapotranspiration will also affect groundwater recharge and groundwater levels, though there have been relatively few studies in the UK and, as yet, no national assessment. Jackson et al. (2011) use 13 different GCMs to investigate changes in recharge in a chalk aquifer in southeast England. The ensemble average suggests a 5% reduction in recharge

by the 2080s, with a range extending from a 26% decrease to a 31% increase. In the projections that suggest more recharge, it may take place over a shorter period than now. There has been little work on the impact of climate change on groundwater levels, though reductions in recharge would be expected to lead to lower groundwater levels.

### *8 River, lake and groundwater temperature, water quality and freshwater ecosystems*

Water temperature is expected to shift with climate change, but there has been little work to examine this in the UK. Worldwide, most projections are based on statistical relationships between air temperature and water temperature: these relationships are stronger at monthly resolution than daily, and weaker again at the annual resolution because water temperature varies much less than air temperature (Webb et al., 2008). George et al. (2007) suggest that in the English Lake District summer lake water temperatures may increase at a faster rate than air temperatures, with slower increases in winter lake water temperature. Van Vliet et al. (2011) demonstrate that, globally, changes in flow can be an important control on changes in water temperature, with reductions in flow leading to greater increases in river temperature; projections of predominantly reduced summer flow (Christensen et al., 2012; Prudhomme et al., 2012) may imply enhanced increases in summer UK river water temperature through the 21st century. Future changes in groundwater temperature remain unclear.

Most chemical reactions and bacteriological processes are faster at higher water temperatures, and water temperature and light levels control the growth rate of algae and many aquatic plants (Wade et al., 2002; Whitehead and Hornberger, 1984; Whitehead et al., 2009) as well as the behaviour of aquatic animals including fish and insects (Durance and Ormerod, 2007). More intense rainfall could result in increased

suspended solids (Lane et al., 2007), increased sediment yields (Wilby et al., 1997) and increases in some nutrients (George et al., 2004). Higher water temperatures may lead to increased growth of algae (Whitehead and Hornberger, 1984) and reduced flows may increase the impact of nutrients from agriculture (Whitehead et al., 2006). In the uplands, higher winter rainfall and increased storminess as well as increased summer drought could exacerbate the acidification of acidified but recovering streams and lakes (Curtis et al., 2014; Whitehead et al., 2009; Wilby, 1994). In urban environments, poorer water quality may be driven by short-duration high-intensity rainfall events (Whitehead et al., 2009). Groundwater quality will be affected by changes in temperature and the rate of recharge, as well as the availability of pollutants and nutrients (Bloomfield et al., 2006; Stuart et al., 2011).

What does this mean for UK freshwater ecosystems? Cold water fish species such as Arctic Charr may be threatened by increases in water temperature, with invasive fish species such as Common Carp *Cyprinus carpio*, European Catfish *Silurus glanis* and Roach *Rutilus rutilus* being more successful (Britton et al., 2010; Winfield et al., 2008). Similar changes already seem to be apparent in the upper Rhone (Daufresne et al., 2004). Ecosystem responses can be complex, and it is important to understand how other factors (such as water quality, light levels and flow) interact with changes in water temperature before making speculative assessments of possible change (Durance and Ormerod, 2009).

Few studies have tried to draw all these possible impacts together to understand their combined effects. Whitehead et al. (2006) use a coupled hydrological and nitrate model to investigate flow, nitrate and ammonia changes in a groundwater-fed river in southeast England, finding increases in nitrate and ammonia over the 21st century, mainly driven by higher temperatures and enhanced microbial activity.

Johnson et al. (2009) bring together climate models, hydrological models and expert opinion to draw a picture of the British river of the future, concentrating on contrasting river systems in southeast and northeast England. They conclude that, with lower flows and higher temperatures, there may be more algal blooms and that ecosystems may change, with wetlands particularly vulnerable. These effects are greater in the river in southeast England. Moss et al. (2011) suggest that increasing water temperatures in shallow lakes may lead to complex responses including an increase in the abundance of planktivorous fish, increased predation of zooplankton, and increasingly severe algal blooms.

### 9 Summary of future impacts

It is clear that the impact of anthropogenic climate change on the water cycle and water environment in the UK could be significant (Table 2); changes in rainfall, evapotranspiration, flows and water temperature all affect water quality, ecosystem form and function, and the occurrence of droughts and floods. Approaches to understanding these effects vary greatly; rainfall, evapotranspiration and river flow have been the subject of detailed numerical assessment, but many studies tend to be either site specific or more conjectural. This affects confidence in the assessments. There is greatest confidence in the assessment of changes in air temperature, though the range of possible changes is great. Confidence in the impact on aquatic ecosystems is lowest, because of the multiple interacting drivers of change and poor understanding of the responses of and interactions between species.

## III Discussion and conclusions

In this paper we have summarized and synthesized research on the impact of anthropogenic climate change on the water cycle and water environment of the UK, with the aim of

providing information that helps decision-makers understand and plan for the particular problems that they may face. We use the term ‘decision-makers’ in a wide sense to include all those who have a role in managing or planning water in the UK at a variety of scales; we know that these people are experts in their own fields but often lack access to credible or consistent information about how they may be affected by climate change. For this reason, our review does not try to anticipate the impacts of change beyond the natural environment or to identify the actions that should be taken to adapt to maintain water supply or to defend against floods. Decisions on such actions must take into account economic and social factors as well as other changes that affect the area in question; without in-depth analysis, recommendations for action can appear facile to the experts involved. We also aimed to identify areas where further research would be valuable, thus helping the research community meet the needs of decision-makers more effectively. Hence we divide our recommendations into two, reflecting the two different audiences for this work, though we note that many people work across science and practice, and that this apparent divide is often fluid. Indeed, this translator role is seen as increasingly important, with the IPCC noting that organizations that bridge science and decision-making ‘play an important role in the communication, transfer and development of climate-related knowledge’ (IPCC, 2014: 23).

We recommend that **decision-makers** should:

- Work to understand how their decisions are affected by weather and climate, which may be sensitive to changes in climate and the projected changes in hydrological system response. Armed with such information, decision-makers should be able to interrogate climate change projections more effectively, understanding when and where climate change affects their work.
  - Find ways to plan that accommodate the uncertainty in climate change projections. Uncertainties in future emissions and how the climate system will respond are accompanied by uncertainties in hydrological response. It is also important to set climate change in its wider context; where the planning horizon is short – perhaps 20 years or less – climate variability will dominate (Hawkins and Sutton, 2011) and for all time horizons other social and economic changes will always be of importance (Wade et al., 2013).
  - Work with academics to set the research agenda (note that we mirror this in our recommendations for researchers). If decision-makers expect research to support their work, they should engage actively from the outset, identifying clearly how research can contribute to their planning and decisions. At the same time, decision-makers need to be realistic about how research can support their decisions, and in particular should not hope that research will solve all their problems; the expertise of decision-makers themselves will be vital in understanding and adapting to the impacts of climate change.
- We recommend that **researchers** should:
- Find ways to be clear about uncertainty and confidence, bearing in mind that the two are readily conflated by non-specialists trying to understand the relevance of research. While some papers have very good discussions of the two, others are less convincing and may leave the reader confused.
  - Work to make sure that the main points of their work are accessible to non-specialists. There are still too many academic papers where important findings may be obfuscated by the language used

**Table 2.** Possible future changes to water in the UK.

Area	Projected change	Confidence	Source
Air temperature	Increasing through the 21st century, with greater increases in summer than winter, and greater increases in the south than in the north.	High confidence in temperature increase; medium confidence in the range and pattern of increase.	Murphy et al. (2009)
Precipitation	Little change in annual average precipitation through the 21st century.	Medium confidence of small change in annual average.	Murphy et al. (2009)
	Increase in winter rainfall over much of the UK by the end of the 21st century but with small decreases in Scotland.	Medium confidence in changes, but low confidence in patterns.	Murphy et al. (2009)
	Decrease in summer rainfall by the end of the 21st century, especially in the south, but with little change in summer rainfall in northern Scotland.	Medium confidence in changes but low confidence in patterns.	Murphy et al. (2009)
	Increases in winter, spring and autumn extreme rainfall by the 2080s. Summer extreme rainfall changes less clear.	Medium confidence in increases in extreme rainfall except in summer, where confidence in changes is low.	Fowler and Ekström (2009); Murphy et al. (2009)
Evapotranspiration	Increases in all seasons through the 21st century.	Low confidence in increase or scale of increase.	Kay et al. (2013a)
River flows	Increases in average winter flows by the 2020s and 2050s.	Medium confidence in increasing winter flow.	Christiensen et al. (2012); Prudhomme et al. (2012)
	Reduced summer flows by the 2020s and the 2050s.	Medium confidence in reduced summer flows.	Christiensen et al. (2012); Prudhomme et al. (2012)
	Increased flood magnitude through the 21st century.	Medium confidence in increased flood magnitude; low confidence in the scale of the increase.	Fowler and Wilby (2010); Fowler et al. (2007b); Kay et al. (2013b, 2013c)
Groundwater levels and recharge	No clear picture.	Low confidence in the response of groundwater to climate change.	Jackson et al. (2014)
River water temperature	Increases in river water temperature through the 21st century.	Medium confidence in increases broadly in line with air temperature increase.	Hannah and Garner (2014)
Groundwater temperature	No information available.	Low confidence in the response of groundwater temperature to climate change.	Jackson et al. (2014)

(continued)



**Table 2.** (continued)

Area	Projected change	Confidence	Source
River water quality	Many different possible responses: increased algal blooms, increased suspended solids, increased flushing after dry periods.	Low confidence in the overall impact of different changes on water quality.	Whitehead et al. (2009)
Groundwater quality	Little information available.	Low confidence in the response of groundwater quality to climate change.	Jackson et al. (2014)
Freshwater ecosystems	Decline in species adapted to cold conditions, with increase in invasive species.	Low confidence in the overall impact of changes.	Durance and Ormerod (2009)

Note: confidence is the subjective view of the authors, based on the level of agreement of studies, their spatial coverage and the degree of quantification.

or the way the results are presented. If researchers want their research to be relevant, they need to find ways to make sure that it can be understood by experts from other fields. This may include alternative presentations that complement the academic paper, though in preparing such alternatives academic standards and rigour must be maintained.

- Work with decision-makers to understand their needs and identify how research can help to meet these needs (note that this mirrors the equivalent recommendation for decision-makers). Engagement from the beginning to the end of a project will yield results that are relevant and have genuine impact, but this requires considerable time from both parties, and may be particularly difficult for decision-makers who are not funded for such activities. Researchers sometimes seem to disengage when the academic paper is submitted, but without continued involvement their research may never be embedded in practice.

We believe that scientific syntheses like this and the LWEC water climate impact report

card (Watts and Anderson, 2013) play a valuable role in bridging the gap between research and practice, and indeed between researchers and practitioners. It would be valuable to expand this water review to cover other areas that have not been considered in detail here, such as snow and snow melt, estuaries, built environments and water use. It would also be useful to consider the specific questions of different geographical areas, such as the uplands and wetlands, and the impact of changes in freshwater temperatures, flows and quality on the marine environment.

Despite the growing body of literature on the impact of climate change on the UK water environment, there are several areas where more research would be of value:

- Changes in historical and future evapotranspiration are still poorly understood (Kay et al., 2013a); as different studies calculate evapotranspiration in different ways, it can be hard for practitioners to ensure consistency between baseline and future projections.
- Droughts are important for water supply, power generation and agriculture. As the droughts with the biggest impact in the

UK tend to be both prolonged and spatially coherent, there are relatively few in the historical record (Marsh et al., 2007, provide a valuable catalogue). This means that historical droughts may provide a poor basis for future planning – even if (say) 1976 represents a difficult drought in one place, it may have been less severe elsewhere and planning for a repeat of this drought may be unnecessarily precautionary in some places but over-cautious in others. Projections of future drought are currently limited by the poor understanding of the drivers of long drought and the apparent inability of climate models to reproduce persistent periods of low rainfall in the UK. Improvements in climate modelling may lead to more reliable drought projections, but in any case it would be valuable both to improve understanding of hydrological response to historical droughts and to investigate the level to which systems, such as water supply systems, are resilient to drought.

- Future flooding also remains a difficult area; intense rainfall (particularly summer convective storms) occurs at scales that cannot readily be resolved by current climate models, which is particularly problematical where future flood risk in small catchments is in question. Improvements to climate models may lead to better projections of intense rainfall events, but it would be valuable also to continue work to understand how catchments respond to different events. Further work to understand the physical processes that lead to widespread flooding (e.g. atmospheric rivers; Lavers et al., 2011) could lead to improved understanding of future floods.
- The impact of climate change on groundwater levels and quality is unclear. Baseline groundwater temperatures are poorly

understood but groundwater contributes much of the summer flow in some rivers and streams, directly influencing water temperature.

- The way that aquatic ecosystems respond to climate change remains uncertain, both because many ecosystems are robust to considerable climate variability and because of the complex relationships between flow, water temperature, light, water quality and ecosystem response. High-frequency monitoring (for example, hourly) may help to understand these complex interactions (Wade et al., 2012). High-quality, long-term monitoring programmes are essential in understanding and explaining change, and long records are especially important in detecting and monitoring change and evaluating the effectiveness of adaptation measures.

Perhaps most importantly, it is apparent that there are few studies of change in water and the water environment at scales that help policy-makers and decision-makers examine responses to the problems they face. Large-scale, regional or countrywide assessments are perhaps inevitably compromised by their general nature, but these are the studies that provide the picture of change that decision-makers find most useful. Consistent approaches applied across wide areas allow for comparison between places, identifying the areas at most risk from different water hazards: this is especially important when funds are limited and must be spent where they can deliver the greatest benefits. The next UK Climate Change Risk Assessment, due in 2017, may present a valuable focus for such work. By working together, decision-makers and researchers can identify targeted research that should have genuine impact in improving understanding of the potential effect of climate change on water in the UK.

## Acknowledgements

This work was steered by the Living With Environmental Change (LWEC) partnership. An earlier, longer draft of this paper formed an LWEC working paper. The authors are grateful to Catherine Wright, Molly Anderson and Neil Veitch from the Environment Agency and to Nikki van Dijk from Atkins Global for their guidance and support. Many people provided helpful input to the approach, including the informal working group comprising Konrad Bishop, Jim Hall, Kathryn Humphrey, Hans Jensen, Nick Reynard, Andrew Wade and Steven Wade. Nigel Arnell, Hayley Fowler and Jason Lowe made valuable comments on an early draft. Thanks to Stephen Maberly for helpful comments on a later draft. Bloomfield, Jackson, Mackay and Stuart publish with the permission of the Executive Director of the British Geological Survey (Natural Environment Research Council). The views expressed in this paper are those of the authors alone, and not the organizations for which they work. We thank an anonymous referee for thorough and careful comments that greatly improved the final paper.

## Funding

This work was funded by the UK Department for Environment, Food and Rural Affairs (Defra) (project WC1052) and supported by the Natural Environment Research Council and the Environment Agency.

## References

- Adger WN, Dessai S, Goulden M, Hulme M, Lorenzoni I, Nelson DR, Naess LO, Wolf J and Wreford A (2009) Are there social limits to adaptation to climate change? *Climatic Change* 93:335–354.
- Alexander LV and Jones PD (2001) Updated precipitation series for the UK and discussion of recent extremes. *Atmospheric Science Letters* 1:142–150.
- Allan RP (2011) Human influence on rainfall. *Nature* 470: 344–345.
- Allen RG, Pereira LS, Raes D and Smith M (1998) Crop evapotranspiration – Guidelines for computing crop water requirements. FAO Irrigation and Drainage Paper 56. Rome: Food and Agriculture Organization.
- Arnell NW, Lowe JA, Brown S, Gosling SN, Gottschalk P, Hinkel J, Lloyd-Hughes B, Nicholls RJ, Osborn TJ, Osborne TM, Rose GA, Smith P and Warren RF (2013) A global assessment of the effects of climate policy on the impacts of climate change. *Nature Climate Change* 3: 512–519.
- Arnell NW, van Vuuren DP and Isaac M (2011) The implications of climate policy for the impacts of climate change on global water resources. *Global Environmental Change* 21: 592–603.
- Bates BC, Kundzewicz ZW, Wu S and Palutikof JP (eds) (2008) Climate change and water. Technical Paper of the Intergovernmental Panel on Climate Change. Geneva: IPCC Secretariat, 210 pp.
- Battarbee RW, Simpson GL, Shilland EM, Flower RJ, Kreiser A, Yang H and Clarke G (2012) Recovery of UK lakes from acidification: An assessment using combined palaeoecological and contemporary diatom assemblage data. *Ecological Indicators* 37: 365–380.
- Bell VA, Kay AL, Cole SJ, Jones RG, Moore RJ and Reynard NS (2012) How might climate change affect river flows across the Thames Basin? An area-wide analysis using the UKCP09 Regional Climate Model ensemble. *Journal of Hydrology* 442–443: 89–104.
- Bennion H, Carvalho L, Sayer CD, Simpson GL and Wischniewski J (2012) Identifying from recent sediment records the effects of nutrients and climate on diatom dynamics in Loch Leven. *Freshwater Biology* 57(10): 2015–2029.
- Blenkinsop S and Fowler HJ (2007) Changes in drought frequency, severity and duration for the British Isles projected by the PRUDENCE regional climate models. *Journal of Hydrology* 342: 50–71.
- Bloomfield JP, Williams RJ, Gooddy DC, Cape JN and Guha P (2006) Impacts of climate change on the fate and behaviour of pesticides in surface and groundwater – a UK perspective. *Science of the Total Environment* 369: 163–177.
- Bradford RB and Marsh TJ (2003) Defining a network of benchmark catchments for the UK. *Proceedings of the Institution of Civil Engineers: Water and Maritime Engineering* 156: 109–116.
- Britton JR, Cucherousset J, Davies GD, Godard MJ and Copp GH (2010) Non-native fishes and climate change: Predicting species responses to warming temperatures in a temperate region. *Freshwater Biology* 55: 1130–1141.
- Burke EJ, Perry RJH and Brown SJ (2010) An extreme value analysis of UK drought and projections of change in the future. *Journal of Hydrology* 388: 131–143.

- Burt TP and Ferranti EJS (2011) Changing patterns of heavy rainfall in upland areas: A case study from northern England. *International Journal of Climatology* 32: 518–532.
- Burt TP and Shahgedanova M (1998) An historical record of evaporation losses since 1815 calculated using long-term observations from the Radcliffe Meteorological Station, Oxford, England. *Journal of Hydrology* 205: 101–111.
- Cahill AE, Aiello-Lammens ME, Fisher-Reid MC, Hua X, Karanewsky CJ, Ryu HY, Sbeglia GC, Spagnolo F, Waldron JB, Warsi O and Wiens JJ (2013). How does climate change cause extinction? *Proceedings of the Royal Society of London B* 280: 20121890.
- Caissie D (2006) The thermal regime of rivers: A review. *Freshwater Biology* 51: 1389–1406.
- Christensen BV, Vidal J-P and Wade SJ (2012) Using UKCP09 probabilistic climate information for UK water resource planning. *Journal of Hydrology* 424–425: 48–67.
- Clews E, Durance I, Vaughan IP and Ormerod SJ (2010) Juvenile salmonid populations in a temperate river system track synoptic trends in climate. *Global Change Biology* 16: 3271–3283.
- Climate Change Risk Assessment (CCRA) (2012) Summary of the key findings from the UK Climate Change Risk Assessment. Available at: [http://randd.defra.gov.uk/Document.aspx?Document=Summary\\_of\\_Key\\_Findings.pdf](http://randd.defra.gov.uk/Document.aspx?Document=Summary_of_Key_Findings.pdf).
- Croxtton PJ, Huber K, Collinson N and Sparks TH (2006) How well do the Central England temperature and the England and Wales precipitation series represent the climate of the UK? *International Journal of Climatology* 26: 2287–2292.
- Curtis CJ, Battarbee RW, Monteith D and Shilland E (2014) The future of upland water ecosystems of the UK in the 21st century: A synthesis. *Ecological Indicators* 37: 412–430.
- Dai A (2011) Drought under global warming: A review. *WIREs Climate Change* 2: 45–65.
- Daufresne M, Roger MC, Capra H and Lamouroux N (2004) Long-term changes within the invertebrate and fish communities of the Upper Rhone River: Effects of climatic factors. *Global Change Biology* 10: 124–140.
- Doornkamp JC, Gregory KJ and Burn AS (eds) (1980) *Atlas of Drought in Britain 1975–1976*. London: Institute of British Geographers.
- Durance I and Ormerod SJ (2007) Climate change effects on upland stream invertebrates over a 25 year period. *Global Change Biology* 13: 942–957.
- Durance I and Ormerod SJ (2009) Trends in water quality and discharge confound long-term warming effects on river macro invertebrates. *Freshwater Biology* 54: 388–405.
- Evans CD, Monteith DT and Cooper DM (2005) Long-term increases in surface water dissolved organic carbon: Observations, possible causes and environmental impacts. *Environmental Pollution* 137: 55–71.
- Evans CD, Monteith DT and Harriman R (2001) Long-term variability in the deposition of marine ions at west coast sites in the UK Acid Waters Monitoring Network: Impacts on surface water chemistry and significance for trend determination. *Science of the Total Environment* 265: 115–129.
- Foley B, Jones ID, Maberley SC and Rippey B (2012) Long term changes in oxygen depletion in a small temperate lake: Effects of climate change and eutrophication. *Freshwater Biology* 57: 278–289.
- Fowler HJ and Ekström M (2009) Multi-model ensemble estimates of climate change impacts on UK seasonal precipitation extremes. *International Journal of Climatology* 29(3): 385–416.
- Fowler HJ and Kilsby CG (2003) A regional frequency analysis of United Kingdom extreme rainfall from 1961 to 2000. *International Journal of Climatology* 23(11): 1313–1334.
- Fowler HJ and Wilby RL (2010) Detecting changes in seasonal precipitation extremes using regional climate model projections: Implications for managing fluvial flood risk. *Water Resources Research* 46: W03525.
- Fowler HJ, Blenkinsop S and Tebaldi C (2007a) Linking climate change modelling to impacts studies: Recent advances in downscaling techniques for hydrological modelling. *International Journal of Climatology* 27: 1547–1578.
- Fowler HJ, Ekström M, Blenkinsop S and Smith AP (2007b) Estimating change in extreme European precipitation using a multimodel ensemble. *Journal of Geophysical Research* 112: D18104.
- Freeman C, Evans CD, Monteith DT, Reynolds B and Fenner N (2001) Export of organic carbon from peat soils. *Nature* 412: 785.
- George DG, Hurley M and Hewitt D (2007) The impact of climate change on the physical characteristics of the

- larger lakes in the English Lake District. *Freshwater Biology* 52: 1467–1666.
- George DG, Maberly SC and Hewitt DP (2004) The influence of the North Atlantic Oscillation on the physical, chemical and biological characteristics of four lakes in the English Lake District. *Freshwater Biology* 49: 760–774.
- Giorgi F, Im E-S, Coppola E, Diffenbaugh NS, Gao XJ, Mariotti L and Shi Y (2011) Higher hydroclimatic intensity with global warming. *Journal of Climate* 24(20): 5309–5324.
- Hall J (2008) Probabilistic climate scenarios may misrepresent uncertainty and lead to bad adaptation decisions. *Hydrological Processes* 21: 1127–1129.
- Hannaford J (2015) Climate-driven changes in UK river flows: A review of the evidence. *Progress in Physical Geography* 39: 29–48.
- Hannaford J and Buys G (2012) Seasonal trends in river flow regimes and extremes in the UK. *Journal of Hydrology* 475: 158–174.
- Hannah and Garner (2015) River water temperature in the United Kingdom: Changes over the 20th century and possible changes over the 21st century. *Progress in Physical Geography* 39: 68–92.
- Hannah DM, Demuth S, van Lanen HAJ, Looser U, Prudhomme C, Rees G, Stahl K and Tallaksen LM (2011) Large-scale river flow archives: Importance, current status and future needs. *Hydrological Processes* 25: 1191–1200.
- Hannah DM, Malcolm IA, Soulsby C and Youngson AF (2008) A comparison of forest and moorland stream microclimate, heat exchanges and thermal dynamics. *Hydrological Processes* 22: 919–940.
- Hawkins E and Sutton R (2009) The potential to narrow uncertainty in regional climate predictions. *Bulletin of the American Meteorological Society* 90(8): 1095–1107.
- Hawkins E and Sutton R (2011) The potential to narrow uncertainty in projections of regional precipitation change. *Climate Dynamics* 37: 407–418.
- Howden NJK, Burt TP, Worrall F, Whelan MJ and Bierzoza M (2010) Nitrate concentrations and fluxes in the River Thames over 140 years (1868–2008): Are increases irreversible? *Hydrological Processes* 24: 2657–2662.
- Intergovernmental Panel on Climate Change (IPCC) (2000) *Emissions Scenarios*. Cambridge: Cambridge University Press.
- Intergovernmental Panel on Climate Change (IPCC) (2013a) Summary for policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 1–27.
- Intergovernmental Panel on Climate Change (IPCC) (2013b) Technical summary. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 33–115.
- Intergovernmental Panel on Climate Change (IPCC) (2014) Summary for policymakers. In: *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press, 1–32.
- Jackson CR, Meister R and Prudhomme C (2011) Modelling the effects of climate change and its uncertainty on UK Chalk groundwater resources from an ensemble of global climate model projections. *Journal of Hydrology* 399: 12–28.
- Jackson et al. (2015) Evidence for changes in historic and future groundwater levels in the UK. *Progress in Physical Geography* 39: 49–67.
- Jarvie HP, Neal C, Juergens MD, Sutton EJ, Neal M, Wickham HD, Hill LK, Harman SA, Davies JLL, Warwick A, Barrett C, Griffiths J, Binley A, Swannack N and McIntyre N (2006) Within-river nutrient processing in Chalk streams: The Pang and Lambourn, UK. *Journal of Hydrology* 330(1–2): 101–125.
- Jenkins GJ, Murphy JM, Sexton DS, Lowe J, Jones P and Kilsby C (2002) UK climate projections: Briefing report. Exeter: Met Office Hadley Centre.
- Jenkins GJ, Perry MC and Prior MJ (2008) The climate of the United Kingdom and recent trends. Exeter: Met Office Hadley Centre.
- Jeppesen E, Mehner T, Winfield IJ, Kangur K, Sarvala J, Gerdeaux D, Rask M, Malmquist HJ, Holmgren K, Volta P, Romo S, Eckmann R, Sandstrom A, Blanco S, Kangur A, Ragnarsson Stabo H, Tarvainen M, Ventela A-M, Sondergaard M, Lauridsen TL and Meerhoff M (2012) Impacts of climate warming on the long-term dynamics of key fish species in 24 European lakes. *Hydrobiologia* 694: 1–39.

- Johnson AC, Acreman MC, Dunbar MJ, Feist SW, Giacomello AM, Gozlan RE, Hinsley SA, Ibbotson AT, Jarvie HP, Jones JI, Longshaw M, Maberly SC, Marsh TJ, Neal C, Newman JR, Nunn MA, Pickup RW, Reynard NS, Sullivan CA, Sumpter JP and Williams RJ (2009) The British river of the future: How climate change and human activity might affect two contrasting river ecosystems in England. *Science of the Total Environment* 407: 4787–4798.
- Jones MR, Fowler HJ, Kilsby CG and Blenkinsop S (2012) An assessment of changes in seasonal and annual extreme rainfall in the UK between 1961 and 2009. *International Journal of Climatology* 33(5): 1178–1194.
- Karoly DJ and Stott PA (2006) Anthropogenic warming of central England temperature. *Atmospheric Science Letters* 7: 81–85.
- Kay AL and Davies HN (2008) Calculating potential evaporation from climate model data: A source of uncertainty for hydrological climate change impacts. *Journal of Hydrology* 358: 221–239.
- Kay AL and Jones RG (2012) Comparison of the use of alternative UKCP09 products for modelling the impacts of climate change on flood frequency. *Climatic Change* 114(2): 211–230.
- Kay AL, Bell VA, Blyth EM, Crooks SM, Davies HN and Reynard NS (2013a) A hydrological perspective on evaporation: Historical trends and future projections in Britain. *Journal of Water and Climate Change* 4(3): 193–208.
- Kay AL, Crooks SM, Davies HN, Prudhomme C and Reynard NS (2013b) Probabilistic impacts of climate change on flood frequency using response surfaces. I: England and Wales. *Regional Environmental Change* 14(3): 1215–1227.
- Kay AL, Crooks SM, Davies HN and Reynard NS (2013c) Probabilistic impacts of climate change on flood frequency using response surfaces. II: Scotland. *Regional Environmental Change* 14(3): 1243–1255.
- Kay AL, Crooks SM, Pall P and Stone DA (2011) Attribution of autumn/winter 2000 flood risk in England to anthropogenic climate change: A catchment-based study. *Journal of Hydrology* 406: 97–112.
- Kay AL, Crooks SM and Reynard NS (2013d) Using response surfaces to estimate impacts of climate change on flood peaks: Assessment of uncertainty. *Hydrological Processes*. doi:10.1002/hyp.10000.
- Kernan MR, Battarbee RW and Moss B (eds) (2010) *Climate Change Impacts on Freshwater Ecosystems*. Chichester: Wiley.
- Kingston DG, Todd MC, Taylor RG, Thompson JR and Arnell NW (2009) Uncertainty in the estimation of potential evapotranspiration under climate change. *Geophysical Research Letters* 36: L20403.
- Lane SN, Tayefi V, Reid SC, Yu D and Hardy RJ (2007) Interactions between sediment delivery, channel change, climate change and flood risk in a temperate upland environment. *Earth Surface Processes and Landforms* 32: 429–446.
- Langan SJ, Johnston L, Donaghy MJ, Youngson AF, Hay DW and Soulsby C (2001) Variation in river water temperatures in an upland stream over a 30-year period. *Science of the Total Environment* 265: 195–207.
- Lavers DA, Allan RP, Villarini G, Lloyd-Hughes B, Brayshaw DJ and Wade AJ (2013) Future changes in atmospheric rivers and their implications for winter flooding in Britain. *Environmental Research Letters* 8: 034010.
- Lavers DA, Allan RP, Wood EF, Villarini G, Brayshaw DJ and Wade AJ (2011) Winter floods in Britain are connected to atmospheric rivers. *Geophysical Research Letters* 38: L23803.
- Lempert RJ, Groves DG, Popper SW and Bankes SC (2006) A general, analytic method for generating robust strategies and narrative scenarios. *Management Science* 52(4): 514–528.
- Marsh T, Cole G and Wilby R (2007) Major droughts in England and Wales 1800–2006. *Weather* 62(4): 87–93.
- Meyer JL, Sale MJ, Mulholland PJ and Poff NL (1999) Impacts of climate change in aquatic ecosystem functioning and health. *Journal of American Water Resources Association* 33(6): 1373–1386.
- Millennium Ecosystem Assessment (2005) *Ecosystems and Human Well-Being: Synthesis*. Washington, DC: Island Press.
- Monteith DT, Evans CD, Henrys PA, Simpson GL and Malcolm IA (2012) Trends in the hydrochemistry of acid-sensitive surface waters in the UK 1988–2008. *Ecological Indicators* 37: 287–303.
- Monteith DT, Evans CD and Patrick ST (2001) Monitoring acid waters in the UK: 1988–1998 trends. *Water, Air and Soil Pollution* 130:1307–1312.
- Monteith DT, Evans CD and Reynolds B (2000) Are temporal variations in the nitrate content of UK upland freshwaters linked to the North Atlantic Oscillation? *Hydrological Processes* 14: 1745–1749.

- Monteith DT, Stoddard JL, Evans CD, de Wit HA, Forsius M, Høgåsen T, Wilander A, Skjelkvåle BL, Jeffries DS, Vuorenmaa J, Keller B, Kopáček J and Vesely J (2007) Dissolved organic carbon trends resulting from changes in atmospheric deposition chemistry. *Nature* 450: 537–540.
- Monteith JL (1965) Evaporation and environment. *Symposium of the Society for Experimental Biology* 19: 205–234.
- Moser SC and Ekstrom JA (2010) A framework to diagnose barriers to climate change adaptation. *Proceedings of the National Academy of Sciences* 107(51): 22026–22031.
- Moss B, Kosten S, Meerhoff M, Battarbee RW, Jeppesen E, Mazzeo N, Havens K, Lacerot G, Liu Z, De Meester L, Paerl H and Scheffer M (2011) Allied attack: Climate change and eutrophication. *Inland Waters* 1: 101–105.
- Murphy JM, Booth BBB, Collins M, Harris GR, Sexton DMH and Webb MJ (2008) A methodology for probabilistic predictions of regional climate change from perturbed physics ensembles. *Philosophical Transactions of the Royal Society A* 365:1993–2028.
- Murphy JM, Sexton DMH, Jenkins GJ, Boorman P, Booth B, Brown K, Clark R, Collins M, Harris G and Kendon L (2009) UK climate projections science report: Climate change projections. Exeter: Met Office Hadley Centre.
- Neal C, Jarvie HP, Williams R, Love A, Neal M, Wickham H, Harman S and Armstrong L (2010) Declines in phosphorus concentration in the upper River Thames (UK): Links to sewage effluent cleanup and extended end-member mixing analysis. *Science of the Total Environment* 408(6):1315–1330.
- New M, Lopez A, Dessai S and Wilby R (2007) Challenges in using probabilistic climate change information for impact assessments: An example from the water sector. *Philosophical Transactions of the Royal Society A* 365: 2117–2131.
- Ormerod SJ (2009) Climate change, river conservation and the adaptation challenge. *Aquatic Conservation: Marine and Freshwater Ecosystems* 19: 609–613.
- Orr HG, Simpson GL, des Clers S, Watts G, Hughes M, Hannaford J, Dunbar MJ, Laizé CLR, Wilby RL, Battarbee RW and Evans R (2014) Detecting changing river temperatures in England and Wales. *Hydrological Processes*. doi: 10.1002/hyp.10181.
- Osborn TJ and Hulme M (2002) Evidence for trends in heavy rainfall events over the United Kingdom. *Philosophical Transactions of the Royal Society A* 360: 1313–1325.
- Pall P, Aina T, Stone DA, Stott PA, Nozawa T, Hilberts AGJ, Lohmann D and Allen MR (2011) Anthropogenic greenhouse gas contribution to flood risk in England and Wales in autumn 2000. *Nature* 470: 382–385.
- Penman HL (1948) Natural evaporation from open water, bare soil and grass. *Proceedings of the Royal Society of London A* 193: 120–145.
- Prudhomme C, Reynard N and Crooks S (2002) Down-scaling of global climate models for flood frequency analysis: Where are we now? *Hydrological Processes* 16: 1137–1150.
- Prudhomme C, Young A, Watts G, Haxton T, Crooks S, Williamson J, Davies H, Dadson S and Allen S (2012) The drying up of Britain? A national estimate of changes in seasonal river flows from 11 Regional Climate Model simulations. *Hydrological Processes* 26: 1115–1118.
- Rahiz M and New M (2013) 21st century drought scenarios for the UK. *Water Resources Management* 27(4): 1039–1061.
- Rockström J, Steffen W, Noone K, Persson Å, Chapin FS III, Lambin E, Lenton TM, Scheffer M, Folke C, Schellnhuber H, Nykvist B, De Wit CA, Hughes T, van der Leeuw S, Rodhe H, Sörlin S, Snyder PK, Costanza R, Svedin U, Falkenmark M, Karlberg L, Corell RW, Fabry VJ, Hansen J, Walker BH, Liverman D, Richardson K, Crutzen C and Foley J (2009) A safe operating space for humanity. *Nature* 461: 472–475.
- Shand P, Edmunds WM, Lawrence AR, Smedley PL and Burke S (2007) The natural (baseline) quality of groundwater in England and Wales. British Geological Survey Research Report No. RR/07/06.
- Sheffield J, Wood EF and Roderick ML (2012) Little change in global drought over the past 60 years. *Nature* 491: 435–438.
- Stern N (2006) *The Economics of Climate Change*. Cambridge: Cambridge University Press.
- Stuart ME, Goody DC, Bloomfield JP and Williams AT (2011) A review of the impact of climate change on future nitrate concentrations in groundwater of the UK. *Science of the Total Environment* 409: 2859–2873.
- van Vliet MTH, Ludwig F, Zwolsman JGG, Weedon GP and Kabat P (2011) Global river temperatures and sensitivity to atmospheric warming and changes

- in river flow. *Water Resources Research* 47: W02544.
- Verdonschot PFM, Hering D, Murphy J, Jähnig SC, Rose NL, Graf W, Brabec K and Sandin L (2010) Climate change and the hydrology and morphology of freshwater ecosystems. In: Kernan MR, Battarbee RW and Moss B (eds) *Climate Change Impacts on Freshwater Ecosystems*. Chichester: Wiley, 65–83.
- Vidal J-P and Wade S (2009) A multimodel assessment of future climatological droughts in the United Kingdom. *International Journal of Climatology* 29: 2056–2071.
- Wade AJ, Palmer-Felgate EJ, Halliday SJ, Skeffington RA, Loewenthal M, Jarvie HJ, Bowes MJ, Greenway GM, Haswell SJ, Bell IM, Joly E, Fallatah A, Neal C, Williams RJ, Gozzard E and Newman JR (2012) Hydrochemical processes in lowland rivers: Insights from in situ, high-resolution monitoring. *Hydrology and Earth System Science* 16: 4323–4342.
- Wade AJ, Whitehead PG, Hornberger GM and Snook D (2002) On modelling the flow controls on macrophytes and epiphyte dynamics in a lowland permeable catchment: The River Kennet, southern England. *Science of the Total Environment* 282–283: 395–417.
- Wade SD, Rance J and Reynard N (2013) The UK Climate Change Risk Assessment, 2012: Assessing the impacts on water resources to inform policy makers. *Water Resources Management* 27(4): 1085–1109.
- Watts G and Anderson M (2013) A climate change report card for water. LWEC report card. Available at: [http://www.lwec.org.uk/sites/default/files/attachments\\_report\\_cards/Water%20Card%20English%20web.pdf](http://www.lwec.org.uk/sites/default/files/attachments_report_cards/Water%20Card%20English%20web.pdf).
- Watts G, Christierson BV, Hannaford J and Lonsdale K (2012) Testing the resilience of water supply systems to long droughts. *Journal of Hydrology* 414–415: 255–267.
- Webb BW and Walling DE (1992) Long term water temperature behaviour and trends in a Devon, UK, river system. *Hydrological Sciences Journal* 37: 567–580.
- Webb BW, Hannah DM, Moore RD, Brown LE and Nobilis F (2008) Recent advances in stream and river temperature research. *Hydrological Processes* 22: 902–918.
- Werritty A (2002) Living with uncertainty: Climate change, river flows and water resource management in Scotland. *Science of the Total Environment* 294(1–3): 29–40.
- Whitehead PG (1990) Modelling nitrate from agriculture into public water supplies. *Philosophical Transactions of the Royal Society B* 329: 403–410.
- Whitehead PG and Hornberger GE (1984) Modelling algal behaviour in the River Thames. *Water Research* 18: 945–953.
- Whitehead PG, Wilby RL, Battarbee RW, Kernan M and Wade AJ (2009) A review of the potential impacts of climate change on surface water quality. *Hydrological Sciences Journal/Journal des Sciences Hydrologiques* 54(1): 101–123.
- Whitehead PG, Wilby RL, Butterfield D and Wade AJ (2006) Impacts of climate change on nitrogen in lowland chalk streams: Adaptation strategies to minimise impacts. *Science of the Total Environment* 365: 260–273.
- Wilby RL (1994) Exceptional weather in the Midlands, UK, during 1988–1990 results in the rapid acidification of an upland stream. *Environmental Pollution* 86: 15–19.
- Wilby RL and Dessai S (2010) Robust adaptation to climate change. *Weather* 65(7): 180–185.
- Wilby RL and Quinn NW (2013) Reconstructing multi-decadal variations in fluvial flood risk using atmospheric circulation patterns. *Journal of Hydrology* 487: 109–121.
- Wilby RL, Beven KJ and Reynard NS (2008) Climate change and fluvial flood risk in the UK: More of the same? *Hydrological Processes* 22: 2511–2523.
- Wilby RL, Dalglish HY and Foster IDL (1997) The impact of weather patterns on contemporary and historic catchment sediment yields. *Earth Surface Processes and Landforms* 22: 353–363.
- Winfield IJ, Fletcher JM and James JB (2008). The Arctic charr (*Salvelinus alpinus*) populations of Windermere, UK: Population trends associated with eutrophication, climate change and increased abundance of roach (*Rutilus rutilus*). *Environmental Biology of Fishes* 83: 25–35.
- Worrall F, Burt TP and Adamson J (2004) Can climate change explain increases in DOC flux from upland peat catchments? *Science of the Total Environment* 326: 95–112.
- Worrall F, Swank WT and Burt TP (2003) Changes in stream nitrate concentrations due to land management practices, ecological succession, and climate: Developing a systems approach to integrated catchment response. *Water Resources Research* 39: 1177.
- Wright JL, Clarke RT, Gunn RJM, Winder JM, Kneebone NT and Davy-Bowker J (2004) Response of the flora



- and macroinvertebrate fauna of a chalk stream site to changes in management. *Freshwater Biology* 48: 894–911.
- Wright RF, Larssen T, Camarero L, Cosby BJ, Ferrier RC, Helliwell R, Forsius M, Jenkins A, Kopáček J, Majer V, Moldan F, Posch M, Rogora M and Schöpp W (2005) Recovery of acidified European surface waters. *Environmental Science and Technology* 39: 64A–72A.
- Young AR (2006) Stream flow simulation within UK ungauged catchments using a daily rainfall-runoff model. *Journal of Hydrology* 320: 155–172.