

Raft-partitioning of calcium channels regulates their function

Article

Accepted Version

Ronzitti, G., Bucci, G., Stephens, G. ORCID: <https://orcid.org/0000-0002-8966-4238> and Chierregatti, E. (2015) Raft-partitioning of calcium channels regulates their function. *Channels*, 9 (4). pp. 169-170. ISSN 1933-6969 doi: <https://doi.org/10.1080/19336950.2015.1063285> Available at <https://centaur.reading.ac.uk/40584/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1080/19336950.2015.1063285>

Publisher: Taylor & Francis

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



This article was downloaded by: [University of Reading]

On: 24 June 2015, At: 02:34


Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Editor-in-Chief
Gerald Cooper
...
Special Features Editor
Andreas P. Davis

Volume 9 • Issue 3 • May/June 2015



Editorial Board

David Allen	Sharon Gordon	Domenico Lupatini	Michael Palmer	Edin Turek
John Allison	Andrew Brown	Brian Martin	David Tsao	Yin-Qing Wang
Ken Buse	Yany Heber	Hana Mikhlik	Concepción Sánchez	Stephen Warren
Francisco González	Yael Kuperman	Yael Tsoref	Shufu Yao	David Whaley
Emmanuel Fassinat	Benji Hildes	Joel Nagler	Mark Wagner	Wolfgang Wolf
William Gifford	Tom Homan	Justin Robinson	Christina Wang	
Haruo Hoshi	Leonard Kanner	Chia-Niuh	Yun-Wah Wong	
V.R. Marder	Yoshitaka Kawa	Kevin Pahl	Shih-Wei Yang	
Luca Di Rienzo	Ramona Lerner	Michael Pehkonen	Jing-Jing Zhang	
Armin Mikhlik	Sheng-Li Luo	Kenan Pehlivan	Julian Zentgraf	
Angela Di Rienzo	Mark Lerner	Dimitris Pournaras	Hiroaki Yano	
David Little	Ben Little	Julian Raza	Chun-Yuan Yeh	
Hilary French	Shari Lyman	Alida Rodriguez	Naohiko Yoshida	
Helen Gorman	Iman Lyman	Patrick	Michael Zentgraf	



Channels

Publication details, including instructions for authors and subscription information:

<http://www.tandfonline.com/loi/kchl20>

Raft-partitioning of calcium channels regulates their function

Giuseppe Ronzitti^a, Giovanna Bucci^b, Gary Stephens^c & Evelina Chieriegatti^d

^a Genethon; Evry, France

^b Department of Physics; University of Oxford; Oxford, UK

^c School of Pharmacy; University of Reading; Whiteknights, Reading, UK

^d Department of Neuroscience and Brain Technologies; Istituto Italiano di Tecnologia; Genoa, Italy

Accepted author version posted online: 23 Jun 2015.



[Click for updates](#)

To cite this article: Giuseppe Ronzitti, Giovanna Bucci, Gary Stephens & Evelina Chieriegatti (2015): Raft-partitioning of calcium channels regulates their function, Channels, DOI: [10.1080/19336950.2015.1063285](https://doi.org/10.1080/19336950.2015.1063285)

To link to this article: <http://dx.doi.org/10.1080/19336950.2015.1063285>

Disclaimer: This is a version of an unedited manuscript that has been accepted for publication. As a service to authors and researchers we are providing this version of the accepted manuscript (AM). Copyediting, typesetting, and review of the resulting proof will be undertaken on this manuscript before final publication of the Version of Record (VoR). During production and pre-press, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal relate to this version also.

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

Raft-partitioning of calcium channels regulates their function

Giuseppe Ronzitti¹, Giovanna Bucci², Gary Stephens³ and Evelina Chiergatti⁴

¹Genethon; Evry, France, ²Department of Physics; University of Oxford; Oxford, UK, ³School of Pharmacy; University of Reading; Whiteknights, Reading, UK, ⁴Department of Neuroscience and Brain Technologies; Istituto Italiano di Tecnologia; Genoa, Italy

Keywords: alpha-synuclein, lipid rafts, calcium channels, plasma membrane

Commentary to: Ronzitti G, et al. Exogenous α -Synuclein Decreases Raft Partitioning of Cav2.2 Channels Inducing Dopamine Release. *J Neurosci* 2014; 34(32):10603-15; <http://dx.doi.org/10.1523/JNEUROSCI.0608-14.2014>

Alpha-synuclein (Syn) is a cytosolic, soluble, and unstructured protein, which may also associate with membranes by acquiring an alpha-helical conformation. High Syn concentrations are present in genetic Parkinson's disease (PD) due to duplication/triplication of *SNCA* gene, and are also associated with polymorphisms found in idiopathic PD cases¹. Syn has been shown to act on several intracellular targets, and to alter synaptic function. The discovery of the presence of Syn in cerebrospinal fluid and of its release from neurons, as demonstrated by numerous studies², suggests that Syn-driven pathology may spread in the brain, although the mechanism of Syn release is still unclear. There is evidence that Syn overexpression modifies the structure of the cellular membrane, however a possible effect of excess Syn on the pre- or post-synaptic terminal from the extracellular *milieu* has yet to be fully analyzed.

In our recent paper³, we show that treatment of isolated rat cortical neurons with recombinant, highly purified, monomeric Syn leads to an increase in KCl-stimulated calcium influx through Ca_v2.2 (N-type) calcium channels. The increase in calcium entry is evident at the presynaptic compartment, as shown by the use of the synaptic vesicles-targeted calcium dye SyGCaMP3, suggesting that Syn may profoundly affect synaptic neurotransmitter release.

The activating effect on calcium channels has been confirmed in primary cultures of superior cervical ganglion neurons (SCGN), a model used to study the biophysical properties of Ca_v2.2 in a physiological context. Syn pre-treatment significantly increases both calcium conductance and voltage sensitivity of the channels. Accordingly, augmented cholinergic transmission has been observed in long-term cultured SCGN after acute Syn application. Ca_v2.2 channels play a fundamental role in the release of dopamine in the striatum, where dopaminergic innervations target and control function. Our study shows that Syn perfusion of striatal brain slices induces a significant increase of dopamine release, an effect confirmed *in vivo* by microdialysis of Syn in rat striatum.

Ca_v2.2 channels are multimeric protein complexes that allow functional coupling between action potentials and calcium entry into nerve terminals, thus controlling neurotransmitter release. Ca_v2.2 regulation occurs by the integration of different pathways involving G protein subunits, synaptic proteins such as syntaxin, and kinases such as PKC. However, we show that Syn does not affect any of these targets; rather, analysis of channel biophysical parameters favours that Syn facilitates Ca_v2.2 transition from a closed to an open state by allowing movement of channel gating charge. This effect can be related to changes occurring in the stiffness of the plasma membrane where the channel is inserted, as dictated by lipid composition and organization of the

membrane. In this regard, the level of cholesterol in the membrane affects its structure, as demonstrated by the use of reagents that either decrease or increase cellular cholesterol content. Our study shows that, although Syn treatment leaves the total cellular cholesterol content unchanged, it decreases the level of the lipid in the plasma membrane. Interestingly, this effect is accompanied by a translocation of $Ca_v2.2$ from high to low cholesterol compartments. Lipid rafts are patches in the membrane with high cholesterol content that constitute ordered lipid domains, compared to the disordered bilayer. We propose that the demonstrated shift of $Ca_v2.2$ channels into non-raft domains, where lower energy is required for membrane deformation, favours channel transition to the open state. The net activity of the channel will then derive from a balance between its localization in the membrane and the relative localization of its regulatory proteins.

Reports in the literature show that cholesterol depletion by methyl- β -cyclodextrin induces an increase in calcium current, as we observe with Syn treatment. Moreover cholesterol loading of neurons causes a clear decrease in calcium current⁴. We show that methyl- β -cyclodextrin causes an increase in calcium entry, but it also decreases synaptic vesicles release by altering membrane curvature⁵; we further show that extracellular Syn, acting only at the plasma membrane level, is able to couple the increase in $Ca_v2.2$ activity and intracellular calcium levels with an increase in neurotransmitter release. Conversely, cholesterol loading causes a concentration-dependent decrease in excitatory postsynaptic potential amplitude in SCGN synapses, consistent with a reduction in calcium current (Figure 1).

The effect of Syn on cholesterol is still unknown, whether inducing an internalization of cholesterol-rich domains, or a change in size/number of lipid rafts; however, the effect of Syn

appears selective for Cav2.2 over the other major presynaptic calcium channel, Cav2.1. This may suggest that insertion and re-localization pathways differ for these channels. Evidence suggests that Cav2.2 and Cav2.1 insert into different 'slots' in central synapses⁶, this difference may extend to re-localization by Syn.

Syn-mediated enhancement of Cav2.2 activity may explain the increase in DA release that precedes degeneration of dopaminergic terminals observed in animal models of PD⁷, and recently it was shown that mutant Syn increases the firing rate of substantia nigra neurons, a subtype of dopaminergic neurons selectively vulnerable to degeneration⁸. The increased concentration of DA may therefore account for the toxicity observed in dopaminergic neurons at later stages of disease.

Accepted Manuscript

1. Venda, L.L., et al. Trends Neurosci 2010; 33: 559-568.
2. Marques, O., et al. Cell death & disease 2012; 3: e350.
3. Ronzitti, G., et al. J Neurosci 2014; 34: 10603-10615.
4. Toselli, M., et al. Biophys J 2005; 89: 2443-2457.
5. Rituper, B., et al. Biochimica et biophysica acta 2013; 1831: 1228-1238.
6. Cao, Y.Q., et al. J Neurosci 2010; 30: 4536-4546.
7. Lam, H.A., et al. J Neurosci Res 2011; 89: 1091-1102.
8. Subramaniam, M., et al. J Neurosci 2014; 34: 13586-13599.




Figure 1. Cartoon showing the effect of cholesterol-loaded methyl-β-cyclodextrin (Chol-MβCD), methyl-β-cyclodextrin (MβCD), and Syn on the plasma membrane cholesterol content and on the activity of Ca_v2.2.