

Possible impacts of a future grand solar minimum on climate: stratospheric and global circulation changes

Article

Accepted Version

Open access prepublished version

Maycock, A. C., Ineson, S., Gray, L. J., Scaife, A. A., Anstey, J. A., Lockwood, M. ORCID: <https://orcid.org/0000-0002-7397-2172>, Butchart, N., Hardiman, S. C., Mitchell, D. M. and Osprey, S. M. (2015) Possible impacts of a future grand solar minimum on climate: stratospheric and global circulation changes. *Journal of Geophysical Research: Atmospheres*, 120 (18). pp. 9043-9058. ISSN 2169-8996 doi: 10.1002/2014JD022022 Available at <https://centaur.reading.ac.uk/40588/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1002/2014JD022022>

To link to this article DOI: <http://dx.doi.org/10.1002/2014JD022022>

Publisher: American Geophysical Union

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Possible impacts of a future Grand Solar Minimum on climate: Stratospheric and global circulation changes

A. C. Maycock^{1,2}, S. Ineson³, L. J. Gray^{2,4}, A. A. Scaife³, J. A. Anstey⁴, M. Lockwood⁵, N. Butchart³, S. C. Hardiman³, D. M. Mitchell⁴, S. M. Osprey²

A. C. Maycock, Centre for Atmospheric Science, University of Cambridge, Cambridge, CB2 1EW, UK. (acm204@cam.ac.uk)

S. Ineson, Met Office Hadley Centre, Met Office, Exeter, UK.

L. J. Gray, Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK.

A. A. Scaife, Met Office Hadley Centre, Met Office, Exeter, UK.

J. A. Anstey, Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK.

M. Lockwood, Department of Meteorology, University of Reading, Reading, UK.

N. Butchart, Met Office Hadley Centre, Met Office, Exeter, UK.

S. C. Hardiman, Met Office Hadley Centre, Met Office, Exeter, UK.

D. M. Mitchell, Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK.

S. M. Osprey, Department of Atmospheric, Oceanic and Planetary Physics, University of Oxford, Oxford, UK.

¹Centre for Atmospheric Science,
University of Cambridge, Cambridge, UK.

²National Centre for Atmospheric Science,
UK.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/2014JD022022

Abstract. It has been suggested that the Sun may evolve into a period of lower activity over the 21st century. This study examines the potential climate impacts of the onset of an extreme ‘Maunder Minimum like’ grand solar minimum using a comprehensive global climate model. Over the second half of the 21st century, the scenario assumes a decrease in total solar irradiance of 0.12% compared to a reference RCP8.5 experiment. The decrease in solar irradiance cools the stratopause (~ 1 hPa) in the annual and global mean by 1.4 K. The impact on global mean near-surface temperature is small (~ -0.1 K), but larger changes in regional climate occur during the stratospheric dynamically active seasons. In Northern hemisphere (NH) winter-time, there is a weakening of the stratospheric westerly jet by up to $\sim 3-4$ m s^{-1} , with the largest changes occurring in January-February. This is accompanied by a deepening of the Aleutian low at the surface and an increase in blocking over northern Europe and the north Pacific. There is also an equator-

³Met Office Hadley Centre, Met Office,
Exeter, UK.

⁴Department of Atmosphere, Ocean and
Planetary Physics, University of Oxford,
Oxford, UK.

⁵Department of Meteorology, University
of Reading, Reading, UK.

ward shift in the Southern hemisphere (SH) midlatitude eddy-driven jet in austral spring. The occurrence of an amplified regional response during winter and spring suggests a contribution from a top-down pathway for solar-climate coupling; this is tested using an experiment in which ultraviolet (200–320 nm) radiation is decreased in isolation of other changes. The results show that a large decline in solar activity over the 21st century could have important impacts on the stratosphere and regional surface climate.

Accepted Article

1. Introduction

Electromagnetic radiation from the Sun is a fundamental source of energy for the terrestrial climate system. Therefore changes in solar activity have the potential to influence global climate. The Sun's output varies on a number of characteristic timescales. In the context of Earth's climate, the most frequently studied of these is the approximately 11-year (Schwabe) solar cycle, which is typically associated with a maximum to minimum change in total solar irradiance (TSI) of $\sim 1 \text{ W m}^{-2}$ or $\sim 0.07\%$ [Gray *et al.*, 2010]. The Sun's output is also known to vary on longer timescales; however, characterising these variations requires a much longer record of solar activity. Direct measurements of sunspot numbers extend back to 1610 [Ribes and Nesme-Ribes, 1993], but proxy records must be used to reconstruct solar activity further back in time. Steinhilber *et al.* [2008] compiled a record of the solar modulation potential, ϕ , for the last 9300 years. This is a measure of the shielding of the Earth from galactic cosmic rays (GCRs) by the Sun's magnetic field, and is derived from cosmogenic radionuclide data from ice cores. Lockwood [2010] showed that, when smoothed to remove the signal of the 11-year cycle, the ϕ record exhibits 'grand maxima' and 'grand minima' with a timescale of $\sim 100\text{-}200$ years. The period of relatively high solar activity over the last ~ 50 years has coincided with a so-called 'grand solar maximum', and the period of low solar activity during the late 17th century, known as the Maunder Minimum (MM), is believed to have coincided with a 'grand solar minimum'. Abreu *et al.* [2008] conducted a statistical analysis of the ϕ record and deduced that the current grand maximum is only likely to persist for up to another 15-36 years, after which the Sun would be expected to evolve towards a state of lower output. It has been sug-

gested that the amplitude and persistence of the recent 11-year solar cycle 23 minimum and the relatively low cycle 24 maximum may be indicative of the onset of a grand solar minimum [Owens *et al.*, 2011; Lockwood, 2011]. However, the timescale and amplitude of such a grand solar minimum is unpredictable and highly uncertain. Barnard *et al.* [2011] used the ϕ record to construct a range of possible future scenarios for solar activity based on past variations (see also Lockwood [2010]). They deduced that there is an $\sim 8\%$ chance of returning to the MM-like levels of solar activity within the next ~ 40 years. However, the current rate of decline is larger than at any other point in the ϕ record, and hence the estimated likelihood for this extreme scenario to occur has been increased to 15-20% [M. Lockwood, pers. comms.].

Given the fundamental role of solar energy in the climate system, a period of low solar activity may have important ramifications for the state of both the stratosphere and troposphere, and it is these aspects which are the focus of this study. It has been found, for example, that colder UK winters tend to occur more frequently during periods of low solar activity [Lockwood *et al.*, 2010]. Jones *et al.* [2012] examined the impact of a range of possible future TSI scenarios on global mean surface temperatures using a simple energy balance climate model. They found that a descent into MM-like conditions over the next ~ 70 years would only decrease global mean surface temperatures by up to ~ 0.2 K, with some uncertainty depending on the assumed reconstruction of past TSI. Feulner and Rahmstorf [2010] reached similar conclusions about the impact on global surface temperature using an intermediate complexity model and two scenarios for a decline in TSI of 0.08% and 0.25% relative to 1950 levels. These results make clear that even a large reduction in solar output would only offset a small fraction of the projected global warm-

ing due to anthropogenic activities. This has been further emphasised by Meehl *et al.* [2013], who used a comprehensive climate model to show that a 0.25% decrease in TSI in the mid-21st century would only offset the projected anthropogenic global warming trend by a few tenths of a degree. In addition to considerations of the impact on global mean climate, where solar influences are thought to be small, there has been considerable interest and debate surrounding mechanisms for an amplified *regional* responses to solar perturbations (see e.g. Gray *et al.* [2010, 2013]). These are broadly categorised as: ‘top-down’ mechanisms, which focus on the impact of changes in ultraviolet (UV) radiation on stratospheric temperatures and ozone and associated changes in the extratropical stratospheric circulation (e.g. Haigh [1996]; Soukharev and Hood [2006]; Frame and Gray [2010]), which can impact on surface weather and climate via stratosphere-troposphere dynamical coupling (e.g. Kuroda and Kodera [2002]; Haigh *et al.* [2005]; Matthes *et al.* [2006]; Ineson *et al.* [2011]); and ‘bottom-up’ mechanisms, which focus on changes in surface heating and sensible and latent heat fluxes over the oceans and associated coupled air-sea feedbacks (e.g. White *et al.* [1997]; Meehl *et al.* [2008, 2009]; Misios and Schmidt [2012]; Scaife *et al.* [2013]). Anet *et al.* [2013] found amplified cooling over the Arctic and a warming over the North Atlantic in response to a 6 W m^{-2} (0.45%) decrease in TSI over the 21st century; the North Atlantic warming was related to a reduction in the weakening of the Atlantic meridional overturning circulation due to climate change, which may be partly related to a top-down pathway [Reichler *et al.*, 2012]. Meehl *et al.* [2013] also found regionally dependent temperature changes in response to a sudden decrease in TSI in the tropical east Pacific, where an initial warm anomaly transitioned to a cold anomaly after around a decade. This is broadly similar to the east Pacific response to the

11-year solar cycle identified in some studies (e.g. Meehl *et al.* [2009]). In addition to the effects of changes in solar irradiance, there has also been discussion around the possible climate impacts of changes in solar energetic particle fluxes. For example, Seppälä *et al.* [2013] analysed reanalysis data and found a stronger Arctic polar vortex under high solar geomagnetic activity and changes in surface temperature that resemble the positive phase of the North Atlantic Oscillation (NAO) [Seppälä *et al.*, 2009]. Such effects will not be considered in this study because solar particles are not currently represented in the climate model employed here; further research is therefore required to test whether this may also play a role in climate in the case of a decline in solar activity.

In this study, we investigate the possible climate impacts of a descent into a deep grand solar minimum over the 21st century using a comprehensive stratosphere-resolving coupled atmosphere-ocean climate model. Whilst some studies have focused on the surface response to a grand solar minimum like forcing, we provide further context by analyzing the effects on the stratosphere and their relationship to the surface changes. The focus of this study is on changes in the stratospheric and tropospheric circulations. A separate paper [Ineson *et al.* [2015]] examines the European winter-time surface response in more detail. The remainder of the paper is structured as follows: Section 2 describes the model and experiments carried out, Section 3 describes the results of the core solar minimum experiment, Section 4 assesses the role of a top-down mechanism for enhanced regional effects, and Section 5 summarises our findings.

2. Methods

2.1. The global climate model (GCM)

Experiments have been conducted using the Met Office's 'high-top' HadGEM2-CC climate model, which is one configuration of the HadGEM2 model suite [Martin *et al.*, 2011]. The model is described in detail by Hardiman *et al.* [2012] and Osprey *et al.* [2013], and participated in the Coupled Model Intercomparison Project Phase 5 (CMIP5; Jones *et al.* [2011]). The model has 60 levels in the vertical domain with an upper boundary at ~ 84 km and is run at N96 horizontal resolution ($1.250^\circ \times 1.875^\circ$). It includes orographic and non-orographic gravity wave drag schemes and simulates a realistic quasi-biennial oscillation (QBO) [Scaife *et al.*, 2002]. The atmosphere is coupled to the Hadley Centre ocean model, which has 40 vertical layers and $1^\circ \times 1^\circ$ resolution (increasing in the tropics), and sea-ice scheme as described by Johns *et al.* [2006]. The model also includes an interactive carbon cycle. The atmospheric model uses the Edwards and Slingo [1996] radiative transfer scheme, which has been updated to use the correlated-k method for calculating transmittances [Cusack *et al.*, 1999]. In the configuration used here, the radiation code has six bands in the shortwave spectral region covering the intervals 200-320 nm, 320-690 nm (ozone only), 320-690 nm (ozone and water vapour), 690-1190 nm, 1190-2380 nm and 2380-10000 nm. The radiation scheme also employs updates to the treatment of shortwave absorption by ozone as described by Zhong *et al.* [2008].

Each experiment consists of a three member ensemble run from 1 December 2005 to 1 January 2100 with atmospheric and oceanic initial conditions taken from three 'historical' all-forcings HadGEM2-CC simulations. All experiments include time-varying well-mixed greenhouse gases (CO_2 , CH_4 , N_2O , chlourofluorocarbons) and aerosols as specified by the

Representative Concentration Pathway 8.5 (RCP8.5) scenario (Meinshausen *et al.* [2011]).

This is a high greenhouse gas forcing scenario in which atmospheric CO₂ concentrations increase from ~380 ppm in 2005 to ~970 ppm in 2100. HadGEM2-CC does not include interactive chemistry, and thus ozone is prescribed as a zonally-averaged latitude-height-time field using the SPARC AC&C ozone dataset (Cionni *et al.* [2011]). This dataset was recommended for use in CMIP5 and includes the recovery of the ozone layer over the 21st century due to declining abundances of ozone depleting substances and a climate change trend according to the SRES A1b greenhouse gas scenario. The original ozone dataset did not include a solar cycle component for the future period, so this was added for the HadGEM2-CC CMIP5 simulations (see Section 2.3 for details). Unless otherwise stated, the figures presented in Sections 3 and 4 show averages over the three ensemble members.

2.2. Specification of TSI and SSI

To explore the possible impacts of a future decline into a grand solar minimum, we use the HadGEM2-CC RCP8.5 experiment submitted to the CMIP5 archive as a baseline (denoted RCP8.5_ref). This experiment assumes a sinusoidal 11-year solar cycle in total solar irradiance (TSI) over the 21st century, with a constant amplitude based on solar cycle 23 and a fixed long-term background (see black line in Figure 1(a)). The spectrally-resolved irradiances are apportioned into the model's radiation bands by integrating the 1 nm fluxes provided for CMIP5 (<http://solarisheppa.geomar.de/cmip5>), which are derived from the Naval Research Laboratory Spectral Solar Irradiance (NRLSSI) model [Wang *et al.*, 2005]. As specified by the dataset, monthly-mean TSI and SSI values are used from 1882 onwards and annual mean values prior to this. The irradiance in the 200-320 nm spectral band in this experiment is shown by the black line in Figure 1(b). For

reference, the 11-year solar max-min change in this spectral band over the 21st century is $\sim 0.7\%$ in the RCP8.5_ref experiment. We note that this represents the smallest change in solar UV irradiance indicated by the current uncertainty range [Ermolli *et al.*, 2013].

The solar perturbation experiment, denoted ‘RCP8.5_solmin’, includes a modified future TSI trend shown by the blue line in Figure 1(a). This scenario is equivalent to the most extreme grand solar minimum case examined by Jones *et al.* [2012] (see their Figure 1) and is based on the analogue forecasts of Barnard *et al.* [2011]. The spectrally-resolved irradiances for this scenario are calculated by extrapolating second-order polynomial regressions of the irradiances in each of the six spectral bands against TSI over the period 1860-2009. The changes in TSI are thus apportioned across the spectrum using the assumption that the NRLSSI spectral data for the historical period would scale for the assumed future solar minimum scenario. The irradiance in the 200-320 nm band in this experiment is shown by the blue line in Figure 1(b). The scenario corresponds to an average reduction in UV irradiance over the period 2050-99 of 0.85% compared to RCP8.5_ref.

2.3. Treatment of solar-ozone response

Since HadGEM2-CC does not include interactive chemistry, stratospheric and tropospheric ozone are prescribed using a modified version of the CMIP5 recommended SPARC AC&C ozone dataset [Cionni *et al.*, 2011]. The modifications include a vertical extrapolation of the ozone data above 1 hPa to coincide with the upper levels of the model domain.

The original CMIP5 ozone dataset did not include a solar cycle component in the future. It has therefore been added by regressing the ozone mixing ratios at each latitude and height for the historical period onto terms representing solar forcing (O_3^{sol}), equiva-

lent effective stratospheric chlorine (O_3^{Cl}), a seasonal cycle (O_3^{seas}), and a residual term (O_3^{res}):

$$O_3(t) = \alpha O_3^{sol}(t) + \beta O_3^{Cl}(t) + O_3^{seas}(t) + O_3^{res}(t) \quad (1)$$

The solar regression term is then added to the ozone field for the future period. A cosine latitude extrapolation of the solar term is also made over high latitudes since the signal in the original dataset only extended to $\pm 60^\circ$ latitude. The magnitude of the solar max-min ozone response in the RCP8.5_ref experiment is $\sim 4\%$ in the tropical upper stratosphere.

Although this magnitude is towards the upper end of estimates from observations, it is still within the plausible range [Gray *et al.*, 2009, 2010].

The solar-ozone response term is also included in the RCP8.5_solmin experiment, with the magnitude adjusted to account for the modified future TSI trend. RCP8.5_solmin therefore includes a representation of the ozone response to solar variability (e.g. Haigh [1994]), which amounts to a decrease in ozone at the tropical stratopause of $\sim 6\%$ for the period 2050-99 (see Figure 2).

A summary of the experimental set-ups is provided in Table 1. For reference, Table 2 gives the differences in shortwave irradiances in the six spectral bands over the period 2050-99. The analysis in Section 3 focuses on the differences between RCP8.5_solmin and RCP8.5_ref for this period and, unless otherwise stated, significance testing is carried out using a two-sided Student's t-test for 3×50 years = 150 years under the assumption that each data point (e.g. a detrended monthly or seasonal mean) can be considered as an independent sample.

In Section 3.3.2, the location of the midlatitude jet is identified by a spline interpolation of the seasonal mean zonal mean zonal wind (\bar{u}) onto a 0.1° latitude grid and locating

the maximum wind speed at 850 hPa between 30-70°. Jet shifts are then computed as the differences between the latitudes of the \bar{u} maxima. The changes in Southern Annular Mode (SAM) index in the same section are measured as the difference in zonal mean mean sea level pressure (MSLP) between 40-60°S and 70-90°S.

3. Results

3.1. Temperature changes

Figure 3(a) shows the differences in annual mean zonal mean shortwave heating rates (K day^{-1}) between the RCP8.5_solmin and RCP8.5_ref experiments for the period 2050-99. The grey shading indicates where the differences are not statistically significant at the 95% confidence level. At the tropical stratopause, the decrease in shortwave heating rates has a peak of $\sim -0.4 \text{ K day}^{-1}$. This localised minimum is partly related to the structure of the imposed ozone changes, which have a peak near the tropical stratopause (see Figure 2). The magnitude of the decrease in heating rate drops off rapidly with increasing latitude and decreasing altitude. There is some hemispheric asymmetry in the heating rate anomalies, with larger changes found in the Northern hemisphere, and also some small localised increases in heating, both of which are also related to the structure of the imposed ozone changes (see Figure 2).

Figure 3(b) shows the corresponding differences in annual mean zonal mean temperature (\bar{T} ; K). The changes in temperature are closely related to the shortwave heating rate response shown in Figure 3(a). There is cooling across most of the stratosphere and mesosphere which peaks at $\sim 1.5 \text{ K}$ near the tropical stratopause. This can be compared to the stratospheric cooling due to climate change in the RCP8.5_ref experiment of $\sim 18 \text{ K}$ at 1 hPa (2060-99 vs. 1960-99). The upper stratospheric cooling is comparable to the

solar max-min temperature change found by Frame and Gray [2010], despite the fact that our TSI perturbation is approximately 1.5 times the typical amplitude of the 11-year solar cycle. However, a recent study by Mitchell *et al.* [2014] has found differences in the detailed magnitude and structure of the upper stratospheric temperature response to the 11-year solar cycle across multiple reanalysis datasets. There is also the suggestion of a weak secondary temperature maximum in the tropical lower stratosphere, similar to that identified in several reanalysis datasets [Crooks and Gray [2005]; Frame and Gray [2010], Mitchell *et al.* [2014]]. Gray *et al.* [2009] suggested that the 11-year cycle in ozone may be an important factor in determining the structure of the temperature response in the tropical lower stratosphere. This would appear to be consistent with the imposed changes in ozone, which include a decrease in tropical lower stratospheric ozone. However, there are substantial uncertainties in estimates of the structure and amplitude of the lower stratospheric temperature signal because this region is strongly influenced by QBO variability and volcanic eruptions and the observational record is not long enough to adequately separate the signals [Chiodo *et al.*, 2014].

Figure 4 shows the vertical profile of differences in annual and global mean \bar{T} (K). The maximum cooling occurs at 1 hPa with a magnitude of 1.2 K and decreases rapidly in magnitude above and below this level. From 10-50 hPa the cooling is roughly constant in height with a magnitude of ~ 0.3 K. There is cooling throughout the troposphere, which increases with altitude from ~ 0.1 K at the surface to ~ 0.25 K near the tropopause. The change in global mean 1.5 m temperature ($\bar{T}_{1.5m}$) for the period 2050-99 is -0.13 K. This is broadly consistent with the energy balance model results of Jones *et al.* [2012].

The results in this section show that evolving into a grand solar minimum over the 21st century has the potential to enhance stratospheric cooling trends due to increasing carbon dioxide concentrations, but as has been highlighted in other recent studies (Jones *et al.* [2011]; Meehl *et al.* [2013]), such a decline would have only a small impact on any anthropogenic global warming trend.

3.2. Stratospheric changes

A solar cycle influence on the high latitude stratosphere has been identified in reanalysis data and climate models (e.g. Kuroda and Kodera [2002], Matthes *et al.* [2006], Ineson *et al.* [2011], Mitchell *et al.* [2014]). Given the inter-hemispheric differences in the generation of planetary wave activity in the troposphere, which partly determines the mean strength and unforced variability of the winter polar vortices, it is perhaps unsurprising that the dynamical responses to external forcings, such as the QBO and solar variability, tend to be different in the two hemispheres (e.g. Anstey and Shepherd [2014]). In the Northern hemisphere (NH), studies have shown a time-averaged solar cycle signal in the high latitude stratosphere consisting of a poleward and downward propagation of zonal wind and temperature anomalies over the winter season [Kuroda and Kodera, 2002]. The main mechanism proposed to explain the propagation and amplification of these anomalies invokes wave-mean-flow interactions (e.g. Kodera *et al.* [2003]; Ineson *et al.* [2011]). In contrast, the extratropical circulation response to an external forcing in the Southern hemisphere is often manifested around the time of the spring break-up of the polar vortex (e.g. Kuroda and Kodera [2005]). We now discuss the stratospheric circulation response to the imposed decline in solar activity.

3.2.1. Northern hemisphere (NH)

Figures 5(a-e) show monthly mean \bar{T} (K) differences between the RCP8.5_solmin and RCP8.5_ref experiments for October to February averaged over the period 2050-99. The shading is as in Figure 3. In the NH, there is a relative warming of the Arctic lower stratosphere in February. Since the direct radiative tendency of the decrease in solar irradiance would be to cool the stratosphere, the high latitude warming is indicative of a dynamical response to the solar perturbation. Further analysis (not shown) shows that there is an increase in wave driving (i.e. Eliassen Palm flux divergence) in the high latitude mid and lower stratosphere (cf. Ineson *et al.* [2011]), particularly in January, and the associated dynamical heating acts against the radiatively driven cooling.

Figures 5(f-j) show equivalent plots to panels (a-e) for differences in zonal mean zonal wind (\bar{u} ; m s^{-1}). The warming of the Arctic polar vortex in boreal winter is coincident with a weakening of the stratospheric westerly jet. There is an easterly anomaly in the region of the jet core (~ 1 hPa) of up to $3\text{-}4 \text{ m s}^{-1}$ in January-February. A weaker easterly anomaly, more confined to the mid and upper stratosphere and the mesosphere, is also present in October-November, but the differences in December are not highly statistically significant, probably in part due to the large inter-annual variability during NH midwinter.

It has been suggested that changes in solar irradiance may impact on the timing of major sudden stratospheric warming events (SSWs; see e.g. Gray *et al.* [2004]), which occur in the Arctic stratosphere during boreal winter. Thus some of the changes in \bar{u} in Figures 5(f-j) may reflect changes in the frequency or timing of SSWs. Figure 6 shows the winter-time distribution of SSWs in each experiment for the period 2050-99 using the Charlton and Polvani [2007] definition based on a temporary reversal of \bar{u} at 10 hPa

and 60°N to easterlies. The results for the RCP8.5_ref experiment have been previously discussed by Mitchell *et al.* [2012]. The histograms suggest that, although the average number of SSWs in RCP8.5_solmin remains similar to that in RCP8.5_ref (0.83 yr⁻¹), there is a slight decrease in the occurrence of SSWs in March and an increase in January-February. However, these changes are not highly statistically significant according to the t-test formulated by Charlton *et al.* [2007], which partly reflects the large inter-decadal variability in SSWs, as noted by Butchart *et al.* [2000]. Consequently longer simulations would be required to make robust conclusions about whether a decline in solar activity would impact on the frequency or timing of SSWs.

3.2.2. Southern hemisphere (SH)

In the Southern hemisphere (SH), the plots of monthly mean \bar{T} in Figures 5(a-c) show a relative warming of the Antarctic lower stratosphere by up to ~ 1 K during the SH dynamically active season (October-December). This reflects a decrease in the equator-to-pole temperature gradient, and is coincident with a weakening of the climatological westerly jet throughout the stratosphere by up to ~ 3 m s⁻¹ (Figures 5(a-c)). In mid-winter (June-August; JJA), there is an easterly anomaly in the midlatitudes (30-60°S) of up to ~ 3 m s⁻¹ (see Figure 7), which reflects a weakening of the westerlies on the equatorward flank of the stratospheric jet. These changes in circulation which extend throughout the stratosphere are in contrast to the response in austral summer (January-February), where the subtropical easterly anomaly is mainly confined to the upper stratosphere and mesosphere.

3.3. Tropospheric changes

Circulation changes in the stratosphere during the dynamically active seasons, such as those described in Section 3.2, can impact on the underlying troposphere via stratosphere-troposphere dynamical coupling (see e.g. Gerber *et al.* [2012] for an overview). Furthermore, a number of studies have shown a potential influence of solar variability on the tropical Pacific ocean and the El Niño Southern Oscillation (ENSO) (e.g. Meehl *et al.* [2009]). We now discuss the changes in the tropospheric state in the grand solar minimum experiment.

3.3.1. Northern hemisphere

Figure 8(a) shows the seasonal mean tropospheric \bar{u} (m s^{-1}) changes in the NH in DJF. The shading denotes the differences between the RCP8.5_solmin and RCP8.5_ref experiments and the contours show the climatology of the latter for reference. The hatching denotes where the differences are not statistically significant at the 95% confidence level. There is a barotropic dipole change in \bar{u} in the region of the midlatitude jet, with a westerly anomaly between 30-45°N and an easterly anomaly between 60-70°N. This feature shows a peak-to-peak \bar{u} dipole change of 0.36 m s^{-1} at 850 hPa. The dipole \bar{u} response is comparable to the climate change signal in DJF in the NH (2060-99 vs. 1960-99), which shows a strengthening of the westerlies in the jet core by 0.5 m s^{-1} , and a weakening of the westerlies on the poleward flank by $\sim 0.9 \text{ m s}^{-1}$ (not shown). Thus, whilst the impact of the decline in solar activity on global near-surface temperature is relatively small, its effects on the midlatitude circulation amount to a considerable fraction of the uncertainty due to future greenhouse gas trends (see also Ineson *et al.* [2015]). A significant NH tropospheric \bar{u} response is not found outside of boreal winter, which suggests a role for a

top-down influence of changes in the stratospheric circulation on mid and high latitude climate.

Figure 8(b) shows a polar stereographic map of the differences in DJF MSLP (hPa) between the RCP8.5_solmin and RCP8.5_ref experiments. The green lines encompass regions where the differences are statistically significant at the 95% confidence level. A more negative Arctic Oscillation (AO) index is characterised by lower pressure in the midlatitudes and higher pressure over the polar cap, which corresponds to a weakening of the climatological equator-to-pole pressure gradient and anomalously easterly flow across Europe and the Atlantic sector [Thompson and Wallace, 1998]. The pattern in Figure 8(b) suggests a more negative AO index, although the response is not highly statistically significant and the structure over the North Atlantic does not strongly resemble the NAO. There is a deepening of the Aleutian low, which has also been identified during solar minimum conditions in observations (e.g. Roy and Haigh [2010]; Gray *et al.* [2013]).

Blocking episodes have been highlighted as an important aspect of variability in the North Atlantic circulation (e.g. Shabbar *et al.* [2001]; Woollings *et al.* [2010b]). Previous studies have identified variations in blocking frequency associated with the 11-year solar cycle [Barriopedro *et al.*, 2008; Woollings *et al.*, 2010a]. The solar-blocking signal identified in these studies consists of an increase in Euro-Atlantic blocking during solar minimum, with the precise magnitude of the changes being somewhat sensitive to the metric used to define solar activity (e.g. F10.7cm radio flux or Open Solar Flux), but is typically around ~8-10% of total blocked days. Figure 9 shows differences in the ensemble mean DJF blocking frequency (as a % of total blocked days) between the RCP8.5_solmin and RCP8.5_ref experiments. The blocking index used here is based on temporary reversals

in the meridional gradient of potential temperature on the dynamical tropopause which must persist for at least 5 days, and is identical to that used by Woollings *et al.* [2010a] and Anstey *et al.* [2013]. The general pattern of an increase in Euro-Atlantic and Pacific blocking at high latitudes and a decrease over the Mediterranean are consistent with the results of previous studies (e.g. Woollings *et al.* [2010a]), but the magnitude of the difference is several times smaller. Like many CMIP5 models, HadGEM2-CC has biases in its representation of NH blocking, the main features of which are a lack of blocking events at high latitudes and too much blocking at lower latitudes [Anstey *et al.*, 2013]. It is possible that the underlying model biases could impact on the simulation of a solar-blocking connection (e.g. Scaife *et al.* [2011]). However, the reanalysis based studies described above have mostly focused on the late 20th century period, when there was an *immediate* correlation between solar variability and the NAO (and, by proxy, blocking events), but the solar-NAO relationship at *zero lag* has been shown to be considerably weaker over a longer record (Roy and Haigh [2010]). Gray *et al.* [2013] showed that over the longer period 1870-2010 the strongest correlation was at lags of 3-4 years, but the HadGEM2-CC model was unable to reproduce this behaviour, leading Scaife *et al.* [2013] to suggest that there may be deficiencies in the representation of midlatitude ocean-atmosphere coupling in the model. Despite these outstanding questions, our results are consistent with the findings of other studies which have highlighted a solar influence on NH blocking and the NAO.

The results in this section show that there is a coherent change in the NH extratropical circulation in response to the decline in solar activity which extends from the upper stratosphere to the surface (cf. e.g. Ineson *et al.* [2011]; Gray *et al.* [2013]).

3.3.2. Southern hemisphere

As was shown in Figure 5, the SH high latitude zonal wind anomalies in December extend throughout the stratosphere, and are accompanied by dipole changes in the troposphere in the region of the midlatitude jet. Figure 10 shows differences in the seasonal mean \bar{u} (m s^{-1}) between the RCP8.5_solmin and RCP8.5_ref experiments for (a) JJA and (b) October-December (OND) seasons. In JJA, there is a small poleward shift in the midlatitude jet, with a peak-to-peak dipole change in \bar{u} at 850 hPa of 0.32 m s^{-1} . The strongest signal is a weakening of the westerlies on the equatorward flank of the jet. In OND, there is an equatorward jet shift of $\sim 0.5^\circ$ latitude, with a peak-to-peak \bar{u} dipole of 0.87 m s^{-1} .

Projected future trends in the position of the SH midlatitude jet have been shown to be sensitive to the recovery of the Antarctic ozone hole and increases in greenhouse gas concentrations [Son et al., 2008]. In austral winter, the trend in jet position is largely determined by the greenhouse gas forcing [Barnes *et al.*, 2014]. In the baseline RCP8.5 experiment, there is a poleward shift in the jet of 2.5° (2.7 m s^{-1} \bar{u} dipole) in JJA and 2.0° (3.5 m s^{-1} \bar{u} dipole) in OND (2060-2099 vs. 1960-1999). The change in jet latitude in OND is coincident with a more negative Southern Annular Mode (SAM) index of $\sim 1.4 \text{ hPa}$, which offsets the positive SAM trend in this season in the RCP8.5_ref experiment by $\sim 25\%$ (see Figure 11).

Interestingly, the circulation changes in JJA are of the opposite sign to what is typically associated with stratosphere-troposphere dynamical coupling (weaker stratospheric westerlies lead to a more equatorward tropospheric jet). Such a seasonal dependence of the sign of the SH jet shift has been identified in other studies. Varma *et al.* [2011] found

that the response to a constant 2 W m^{-2} reduction in TSI consisted of a poleward jet shift in JJA and an equatorward jet shift in DJF in a coupled model without a well-resolved stratosphere. The annual mean response was dominated by the signal in DJF (i.e. an equatorward jet shift). Thresher [2002] found observational evidence for a seasonal cycle in the SH surface solar response over the late 20th century, which would be consistent with the findings of Varma *et al.* [2011]. However, the findings of modeling studies may be sensitive to e.g. the inclusion of a well-resolved stratosphere, and thus this highlights the need for further research to better understand the response of the SH circulation to stratospheric changes and its dependence on season.

3.3.3. Tropics

In addition to the proposed top-down mechanisms for an amplified surface response to solar forcing in the midlatitudes, some studies have proposed an additional ‘bottom-up’ mechanism operating in the tropical Pacific. This involves coupled air-sea feedbacks in response to small changes in surface heating and results in an anomalous sea surface temperature (SST) pattern that resembles ENSO [White *et al.*, 1997; Meehl *et al.*, 2008, 2009]). However, there has been some disagreement as to whether the observed response corresponds to a warm or cold ENSO phase at solar maximum, and Roy and Haigh [2010] further showed that the apparent solar-ENSO connection could be due to aliasing onto unconnected ENSO variations.

As described in Section 1, Meehl *et al.* [2013] identified an ENSO-like response in model simulations of a persistent solar minimum, with relatively warm east Pacific SSTs during the first decade after a reduction in TSI was imposed, followed by colder SSTs in the second decade. However, the inter-decadal changes were not found to be highly statistically

significant. Figure 12 shows the differences in DJF sea surface temperatures between the RCP8.5_solmin and RCP8.5_ref experiments. There is weak cooling (~ 0.1 K) across much of the tropical Pacific; the change in area averaged (15°N - 15°S , 150°E - 90°W) temperature is -0.075 K. However, there is no indication of a local amplification in the ENSO region. Our simulations therefore do not lend support to the existence of a solar-ENSO connection. This is in contrast to the results of Meehl *et al.* [2013], although their simulated ENSO-like response to a persistent solar minimum was weaker than that found for 11-year cycle variations [Meehl *et al.*, 2009]. However, the experiments do show enhanced cooling over the north Pacific, which is consistent with the deepened Aleutian low (e.g. Schneider and Cornuelle [2005]). There is also a band of stronger cooling across the SH midlatitudes, which may be partly related to changes in the midlatitude jet (see Section 3.3.2).

4. Sensitivity to UV forcing

The RCP8.5_solmin experiment shows enhanced regional surface responses to a decline in solar activity, particularly in the mid and high latitudes. As described in Section 1, one proposed mechanism for such localised effects involves the impact of changes in shortwave heating rates on the stratospheric circulation and subsequent surface impacts via stratosphere-troposphere coupling. The potential for this mechanism to contribute to the response to solar forcing in the Northern hemisphere was demonstrated by Ineson *et al.* [2011]. They imposed a perturbation in 200-320 nm radiation (i.e. in UV radiation alone) in a model and found a more negative NAO index under solar minimum conditions.

To identify whether a similar mechanism may also be operating here, we conduct a further experiment (RCP8.5_uvmin) in which the 200-320 nm irradiance is reduced by

6.4% in isolation of any other changes (e.g. ozone, visible radiation); this enables a separation of a pure ‘top-down’ influence from a decline in solar activity. This is a highly idealised experiment, in which the imposed UV perturbation is considerably larger than in RCP8.5_solmin, and other effects, such as the solar-ozone response, are neglected. Nevertheless, it allows us to make an assessment of at least one pathway that may be contributing to the RCP8.5_solmin results discussed in the previous sections and to elucidate more generally the role of the top-down pathway for solar-climate coupling. The same experimental protocol as described in Section 2 is carried out, with a reduction in 200-320 nm radiation (and by definition in TSI) of 1.75 W m^{-2} over the 2050-99 period.

Figure 13(a) shows the December-February mean difference in \bar{u} between the RCP8.5_uvmin and the RCP8.5_ref experiments over 2050-99. There is a weakening of the stratospheric jet by up to 3 m s^{-1} and poleward shift in the tropospheric jet. This is qualitatively similar to the response in the RCP8.5_solmin experiment (Figures 5 and 8), but about 20-30% larger, and is consistent with a top-down pathway which contributes to the amplified regional climate responses.

Figure 13(b) shows the DJF mean sea level pressure differences between the RCP8.5_uvmin and RCP8.5_ref experiments. The changes in tropospheric \bar{u} are commensurate with a more negative NAO index and a deepening of the Aleutian low, again with the amplitude of the changes being slightly larger than in RCP8.5_solmin. Conversely in the SH (not shown), there is no indication of an enhanced tropospheric response during OND, as was found in RCP8.5_solmin in Section 3.3.2.

These results suggest that changes in UV irradiance and a top-down influence are likely to be contributing to the enhanced NH surface response discussed in Section 3.3, but that

the enhanced response in the SH in OND may be related to other processes such as the ozone response or changes in visible irradiance. Future studies should therefore aim to elucidate the roles of these mechanisms in driving the SH response to solar forcing.

5. Summary and discussion

A comprehensive coupled atmosphere-ocean global climate model with a well-resolved stratosphere (HadGEM2-CC) has been used to investigate the possible impacts of evolving into a period of very low solar activity over the 21st century. The assumed scenario is akin to what may have occurred during the Maunder Minimum (MM) in the late 17th century. The RCP8.5_solmin experiment assumes a mean decrease in TSI of $\sim 0.12\%$ over the second half of the 21st century and includes a change in UV irradiance (200-320 nm) of -0.85% , along with a representation of the solar cycle impact on stratospheric ozone.

The key conclusions of the study for projections of global mean climate are:

- A return to MM-like levels of solar activity would enhance the anticipated stratospheric cooling trend due to increasing atmospheric carbon dioxide concentrations. The maximum cooling at the stratopause is ~ 1.2 K, which can be compared to the projected cooling due to climate change in the RCP8.5 scenario of ~ 18 K.
- The change in global mean near-surface temperature over the second half of the 21st century is $O(0.1$ K), confirming the findings of earlier studies which have shown that a large decrease in solar activity would do little to offset the projected anthropogenic global warming trend (cf. Feulner and Rahmstorf [2010]; Jones *et al.* [2011]; Meehl *et al.* [2013]; Anet *et al.* [2013]).

In the NH during boreal winter, the main features of the response to the solar minimum consist of:

• A warmer polar lower stratosphere and slight weakening ($<4\text{ m s}^{-1}$) of the polar vortex, with the largest changes occurring in January-February.

• Dipole changes in NH \bar{u} in DJF in the region of the midlatitude jet. The changes in the large-scale circulation suggest a more negative Arctic Oscillation index, but the pattern over the North Atlantic does not strongly resemble the NAO.

• Changes in the occurrence of NH tropospheric blocking events, with an increase over Northern Europe and the North Pacific and a decrease over Southern Europe. The magnitude of this change is smaller than has been suggested in studies using reanalysis data for the recent past, but the patterns are similar [Woollings *et al.*, 2010a].

• A further sensitivity experiment which only included changes in 200-320 nm (UV) radiation indicates that the enhanced NH regional responses are at least partly driven by changes in UV irradiance and a top-down pathway.

A separate paper [Ineson *et al.* [2015]] describes the European winter-time surface response and land surface temperature changes in more detail.

In the SH during austral winter and spring, we find that the decrease in solar activity leads to:

• A relative warming of the Antarctic stratosphere during June-December. This is coincident with a weakening of the background stratospheric westerly jet of up to 3 m s^{-1} .

• An equatorward shift in the Southern hemisphere (SH) tropospheric midlatitude jet by $\sim 0.5^\circ$ and a more negative Southern Annular Mode index of $\sim 1.4\text{ hPa}$ in October-December (OND).

Finally, in contrast to earlier studies (e.g. Meehl *et al.* [2009]), we find no evidence of an enhanced sea surface temperature response over the tropical Pacific that would be

suggestive of an impact on ENSO. Our experiment therefore does not lend support to the existence of a solar-ENSO connection.

It is projected that over the 21st century there will be significant changes in the tropospheric circulation due to the combined effects of ozone recovery and increasing greenhouse gas concentrations (e.g. Wilcox *et al.* [2012], Scaife *et al.* [2012]). Our experiment has shown that, although any impact on global mean surface temperature can be expected to be small, uncertainties in future solar forcing should be considered in projections of regional high latitude climate change. It is also important to note that, although some studies have presented arguments for a future decline in solar output (e.g. Barnard *et al.* [2011]; Abreu *et al.* [2008]), the CMIP5 integrations assumed no trend in solar activity over the future period. It is therefore important that more scenarios which reflect the range of possible future changes in solar activity should be generated for use in studies of 21st century climate. We further emphasise that the recommended representation of spectral solar irradiance in CMIP5 was based on the Wang *et al.* [2005] dataset, which is at the lower end of the estimated range for UV variability [Ermolli *et al.*, 2013]. We therefore highlight the need for alternate scenarios which better reflect the current understanding of SSI variability for use in future model intercomparisons.

Acknowledgments. LJG and ACM were supported by the National Centre for Atmospheric Science's Climate Directorate. ACM also acknowledges support from the ERC ACCI project no. 267760 and an AXA Postdoctoral Fellowship. SI and AAS were supported by the Joint DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). LJG and JAA were supported by a grant from the National Environmental Research Council. The model runs were conducted at the University of Oxford and the

UK Met Office using the Met Office MONSooN supercomputer: the data are not currently available to external users.

References

- Abreu, J. A., Beer, J., Steinhilber, F., Tobias, S. M., , and Weiss, N. O. (2008). For how long will the current grand maximum of solar activity persist? *Geophys. Res. Lett.*, **35**, L20109.
- Anet, J. G., Rozanov, E. V., Muthers, S., Peter, T., Brönnimann, S., Arfeuille, F., Beer, J., Shapiro, A. I., Raible, C. C., Steinhilber, F., and Schmutz, W. K. (2013). Impact of a potential 21st century ‘grand solar minimum’ on surface temperatures and stratospheric ozone. *Geophys. Res. Lett.*, **40**, 4420–4425.
- Anstey, J. and Shepherd, T. G. (2014). High-latitude influence of the quasi-biennial oscillation. *Q. J. Roy. Met. Soc.*, **140**, 1–21.
- Anstey, J. A., Davini, P., Gray, L. J., Woollings, T. J., Butchart, N., Cagnazzo, C., Christiansen, B., Hardiman, S. C., Osprey, S. M., and Yang, S. (2013). Multi-model analysis of Northern Hemisphere winter blocking: Model biases and the role of resolution. *J. Geophys. Res.*, **118**, 3956–3971.
- Barnard, L., Lockwood, M., Hapgood, M. A., Owens, M. J., Davis, C. J., and Steinhilber, F. (2011). Predicting space climate change. *Geophys. Res. Lett.*, **38**, L16103.
- Barnes, E. A., Barnes, N. W., and Polvani, L. M. (2014). Delayed southern hemisphere climate change induced by stratospheric ozone recovery, as projected by the CMIP5 models. *J. Clim.*, **27**, 852–867.

- Barriopedro, D., Garcia-Herrera, R., and Huth, R. (2008). Solar modulation of Northern Hemisphere winter blocking. *J. Geophys. Res.*, **113**, D14118.
- Butchart, N., Austin, J., Knight, J. R., Scaife, A. A., and Gallani, M. L. (2000). The response of the stratospheric climate to projected changes in the concentrations of well-mixed greenhouse gases from 1992 to 2051. *J. Clim.*, **13**, 2142–2159.
- Charlton, A. J. and Polvani, L. (2007). A new look at stratospheric sudden warmings. Part I. Climatology and modeling benchmarks. *J. Clim.*, **20**, 449–469.
- Charlton, A. J., Polvani, L. M., Perlwitz, J., Sassi, F., Manzini, E., Shibata, K., Pawson, S., Nielsen, J. E., and Rind, D. (2007). A new look at stratospheric sudden warmings. Part II. Evaluation of numerical model simulations. *J. Climate*, **20**, 470–488.
- Chiodo, G., Marsh, D. R., Garcia-Herrera, R., Calvo, N., and Garcia, J. A. (2014). On the detection of the solar signal in the tropical stratosphere. *Atmos. Chem. Phys.*, **14**, 5251–5269.
- Cionni, I., Eyring, V., Lamarque, J.-F., Randel, W. J., Stevenson, D. S., Wu, F., Bodeker, G. E., Shepherd, T. G., Shindell, D. T., and Waugh, D. W. (2011). Ozone database in support of CMIP5 simulations: Results and corresponding radiative forcing. *Atmos. Chem. Phys.*, **11**, 11,267–11,292.
- Crooks, S. and Gray, L. J. (2005). Characterization of the 11-year solar signal using a multiple regression analysis of the ERA-40 dataset. *J. Clim.*, **18**, 996–1015.
- Cusack, S., Edwards, J. M., and Crowther, M. J. (1999). Investigating k distribution methods for parameterizing gaseous absorption in the Hadley Centre Climate Model. *J. Geophys. Res.*, **104**, 2051–2057.

- Edwards, J. M. and Slingo, A. (1996). Studies with a flexible new radiation code. I: Choosing a configuration for a large-scale model. *Q. J. Roy. Meteorol. Soc.*, **122**, 689–719.
- Ermolli, I., Matthes, K., de Wit, T. D., Krivova, N. A., Tourpali, K., Weber, M., Unruh, Y. C., Gray, L., Langematz, U., Pilewskie, P., Rozanov, E., Schmutz, W., Shapiro, A., Solanki, S. K., Thuillier, G., and Woods, T. N. (2013). Recent variability of the solar spectral irradiance and its impact on climate modelling. *Atmos. Chem. Phys.*, **13**, 3945–3977.
- Feulner, G. and Rahmstorf, S. (2010). On the effect of a new grand minimum of solar activity on the future climate on Earth. *Geophys. Res. Lett.*, **37**, L05707.
- Frame, T. H. A. and Gray, L. J. (2010). The 11-yr solar cycle in ERA-40 data: An update to 2008. *J. Climate*, **23**, 2213–2222.
- Gerber, E. P., Butler, A., Calvo, N., Charlton-Perez, A. J., Giorgetta, M., Manzini, E., Perlwitz, J., Polvani, L. M., and Sassi, F. (2012). Assessing and understanding the impact of stratospheric dynamics and variability on the Earth system. *Bull. Amer. Met. Soc.*, **93**, 845–859.
- Gray, L. J., Crooks, S., Pascoe, C., Sparrow, S., and Palmer, M. (2004). Solar and QBO influences on the timing of stratospheric sudden warmings. *J. Atmos. Sci.*, **61**, 2776–2796.
- Gray, L. J., Rumbold, S., and Shine, K. P. (2009). Stratospheric temperatures and radiative forcing response to 11-year solar cycle changes in irradiance and ozone. *J. Atmos. Sci.*, **66**, 2402–2417.

Gray, L. J., Beer, J., Geller, M., Haigh, J. D., Lockwood, M., Matthes, K., Cubasch, U., Fleitmann, D., Harrison, G., Hood, L., Luterbacher, J., Meehl, G. A., Shindell, D., van Geel, B., and White, W. (2010). Solar influences on climate. *Rev. Geophys.*, **48**, RG4001.

Gray, L. J., Scaife, A. A., Mitchell, D. M., Osprey, S. M., Ineson, S., Hardiman, S., Butchart, N., Knight, J., Sutton, R., and Kodera, K. (2013). A lagged response to the 11-year solar cycle in observed winter Atlantic/European weather patterns. *J. Geophys. Res.*, **118**, 13,405–13,420.

Haigh, J. D. (1994). The role of stratospheric ozone in modulating the solar radiative forcing of climate. *Nature*, **370**, 544–546.

Haigh, J. D. (1996). The impact of solar variability on climate. *Science*, **272**, 981–984.

Haigh, J. D., Blackburn, M., and Day, R. (2005). The response of tropospheric circulation to perturbations in lower-stratospheric temperature. *J. Climate*, **18**, 3672–3685.

Hardiman, S. C., Butchart, N., Osprey, S. M., Gray, L. J., Bushell, A. C., and Hinton, T. J. (2012). The effect of a well-resolved stratosphere on surface climate: Differences between CMIP5 simulations with high and low top versions of the Met Office climate model. *J. Climate*, **25**, 7083–7099.

Ineson, S., Scaife, A., Knight, J. R., Manners, J. C., Dunstone, N. J., Gray, L. J., and Haigh, J. D. (2011). Solar forcing of winter climate variability in the Northern Hemisphere. *Nature Geoscience*, **4**, 753–757.

Johns, T. C., Durman, C. F., Banks, H. T., Roberts, M. J., McLaren, A. J., Ridley, J. K., Senior, C. A., Williams, K. D., Jones, A., Rickard, G. J., Cusack, S., Ingram, W. J., Crucifix, M., Sexton, D. M. H., Joshi, M. M., Dong, B.-W., Spencer, H., Hill, R. S. R.,

Gregory, J. M., Keen, A. B., Pardaens, A. K., Lowe, J. A., Bodas-Salcedo, A., Stark, S.,
, and Searl, Y. (2006). The new Hadley Centre climate model (HadGEM1): Evaluation
of coupled simulations. *J. Climate*, **19**, 1327–1353.

Jones, C. D., Hughes, J. K., Bellouin, N., Hardiman, S. C., Jones, G. S., Knight, J., Lid-
dicat, S., O'Connor, F. M., Andres, R. J., Bell, C., Boo, K.-O., Bozzo, A., Butchart,
N., Cadule, P., Corbin, K. D., Doutriaux-Boucher, M., Friedlingstein, P., Gornall, J.,
Gray, L., Halloran, P. R., Hurtt, G., Ingram, W. J., Lamarque, J.-F., Law, R. M.,
Meinshausen, M., Osprey, S., Palin, E. J., Chini, L. P., Raddatz, T., Sanderson, M. G.,
Sellar, A. A., Schurer, A., Valdes, P., Wood, N., Woodward, S., Yoshioka, M., and
Zerroukat, M. (2011). The HadGEM2-ES implementation of CMIP5 centennial simu-
lations. *Geosci. Model Dev.*, **4**, 543–570.

Jones, G. S., Lockwood, M., and Stott, P. A. (2012). What influence will future solar
activity changes over the 21st century have on projected global near surface temperature
changes? *J. Geophys. Res.*, **117**, D05103.

Kodera, K., Matthes, K., Shibata, K., Langematz, U., and Kuroda, Y. (2003). Solar
impact on the lower mesospheric subtropical jet: A comparative study with general
circulation model simulations. *Geophys. Res. Letts.*, **30**(6), 1315.

Kuroda, Y. and Kodera, K. (2002). Effect of solar activity on the Polar-night Jet Os-
cillation in the Northern and Southern hemisphere winter. *J. Met. Soc. Japan*, **80**,
973–984.

Kuroda, Y. and Kodera, K. (2005). Solar cycle modulation of the Southern Annular
Mode. *J. Geophys. Res.*, **32**, L13802.

Lean, J. (2000). Evolution of the Sun's Spectral Irradiance since the Maunder Minimum. *Geophys. Res. Lett.*, **27**, 2425–2428.

Lockwood, M. (2010). Solar change and climate: an update in the light of the current exceptional solar minimum. *Proc. R. Soc. A*, **466**, 303–329.

Lockwood, M. (2011). Was spectral solar irradiance lower during the recent low sunspot minimum? *J. Geophys. Res.*, **116**, D16103.

Lockwood, M., Harrison, R. G., Woollings, T., and Solanki, S. K. (2010). Are cold winters in Europe associated with low solar activity? *Environ. Res. Lett.*, **5**, 024001.

Marshall, J., Kushnir, Y., Battisti, D., Chang, P., Czaja, A., Dickson, R., McCartney, M., Saravanan, R., and Visbeck, M. (2001). North Atlantic Climate Variability: phenomena, impacts and mechanisms. *Inter. Jour. Climatology*, **21**, 1863–1898.

Martin, G. M., Bellouin, N., Collins, W. J., Culverwell, I. D., Halloran, P. R., Hardiman, S. C., Hinton, T. J., Jones, C. D., McDonald, R. E., McLaren, A. J., O'Connor, F. M., Roberts, M. J., Rodriguez, J. M., Woodward, S., Best, M. J., Brooks, M. E., Brown, A. R., Butchart, N., Dearden, C., Derbyshire, S. H., Dharsssi, I., Doutriaux-Boucher, M., Edwards, J. M., Falloon, P. D., Gedney, N., Gray, L. J., Hewitt, H. T., Hobson, M., Huddleston, M. R., Hughes, J., Ineson, S., Ingram, W. J., James, P. M., Johns, T. C., Johnson, C. E., Jones, A., Jones, C. P., Joshi, M. M., Keen, A. B., Liddicoat, S., Lock, A. P., Maidens, A. V., Manners, J. C., Milton, S. F., Rae, J. G. L., Ridley, J. K., Sellar, A., Senior, C. A., Totterdell, I. J., Verhoef, A., Vidale, P. L., and Wiltshire, A. (2011). The HadGEM2 family of Met Office Unified Model Climate configurations. *Geosci. Model Dev.*, **4**, 723–757.

- Matthes, K., Kuroda, Y., Kodera, K., and Langematz, U. (2006). Transfer of the solar signal from the stratosphere to the troposphere: Northern winter. *J. Geophys. Res.*, **111**, D06108.
- Meehl, G. A., Arblaster, J. M., Branstator, G. W., and van Loon, H. (2008). A coupled air-sea response mechanism to solar forcing in the Pacific region. *J. Climate*, **21**, 2883–2897.
- Meehl, G. A., Arblaster, J. M., Matthes, K., Sassi, F., and van Loon, H. (2009). Amplifying the Pacific climate system response to a small 11-year solar cycle forcing. *Science*, **325**, 1114.
- Meehl, G. A., Arblaster, J. M., and Marsh, D. R. (2013). Could a future grand solar minimum like the maunder minimum stop global warming? *Geophys. Res. Letts.*, **40**, 1789–1793.
- Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J.-F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., and Riahi, K. (2011). The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climatic Change*, **109**, 213–241.
- Misios, S. and Schmidt, H. (2012). Mechanisms involved in the amplifications of the 11-year solar cycle signal in the tropical Pacific ocean. *J. Clim.*, **25**, 5102–5118.
- Mitchell, D. M., Gray, L. J., Butchart, N., Hardiman, S., Charlton-Perez, A. J., and Watson, P. (2012). The Effect of Climate Change on the Variability of the Northern Hemisphere Stratospheric Polar Vortex. *J. Atmos. Sci.*, **69**, 2608–2618.
- Osprey, S. M., Gray, L. J., Hardiman, S. C., Butchart, N., and Hinton, T. J. (2013). Stratospheric variability in the 20th century CMIP5 simulations of the Met Office cli-

mate model: High-top versus low-top. *J. Clim.*, **26**, 1595–1606.

Owens, M. J., Lockwood, M., Barnard, L., and Davis, C. J. (2011). Solar cycle 24: Implications for energetic particles and the probability of a new Maunder Minimum.

Geophys. Res. Lett., **38**, L19106.

Reichler, T., Kim, J., Manzini, E., and Kröger, J. (2012). A stratospheric connection to Atlantic climate variability. *Nature Geoscience*, **5**, 783–787.

Ribes, J. C. and Nesme-Ribes, E. (1993). The solar sunspot cycle in the Maunder Minimum AD 1645 to AD 1715. *Astron. Astrophys.*, **276**, 549–563.

Roy, I. and Haigh, J. D. (2010). Solar cycle signals in sea level pressure and sea surface temperature. *Atmos. Chem. Phys.*, **10**, 3147–3153.

Scaife, A. A., Butchart, N., Warner, C. D., and Swinbank, R. (2002). Impact of a spectral gravity wave parameterization on the stratosphere in the Met Office Unified Model. *J.*

Atmos. Sci., **59**, 1473–1489.

Scaife, A. A., Copsey, D., Gordon, C., Harris, C., Hinton, T., Keeley, S., O'Neill, A., Roberts, M., and Williams, K. (2011). Improved Atlantic winter blocking in a climate

model. *Geophys. Res. Letts.*, **38**, L23703.

Scaife, A. A., Spanghel, T., Fereday, D. R., Cubasch, U., Langematz, U., Akiyoshi, H., Bekki, S., Braesicke, P., Butchart, N., Chipperfield, M. P., Gettelman, A., Hardiman,

S. C., Michou, M., Rozanov, E., and Shepherd, T. G. (2012). Climate change projections and stratosphere-troposphere interaction. *Clim. Dyn.*, **38**, 2089–2097.

Scaife, A. A., Ineson, S., Knight, J. R., Gray, L. J., Kodera, K., and Smith, D. M. (2013).

A mechanism for lagged North Atlantic climate response to solar variability. *Geophys.*

Res. Letts., **40**, 434–439.

- Schmidt, H., Rast, S., Bunzel, F., Esch, M., Giorgetta, M., Kinne, S., Krismer, T., Stenchikov, G., Timmreck, C., Tomassini, L., and Walz, M. (2013). Response of the middle atmosphere to anthropogenic and natural forcings in the CMIP5 simulations with the Max Planck Institute Earth system model. *J. Adv. Model. Earth Syst.*, **5**, 98–116.
- Schneider, N. and Cornuelle, B. D. (2005). The forcing of the Pacific Decadal Oscillation. *J. Clim.*, **18**, 4355–4373.
- Seppälä, A., Randall, C. E., Clilverd, M. A., Rozanov, E., and Rodger, C. J. (2009). Geomagnetic activity and polar surface air temperature variability. *J. Geophys. Res.*, **114**, A10312.
- Seppälä, A., Lu, H., Clilverd, M. A., and Rodger, C. J. (2013). Geomagnetic activity signatures in wintertime stratosphere wind, temperature, and wave response. *J. Geophys. Res.*, **118**, 2169–2183.
- Shabbar, A., Huang, J., and Higuchi, K. (2001). The relationship between the wintertime North Atlantic Oscillation and blocking episodes in the North Atlantic. *Int. J. Clim.*, **21**, 355–369.
- Son, S.-W., Polvani, L. M., Waugh, D. W., Akiyoshi, K., Garcia, R., Kinnison, D., Pawson S., Rozanov, E., Shepherd, T. G., Shibata, K. (2008). The Impact of Stratospheric Ozone Recovery on the Southern Hemisphere Westerly Jet. *Science*, **320**, 1486-1489.
- Soukharev, B. E. and Hood, L. L. (2006). Solar cycle variation of stratospheric ozone: Multiple regression analysis of long-term satellite data sets and comparisons with models. *J. Geophys. Res.*, **111**, D20314.

Steinhilber, F., Abreu, J. A., and Beer, J. (2008). Solar modulation during the Holocene.

Astrophys. Space Sci. Trans., **4**, 1–6.

Thompson, D. W. J. and Wallace, J. M. (1998). The Arctic Oscillation signature in the wintertime Geopotential Height and Temperature fields. *Geophys. Res. Lett.*, **25**, 1297–1300.

Thresher, R. E. (2002). Solar correlates of Southern Hemisphere mid-latitude climate variability. *Int. Jour. Clim.*, **22**, 901–915.

Varma, V., Prange, M., Lamy, F., Merkel, U., and Schulz, M. (2011). Solar-forced shifts of the Southern hemisphere westerlies during the Holocene. *Climate of the Past*, **7**, 339–347.

Wang, Y.-M., Lean, J. L., and Shelley, N. R. (2005). Modeling the Sun's magnetic field and irradiance since 1713. *J. Astrophys.*, **625**, 522–538.

White, W. B., Lean, J., Cayan, D. R., and Dettinger, M. D. (1997). Response of global upper ocean temperature to changing solar irradiance. *J. Geophys. Res.*, **102**, 3255–3266.

Wilcox, L. J., Charlton-Perez, A. J., and Gray, L. J. (2012). Trends in Austral jet position in ensembles of high- and low-top CMIP5 models. *J. Geophys. Res.*, **117**, D13115.

Woollings, T. J., Lockwood, M., Masato, G., Bell, C. J., and Gray, L. J. (2010a). Enhanced signature of solar variability in Eurasian winter climate. *Geophys. Res. Lett.*, **37**, L20805.

Woollings, T. J., Hannachi, A., Hoskins, B. J., and Turner, A. (2010b). A regime view of the North Atlantic Oscillation and its response to anthropogenic forcing. *J. Clim.*, **23**, 1291–1307.

Zhong, W. Y., Osprey, S. M., Gray, L. J., and Haigh, J. D. (2008). Influence of the prescribed solar spectrum on calculations of atmospheric temperature. *Geophys. Res. Lett.*, **35**, L22813.

Accepted Article

Table 1. Details of the solar irradiance and prescribed ozone fields used in the three sets of experiments described in this study.

Experiment	Solar Irradiance	Prescribed ozone
RCP8.5_ref	Assumes a constant amplitude sinusoidal 11-year solar cycle over 21st century. Following CMIP5 recommendations, SSI is specified according to Wang <i>et al.</i> [2005].	CMIP5 recommended SPARC ozone dataset [Cionni <i>et al.</i> , 2011] with solar cycle regression term included (see Osprey <i>et al.</i> 2013).
RCP8.5_solmin	Same as in RCP8.5_ref, but assumes a large ($\sim 0.12\%$) transient decrease in TSI over the 21st century which is distributed across the model's 6 spectral bands.	Same as in RCP8.5_ref, but solar cycle regression term altered to be consistent with assumed future TSI trend.

Table 2. Differences in irradiances for the six HadGEM2-CC radiation bands averaged over the period 2050-99. Units are W m^{-2} .

	200-320 nm	320-690 nm (ozone only)	320-690 nm (ozone and water vapour)	690-1190 nm	1190- 2380 nm	2380- 10000 nm	TSI
RCP8.5_solmin minus RCP8.5_ref [W m^{-2}]	-0.23	-0.39	-0.30	-0.42	-0.24	-0.05	-1.63




Figure 1. Timeseries of (a) TSI (W m^{-2}) and (b) the irradiance in the 200-320 nm spectral band for the RCP8.5.ref (black) and RCP8.5.solmin (blue) cases. The mean % differences for the period 2050-99 are shown in the legends.

Difference in annual-mean ozone [%] RCP8.5_solmin - RCP8.5_ref, 2050-2099




Figure 2. The % differences in ozone between the RCP8.5_solmin and the RCP8.5_ref experiments for the period 2050-99. The changes are based on the CMIP5 recommended ozone database described by Cionni *et al.* [2011] (see also Schmidt *et al.* [2013]). No changes are imposed at pressures higher than ~100 hPa.




Figure 3. The difference in (a) annual mean zonal mean shortwave heating rates (K day^{-1}) and (b) annual mean zonal mean temperature (\bar{T}) between the RCP8.5_solmin and RCP8.5_ref experiments for the period 2050-99. Grey shading indicates where the differences are not statistically significant at the 95% confidence level. The contour interval is 0.1 K day^{-1} and 0.25 K, respectively.




Figure 4. The difference in global mean annual mean \bar{T} (K) between the RCP8.5_solmin and RCP8.5_ref experiments for the period 2050-99. The dashed lines show the 95% confidence intervals.




Figure 5. The difference in monthly mean (a-e) \bar{T} (K) and (f-j) \bar{u} (ms^{-1}) for the period 2050-99 between the RCP8.5_solmin and RCP8.5_ref experiments. The panels show data for October, November, December, January and February, respectively. The grey shading denotes regions where the differences are not statistically significant at the 95% confidence level.




Figure 6. The winter-time distribution of Arctic stratospheric sudden warming events (yr^{-1}) in the RCP8.5_ref (black) and RCP8.5_solmin (blue) experiments for the period 2050-99.




Figure 7. As in Figures 5(f-j) but for the Southern hemisphere in May, June, July, August and September, respectively.




Figure 8. The differences in seasonal mean DJF (a) \bar{u} (m s^{-1}) and (b) MSLP (hPa) for the period 2050-99 between the RCP8.5_solmin and RCP8.5_ref experiments. Note that the shading intervals are not constant and that data are only shown for the Northern hemisphere. The solid contours in (a) denote the 2050-99 \bar{u} climatology in the RCP8.5_ref experiment and are marked at $\pm 5, 10, 20, 30, 40$ and 50 m s^{-1} . The hatching in (a) denotes regions where the differences are not statistically significant at the 95% confidence level. The green lines in (b) encompass regions where the differences are statistically significant at the 95% confidence level.

RCP8.5_solmin - RCP8.5_ref

DJF Δ blocking, 2050-2099




Figure 9. The difference in ensemble mean DJF blocking frequency (as a % of blocked days) between the RCP8.5_solmin and RCP8.5_ref experiments for the period 2050-99. Blocking events are defined using a metric based on potential temperature on the dynamical tropopause and is consistent with that used by Woollings *et al.* [2010a]. The stippling indicates where the differences are significant at the 95% confidence level.




Figure 10. As in Figure 8, but for the Southern hemisphere (SH) for (a) JJA and (b) OND seasons. The contours denote the 2050-99 climatology in the RCP8.5_ref experiment. The hatching denotes regions where differences are not statistically significant at the 95% confidence level.

Change in SAM index 2060-2099 minus 1960-1999




Figure 11. The difference in SAM index (hPa) between 2060-2099 and 1960-1999 for the RCP8.5 and RCP8.5_solmin experiments compared to the historical experiment. The SAM index is defined as the difference in zonally averaged MSLP between 40-60°S and 70-90°S. The whiskers show 5-95% confidence intervals.




Figure 12. The difference in DJF sea surface temperature (K) for the period 2050-99 between the RCP8.5_solmin and RCP8.5_ref experiments. The solid black contours denote 0.1K intervals. The light grey shading denotes regions that are not statistically significant at the 95% confidence level.




Figure 13. As in Figure 8, but for the differences between the RCP8.5_uvmin and RCP8.5_ref experiments. Note that data are only shown for the Northern hemisphere.