

A moment problem for random discrete measures

Article

Accepted Version

Kondratiev, Y. G., Kuna, T. and Lytvynov, E. (2015) A moment problem for random discrete measures. *Stochastic Processes and their Applications*, 125 (9). pp. 3541-3569. ISSN 0304-4149 doi: <https://doi.org/10.1016/j.spa.2015.03.007> Available at <https://centaur.reading.ac.uk/41058/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1016/j.spa.2015.03.007>

To link to this article DOI: <http://dx.doi.org/10.1016/j.spa.2015.03.007>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A moment problem for random discrete measures

Yuri G. Kondratiev

Fakultät für Mathematik, Universität Bielefeld, Postfach 10 01 31, D-33501 Bielefeld, Germany; NPU, Kyiv, Ukraine

e-mail: kondrat@mathematik.uni-bielefeld.de

Tobias Kuna

University of Reading, Department of Mathematics, Whiteknights, PO Box 220, Reading RG6 6AX, U.K.

e-mail: t.kuna@reading.ac.uk

Eugene Lytvynov

Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, U.K.

e-mail: e.lytvynov@swansea.ac.uk

Corresponding author: Eugene Lytvynov

Department of Mathematics, Swansea University, Singleton Park, Swansea SA2 8PP, U.K.

e-mail: e.lytvynov@swansea.ac.uk

Tel.: +44 1792 602156

AMS 2000 subject classifications. Primary 60G55, 60G57; secondary 44A60, 60G51.

Key words and phrases. Discrete random measure, moment problem, point process, random measure.

Abstract

Let X be a locally compact Polish space. A random measure on X is a probability measure on the space of all (nonnegative) Radon measures on X . Denote by $\mathbb{K}(X)$ the cone of all Radon measures η on X which are of the form $\eta = \sum_i s_i \delta_{x_i}$, where, for each i , $s_i > 0$ and δ_{x_i} is the Dirac measure at $x_i \in X$. A random discrete measure on X is a probability measure on $\mathbb{K}(X)$. The main result of the paper states a necessary and sufficient condition (conditional upon a mild *a priori* bound) when a random measure μ is also a random discrete measure. This condition is formulated solely in terms of moments of the random measure μ . Classical examples of random discrete measures are completely random measures and additive subordinators, however, the main result holds independently of any independence property. As a corollary, a characterisation via a moments is given when a random measure is a point process.

1 Introduction

Let X be a locally compact Polish space, and let $\mathcal{B}(X)$ denote the associated Borel σ -algebra. For example, X can be the Euclidean space \mathbb{R}^d , $d \in \mathbb{N}$. Let $\mathbb{M}(X)$ denote the space of all (nonnegative) Radon measures on $(X, \mathcal{B}(X))$. The space $\mathbb{M}(X)$ is equipped with the vague topology. Let $\mathcal{B}(\mathbb{M}(X))$ denote the Borel σ -algebra on $\mathbb{M}(X)$.

Let us define the *cone of (nonnegative) discrete Radon measures on X* by

$$\mathbb{K}(X) := \left\{ \eta = \sum_i s_i \delta_{x_i} \in \mathbb{M}(X) \mid s_i > 0, x_i \in X \right\}.$$

Here δ_{x_i} denotes the Dirac measure with mass at x_i . In the above representation, the atoms x_i are assumed to be distinct, i.e., $x_i \neq x_j$ for $i \neq j$, and their total number is at most countable. By convention, the cone $\mathbb{K}(X)$ contains the null mass $\eta = 0$, which is represented by the sum over an empty set of indices i . As shown in [9], $\mathbb{K}(X) \in \mathcal{B}(\mathbb{M}(X))$.

A *random measure on X* is a measurable mapping $\xi : \Omega \rightarrow \mathbb{M}(X)$, where (Ω, \mathcal{F}, P) is a probability space, see e.g. [6, 7, 10]. A random measure which takes values in $\mathbb{K}(X)$ with probability one will be called a *random discrete measure*. We will give results which characterize when a random measure is a random discrete measure in terms of its moments.

Let us recall the classical characterization of a completely random measure by Kingman [7, 12]. A random measure ξ is called *completely random* if, for any mutually disjoint sets $A_1, \dots, A_n \in \mathcal{B}(X)$, the random variables $\xi(A_1), \dots, \xi(A_n)$ are independent. Kingman's theorem states that every completely random measure ξ can be represented as $\xi = \xi_d + \xi_f + \xi_r$. Here ξ_d , ξ_f , ξ_r are independent completely random measures such that: ξ_d is a deterministic measure on X without atoms; ξ_f is a random measure with fixed (non-random) atoms, that is there exists a deterministic countable collection of points $\{x_i\}$ in X and non-negative independent random numbers $\{a_i\}$ with $\xi_f = \sum_i a_i \delta_{x_i}$; finally the most essential part ξ_r is an extended marked Poisson process which has no fixed atoms, in particular with probability one ξ_r is of the form $\sum_j b_j \delta_{y_j}$, where $\{b_j\}$ are non-negative random numbers and $\{y_j\}$ are random points in X .

Thus, by Kingman's result a completely random measure is a random discrete measure up to a non-random component. If one drops the assumption that the random measure is completely random, one cannot expect anymore to concretely characterize the distribution of ξ . Thus, a natural appropriate question is to ask when a random measure is a random discrete measure. One may be tempted to replace the assumption of complete randomness by a property of a sufficiently strong decay of correlation. However, the result of this paper shows that such an assumption cannot be sufficient.

Note that, in most interesting examples of completely random measures, the set of atoms of the random discrete measure is almost surely dense in X . A study of countable dense random subsets of X leads to "situations in which probabilistic statements about

such sets can be uninformative” [11], see also [2]. It is the presence of the weights s_i in the definition of a random discrete measure that makes a real difference.

An important characteristic of a random measure is its moment sequence. We say that a random measure ξ has finite moments of all orders if, for each $n \in \mathbb{N}$ and all bounded subsets $A \in \mathcal{B}(X)$,

$$\mathbb{E}[\xi(A)^n] < \infty.$$

Then, the n -th moment measure of ξ is the unique symmetric measure $M^{(n)} \in \mathbb{M}(X^n)$ defined by the following relation

$$\forall A_1, \dots, A_n \in \mathcal{B}(X) : \quad M^{(n)}(A_1 \times \dots \times A_n) := \mathbb{E}[\xi(A_1) \cdots \xi(A_n)].$$

We also set $M^{(0)} := \mathbb{E}(1) = 1$. The $(M^{(n)})_{n=0}^\infty$ is called the *moment sequence of the random measure* ξ .

The main result of this paper is a solution of the following problem: *Assume that ξ is a random measure on X whose moment sequence $(M^{(n)})_{n=0}^\infty$ is known and satisfies a mild a priori bound. Give a necessary and sufficient condition, in terms of the moments $(M^{(n)})_{n=0}^\infty$, for ξ to be a random discrete measure, i.e., for the distribution of the random measure ξ to be concentrated on $\mathbb{K}(X)$.*

As a consequence of our main result we also obtain a solution of the (infinite dimensional) moment problem on $\mathbb{K}(X)$. Since we will only use the distribution of a random measure on $\mathbb{M}(X)$, in what follows, under a random measure we will always understand a probability measure on $\mathbb{M}(X)$, and under a random discrete measure a probability measure concentrated on the subset $\mathbb{K}(X)$.

In Section 4 we state three corollaries of the main result. In Corollary 19, *we give a necessary and sufficient condition for a sequence of Radon measures, $(M^{(n)})_{n=0}^\infty$, to be the moment sequence of a random discrete measure*, cf. Section 4 for details. In Corollary 20, *we give a necessary and sufficient condition, in terms of the moments $(M^{(n)})_{n=0}^\infty$, for a random measure to be a simple point process*. In Corollary 21, we relate the previous corollary to the analogous result in terms of the so-called *generalized correlation functions*. (These results are also conditional upon an *a priori* bound satisfied by $(M^{(n)})_{n=0}^\infty$.)

Our main result is very different in spirit and technique to the known results about the localization of measures on cones. As far as we know, all known techniques require the cone under consideration to be closed, cf. [17], but $\mathbb{K}(X)$ is dense in $\mathbb{M}(X)$, cf. the proof of separability in Proposition A2.5.III in [6].

In order to describe the main result more precisely we have to introduce some further notation. Let $i_1, \dots, i_k \in \mathbb{N}$ with $i_1 + \dots + i_k = n$. Denote by M_{i_1, \dots, i_k} the restriction of $M^{(n)}$ to the following subset of X^n

$$\left\{ \left(\underbrace{x_1, \dots, x_1}_{i_1}, x_2, \dots, \underbrace{x_k, \dots, x_k}_{i_k} \right) \in X^n \mid x_i \neq x_j \text{ for } i \neq j \right\}.$$

Denote by $X_{\hat{0}}^{(k)}$ the collection of points $(x_1, \dots, x_k) \in X^k$ whose coordinates are all different. We consider M_{i_1, \dots, i_k} as a measure on $X_{\hat{0}}^{(k)}$, cf. Section 2 for details.

It is clear that a result of the type we wish to derive can only hold under an appropriate estimate on the growth of the measures $M^{(n)}$. Below we will assume that the following conditions are satisfied, see also Remark 5:

(C1) For each $\Lambda \in \mathcal{B}_c(X)$, there exists a constant $C_\Lambda > 0$ such that

$$M^{(n)}(\Lambda^n) \leq C_\Lambda^n n!, \quad n \in \mathbb{N}. \quad (1)$$

Here $\mathcal{B}_c(X)$ denotes the collection of all sets from $\mathcal{B}(X)$ which have compact closure.

(C2) For each $\Lambda \in \mathcal{B}_c(X)$, there exists a constant $C'_\Lambda > 0$ such that

$$M^{(n)}(\Lambda_{\hat{0}}^{(n)}) \leq (C'_\Lambda)^n n!, \quad \forall n \in \mathbb{N} \quad (2)$$

and for any sequence $\{\Lambda_k\}_{k=1}^\infty \in \mathcal{B}_c(X)$ such that $\Lambda_k \downarrow \emptyset$, we have $C'_{\Lambda_k} \rightarrow 0$ as $k \rightarrow \infty$.

We fix a sequence $(\Lambda_l)_{l=1}^\infty$ of compact subsets of X such that $\Lambda_1 \subset \Lambda_2 \subset \Lambda_3 \subset \dots$ and $\bigcup_{l=1}^\infty \Lambda_l = X$. For example, in the case $X = \mathbb{R}^d$, one may choose $\Lambda_l = [-l, l]^d$.

Theorem 1. *Let μ be a random measure on X , i.e., a probability measure on $(\mathbb{M}(X), \mathcal{B}(\mathbb{M}(X)))$. Assume that μ has finite moments, and let $(M^{(n)})_{n=0}^\infty$ be its moment sequence. Further assume that conditions (C1) and (C2) are satisfied. Then μ is a random discrete measure, i.e., $\mu(\mathbb{K}(X)) = 1$, if and only if the moment sequence $(M^{(n)})_{n=0}^\infty$ satisfies the following conditions:*

(i) For any $n \in \mathbb{N}$, $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$, and $\mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_+^n$, let

$$\xi_{\mathbf{i}}^\Delta = \xi_{i_1, \dots, i_n}^\Delta := \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta). \quad (3)$$

cf. for more details (30). Here $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$.

Then the sequence $(\xi_{\mathbf{i}}^\Delta)_{\mathbf{i} \in \mathbb{Z}_+^n}$ is positive definite, i.e., for $N \in \mathbb{N}$ and any finite sequence of complex numbers indexed by elements of \mathbb{Z}_+^n , $(z_{\mathbf{i}})_{\mathbf{i} \in \mathbb{Z}_+^n, |\mathbf{i}| \leq N}$, we have

$$\sum_{\substack{i_1, \dots, i_n=1 \\ j_1, \dots, j_n=1}}^N \xi_{i_1+j_1, \dots, i_n+j_n}^\Delta z_{i_1, \dots, i_n} \overline{z_{j_1, \dots, j_n}} \geq 0.$$

Here $|\mathbf{i}| := \max\{i_1, \dots, i_n\}$.

(ii) For each $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$ of the form $\Delta = (\Lambda_l)_{\hat{0}}^{(n)}$ with $l \in \mathbb{N}$, set

$$r_i^\Delta := \xi_{i,0,0,\dots,0}^\Delta, \quad i \in \mathbb{Z}_+. \quad (4)$$

Then, for any finite sequence of complex numbers, $(z_n)_{n=0}^N$, we have

$$\sum_{i,j=0}^N r_{i+j+1}^\Delta z_i \bar{z}_j \geq 0, \quad (5)$$

and furthermore

$$\sum_{k=1}^{\infty} (D_{k-1}^\Delta D_k^\Delta)^{-1} \det \begin{bmatrix} r_1^\Delta & r_2^\Delta & \cdots & r_k^\Delta \\ r_2^\Delta & r_3^\Delta & \cdots & r_{k+1}^\Delta \\ \vdots & \vdots & \ddots & \vdots \\ r_k^\Delta & r_{k+1}^\Delta & \cdots & r_{2k-1}^\Delta \end{bmatrix}^2 = \infty, \quad (6)$$

where

$$D_k := \det \begin{bmatrix} r_0^\Delta & r_1^\Delta & \cdots & r_k^\Delta \\ r_1^\Delta & r_2^\Delta & \cdots & r_{k+1}^\Delta \\ \vdots & \vdots & \ddots & \vdots \\ r_k^\Delta & r_{k+1}^\Delta & \cdots & r_{2k}^\Delta \end{bmatrix}, \quad k \in \mathbb{Z}_+.$$

Let us now briefly describe the strategy we follow in this paper. Denote $\mathbb{R}_+ := (0, \infty)$. We introduce a logarithmic metric on \mathbb{R}_+ : for $a, b \in \mathbb{R}_+$, $\text{dist}(a, b) := |\ln(\frac{a}{b})|$. Then \mathbb{R}_+ becomes a locally compact Polish space. Thus, $Y := X \times \mathbb{R}_+$ is also a locally compact Polish space. We consider the configuration space $\Gamma(Y)$, i.e., the space of all locally finite subsets of Y . This space is also equipped with the vague topology. A (simple) point process in Y is a probability measure on $(\Gamma(Y), \mathcal{B}(\Gamma(Y)))$. A point process is (uniquely) characterized by its *correlation measure* (also called the *factorial moments measures*), see e.g. [6].

Let μ be a random discrete measure on X . It is often convenient to interpret μ as a point process in Y . More precisely, take any discrete Radon measure $\eta = \sum_i s_i \delta_{x_i} \in \mathbb{K}(X)$ and set

$$\mathcal{E}\eta := \sum_i \delta_{(x_i, s_i)}.$$

As easily seen $\mathcal{E}\eta \in \Gamma(Y)$. Furthermore, it can be shown that the mapping $\mathcal{E} : \mathbb{K}(X) \rightarrow \Gamma(Y)$ is measurable, see [9]. (Note, however, that the range of the mapping \mathcal{E} is not the whole space $\Gamma(Y)$, see the definition in equation (11) below.) We denote $\nu := \mathcal{E}(\mu)$, i.e., the pushforward of μ under \mathcal{E} . Thus, ν is a point process in Y . Hence, one can study the random discrete measure μ through the point process ν .

Our strategy to solve the main problem is to first construct the point process ν associated to the searched random discrete measure μ . An important step along

the way here is to solve the following problem, which is of independent interest in itself: *How can one recover the correlation measure of the associated point process ν from the moment sequence $(M^{(n)})_{n=0}^{\infty}$ of a random discrete measure μ ?* A solution to this problem is given in Section 2. Our approach is significantly influenced by the paper of Rota and Wallstrom [16], which combines ideas of (stochastic) integration and combinatorics. Additionally, to find the correlation measure of ν concretely, one has to solve a sequence of finite-dimensional moment problems. A solution to the main problem is given in Section 3 and the consequence for the moment problem on $\mathbb{K}(X)$ is discussed in Section 4.

Beside completely random measures or additive subordinators in the case $X = \mathbb{R}_+$ (in particular, Lévy processes which are subordinators), we would like to mention the gamma measure, the spatial version of the gamma process, see e.g. [8, 18, 19] for interesting properties. The gamma measure is the completely random discrete measure μ on $X = \mathbb{R}^d$ for which $\mathcal{E}(\mu) = \nu$ is the Poisson point process in $\mathbb{R}^d \times \mathbb{R}_+$ with intensity measure $dx s^{-1}e^{-s} ds$. Note that Gibbs perturbations of the gamma measure have been studied in [9]. These are also random discrete measures which have a.s. a dense set of atoms.

2 Recovering the correlation measure of ν

A partition of a nonempty set Z is any finite collection $\pi = \{A_1, \dots, A_k\}$, where A_1, \dots, A_k are mutually disjoint nonempty subsets of Z such that $Z = \bigcup_{i=1}^k A_i$. The sets A_1, \dots, A_k are called blocks of the partition π .

For each $n \in \mathbb{N}$, denote by $\Pi(n)$ the set of all partitions of the set $\{1, 2, \dots, n\}$. For each partition $\pi = \{A_1, \dots, A_k\} \in \Pi(n)$, we denote by $X_\pi^{(n)}$ the subset of X^n which consists of all $(x_1, \dots, x_n) \in X^n$ such that, for any $1 \leq i < j \leq n$, $x_i = x_j$ if and only if i and j belong to the same block of the partition π , say A_l . For example, for the so-called zero partition $\widehat{0} = \{\{1\}, \{2\}, \dots, \{n\}\}$, the set $X_{\widehat{0}}^{(n)}$ consists of all points $(x_1, \dots, x_n) \in X^n$ whose coordinates are all different. For the so-called one partition $\widehat{1} = \{\{1, 2, \dots, n\}\}$, the set $X_{\widehat{1}}^{(n)}$ consists of all points $(x_1, \dots, x_n) \in X^n$ such that $x_1 = x_2 = \dots = x_n$. Clearly, the collection of sets $X_\pi^{(n)}$ with π running over $\Pi(n)$ forms a partition of X^n .

Let $m^{(n)}$ be any nonnegative Radon measure on X^n , i.e., $m^{(n)} \in \mathbb{M}(X^n)$. For each partition $\pi \in \Pi(n)$, we denote by $m_\pi^{(n)}$ the restriction of the measure $m^{(n)}$ to the set $X_\pi^{(n)}$. Note that we may also consider $m_\pi^{(n)}$ as a measure on X^n by setting

$$m_\pi^{(n)}(X^n \setminus X_\pi^{(n)}) := 0.$$

Then we get

$$m^{(n)} = \sum_{\pi \in \Pi(n)} m_\pi^{(n)}.$$

Let us fix a partition $\pi = \{A_1, A_2, \dots, A_k\} \in \Pi(n)$. Here and below, we will always assume that the blocks of the partition are enumerated so that

$$\min A_1 < \min A_2 < \dots < \min A_k.$$

We denote by $|\pi|$ the number of blocks in the partition π . We construct a measurable, bijective mapping, for $k = |\pi|$,

$$B_\pi : X_\pi^{(n)} \rightarrow X_{\widehat{0}}^{(k)}$$

as follows. For any $(x_1, \dots, x_n) \in X_\pi^{(n)}$, we set

$$B_\pi(x_1, \dots, x_n) = (y_1, \dots, y_k),$$

where, for $i = 1, 2, \dots, k$, $y_i = x_j$ for a $j \in A_i$ (recall that $x_j = x_{j'}$ for all $j, j' \in A_i$). Note that, if $\pi = \widehat{0}$, then B_π is just the identity mapping. We denote by $B_\pi(m_\pi^{(n)})$ the pushforward of the measure $m_\pi^{(n)}$ under B_π .

Let us now additionally assume that the initial measure $m^{(n)}$ is symmetric, i.e., the measure $m^{(n)}$ remains invariant under the natural action of permutations $\sigma \in \mathfrak{S}_n$ on X^n . (Here \mathfrak{S}_n denotes the symmetric group of degree n .) For a partition π as in the above paragraph, we set, for each $l = 1, 2, \dots, k$, $i_l := |A_l|$, the number of elements of the block A_l . Note that $i_1 + i_2 + \dots + i_k = n$. Since $m^{(n)}$ is symmetric, it is clear that the measure $B_\pi(m_\pi^{(n)})$ is completely identified by the numbers i_1, \dots, i_k . That is, if $\pi' = \{A'_1, \dots, A'_k\}$ is another partition from $\Pi(n)$ with $|A'_l| = i_l$, $l = 1, \dots, k$, then $B_\pi(m_\pi^{(n)}) = B_{\pi'}(m_{\pi'}^{(n)})$. Hence, we will denote

$$m_{i_1, \dots, i_k} := B_\pi(m_\pi^{(n)}), \tag{7}$$

and we may assume, without loss of generality, that in formula (7) the partition $\pi = \{A_1, \dots, A_k\}$ is given by

$$A_1 = \{1, \dots, i_1\}, A_2 = \{i_1 + 1, \dots, i_1 + i_2\}, A_3 = \{i_1 + i_2 + 1, \dots, i_1 + i_2 + i_3\}, \dots \tag{8}$$

Note that, since $m^{(n)}$ is a Radon measure on X^n , each measure m_{i_1, \dots, i_k} is a Radon measure on $X_{\widehat{0}}^{(k)}$, i.e., for each $\Delta \in \mathcal{B}_c(X_{\widehat{0}}^{(k)})$, we have $m_{i_1, \dots, i_k}(\Delta) < \infty$. Here $\mathcal{B}_c(X_{\widehat{0}}^{(k)})$ denotes the collection of all sets $\Delta \in \mathcal{B}(X_{\widehat{0}}^{(k)})$ which have a compact closure in X^k , and $\mathcal{B}(X_{\widehat{0}}^{(k)})$ is the trace σ -algebra of $\mathcal{B}(X^k)$ on $X_{\widehat{0}}^{(k)}$. Thus, a given sequence of symmetric Radon measures $m^{(n)}$ on X^n , $n \in \mathbb{N}$, uniquely identifies a sequence of Radon measures m_{i_1, \dots, i_k} on $X_{\widehat{0}}^{(k)}$, where $i_1, \dots, i_k \in \mathbb{N}$, $k \in \mathbb{N}$. Note that this sequence is symmetric in the entries i_1, \dots, i_k , i.e., for any permutation $\sigma \in \mathfrak{S}_k$,

$$dm_{i_1, \dots, i_k}(x_1, \dots, x_k) = dm_{i_{\sigma(1)}, \dots, i_{\sigma(k)}}(x_{\sigma(1)}, \dots, x_{\sigma(k)}).$$

As easily seen the converse implication is also true, i.e., any sequence of Radon measures m_{i_1, \dots, i_k} on $X_0^{(k)}$, with $i_1, \dots, i_k \in \mathbb{N}$ and $k \in \mathbb{N}$ which is symmetric in the entries i_1, \dots, i_k uniquely identifies a sequence of symmetric Radon measures $m^{(n)}$ on X^n , $n \in \mathbb{N}$.

Let now μ be a random discrete measure on X which has finite moments, and let $(M^{(n)})_{n=0}^\infty$ be its moment sequence. Clearly, each $M^{(n)}$ is a symmetric measure on X^n . Below we will deal with the measures M_{i_1, \dots, i_k} derived from the moment sequence $(M^{(n)})_{n=0}^\infty$.

For each $n \in \mathbb{N}$, we denote by $C_0(X^n)$ the space of all continuous functions on X^n with compact support equipped with the natural topology of uniform convergence on compact sets from X^n . Clearly, for each $f^{(n)} \in C_0(X^n)$, the function

$$\mathbb{M}(X) \ni \eta \mapsto \langle \eta^{\otimes n}, f^{(n)} \rangle := \int_{X^n} f^{(n)}(x_1, \dots, x_n) d\eta(x_1) \cdots d\eta(x_n)$$

is measurable. By the dominated convergence theorem it also holds that

$$\int_{X^n} f^{(n)}(x_1, \dots, x_n) dM^{(n)}(x_1, \dots, x_n) = \int_{\mathbb{M}(X)} \langle \eta^{\otimes n}, f^{(n)} \rangle d\mu(\eta). \quad (9)$$

Consider the locally compact Polish space $Y = X \times \mathbb{R}_+$ (see Introduction), and consider the configuration space $\Gamma(Y)$. Recall that

$$\Gamma(Y) := \{\gamma \subset Y \mid |\gamma \cap \Lambda| < \infty \text{ for each compact } \Lambda \subset Y\}.$$

Here, $|\gamma \cap \Lambda|$ denotes the number of points in the set $\gamma \cap \Lambda$. One usually identifies a configuration $\gamma = \{y_i\} \in \Gamma(Y)$ with a Radon measure $\gamma = \sum_i \delta_{y_i}$. Thus, we get the inclusion $\Gamma(Y) \subset \mathbb{M}(Y)$ and we denote by $\mathcal{B}(\Gamma(Y))$ the trace σ -algebra of $\mathcal{B}(\mathbb{M}(Y))$ on $\Gamma(Y)$.

Denote by $\Gamma_p(Y)$ the set of so-called *pinpointing configurations in Y* . By definition, $\Gamma_p(Y)$ consists of all configurations $\gamma \in \Gamma(Y)$ such that if $(x_1, s_1), (x_2, s_2) \in \gamma$ and $(x_1, s_1) \neq (x_2, s_2)$, then $x_1 \neq x_2$. Thus, a configuration $\gamma \in \Gamma_p(Y)$ cannot contain two points (x, s_1) and (x, s_2) with $s_1 \neq s_2$. For each $\gamma \in \Gamma_p(Y)$ and $\Lambda \in \mathcal{B}_c(X)$, we define a *local mass* by

$$\mathfrak{M}_\Lambda(\gamma) := \int_Y \chi_\Lambda(x) s d\gamma(x, s) = \sum_{(x, s) \in \gamma} \chi_\Lambda(x) s \in [0, \infty]. \quad (10)$$

Here χ_Λ denotes the indicator function of the set Λ . The set of *pinpointing configurations with finite local mass* is then defined by

$$\Gamma_{pf}(Y) := \{\gamma \in \Gamma_p(Y) \mid \mathfrak{M}_\Lambda(\gamma) < \infty \text{ for each compact } \Lambda \subset X\}. \quad (11)$$

As easily seen, $\Gamma_{pf}(Y) \in \mathcal{B}(\Gamma(Y))$ and we denote by $\mathcal{B}(\Gamma_{pf}(Y))$ the trace σ -algebra of $\mathcal{B}(\Gamma(Y))$ on $\Gamma_{pf}(Y)$.

We construct a bijective mapping $\mathcal{E} : \mathbb{K}(X) \rightarrow \Gamma_{pf}(Y)$ by setting, for each $\eta = \sum_i s_i \delta_{x_i} \in \mathbb{K}(X)$, $\mathcal{E}\eta := \{(x_i, s_i)\}$. By [9, Theorem 6.2], we have

$$\mathcal{B}(\Gamma_{pf}(Y)) = \{\mathcal{E}A \mid A \in \mathcal{B}(\mathbb{K}(X))\}.$$

Hence, both \mathcal{E} and its inverse \mathcal{E}^{-1} are measurable mappings.

We denote by $\nu := \mathcal{E}(\mu)$ the pushforward of the measure μ under the mapping \mathcal{E} . Thus ν is a probability measure on $\Gamma_{pf}(Y)$, in particular, it is a point process in Y .

Let $\Gamma_0(Y)$ denote the space of all finite configurations in Y :

$$\Gamma_0(Y) := \{\gamma \subset Y \mid |\gamma| < \infty\}.$$

Note that $\Gamma_0(Y) = \bigcup_{n=0}^{\infty} \Gamma^{(n)}(Y)$, where $\Gamma^{(n)}(Y)$ is the space of all n -point configurations (subsets) in Y . Clearly, $\Gamma_0(Y) \subset \Gamma(Y)$, and we denote by $\mathcal{B}(\Gamma_0(Y))$ the trace σ -algebra of $\mathcal{B}(\Gamma(Y))$ on $\Gamma_0(Y)$. The σ -algebra $\mathcal{B}(\Gamma_0(Y))$ admits the following description: for each $n \in \mathbb{N}$, $\Gamma^{(n)}(Y) \in \mathcal{B}(\Gamma_0(Y))$ and the restriction of $\mathcal{B}(\Gamma_0(Y))$ to $\Gamma^{(n)}(Y)$ coincides (up to a natural isomorphism) with the collection of all symmetric (i.e., invariant under the action of $\sigma \in \mathfrak{S}_n$) Borel-measurable subsets of $Y_{\hat{0}}^{(n)}$. The *correlation measure of the point process* ν is defined as the (unique) measure ρ on $(\Gamma_0(Y), \mathcal{B}(\Gamma_0(Y)))$ which satisfies

$$\int_{\Gamma(Y)} \sum_{\lambda \in \gamma} G(\lambda) d\nu(\gamma) = \int_{\Gamma_0(Y)} G(\lambda) d\rho(\lambda) \quad (12)$$

for each measurable function $G : \Gamma_0(Y) \rightarrow [0, \infty]$. In formula (12), the summation $\sum_{\lambda \in \gamma}$ is over all finite subsets λ of γ .

For each $n \in \mathbb{N}$, we denote by $\rho^{(n)}$ the restriction of the measure ρ to $\Gamma^{(n)}(Y)$. By (12), the measure $\rho^{(n)}$ can be identified with the symmetric measure on $Y_{\hat{0}}^{(n)}$ which satisfies

$$\begin{aligned} & \int_{\Gamma_{pf}(Y)} \sum_{\{(x_1, s_1), \dots, (x_n, s_n)\} \subset \gamma} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\nu(\gamma) \\ &= \int_{Y_{\hat{0}}^{(n)}} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) \end{aligned} \quad (13)$$

for each symmetric measurable function $f^{(n)} : Y_{\hat{0}}^{(n)} \rightarrow [0, \infty]$. Since $\nu(\Gamma_p(Y)) = 1$, the measure $\rho^{(n)}$ is concentrated on the smaller set

$$\mathcal{V}_n := \{(x_1, s_1, \dots, x_n, s_n) \in Y^n \mid (x_1, \dots, x_n) \in X_{\hat{0}}^{(n)}\}. \quad (14)$$

The following theorem gives a three-step way of recovering the correlation measure ρ of the point process $\nu = \mathcal{E}(\mu)$ directly from the moment sequence $(M^{(n)})_{n=0}^{\infty}$.

Theorem 2. Let μ be a random discrete measure on X which has finite moments. Let $(M^{(n)})_{n=0}^\infty$ be the moment sequence of μ , and assume that condition (C1) is satisfied.

(i) For each $n \in \mathbb{N}$ and $\Delta \in \mathcal{B}_c(X_0^{(n)})$, there exists a unique finite measure $\xi_\Delta^{(n)}$ on $(\mathbb{R}_+)^n$ which solves the moment problem

$$\int_{(\mathbb{R}_+)^n} s_1^{i_1} \cdots s_n^{i_n} d\xi_\Delta^{(n)}(s_1, \dots, s_n) = \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta), \quad (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \quad (15)$$

(ii) For each $n \in \mathbb{N}$, there exists a unique measure $\xi^{(n)}$ on \mathcal{V}_n which satisfies

$$\xi_\Delta^{(n)}(A) = \int_{\mathcal{V}_n} \chi_\Delta(x_1, \dots, x_n) \chi_A(s_1, \dots, s_n) d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n) \quad (16)$$

for all $\Delta \in \mathcal{B}_c(X_0^{(n)})$ and $A \in \mathcal{B}((\mathbb{R}_+)^n)$.

(iii) For each $n \in \mathbb{N}$, let $\rho^{(n)}$ be the measure on \mathcal{V}_n given by

$$d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) := (s_1 \cdots s_n)^{-1} d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n). \quad (17)$$

Then $\rho^{(n)}$ is the restriction of the correlation measure ρ of the point process $\nu = \mathcal{E}(\mu)$ to $\Gamma^{(n)}(X)$.

Remark 3. Note that, by the definition of a correlation measure, one always has $\rho(\emptyset) = 1$. Note also that $\rho^{(n)}$ is related to $(M^{(n)})_{n=0}^\infty$ via a moment problem, because, as shown in the proof, the following relation holds, for any measurable function $g^{(n)} : X_0^{(n)} \rightarrow [0, \infty]$,

$$\begin{aligned} & \int_{\mathcal{V}_n} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n} d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \\ &= \frac{1}{n!} \int_{X_0^{(n)}} g^{(n)}(x_1, \dots, x_n) dM_{i_1, \dots, i_n}(x_1, \dots, x_n). \end{aligned} \quad (18)$$

Proof of Theorem 2. We start the proof with derivation of the following bound.

Lemma 4. Assume that, for each $n \in \mathbb{N}$, $m^{(n)}$ is a symmetric measure on X^n . Assume that, for each $\Lambda \in \mathcal{B}_c(X)$, there exists a constant $C_\Lambda > 0$ such that $m^{(n)}(\Lambda^n) \leq C_\Lambda^n n!$ for all $n \in \mathbb{N}$. Then, for any $i_1, \dots, i_n \in \mathbb{N}$, $n \in \mathbb{N}$, and $\Lambda \in \mathcal{B}_c(X)$,

$$\frac{1}{n!} m_{i_1, \dots, i_n}(\Lambda_0^{(n)}) \leq i_1! \cdots i_n! C_\Lambda^{i_1 + \cdots + i_n}.$$

Proof. Fix any $i_1, \dots, i_n \in \mathbb{N}$ and $\Lambda \in \mathcal{B}_c(X)$. Abbreviate $I := i_1 + \dots + i_n$. Let $\pi = \{A_1, \dots, A_n\} \in \Pi(I)$ be as in (8). By the construction of the measure m_{i_1, \dots, i_n} , we get

$$\begin{aligned}
& m_{i_1, \dots, i_n}(\Lambda_{\hat{0}}^{(n)}) \\
&= \int_{X_\pi^{(I)}} \chi_{\Lambda^n}(x_1, x_{i_1+1}, \dots, x_{i_1+\dots+i_{n-1}+1}) dm^{(I)}(x_1, \dots, x_I) \\
&= \int_{X^I} \chi_{\Lambda^I \cap X_\pi^{(I)}}(x_1, \dots, x_I) dm^{(I)}(x_1, \dots, x_I) \\
&= \int_{X^I} \chi_{\Lambda_\pi^{(I)}} dm^{(I)} \\
&= \int_{X^I} \text{Sym}_I \chi_{\Lambda_\pi^{(I)}} dm^{(I)}. \tag{19}
\end{aligned}$$

Here, for a function $f^{(k)} : X^k \rightarrow \mathbb{R}$, $\text{Sym}_k f^{(k)}$ denotes its symmetrization:

$$(\text{Sym}_k f^{(k)})(x_1, \dots, x_k) := \sum_{\sigma \in \mathfrak{S}_k} \frac{1}{k!} f(x_{\sigma(1)}, \dots, x_{\sigma(k)}).$$

Let $\psi \in \Pi(I)$ be a partition having exactly n blocks:

$$\psi = \{B_1, \dots, B_n\}.$$

Set $j_l := |B_l|$, $l = 1, \dots, n$. Denote by Ψ_{i_1, \dots, i_n} the set of all such partitions ψ which satisfy

$$(i_1, \dots, i_n) = (j_{\sigma(1)}, \dots, j_{\sigma(n)})$$

for some permutation $\sigma \in \mathfrak{S}_n$. An easy combinatoric argument shows that the number N_{i_1, \dots, i_n} of all partitions in Ψ_{i_1, \dots, i_n} is equal to

$$N_{i_1, \dots, i_n} = \frac{I!}{i_1! \dots i_n! r_1! r_2! r_3! \dots}. \tag{20}$$

Here for $l = 1, 2, 3, \dots$, r_l denotes the number of coordinates in the vector (i_1, i_2, \dots, i_n) which are equal l . In particular,

$$r_1 + r_2 + r_3 + \dots = n,$$

which implies

$$r_1! r_2! r_3! \dots \leq n!.$$

Therefore,

$$N_{i_1, \dots, i_n} \geq \frac{I!}{i_1! \dots i_n! n!}. \tag{21}$$

For each $\psi \in \Psi_{i_1, \dots, i_n}$,

$$\text{Sym}_I \chi_{\Lambda_\psi^{(I)}} = \text{Sym}_I \chi_{\Lambda_\pi^{(I)}}.$$

Hence, by (19) and (21),

$$\begin{aligned} & \frac{1}{n!} m_{i_1, \dots, i_n}(\Lambda_0^{(n)}) \\ &= \frac{1}{n! N_{i_1, \dots, i_n}} \sum_{\psi \in \Psi_{i_1, \dots, i_n}} \int_{X^I} \chi_{\Lambda_\psi^{(I)}} dm^{(I)} \\ &\leq \frac{i_1! \cdots i_n!}{I!} \int_{X^I} \sum_{\psi \in \Psi_{i_1, \dots, i_n}} \chi_{\Lambda_\psi^{(I)}} dm^{(I)} \\ &\leq \frac{i_1! \cdots i_n!}{I!} m^{(I)}(\Lambda^I) \\ &\leq i_1! \cdots i_n! C_\Lambda^I. \end{aligned}$$

□

To prove statements (i)–(iii) of the theorem, let us first carry out some considerations. Note that, for each $n \in \mathbb{N}$ and each measurable function $f^{(n)} : X^n \rightarrow [0, \infty]$, the functional

$$\mathbb{K}(X) \ni \eta \mapsto \langle \eta^{\otimes n}, f^{(n)} \rangle \in [0, \infty]$$

is measurable and

$$\int_{\mathbb{K}(X)} \langle \eta^{\otimes n}, f^{(n)} \rangle d\mu(\eta) = \int_{X^n} f^{(n)} dM^{(n)}. \quad (22)$$

As easily seen, equality (13) can be extended to the class of all measurable (not necessarily symmetric) functions $f^{(n)} : \mathcal{Y}_n \rightarrow [0, \infty]$ as follows:

$$\begin{aligned} & \int_{\Gamma_{pf}(Y)} \frac{1}{n!} \sum_{\substack{(x_1, s_1), \dots, (x_n, s_n) \in \gamma \\ x_1, \dots, x_n \text{ different}}} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\nu(\gamma) \\ &= \int_{\mathcal{Y}_n} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \end{aligned} \quad (23)$$

If we extend the function $f^{(n)}$ by zero to the whole space Y^n , we can rewrite (23) in the equivalent form:

$$\begin{aligned} & \int_{\Gamma_{pf}(Y)} \frac{1}{n!} \sum_{(x_1, s_1), \dots, (x_n, s_n) \in \gamma} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\nu(\gamma) \\ &= \int_{\mathcal{Y}_n} f^{(n)}(x_1, s_1, \dots, x_n, s_n) d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \end{aligned} \quad (24)$$

In particular, for any measurable function $g^{(n)} : X^n \rightarrow [0, \infty]$ which vanishes outside $X_{\hat{0}}^{(n)}$ and any $i_1, \dots, i_n \in \mathbb{N}$, we get

$$\begin{aligned} & \int_{\Gamma_{pf}(Y)} \frac{1}{n!} \sum_{(x_1, s_1), \dots, (x_n, s_n) \in \gamma} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n} d\nu(\gamma) \\ &= \int_{\mathcal{Y}_n} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n} d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \end{aligned} \quad (25)$$

For each function $f : X_{\pi}^{(i_1 + \dots + i_n)} \rightarrow \mathbb{R}$ one can define a function $X_{\hat{0}}^{(n)} \rightarrow \mathbb{R}$ via $B_{\pi} : X_{\pi}^{(i_1 + \dots + i_n)} \rightarrow X_{\hat{0}}^{(n)}$. Now we will describe the opposite procedure. For simplicity of notation, we will write below

$$\mathcal{I}_n(x_1, \dots, x_n) := \chi_{X_{\hat{1}}^{(n)}}(x_1, \dots, x_n), \quad (x_1, \dots, x_n) \in X^n.$$

Thus, $\mathcal{I}_n(x_1, \dots, x_n)$ is equal to 1 if $x_1 = x_2 = \dots = x_n$, and is equal to zero otherwise. For $i_1, \dots, i_n \in \mathbb{N}$, we define a function $\mathcal{I}_{i_1, \dots, i_n} : X^{i_1 + \dots + i_n} \rightarrow \{0, 1\}$ by setting

$$\begin{aligned} & \mathcal{I}_{i_1, \dots, i_n}(x_1, \dots, x_{i_1 + \dots + i_n}) \\ &:= \mathcal{I}_{i_1}(x_1, \dots, x_{i_1}) \mathcal{I}_{i_2}(x_{i_1+1}, \dots, x_{i_1+i_2}) \cdots \mathcal{I}_{i_n}(x_{i_1 + \dots + i_{n-1} + 1}, \dots, x_{i_1 + \dots + i_n}). \end{aligned}$$

For a measurable function $g^{(n)} : X^n \rightarrow [0, \infty)$ which vanishes outside $X_{\hat{0}}^{(n)}$, we define a measurable function $\mathcal{R}_{i_1, \dots, i_n} g^{(n)} : X^{i_1 + \dots + i_n} \rightarrow [0, \infty]$ by

$$\begin{aligned} & (\mathcal{R}_{i_1, \dots, i_n} g^{(n)})(x_1, \dots, x_{i_1 + \dots + i_n}) \\ &:= g^{(n)}(x_1, x_{i_1+1}, x_{i_1+i_2+1}, \dots, x_{i_1 + \dots + i_{n-1} + 1}) \mathcal{I}_{i_1, \dots, i_n}(x_1, \dots, x_{i_1 + \dots + i_n}). \end{aligned} \quad (26)$$

Note that the function $\mathcal{R}_{i_1, \dots, i_n} g^{(n)}$ vanishes outside the set $X_{\pi}^{(i_1 + \dots + i_n)}$, where $\pi = \{A_1, \dots, A_n\}$ with the sets A_1, \dots, A_n being as in (8). For each $\eta \in \mathbb{K}(X)$,

$$\begin{aligned} & \langle \eta^{\otimes(i_1 + \dots + i_n)}, \mathcal{R}_{i_1, \dots, i_n} g^{(n)} \rangle \\ &= \sum_{(x_1, s_1), \dots, (x_I, s_I) \in \mathcal{E}(\eta)} (\mathcal{R}_{i_1, \dots, i_n} g^{(n)})(x_1, \dots, x_I) s_1 \cdots s_I \\ &= n! \sum_{\{(x_1, s_1), \dots, (x_n, s_n)\} \subset \mathcal{E}(\eta)} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n}. \end{aligned} \quad (27)$$

Here we write $I = i_1 + \dots + i_n$ to save the space. By (25), (27), and the definition of the measure ν , we get

$$\frac{1}{n!} \int_{\mathbb{K}(X)} \langle \eta^{\otimes(i_1 + \dots + i_n)}, \mathcal{R}_{i_1, \dots, i_n} g^{(n)} \rangle d\mu(\eta)$$

$$= \int_{\mathcal{V}_n} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \dots s_n^{i_n} d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n).$$

Hence, by (22),

$$\begin{aligned} & \int_{\mathcal{V}_n} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \dots s_n^{i_n} d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \\ &= \frac{1}{n!} \int_{X^{i_1+\dots+i_n}} \mathcal{R}_{i_1, \dots, i_n} g^{(n)} dM^{(i_1+\dots+i_n)} \\ &= \frac{1}{n!} \int_{X_\pi^{(i_1+\dots+i_n)}} \mathcal{R}_{i_1, \dots, i_n} g^{(n)} dM^{(i_1+\dots+i_n)}, \end{aligned}$$

where the partition π is as above. From here we conclude that equality (18) holds. We define a symmetric measure $\xi^{(n)}$ on \mathcal{V}_n by setting

$$d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n) := s_1 \dots s_n d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \quad (28)$$

Then, equality (18) can be rewritten as follows:

$$\begin{aligned} & \int_{\mathcal{V}_n} g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \dots s_n^{i_n} d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n). \\ &= \frac{1}{n!} \int_{X_0^{(n)}} g^{(n)}(x_1, \dots, x_n) dM_{i_1+1, \dots, i_n+1}(x_1, \dots, x_n), \quad (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \end{aligned} \quad (29)$$

For any $\Delta \in \mathcal{B}_c(X_0^{(n)})$, let $\xi_\Delta^{(n)}$ be the finite measure on $(\mathbb{R}_+)^n$ which satisfies (16). Denote

$$\xi_{\mathbf{i}}^\Delta = \xi_{i_1, \dots, i_n}^\Delta := \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta), \quad \mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \quad (30)$$

Then, by (29) and (30),

$$\xi_{\mathbf{i}}^\Delta = \int_{(\mathbb{R}_+)^n} s_1^{i_1} \dots s_n^{i_n} d\xi_\Delta^{(n)}(s_1, \dots, s_n), \quad \mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \quad (31)$$

Thus, $(\xi_{\mathbf{i}}^\Delta)_{\mathbf{i} \in \mathbb{Z}_+^n}$ is the moment sequence of the finite measure $\xi_\Delta^{(n)}$.

Choose any $\Lambda \in \mathcal{B}_c(X)$ such that $\Delta \subset \Lambda_0^{(n)}$. By formulas (1), (30) and Lemma 4,

$$\begin{aligned} \xi_{i_1, \dots, i_n}^\Delta &\leq \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Lambda_0^{(n)}) \\ &\leq (i_1 + 1)! \dots (i_n + 1)! C_\Lambda^{i_1+\dots+i_n+n} \\ &\leq (i_1 + \dots + i_n + n)! C_\Lambda^{i_1+\dots+i_n+n}, \quad (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \end{aligned} \quad (32)$$

We are now ready to finish the proof of the theorem. Since $(\xi_{\mathbf{i}}^\Delta)_{\mathbf{i} \in \mathbb{Z}_+^n}$ is the moment sequence of the finite measure $\xi_\Delta^{(n)}$ on $(\mathbb{R}_+)^n$, and since this moment sequence satisfies

estimate (32), we conclude from e.g. [4, Chapter 5, Subsec. 2.1, Examples 2.1, 2.2] that the moment sequence $(\xi_i^\Delta)_{i \in \mathbb{Z}_+^n}$ uniquely identifies the measure $\xi_\Delta^{(n)}$. Hence, statement (i) holds. Next, equality (16) evidently holds by (29). Note also that the values of the measure $\xi^{(n)}$ on the sets of the form

$$\{(x_1, s_1, \dots, x_n, s_n) \in \mathcal{V}_n \mid (x_1, \dots, x_n) \in \Delta, (s_1, \dots, s_n) \in A\}$$

where $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$ and $A \in \mathcal{B}((\mathbb{R}_+)^n)$, completely identify the measure $\xi^{(n)}$ on \mathcal{V}^n . Thus, statement (ii) holds. Finally, statement (iii) trivially follows from (28). \square

3 A characterization of random discrete measure in terms of moments

In this section, we assume that μ is a random measure on X which has finite moments. Let $(M^{(n)})_{n=0}^\infty$ be its moment sequence. We assume additionally to condition (C1) that condition (C2) is satisfied.

Remark 5. Assumption (C2) is usually satisfied by a measure μ being concentrated on the cone $\mathbb{K}(X)$. In the latter case, by the proof of Theorem 2, we have

$$\begin{aligned} M^{(n)}(\Lambda_{\hat{0}}^{(n)}) &= M_{1, \dots, 1}(\Lambda_{\hat{0}}^{(n)}) = n! \xi^{(n)}(\mathcal{V}_n \cap (\Lambda \times \mathbb{R}_+)^n) \\ &= n! \int_{\mathcal{V}_n \cap (\Lambda \times \mathbb{R}_+)^n} s_1 \cdots s_n d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n), \end{aligned}$$

so that estimate (2) becomes

$$\int_{\mathcal{V}_n \cap (\Lambda \times \mathbb{R}_+)^n} s_1 \cdots s_n d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) \leq (C'_\Lambda)^n.$$

For example, in the case of the gamma measure (see Introduction), we have

$$\int_{\mathcal{V}_n \cap (\Lambda \times \mathbb{R}_+)^n} s_1 \cdots s_n d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) = \frac{1}{n!} \left(\int_\Lambda dx \right)^n,$$

so condition (C2) is trivially satisfied.

Note also that one should not expect that the constant C_Λ in estimate (C1) becomes small as set Λ shrinks to an empty set. This, for example, is not even true in the case of the gamma measure. Indeed,

$$M^{(n)}(\Lambda) = \prod_{k=0}^{n-1} \left(\int_\Lambda dx + k \right).$$

($M^{(n)}(\Lambda)$ is the n -th moment of the gamma distribution with parameter $\int_\Lambda dx$.) For each n , this decays at most like $\int_\Lambda dx$ and hence C_Λ cannot decrease to zero.

Recall that before Theorem 1 we fixed a sequence $(\Lambda_l)_{l=1}^\infty$ of compact subsets of X such that $\Lambda_1 \subset \Lambda_2 \subset \Lambda_3 \subset \dots$ and $\bigcup_{l=1}^\infty \Lambda_l = X$.

Proof of Theorem 1. Assume that $\mu(\mathbb{K}(X)) = 1$ and let us show that conditions (i) and (ii) are satisfied. Let $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$. It follows from the proof of Theorem 2 (see in particular formula (31)) that the sequence $(\xi_i^\Delta)_{i \in \mathbb{Z}_+^n}$ is the moment sequence of the finite measure $\xi_\Delta^{(n)}$. Hence, condition (i) is indeed satisfied (see e.g. [4, Chapter 5, Subsec. 2.1]).

Next, let $\Delta \in \mathcal{B}(X_{\hat{0}}^{(n)})$ be of the form $\Delta = (\Lambda_l)_{\hat{0}}^{(n)}$. Clearly, $(r_i^\Delta)_{i=0}^\infty$ is the moment sequence of the first coordinate projection of the measure $\xi_\Delta^{(n)}$, which we denote by $P_1 \xi_\Delta^{(n)}$. The measure $P_1 \xi_\Delta^{(n)}$ is concentrated on $[0, \infty)$, hence (5) follows (see e.g. [1, Chapter 2, Subsec. 6.5]). By (C1), (16), (29), Lemma 4 and as $\Delta = (\Lambda_l)_{\hat{0}}^{(n)}$

$$\begin{aligned}
r_i^\Delta &= \int_{(\mathbb{R}_+)^n} s_1^i d\xi_\Delta^{(n)}(s_1, \dots, s_n) \\
&= \int_{\mathcal{Y}_n} \chi_\Delta(x_1, \dots, x_n) s_1^i d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n) \\
&= \frac{1}{n!} \int_{X_{\hat{0}}^{(n)}} \chi_\Delta(x_1, \dots, x_n) dM_{i+1,1,1,\dots,1}(x_1, \dots, x_n) \\
&= \frac{1}{n!} M_{i+1,1,1,\dots,1}((\Lambda_l)_{\hat{0}}^{(n)}) \\
&\leq (i+1)! C_\Lambda^{n+i}, \quad i \in \mathbb{Z}_+.
\end{aligned} \tag{33}$$

Hence, by the Carleman criterion (see e.g. [1]), the measure $P_1 \xi_\Delta^{(n)}$ is the unique measure on \mathbb{R} which has moments $(r_i^\Delta)_{i=0}^\infty$. Therefore, by [1, formula (4) in Chapter I, Sect.1; Chapter II, Subsec. 4.1; Theorem 2.5.3], formula (6) follows from the fact that the measure $P_1 \xi_\Delta^{(n)}$ has no atom at point 0. Thus, condition (ii) is satisfied.

Remark 6. Note that, in this part of the proof, we have not used condition (C2).

Let us now prove the converse statement. So, we assume that $(M^{(n)})_{n=0}^\infty$ is a sequence of symmetric Radon measures fulfilling (C1), (C2), (i), (ii) and $M^{(0)} = 1$. That $(M^{(n)})_{n=0}^\infty$ is the moment sequence of a probability measure μ on $(\mathbb{M}(X), \mathcal{B}(\mathbb{M}(X)))$ will only be used in Lemma 15. We will show the existence of a measure μ' with $\mu'(\mathbb{K}(X)) = 1$ which has as its moments $(M^{(n)})_{n=0}^\infty$. Finally, we will argue that, if $(M^{(n)})_{n=0}^\infty$ is the moment sequence of a probability measure μ , then by the uniqueness of solution of the moment problem $\mu = \mu'$ and hence $\mu(\mathbb{K}(X)) = 1$.

Fix any $n \in \mathbb{N}$ and $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$. Choose $\Lambda \in \mathcal{B}_c(X)$ such that $\Delta \subset \Lambda_{\hat{0}}^{(n)}$. By (C1), (3), and Lemma 4,

$$\xi_{i_1, \dots, i_n}^\Delta = \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta)$$

$$\begin{aligned}
&\leq \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Lambda_{\hat{0}}^{(n)}) \\
&\leq (i_1 + 1)! \cdots (i_n + 1)! C_{\Lambda}^{i_1 + \dots + i_n + n} \\
&\leq (i_1 + \dots + i_n + n)! C_{\Lambda}^{i_1 + \dots + i_n + n}, \quad (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \tag{34}
\end{aligned}$$

Furthermore, by condition (i), the sequence $(\xi_{\mathbf{i}}^{\Delta})_{\mathbf{i} \in \mathbb{Z}_+^n}$ is positive definite. Hence, using e.g. [4, Chapter 5, Subsec. 2.1, Examples 2.1, 2.2], we conclude that there exists a unique measure $\xi_{\Delta}^{(n)}$ on \mathbb{R}^n such that $(\xi_{\mathbf{i}}^{\Delta})_{\mathbf{i} \in \mathbb{Z}_+^n}$ is its moment sequence, i.e.,

$$\xi_{\mathbf{i}}^{\Delta} = \int_{\mathbb{R}^n} s_1^{i_1} \cdots s_n^{i_n} d\xi_{\Delta}^{(n)}(s_1, \dots, s_n), \quad \mathbf{i} = (i_1, \dots, i_n) \in \mathbb{Z}_+^n. \tag{35}$$

Lemma 7. *Let $n \in \mathbb{N}$. Let $\{\Delta_k\}_{k=1}^{\infty}$ be a sequence of disjoint sets from $\mathcal{B}_c(X_{\hat{0}}^{(n)})$. Denote $\Delta := \bigcup_{k=1}^{\infty} \Delta_k$ and assume that $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$. We then have*

$$\sum_{k=1}^{\infty} \xi_{\Delta_k}^{(n)} = \xi_{\Delta}^{(n)}. \tag{36}$$

Proof. Fix any $i_1, \dots, i_n \in \mathbb{Z}_+$. Since $M^{(i_1 + \dots + i_n)}$ is a measure, we easily get

$$\begin{aligned}
&\sum_{k=1}^{\infty} \int_{\mathbb{R}^n} s_1^{i_1} \cdots s_n^{i_n} d\xi_{\Delta_k}^{(n)}(s_1, \dots, s_n) \\
&= \sum_{k=1}^{\infty} \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta_k) \\
&= \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta) \\
&= \int_{\mathbb{R}^n} s_1^{i_1} \cdots s_n^{i_n} d\xi_{\Delta}^{(n)}(s_1, \dots, s_n).
\end{aligned}$$

Hence, the measures $\sum_{k=1}^{\infty} \xi_{\Delta_k}^{(n)}$ and $\xi_{\Delta}^{(n)}$ have the same moments. The measure $\xi_{\Delta}^{(n)}$ fulfils the Carleman bound, hence it is uniquely identified by its moments. So (36) holds. \square

Lemma 8. *For any $n \in \mathbb{N}$ and $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$, the measure $\xi_{\Delta}^{(n)}$ is concentrated on $(\mathbb{R}_+)^n$.*

Proof. Fix any $l \in \mathbb{N}$ and set $\Delta = (\Lambda_l)_{\hat{0}}^{(n)}$. By (4) and (35),

$$r_i^{\Delta} = \int_{\mathbb{R}^n} s_1^i d\xi_{\Delta}^{(n)}(s_1, \dots, s_n), \quad i \in \mathbb{Z}_+.$$

Thus, the numbers $(r_i^\Delta)_{i=0}^\infty$ form the moment sequence of the first coordinate projection of the measure $\xi_\Delta^{(n)}$, which we denote, as above, by $P_1\xi_\Delta^{(n)}$. As easily follows from (34) and the Carleman criterion, the measure $P_1\xi_\Delta^{(n)}$ is uniquely identified by its moment sequence. Then, by (5), the measure $P_1\xi_\Delta^{(n)}$ is concentrated on $[0, \infty)$, and by (6), $(P_1\xi_\Delta^{(n)})(\{0\}) = 0$, see [1]. Therefore, the measure $P_1\xi_\Delta^{(n)}$ is concentrated on \mathbb{R}_+ . Evidently, for any $(i_1, \dots, i_n) \in \mathbb{Z}_+^n$ and any $\sigma \in \mathfrak{S}_n$, we get

$$\begin{aligned} & \int_{\mathbb{R}^n} s_{\sigma(1)}^{i_1} \cdots s_{\sigma(n)}^{i_n} d\xi_\Delta^{(n)}(s_1, \dots, s_n) \\ &= \int_{\mathbb{R}^n} s_1^{i_{\sigma^{-1}(1)}} \cdots s_n^{i_{\sigma^{-1}(n)}} d\xi_\Delta^{(n)}(s_1, \dots, s_n) \\ &= \frac{1}{n!} M_{i_{\sigma^{-1}(1)+1}, \dots, i_{\sigma^{-1}(n)+1}}(\Delta) \\ &= \frac{1}{n!} M_{i_1+1, \dots, i_n+1}(\Delta) \\ &= \int_{\mathbb{R}^n} s_1^{i_1} \cdots s_n^{i_n} d\xi_\Delta^{(n)}(s_1, \dots, s_n). \end{aligned}$$

Hence, the measure $\xi_\Delta^{(n)}$ is symmetric on \mathbb{R}^n . Therefore, for each $j = 1, \dots, n$, the j -th coordinate projection of $\xi_\Delta^{(n)}$ is concentrated on \mathbb{R}_+ . This implies that the measure $\xi_\Delta^{(n)}$ is concentrated on $(\mathbb{R}_+)^n$.

Now, fix an arbitrary $\Delta \in \mathcal{B}_c(X_\emptyset^{(n)})$. Choose $l \in \mathbb{N}$ large enough so that $\Delta \subset (\Lambda_l)_\emptyset^{(n)} =: \Delta'$. Then, by Lemma 7,

$$\xi_\Delta^{(n)}(\mathfrak{C}(\mathbb{R}_+)^n) \leq \xi_{\Delta'}^{(n)}(\mathfrak{C}(\mathbb{R}_+)^n) = 0.$$

Here $\mathfrak{C}(\mathbb{R}_+)^n$ denotes the complement of $(\mathbb{R}_+)^n$. Thus, the measure $\xi_\Delta^{(n)}$ is concentrated on $(\mathbb{R}_+)^n$. \square

Lemma 9. *For each $n \in \mathbb{N}$, there exists a unique measure $\xi^{(n)}$ on \mathcal{Y}_n which satisfies (16) for all $\Delta \in \mathcal{B}_c(X_\emptyset^{(n)})$ and $A \in \mathcal{B}((\mathbb{R}_+)^n)$.*

Proof. For each $\Delta \in \mathcal{B}_c(X^n)$, we define a measure $\xi_\Delta^{(n)}$ on $(\mathbb{R}_+)^n$ by

$$\xi_\Delta^{(n)} := \xi_{\Delta \cap X_\emptyset^{(n)}}^{(n)}.$$

(Note that $\Delta \cap X_\emptyset^{(n)} \in \mathcal{B}_c(X_\emptyset^{(n)})$.) The statement analogous to Lemma 7 holds for $\mathcal{B}_c(X^n)$. So, it suffices to prove that there exists a unique measure $\xi^{(n)}$ on Y^n which satisfies

$$\xi^{(n)}(\Delta \times A) = \xi_\Delta^{(n)}(A), \quad \Delta \in \mathcal{B}_c(X^n), \quad A \in \mathcal{B}((\mathbb{R}_+)^n). \quad (37)$$

But this follows from the fact that, for each $\Delta \in \mathcal{B}_c(X^n)$, $\xi_\Delta^{(n)}$ is a measure on $(\mathbb{R}_+)^n$ and from Lemma 7, see e.g. Remark (3), p. 66 in [12]. \square

Let us now recall a result from [15, Corollary 1] (see also [5, 13]) about existence of a unique point process with a given correlation measure. We will adopt this result to the case of the locally compact Polish space $Y = X \times \mathbb{R}_+$.

Let ρ be a measure on $(\Gamma_0(Y), \mathcal{B}(\Gamma_0(Y)))$. We assume that ρ satisfies the conditions (LB) and (PD) introduced below.

(LB) *Local bound:* For any $\Lambda \in \mathcal{B}_c(X)$ and $A \in \mathcal{B}_c(\mathbb{R}_+)$, there exists a constant $\text{const}_{\Lambda, A} > 0$ such that

$$\rho^{(n)}((\Lambda \times A)^n \cap \mathcal{V}_n) \leq \text{const}_{\Lambda, A}^n, \quad n \in \mathbb{N},$$

and for any sequence $\Lambda_k \in \mathcal{B}_c(X)$ such that $\Lambda_k \downarrow \emptyset$ and $A \in \mathcal{B}_c(\mathbb{R}_+)$, we have $\text{const}_{\Lambda_k, A} \rightarrow 0$ as $k \rightarrow \infty$.

To formulate condition (PD) we first need to give some definitions. For any measurable functions $G_1, G_2 : \Gamma_0(Y) \rightarrow \mathbb{R}$, we define their \star -product as the measurable function $G_1 \star G_2 : \Gamma_0(Y) \rightarrow \mathbb{R}$ given by

$$G_1 \star G_2(\lambda) := \sum_{\substack{\lambda_1 \subset \lambda, \lambda_2 \subset \lambda \\ \lambda_1 \cup \lambda_2 = \lambda}} G_1(\lambda_1) G_2(\lambda_2), \quad \lambda \in \Gamma_0(Y). \quad (38)$$

We denote by \mathcal{S} the class of all functions $G : \Gamma_0(Y) \rightarrow \mathbb{R}$ which satisfy the following assumptions:

- (i) There exists $N \in \mathbb{N}$ such that $G^{(n)} := G \upharpoonright \Gamma^{(n)}(Y) = 0$ for all $n > N$.
- (ii) For each $n = 1, \dots, N$, the function $G^{(n)} := G \upharpoonright \Gamma^{(n)}(Y)$ can be identified with a finite linear combination of functions of the form

$$\text{Sym}_n(\chi_{B_1} \otimes \cdots \otimes \chi_{B_n}),$$

where for $i = 1, \dots, n$ $B_i = \Lambda_i \times A_i$ with $\Lambda_i \in \mathcal{B}_c(X)$ and $A_i \in \mathcal{B}_c(\mathbb{R}_+)$, Sym_n denotes the operator of symmetrization of a function, and

$$(\chi_{B_1} \otimes \cdots \otimes \chi_{B_n})(x_1, s_1, \dots, x_n, s_n) := \chi_{B_1}(x_1, s_1) \cdots \chi_{B_n}(x_n, s_n),$$

where $(x_1, s_1), \dots, (x_n, s_n) \in Y$ with $(x_i, s_i) \neq (x_j, s_j)$ if $i \neq j$.

It is evident that each function $G \in \mathcal{S}$ is bounded and integrable with respect to the measure ρ , and for any $G_1, G_2 \in \mathcal{S}$, we have $G_1 \star G_2 \in \mathcal{S}$.

(PD) *\star -positive definiteness:* For each $G \in \mathcal{S}$, we have

$$\int_{\Gamma_0(Y)} G \star G d\rho \geq 0. \quad (39)$$

Remark 10. Assume that ρ is the correlation measure of a point process ν , and assume that ρ satisfies (LB). Let us briefly explain why condition (PD) must then be satisfied. For a function $G \in \mathcal{S}$, we denote

$$(KG)(\gamma) := \sum_{\lambda \in \gamma} G(\lambda), \quad \gamma \in \Gamma(Y),$$

where $\lambda \in \gamma$ means that $\lambda \subset \gamma$ and $\lambda \in \Gamma_0(Y)$. Then, by (12),

$$\int_{\Gamma_0(Y)} G d\rho = \int_{\Gamma(Y)} KG d\nu.$$

Furthermore, an easy calculation shows that, for any $G_1, G_2 \in \mathcal{S}$, we have

$$K(G_1 \star G_2) = KG_1 \cdot KG_2,$$

where the \cdot in the above equality denotes the pointwise multiplication. Hence, in this case, formula (39) becomes

$$\int_{\Gamma(Y)} (KG)^2 d\nu \geq 0.$$

Theorem 11 ([15]). *Let ρ be a measure on $(\Gamma_0(Y), \mathcal{B}(\Gamma_0(Y)))$ with*

$$\rho(\Gamma^{(0)}(Y)) = 1 \tag{40}$$

which satisfies the conditions (LB) and (PD). Then there exists a unique point process ν in Y whose correlation measure is ρ .

For each $n \in \mathbb{N}$, we now define a measures $\rho^{(n)}$ on \mathcal{Y}_n by

$$d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) := (s_1 \cdots s_n)^{-1} d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n). \tag{41}$$

Note that $\rho^{(n)}$ is a symmetric measure on \mathcal{Y}_n . We next define a measure ρ on $(\Gamma_0(Y), \mathcal{B}(\Gamma_0(Y)))$ by requiring that, for each $n \in \mathbb{N}$, the restriction of the measure ρ to $\Gamma^{(n)}(Y)$ be equal to $\rho^{(n)}$, i.e., for each measurable function $G : \Gamma_0(Y) \rightarrow [0, \infty]$

$$\int_{\Gamma^{(n)}(Y)} G(\lambda) d\rho(\lambda) = \int_{\mathcal{Y}_n} G(\{x_1, s_1, \dots, x_n, s_n\}) d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n). \tag{42}$$

For $n = 0$ we define ρ by (40). A crucial part of the proof of Theorem 1 is the following theorem.

Theorem 12. *Let the measure ρ on $(\Gamma_0(Y), \mathcal{B}(\Gamma_0(Y)))$ be defined by (40)–(42). Then there exists a unique point process ν in Y whose correlation measure is ρ .*

In view of Theorem 11, it suffices to prove that ρ satisfies (LB) and (PD). We split the proof into several lemmas.

Lemma 13. *The measure ρ defined by (40)–(42) satisfies (LB).*

Proof. Let $A \in \mathcal{B}_c(\mathbb{R}_+)$ and let $C := \sup_{s \in A} s^{-1}$. Then we see by (41) that

$$\rho^{(n)}((\Lambda \times A)^n \cap \mathcal{V}_n) \leq C^n \xi^{(n)}((\Lambda \times \mathbb{R}_+)^n \cap \mathcal{V}_n) \quad (43)$$

for each $\Lambda \in \mathcal{B}_c(X)$. By (3), (35), Lemma 9,

$$\begin{aligned} \xi^{(n)}((\Lambda \times \mathbb{R}_+)^n \cap \mathcal{V}_n) &= \xi_{\Lambda_0^{(n)}}^{(n)}((\mathbb{R}_+)^n) \\ &= \xi_{0, \dots, 0}^{\Lambda_0^{(n)}} \\ &= \frac{1}{n!} M_{1, \dots, 1}(\Lambda_0^{(n)}) \\ &= \frac{1}{n!} M^{(n)}(\Lambda_0^{(n)}). \end{aligned} \quad (44)$$

Condition (LB) now follows from (43) and (44), and condition (C2). \square

In order to prove (PD), we rewrite this condition for non-symmetrized product of spaces. We denote

$$\Phi(Y) := \bigcup_{n=0}^{\infty} \Phi^{(n)}(Y),$$

where the set $\Phi^{(0)}(Y)$ contains just one element, and for $n \in \mathbb{N}$, $\Phi^{(n)}(Y) := \mathcal{V}_n$. We define a σ -algebra $\mathcal{B}(\Phi(Y))$ on $\Phi(Y)$ so that, for each $n = 0, 1, 2, \dots$, $\Phi^{(n)}(Y) \in \mathcal{B}(\Phi(Y))$ and for each $n \in \mathbb{N}$, the restriction of $\mathcal{B}(\Phi(Y))$ to $\Phi^{(n)}(Y)$ coincides with $\mathcal{B}(\mathcal{V}_n)$. We can treat ρ as a measure on $\Phi(Y)$, so that $\rho(\Phi^{(0)}(Y)) = 1$ and, for $n \in \mathbb{N}$, the restriction of ρ to $\Phi^{(n)}(Y)$ is $\rho^{(n)}$. We call a function $G : \Phi(Y) \rightarrow \mathbb{R}$ symmetric if, for each $n \in \mathbb{N}$, the restriction of G to $\Phi^{(n)}(Y)$ is a symmetric function. Clearly, each function G on $\Gamma_0(Y)$ identifies a symmetric function on $\Phi(Y)$, for which we preserve the notation G . Furthermore, for an integrable function G , we then have $\int_{\Gamma_0(Y)} G d\rho = \int_{\Phi(Y)} G d\rho$.

Let $m, n \in \mathbb{N}$. We denote by $\text{Pair}(m, n)$ the collection of all subsets \varkappa of the set

$$\{1, 2, \dots, m\} \times \{m+1, m+2, \dots, m+n\}$$

such that, if $(\alpha_i, \beta_i), (\alpha_j, \beta_j) \in \varkappa$ and $(\alpha_i, \beta_i) \neq (\alpha_j, \beta_j)$, then $\alpha_i \neq \alpha_j$ and $\beta_i \neq \beta_j$. By definition, an empty set is an element of $\text{Pair}(m, n)$. For $\varkappa \in \text{Pair}(m, n)$, we denote by $|\varkappa|$ the number of elements of the set \varkappa . In words, this means that we build $|\varkappa|$ pairs between elements in $\{1, 2, \dots, m\}$ and in $\{m+1, m+2, \dots, m+n\}$. Each element can be member of only one pair.

Let $G_1^{(m)} : \mathcal{V}_m \rightarrow \mathbb{R}$, $G_2^{(n)} : \mathcal{V}_n \rightarrow \mathbb{R}$, and let $\varkappa = \{(\alpha_i, \beta_i)\} \in \text{Pair}(m, n)$. We define a function $(G_1^{(m)} \otimes G_2^{(n)})_{\varkappa} : \mathcal{V}_{m+n-|\varkappa|} \rightarrow \mathbb{R}$ as follows. Relabel so that

$$\beta_1 < \beta_2 < \dots < \beta_{|\varkappa|}.$$

Then $(G_1^{(m)} \otimes G_2^{(n)})_{\varkappa}(y_1, \dots, y_{m+n-|\varkappa|})$ is defined as follows. Take

$$G_1^{(m)}(y_1, \dots, y_m)G_2^{(n)}(z_{m+1}, \dots, z_{m+n}).$$

For each $i \in \{1, \dots, |\varkappa|\}$, replace the variable z_{β_i} with y_{α_i} . After this, the variables z_j with $j \in \{m+1, \dots, m+n\} \setminus \{\beta_1, \dots, \beta_{|\varkappa|}\}$ (these are the remaining z_j) are consecutively set to the values $y_{m+1}, y_{m+2}, \dots, y_{m+n-|\varkappa|}$. Here, $y_l := (x_l, s_l)$. In words, this means that $G_1^{(m)}$ and $G_2^{(n)}$ share some of the y_i variables whose indices and positions in the arguments of $G_1^{(m)}$ and $G_2^{(n)}$ are described by the pairs in \varkappa .

For example, for $m = 3$, $n = 4$, $\varkappa = \{(3, 5), (2, 6)\}$, we have

$$(G_1^{(3)} \otimes G_2^{(4)})_{\varkappa}(y_1, y_2, y_3, y_4, y_5) = G_1^{(3)}(y_1, y_2, y_3)G_2^{(4)}(y_4, y_3, y_2, y_5), \quad (y_1, y_2, y_3, y_4, y_5) \in \mathcal{Y}_5.$$

Let us interpret $G_1^{(m)} : \mathcal{Y}_m \rightarrow \mathbb{R}$ and $G_2^{(n)} : \mathcal{Y}_n \rightarrow \mathbb{R}$ as functions defined on $\Phi(Y)$ which vanish outside $\Phi^{(m)}(Y)$ and $\Phi^{(n)}(Y)$, respectively. We then define a function

$$G_1^{(m)} \diamond G_2^{(n)} : \Phi(Y) \rightarrow \mathbb{R}$$

by

$$G_1^{(m)} \diamond G_2^{(n)} := \sum_{\varkappa \in \text{Pair}(m, n)} \frac{(m+n-|\varkappa|)!}{m!n!} (G_1^{(m)} \otimes G_2^{(n)})_{\varkappa}. \quad (45)$$

In the above formula, each $(G_1^{(m)} \otimes G_2^{(n)})_{\varkappa}$ is also treated as a function on $\Phi(Y)$.

Note that a function $G_1^{(0)} : \Phi^{(0)}(Y) \rightarrow \mathbb{R}$ is just a real number and we set, for each function $G_2 : \Phi(Y) \rightarrow \mathbb{R}$,

$$G_1^{(0)} \diamond G_2 = G_2 \diamond G_1^{(0)} := G_1^{(0)} \cdot G_2. \quad (46)$$

Extending formulas (45), (46) by linearity, we define, for any functions $G_1, G_2 : \Phi(Y) \rightarrow \mathbb{R}$, their \diamond -product $G_1 \diamond G_2$ as a function on $\Phi(Y)$.

Lemma 14. *Assume that G_1 and G_2 are symmetric functions on $\Phi(Y)$ which vanish outside the set $\bigcup_{n=0}^N \Phi^{(n)}(Y)$ for some $N \in \mathbb{N}$. Then*

$$\int_{\Phi(Y)} G_1 \star G_2 d\rho = \int_{\Phi(Y)} G_1 \diamond G_2 d\rho,$$

provided the integrals in the above formulas make sense.

Proof. It suffices to consider the case where $G_1 = G_1^{(m)} : \mathcal{Y}_m \rightarrow \mathbb{R}$, $G_2 = G_2^{(n)} : \mathcal{Y}_n \rightarrow \mathbb{R}$ for some $m, n \in \mathbb{N}$. Using (38), we have

$$\int_{\Phi(Y)} G_1^{(m)} \star G_2^{(n)} d\rho$$

$$= \sum_{k=0}^{m \wedge n} \sum_{\substack{(\theta_1, \theta_2, \theta_3) \in \mathcal{P}_3(m+n-k) \\ |\theta_1|=m-k, |\theta_2|=k, |\theta_3|=n-k}} \int_{\mathcal{Y}_{m+n-k}} G_1^{(m)}(y_{\theta_1}, y_{\theta_2}) G_2^{(n)}(y_{\theta_2}, y_{\theta_3}) d\rho^{(m+n-k)}(y_1, \dots, y_{m+n-k}).$$

Here $\mathcal{P}_3(m+n-k)$ denotes the set of all ordered partitions $(\theta_1, \theta_2, \theta_3)$ of the set $\{1, \dots, m+n-k\}$ into three parts, $|\theta_i|$ denotes the number of elements in block θ_i , and, for block $\theta_i = \{r_1, r_2, \dots, r_{|\theta_i|}\}$, y_{θ_i} denotes $y_{r_1}, y_{r_2}, \dots, y_{r_{|\theta_i|}}$. Evidently, the set $\mathcal{P}_3(m+n-k)$ contains $\frac{(m+n-k)!}{(m-k)!(n-k)!k!}$ elements $(\theta_1, \theta_2, \theta_3)$ such that $|\theta_1| = m-k$, $|\theta_2| = k$, $|\theta_3| = n-k$. Hence

$$\begin{aligned} \int_{\Phi(Y)} G_1^{(m)} \star G_2^{(n)} d\rho &= \sum_{k=0}^{m \wedge n} \frac{(m+n-k)!}{(m-k)!(n-k)!k!} \\ &\times \int_{\mathcal{Y}_{m+n-k}} G_1^{(m)}(x_1, \dots, x_m) G_2^{(n)}(x_{m-k+1}, \dots, x_{m+n-k}) d\rho^{(m+n-k)}(x_1, \dots, x_{m+n-k}). \end{aligned} \quad (47)$$

On the other hand, by (45),

$$\int_{\Phi(Y)} G_1^{(m)} \diamond G_2^{(n)} d\rho = \sum_{k=0}^{m \wedge n} \frac{(m+n-k)!}{m!n!} \sum_{\substack{\varkappa \in \text{Pair}(m,n) \\ |\varkappa|=k}} \int_{\mathcal{Y}_{m+n-k}} (G_1^{(m)} \otimes G_2^{(n)})_{\varkappa} d\rho^{(m+n-k)}.$$

An easy combinatoric argument shows that there are

$$\frac{m!}{(m-k)!k!} \times \frac{n!}{(n-k)!k!} \times k! = \frac{m!n!}{(m-k)!(n-k)!k!}$$

elements $\varkappa \in \text{Pair}(m, n)$ such that $|\varkappa| = k$. Hence

$$\begin{aligned} \int_{\Phi(Y)} G_1^{(m)} \diamond G_2^{(n)} d\rho &= \sum_{k=0}^{m \wedge n} \frac{(m+n-k)!}{m!n!} \times \frac{m!n!}{(m-k)!(n-k)!k!} \\ &\times \int_{\mathcal{Y}_{m+n-k}} G_1^{(m)}(x_1, \dots, x_m) G_2^{(n)}(x_{m-k+1}, \dots, x_{m+n-k}) d\rho^{(m+n-k)}(x_1, \dots, x_{m+n-k}). \end{aligned} \quad (48)$$

By (47) and (48) the lemma follows. \square

We denote

$$\Psi(X) := \bigcup_{n=0}^{\infty} \Psi^{(n)}(X),$$

where the set $\Psi^{(0)}(X)$ contains one element, and for $n \in \mathbb{N}$, $\Psi^{(n)}(X) := X^n$. Analogously to $\mathcal{B}(\Phi(Y))$, we define the σ -algebra $\mathcal{B}(\Psi(X))$. We next define a measure M on $(\Psi(X), \mathcal{B}(\Psi(X)))$ so that $M(\Psi^{(0)}(X)) := M^{(0)} = 1$ and, for $n \in \mathbb{N}$, the restriction of M to $\Psi^{(n)}(X)$ is $M^{(n)}$. For any functions $F_1^{(m)}$ and $F_2^{(n)}$ on $\Psi^{(m)}(X)$ and $\Psi^{(n)}(X)$, respectively, their tensor product $F_1^{(m)} \otimes F_2^{(n)}$ is a function on $\Psi^{(m+n)}(X)$. (In the case where either m or n is equal to zero, the tensor product becomes a usual product.) Extending the tensor product by linearity, we define, for any functions F_1 and F_2 on $\Psi(X)$, their tensor product $F_1 \otimes F_2$ as a function on $\Psi(X)$.

The following lemma shows that the measure M on $\Psi(X)$ is \otimes -positive definite.

Lemma 15. *Assume that a function F on $\Psi(X)$ vanishes outside a set $\bigcup_{n=0}^N \Psi^{(n)}(X)$ for some $N \in \mathbb{N}$. Assume that the function $F \otimes F$ is integrable with respect to M . Then*

$$\int_{\Psi(X)} F \otimes F dM \geq 0. \quad (49)$$

Proof. The result immediately follows from

$$\int_{\mathbb{M}(X)} \langle \eta^{\otimes n}, F^{(n)} \rangle d\mu(\eta) = \int_{X^n} F^{(n)} dM^{(n)}.$$

□

Let a function $g^{(n)} : X_{\hat{0}}^{(n)} \rightarrow \mathbb{R}$ be bounded, measurable, and having support from $\mathcal{B}_c(X_{\hat{0}}^{(n)})$. For $i_1, \dots, i_n \in \mathbb{N}$, we set

$$G^{(n)}(x_1, s_1, \dots, x_n, s_n) := g^{(n)}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n}, \quad (x_1, s_1, \dots, x_n, s_n) \in \mathcal{V}_n. \quad (50)$$

We extend the function $g^{(n)}$ by zero to the whole space X^n . We define a function $\mathcal{R}_{i_1, \dots, i_n} g^{(n)} : X^{i_1 + \dots + i_n} \rightarrow \mathbb{R}$ by using formula (26). We denote $G^{(0)} : \Psi^{(0)}(X) \rightarrow \mathbb{R}$

$$\mathcal{K}G^{(n)} := \frac{1}{n!} \mathcal{R}_{i_1, \dots, i_n} g^{(n)}. \quad (51)$$

We denote by \mathcal{Q} the class of all functions on $\Phi(Y)$ which are finite sums of functions of form (50). Extending \mathcal{K} by linearity, we define, for each $G \in \mathcal{Q}$, $\mathcal{K}G$ as a function on $\Psi(X)$.

Lemma 16. *For each $G \in \mathcal{Q}$, we have*

$$\int_{\Phi(Y)} G d\rho = \int_{\Psi(X)} \mathcal{K}G dM. \quad (52)$$

Proof. Let $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$ and let $G^{(n)}$ be given by (50) with $g^{(n)} = \chi_{\Delta}$. By Lemma 9 and formulas (3), (35), (41), and (51),

$$\begin{aligned}
\int_{\mathcal{V}_n} G^{(n)} d\rho^{(n)} &= \int_{\mathcal{V}_n} \chi_{\Delta}(x_1, \dots, x_n) s_1^{i_1} \cdots s_n^{i_n} d\rho^{(n)}(x_1, s_1, \dots, x_n, s_n) \\
&= \int_{\mathcal{V}_n} \chi_{\Delta}(x_1, \dots, x_n) s_1^{i_1-1} \cdots s_n^{i_n-1} d\xi^{(n)}(x_1, s_1, \dots, x_n, s_n) \\
&= \int_{(\mathbb{R}_+)^n} s_1^{i_1-1} \cdots s_n^{i_n-1} d\xi_{\Delta}^{(n)}(s_1, \dots, s_n) \\
&= \xi_{i_1-1, \dots, i_n-1}^{\Delta} \\
&= \frac{1}{n!} M_{i_1, \dots, i_n}(\Delta) \\
&= \int_{X^{i_1+\dots+i_n}} \frac{1}{n!} \mathcal{R}_{i_1, \dots, i_n} \chi_{\Delta} dM^{(i_1+\dots+i_n)} \\
&= \int_{\Psi(X)} \mathcal{K}G^{(n)} dM.
\end{aligned}$$

From here it easily follows by linearity and approximation that formula (52) holds for each $G \in \mathcal{Q}$. \square

Lemma 17. *For each $G \in \mathcal{Q}$,*

$$\int_{\Phi(Y)} G \diamond G d\rho \geq 0.$$

Proof. Let functions $g_1^{(m)} : X_{\hat{0}}^{(m)} \rightarrow \mathbb{R}$ and $g_2^{(n)} : X_{\hat{0}}^{(n)} \rightarrow \mathbb{R}$ be bounded, measurable, and having support from $\mathcal{B}_c(X_{\hat{0}}^{(m)})$ and $\mathcal{B}_c(X_{\hat{0}}^{(n)})$, respectively. Let $i_1, \dots, i_m, j_1, \dots, j_n \in \mathbb{N}$. Let

$$\begin{aligned}
G_1^{(m)}(x_1, s_1, \dots, x_m, s_m) &:= g_1^{(m)}(x_1, \dots, x_m) s_1^{i_1} \cdots s_m^{i_m}, \quad (x_1, s_1, \dots, x_m, s_m) \in \mathcal{V}_m \\
G_2^{(n)}(x_1, s_1, \dots, x_n, s_n) &:= g_2^{(n)}(x_1, \dots, x_n) s_1^{j_1} \cdots s_n^{j_n}, \quad (x_1, s_1, \dots, x_n, s_n) \in \mathcal{V}_n.
\end{aligned}$$

Then, by (26) and (51),

$$\begin{aligned}
&(\mathcal{K}G_1^{(m)} \otimes \mathcal{K}G_2^{(n)})(x_1, \dots, x_{i_1+\dots+i_m+j_1+\dots+j_n}) \\
&= \frac{1}{m! n!} (\mathcal{R}_{i_1, \dots, i_m} g_1^{(m)} \otimes \mathcal{R}_{j_1, \dots, j_n} g_2^{(n)})(x_1, \dots, x_{i_1+\dots+i_m+j_1+\dots+j_n}) \\
&= \frac{1}{m! n!} g_1^{(m)}(x_1, x_{i_1+1}, \dots, x_{i_1+\dots+i_{m-1}+1}) \\
&\quad \times g_2^{(n)}(x_{i_1+\dots+i_m+1}, x_{i_1+\dots+i_m+j_1+1}, \dots, x_{i_1+\dots+i_m+j_1+\dots+j_{n-1}+1}) \\
&\quad \times \mathcal{I}_{i_1, \dots, i_m}(x_1, \dots, x_{i_1+\dots+i_m}) \mathcal{I}_{j_1, \dots, j_n}(x_{i_1+\dots+i_m+1}, \dots, x_{i_1+\dots+i_m+j_1+\dots+j_n}). \tag{53}
\end{aligned}$$

Define for $(x_1, x_{i_1+1}, \dots, x_{i_1+\dots+i_{m-1}+1})$ and

$$(x_{i_1+\dots+i_m+1}, x_{i_1+\dots+i_m+j_1+1}, \dots, x_{i_1+\dots+i_m+j_1+\dots+j_{n-1}+1})$$

the number α_1 as the lowest index j such that there exists a

$$\beta_1 \in \{i_1 + \dots + i_m + 1, i_1 + \dots + i_m + j_1 + 1, \dots, i_1 + \dots + i_m + j_1 + \dots + j_{n-1} + 1\}$$

with $x_j = x_{j'}$. Define $(\alpha_i, \beta_i)_i$ for $i > 1$ analogously. In this way one produces a $\varkappa \in \text{Pair}(m, n)$. Then $(\mathcal{I}_{i_1, \dots, i_m} \otimes \mathcal{I}_{j_1, \dots, j_n})_{\varkappa}$ is of the form $\mathcal{I}_{l_1, \dots, l_{m+n-k}}$ for appropriate l_1, \dots, l_k and $k = |\varkappa|$. By (45), (50)–(53) and recalling that the measure M is symmetric on each $\Psi^{(k)}(X)$,

$$\int_{\Psi(X)} \mathcal{K}G_1^{(m)} \otimes \mathcal{K}G_2^{(n)} dM = \int_{\Psi(X)} \mathcal{K}(G_1^{(m)} \diamond G_2^{(n)}) dM.$$

Hence, for any $G_1, G_2 \in \mathcal{Q}$,

$$\int_{\Psi(X)} \mathcal{K}G_1 \otimes \mathcal{K}G_2 dM = \int_{\Psi(X)} \mathcal{K}(G_1 \diamond G_2) dM. \quad (54)$$

(Note that $G_1 \diamond G_2 \in \mathcal{Q}$.) Hence, by Lemma 15 and (54), for each $G \in \mathcal{Q}$

$$\int_{\Psi(X)} \mathcal{K}(G \diamond G) dM \geq 0.$$

Now the result follows from Lemma 16. \square

Next we extend the result of Lemma 17 to a more general class of functions G by approximation.

Lemma 18. *Let $\Lambda \in \mathcal{B}_c(X)$. Let a function $G : \Phi(Y) \rightarrow \mathbb{R}$ be of the form*

$$G = G^{(0)} + \sum_{j=1}^J G_j^{(n_j)}, \quad (55)$$

where $G^{(0)} : \Phi^{(0)}(Y) \rightarrow \mathbb{R}$, $J \in \mathbb{N}$, and each function $G_j^{(n_j)} : \Phi^{(n_j)}(Y) \rightarrow \mathbb{R}$ is of the form

$$G_j^{(n_j)}(x_1, s_1, \dots, x_{n_j} s_{n_j}) = g_j^{(n_j)}(x_1, \dots, x_{n_j}) f_j^{(n_j)}(s_1, \dots, s_{n_j}) s_1 \cdots s_{n_j}. \quad (56)$$

Here $n_j \in \mathbb{N}$, the functions $g_j^{(n_j)}$ and $f_j^{(n_j)}$ are measurable and bounded and each function $g_j^{(n_j)}$ vanishes outside the set $\Lambda_0^{(n_j)}$. Then

$$\int_{\Phi(Y)} G \diamond G d\rho \geq 0. \quad (57)$$

Proof. Let $N := \max\{n_1, n_2, \dots, n_J\}$. For each $n \in \{1, 2, \dots, N\}$, we define a measure $\zeta_{n,N}$ on $(\mathbb{R}_+)^n$ by

$$\zeta_{n,N} := \sum_{i=n}^{2N} P_n \xi_{\Delta_i}^{(i)}. \quad (58)$$

Here $\Delta_i := \Lambda_0^{(i)}$ and $P_n \xi_{\Delta_i}^{(i)}$ denotes the projection of the (symmetric) measure $\xi_{\Delta_i}^{(i)}$ onto its first n coordinates. Note that $\zeta_{n,N}$ is a symmetric measure on $(\mathbb{R}_+)^n$. We next define a measure $Z_{n,N}$ on $(\mathbb{R}_+)^n$ by

$$dZ_{n,N}(s_1, \dots, s_n) := d\zeta_{n,N}(s_1, \dots, s_n) \sum_{A \in \mathcal{P}(n)} \prod_{j \in A} s_j. \quad (59)$$

Here $\mathcal{P}(n)$ denotes the power set of $\{1, \dots, n\}$ and $\prod_{j \in \emptyset} s_j := 1$. Clearly, $Z_{n,N}$ is also a symmetric measure. By (35), (58), and (59), the moments of the measure $Z_{n,N}$ are given by

$$\int_{(\mathbb{R}_+)^n} s_1^{i_1} \cdots s_n^{i_n} dZ_{n,N}(s_1, \dots, s_n) = \sum_{i=n}^{2N} \sum_{A \in \mathcal{P}(n)} \xi_{i_1 + \chi_A(1), \dots, i_n + \chi_A(n), 0, \dots, 0}^{\Delta_i} \quad (i_1, \dots, i_n) \in \mathbb{Z}_+^n.$$

Hence, by (34),

$$\int_{(\mathbb{R}_+)^n} s_1^{i_1} \cdots s_n^{i_n} dZ_{n,N}(s_1, \dots, s_n) \leq (2N - n - 1) 2^n (i_1 + \cdots + i_n + n + 2N)! C_{\Lambda}^{i_1 + \cdots + i_n + n + 2N}. \quad (60)$$

By (60) and [4, Chapter 5, Subsec. 2.1, Examples 2.1, 2.2], the set of polynomials is dense in $L^2((\mathbb{R}_+)^n, dZ_{n,N})$.

For each $j = 1, \dots, J$, we clearly have $f_j^{(n_j)} \in L^2((\mathbb{R}_+)^{n_j}, dZ_{n_j,N})$. Hence, there exists a sequence of polynomials $(p_{j,k}^{(n_j)})_{k=1}^{\infty}$ such that

$$p_{j,k}^{(n_j)} \rightarrow f_j^{(n_j)} \text{ in } L^2((\mathbb{R}_+)^{n_j}, dZ_{n_j,N}) \text{ as } k \rightarrow \infty. \quad (61)$$

Set $G_k := G^{(0)} + \sum_{j=1}^J G_{j,k}^{(n_j)}$, where

$$G_{j,k}^{(n_j)}(x_1, s_1, \dots, x_{n_j}, s_{n_j}) := g_j^{(n_j)}(x_1, \dots, x_{n_j}) p_{j,k}^{(n_j)}(s_1, \dots, s_{n_j}) s_1 \cdots s_{n_j}.$$

We then have $G_k \in \mathcal{Q}$ for each $k \in \mathbb{N}$. By Lemma 17,

$$\int_{\Phi(Y)} G_k \diamond G_k d\rho \geq 0, \quad k \in \mathbb{N}. \quad (62)$$

We claim that

$$\int_{\Phi(Y)} G_k \diamond G_k d\rho \rightarrow \int_{\Phi(Y)} G \diamond G d\rho \text{ as } k \rightarrow \infty. \quad (63)$$

Indeed, let us fix any $i, j \in \{1, \dots, J\}$ and any $\varkappa \in \text{Pair}(n_i, n_j)$ with $|\varkappa| = l$, and prove that

$$\int_{\mathcal{Y}_{n_i+n_j-l}} (G_{i,k}^{(n_i)} \otimes G_{j,k}^{(n_j)})_{\varkappa} d\rho^{(n_i+n_j-l)} \rightarrow \int_{\mathcal{Y}_{n_i+n_j-l}} (G_i^{(n_i)} \otimes G_j^{(n_j)})_{\varkappa} d\rho^{(n_i+n_j-l)} \quad \text{as } k \rightarrow \infty. \quad (64)$$

For simplicity of notation, let us assume that \varkappa is of the form

$$\{(n_i - l + 1, n_i + 1), (n_i - l + 2, n_i + 2), (n_i - l + 3, n_i + 3) \dots, (n_i, n_i + l)\}.$$

Then

$$\begin{aligned} & \int_{\mathcal{Y}_{n_i+n_j-l}} (G_{i,k}^{(n_i)} \otimes G_{j,k}^{(n_j)})_{\varkappa} d\rho^{(n_i+n_j-l)} \\ &= \int_{\mathcal{Y}_{n_i+n_j-l}} g_i^{(n_i)}(x_1, \dots, x_{n_i}) p_{i,k}^{(n_i)}(s_1, \dots, s_{n_i}) \\ & \quad \times g_j^{(n_j)}(x_{n_i-l+1}, x_{n_i-l+2}, \dots, x_{n_i+n_j-l}) p_{j,k}^{(n_j)}(s_{n_i-l+1}, s_{n_i-l+2}, \dots, s_{n_i+n_j-l}) \\ & \quad \times s_{n_i-l+1} s_{n_i-l+2} \cdots s_{n_i} d\xi^{(n_i+n_j-l)}(x_1, s_1, \dots, x_{n_i+n_j-l}, s_{n_i+n_j-l}). \end{aligned} \quad (65)$$

Hence, there exists $C > 0$ such that

$$\begin{aligned} & \left| \int_{\mathcal{Y}_{n_i+n_j-l}} (G_{i,k}^{(n_i)} \otimes G_{j,k}^{(n_j)})_{\varkappa} d\rho^{(n_i+n_j-l)} - \int_{\mathcal{Y}_{n_i+n_j-l}} (G_i^{(n_i)} \otimes G_j^{(n_j)})_{\varkappa} d\rho^{(n_i+n_j-l)} \right| \\ & \leq \int_{\mathcal{Y}_{n_i+n_j-l}} |g_i^{(n_i)}(x_1, \dots, x_{n_i}) g_j^{(n_j)}(x_{n_i-l+1}, x_{n_i-l+2}, \dots, x_{n_i+n_j-l})| \\ & \quad \times |p_{i,k}^{(n_i)}(s_1, \dots, s_{n_i}) - f_i^{(n_i)}(s_1, \dots, s_{n_i})| \\ & \quad \times |p_{j,k}^{(n_j)}(s_{n_i-l+1}, s_{n_i-l+2}, \dots, s_{n_i+n_j-l})| \\ & \quad \times s_{n_i-l+1} s_{n_i-l+2} \cdots s_{n_i} d\xi^{(n_i+n_j-l)}(x_1, s_1, \dots, x_{n_i+n_j-l}, s_{n_i+n_j-l}) \\ & \leq C \int_{\mathcal{Y}_{n_i+n_j-l}} \chi_{\Lambda_0^{(n_i+n_j-l)}}(x_1, \dots, x_{n_i+n_j-l}) \\ & \quad \times |p_{i,k}^{(n_i)}(s_1, \dots, s_{n_i}) - f_i^{(n_i)}(s_1, \dots, s_{n_i})| \\ & \quad \times |p_{j,k}^{(n_j)}(s_{n_i-l+1}, s_{n_i-l+2}, \dots, s_{n_i+n_j-l})| \\ & \quad \times s_{n_i-l+1} s_{n_i-l+2} \cdots s_{n_i} d\xi^{(n_i+n_j-l)}(x_1, s_1, \dots, x_{n_i+n_j-l}, s_{n_i+n_j-l}) \\ & \leq C \left(\int_{\mathcal{Y}_{n_i+n_j-l}} \chi_{\Lambda_0^{(n_i+n_j-l)}}(x_1, \dots, x_{n_i+n_j-l}) \right. \\ & \quad \left. \times |p_{i,k}^{(n_i)}(s_1, \dots, s_{n_i}) - f_i^{(n_i)}(s_1, \dots, s_{n_i})|^2 \right) \end{aligned}$$

$$\begin{aligned}
& \times s_{n_i-l+1} s_{n_i-l+2} \cdots s_{n_i} d\xi^{(n_i+n_j-l)}(x_1, s_1, \dots, x_{n_i+n_j-l}, s_{n_i+n_j-l}) \Big)^{1/2} \\
& \times \left(\int_{\mathcal{Y}_{n_i+n_j-l}} \chi_{\Lambda_{\bar{0}}^{(n_i+n_j-l)}}(x_1, \dots, x_{n_i+n_j-l}) \right. \\
& \times |p_{j,k}^{(n_j)}(s_{n_i-l+1}, s_{n_i-l+2}, \dots, s_{n_i+n_j-l})|^2 \\
& \left. \times s_{n_i-l+1} s_{n_i-l+2} \cdots s_{n_i} d\xi^{(n_i+n_j-l)}(x_1, s_1, \dots, x_{n_i+n_j-l}, s_{n_i+n_j-l}) \right)^{1/2} \\
& \leq C \|p_{i,k}^{(n_i)} - f_i^{(n_i)}\|_{L^2((\mathbb{R}_+)^{n_i}, dZ_{n_i, N})} \|p_{j,k}^{(n_j)}\|_{L^2((\mathbb{R}_+)^{n_j}, dZ_{n_j, N})} \rightarrow 0 \text{ as } k \rightarrow \infty, \quad (66)
\end{aligned}$$

where we used the Cauchy inequality and (61). Analogously,

$$\left| \int_{\mathcal{Y}_{n_i+n_j-l}} (G_i^{(n_i)} \otimes G_{j,k}^{(n_j)})_{\neq} d\rho^{(n_i+n_j-l)} - \int_{\mathcal{Y}_{n_i+n_j-l}} (G_i^{(n_i)} \otimes G_j^{(n_j)})_{\neq} d\rho^{(n_i+n_j-l)} \right| \rightarrow 0 \text{ as } k \rightarrow \infty. \quad (67)$$

By (66) and (67), formula (64) follows. Formula (63) follows from (64). Now, the lemma follows from (62) and (63). \square

Proof of Theorem 12. As a special case of Lemma 18, formula (57) holds for each function $G \in \mathcal{S}$. Hence, by Lemma 14, the measure ρ satisfies condition (PD). Thus, Theorem 12 is proven. \square

Since the correlation measure ρ of the point process ν from Theorem 12 is concentrated on $\Phi(Y)$, the point process ν is concentrated on $\Gamma_p(Y)$, the set of pinpointing configurations in Y , see e.g. [15, Corollary 1]. Recalling formula (10), one sees that for each $\Lambda \in \mathcal{B}_c(X)$,

$$\begin{aligned}
\int_{\Gamma_p(Y)} \mathfrak{M}_\Lambda d\nu &= \int_{\Gamma_p(Y)} \sum_{(x,s) \in \gamma} \chi_\Lambda(x) s d\nu(\gamma) \\
&= \int_Y \chi_\Lambda(x) s d\rho^{(1)}(x, s) \\
&= \int_Y \chi_\Lambda(x) d\xi^{(1)}(x, s) < \infty. \quad (68)
\end{aligned}$$

Hence, $\mathfrak{M}_\Lambda < \infty$ ν -a.s., and therefore $\nu(\Gamma_{pf}(Y)) = 1$, cf. (11) for the definition of $\Gamma_{pf}(Y)$. Recall the bijective mapping $\mathcal{E} : \mathbb{K}(X) \rightarrow \Gamma_{pf}(Y)$. As already discussed in Section 2, the inverse mapping \mathcal{E}^{-1} is measurable. So we can define a probability measure μ' on $\mathbb{K}(X)$ as the pushforward of ν under \mathcal{E}^{-1} . Thus, to finish the proof of Theorem 1, it suffices to show that $\mu = \mu'$.

Let $\Lambda \in \mathcal{B}_c(X)$. Recall that, for any $i_1, \dots, i_k \in \mathbb{N}$, $k \in \mathbb{N}$,

$$\int_{\mathcal{Y}_k} \chi_{\Lambda_{\bar{0}}^{(k)}}(x_1, \dots, x_k) s_1^{i_1} \cdots s_k^{i_k} d\rho^{(k)}(x_1, s_1, \dots, x_k, s_k) < \infty.$$

Hence, using the definition of a correlation measure (analogously as in (68)), we easily see that, for each $n \in \mathbb{N}$,

$$\int_{\Gamma_{pf}(Y)} \left(\sum_{(x,s) \in \gamma} \chi_{\Lambda}(x)s \right)^n d\nu(\gamma) < \infty.$$

Therefore, for each $n \in \mathbb{N}$,

$$\int_{\mathbb{K}(X)} \eta(\Lambda)^n d\mu'(\eta) < \infty.$$

Here $\eta(\Lambda) := \langle \eta, \chi_{\Lambda} \rangle$, i.e., the η -measure of Λ . Hence, μ' has finite moments. We denote by $(M_{\mu'}^{(n)})_{n=0}^{\infty}$ the moment sequence of the random discrete measure μ' . By Theorem 2 and the construction of the measure ρ , it follows that

$$M'_{i_1, \dots, i_n} = M_{i_1, \dots, i_n}, \quad i_1, \dots, i_n \in \mathbb{N}, \quad n \in \mathbb{N}, \quad (69)$$

where the measures M'_{i_1, \dots, i_n} are defined analogously to M_{i_1, \dots, i_n} , by starting with the moment sequence $(M_{\mu'}^{(n)})_{n=0}^{\infty}$, rather than $(M^{(n)})_{n=0}^{\infty}$. By virtue of (69), the moment sequence $(M_{\mu'}^{(n)})_{n=0}^{\infty}$ coincides with the moment sequence $(M^{(n)})_{n=0}^{\infty}$.

Now, fix any sets $\Lambda_1, \dots, \Lambda_n \in \mathcal{B}_c(X)$. For any $i_1, \dots, i_n \in \mathbb{Z}_+$, we get

$$\begin{aligned} & \int_{\mathbb{K}(X)} \eta(\Lambda_1)^{i_1} \cdots \eta(\Lambda_n)^{i_n} d\mu'(\eta) \\ &= \int_{X^{i_1 + \dots + i_n}} (\chi_{\Lambda_1}^{\otimes i_1} \otimes \cdots \otimes \chi_{\Lambda_n}^{\otimes i_n})(x_1, \dots, x_{i_1 + \dots + i_n}) dM^{(i_1 + \dots + i_n)}(x_1, \dots, x_{i_1 + \dots + i_n}). \\ &= \int_{\mathbb{M}(X)} \eta(\Lambda_1)^{i_1} \cdots \eta(\Lambda_n)^{i_n} d\mu(\eta). \end{aligned} \quad (70)$$

By (C1), (70), and the Carleman criterion, the joint distribution of the random variables $\eta(\Lambda_1), \dots, \eta(\Lambda_n)$ under μ' coincides with the joint distribution of the random variables $\eta(\Lambda_1), \dots, \eta(\Lambda_n)$ under μ . But it is well known (see e.g. [10]) that $\mathcal{B}(\mathbb{M}(X))$ coincides with the minimal σ -algebra on $\mathbb{M}(X)$ with respect to which each function $\eta \mapsto \eta(\Lambda)$ with $\Lambda \in \mathcal{B}_c(X)$, is measurable. Therefore, we indeed get the equality $\mu = \mu'$. \square

4 Moment problem on $\mathbb{K}(X)$

As a consequence of our results, we will now present a solution of the moment problem on $\mathbb{K}(X)$. Consider a sequence $(M^{(n)})_{n=0}^{\infty}$, where $M^{(0)} = 1$ and for each $n \in \mathbb{N}$, $M^{(n)} \in \mathbb{M}(X^n)$ is symmetric. Analogously to the proof of Theorem 1, we define the measure M on $\Psi(X)$. Denote by \mathcal{F} the space of all measurable, bounded functions

$F : \Psi(X) \rightarrow \mathbb{R}$ such that F vanishes outside a set $\Psi^{(0)}(X) \cup \left(\bigcup_{n=1}^N \Lambda^n\right)$ where $N \in \mathbb{N}$ and $\Lambda \in \mathcal{B}_c(X)$. We will say that the sequence $(M^{(n)})_{n=0}^\infty$ is *positive definite* if, for each $F \in \mathcal{F}$, (49) holds. Clearly, if $(M^{(n)})_{n=0}^\infty$ is the moment sequence of a random measure μ , then it is positive definite.

Corollary 19. *Consider a sequence $(M^{(n)})_{n=0}^\infty$, where $M^{(0)} = 1$ and for each $n \in \mathbb{N}$, $M^{(n)} \in \mathbb{M}(X^n)$ is symmetric. Assume that $(M^{(n)})_{n=0}^\infty$ satisfies conditions (C1) and (C2). Then $(M^{(n)})_{n=0}^\infty$ is the moment sequence of a random discrete measure on X if and only if $(M^{(n)})_{n=0}^\infty$ is positive definite and satisfies conditions (i) and (ii) of Theorem 1.*

Proof. The result immediately follows from Theorem 1 and its proof because the existence of μ was only used in Lemma 15. The assertion of this lemma is nothing else but the positive definiteness of $(M^{(n)})_{n=0}^\infty$. \square

We also obtain a characterization of point processes in terms of their moments.

Corollary 20. (i) *Let μ be a random measure on X , i.e., a probability measure on $(\mathbb{M}(X), \mathcal{B}(\mathbb{M}(X)))$. Assume that μ has finite moments, and let $(M^{(n)})_{n=0}^\infty$ be its moment sequence. Further assume that conditions (C1) and (C2) are satisfied. Then μ is a simple point process, i.e., $\mu(\Gamma(X)) = 1$, if and only if, for any $n \in \mathbb{N}$ and any $i_1, \dots, i_n \in \mathbb{N}$, we have $M_{i_1, \dots, i_n} = M_{1, \dots, 1}$, i.e., for each $\Delta \in \mathcal{B}(X_0^{(n)})$,*

$$M_{i_1, \dots, i_n}(\Delta) = M_{1, \dots, 1}(\Delta), \quad i_1, \dots, i_n \in \mathbb{N}. \quad (71)$$

In the latter case, the correlation measure ρ of μ is given by

$$\rho^{(n)}(\Delta) = \frac{1}{n!} M^{(n)}(\Delta), \quad \Delta \in \mathcal{B}(X_0^{(n)}), \quad (72)$$

where $\rho^{(n)}$ is the restriction of ρ to $\Gamma^{(n)}(X)$, $\rho^{(n)}$ being identified with a measure on $X_0^{(n)}$.

(ii) *Consider a sequence $(M^{(n)})_{n=0}^\infty$, where $M^{(0)} = 1$ and for each $n \in \mathbb{N}$, $M^{(n)} \in \mathbb{M}(X^n)$ is symmetric. Assume that $(M^{(n)})_{n=0}^\infty$ satisfies conditions (C1) and (C2). Then $(M^{(n)})_{n=0}^\infty$ is the moment sequence of a simple point process in X if and only if $(M^{(n)})_{n=0}^\infty$ is positive definite and (71) holds.*

Proof. As easily seen, it suffices to prove only part (i). Assume that μ is a point process in X . Hence, μ is a random discrete measure on X . The corresponding point process $\nu = \mathcal{E}(\mu)$ is concentrated on

$$\Gamma(X \times \{1\}) = \{ \{(x, 1)\}_{x \in \gamma} \mid \gamma \in \Gamma(X) \}.$$

Hence, $\Gamma(X \times \{1\})$ can naturally be identified with $\Gamma(X)$, and under this identification we get $\mu = \nu$. Furthermore, the correlation measure ρ of μ coincides with the correlation

measure of ν , provided we have identified $\Gamma_0(X)$ with $\Gamma_0(X \times \{1\})$. Now, formulas (71), (72) follow from Theorem 2.

Next, assume that μ is a random measure which satisfies (71). Hence, for any $n \in \mathbb{N}$ and $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$, we get

$$\xi_{i_1, \dots, i_n}^\Delta = \xi_{0, \dots, 0}^\Delta, \quad i_1, \dots, i_n \in \mathbb{Z}_+.$$

In other words, for each $n \in \mathbb{N}$ and $\Delta \in \mathcal{B}_c(X_{\hat{0}}^{(n)})$, the moment sequence $\xi_{\mathbf{i}}^\Delta$ is constant and thus the measure $\xi_{\Delta}^{(n)}$ is concentrated at one point, $(1, \dots, 1)$. Hence, conditions (i) and (ii) Theorem 1 are satisfied, and so μ is a random discrete measure. Consequently, by (16) and (17), the measure $\rho^{(n)}$ is concentrated on the set

$$\{(x_1, 1, \dots, x_n, 1) \mid (x_1, \dots, x_n) \in X_{\hat{0}}^{(n)}\}.$$

Therefore, the point process $\nu = \mathcal{E}(\mu)$ is concentrated on $\Gamma(X \times \{1\})$. Hence, μ is a point process in X . \square

Let us now assume that $X = \mathbb{R}^d$, or more generally, that X is a connected C^∞ Riemannian manifold. Let $\mathcal{D}(X) := C_0^\infty(X)$ be the space of smooth, compactly supported, real-valued functions on X , equipped with the nuclear space topology, see e.g. [4] for detail. Let $\mathcal{D}'(X)$ be its dual space, and let $\mathcal{C}(\mathcal{D}'(X))$ be the cylinder σ -algebra on it. Note that $\mathbb{M}(X) \subset \mathcal{D}'(X)$ and the trace σ -algebra of $\mathcal{C}(\mathcal{D}'(X))$ on $\mathbb{M}(X)$ coincides with $\mathcal{B}(\mathbb{M}(X))$.

For $\omega \in \mathcal{D}'(X)$ and $\varphi \in \mathcal{D}(X)$, we denote by $\langle \omega, \varphi \rangle$ their dual pairing. Following [5], we inductively define Wick polynomials on $\mathcal{D}'(X)$ by

$$\begin{aligned} \langle : \omega :, \varphi \rangle &:= \langle \omega, \varphi \rangle, \quad \varphi \in \mathcal{D}(X) \\ \langle : \omega^{\otimes n} :, \varphi_1 \otimes \dots \otimes \varphi_n \rangle &:= \frac{1}{n^2} \left[\sum_{i=1}^n \langle \omega, \varphi_i \rangle \langle : \omega^{\otimes (n-1)} :, \varphi_1 \otimes \dots \otimes \check{\varphi}_i \otimes \dots \otimes \varphi_n \rangle \right. \\ &\quad \left. - 2 \sum_{1 \leq i < j \leq n} \langle \omega, \varphi_i \rangle \langle : \omega^{\otimes (n-1)} :, \varphi_1 \otimes \dots \otimes (\varphi_j \varphi_i) \otimes \dots \otimes \check{\varphi}_j \otimes \dots \otimes \varphi_n \rangle \right], \\ \varphi_1, \dots, \varphi_n &\in \mathcal{D}(X), \quad n \geq 2, \end{aligned} \tag{73}$$

where $\check{\varphi}_i$ denotes that the factor φ_i is absent in the tensor product.

Let μ be a probability measure on $(\mathcal{D}'(X), \mathcal{C}(\mathcal{D}'(X)))$ which has finite moments. For each $n \in \mathbb{N}$, we consider the function

$$\begin{aligned} (\mathcal{D}'(X))^{\otimes n} &\rightarrow \mathbb{R} \\ \varphi_1 \otimes \dots \otimes \varphi_n &\mapsto \int_{\mathcal{D}'(X)} \langle : \omega^{\otimes n} :, \varphi_1 \otimes \dots \otimes \varphi_n \rangle d\mu(\omega). \end{aligned} \tag{74}$$

These functions form a proper generalization of the correlation measure of a point process. Let us call them *generalized correlation functions of the measure μ* . The above results can be rewritten in terms of conditions on the generalized correlation functions.

Corollary 21. *Assume that, for each $n \in \mathbb{N}$, the generalized correlation function defined in (74) associated to a measure μ on $\mathcal{D}'(X)$ can be represented via a (positive) measure $\rho^{(n)}$ on $(X_{\hat{0}}^{(n)}, \mathcal{B}(X_{\hat{0}}^{(n)}))$, that is, for any $\varphi_1, \dots, \varphi_n \in \mathcal{D}(X)$,*

$$\int_{\mathcal{D}'(X)} \langle : \omega^{\otimes n} :, \varphi_1 \otimes \dots \otimes \varphi_n \rangle d\mu(\omega) = \int_{X_{\hat{0}}^{(n)}} \varphi_1(x_1) \dots \varphi_n(x_n) d\rho^{(n)}(x_1, \dots, x_n). \quad (75)$$

Furthermore, assume that the measures $\rho^{(n)}$ satisfy condition (C2) in the sense that $M^{(n)}$ is replaced with $\rho^{(n)}$ in the formulation of (C2). Then, μ is a point process, i.e., $\mu(\Gamma(X)) = 1$.

Proof. Using (73), one can easily derive by induction a representation of a monomial $\langle \omega, \varphi_1 \rangle \dots \langle \omega, \varphi_n \rangle$ through Wick polynomials. This formula and (75) imply that, for each $n \in \mathbb{N}$, there exists a (positive) measure $M^{(n)}$ on X^n such that

$$\int_{\mathcal{D}'(X)} \langle \omega, \varphi_1 \rangle \dots \langle \omega, \varphi_n \rangle d\mu(\omega) = \int_{X^n} \varphi_1(x_1) \dots \varphi_n(x_n) dM^{(n)}(x_1, \dots, x_n).$$

Furthermore, formulas (71), (72) hold, because each summand in the representation of a monomial through Wick polynomials corresponds to a particular sub-diagonal $X_{\pi}^{(n)}$ of X^n . (We leave details of these calculations to the interested reader.)

By the assumption of the corollary, the sequence $(M^{(n)})_{n=0}^{\infty}$ with $M^{(0)} = 1$ satisfies (C2). Furthermore, (C2) and (71) easily imply that $(M^{(n)})_{n=0}^{\infty}$ satisfies (C1). Since $(M^{(n)})_{n=0}^{\infty}$ is the moment sequence of a probability measure, it is positive definite. Hence, the statement follows from Corollary 20, (ii). \square

Remark 22. In fact, Corollary 21 is essentially already contained in [5] and [15, Corollary 1], though not presented as an independent result. If we do not assume *a priori* the existence of a measure μ , then we have additionally to assume that the generalised correlation functions have to fulfil the condition (PD). Note that Theorem 11, taken from [15], was used in order to obtain the point process in Y (Theorem 12), which in turn, was used to construct the random discrete measure on X . Hence, it is not surprising that we get a comparable result in the special case where instead of a random discrete measure on X , one actually wants to characterize a point process in X .

Acknowledgements

We are grateful to the referee for his/her careful reading of the manuscript and a number of helpful suggestions for improvement in the article.

The authors acknowledge the financial support of the SFB 701 “Spectral structures and topological methods in mathematics” (Bielefeld University) and the Research Group “Stochastic Dynamics: Mathematical Theory and Applications” (Center for Interdisciplinary Research, Bielefeld University). The authors would like to thank Ilya Molchanov for fruitful discussions.

References

- [1] Akhiezer, N.I.: The classical moment problem and some related questions in analysis. Hafner Publishing Co., New York, 1965.
- [2] Aldous, D.J., Barlow, M.T.: On countable dense random sets. Seminar on Probability, XV (Univ. Strasbourg, Strasbourg, 1979/1980), pp. 311–327, Lecture Notes in Math., 850, Springer, Berlin–New York, 1981.
- [3] Bauer, H.: Measure and integration theory. Walter de Gruyter & Co., Berlin, 2001.
- [4] Berezansky, Y.M., Kondratiev, Y.G.: Spectral methods in infinite-dimensional analysis. Vol. 2, Kluwer Academic Publishers, Dordrecht, 1995.
- [5] Berezansky, Y.M., Kondratiev, Y.G., Kuna, T., Lytvynov, E.: On a spectral representation for correlation measures in configuration space analysis. *Methods Funct. Anal. Topology* 5 (1999), no. 4, 87–100.
- [6] Daley, D. J., Vere-Jones, D.: An introduction to the theory of point processes. Vol. I. Elementary theory and methods. Second edition. Springer-Verlag, New York, 2003.
- [7] Daley, D. J., Vere-Jones, D.: An introduction to the theory of point processes. Vol. II. General theory and structure. Second edition. Springer, New York, 2008.
- [8] Gel’fand, I.M., Graev, M.I., Vershik, A.M.: Models of representations of current groups. *Representations of Lie groups and Lie algebras* (Budapest, 1971), 121179, Akad. Kiadó, Budapest, 1985.
- [9] Hagedorn, D., Kondratiev, Y., Pasurek, T., Rökner, M.: Gibbs states over the cone of discrete measures. *J. Funct. Anal.* 264 (2013), 2550–2583.
- [10] Kallenberg, O.: Random measures. Fourth edition. Akademie-Verlag, Berlin; Academic Press, London, 1986.
- [11] Kendall, W.S.: Stationary countable dense random sets. *Adv. in Appl. Probab.* 32 (2000), 86–100.

- [12] Kingman, J.F.C.: Completely random measures. *Pacific J. Math.* 21 (1967), 59–78.
- [13] Kondratiev, Y.G., Kuna, T.: Harmonic analysis on configuration space. I. General theory. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.* 5 (2002), 201–233.
- [14] Lenard, A.: Correlation functions and the uniqueness of the state in classical statistical mechanics. *Comm. Math. Phys.* 30 (1973), 35–44.
- [15] Lytvynov, E, Mei, L.: On the correlation measure of a family of commuting Hermitian operators with applications to particle densities of the quasi-free representations of the CAR and CCR. *J. Funct. Anal.* 245 (2007), 62–88.
- [16] Rota, G.-C., Wallstrom, T.: Stochastic integrals: a combinatorial approach. *Ann. Probab.* 25 (1997), 1257–1283.
- [17] Shifrin, S. N.: Infinite-dimensional smooth symmetric power moment problem for nuclear spaces. *S. N. Shifrin Ukrainskii Matematicheskii Zhurnal*, 28 (1976) 793–802.
- [18] Tsilevich, N, Vershik, A, Yor, M.: An infinite-dimensional analogue of the Lebesgue measure and distinguished properties of the gamma process. *J. Funct. Anal.* 185 (2001), 274–296.
- [19] Vershik, A.M.; Gel’fand, I.M., Graev, M.I.: Commutative model of the representation of the group of flows $SL(2, \mathbf{R})^X$ connected with a unipotent subgroup. *Funct. Anal. Appl.* 17 (1983), 80–82.