

Diet and herding strategies in a changing environment: stable isotope analysis of Bronze Age and Late Antique skeletal remains from Ya'amūn, Jordan

Article

Accepted Version

Sandias, M. and Müldner, G. ORCID: <https://orcid.org/0000-0002-4513-9263> (2015) Diet and herding strategies in a changing environment: stable isotope analysis of Bronze Age and Late Antique skeletal remains from Ya'amūn, Jordan. *Journal of Archaeological Science*, 63. pp. 24-32. ISSN 0305-4403 doi: 10.1016/j.jas.2015.07.009 Available at <https://centaur.reading.ac.uk/42502/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://www.sciencedirect.com/science/article/pii/S030544031500237X>

To link to this article DOI: <http://dx.doi.org/10.1016/j.jas.2015.07.009>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Diet and herding strategies in a changing environment: stable isotope**
2 **analysis of Bronze Age and Late Antique skeletal remains from**
3 **Ya‘amūn, Jordan**

4 Michela Sandias^{a*} and Gundula Müldner^a

5 ^aDepartment of Archaeology, University of Reading, Whiteknights PO Box 227, Reading, U.K.

6 * Corresponding Author. Present address: Via del Cestello 2/2, 40124, Bologna, Italy, email:
7 m.sandias@gmail.com

8

9 **Abstract**

10 Carbon and nitrogen stable isotope ratios of 45 human and 23 faunal bone collagen samples were
11 measured to study human diet and the management of domestic herbivores in past Jordan,
12 contrasting skeletal remains from the Middle and Late Bronze Age and the Late Roman and
13 Byzantine periods from the site of Ya‘amūn near Irbid. The isotope data demonstrate that the
14 management of the sheep and goats changed over time, with the earlier animals consuming more
15 plants from semi-arid habitats, possibly because of transhumant herding strategies. The isotope
16 data for fish presented here are the first from archaeological contexts from the Southern Levant.
17 Although fish of diverse provenance was available at the site, human diet was predominately based
18 on terrestrial resources and there was little dietary variability within each time-period. Isotopic
19 variation between humans from different time-periods can mostly be explained by ‘baseline shifts’
20 in the available food sources; however, it is suggested that legumes may have played a more
21 significant role in Middle and Late Bronze Age diet than later on.

22 **Keywords:** carbon and nitrogen isotopes; bone collagen; Bronze Age; Roman period; Byzantine
23 period; fish;

24 **Highlights**

25 $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values were employed to reconstruct herd management and human diet in past Jordan
26 Bronze Age ovicaprids consumed more plants from semi-arid habitats than Late Roman/Byzantine
27 animals
28 The data suggest drier conditions or transhumant herd management in the Bronze Age
29 Differences in the human data can mostly be explained by environmental changes.

30

31 **1. Introduction**

32 The archaeology of the Near East is one of the areas where carbon and nitrogen isotope analysis has
33 made a considerable impact in recent years, contributing, for example, to our understanding of early
34 husbandry practices (Makarewicz and Tuross, 2012; Pearson *et al.*, 2007;), the heterogeneous
35 origins of people and animals (Hartman *et al.*, 2013, Thompson *et al.* 2008), the contribution to
36 human diet of specific food resources (Lösch, *et al.*, 2006, Richards *et al.*, 2003, Thompson *et al.*,
37 2005) and the development of social complexity (Makarewicz, 2013, Pearson *et al.*, 2013). One of
38 the key strengths of the method is where it permits diachronic comparisons of isotope data from
39 individual sites, enabling the tracing of continuity or change of subsistence strategies and
40 environmental contexts through time. Using this approach, carbon stable isotope analysis of dental
41 enamel of Bronze Age and Byzantine burials from the sites of Ya‘amūn, Sa‘ad and Yasieleh in
42 North Jordan suggested remarkable homogeneity in diet across time and sites (Al-Shorman, 2003,
43 2004). A complementary study of carbon and nitrogen stable isotope ratios in dentinal collagen of
44 human teeth recovered from Ya‘amūn, Sa‘ad and Yasieleh also indicated continuity (King, 2001).
45 Because of the samples chosen, both studies explored diet only over the relatively short period of

tooth formation in childhood and early adolescence and the lack of site- and period-specific faunal baseline data did not allow monitoring for differences in environmental settings in this ecologically diverse region. Building on this previous research, the present study adds new evidence from carbon and nitrogen stable isotope analysis of bone collagen of human and faunal remains from Ya‘amūn, in an attempt to reconstruct animal and human long-term adult diet at the site during two profoundly different time periods. We aim to explore how Bronze Age and Late Antiquity consumption profiles reflect changes in landscape exploitation and economic strategies, which are of great significance for understanding the way of life in northern Jordan in the past.

2. The site of Ya‘amūn

Jordan’s territory is characterised by high variability in vegetation, physiography, hydrology, and climate. Four of the five vegetation regions identified in the Middle East, the Mediterranean region, the Irano-Turanian steppe, the Saharo-Arabian region and the Sudanian region are present in Jordan (Zohary, 1973; Palmer 2013, Figure 1a). Ya‘amūn is located in the northern part of the Western Highlands at about eight hundred metres above sea level, 23 km southeast of Irbid. Here, current mean annual precipitation is ~400mm (Cordova, 2007). However, within few tens of kilometres of Ya‘amūn the amount of rainfall decreases drastically to less than 200 mm, with the dry Jordan River Valley to the west and the steppe and desert landscapes to the east (Figure 1b). The region around Ya‘amūn therefore presents as a mosaic of ecosystems, where oak forests, Mediterranean low vegetation and steppe habitats are in relative close proximity (Al-Eisawi, 1985).

Occupation at Ya‘amūn spanned from the Early Bronze Age, beginning at c. 3600 cal BCE to the Ayyubid-Mamluk period, thirteenth to sixteenth century CE (dates as in Adams, 2008). Since the first season of excavation, several tombs of variable type, chamber and shaft tombs as well as natural caves used for burial have been identified and excavated (Renfro and Cooper, 2000, Rose, 2002, Rose *et al.*, 2007, Rose *et al.*, 2003). As most of these have been robbed in modern times

and/or reused during the Islamic and later periods a detailed description of burial rites is not available (Rose, 2005). Excavations of the tombs and of the Bronze Age settlement have produced Mycenaean and Cypriot ceramic sherds, Egyptian scarabs and a Mittanian cylinder seal, which demonstrate trade and contact with different areas of the Mediterranean (Rose, 2001). During Late Antiquity (approx. 4th– 7th century CE (Watson, 2008)), Ya‘amūn was a thriving agricultural settlement which contributed to the evidently booming economy in the Late Roman and particularly the Byzantine period (Cameron, 1993, Freeman, 2008, Kennedy, 2007, Parker 1999, Rosen, 2007). The prosperity of Ya‘amūn is shown by various olive and wine presses and by numerous carved water cisterns (El-Najjar and Rose, 2003, Rose *et al.*, 2007). Furthermore, the mosaics of the Ya‘amūn Byzantine church are of equally high quality as those in contemporaneous churches of the nearby Decapolis cities (El-Najjar *et al.*, 2001, Rose *et al.*, 2007).

3. Diet and environmental reconstruction by carbon and nitrogen stable isotope analysis of bone collagen

The ratios of the stable isotopes of carbon ($^{13}\text{C}/^{12}\text{C}$) and nitrogen ($^{15}\text{N}/^{14}\text{N}$) are amongst the most frequently measured in ancient skeletal remains for diet and environmental reconstruction. These ratios are conventionally referred to relative to a standard as $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, respectively. Due to isotope partitioning, isotopes of the same element are unequally distributed in different types of soils, and in different water bodies, plants and animals (Ehleringer and Rundel, 1989, Hoefs, 2009). As a result, organisms from different ecosystems can have distinct isotopic signatures and, similarly, their isotopic composition may vary according to their trophic level (Kelly, 2000). Feeding experiments have shown that the stable isotope ratios measured in human and animal bone collagen reflect the isotopic composition of plant and animal foods, and especially of the dietary protein (Ambrose and Norr, 1993, Tieszen and Fagre, 1993, Froehle *et al.*, 2010), consumed over several years of life (Hedges *et al.*, 2007).

94 Photosynthesis is the main source of carbon isotopic variation in terrestrial ecosystems. According
95 to their photosynthetic pathway, terrestrial plants fall into two main groups, C₃ and C₄ plants. C₃
96 plants are characteristic of the temperate environments and include most plants used for human
97 consumption such as wheat, barley, most fruits, legumes and nuts. In contrast, C₄ plants are adapted
98 to high light intensity, high temperatures and frequent water shortages. They include many tropical
99 grasses and the cultural crops millet, sorghum, maize and sugar cane (van der Merwe, 1989).
100 During photosynthesis, C₃ plants tend to incorporate less ¹³C by discriminating more than C₄ plants
101 against ¹³CO₂. The tissues of C₃ plants therefore have lower δ¹³C values than C₄ plants. Mean δ¹³C
102 values are −26 ‰ and −12.5 ‰ for C₃ and C₄ plants, respectively (Smith and Epstein, 1971). These
103 differences in the isotope values of the plants are reflected in the isotopic signature of the
104 consumers, although absolute values are slightly changed by different metabolic pathways (DeNiro
105 and Epstein, 1978 and studies summarised in Ambrose and Norr 1993). The carbon isotope
106 composition of bone collagen therefore provides an indication of the relative contributions of C₃
107 and C₄ plants to the diet of herbivores and, in turn, may give indication of their abundance in the
108 environment. For the humans, who in most cases eat an omnivorous diet, δ¹³C values can reflect
109 either direct consumption of plants or the isotopic composition of meat and dairy products derived
110 from C₃- or C₄-fed animals (Ambrose, 1993). δ¹³C values are also used to identify consumption of
111 marine resources as marine foods have substantially higher δ¹³C than terrestrial C₃ foods (Richards
112 and Hedges, 1999, Schoeninger and DeNiro, 1984). However, within the environmental context of
113 Jordan, it is the variable reliance on C₃ and C₄ plants that is likely to explain most carbon isotope
114 variation in animals and humans.

115 Bone collagen δ¹⁵N values are mainly used as an indicator of the trophic position of an organism in
116 the foodweb, an attribute which is based on the enrichment of consumer tissues in ¹⁵N with each
117 step up the food chain (DeNiro and Epstein, 1981, Minagawa Wada, 1984, Schoeninger and
118 DeNiro, 1984), breastfeeding mammals being at the top of this sequence (Fuller *et al.*, 2006).

119 Although trophic level enrichments are consistently observed in modern foodwebs, estimating the
120 relative contributions of plant and animal protein in omnivorous diets is complicated by a number
121 of factors, not the least uncertainty about the exact mechanisms behind the trophic level effect and
122 how the diet-tissue spacing is affected by various nutritional and metabolic factors (Vanderklift and
123 Ponsard, 2003, Caut *et al.*, 2009). While a full trophic offset in collagen stable isotope studies is
124 commonly estimated as between 3 and 5‰ (Bocherens and Drucker, 2003), higher values in
125 humans have also been suggested (Hedges and Reynard, 2007, O'Connell *et al.*, 2012). It is
126 generally acknowledged that, because most plants are relatively low in protein, their contribution to
127 the diet will be underrepresented in the collagen stable isotope signal; however, another issue that
128 has been raised more recently is the fact that the isotopic composition of plant foods usually needs
129 to be estimated from the bone collagen values of domestic herbivores which may not always give a
130 truthful reflection of the plants used for human consumption (Fraser *et al.*, 2013). Despite these
131 issues, studies have shown that the use of $\delta^{15}\text{N}$ as a broad indicator of the level of animal protein
132 consumption in humans is overall sound, even though they cannot distinguish between different
133 foods of animal origin such as between meat and dairy products (O'Connell and Hedges, 1999,
134 Petzke *et al.*, 2005).

135 Although bone stable isotope data are primarily a reflection of diet, isotope analysis of faunal,
136 specifically herbivore remains has also been used to indirectly reconstruct environmental
137 conditions, the underlying principle being that herbivore data provide an averaged isotope value for
138 the local vegetation, the isotopic composition of which will vary according to a number of
139 environmental and climatic factors (Hedges *et al.*, 2004, van Klinken *et al.*, 1994). Most obviously,
140 herbivore $\delta^{13}\text{C}$ values may give information about the proportion of C_3 *versus* aridity-adapted C_4
141 plants in an area – although the feeding preferences of individual species must also be taken into
142 account (Ambrose and DeNiro, 1986, Hartman *et al.*, 2013). Variations in temperature or rainfall
143 patterns, among others, also affect the isotopic composition of C_3 plants and may therefore be

144 traceable in herbivore tissues (Hartman and Danin, 2010, van Klinken *et al.*, 1994). For nitrogen
145 isotope ratios, a marked inverse correlation has been demonstrated between bone collagen $\delta^{15}\text{N}$
146 values and rainfall, so that herbivores living in arid environments show elevated $\delta^{15}\text{N}$ values
147 (Ambrose and DeNiro, 1986, Heaton *et al.*, 1986). While physiological mechanisms have been
148 proposed in explanation (Ambrose, 1991, Sealy *et al.*, 1987), it is now thought most likely that the
149 'rainfall effect' is due to the ^{15}N -enrichment of plants, through the effects of denitrification and
150 ammonia volatilization in the soil (Hartman, 2011, Heaton, 1987, Murphy and Bowman, 2006,
151 Schwarcz *et al.*, 1999). A linkage has also been found between high $\delta^{15}\text{N}$ values in cultivated plants
152 and use of manure on fields (Bogaard *et al.*, 2007, Fraser *et al.*, 2011) and should be kept in mind
153 when comparing the $\delta^{15}\text{N}$ values of humans and their domestic animals.

154

155 **4. Materials and Methods**

156 The excavation of the Ya'amūn tombs led to the identification of Middle Bronze Age, Late Bronze
157 Age, Late Roman and Byzantine burials, with dating achieved through a systematic study of pottery
158 fragments and grave goods, and of the architectural features (Al-Shorman, 2004, Barnes, 2003,
159 Burke and Rose, 2001, El-Najjar and Rose, 2003, El-Najjar *et al.*, 2001, Rose *et al.*, 2007). From
160 the skeletal remains made accessible at the Department of Archaeology and Anthropology of the
161 University of Yarmouk (Irbid, Jordan), it was possible to sample 45 individuals. Of these, 22 date to
162 the Middle-Late Bronze Age, while 23 date to the Late Roman-Byzantine period. A further 23
163 samples were obtained from the remains of adult domestic ungulates and fishbone recovered from
164 well-dated contexts. The entire assemblage was highly fragmented. In the case of the human
165 remains, this prevented the assessment of age and sex. For the faunal remains, it was usually
166 problematic to distinguish between sheep (*Ovis aries*) and goats (*Capra hircus*) on morphological
167 grounds. These are therefore collectively referred to as domestic ovicaprids or as sheep/goats (for

168 more information on sampling processing and analysis see Supplementary Information). Isotopic
169 differences between the two species on account of their feeding ecology are not expected in the
170 environmental context of the southern Levant (see Hartman et al., 2013: S1).

171 **5. Results**

172 The $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ isotope data and quality indicators of the faunal and human bone samples from
173 Ya‘amūn are reported in Tables S1 and S2 (see Supplementary Materials), respectively. Fifty (18
174 faunal and 32 humans samples) out of 68 bone specimens yielded good quality collagen (Ambrose
175 1990; DeNiro 1985). Descriptive statistics for each of the sample groups are presented in Table 1.
176 The highest variability was found amongst the Bronze Age (MBA and LBA combined) animals,
177 while the Late Antique (LR-Byz) sheep/goats presented the lowest. The δ -values of the domestic
178 herbivores, illustrate a clear difference in the diet of Bronze Age and Late Antique sheep/goats with
179 apparently decreased $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in the later periods (Figure 2). Although the difference is
180 statistically significant for $\delta^{15}\text{N}$ only (Independent Samples Mann-Whitney test with exact
181 probabilities to account for small sample sizes: $U=11.5$, $p=0.537$ for $\delta^{13}\text{C}$; $U=1.0$, $p=0.009$ for
182 $\delta^{15}\text{N}$), the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the MBA/LBA sheep-goats are clearly correlated ($r^2=0.75$).
183 While the later ovicaprids had a more monotonous diet, the two LR/Byz *Bos* specimens plot
184 relatively far apart, with a 3.2‰ difference in $\delta^{13}\text{C}$ and a 4.3‰ difference in $\delta^{15}\text{N}$ values. In Figure
185 3, the human values are compared to the terrestrial fauna. No statistically significant differences
186 were found between human isotope values from MBA and LBA (Independent samples Mann-
187 Whitney test, $U=22.5$, $p=0.536$ for $\delta^{13}\text{C}$ and $U=18.0$, $p=1.0$ for $\delta^{15}\text{N}$), and Late Roman and
188 Byzantine Ya‘amūn ($U=8.5$, $p=0.667$ for $\delta^{13}\text{C}$ and $U=4.5$, $p=1.83$ for $\delta^{15}\text{N}$). Consequently these
189 were pooled into one Bronze Age and one Late Antique group. The differences between the Bronze
190 Age and Late Antique humans are statistically significant for both carbon and nitrogen

191 (Independent-sample Mann-Whitney test, $U=49.0$, $p=0.002$ for $\delta^{13}\text{C}$ and $U=55.5$, $p=0.005$ for
192 $\delta^{15}\text{N}$).

193

194 **6. Discussion**

195 *6.1 The fauna from Ya'amūn*

196 Overall, the range of $\delta^{13}\text{C}$ values of the domestic ungulates from Ya'amūn indicates that they were
197 grazing in an environment where C_3 vegetation dominates over C_4 vegetation. These results are in
198 agreement with the data presented previously by Al-Shorman (2004, 2003) and King (2001) on the
199 human diet and are also consistent with the vegetation composition around the site in modern times
200 (Al-Eisawi 1996, Shomer-Ilan *et al.*, 1981, Vogel *et al.*, 1986, Winter, 1981). Nevertheless, the
201 most positive $\delta^{13}\text{C}$ (-16.6‰ for sheep/goat and -15.6‰ for cattle, Table 1) indicate the inclusion of
202 varying and sometimes substantial amounts of C_4 plants in the animals' diet. These higher $\delta^{13}\text{C}$
203 values are correlated with raised $\delta^{15}\text{N}$ values which are also consistent with grazing in arid regions
204 (Hartman and Danin, 2010, Heaton, 1987, see section 3 above).

205 The differences observed in the sheep/goat data from the Bronze Age and Late Antiquity (Figure 2)
206 are consistent with suggestions that climate in the MBA and LBA in the Southern Levant was dry
207 and similar to present conditions, while sedimentological and palynological evidence as well as
208 speleothem oxygen isotope analyses unequivocally indicate wetter conditions in the Roman and
209 Byzantine periods (Bookman *et al.*, 2004; Enzel *et al.*, 2003; Neumann *et al.*, 2007; Orland *et al.*,
210 2009; Finné *et al.*, 2011; Rambeau and Black 2011). Nevertheless, if climatic conditions in the
211 MBA/LBA were indeed similar to the present day and even more so if, as has also been suggested,
212 they were slightly wetter (see Rambeau and Black 2011: 99), it is unlikely that the MBA/LBA
213 sheep/goats with elevated carbon and/or nitrogen isotope values (at least half of what is, admittedly,
214 a small sample) could have acquired these by freely foraging in the immediate hinterland of the site.

215 Ya'amūn itself is situated in the xeric Mediterranean phytogeographic zone which receives modest
216 rainfall (currently ~400mm/year) and therefore has little C₄ vegetation, which only becomes notable
217 in areas with less than ~350mm annual precipitation . The observed δ -values are therefore
218 consistent with animals grazing in much drier environments and these indeed provide the best
219 parallels among published data. According to the large reference data-set compiled by Hartman *et*
220 *al.* (2013), the MBA/LBA sheep/goats from Ya'amūn are most consistent with ovicaprids feeding
221 in the desert zone, although similar isotope values have also been produced for goats from a semi-
222 arid steppe environment (Makarewicz and Tuross, 2012). In any case, the Bronze Age herbivore
223 data from Ya'amūn suggest that sheep/goat were herded away from the site for at least part of the
224 year. The isotope values for the sheep/goat from Ya'amūn are almost identical to another small
225 data-set of MBA sheep from Tell Al-Husn, only a few kilometres to the north (Al-Bashaireh and
226 Al-Muheisen, 2011). Here, elevated isotope values were only observed in three very young
227 individuals and the ¹⁵N-enrichment was consequently attributed to a 'suckling effect'; however, the
228 consumption of mother ewes' milk cannot easily explain the relatively large difference in $\delta^{13}\text{C}$
229 between the young and adult sheep (Balasse *et al.*, 1999), which approaches 2‰. Instead, the bones
230 of the suckling sheep may record a seasonal shift in the diet of the mothers, during the gestation
231 period and early life of their lambs, to include ¹³C-enriched graze. Transhumance is standard
232 practice amongst the traditional pastoralists in the southern Levant who, usually, spend the months
233 between November and April leading herds to the green pastures that develop in arid and semi-arid
234 areas during the rainfall season. This system has the added advantage of keeping the animals away
235 from agricultural fields during the crop growing season (Levy, 1983, Safrai, 2004, 93). Texts dated
236 to the Middle and Late Bronze periods of Syria-Palestine and Anatolia describe various strategies
237 for managing domestic herbivores (Liverani, 1988, 374, 437), and how, for instance, the sheep
238 owned by the elites spent most of the year away from the city to return only for the shearing season
239 (Snell, 1997, 72, 126). Another scenario is that animals raised at sites in the more arid zones were

240 brought to Ya‘amūn, which fits with artefactual evidence that the site was part of a network of
 241 commercial exchanges connecting north Jordan with the wider eastern Mediterranean (Bourke *et*
 242 *al.*, 2006, Strange, 2008), and would also explain why only part of the MBA/LBA sheep/goat
 243 sample exhibit the raised stable isotope values.

244 In contrast, the Late Roman/Byzantine animals were feeding almost entirely, if not exclusively, on
 245 C₃ plants and their $\delta^{15}\text{N}$ values are significantly lower than in the earlier animals. Assuming, as
 246 Hartman *et al.* (2013: 4372) argue, that the wetter climatic conditions in Late Antiquity did not
 247 significantly shift the boundaries of environmental zones in the region, this suggests that the
 248 sheep/goats of this later period were more restricted in their mobility. They would have grazed on
 249 unused fields and between the orchards in the surroundings of Ya‘amūn where conditions were
 250 relatively moist because of wide-ranging irrigation. It is known that vine leaves and trimmings were
 251 sometimes used as animal fodder (Horden and Purcell, 2000, 214) and these must have been
 252 abundant at Ya‘amūn, where vine cultivation was extensive (see above).

253 There are too few cattle data to attempt reconstructing husbandry regimes, but it is nevertheless
 254 interesting to observe that the two Late Antique cattle have sharply diverging $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values
 255 (Figure 2). While one plots close to the contemporaneous sheep/goats, the other has substantially
 256 higher $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values, suggesting that it fed in a much drier environmental zone and/or an
 257 area where C₄ fodder crops had been adopted (Copley *et al.*, 2004). The animal therefore suggests
 258 that livestock was brought to Late Antique Ya‘amūn from the outside.

259 Ya‘amūn freshwater and marine fish isotope values (Table S1, Figure 4) are comparable with
 260 results from archaeological fish from Greece (Vika & Theodoropoulou 2012). The values for the
 261 marine species, in particular, add to a growing corpus of data that show fish from the Mediterranean
 262 to be ^{13}C -enriched and ^{15}N -depleted compared to animals from the North Atlantic. The observed
 263 differences in $\delta^{15}\text{N}$ between the fish samples can be explained by differences in trophic level

(Froese and Pauly, 2014) and variation in the isotopic composition of aquatic primary producers (Ambrose 1993; France 1995; Fuller 2012b; Katzenberg & Weber 1999; Schoeninger & DeNiro 1984). Particularly notable is the stark difference in $\delta^{13}\text{C}$ between the two LR/Byz samples of the freshwater genus *Tilapia*, which exceeds 10‰ and indicates that these specimens lived in very different environments. The $\delta^{13}\text{C}$ values for modern *Tilapia* from Lake Tiberias (Zohary *et al.*, 1994), once corrected for the offset between muscle and bone collagen as well as lipid content, are very close to the relatively ^{13}C -depleted value observed for YMNTfb100. An origin from the river Jordan or the Lake itself therefore appears plausible. The same may then be true for the specimen of *Clariidae* (catfish, YMNTfb102), as variation between the two could mostly be accounted for by differences in trophic level. *Tilapia* are also known to inhabit coastal rivers, including those of the southern Levant (Van Neer *et al.*, 2000) and a significant input from marine biomass could explain why specimen YMNTfb101 is substantially ^{13}C -enriched over YMNTfb100. Nevertheless, Vika and Theodoropoulou (2012) have observed similar carbon isotope values for freshwater fish from Greece. *Tilapia* are bottom feeders and it is possible that their foodwebs are ^{13}C -enriched by poor availability of CO_2 in warm and stagnant waters (France, 1995, Hecky and Hesslein, 1995), perhaps combined with a hard water effect in a region where marine carbonate is the dominant geological substrate (Day, 1996). Further isotope studies on freshwater fish from the Levant will be necessary to test this hypothesis.

6.2 The inhabitants of Ya'amūn

The isotope values show that terrestrial C_3 -derived resources dominated human diet at Ya'amūn. The isotopic differences between Bronze Age and Late Antique humans mirror those between the sheep/goats from both periods and suggest that much of what first appears as dietary variation can instead be explained by a baseline-shift in the isotopic composition of available food sources (Figure 3). Because of the impact of water availability on carbon isotope discrimination of C_3 plants

288 during photosynthesis (Farquhar *et al.*, 1989), it is likely that the plant foods available at Ya‘amūn
289 in the Bronze Age were also ^{13}C -enriched over those cultivated under the wetter conditions of the
290 Late Roman/Byzantine period. It should be noted that the same cannot necessarily be assumed for
291 plant $\delta^{15}\text{N}$ values as these are determined by numerous complex mechanisms (Evans, 2001,
292 Högberg, 1997). Cultivated fields especially may be subject to additional measures such as the
293 application of animal fertilizers (Fraser *et al.*, 2011). Despite significant differences in the isotopic
294 composition of the bone collagen, human diet between the two time periods may therefore not have
295 varied greatly in terms of the actual staple foods consumed and their relative proportions.

296 In this context, the relatively large difference in the average $\delta^{15}\text{N}$ human-herbivore offset
297 ($\Delta^{15}\text{N}_{\text{human-herbivore}}$), which is 3.5‰ in the Late Antique but only 0.3‰ in the Bronze Age sample,
298 requires some discussion. If herbivore $\delta^{15}\text{N}$ were used to estimate the nitrogen isotope composition
299 of plants consumed by humans and the spacing between humans and animals was therefore taken as
300 an indicator for the relative contributions of plant and animal protein to the human diet (Hedges and
301 Reynard, 2007), these data would suggest that Bronze Age diet was almost entirely based on plant
302 foods, while humans in Late Antiquity habitually consumed large amounts of animal protein (~
303 70% of the dietary protein according to Hedges and Reynard’s (2007) ‘standard model’, even if a
304 generous trophic level offset of +5‰ is used). Either of these extreme scenarios seems unlikely in
305 light of evidence for diet in these periods available from other sources (Grigson, 1998, 256; Safrai,
306 2004, 96).

307 The human isotope data from MBA/LBA Ya‘amūn are almost identical to those obtained from
308 EBA/MBA Tell Al-Husn (mean $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values ± 1 s.d.: $-18.5 \pm 0.4\text{‰}$ and 8.7 ± 0.8 , $n=10$), and
309 suggest that the subsistence regime reflected was typical for the wider region (Al-Bashaireh and Al-
310 Muheisen, 2011). It cannot be denied that staple foods derived from C_3 cereals such as wheat and
311 barley made up the bulk of the diet at this time (Zohary and Hopf, 2000; see Snell, 1997; for Jordan

312 see Bourke *et al.*, 2003; McNicoll *et al.*, 1992; Tubb 1988; Tubb *et al.*, 1997). Nevertheless, the
313 faunal assemblage from BA Ya‘amūn itself as well as nearby sites such as Pella (Bourke *et al.*,
314 1994, Bourke *et al.*, 1998) demonstrate that sheep and goat husbandry was well-established in the
315 region during the Bronze Age. While the emphasis may have been on wool production processing
316 of surplus dairy and the consumption of meat from animals that had outlived their usefulness would
317 have been an integrated part of this economy (Grigson, 1998, 256). If it is therefore unlikely that the
318 human diet was almost entirely plant-based, the relatively low $\delta^{15}\text{N}$ values in the BA humans
319 compared to the animal data need additional explanation. Pulses, especially lentils, were important
320 protein-rich foods and a means of crop and diet diversification in the entire Mediterranean area
321 since the beginning of agriculture (Grigson, 1998, Horden and Purcell, 2000, 203). Like other
322 leguminous plants, they are able to fix nitrogen directly from the atmosphere and, as a result, are
323 habitually ^{15}N -depleted compared to other crops. Their regular inclusion in the diet would therefore
324 have the effect of lowering the $\delta^{15}\text{N}$ of human consumers, potentially masking the consumption of
325 animal products (Fraser *et al.*, 2013).

326 If the human-herbivore $\delta^{15}\text{N}$ offset therefore likely underrepresents the role of animal products in
327 the Bronze Age diet at least to an extent, the opposite may be true for the Late Roman/Byzantine
328 sample. It is well-established that cereals, in the form of wheat or barley bread, contributed the
329 majority of the daily caloric intake in Roman-period Palestine, while the degree to which animal
330 products were part of everyday diet varied according to wealth and, particularly for fish, to the
331 distance from the place of production (Broshi, 1986, Dar, 1995, Garnsey, 1999:16, Safrai 2004,
332 Wilkins and Hill, 2006). Ya‘amūn in Late Antiquity was the site of intensive agriculture, and
333 manuring was widely practiced in Romano-Byzantine Palestine (Almagro, 2007). The herbivore
334 $\delta^{15}\text{N}$ may therefore well underestimate the nitrogen isotope composition of the cultivated plants (see
335 Fraser *et al.*, 2013). Written sources from the region also describe a system of crop rotation where
336 wheat fields were periodically turned over to leguminous plants and used as animal pasture (Safrai,

2004, 98). Alternatively, dietary diversification may be responsible for the human-herbivore spacing. In the Roman period, domestic fowl and chicken especially gained economic importance in the Southern Levant and it has been suggested that egg consumption was considerable (Safrai 2004: 101-102). Quantitative data from Ya‘amūn do not exist, but at nearby Pella, reliance on poultry, mostly chicken, increases sharply during the Byzantine period (McNicoll *et al.*, 1982, 110). The isotopic composition of eggs depends on the diet of the chicken (Hobson, 1995); however, because of their omnivorous feeding ecology, these can be significantly ^{15}N -enriched over herbivores (Müldner and Richards, 2007).

There are numerous sources emphasizing the importance of fish and fish products in Roman and Byzantine Palestine (Garnsey, 1999, Lev-Tov, 2003, Marzano, 2013, Purcell, 1995, Van Neer and Parker, 2008). Based on the fishbone isotope data assembled here, there is little conclusive evidence that fish made any measurable contribution to Roman-Byzantine diet at Ya‘amūn. Lack of isotopic separation between freshwater fish and terrestrial animals makes it very difficult to convincingly demonstrate the consumption of freshwater fish, and although the $\delta^{15}\text{N}$ values of the Late Antique humans could theoretically be explained by small-scale consumption of higher trophic level marine fish (such as the specimen of *Sciaenidae*, YMNfb88), Ya‘amūn’s inland location and the fact that YMNfb88 (which actually dates to the MBA) plots at the top end of $\delta^{15}\text{N}$ values measured for Mediterranean fish to date (and is therefore not necessarily representative of any marine fish that reached the site), make significant consumption of marine protein at the site very unlikely .

The isotope data from Ya‘amūn are very similar to those from other Late Antique sites in the Levant (Al-Bashaireh and Al-Muheisen, 2011, Fuller *et al.*, 2012a, Gregoricka and Sheridan, 2013),, suggesting again a similar subsistence base for the wider region. Unlike some other bone collagen data-sets (Bourbou *et al.*, 2011, Iacumin *et al.*, 1998, Thompson *et al.*, 2008), the sample from Ya‘amūn does not have any statistical outliers that would suggest that individuals moved from ecologically different regions, although the sample size is too small for any far-reaching

362 conclusions and carbon and nitrogen isotopes are not well suited to identify migrants in a
363 population in any case.

364

365 **7. Conclusions**

366 Despite the small sample sizes which are, unfortunately, a common limitation of bone isotope
367 investigations in arid and semi-arid regions, this study has established a number of clear trends. Of
368 particular importance are differences in animal husbandry between the Middle and Late Bronze Age
369 and Late Antiquity, which involved Bronze Age sheep/goats spending at least part of the year in
370 arid or semi-arid regions, while Romano-Byzantine animals evidently stayed in the same
371 phytogeographic zone. The reason behind this significant economic change may be the greater
372 abundance of suitable fodder in the slightly wetter climate of Late Antiquity or else the need to keep
373 the human workforce on-site to concentrate on other agricultural tasks, including the work-intensive
374 viticulture (Horden and Purcell, 2000, 215). The cattle data from this period nevertheless show that
375 the site was still connected to the drier regions to the East and South, possibly reflecting the move
376 to expand agricultural production to the more marginal areas in the Byzantine period (Watson,
377 2008). As expected based on the results of previous studies, the human diet in both periods was
378 based almost exclusively on C₃-based resources. While most isotopic differences between the
379 human groups can be explained in terms of a baseline shift due to climatic change between the two
380 periods, the Middle and Late Bronze Age inhabitants of Ya‘amūn may have consumed a greater
381 proportion of leguminous plants, while the diet in Late Antiquity could have included a wider range
382 of foods. Alternatively, their isotopic data may reflect the documented agricultural intensification in
383 this period. Neither freshwater nor marine fish seem to have contributed significantly to the food
384 intake of the sampled individuals. Overall, this study illustrates the need to analyse coeval faunal
385 remains for human palaeodietary studies and confirms the great value of carbon and nitrogen stable

386 isotope analysis of herbivores for reconstructing environmental conditions in relation to changes in
387 geographical location and climate.

388 **Acknowledgements**

389 This study, part of MS's PhD, was financed by Leverhulme Trust as part of the project "Water Life
390 and Civilisation". Thanks go to Prof. Steven J. Mithen, University of Reading, for help and advice
391 during PhD supervision. We are grateful to Prof. Mahmoud El-Najjar †, Dr Abdulla Al-Shorman,
392 Dr Mohammad Al-Rousan and Dr Ammar Al-Obiedat of the Institute of Archaeology and
393 Anthropology of Yarmouk University, Irbid, Jordan, to Prof Jerome Rose of the University of
394 Arkansas, USA, for access to the skeletal material and for providing assistance and advice during
395 sampling. We gratefully acknowledge the Department of Antiquities of Jordan as well as staff of
396 Council for British Research in the Levant in Amman and especially Prof. Bill Finlayson for
397 support during the 2006 fieldwork. Thanks go to Prof. Wim Van Neer, University of Leuven, for
398 the identification of the fish vertebrae, Sarah Lambert-Gates (Reading) and Carlos H. Caracciolo
399 (INGV, Bologna) for production of maps, and two anonymous reviewers for their constructive and
400 helpful comments.

401

402 **References**

- 403 Adams, R. B. (ed.) 2008. *Jordan, An Archaeological Reader*, London: Equinox.
404 Al-Bashaireh, K. and Al-Muheisen, Z. 2011. Subsistence strategies and palaeodiet of Tell al-Husn,
405 northern Jordan: nitrogen and carbon stable isotope evidence and radiocarbon dates. *Journal*
406 *of Archaeological Science*, 38, 2606-2612.
407 Al-Eisawi, D. M. 1985. Vegetation in Jordan. In: Hadidi, A. (ed.) *Studies in the History and*
408 *Archaeology of Jordan 2*. Amman: Department of Antiquities.
409 Al-Eisawi, D. M. 1996. *Vegetation of Jordan*, Cairo, UNESCO: Regional Office for Science and
410 Technology for the Arab States.

- 411 Al-Shorman, A. 2003. Byzantine palaeodiet and social status at Sa'ad and Yasideh in Northern
412 Jordan. *Athena Review*, 3, 60-63.
- 413 Al-Shorman, A. 2004. Stable carbon isotope analysis of human tooth enamel from the Bronze Age
414 cemetery of Ya'amoun in Northern Jordan. *Journal of Archaeological Science*, 31, 1693-
415 1698.
- 416 Almagro, A. 2007. Soil Improvement and Agricultural Pesticides in Antiquity: Examples from
417 Archaeological Research in Israel. In: Conan, M. (ed.) *Middle East Garden Traditions:
418 Unity and Diversity*. Washington D.C.: Dumbarton Oaks.
- 419 Ambrose, S. H. 1990. Preparation and characterisation of bone and tooth collagen for isotopic
420 analysis. *Journal of Archaeological Science*, 17, 431-451.
- 421 Ambrose, S. H. 1991. Effects of diet, climate and physiology on nitrogen isotopes abundances in
422 terrestrial foodwebs. *Journal of Archaeological Science*, 18, 293-317.
- 423 Ambrose, S. H. 1993. Isotopic analysis of palaeodiets: methodological and interpretative
424 considerations. In: Sandford, M. K. (ed.) *Investigation of Ancient Human Tissue: Chemical
425 Analysis in Anthropology*. Langthorne: Gordon and Breach, 59-130.
- 426 Ambrose, S. H. and DeNiro, M. J. 1986. The isotopic ecology of East African mammals. *Oecologia*,
427 69, 395-406.
- 428 Ambrose, S. H. and Norr, L. 1993. Experimental evidence for the relationship of the carbon isotope
429 ratios of whole diet and dietary protein to those of bone collagen and carbonate. In:
430 Lambert, J. B. & Grupe, G. (eds.) *Prehistoric Human Bone. Archaeology at the Molecular
431 Level*. Berlin: Springer-Verlag, 1-37.
- 432 Balasse, M., Bocherens, H. and Mariotti, A. 1999. Intra-bone variability of collagen and apatite
433 composition used as evidence of a change of diet. *Journal of Archaeological Science*, 26,
434 593-598.
- 435 Barnes, J. S. 2003. *Tell Ya'amūn: Mortuary, Biological and Cultural Characteristics. Unpublished
436 MA dissertation*. Master of Arts MA Dissertation, University of Arkansas.
- 437 Bocherens, H. and Drucker, D. 2003. Trophic level isotopic enrichment of carbon and nitrogen in
438 bone collagen: case studies from recent and ancient terrestrial ecosystems. *International
439 Journal of Osteoarchaeology*, 13, 46-53.
- 440 Bogaard, A., Heaton, T. H. E., Poulton, P. and Merbach, I. 2007. The impact of manuring on
441 nitrogen isotope ratios in cereals: archaeological implications for reconstruction of diet and
442 crop management practices. *Journal of Archaeological Science*, 34, 335-343.
- 443 Bookman, R., Y. Enzel, a. Agnon, and M. Stein. 2004. 'Late Holocene Lake Levels of the Dead
444 Sea.' *Bulletin of the Geological Society of America* 116, 555-571.
- 445 Bourbou, C., Fuller, B. T., Garvie-Lok, S. J. and Richards, M. P. 2011. Reconstructing the diets of
446 Greek Byzantine populations (6th–15th centuries AD) using carbon and nitrogen stable
447 isotope ratios. *American Journal of Physical Anthropology*, 146, 569-581.
- 448 Bourke, S., Sparks, R. and Schroder, M. 2006. Pella in the Middle Bronze Age. In: Fischer, P. M.
449 (ed.) *The Chronology of the Jordan Valley During the Middle and Late Bronze Ages : Pella,
450 Tell Abu Al-Kharaz, and Tell Deir 'Alla*. Wien: Verlag der Österreichischen Akademie der
451 Wissenschaften, 9-58.
- 452 Bourke, S., Sparks, R., Sowada, K. N. and Mairs, L. 1994. Preliminary report of the University of
453 Sydney's fourteenth season of excavations at Pella (Tabaqat Fahl) in 1992. *Annual of the
454 Department of Antiquities of Jordan*, 38, 81-126.

- 455 Bourke, S., Sparks, R., Sowada, K. N., McLaren, P. B. and Mairs, L. 1998. Preliminary report on the
456 University of Sydney's sixteenth and seventeenth seasons of excavations at Pella (Tabaqat
457 Fahl) in 1994/95. *Annual of the Department of Antiquities of Jordan*, 42, 179-211.
- 458 Bourke, S. J., Sparks, R. T., McLaren, B., Sowada, K., Mairs, L. D., Meadows, J., Hikade, T. and
459 Reade, W. 2003. Preliminary reports on the University of Sydney's eighteenth and
460 nineteenth seasons of excavations at Pella (Tabaqat Fahl) in 1996/97. *Annual of the*
461 *Department of Antiquities of Jordan*, 47, 335-388.
- 462 Broshi, M. 1986. The diet of Palestine in the Roman Period - Introductory notes. *Israel Museum*
463 *Journal*, 46, 41-56.
- 464 Burke, D. L. and Rose, J. C. 2001. Les morts de Ya'amūn. *Le Monde de la Bible*, 133, 56.
- 465 Cameron, A. 1993. *The Mediterranean World in Late Antiquity*, London, Routledge.
- 466 Caut, S., Angulo, E. and Courchamp, F. 2009. Variation in discrimination factors (Delta N-15 and
467 Delta C-13): the effect of diet isotopic values and applications for diet reconstruction.
468 *Journal of Applied Ecology*, 46, 443-453.
- 469 Copley, M. S., Jim, S., Jones, V., Rose, P., Clapham, A., Edwards, D. N., Horton, M., Rowley-
470 Conwy, P. and Evershed, R. P. 2004. Short- and long-term foraging and foddering strategies
471 of domesticated animals from Qasr Ibrim, Egypt. *Journal of Archaeological Science*, 31,
472 1273-1286.
- 473 Cordova, C. E. 2007. *Millennial Landscape Change in Jordan*, Tucson, The University of Arizona
474 Press.
- 475 Dar, S. 1995. Food and archaeology in Romano-Byzantine Palestine. In: Wilkins, J., Harvey, D. &
476 Dobson, M. (eds.) *Food in Antiquity*. Exeter: University of Exeter Press.
- 477 Day, S. P. 1996. Dogs, Deer and Diet at Star Carr: a Reconsideration of C-isotope Evidence from
478 Early Mesolithic Dog Remains from the Vale of Pickering, Yorkshire, England. *Journal of*
479 *Archaeological Science*, 23, 783-787.
- 480 DeNiro, M. J. 1985. Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios
481 in relation to palaeodietary reconstruction. *Nature*, 317, 806-809.
- 482 DeNiro, M. J. and Epstein, S. 1978. Influence of diet on the distribution of carbon isotopes in
483 animals. *Geochimica et Cosmochimica Acta*, 42, 495-506.
- 484 DeNiro, M. J. and Epstein, S. 1981. Influence of diet on the distribution of nitrogen isotopes in
485 animal. *Geochimica et Cosmochimica Acta*, 45, 341-351.
- 486 Ehleringer, J. R. and Rundel, P. W. 1989. Stable isotopes: history, units and instrumentation. In:
487 Rundel, P. W., Ehleringer, J. R. & Nagy, K. A. (eds.) *Stable Isotopes in Ecological*
488 *Research*. New York: Springer-Verlag, pp.1-15.
- 489 El-Najjar, M. and Rose, J. C. 2003. Preliminary report of the 2003 field season at Ya'mun by the
490 joint Yarmouk University/University of Arkansas Project. *Annual of the Department of*
491 *Antiquities of Jordan*, 47, 491-492.
- 492 El-Najjar, M., Rose, J. C., Atallah, N., Turshan, N., Khasawneh, N. and Burke, D. L. 2001. First
493 season of excavation at Ya'mun (1999). *Annual of the Department of Antiquities of Jordan*,
494 45, 413-417.
- 495 Enzel, Yehouda, Revital Bookman, David Sharon, Haim Gvirtzman, Uri Dayan, Baruch Ziv, and
496 Mordechai Stein. 2003. 'Late Holocene Climates of the Near East Deduced from Dead Sea
497 Level Variations and Modern Regional Winter Rainfall.' *Quaternary Research*, 60, 263-73.

- 498 Evans, D. R. 2001. Physiological mechanisms influencing plant nitrogen isotope composition.
499 *Trends in Plant Science*, 6, 121-126.
- 500 Farquhar, G. D., Ehleringer, J. R. and Hubick, K. T. 1989. Carbon isotope discrimination and
501 photosynthesis. *Annual Review of Plant Physiology: Plant Molecular Biology*, 40.
- 502 Finné, M., Holmgren, K., Sundqvist, H. S., Weiberg, E. and Lindblom, M. 2011. Climate in the
503 eastern Mediterranean, and adjacent regions, during the past 6000 years – A review. *Journal*
504 *of Archaeological Science*, 38, 3153-3173.
- 505 France, R. 1995. Differentiation between littoral and pelagic food webs in lakes using stable carbon
506 isotopes. *Limnology and Oceanography*, 40, 1310-1313.
- 507 Fraser, R. A., Bogaard, A., Heaton, T., Charles, M., Jones, G., Christensen, B. T., Halstead, P.,
508 Merbach, I., Poulton, P. R., Sparkes, D. and Styring, A. K. 2011. Manuring and stable
509 nitrogen isotope ratios in cereals and pulses: towards a new archaeobotanical approach to
510 the inference of land use and dietary practices. *Journal of Archaeological Science*, 38, 2790-
511 2804.
- 512 Fraser, R. A., Bogaard, A., Schäfer, M., Arbogast, R. and Heaton, T. H. E. 2013. Integrating
513 botanical, faunal and human stable carbon and nitrogen isotope values to reconstruct land
514 use and palaeodiet at LBK Vaihingen an der Enz, Baden-Württemberg. *World Archaeology*,
515 45, 492-517.
- 516 Freeman, P. 2008. The Roman Period. In: Adams, R. B. (ed.) *Jordan. An Archaeological Reader*.
517 London: Equinox, 413-441.
- 518 Froehle, A. W., Kellner, C. M. and Schoeninger, M. J. 2010. Effect of diet and protein source on
519 carbon stable isotope ratios in collagen: follow up to Warinner and Tuross (2009). *Journal*
520 *of Archaeological Science*, 37, 2662.
- 521 Froese, R. and Pauly, D. 2014. *Fish Base*. [Online]. Available: <http://www.fishbase.org> [Accessed
522 20/05/2014].
- 523 Fuller, B. T., De Cupere, B., Marinova, E., Van Neer, W., Waelkens, M. and Richards, M. P. 2012a.
524 Isotopic reconstruction of human diet and animal husbandry practices during the Classical-
525 Hellenistic, imperial, and Byzantine periods at Sagalassos, Turkey. *American Journal of*
526 *Physical Anthropology*, 149, 157-171.
- 527 Fuller, B. T., Müldner, G., Van Neer, W., Ervynck, A. and Richards, M. P. 2012b. Carbon and
528 nitrogen stable isotope ratio analysis of freshwater, brackish and marine fish from Belgian
529 archaeological sites (1st and 2nd millennium AD). *Journal of Analytical Atomic*
530 *Spectrometry*, 27, 807-820.
- 531 Fuller, B. T., Fuller, J. L., Harris, D. A. and Hedges, R. E. M. 2006. Detection of breastfeeding and
532 weaning in modern human infants with carbon and nitrogen stable isotope ratios. *American*
533 *Journal of Physical Anthropology*, 129, 279-293.
- 534 Garcia-Guixé, E., Subirà, M. E., Marlasca, R. and Richards, M. P. 2010. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in ancient
535 and recent fish bones from the Mediterranean Sea. *Journal of Nordic Archaeological*
536 *Science*, 17, 83-92.
- 537 Garnsey, P. 1999. *Food and society in classical antiquity*, Cambridge, Cambridge University Press.
- 538 Gregoricka, L. A. and Sheridan, S. G. 2013. Ascetic or affluent? Byzantine diet at the monastic
539 community of St. Stephen's, Jerusalem from stable carbon and nitrogen isotopes. *Journal of*
540 *Anthropological Archaeology*, 32, 63-73.

541 Grigson, C. 1998. Plough and pasture in the early economy of the Southern Levant. *In*: Levy, T. E.
542 (ed.) *The Archaeology of Society in the Holy Land*. London: Continuum, 245-268.

543 Hartman, G. 2011. Are elevated $\delta^{15}\text{N}$ values in herbivores in hot and arid environments caused by
544 diet or animal physiology? *Functional Ecology*, 25, 122-131.

545 Hartman, G., Bar-Oz, G., Bouchnick, R. and Reich, R. 2013. The pilgrimage economy of Early
546 Roman Jerusalem (1st century BCE-70 CE) reconstructed from the $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ values of
547 goat and sheep remains. *Journal of Archaeological Science*, 40, 4369-4376.

548 Hartman, G. and Danin, A. 2010. Isotopic values of plants in relation to water availability in the
549 Eastern Mediterranean region. *Oecologia*, 162, 837-852.

550 Heaton, T. H. E. 1987. The $^{15}\text{N}/^{14}\text{N}$ ratios of plants in South Africa and Namibia: relationship to
551 climate and coastal/saline environments. *Oecologia*, 74, 236-246.

552 Heaton, T. H. E., Vogel, J. C., la Chevallierie, G. v. and Collet, G. 1986. Climatic influence on the
553 isotopic composition of bone nitrogen. *Nature*, 322, 822-823.

554 Hecky, R. E. and Hesslein, R. H. 1995. Contribution of benthic algae to lake food webs as revealed
555 by stable isotope analysis. *Journal of the North American Benthological Society*, 14, 631-
556 653.

557 Hedges, R. E. M., Clement, J. G., Thomas, C. D. L. and O'Connell, T. C. 2007. Collagen turnover in
558 the adult femoral mid-shaft: modeled from anthropogenic radiocarbon tracer measurements.
559 *American Journal of Physical Anthropology*, 133, 808-816.

560 Hedges, R. E. M. and Reynard, L. M. 2007. Nitrogen isotopes and the trophic level of humans in
561 archaeology. *Journal of Archaeological Science*, 34, 1240-1251.

562 Hedges, R. E. M., Stevens, R. E. and Richards, M. 2004. Bone as a stable isotope archive for local
563 climatic information. *Quaternary Science Reviews*, 23, 959-965.

564 Hobson, K. A. 1995. Reconstructing Avian Diets Using Stable-Carbon And Nitrogen Isotope
565 Analysis Of Egg Components - Patterns Of Isotopic Fractionation And Turnover. *Condor*,
566 97, 752-762.

567 Hoefs, J. 2009. *Stable Isotope Geochemistry*, Berlin et al., Springer.

568 Högborg, P. 1997. Tansley Review No. 95. ^{15}N natural abundance in soil-plant systems. *New*
569 *Phytologist*, 137, 179-203.

570 Horden, P. and Purcell, N. 2000. *The Corrupting Sea: a Study of Mediterranean History*, Oxford,
571 Blackwell.

572 Iacumin, P., Bocherens, H., Chaix, L. and Marioth, A. 1998. Stable carbon and nitrogen isotopes as
573 dietary indicators of ancient Nubian populations (Northern Sudan). *Journal of*
574 *Archaeological Science*, 25, 293-301.

575 Katzenberg, M. A. and Weber, A. 1999. Stable isotope ecology and palaeodiet in the Lake Baikal
576 region of Siberia. *Journal of Archaeological Science*, 26, 651-659.

577 Kelly, J. F. 2000. Stable isotopes of carbon and nitrogen in the study of avian and mammalian
578 trophic ecology. *Canadian Journal of Zoology*, 78, 1-27.

579 Kennedy, D. 2007. *Gerasa and the Decapolis. A "Virtual Island" in Northwest Jordan*, London,
580 Duckworth.

581 King, M. 2001. *Analysis of diet in Byzantine Jordan: isotopic evidence in human dentine*
582 *(Contribution to the Bioarchaeology of the Levant)*. Unpublished MA dissertation. Master of
583 Arts MA dissertation, University of Arkansas.

584 Levy, T. E. 1983. The emergence of specialized pastoralism in the southern Levant. *World*
585 *Archaeology*, 15, 15-36.

586 Liverani, M. 1988. *Antico Oriente*, Bari, Editori Laterza.

587 Lösch, S., Grupe, G. and Peters, J. 2006. Stable isotopes analysis and dietary adaptations in humans
588 and animals at Pre-Pottery Neolithic Nevalı Çori, Southeast Anatolia. *American Journal of*
589 *Physical Anthropology*, 131, 181-193.

590 Makarewicz, C. 2013. More than meat: diversity in caprine harvesting strategies and the emergence
591 of complex production systems during the Late Pre-Pottery Neolithic B. *Levant*, 45, 236-
592 261.

593 Makarewicz, C. and Tuross, N. 2012. Finding fodder and tracking transumance: isotopic detection
594 of goat domestication processes in the Near East. *Current Anthropology*, 53, 495-505.

595 Marzano, A. 2013. *Harvesting the Sea: The Exploitation of Marine Resources in the Roman*
596 *Mediterranean*, Oxford, Oxford University Press.

597 McNicoll, A., Smith, R. H. and Hennessy, B. 1982. *Pella in Jordan 1, An interim report on the joint*
598 *University of Sydney and The College of Wooster excavations at Pella 1979-1981*, Canberra,
599 Australian National Gallery.

600 McNicoll, A. W., Edwards, P. C., Hanbury-Tenison, J., Hennessy, J. B., Potts, T. F., Smith, R. H.,
601 Walmsley, A. and Watson, P. (eds.) 1992. *Pella In Jordan 2. The second interim report on*
602 *the joint University of Sydney and The College of Wooster excavations at Pella 1982-1985*,
603 Sydney: Mediterranean Archaeology.

604 Minagawa, M. and Wada, E. 1984. Stepwise enrichment of ^{15}N along food chains: further evidence
605 and the relation between $\delta^{15}\text{N}$ and animal age. *Geochimica et Cosmochimica Acta*, 48,
606 1135-1140.

607 Müldner, G. and Richards, M. P. 2007. Diet and diversity at Later Medieval Fishergate: the isotopic
608 evidence. *American Journal of Physical Anthropology*, 134, 162-174.

609 Murphy, B. P. and Bowman, D. M. J. S. 2006. Kangaroo metabolism does not cause the relationship
610 between bone collagen $\delta^{15}\text{N}$ and water availability. *Functional Ecology*, 20, 1062-1069.

611 Neumann, Frank Harald, Elisa J. Kagan, Markus J. Schwab, and Mordechai Stein. 2007.
612 'Palynology, Sedimentology and Palaeoecology of the Late Holocene Dead Sea.'
613 *Quaternary Science Reviews* 26, 1476-98.

614 O'Connell, T. C. and Hedges, R. E. M. 1999. Investigation into the effect of diet on modern human
615 hair isotopic values. *American Journal of Physical Anthropology*, 108, 409-425.

616 O'Connell, T. C., Kneale, C. J., Tasevska, N. and Khunle, G. G. C. 2012. The diet-body offset in
617 human nitrogen isotopic values: A controlled dietary study. *American Journal of Physical*
618 *Anthropology*, 149, 426-434.

619 Orland, Ian J., Miryam Bar-Matthews, Noriko T. Kita, Avner Ayalon, Alan Matthews, and John W.
620 Valley. 2009. 'Climate Deterioration in the Eastern Mediterranean as Revealed by Ion
621 Microprobe Analysis of a Speleothem That Grew from 2.2 to 0.9 Ka in Soreq Cave, Israel.'
622 *Quaternary Research* 71, 27-35.

623 Palmer, C. 2013. Biogeography. In: Ababsa, M. (Ed.), *Atlas of Jordan: History, Territories and*
624 *Society*. Presses de l'Ifpo: Beyrouth, Retrieved from <http://books.openedition.org/ifpo/4871>.

625 Parker, T. S. 1999. An Empire's new Holy Land: the Byzantine Period. *Near Eastern Archaeology*,
626 62, 134-180.

- Pearson, J. A., Buitenhuis, H., Hedges, R. E. M., Martin, L., Russel, N. and Twiss, K. C. 2007. New light on early caprine herding strategies from isotope analysis: a case study from Neolithic Anatolia. *Journal of Archaeological Science*, 34, 2170-2179.
- Pearson, J. A., Grove, M., Özbek, M. and Hongo, H. 2013. Food and social complexity at Çayönü Tepesi, southeastern Anatolia: stable isotope evidence of differentiation in diet according to burial practice and sex in the early Neolithic *Journal of Anthropological Archaeology*, 32, 180-189.
- Petzke, K. J., Boeing, H. and Metges, C. C. 2005. Choice of dietary protein of vegetarians and omnivores is reflected in their hair protein ¹³C and ¹⁵N abundance. *Rapid Communications in Mass Spectrometry*, 19, 1392-1400.
- Purcell, N. 1995. Eating fish: the paradoxes of seafood. In: Wilkins, J., Harvey, D. & Dobson, M. (eds.) *Food in Antiquity*. Exeter: University of Exeter Press.
- Rambeau, C. and Black, S. 2011. Palaeoenvironments of the Southern Levant 5,000 BP to present: linking geological and archaeological records. In: Mithen, S. & Black, E. (eds.) *Water, Life and Civilisation. Climate, Environment and Society in the Jordan Valley*. Cambridge: Cambridge University Press.
- Renfro, B. and Cooper, H. K. 2000. Ya'amün in Egan, V., Bikai P.M., Zamora, K. (eds) *Archaeology in Jordan*. *American Journal of Archaeology*, 104, 581.
- Richards, M. P. and Hedges, R. E. M. 1999. Stable isotope evidence for similarities in the types of marine foods used by Late Mesolithic humans at sites along the Atlantic coast of Europe. *Journal of Archaeological Science*, 26, 717-722.
- Richards, M. P., Pearson, J. A., Molleson, T. I., Russell, N. and L., M. 2003. Stable isotope evidence of diet at Neolithic Çatalhöyük, Turkey. *Journal of Archaeological Science*, 30, 67-76.
- Rose, J. C. 2001. *Tell Ya'amün 2001 Excavations*. [Online]. University of Arkansas. Available: <http://cavern.uark.edu/~jcrose/yaamun1/> [Accessed 21/05/2014].
- Rose, J. C. 2002. *Ya'amün 2002 Report* [Online]. University of Arkansas. Available: <http://cavern.uark.edu/~jcrose/yaamun02/yaamun2.html> [Accessed 21/05/2014].
- Rose, J. C. 2005. *Tell Ya'amün: The 2005 Season* [Online]. University of Arkansas. Available: <http://cavern.uark.edu/~jcrose/fieldseason2005/> [Accessed 21/05/2014].
- Rose, J. C., El-Najjar, M. and Burke, D. L. 2007. Trade and the acquisition of wealth in rural Late Antique North Jordan. In: Harahsheh, R., Fakhoury, Q., Taher, H. & Khouri, S. (eds.) *Studies in the History and Archaeology of Jordan*. Amman: Department of Antiquities of Jordan, 61-70.
- Rose, J. C., El-Najjar, M. Y., Hunton, C. I. and Rolf, K. 2003. Tell Ya'amün in Savage, S.H., Zamora, K.A, Keller, D.L. (eds) *Archaeology in Jordan*. *American Journal of Archaeology*, 107, 457-458.
- Rosen, A. M. 2007. *Civilizing Climate. Social Responses to Climate Change in the Ancient Near East*, Lanham, MD, Altamira Press.
- Safrai, Z. 2004. *The Economy of Roman Palestine*, London, Taylor and Francis.
- Schoeninger, M. J. and DeNiro, M. J. 1984. Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica et Cosmochimica Acta*, 48, 625-639.
- Schwarcz, H. P., Dupras, T. L. and Fairgrieve, S. I. 1999. ¹⁵N enrichment in the Sahara: in search of a global relationship. *Journal of Archaeological Science*, 26, 629-636.

- 671 Sealy, J. C., van der Merwe, N., Lee Thorp, J. A. and Lanham, J. L. 1987. Nitrogen isotopic ecology
672 in southern Africa: implications for environmental and dietary tracing. *Geochimica et*
673 *Cosmochimica Acta*, 51, 2707-2717.
- 674 Shomer-Ilan, A., Nissenbaum, A. and Waisel, Y. 1981. Photosynthetic pathways and the ecological
675 distribution of the Chenopodiaceae in Israel. *Oecologia*, 48, 244-248.
- 676 Smith, B. N. and Epstein, S. 1971. Two categories of $^{13}\text{C}/^{12}\text{C}$ ratios for higher plants. *Plant*
677 *Physiology*, 47, 380-384.
- 678 Snell, D. C. 1997. *Life in the Ancient Near East*, New Haven and London, Yale University Press.
- 679 Strange, J. 2008. The Late Bronze Age. In: Adams, R. B. (ed.) *Jordan. An Archaeological Reader*.
680 London: Equinox, 281-310.
- 681 Thompson, A. H., Chaix, L. and Richards, M. P. 2008. Stable isotopes and diet in Ancient Kerma,
682 Upper Nubia (Sudan). *Journal of Archaeological Science*, 35, 376-387.
- 683 Thompson, A. H., Richards, M. P., Shortland, A. and Zakrzewski, S. R. 2005. Isotopic palaeodiet
684 studies of Ancient Egyptian fauna and humans. *Journal of Archaeological Science*, 32, 451-
685 463.
- 686 Tieszen, L. L. and Fagre, T. 1993. Effect of diet quality and composition on the isotopic
687 composition of respiratory CO_2 , bone collagen, bioapatite and soft tissues. In: Lambert, J. B.
688 & Grupe, G. (eds.) *Prehistoric Human Bone, Archaeology at the Molecular Level*. Berlin:
689 Springer, 121-155.
- 690 Tubb, J. N. 1988. Tell es-Sa'idiyeh: preliminary report on the first three seasons of renewed
691 excavations. *Levant*, 20, 23-88.
- 692 Tubb, J. N., Dorrell, P. G. and Cobbing, F. J. 1997. Interim report of the ninth season (1996) of
693 excavations at Tell es-Sa'idiyeh, Jordan. *Palestine Exploration Quarterly*, 129, 54-77.
- 694 van der Merwe, N. J. 1989. Natural variations in ^{13}C concentration and its effect on environmental
695 reconstruction using $^{13}\text{C}/^{12}\text{C}$ ratios in animal bones. In: Price, T. D. (ed.) *The Chemistry of*
696 *Prehistoric Human Bone*. Cambridge: Cambridge University Press, 105-125.
- 697 van Klinken, G. J., van der Plicht, H. and Hedges, R. E. M. 1994. Bone $^{13}\text{C}/^{12}\text{C}$ ratios reflect
698 (palaeo-) climatic variations. *Geophysical Research Letters*, 21, 445-448.
- 699 Van Neer, W. and Parker, T. S. 2008. First archaeozoological evidence for *haimation*, the "invisible"
700 *garum*. *Journal of Archaeological Science*, 35, 1821-1827.
- 701 Van Neer, W., Wildekamp, R., Waelkens, M., Arndt, A. and Volckaert, F. 2000. Fish as indicators
702 of trade relationships in Roman times: the example of Sagalassos, Turkey. In: Mashkour,
703 M., Choyke, A. M., Buitenhuis, H. & Poplin, F. (eds.) *International Symposium on the*
704 *Archaeozoology of Southwestern Asia and Adjacent Areas (4th: 1998 : Paris, France):*
705 *Archaeozoology of the Near East IVB. Proceedings of the fourth international symposium*
706 *on the archaeozoology of south western Asia and adjacent areas*. Groeningen: Centre for
707 Archaeological Research and Consultancy, 206-215.
- 708 Vanderklift, M. A. and Ponsard, S. 2003. Sources of variation in consumer-diet $\delta^{15}\text{N}$ enrichment: a
709 meta-analysis. *Oecologia*, 136, 169-182.
- 710 Vika, E. and Theodoropoulou, T. 2012. Re-investigating fish consumption in Greek antiquity:
711 results from $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ analysis from fish bone collagen. *Journal of Archaeological*
712 *Science*, 39, 1618-1627.
- 713 Vogel, J. C., Fuls, A. and Danin, A. 1986. Geographical and environmental distribution of C_3 and C_4
714 grasses in the Sinai, Negev, and Judean deserts. *Oecologia*, 70, 258-265.

715 Watson, P. M. 2008. The Byzantine Period. *In*: Adams, R. B. (ed.) *Jordan. An Archaeological*
 716 *Reader*. London: Equinox, 443-482.
 717 Wilkins, J. M. and Hill, S. 2006. *Food in the Ancient World*, Oxford, Blackwell.
 718 Winter, K. 1981. C₄ plants of high biomass in arid regions of Asia - Occurrence of C₄
 719 photosynthesis in Chenopodiaceae and Polygonaceae from the Middle East and USSR.
 720 *Oecologia*, 48, 100-106.
 721 Zohary, D. and Hopf, M. 2000. *Domestication of plants in the Old World. Third Edition*, Oxford,
 722 Oxford University Press.
 723 Zohary, M. 1973. *Geobotanical foundations of the Middle East*, Stuttgart, Gustav Fischer Verlag.
 724 Zohary, T., Erez, J., Gophen, M., Berman-Frank, I. and Stiller, M. 1994. Seasonality of stable
 725 carbon isotopes within the pelagic food web of Lake Kinneret. *Limnology and*
 726 *Oceanography*, 39, 1030-1043.
 727

728

Table 1. Descriptive statistics (group sizes, mean $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$, standard deviations and minimum and maximum values) for human and faunal samples from Ya'amūn

Group	n	mean $\delta^{13}\text{C}$ (‰) (min – max)	s.d.	mean $\delta^{15}\text{N}$ (‰) (min – max)	s.d.
MB and LB sheep/goats	6	-18.6 (-20.1 – -16.6)	1.4	8.5 (5.8 – 11.4)	1.9
LR/Byz sheep/goats	5	-19.3 (-20.1 – -19.6)	0.5	5.3 (4.0 – 6.4)	0.9
LR/Byz cattle	2	-18.8 – -15.6		5.4 – 9.7	
MB and LB humans	15	-18.8 (-19.4 – -18.3)	0.3	8.8 (7.4 – 9.8)	0.7
LR/Byz humans	17	-19.1 (-19.8 – -18.5)	0.3	8.1 (7.3 – 9.0)	0.6
Freshwater fish	3	-16.8 (-21.5 – -10.3)	5.8	6.9 (6.1 – 9.3)	2.0
Marine fish	2	-7.2 (-9.7 0 -4.7)	3.5	10.4 (8.2 – 12.6)	3.1

Ref: JASC14-754

Title: Diet and herding strategies in a changing environment: stable isotope analysis of Bronze Age and Late Antique skeletal remains from Ya'amūn, North Jordan

Figure Legends


Figure 1. Map of Jordan with sites mentioned in the text superimposed on **(1.a)** the modern phytogeographic zones (based on data from Zohary 1973 and Al-Eisawi 1985, redrawn from Cordova 2007, Figure P.1, and Palmer 2013, Figure I.18) and **(1.b)** a rainfall map of modern Jordan (redrawn from Kennedy 2007, fig. 3.4a).

Figure 2. Carbon and nitrogen isotope data of Mid- and Late Bronze Age and Late Roman/Byzantine sheep/goats from Ya'amūn. $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of the Bronze Age sheep/goats are positively correlated (Pearson's $r=0.87$, $p=0.026$) and therefore consistent with varying consumption of plants from arid and semi-arid environments.


Figure 3. Carbon and nitrogen isotope data of Mid- and Late Bronze Age and Late Roman/Byzantine humans from Ya'amūn in comparison with mean values ($\pm 1\text{sd}$) for sheep/goats from these time periods. The variation in human values to a large extent mirrors that observed in the fauna.

Figure 4. Carbon and nitrogen isotope data for humans (individual data) and sheep/goats (mean values $\pm 1\text{sd}$) from Ya'amūn in comparison with $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values for marine (one specimen each of family *Scienidae* and *Mugilidae*) and freshwater fish (two specimens of *Tilapia* sp. and one of *Clarias* (catfish)) from the same site.


Figure




Figure




Figure



Figure



Figure



1 **Supplementary Information**

2 **Diet and herding strategies in a changing environment: stable isotope analysis of** 3 **Bronze Age and Late Antique skeletal remains from Ya'amūn, Jordan**

4 **Sample Preparation and Analytical Methods**

5 Cortical bone from the diaphysis of long bones, taken from areas devoid of any
6 pathological lesions, was the preferred sampling material. Samples consisted of bone
7 chunks weighing between 200 and 300 mg. All outer surfaces were abraded with the aid of
8 a drill before collagen extraction was carried out following the Longin ((1971) method
9 modified according to recommendations by Collins and Galley (1998). . Briefly, bone
10 chunks were demineralised in 0.5 M in the fridge for several days, after which they were
11 rinsed to neutrality with ultrapure water (Milli-Q®). The samples were then placed in a pH3
12 HCl solution and gelatinised in a heater-block at 70 degrees C for 48 hours. Acid insoluble
13 residues were removed with the aid of an Ezee®-filter (60-90µm, Elkay) and the
14 remaining solutions were frozen and then freeze-dried for 48h. Aliquots of between 0.9 and
15 1.1 mg of freeze-dried 'collagen' were weighed in duplicates into ultraclean tin capsules.

16 Carbon and nitrogen stable isotope compositions of samples were determined by analysis
17 on a Europa Geo 20-20 Continuous Flow Isotope Ratio Mass Spectrometer (CF-IRMS)
18 interfaced with Sercon® elemental analyser (EA) in the School of Human and
19 Environmental Sciences, University of Reading, UK. All $\delta^{13}\text{C}$ values are expressed relative
20 to Pee Dee Belemnite (V-PDB) and while the $\delta^{15}\text{N}$ values are referred to atmospheric
21 nitrogen (AIR). The analytical error was calculated from repeat analysis of internal collagen
22 standards included in each run and was determined at $\pm 0.2\text{‰}$ (1sd) or better for $\delta^{13}\text{C}$,
23 and $\delta^{15}\text{N}$ measurements. Internal working standards which were calibrated to
24 internationally certified reference materials included the amino acid methionine (Elemental
25 Microanalysis/MethR), powdered Bovine Liver Standard (NIST1577a/BLS) and a batch of
26 pork gelatine prepared at the Reading stable isotope laboratory ("Reading Pork
27 Gelatine"/RPG).

28 Collagen samples were considered of acceptable quality when having an atomic C:N ratio
29 between 2.9 and 3.6, $\% \text{C} \geq 13\%$ and $\% \text{N} \geq 4.8\%$ (Ambrose, 1990, DeNiro, 1985).
30 Samples that yielded less than 1% collagen were still regarded as acceptable if they

fulfilled these criteria and if their $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ ratios were not unusual within the population context (van Klinken 1999). Inferential statistics were computed with SPSS v.19.

References

- Ambrose, S. H. 1990. Preparation and characterisation of bone and tooth collagen for isotopic analysis. *Journal of Archaeological Science*, 17, 431-451.
- Collins, M.J., Galley, P. 1998. Towards an Optimal Method of Archaeological Collagen Extraction: The Influence of pH and Grinding. *Ancient Biomolecules* 2: 209-222.
- DeNiro, M. J. 1985. Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature*, 317, 806-809.
- van Klinken, G.J. 1999. Bone Collagen Quality Indicators for Palaeodietary and Radiocarbon Measurements. *Journal of Archaeological Science* 26: 687-695.

45 **Table S1.** Carbon and nitrogen stable isotope data and collagen quality indicators of
 46 faunal bone samples from Ya'amūn. Archaeological dates are abbreviated as follows:
 47 MBA=Middle Bronze Age; LBA=Late Bronze Age; LR/Byz=Late Roman/Byzantine. Nil
 48 refers to samples which yielded no collagen for analysis.

Species	Sample Code	Date	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	%C	%N	C:N	%Coll
<i>Bos</i>	YMN fb 084	MBA						Nil
Sheep/Goat	YMN fb 085	MBA						Nil
Sheep/Goat	YMN fb 086	MBA	-19.7	8.5	40.3	14.0	3.4	1.3
Sheep/Goat	YMN fb 087	MBA	-16.6	11.4	19.5	6.8	3.4	2.6
Fish (<i>Sciaenidae</i>).	YMN fb 088	MBA	-9.7	12.6	40.0	14.3	3.3	6.6
Sheep/Goat	YMN fb 089	MBA	-19.4	7.3	41.1	13.8	3.5	0.8
Sheep/Goat	YMN fb 090	MBA						Nil
Sheep/Goat	YMN fb 074	LBA						Nil
Sheep/Goat	YMN fb 075	LBA	-20.1	5.8	42.8	15.0	3.3	3.1
Sheep/Goat	YMN fb 076	LBA	-17.3	8.8	42.8	15.2	3.3	2.7
Sheep/Goat	YMN fb 077	LBA	-18.2	9.1	42.6	15.1	3.3	7.2
Sheep/Goat	YMN fb 078	LBA						Nil
Sheep/Goat	YMN fb 092	LR/Byz	-19.3	5.0	41.6	15.0	3.2	9.0
Sheep/Goat	YMN T fb 093	LR/Byz	-18.8	5.7	39.0	14.2	3.2	12.0
<i>Bos</i>	YMN T fb 094	LR/Byz	-18.8	5.4	42.3	15.2	3.3	3.4
<i>Bos</i>	YMN T fb 095	LR/Byz	-15.6	9.7	44.1	16.0	3.2	16.0
Sheep/Goat	YMN T fb 096	LR/Byz	-20.1	6.4	38.0	13.3	3.3	4.1
Sheep/Goat	YMN T fb 097*	LR/Byz	-18.9	4.0	43.6	16.0	3.2	18.2
Sheep/Goat	YMN T fb 098	LR/Byz	-19.6	5.4	41.1	14.7	3.3	9.2
Fish (<i>Mugilidae</i>)	YMN T fb 099	LR/Byz	-4.7	8.2	43.1	15.8	3.2	12.0
Fish (<i>Tilapia</i> sp.)	YMN T fb 100	LR/Byz	-21.5	6.1	43.2	15.7	3.2	10.0
Fish (<i>Tilapia</i> sp.)	YMN T fb 101	LR/Byz	-10.3	5.4	42.4	15.5	3.2	9.4
Fish (<i>Clariidae</i>)	YMN T fb 102	LR/Byz	-18.6	9.3	41.2	14.9	3.2	6.0

49

50

51 **Table S2.** Carbon and nitrogen stable isotope data and collagen quality indicators of
52 human bone samples from Ya'amūn. Archaeological dates are abbreviated as follows:
53 MBA=Middle Bronze Age; LBA=Late Bronze Age; LR=Late Roman; Byz= Byzantine. Nil
54 refers to samples which yielded no collagen for analysis.

Sample Code	Date	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	%C	%N	C:N	%Coll
YMN hb 001	MBA	-19.4	8.6	40.4	14.0	3.4	0.9
YMN hb 002	MBA	-19.1	8.4	42.2	14.8	3.3	4.0
YMN hb 003	MBA	-18.9	9.5	42.2	14.7	3.3	2.0
YMN hb 004	MBA	-19.2	7.4	41.5	13.9	3.5	0.6
YMN hb 005	MBA						Nil
YMN hb 006	MBA	-19.0	8.4	41.6	13.5	3.6	1.3
YMN hb 007	MBA						Nil
YMN hb 008	MBA						Nil
YMN hb 009	MBA	-18.5	9.1	41.0	14.6	3.3	4.3
YMN hb 010	MBA	-18.3	9.4	43.4	15.6	3.2	9.3
YMN hb 011	MBA	-18.4	9.7	43.8	15.7	3.3	7.5
YMN hb 012	MBA						Nil
YMN hb 013	MBA	-18.6	8.8	42.9	15.4	3.3	9.0
YMN hb 014	MBA	-18.9	8.1	43.3	15.7	3.2	10.0
YMN hb 015	MBA	-18.3	9.8	42.4	15.3	3.2	5.1
YMN hb 016	MBA	-18.8	8.6	43.0	15.7	3.2	14.5
YMN hb 017	LBA						Nil
YMN hb 018	LBA						Nil
YMN hb 019	LBA	-18.7	9.0	42.8	15.4	3.2	6.0
YMN hb 020	LBA						Nil
YMN hb 021	LBA	-18.7	9.5	42.0	15.1	3.2	2.8
YMN hb 022	LBA	-18.5	8.1	39.0	13.8	3.3	4.0
YMN hb 023	LR						Nil
YMN hb 024	LR	-19.1	8.5	41.4	15.1	3.2	10.0
YMN hb 025	LR	-18.7	9.0	42.8	15.3	3.3	10.0
YMN hb 026	LR	-19.5	7.7	40.9	14.1	3.4	1.3
YMN hb 027*	Byz	-18.5	8.6	43.0	15.5	3.2	17.0
YMN hb 028	Byz	-19.1	7.3	38.5	13.5	3.3	3.5
YMN hb 029	Byz	-19.3	7.4	39.1	13.6	3.4	3.6
YMN hb 030	Byz	-19.2	8.0	35.9	12.5	3.3	2.6
YMN hb 031	Byz	-19.8	6.8	42.0	14.9	3.3	3.6
YMN hb 032	Byz	-19.1	7.7	43.4	15.3	3.3	3.3
YMN hb 033	Byz	-19.5	7.8	28.9	9.9	3.4	3.7
YMN hb 034	LR/Byz	-19.0	7.6	33.0	11.7	3.3	5.8
YMN hb 035	LR/Byz	-19.2	8.6	38.3	13.5	3.3	7.2
YMN hb 036	LR/Byz						Nil
YMN hb 037	LR/Byz						Nil
YMN hb 038	LR/Byz	-19.2	8.3	43.0	15.0	3.4	6.0
YMN hb 039	LR/Byz	-19.2	8.1	41.6	14.3	3.4	4.2
YMN hb 040	LR/Byz						Nil
YMN hb 041	LR/Byz						Nil
YMN hb 042	LR/Byz						Nil
YMN hb 043	LR/Byz	-18.7	9.0	43.4	15.6	3.2	6.8
YMN hb 044*	LR/Byz	-19.2	8.4	43.7	15.7	3.2	19.5
YMN hb 045	LR/Byz	-19.1	8.7	42.8	15.1	3.3	11.5

