

# *Atmosphere-ocean coupled processes in the Madden-Julian oscillation*

Article

Accepted Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

DeMott, C. A., Klingaman, N. P. and Woolnough, S. J. (2015) Atmosphere-ocean coupled processes in the Madden-Julian oscillation. *Reviews of Geophysics*, 53 (4). pp. 1099-1154. ISSN 8755-1209 doi: <https://doi.org/10.1002/2014RG000478> Available at <http://centaur.reading.ac.uk/43966/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/2014RG000478>

Publisher: American Geophysical Union

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1 **Thermal and pressure stability of myrosinase enzymes from black mustard (*Brassica nigra***  
2 **L. W.D.J Koch. var. *nigra*), brown mustard (*Brassica juncea* L. Czern. var. *juncea*) and**  
3 **yellow mustard (*Sinapsis alba* L. Subsp *Maire*) seeds**

4

5

6

7 Olukayode Adediran Okunade, Sameer Khalil Ghawi, Lisa Methven, Keshavan Niranjana

8 Department of Food and Nutritional Sciences, University of Reading, Whiteknights,

9 P.O Box 226, Reading, RG6 6AP, UK

10

11

12 **Abbr. Running Title:**

13 **Thermal and pressure stability of myrosinase from black, brown and yellow mustard**

14

15 Corresponding Author;

16 Olukayode Adediran Okunade

17 Department of Food and Nutritional Sciences, University of Reading, Whiteknights,

18 P.O Box 226, Reading, RG6 6AP, UK E-mail: fc030053@reading.ac.uk

19

20

21

22

23

24

25

26

27

28 **Abstract**

29 This study investigates the effects of temperature and pressure on inactivation of  
30 myrosinase extracted from black, brown and yellow mustard seeds.

31 Brown mustard had higher myrosinase activity (2.75 un/mL) than black (1.50 un/mL) and  
32 yellow mustard (0.63 un/mL).

33 The extent of enzyme inactivation increased with pressure (600-800 MPa) and temperature  
34 (30-70 °C) for all the mustard seeds. However, at combinations of lower pressures (200-400  
35 MPa) and high temperatures (60-80 °C), there was less inactivation. For example,  
36 application of 300 MPa and 70 °C for 10 minutes retained 20%, 80% and 65% activity in  
37 yellow, black and brown mustard, respectively, whereas the corresponding activity  
38 retentions when applying only heat (70 °C, 10min) were 0%, 59% and 35%. Thus,  
39 application of moderate pressures (200-400 MPa) can potentially be used to retain  
40 myrosinase activity needed for subsequent glucosinolate hydrolysis.

41

42

43

44 **Keywords:** Processing, Myrosinase activity, Inactivation, Mustard seed.

45

46

47

48

49

## 50        **1    Introduction**

51    Mustard plant belongs to the *Brassicaceae* family, the dry seeds being the main part used in  
52    food processing (canned mustard leaves are available). Common types of mustard are  
53    yellow mustard (*Sinapsis alba*), brown (*Brassica juncea*) and black mustard (*Brassica nigra*).  
54    Mustard has a rich chemical composition and its seed flour is widely used in food processing  
55    (Abul-Fadl, El-Badry, & Ammar, 2011; Wanasundara, 2008). Mustard is also used for its spicy  
56    flavour, produced from the hydrolysis of glucosinolates by myrosinase enzymes  
57    (Wanasundara, 2008). Mustard seed is widely used as a condiment, however, its  
58    advantageous chemical composition and relatively low price offer wide possibilities for  
59    utilization as additives in human food and in animal feeds (Abul-Fadl, El-Badry, & Ammar,  
60    2011; Wanasundara, 2008).

61    The glucosinolates (thioglucosides) in mustard seeds are pseudo-thioglucosides containing  
62    nitrogen and sulphur. Myrosinase enzymes (thioglucoside) are glucohydrolases (EC 3.2.3.1)  
63    (Bones & Rossiter, 1996; Fahey, Zalcmann, & Talalay, 2001; Thangstad & Bones, 1991).  
64    According to Thangstad *et al.* (1991) and Bones *et al.* (1996), glucosinolates and myrosinase  
65    enzymes coexist in segregated compartments of the plant. After plant tissue damage,  
66    glucosinolates are hydrolysed to produce a variety of compounds; some of which are  
67    bioactive (isothiocyanates, indoles) (Bongoni, Verkerk, Steenbekkers, Dekker, & Stieger,  
68    2014), thiocyanates, oxazolidine-2-thiones and others of which are potentially toxic (nitriles,  
69    epithionitriles) by myrosinase enzymes. The nature of the hydrolysis products depends on  
70    the structure of glucosinolate and the reaction conditions (Fahey, Zalcmann, & Talalay,  
71    2001; Lambrix, Reichelt, Mitchell-Olds, Kliebenstein, & Gershenzon, 2001). These  
72    compounds are of immense interest in food processing (taste, aroma, and flavour  
73    attributes) and human health (anticarcinogenic and antimicrobial properties) (Drewnowski

74 & Gomez-Carneros, 2000; Fahey, Zhang, & Talalay, 1997; Johnson, Koh, Wang, Yu, & Yuan,  
75 2010; Tang & Zhang, 2004; Wanasundara, 2008).

76 Myrosinase enzymes are significantly inactivated at normal cooking temperatures regardless  
77 of the method used (Ludikhuyze, Ooms, Weemaes, & Hendrickx, 1999; Van Eylen, Oey,  
78 Hendrickx, & Van Loey, 2007; Yen & Wei, 1993), although, the same processing conditions  
79 rarely affect glucosinolates if leaching out can be avoided (Oerlemans, Barrett, Suades,  
80 Verkerk, & Dekker, 2006). Microflora in the human gut can hydrolyse glucosinolates into  
81 bioactive compounds, but the yield is much lower compared to that resulting from plant  
82 myrosinases (Conaway, Getahun, Liebes, Pusateri, Topham, Botero-Omary, *et al.*, 2001).  
83 Hence, the control of myrosinase activity is important to determine the bioavailability of  
84 hydrolysis products.

85 Myrosinase enzymes in *Brassica* vegetables are known to exhibit varying degree of thermal  
86 stability (Ghawi, Methven, Rastall, & Niranjana, 2012; Ludikhuyze, Ooms, Weemaes, &  
87 Hendrickx, 1999; Matusheski, Juvik, & Jeffery, 2003; Van Eylen, Indrawati, Hendrickx, & Van  
88 Loey, 2006; Yen & Wei, 1993). Yellow mustard and rapeseed myrosinases are known to have  
89 the highest thermal stability of *Brassica* plant species (Kozłowska, Nowak, & Nowak, 1983;  
90 Ludikhuyze, Ooms, Weemaes, & Hendrickx, 1999; Pérez, Barrientos, Román, & Mahn, 2014;  
91 Van Eylen, Indrawati, Hendrickx, & Van Loey, 2006; Van Eylen, Oey, Hendrickx, & Van Loey,  
92 2007; Verkerk & Dekker, 2004; Yen & Wei, 1993). The use of high pressure processing (HPP)  
93 as an alternative to thermal processing has been suggested and it has been found to reduce  
94 thermal inactivation of certain enzymes, including myrosinase (Hendrickx, Ludikhuyze, Van  
95 den Broeck, & Weemaes, 1998; Van Eylen, Oey, Hendrickx, & Van Loey, 2008). Although a  
96 narrow range of thermal and pressure stability in *Brassica* vegetables has been reported, no

97 data is yet available for the thermal and pressure inactivation of brown and black mustard  
98 seed myrosinase enzymes; and yet, they may be more stable than myrosinase in other  
99 *Brassic*as as they are related to yellow mustard which has already been shown to have  
100 higher heat stability (Van Eylen *et al.*, 2006; 2008).

101 In recent studies, it was found that addition of an exogenous source of myrosinase (Daikon  
102 radish root and mustard seeds) to processed *Brassica* can reinitiate the hydrolysis of  
103 glucosinolates (Dosz & Jeffery, 2013; Ghawi, Methven, & Niranjana, 2013). Hence, evaluating  
104 other sources of myrosinase that are more stable under processing conditions is of  
105 importance.

106 The hypothesis of this study were that (1) myrosinase from different mustard species differ  
107 in thermal and pressure stability and (2) lower pressure processing can be used to decrease  
108 thermal inactivation rate of mustard myrosinase. The study aimed to investigate thermal,  
109 pressure and combined thermal and pressure inactivation of myrosinase enzymes from  
110 black, brown and yellow mustard seeds in order to ascertain the possible extent of  
111 degradation that would occur under food processing conditions.

112

113

114

115

116

117

## 118        **2    Materials and methods**

### 119        **2.1   Sample preparation and myrosinase extraction**

120            Yellow mustard (*Sinapsis alba* L. *subsp maire*), black mustard (*Brassica nigra* L. W.D.J  
121 Koch. *var. nigra*) and brown mustard (*Brassica juncea* L. Czern. *var. juncea*) were obtained  
122 from the I.P.K Gene bank (Gatersleben, Germany). All samples were obtained as dried  
123 seeds.

124            Myrosinase enzyme extraction was done as described by Ghawi *et al.* (2012). The process  
125 involved grinding dry mustard seeds in a coffee grinder. 10mL of buffer (Tris HCl 0.2M, pH  
126 7.5 containing EDTA 0.5mM, dithiothreitol 1.5mM and 0.4g Polyvinylpolypyrrolidone) was  
127 then added to 0.5g powdered mustard and blended on ice (15 minutes) and then  
128 centrifuged (11,738 ×g) for 15 minutes at 4 °C. The supernatant was filtered (0.45µm). The  
129 filtrate was made up to 10mL using the buffer solution and 90% precipitation of protein was  
130 achieved using 6.2g ammonium sulphate with slow blending on ice for 30 minutes. The  
131 mixture was then centrifuged (13,694 ×g) for 15 minutes at 4 °C. The pellet obtained was  
132 suspended in 2mL of 10mM Tris HCl buffer, pH 7.5.

133            The mix was extensively dialysed at low temperature (4 °C) using cellulose membrane  
134 (Medicell International Ltd, Molecular weight cut-off 12,000-14,000 Da) and 10mM Tris HCl  
135 buffer for 24 hours to remove excess ammonium and sulphate ions and centrifuged (11,738  
136 ×g) at 4 °C for 15 minutes to remove insoluble materials. Finally, the supernatant was frozen  
137 (-80 °C) and then lyophilised, the resulting powder was stored at -20 °C until further analysis.

### 138        **2.2   Myrosinase enzyme activity assay**

139            Myrosinase activity was measured according to the coupled enzymatic procedure with  
140 some modifications (Gatfield & Sand, 1983; Ghawi, Methven, Rastall, & Niranjana, 2012;



141 Wilkinson, Rhodes, & Fenwick, 1984). A D-glucose determination kit was used (R-Biopharm  
142 Rhone, Heidelberg, Germany). The reaction mixture (1.51mL) included 0.5mL NADP/ATP, 10  
143  $\mu$ L hexokinase/glucose-6-phosphate dehydrogenase, 0.9mL of water containing ascorbic  
144 acid 7mM (cofactor) and 50  $\mu$ L sample. The mix was allowed to equilibrate for 5 minutes  
145 and 50  $\mu$ L sinigrin solution (0.6M) was added. The change in absorbance due to the  
146 formation of NADP was measured at 340nm. Myrosinase activity was determined from the  
147 initial linear rate of increase in the curve of absorbance against reaction time. A standard  
148 myrosinase enzyme (Sigma Aldrich, UK) was used to establish the calibration curve of  
149 absorbance against concentration. One unit (un) of myrosinase was defined as the amount  
150 of enzyme that produces 1  $\mu$ mol of glucose per minute at 25 °C and pH 7.5.

### 151 **2.3 Heat treatment**

152 Thermal inactivation was done under isothermal conditions at different temperatures,  
153 between 10-80 °C. 150  $\mu$ L sample (25mg lyophilised powder/mL de-ionised water) was  
154 pipetted into clean durham tube (6.5mm internal diameter, 1mm thickness and 30mm in  
155 length, Fischer Scientific, Loughborough, U.K) and sealed. These were carefully placed in  
156 heated water bath fitted with a thermometer for pre-set time of 5 and 10 minutes at  
157 different temperature (10-80 °C). After each determination, the samples were quickly  
158 immersed in an ice bath and the enzyme activity was measured not later than an hour after  
159 each heat treatment. Treatments were done in triplicate.

### 160 **2.4 Pressure treatment**

161 Pressure treatments were performed between 100-900 MPa using a high pressure  
162 unit (37mm diameter and 246mm length Food Lab 300 Stansted Fluid Power, Stansted, UK).  
163 1, 2-Propanediol (30%) (sigma-Aldrich, Poole, U.K) was used as the pressure transmitting

164 fluid. The processing temperature was controlled by liquid circulation in the outer jacket of  
165 the high pressure vessel. 200  $\mu$ L samples (25mg/mL) were placed in flexible polyethylene  
166 bags (LDPE) and air was carefully removed from the bags before sealing. Pressure treatment  
167 at different levels for pre-set time of 5 and 10 minutes was applied with temperature  
168 controlled at 15°C. Samples were removed from the vessel and rapidly cooled in an ice bath  
169 and the enzyme activity was measured not later than an hour after pressure treatment.

170 Combined pressure and temperature treatments were performed using a combination of  
171 high pressure (600-800MPa) with moderate temperature (30-70 °C) for 10 minutes and low  
172 pressure (200- 400MPa) with slightly higher temperature (60-80 °C) for different pre-set  
173 times of 5, 10 and 15 minutes. Pressure build up is usually accompanied with increase in  
174 temperature due to adiabatic heating. About 3-5 minutes was needed to reach equilibrium  
175 (desired temperature and pressure) and this was added to the holding time. All treatments  
176 were done in triplicate.

## 177 **2.5 Protein assay**

178 Total protein content (unprocessed samples) was determined using the Bradford  
179 procedure (Bradford, 1976). This is based on formation of a complex with Brilliant Blue G.  
180 The samples (25mg/mL) were mixed with the reagent and the absorbance was measured at  
181 595nm after 20 minutes of incubation at 23 °C. Bovine serum albumin BSA (0 - 1.4mg/mL)  
182 (Sigma Aldrich, UK) was used in constructing a standard curve.

## 183 **2.6 Statistical analysis**

184 The statistical differences between the values obtained under different experimental  
185 conditions were established by undertaking ANOVA followed by Tukey's HSD multiple

186 pairwise comparison test using SPSS software (PASW Statistics 17.0, IBM, UK). Differences  
187 were considered significant at  $P < 0.05$ .

188

189

190

191

192

193

194

195

196

197

198

199

200

## 201 **3 Results and discussion**

### 202 **3.1 Enzyme activity, protein content and specific enzyme activity of myrosinase** 203 **enzymes from black, brown and yellow mustard seeds**

204 Table 1 shows the enzyme activity, protein content and specific enzyme activity of  
205 myrosinase enzyme from black, brown and yellow mustard. Brown mustard had higher  
206 myrosinase activity (2.75 un/mL) than black mustard (1.50 un/mL) and yellow mustard had  
207 the least myrosinase activity (0.63 un/mL). The protein content of all studied samples was  
208 similar. Myrosinase from brown mustard had the highest proportion of specific activity  
209 (2.04un/mg) and yellow mustard the lowest (0.48un/mg). The differences between mustard  
210 species in overall enzyme activity were not related to differences in protein content and  
211 hence, differences between species in specific activity prevailed. Variations in myrosinase  
212 enzyme activity within and between *Brassica* species have been reported previously and  
213 have been attributed to genetic and/or environmental factors (agronomic and climatic  
214 conditions) (Pocock, Heaney, Wilkinson, Beaumont, Vaughan, & Fenwick, 1987; Wilkinson,  
215 Rhodes, & Fenwick, 1984). In addition, Rask *et al.* (2000) had also reported that the  
216 difference in thermal stability of myrosinase in *Brassica* plants was probably due to the  
217 existence of different isoforms of myrosinase, where some of them interact with  
218 myrosinase-binding proteins (a group of proteins found in *Brassica* plants) to form  
219 complexes that may improve stability. It is therefore possible that the different species of  
220 mustard may have genetic differences, or have adapted to different environmental  
221 conditions that have resulted in different isoforms of myrosinase. It has also been suggested  
222 that myrosinase activity for similar *Brassica* samples may vary between studies merely due  
223 to the different protocols employed (Piekarska, Kusznerewicz, Meller, Dziedziul, Namiesnik,  
224 & Bartoszek, 2013). In this study, the observed differences in enzyme activity between the

225 three species cannot be specifically attributed to one or more of the above factors because,  
226 even though the seeds were obtained from the same gene bank, they may not have been  
227 produced under strictly controlled conditions for drawing such inferences. However, the  
228 main purpose of this study is to evaluate the effects of processing on enzyme activity  
229 retention, which can robustly be undertaken for a given seed variety by normalising the  
230 enzyme activity after processing with the corresponding initial activity.

### 231 **3.2 Effect of temperature on black, brown and yellow mustard seed myrosinase**

232 Figure 1 shows the effect of temperature and exposure time on myrosinase activity at  
233 atmospheric pressure. Exposure time of 10 minutes at temperature ranging between 10-80  
234 °C were employed. Recent studies have shown that myrosinase enzymes extracted from  
235 broccoli are stable up to 70 °C by blanching intact broccoli florets (Pérez, Barrientos, Román,  
236 & Mahn, 2014) whereas the enzyme extracted from yellow mustard is stable up to 60 °C  
237 (Ludikhuyze, Ooms, Weemaes, & Hendrickx, 1999; Van Eylen, Indrawati, Hendrickx, & Van  
238 Loey, 2006). Results of heating for 5 minutes (data not included) showed no significant  
239 inactivation up to 60 °C for black mustard myrosinase whereas the myrosinase from brown  
240 and yellow mustard showed activity loss of approximately 28% and 17% respectively.  
241 Myrosinase activity decreased significantly at above 60 °C in all the mustards studied.

242 When the exposure time was 10 minutes, the same trend, as observed for 5 minutes  
243 processing time, was observed in the case for both black and brown mustard. However,  
244 yellow mustard myrosinase was only substantially stable up to 50 °C (27% loss in activity)  
245 and lost about 79% of its activity at 60 °C. Heating up to 70 °C led to about 41% loss in  
246 activity for black mustard, 65% for brown mustard while there was no myrosinase activity

247 for yellow mustard. At 80 °C, there was no significant myrosinase activity in the case of all  
248 three seeds.

249 These results are similar to those in previous studies; Van Eylen *et al.* (2006 & 2008)  
250 concluded that the inactivation of myrosinase extracted from yellow mustard occurs at  
251 temperatures above 60 °C at an exposure time of 10 minutes. Stoin *et al.* (2009) had earlier  
252 reported that myrosinase from black mustard exhibited maximum activity at temperature  
253 ranging from 45-50 °C and even at a range of 70-85 °C, a small amount of enzyme activity  
254 could be observed. However, the authors suggested that activity at temperature above 85  
255 °C may be from other sources for example thermostable desulphatase enzyme using sinigrin  
256 as a substrate. The current study has concluded that myrosinases from black and brown  
257 mustard are fully inactivated at temperatures above 80 °C and that myrosinase from these  
258 mustard sources is more stable than that from yellow mustard.

259 Comparing to heat stability from other *Brassica* sources, studies have shown that in both  
260 white and red cabbage, 90% loss in myrosinase activity was observed after heating at 70 °C  
261 for 30 minutes (Yen and Wei, 1993), while Matusheski *et al.* (2004) discovered high  
262 sulforaphane content in broccoli after treating at 60 °C for 10 minutes, implying that the  
263 myrosinase in broccoli was intact at 60 °C. However, in both studies, intact cabbage and  
264 broccoli were used, therefore, actual temperatures the myrosinase was exposed to may  
265 have been lower. In other studies, only rapeseed has been shown to have a higher  
266 inactivation temperature compared to yellow mustard, where inactivation typically occurs  
267 above 60 °C at a holding time of 10 minutes (Kozłowska, Nowak, & Nowak, 1983; Stoin,  
268 Pirsan, Radu, Poiana, Alexa, & Dogaru, 2009; Van Eylen, Indrawati, Hendrickx, & Van Loey,  
269 2006; Van Eylen, Oey, Hendrickx, & Van Loey, 2008). Overall, myrosinase enzyme is

270 temperature sensitive and thermal treatments like blanching or heat processing are known  
271 to cause a decrease in myrosinase activity (Ghawi, Methven, Rastall, & Niranjana, 2012;  
272 Ludikhuyze, Ooms, Weemaes, & Hendrickx, 1999; Van Eyle, Indrawati, Hendrickx, & Van  
273 Loey, 2006; Van Eyle, Oey, Hendrickx, & Van Loey, 2008). Inactivation of myrosinase leads  
274 to decrease in formation of beneficial hydrolysis products from glucosinolate-myrosinase  
275 hydrolysis, although myrosinase inactivation in *Brassicac*s can be used to control sensory  
276 characteristics in *Brassicac* vegetables.

### 277 **3.3 Effect of pressure treatment on black, brown and yellow mustard seed** 278 **myrosinase**

279 Figure 2 depicts pressure inactivation of myrosinase from yellow, black and brown  
280 mustard seeds. Pressure stability was determined at 10 minutes exposure time at pressures  
281 ranging from 100-900 MPa at 15 °C. Myrosinase was substantially stable at 600 MPa with  
282 about 19% loss in activity for brown mustard after 5 minutes exposure time (data not  
283 included) while loss in activity for black mustard was 31%. Yellow mustard myrosinase was  
284 observed to be notably stable up to 500 MPa (14% loss in activity) and there was 79% loss in  
285 myrosinase activity at 600 MPa. At 700 MPa, there was 50 and 60% loss in myrosinase  
286 activity for black and brown mustard respectively. However, at 700 MPa, myrosinase from  
287 yellow mustard was completely inactivated, whilst there was no myrosinase activity at 900  
288 MPa for black and brown mustard.

289 Pressure treatment for 10 minutes showed similar trend to those observed for 5 minutes.  
290 Myrosinase activity decreased significantly above 600 MPa for both black and brown  
291 mustard. At 800 MPa, the loss in enzyme activity was over 70% for both black and brown  
292 mustard and there was no enzyme activity at 900 MPa. Yellow mustard myrosinase was

293 however only considerably stable up to 500 MPa (21% loss in activity) and at 600 MPa,  
294 about 79% enzyme activity was lost. Van eylen *et al.* (2006) reported that myrosinase from  
295 yellow mustard was inactivated at pressures above 600 MPa, however, this is not in  
296 agreement with the current study where 79% loss in activity was observed at 600 MPa  
297 processing. Compared to some other *Brassic*as, myrosinase from mustard have much higher  
298 pressure stability. Pressure stabilities of myrosinase from other *Brassica* types have only  
299 been reported for broccoli (300-500 MPa), green cabbage (250-300 MPa) and yellow  
300 mustard (above 600 MPa) (Ghawi, Methven, Rastall, & Niranjan, 2012; Ludikhuyze, Ooms,  
301 Weemaes, & Hendrickx, 1999; Van Eylen, Oey, Hendrickx, & Van Loey, 2008).

### 302 **3.4 Effect of combined temperature and pressure treatment on black, brown and** 303 **yellow mustard seed myrosinase**

304 Combined high pressure and temperature stability of myrosinase from brown, black and  
305 yellow mustard seed (Table 2) was studied at a temperature of 30-70 °C and pressure of 600  
306 -800 MPa for an exposure time of 10 minutes. At low temperatures (30-40 °C) and 600 MPa,  
307 it was observed that loss in activity was about 30% for black mustard, 20% for brown and  
308 50% for yellow mustard, respectively. Yellow mustard myrosinase showed no activity at 700  
309 and 800 MPa. An increase in temperature up to 70 °C led to approximately 60% loss in  
310 myrosinase activity at 600 MPa for black mustard and 70% for brown mustard. Overall,  
311 there was a gradual loss in myrosinase activity as the temperature and pressure gradually  
312 increased. This trend is in agreement with previous studies (Van Eylen, Oey, Hendrickx, &  
313 Van Loey, 2008), where applying high pressure (over 600 MPa) increased thermal  
314 inactivation rate.



315 At a combined pressure of 800 MPa and 70 °C, there was no myrosinase activity in any of  
316 the mustard samples studied. This indicates a synergistic effect of high pressure (600-800  
317 Mpa) on thermal inactivation of myrosinase in mustard seeds. However, Ghawi *et al.* (2012)  
318 reported a synergistic effect at lower pressure level in the case of green cabbage  
319 myrosinase. Pressure stability of myrosinase from *Brassicac*s is not widely reported. It is clear  
320 that myrosinase is inactivated at combined high pressure and temperature, so applying  
321 lower pressure and temperature could be more beneficial in retaining myrosinase activity  
322 and enabling formation of hydrolysis products.

323 At low pressure (200-300 MPa), it was observed that myrosinase enzyme activity was  
324 notably stable at 60 °C for black mustard while significant decrease in activity was observed  
325 for brown and yellow mustard (30% and 50%). In earlier studies (Van Eylen, Indrawati,  
326 Hendrickx, & Van Loey, 2006) an antagonistic effect of low pressure (200-300 MPa) on  
327 thermal inactivation of myrosinase in broccoli juice was reported, while Ghawi *et al.* (2012)  
328 reported a synergistic effect of pressure on thermal inactivation of myrosinase in green  
329 cabbage. In this study, an antagonistic effect of low pressure on thermal inactivation of  
330 mustard seed myrosinase was observed. The loss in myrosinase activity was lower using  
331 combined low pressure and temperature than the application of only thermal treatment.  
332 For black and brown mustard myrosinase, activity retention at 75 °C and 200-300 MPa for 10  
333 minutes processing time was above 70% and 55% respectively . Whereas without pressure,  
334 activity retention at 70 °C was approximately 59% for black and 35% for brown mustard  
335 myrosinase. Thermal processing of black and brown mustard myrosinase at 80 °C led to full  
336 inactivation, however, application of low pressure (200-300 MPa) at 80 °C retained  
337 considerable levels of the activity, about 50% for black and 40% for brown mustard. At 400

338 MPa, 80 °C and 10 minutes processing time, myrosinase activity was observed for black  
339 (30%) and brown mustard (20%). Similarly, combining low pressure (200-300 MPa) with  
340 thermal treatment at 70 °C retained 20% activity of yellow mustard myrosinase, whereas,  
341 there was no myrosinase activity under thermal processing for yellow mustard myrosinase  
342 at the same temperature. However, at higher temperature levels, there was no protective  
343 effect of pressure on thermal inactivation for yellow mustard myrosinase.

344 The differences in initial activity between the mustard species, where brown had the highest  
345 activity and yellow the least, led to similar trends in enzyme stability with temperature and  
346 pressure, where myrosinase from yellow mustard was the least stable. As discussed earlier,  
347 the differences in stability between the mustard species might have resulted from genetic  
348 differences or responses to different environmental challenges.

349

350

351

352

353

354

355

356

357

## 358 **4 Conclusion**

359 Myrosinase from different mustard species varied in terms of specific enzyme activity as  
360 well as temperature and pressure stability. Brown mustard myrosinase had the highest  
361 overall myrosinase activity and specific activity. Brown and black mustard myrosinase were  
362 more resistant to pressure and thermal treatment than myrosinase from yellow mustard.  
363 Combined high pressure-thermal treatment (up to 70 °C and 800 MPa) completely  
364 inactivated myrosinase from the mustards studied. However, at low pressure (200-400  
365 MPa), inactivation temperature increased in the mustard samples studied with lower rate of  
366 loss in myrosinase activity compared to any of thermal, pressure and combined high  
367 pressure-thermal treatment. This difference in myrosinase stability could be utilized to  
368 control the hydrolysis level of glucosinolates when mustard seeds are used as a condiment  
369 along with cooked *Brassica* vegetables. This could have important health implications  
370 through increasing the delivery of bioactive isothiocyanates from the *Brassica*. In addition,  
371 controlling enzyme activity can also be used to regulate sensory attributes of *Brassica*  
372 vegetables.

## 373 **Acknowledgments**

374 The authors thank The Federal Polytechnic, Ado Ekiti, Ekiti State, Nigeria and The Tertiary  
375 Education Trust Fund (**TETFUND**), Nigeria, for supporting the first named author, Dr Carol  
376 Wagstaff, Department of Food and Nutritional Sciences, University of Reading,  
377 Whiteknights, Reading, UK for her technical advice and IPK Gene Bank, Gatersleben,  
378 Germany for providing the mustard seeds used in this study.

379

380 **References**

- 381 Abul-Fadl, M. M., El-Badry, N., & Ammar, M. S. (2011). Nutritional and chemical evaluation for two  
382 different varieties of mustard seeds. *World Applied Sciences Journal*, 15(9), 1225-1233.
- 383 Bones, A. M., & Rossiter, J. T. (1996). The myrosinase-glucosinolate system, its organisation and  
384 biochemistry. *Physiologia Plantarum*, 97(1), 194-208.
- 385 Bongoni, R., Verkerk, R., Steenbekkers, B., Dekker, M., & Stieger, M. (2014). Evaluation of Different  
386 Cooking Conditions on Broccoli (*Brassica oleracea* var. *italica*) to Improve the Nutritional Value  
387 and Consumer Acceptance. *Plant Foods for Human Nutrition*, 69(3), 228-234.
- 388 Bradford, M. M. (1976). Rapid and sensitive method for quantification of micrograms quantities of  
389 protein utilizing principle of protein-dye binding. *Analytical Biochemistry*, 72(1-2), 248-254.
- 390 Conaway, C. C., Getahun, S. M., Liebes, L. L., Pusateri, D. J., Topham, D. K. W., Botero-Omary, M., &  
391 Chung, F. L. (2001). Disposition of glucosinolates and sulforaphane in humans after ingestion of  
392 steamed and fresh broccoli (vol 38, pg 177, 2001). *Nutrition and Cancer-an International Journal*,  
393 41(1-2), 196-196.
- 394 Dosz, E. B., & Jeffery, E. H. (2013). Modifying the processing and handling of frozen broccoli for  
395 increased sulforaphane formation. *Journal of Food Science*, 78(9), H1459-H1463.
- 396 Drewnowski, A., & Gomez-Carneros, C. (2000). Bitter taste, phytonutrients, and the consumer: a  
397 review. *American Journal of Clinical Nutrition*, 72(6), 1424-1435.
- 398 Fahey, J. W., Zalcmann, A. T., & Talalay, P. (2001). The chemical diversity and distribution of  
399 glucosinolates and isothiocyanates among plants. *Phytochemistry*, 56(1), 5-51.
- 400 Fahey, J. W., Zhang, Y. S., & Talalay, P. (1997). Broccoli sprouts: An exceptionally rich source of  
401 inducers of enzymes that protect against chemical carcinogens. *Proceedings of the National  
402 Academy of Sciences of the United States of America*, 94(19), 10367-10372.
- 403 Gatfield, I. L., & Sand, T. (1983). A coupled enzymatic procedure for the determination of myrosinase  
404 activity. *Lebensmittel-Wissenschaft & Technologie*, 16(2), 73 - 75.
- 405 Ghawi, S. K., Methven, L., & Niranjana, K. (2013). The potential to intensify sulforaphane formation in  
406 cooked broccoli (*Brassica oleracea* var. *italica*) using mustard seeds (*Sinapsis alba*). *Food  
407 Chemistry*, 138, 1734 - 1741.
- 408 Ghawi, S. K., Methven, L., Rastall, R. A., & Niranjana, K. (2012). Thermal and high hydrostatic pressure  
409 inactivation of myrosinase from green cabbage: A kinetic study. *Food Chemistry*, 131(4), 1240-  
410 1247.
- 411 Gow-Chin, Y., & Que-King, W. (1993). Myrosinase activity and total glucosinolate content of  
412 cruciferous vegetables, and some properties of cabbage myrosinase in Taiwan. *Journal of the  
413 Science of Food and Agriculture*, 61(4), 471-475.
- 414 Hendrickx, M., Ludikhuyze, L., Van den Broeck, L., & Weemaes, C. (1998). Effects of high pressure on  
415 enzymes related to food quality. *Trends in Food Science and Technology*, 9(5), 197 - 203.
- 416 Johnson, S., Koh, W.-P., Wang, R.-W., Yu, M. C., & Yuan, J.-M. (2010). Dietary isothiocyanate intake in  
417 relation to reduced risk of hepatocellular carcinoma: Findings from the Singapore Chinese Health

- 418 Study. *Proceedings of the American Association for Cancer Research Annual Meeting*, 51, 687-  
419 687.
- 420 Kozłowska, H. J., Nowak, H., & Nowak, J. (1983). Characterization of myrosinase in Polish varieties of  
421 rapeseed. *Journal of the Science of Food and Agriculture*, 34(11), 1171-1178.
- 422 Lambrix, V., Reichelt, M., Mitchell-Olds, T., Kliebenstein, D. J., & Gershenzon, J. (2001). The  
423 Arabidopsis epithiospecifier protein promotes the hydrolysis of glucosinolates to nitriles and  
424 influences *Trichoplusia ni* herbivory. *Plant Cell*, 13(12), 2793-2807.
- 425 Ludikhuyze, L., Ooms, V., Weemaes, C., & Hendrickx, M. (1999). Kinetic study of the irreversible  
426 thermal and pressure inactivation of myrosinase from broccoli (*Brassica oleracea* L-Cv. Italica).  
427 *Journal of Agricultural and Food Chemistry*, 47(5), 1794-1800.
- 428 Matusheski, N. V., Juvik, J. A., & Jeffery, E. H. (2003). Sulforaphane content and bioactivity of broccoli  
429 sprouts are enhanced by heat processing: A role for epithiospecifier protein. *Faseb Journal*,  
430 17(4), A377-A377.
- 431 Oerlemans, K., Barrett, D. M., Suades, C. B., Verkerk, R., & Dekker, M. (2006). Thermal degradation  
432 of glucosinolates in red cabbage. *Food Chemistry*, 95(1), 19-29.
- 433 Pérez, C., Barrientos, H., Román, J., & Mahn, A. (2014). Optimization of a blanching step to maximize  
434 sulforaphane synthesis in broccoli florets. *Food Chemistry*, 145(0), 264-271.
- 435 Piekarska, A., Kuszniereicz, B., Meller, M., Dziedziul, K., Namiesnik, J., & Bartoszek, A. (2013).  
436 Myrosinase activity in different plant samples; optimisation of measurement conditions for  
437 spectrophotometric and pH-stat methods. *Industrial Crops and Products*, 50, 58-67.
- 438 Pocock, K., Heaney, R. K., Wilkinson, A. P., Beaumont, J. E., Vaughan, J. G., & Fenwick, G. R. (1987).  
439 Changes in myrosinase activity and isoenzyme pattern, glucosinolate content and the cytology of  
440 myrosin cells in the leaves of heads of 3 cultivars of English white cabbage. *Journal of the Science  
441 of Food and Agriculture*, 41, 245 - 257.
- 442 Rask, L., Andreasson, E., Ekbohm, B., Eriksson, S., Pontoppidan, B., & Meijer, J. (2000). Myrosinase:  
443 gene family evolution and herbivore defense in Brassicaceae. *Plant Molecular Biology*, 42(1), 93-  
444 113.
- 445 Stoin, D., Pirsan, P., Radu, F., Poiana, M.-A., Alexa, E., & Dogaru, D. (2009). Studies regarding the  
446 myrosinase enzymatic activity from black mustard (*Brassica nigra*) seeds. *Journal of Food  
447 Agriculture & Environment*, 7(1), 44-47.
- 448 Tang, L., & Zhang, Y. S. (2004). Isothiocyanates in the chemoprevention of bladder cancer. *Current  
449 Drug Metabolism*, 5(2), 193-201.
- 450 Thangstad, O. P., & Bones, A. (1991). Myrosin cells in developing seedlings. Distribution and  
451 occurrence. *Physiologia Plantarum*, 82(3), B6-B6.
- 452 Van Eylen, D., Indrawati, Hendrickx, M., & Van Loey, A. (2006). Temperature and pressure stability of  
453 mustard seed (*Sinapis alba* L.) myrosinase. *Food Chemistry*, 97(2), 263-271.
- 454 Van Eylen, D., Oey, I., Hendrickx, M., & Van Loey, A. (2007). Kinetics of the stability of broccoli  
455 (*Brassica oleracea* cv. Italica) myrosinase and isothiocyanates in broccoli juice during  
456 pressure/temperature treatments. *Journal of Agricultural and Food Chemistry*, 55(6), 2163-2170.

- 457 Van Eylen, D., Oey, I., Hendrickx, M., & Van Loey, A. (2008). Behavior of mustard seed (*Sinapis alba*  
458 L.) myrosinase during temperature/pressure treatments: a case study on enzyme activity and  
459 stability. *European Food Research and Technology*, 226(3), 545-553.
- 460 Verkerk, R., & Dekker, M. (2004). Glucosinolates and myrosinase activity in red cabbage (*Brassica*  
461 *oleracea* L. var. *Capitata* f. *rubra* DC.) after various microwave treatments. *Journal of Agricultural*  
462 *and Food Chemistry*, 52(24), 7318-7323.
- 463 Wanasundara, J. (2008). Mustard as an Ingredient in Food Processing: Current Uses and the  
464 Potential. In *Mustard Grower, Mar 2008 Issue* Mar. 2008 ed., (pp. 3-5). Canada: Saskatchewan  
465 Mustard Development Commission.
- 466 Wilkinson, A. P., Rhodes, M. J. C., & Fenwick, G. R. (1984). Myrosinase activity of cruciferous  
467 vegetables. *Journal of the Science of Food and Agriculture*, 35, 543 - 552.
- 468 Yen, G. C., & Wei, Q. K. (1993). Myrosinase activity and total glucosinolate content of cruciferous  
469 vegetables, and some properties of cabbage myrosinase in Taiwan. *Journal of the Science of Food*  
470 *and Agriculture*, 61(4), 471-475.
- 471
- 472

473 **Figure and table captions**

474 Figure 1: Effect of thermal processing on relative myrosinase activity in black, brown and  
475 yellow mustard seeds; where temperature exposure time was 10 minutes. (■) brown  
476 mustard; (◆) black mustard; (▲) yellow mustard. (A – enzyme activity after thermal  
477 treatment, A<sub>o</sub> – Initial enzyme activity). Error bars represent standard errors of the means.

478  
479 Figure 2: Effect of pressure on relative myrosinase activity in black, brown and yellow  
480 mustard seeds. Pressure holding time was 10 minutes and processing temperature was  
481 controlled at 15 °C. (■) brown mustard; (◆) black mustard; (▲) yellow mustard. (A – enzyme  
482 activity after pressure treatment, A<sub>o</sub> – Initial enzyme activity). Error bars represent standard  
483 errors of the means.

484  
485 Table 1: Myrosinase activity, protein content and specific activity of yellow, brown and black  
486 mustard seeds. (\*un is activity units defined in section 2.2, lines 150-152).

487  
488 Table 2: Combined temperature and high pressure inactivation of myrosinase from black,  
489 brown and yellow mustard seeds at 10 minutes holding time.

490  
491 Table 3: Effects of combined low pressure and temperature processing on myrosinase  
492 activity in black, brown and yellow mustard seeds at 5, 10 and 15 minutes holding time.

493

494

495

496

497

498

499

500

501

502


503

504

505

506

507  
508  
509  
510



511

512 Error bars represent standard errors of the means

513 Figure 1

514

515

516

517

518

519

520




521

522

523

524



525

526 Error bars represent standard errors of the means

527 Figure 2

528

529

530

531

532

533

534

535

536

537

538

539

Table 1.

| Mustard Seed | Myrosinase Activity (un/mL) | Protein Content (mg/mL) | Specific Activity (un/mg) |
|--------------|-----------------------------|-------------------------|---------------------------|
| Black        | 1.5±0.00                    | 1.21±0.01               | 1.24                      |
| Brown        | 2.75±0.22                   | 1.34±0.05               | 2.04                      |
| Yellow       | 0.63±0.00                   | 1.32±0.01               | 0.48                      |

Table 2

| Pressure<br>(MPa) | Temperature<br>(°C) | Relative Enzyme Activity (A/A <sub>0</sub> ) |                        |                        |
|-------------------|---------------------|--|------------------------|------------------------|
|                   |                     | Black  | Brown                  | Yellow                 |
| 600               | 30                  | 0.7 ±0.00 <sup>a</sup>                       | 0.8 ±0.00 <sup>a</sup> | 0.5 ±0.00 <sup>a</sup> |
|                   | 40                  | 0.7 ±0.00 <sup>a</sup>                       | 0.8 ±0.00 <sup>a</sup> | 0.5 ±0.00 <sup>a</sup> |
|                   | 50                  | 0.7 ±0.04 <sup>a</sup>                       | 0.5 ±0.00 <sup>b</sup> | 0.4 ±0.00 <sup>b</sup> |
|                   | 60                  | 0.7 ±0.00 <sup>a</sup>                       | 0.4 ±0.00 <sup>c</sup> | 0.2 ±0.00 <sup>c</sup> |
|                   | 70                  | 0.4 ±0.00 <sup>b</sup>                       | 0.3 ±0.02 <sup>d</sup> | -                      |
| 700               | 30                  | 0.4 ±0.00 <sup>a</sup>                       | 0.4 ±0.00 <sup>a</sup> | -                      |
|                   | 40                  | 0.4 ±0.00 <sup>a</sup>                       | 0.4 ±0.00 <sup>a</sup> | -                      |
|                   | 50                  | 0.3 ±0.04 <sup>b</sup>                       | 0.3 ±0.02 <sup>b</sup> | -                      |
|                   | 60                  | 0.2 ±0.04 <sup>c</sup>                       | 0.1 ±0.00 <sup>c</sup> | -                      |
|                   | 70                  | 0.2 ±0.00 <sup>c</sup>                       | 0.1 ±0.02 <sup>c</sup> | -                      |
| 800               | 30                  | 0.3 ±0.00 <sup>a</sup>                       | 0.2 ±0.00 <sup>a</sup> | -                      |
|                   | 40                  | 0.3 ±0.00 <sup>a</sup>                       | 0.2 ±0.00 <sup>a</sup> | -                      |
|                   | 50                  | 0.2 ±0.00 <sup>b</sup>                       | 0.1 ±0.00 <sup>b</sup> | -                      |
|                   | 60                  | 0.1 ±0.00 <sup>c</sup>                       | 0.1 ±0.00 <sup>b</sup> | -                      |
|                   | 70                  | -  | -                      | -                      |

Values not sharing a common letter are significantly different at P<0.05.

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554 Table 3.

| P<br>(MPa) | T(°C) | Relative Enzyme Activity (A/A <sub>0</sub> ) |                       |                       |                           |                       |                       |                           |                       |                       |
|------------|-------|--|-----------------------|-----------------------|---------------------------|-----------------------|-----------------------|---------------------------|-----------------------|-----------------------|
|            |       | Black  |                       |                       | Brown                     |                       |                       | Yellow                    |                       |                       |
|            |       | Processing time (Minutes)                    |                       |                       | Processing time (Minutes) |                       |                       | Processing time (Minutes) |                       |                       |
|            |       | 5  | 10                    | 15                    | 5                         | 10                    | 15                    | 5                         | 10                    | 15                    |
| 200        | 60    | 0.9±0.07 <sup>a</sup>                        | 0.9±0.00 <sup>a</sup> | 0.9±0.07 <sup>a</sup> | 0.7±0.07 <sup>a</sup>     | 0.7±0.07 <sup>a</sup> | 0.7±0.00 <sup>a</sup> | 0.6±0.00 <sup>a</sup>     | 0.5±0.11 <sup>a</sup> | 0.5±0.11 <sup>a</sup> |
|            | 65    | 0.9±0.05 <sup>a</sup>                        | 0.9±0.05 <sup>a</sup> | 0.9±0.05 <sup>a</sup> | 0.7±0.03 <sup>a</sup>     | 0.7±0.03 <sup>a</sup> | 0.7±0.00 <sup>a</sup> | 0.4±0.00 <sup>a</sup>     | 0.4±0.00 <sup>a</sup> | 0.4±0.00 <sup>a</sup> |
|            | 70    | 0.8±0.00 <sup>b</sup>                        | 0.8±0.00 <sup>b</sup> | 0.8±0.00 <sup>b</sup> | 0.7±0.00 <sup>a</sup>     | 0.7±0.00 <sup>a</sup> | 0.6±0.05 <sup>b</sup> | 0.2±0.00 <sup>b</sup>     | 0.2±0.00 <sup>b</sup> | 0.2±0.00 <sup>b</sup> |
|            | 75    | 0.8±0.00 <sup>b</sup>                        | 0.8±0.00 <sup>b</sup> | 0.7±0.05 <sup>c</sup> | 0.6±0.00 <sup>b</sup>     | 0.6±0.03 <sup>b</sup> | 0.6±0.03 <sup>b</sup> | -                         | -                     | -                     |
|            | 80    | 0.6±0.00 <sup>c</sup>                        | 0.6±0.05 <sup>c</sup> | 0.5±0.08 <sup>d</sup> | 0.4±0.03 <sup>c</sup>     | 0.4±0.00 <sup>c</sup> | 0.4±0.03 <sup>c</sup> | -                         | -                     | -                     |
| 300        | 60    | 0.9±0.07 <sup>a</sup>                        | 0.9±0.07 <sup>a</sup> | 0.9±0.07 <sup>a</sup> | 0.7±0.07 <sup>a</sup>     | 0.7±0.00 <sup>a</sup> | 0.7±0.07 <sup>a</sup> | 0.5±0.11 <sup>a</sup>     | 0.5±0.11 <sup>a</sup> | 0.5±0.11 <sup>a</sup> |
|            | 65    | 0.9±0.05 <sup>a</sup>                        | 0.9±0.05 <sup>a</sup> | 0.9±0.05 <sup>a</sup> | 0.7±0.03 <sup>a</sup>     | 0.7±0.03 <sup>a</sup> | 0.7±0.00 <sup>a</sup> | 0.3±0.11 <sup>a</sup>     | 0.3±0.11 <sup>a</sup> | 0.3±0.11 <sup>a</sup> |
|            | 70    | 0.8±0.05 <sup>b</sup>                        | 0.8±0.05 <sup>b</sup> | 0.8±0.05 <sup>b</sup> | 0.7±0.00 <sup>a</sup>     | 0.6±0.00 <sup>b</sup> | 0.6±0.03 <sup>b</sup> | 0.2±0.00 <sup>b</sup>     | 0.2±0.00 <sup>b</sup> | 0.2±0.00 <sup>b</sup> |
|            | 75    | 0.7±0.05 <sup>c</sup>                        | 0.7±0.05 <sup>c</sup> | 0.7±0.05 <sup>c</sup> | 0.6±0.03 <sup>b</sup>     | 0.6±0.00 <sup>b</sup> | 0.6±0.03 <sup>b</sup> | -                         | -                     | -                     |
|            | 80    | 0.5±0.00 <sup>d</sup>                        | 0.5±0.00 <sup>d</sup> | 0.5±0.00 <sup>d</sup> | 0.4±0.00 <sup>c</sup>     | 0.4±0.00 <sup>c</sup> | 0.4±0.07 <sup>c</sup> | -                         | -                     | -                     |
| 400        | 60    | 0.8±0.00 <sup>a</sup>                        | 0.7±0.00 <sup>a</sup> | 0.7±0.07 <sup>a</sup> | 0.6±0.07 <sup>a</sup>     | 0.6±0.00 <sup>a</sup> | 0.6±0.07 <sup>a</sup> | 0.4±0.00 <sup>a</sup>     | 0.4±0.00 <sup>a</sup> | 0.3±0.11 <sup>a</sup> |
|            | 65    | 0.7±0.00 <sup>b</sup>                        | 0.7±0.00 <sup>a</sup> | 0.7±0.00 <sup>a</sup> | 0.6±0.00 <sup>a</sup>     | 0.6±0.00 <sup>a</sup> | 0.6±0.03 <sup>a</sup> | 0.3±0.11 <sup>a</sup>     | 0.3±0.11 <sup>a</sup> | 0.2±0.00 <sup>a</sup> |
|            | 70    | 0.7±0.00 <sup>b</sup>                        | 0.7±0.00 <sup>a</sup> | 0.7±0.00 <sup>a</sup> | 0.6±0.03 <sup>a</sup>     | 0.6±0.03 <sup>a</sup> | 0.6±0.07 <sup>a</sup> | 0.1±0.11 <sup>b</sup>     | -                     | -                     |
|            | 75    | 0.5±0.05 <sup>c</sup>                        | 0.5±0.05 <sup>b</sup> | 0.5±0.05 <sup>b</sup> | 0.4±0.03 <sup>b</sup>     | 0.4±0.03 <sup>b</sup> | 0.4±0.05 <sup>b</sup> | -                         | -                     | -                     |
|            | 80    | 0.3±0.05 <sup>d</sup>                        | 0.3±0.00 <sup>c</sup> | 0.3±0.00 <sup>c</sup> | 0.2±0.00 <sup>c</sup>     | 0.2±0.03 <sup>c</sup> | 0.2±0.05 <sup>c</sup> | -                         | -                     | -                     |

555 Values not sharing a common letter are significantly different at P&lt;0.05.

556