

Fringes of the empire: diet and cultural change at the Roman to post-Roman transition in NW Iberia

Article

Accepted Version

López-Costas, O. and Muldner, G. ORCID:
<https://orcid.org/0000-0002-4513-9263> (2016) Fringes of the empire: diet and cultural change at the Roman to post-Roman transition in NW Iberia. *American Journal of Physical Anthropology*. ISSN 1096-8644 doi:
<https://doi.org/10.1002/ajpa.23016> Available at
<https://centaur.reading.ac.uk/66226/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/ajpa.23016>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



**Fringes of the Empire: Diet and cultural change at the
Roman to post-Roman Transition in NW Iberia**

Journal:	<i>American Journal of Physical Anthropology</i>
Manuscript ID	AJPA-2015-00393.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	López-Costas, Olalla; Universidade de Santiago de Compostela, Group Earth System Sciences, Departamento de Xeografía-Facultade de Xeografía e Historia; Stockholm University, Archaeological Research Laboratory. Department of Archaeology and Classical Studies, Wallenberglaboratoriet; Universidad de Granada, Laboratory of Anthropology, Department of Legal Medicine, Toxicology and Physical Anthropology, Faculty of Medicine Müldner, Gundula; University of Reading, Department of Archaeology
Key Words:	millet, saltmarsh influence, Migration period, stable isotopes, paleodiet
Subfield: Please select your first choice in the first field.:	Bioarchaeology [including forensics], Human biology [living humans; behavior, ecology, physiology, anatomy]

SCHOLARONE™
Manuscripts

1
2
3 1 **Fringes of the Empire: Diet and cultural change at the Roman to post-Roman**4
5 2 **Transition in NW Iberia**6
7
8 3 Olalla López-Costas^{1,2,3*} and Gundula Müldner⁴9
10
11 4 ¹Group Earth System Sciences, Departamento de Xeografía, Facultade de Xeografía e Historia,
12 5 Universidade de Santiago de Compostela, Santiago de Compostela, 15782, Spain.13
14 6 ²Archaeological Research Laboratory, Stockholm University, Wallenberglaboratoriet, SE-10691
15 7 Stockholm, Sweden.16
17 8 ³Laboratory of Anthropology, Department of Legal Medicine, Toxicology and Physical Anthropology,
18 9 Faculty of Medicine, University of Granada, Granada 18012, Spain19
20 10 ⁴Department of Archaeology, University of Reading, Whiteknights, PO Box 227, Reading, RG6 6AB,
21 11 UK

22 12

23
24 13 Number of text pages: 26

25 14 Number of figures: 7

26 15 Number of tables: 7

27
28
29 16 **Abbreviated title:** Isotopic diet study in Roman and post-Roman A Lanzada30
31 17 **Key words** (3-5): millet; saltmarsh influence; Migration period; stable isotopes;
32 18 paleodiet33
34 19 *** corresponding author:** Olalla López-Costas35 20 Adress: Current adress: Stockholm University Arkeologiska forsknings laboratoriet
36 21 (AFL) SE-10691 Stockholm, Sweden.

37 22 Telephone number: +468164590

38 23 E.mail addresses: olallalc@gmail.com , olalla.lopez@usc.es ,

39 24 olalla.lopez.costas@arklab.su.se

40
41
42 2543
44 26 **Grant sponsorship:** OLC is funded by Plan Galego I2C mod.A (Xunta de Galicia). The
45 27 research was funded by projects“Antropoloxía dos restos óseos humanos de Galicia”
46 28 (Dirección Xeral de Patrimonio Histórico), GPC2014/009 and R2014/001 of Xunta de
47 29 Galicia.48
49
50 3051
52
53
54 31

ABSTRACT (250)**Objectives**

A growing number of paleodiet investigations over recent years have begun to reveal the stark dietary differences that existed between regions of the Roman Empire, as well as significant changes in subsistence strategies after its fall. The present study explores the dietary changes at the Roman to post-Roman (Germanic) transition in the Northwest Iberian Peninsula, in order to improve our understanding of the changes that occurred at end of the Roman Empire in different regions across Europe and to also consider the influence of climate had on them.

Materials and Methods

We present the carbon and nitrogen stable isotope investigation in bone collagen from A Lanzada, NW Spain (100-700 AD), which was an important commercial, coastal settlement. A human sample of 59 individuals, 6 of them subadults, is compared with 31 faunal specimens, which include a number of marine fish.

Results

Isotope data for the terrestrial fauna reveal the influence of the sea on the local isotope baseline. Analysis of the human samples indicates a mixed marine-terrestrial diet. A shift in mean human $\delta^{13}\text{C}$ values from -16.7 to -14.3‰ provides clear evidence for a significant change in diet in the post-Roman period, probably through the intensification of both marine resources exploitation and C_4 -plant consumption (presumably millet).

Discussion

A deterioration of paleoenvironmental conditions, together with a poor socioeconomic situation and the arrival of new people, the *Sueves*, who brought a new political and socioeconomic system have been discussed as the main causes for the dietary modification in post-Roman times.

1
2
3 56 **Introduction**
4
5

6 57 The study of diet and foodways and their change over time offers important
7
8 58 insights into human societies and individuals in terms of their social and economic
9
10 59 structure, health status and standards of living (Mintz and Du Bois 2002). Dietary
11
12 60 change may be triggered by historical events, such as foreign invasions, but more often
13
14 61 reflect general changes of the social, cultural, economic or environmental conditions
15
16 62 humans lived in. Methodologies, such as the analysis of stable isotopes performed on
17
18 63 human and faunal bone collagen, provide information on the major sources of protein
19
20 64 intake (Jim et al. 2004) during the life of an individual (e.g. the review articles by
21
22 65 Schwarcz and Schoeninger 1991; Sealy 2001) depending on the turnover of the
23
24 66 analysed tissue (Hedges et al. 2007). These are a particularly efficient way of tracing the
25
26 67 history of human diet and its changes over time. Recently, routine applications of this
27
28 68 type of analysis have provided abundant data about the diet and subsistence strategies of
29
30 69 ancient populations (e.g. in Spain Alexander et al. 2015; Arias and Schulting 2010;
31
32 70 Davis 2002; Fuller et al. 2010; García-Guixé et al. 2009; García et al. 2004; García
33
34 71 Guixé et al. 2006; López-Costas et al. 2015; Munde 2009; Salazar-García et al. 2014;
35
36 72 Van Strydonck et al. 2002; Van Strydonck et al. 2005). Consequently, it has now
37
38 73 become established as a key tool for understanding the relationship between humans
39
40 74 and their cultural and natural environment in the past.
41
42
43
44
45

46 75 The end of the Roman Empire and the Migration Period (5th to 8th centuries AD)
47
48 76 had a profound impact on the European landscape, laying the foundations for the
49
50 77 political geography of many modern European countries (Musset 1975). The Migration
51
52 78 period has been relatively unexplored in Spanish archaeology compared to other
53
54 79 periods; however, the interest in it has increased more recently. This is probably due to
55
56 80 the realization that the demise of the Roman Empire had far-reaching consequences for
57
58
59
60

1
2
3 81 the way of life and standards of living of the local populations, due to the breakdown of
4
5 82 the old political and economic structure, the Germanic invasions and the accompanying
6
7 83 climatic downturn which occurred at the end of the Roman Warm Period (among others
8
9 84 Büntgen et al. 2011; Martínez-Cortizas et al. 1999; Mighall et al. 2006). Previous
10
11 85 paleodietary studies of the Roman to post-Roman transition in Central and South-
12
13 86 eastern Europe have detected an increase in the consumption of C₄ plant products, most
14
15 87 probably millet, which may have been related to the arrival of ethnic groups with
16
17 88 different cuisines and traditions (Hakenbeck et al. 2010; Lightfoot et al. 2012).

18
19
20
21 89 The Iberian Peninsula is considered an essential region in the Imperial economy,
22
23 90 especially during Late Roman times (Kulikowski 2004). Despite being conquered
24
25 91 relatively late (end of the 1st century BC) (Syme 1934), the Roman province of
26
27 92 *Gallaecia*, to the Northwest, played an essential role in the Atlantic trade routes and in
28
29 93 mining, among others. The historical importance of Northwest Iberia extended beyond
30
31 94 the end of the Roman Empire to the Migration period, also known as the ‘Germanic
32
33 95 invasions’ in Spain. One of the main reasons is that the Iberian Peninsula, and the
34
35 96 Northwest in particular, was one of the first areas where the foreign populations
36
37 97 permanently settled (Kulikowski 2004; Musset 1975). The historical relevance of the
38
39 98 area contrasts with the poor knowledge of features related to everyday life, such as diet.
40
41 99 The historiography and archaeological evidence about these topics is scarce for Roman
42
43 100 times and dramatically decreases with the arrival of Germanic peoples (Díaz 2011).

44
45
46
47
48
49 101 In *Gallaecia* the overthrow of Roman rule by a Germanic group known as the
50
51 102 *Sueves* in the early 5th century AD, is regarded as the end of the Roman period and the
52
53 103 beginning of Early Middle Ages (see Castellanos and Viso 2005; Díaz 2011). The
54
55 104 *Sueves* permanently settled in the area of modern day Galicia and North Portugal,
56
57 105 creating a Kingdom that lasted almost two centuries (AD 409-585). Recent
58
59
60

1
2
3 106 palaeoenvironmental studies suggest that the landscape underwent a number of
4
5 107 significant human-made transformations during this time, through deforestation and
6
7 108 new modes of agriculture and animal husbandry, which still shape much of the
8
9 109 environment today (Ferro-Vázquez et al. 2014; Martínez Cortizas et al. 2005). These
10
11 110 changes could have implications for the subsistence economy during the transition from
12
13 111 Roman to post-Roman times, which may have left an imprint in human remains.
14
15

16
17 112 The present study explores the dietary changes at the Roman to post-Roman
18
19 113 transition in the Northwest Iberian Peninsula, in order to improve our understanding of
20
21 114 the end of the Roman Empire at the western fringe of Europe, paying particular
22
23 115 attention to the possible influence of climate. In order to address this aim, we carried out
24
25 116 a programme of carbon and nitrogen stable isotope analyses of bone collagen on
26
27 117 samples from the coastal cemetery site of A Lanzada (Galicia, NW Iberia) (100-700
28
29 118 AD) (Blanco Freijeiro et al. 1967), where two funerary areas have been described: one
30
31 119 from the Roman (2nd to 4th centuries AD) and the other from the post-Roman (5th to 7th
32
33 120 centuries AD) times (López-Costas 2015).
34
35
36

37 38 121 **THE COASTAL NECROPOLIS OF A LANZADA**

39

40 122 The archaeological site of A Lanzada is located in the province of Pontevedra
41
42 123 (Galicia, NW Spain) on a small headland just south of the O Grove peninsula, between
43
44 124 the estuaries of Arousa and Pontevedra, two of the so-called *Rias Baixas* (see Fig. 1).
45
46 125 Since the area is surrounded by the sea, a range of different coastal ecosystems occur in
47
48 126 the immediate vicinity and the position of the site once offered excellent control over
49
50 127 the maritime traffic along the coast (see Fig. 1). The soils are not particularly well suited
51
52 128 for agriculture due to their sandy texture, low water retention and high input of sea-
53
54 129 spray, and they are subject to constant winds and soil erosion; however, further inland,
55
56
57
58
59
60

1
2
3 130 the quality of the soils improves (López-Costas et al. 2016). Fish and seafood are
4
5 131 abundant in the area and good pasture for domestic animals is provided by the many
6
7 132 halophytic grasses and shrubs that can be found locally on saltmarshes (Valdés-Bermejo
8
9 133 and Silva Pando 1986).

10
11
12 134 Probably because of its strategic location, A Lanzada area has been occupied
13
14 135 since prehistoric times. Remains from a Bronze to Iron Age settlement, a Roman and
15
16 136 post-Roman cemetery with possible traces of a settlement, as well as a defensive tower
17
18 137 and a Romanesque church from the medieval period have been found (Fariña Busto and
19
20 138 Filgueira Valverde 1974) (see Fig. 2). Recent excavations have uncovered salting
21
22 139 facilities for seafood dating back to the Iron Age and historical sources suggest that a
23
24 140 nearby marshland was intensively exploited for salt production during the Late Roman
25
26 141 and Medieval periods (Rodríguez Martínez et al. 2011). A significant amount of
27
28 142 imported pottery and other foreign artefacts dating from the 5th century BC to the 6th
29
30 143 century AD, attests to A Lanzada lasting importance in the long-distance trade network
31
32 144 even after the end of the Roman phase (González Ruibal 2004; Naveiro López 1991).
33
34 145 Additionally, a large midden deposit, which consisted mainly of faunal remains
35
36 146 presumably dated from Iron to Roman period, was found in the area (Rodríguez
37
38 147 Martínez et al. 2011).

39
40
41
42
43
44 148 The main evidence of occupation from the Roman and post-Roman period is a
45
46 149 cemetery located at the Eastern edge of the headland (Fig. 2), which was partially
47
48 150 excavated during several campaigns between 1949 and 1963 (Blanco Freijeiro et al.
49
50 151 1967; Filgueira Valverde and Blanco Freijeiro 1962) and from 1975 to 1977 (Fariña
52
53 152 Busto 1975). Although many areas were intensively investigated, the available
54
55 153 documentation is of varied quality and only part of the results from the graves were
56
57 154 formally published (Blanco Freijeiro et al. 1961; Blanco Freijeiro et al. 1967; Carro

1
2
3 155 Otero et al. 1987; Filgueira Valverde and Blanco Freijeiro 1962). The preservation of
4
5 156 the human bones was good and the archaeologists also collected some faunal remains
6
7 157 from the graves.
8
9

10
11 158 The burials belong to two funerary areas with different dating based on
12
13 159 archaeological evidence and radiocarbon dates (Fig. 2) (López-Costas 2015),
14
15 160 (information about individual burials is summarised in Table 3). Of the 85 skeletons,
16
17 161 including several sub-adults, which have been recovered, 59 were selected for this
18
19 162 study, based on suitable skeletal preservation and avoiding non-adults under 12 years
20
21 163 old.
22
23

24 164 The first phase comprises the burials from the northern cemetery area (Fig.2),
25
26 165 which are dated to the Roman period (2nd to 4th centuries AD). The bodies were
27
28 166 predominantly aligned south-north and three main types of tombs have been recorded:
29
30 167 (1) simple earth graves or simple trench burials (see Fig. 3.A), (2) earth grave
31
32 168 surrounded by stones, (3) *Tegulae* grave (*cappuccina*) and/or earth grave with an *imbrex*
33
34 169 (curved tile) under the head (see Fig. 3.A). A detailed description of the burial rites has
35
36 170 been published elsewhere (López-Costas 2015). About 50% of the individuals were
37
38 171 accompanied by dress accessories (e.g. hob-nails), which suggests that they had been
39
40 172 buried clothed. Grave goods (pottery, glass) were also frequently present and at least
41
42 173 three skeletons were accompanied by coins, which dated between 213 and 325 AD.
43
44 174 While the majority of burials were inhumations, two cremation burials also belong to
45
46 175 this phase, but no human remains were collected from them. Following Toynbee's
47
48 176 (1971) classification, the burials could be interpreted as a lower-middle class Roman
49
50 177 cemetery.
51
52
53
54
55
56
57
58
59
60

1
2
3 178 The second phase of the cemetery was excavated during the 1970s. According to
4
5 179 the photographs it was located south of the previous excavations (Fig. 2) and it was
6
7 180 characterised by either simple earth graves or graves constructed from stone slabs (Fig.
8
9 181 3.B) (López-Costas 2015). Three of the stone slab coffins were used for multiple
10
11 182 consecutive burials (Fig. 3B). All skeletons were west-east oriented, laid in supine
12
13 183 position and there was a total absence of grave goods, although scallop shells found
14
15 184 near some of the bodies may have been deliberately placed. This second phase has been
16
17 185 dated to the post-Roman or Migration period, 5th to 7th centuries AD. Although stone
18
19 186 slab tombs have also been found in earlier Roman cemeteries in other parts of the
20
21 187 Empire (e.g. Philpott 1991), this mode of burial is known as typical for the Germanic
22
23 188 kingdoms of NW Iberia, where it has also been associated with Christian burials (Fariña
24
25 189 Busto and Suarez Otero 1997).

30 MATERIAL AND METHODS

31
32
33 191 Samples for isotope analysis were taken from a total of 59 humans and 31
34
35 192 animals. The faunal set comprised bone and tooth samples mainly of cattle (11),
36
37 193 sheep/goat (9) and pig (3) that were taken from the burials themselves, although they
38
39 194 may originally have belonged to the midden deposit from the Iron and Roman Age. We
40
41 195 also included seven marine animals, six bony fish and a dolphin, obtained from 15th
42
43 196 century contexts at the site of Ponte do Burgo at Pontevedra, ca. 20 km south-east of A
44
45 197 Lanzada (animal species information are summarized in Table 1), in order to trace the
46
47 198 isotopic background of the marine resources. The human sample was made up of 59
48
49 199 individuals: 38 from the first phase of the cemetery and 20 from the second, later phase,
50
51 200 as well as a single medieval (10th century AD) skeleton found in the western part of the
52
53 201 site. The male/female ratio of the human sample was almost 1:1 (29/24 and 6
54
55 202 subadults). The remains of sub-adults (with an age-at-death over 12 years in order to
56
57
58
59
60

1
2
3 203 avoid the breast-feeding effect; see for example Fuller et al. 2006) were included if they
4
5 204 were well preserved. Since the mode of excavation used from the 1950s-1970 implied
6
7 205 the incomplete retrieval of the skeletons and, specifically, the lack of smaller bones like
8
9 206 ribs, samples were taken from various skeletal elements (table 3). The human remains
10
11 207 were examined morphologically using standard methods for human osteological
12
13 208 analysis and reference database containing other Iberian populations. Detailed results of
14
15 209 osteological and paleopathological analysis are presented elsewhere (López-Costas
16
17 210 2012).

20
21 211 The isotopic analyses were carried out at the Department of Archaeology at the
22
23 212 University of Reading (U.K.). Bone collagen was extracted following the method
24
25 213 described by Longin (1971) with modifications recommended by Collins and Galley
26
27 214 (1998), according to the protocol described in Britton et al. (2008). No ultrafiltration
28
29 215 was applied (Pestle et al. 2014). The degree of preservation has been addressed by
30
31 216 Pyrolysis GC/MS, finding no relationship between the molecular and isotopic ($\delta^{13}\text{C}$ and
32
33 217 $\delta^{15}\text{N}$) composition of extracted bone collagen (Kaal et al. 2016).

34
35
36
37 218 Carbon and nitrogen stable isotope ratios were measured in duplicate on a
38
39 219 Europa 20-20 isotope ratio mass spectrometer coupled to a Sercon elemental analyzer.
40
41 220 Analytical error was calculated by repeated analyses of internal standards and was
42
43 221 $\pm 0.2\text{‰}$ or less for both elements (1 s.d.- standard deviation). Statistical analysis of the
44
45 222 data was made using the program SPSS 16. The comparisons among two groups were
46
47 223 computed by Student's t-test (t-test), or Mann-Whitney U test (M-W-test) whenever the
48
49 224 size of one of the series was equal or smaller than 15 individuals. The non-parametric
50
51 225 Kruskal-Wallis (K-W test) test was performed for comparisons of more than two
52
53 226 series/groups, since at least one series had less than 15 individuals (e.g. comparisons
54
55 227 among three groups of animals or different age groups in humans).

1
2
3 228 **RESULTS AND DISCUSSION**

4
5
6 229 Even though the low pH of Galician soils commonly causes poor skeletal
7
8 230 preservation (López-Costas 2012; López-Costas et al. 2016), the collagen obtained from
9
10 231 the A Lanzada samples was well preserved with an average yield of 8.3 ± 5.1 wt%
11
12 232 (1s.d.). The carbon (35.7 ± 6.8 wt% C) and nitrogen (12.6 ± 2.6 wt% N) contents
13
14 233 indicate that the majority of the samples had good preservation. Samples with lower
15
16 234 contents have not atypical values in other parameters and all human and animal C/N ratios
17
18 235 were between 3.1 and 3.6. The individual data and summary statistics are given in
19
20 236 Tables 1-7. Data are plotted in Figures 4-6.

21
22
23
24 237 **Faunal isotope data**

25
26
27 238 Since this is the first stable isotope study on material from the North-West
28
29 239 Iberian coast, the analysis of animal remains is important in order to map the $\delta^{13}\text{C}$ and
30
31 240 $\delta^{15}\text{N}$ baseline variations in the A Lanzada environment. The terrestrial animals have a
32
33 241 relatively wide range of carbon and nitrogen isotope values of -19.9 ± 1.2 ‰ (-21.5 , -
34
35 242 16.4 ‰) and 7.8 ± 1.8 ‰ (4.5 , 11.7 ‰), respectively (see Table 2 for basic summary
36
37 243 statistics). No systematic differences were found between herbivores and omnivores
38
39 244 (M-W-test $U=41.500$, $p=0.40$ for $\delta^{13}\text{C}$; $U=48.000$, $p=0.17$ for $\delta^{15}\text{N}$), or between the three
40
41 245 most common animal groups, cattle, sheep-goat and pigs ($\delta^{13}\text{C}$: Kruskal=Wallis test
42
43 246 $H_{(2)}=2.85$, $p=0.240$; $\delta^{15}\text{N}$: K-W test $H_{(2)}=2.95$, $p=0.229$). The different groups of
44
45 247 animals are discussed below.

46
47
48
49
50 248 **Herbivores.** The $\delta^{13}\text{C}$ of the terrestrial herbivores range over almost 3‰ (-
51
52 249 21.5 ‰ to -18.8 ‰; -20.2 ± 0.7 ‰, average \pm standard deviation), excluding the outlier
53
54 250 906a (see below). These values are similar to those observed at the Balearic sites (Fuller
55
56 251 et al. 2010; García et al. 2004) and indicate a diet based on C_3 -plants. High carbon

1
2
3 252 isotope values in some of the animals (with $\delta^{13}\text{C} > \sim -20\text{‰}$) are consistent with the
4
5 253 presence of small amounts of C_4 plants or seaweed, in the food chain, or alternatively
6
7 254 with grazing on saline pastures (see discussion below). Consumption of such resources
8
9 255 is particularly evident in the case of outlier 906a (Fig. 4). The sample belonged to a
10
11 256 young sheep or goat (less than 24 months) with elevated $\delta^{13}\text{C}$ (-17.7‰) and $\delta^{15}\text{N}$
12
13 257 (11.2‰). According to the age at death, part of the enrichment could be related to the
14
15 258 suckling effect (Balasse and Tresset 2002), but the input of a ^{13}C -enriched supplement
16
17 259 fodder is also likely. The fact that this signal is particularly evident in a young animal
18
19 260 could suggest that it was a seasonal rather than year-round supplement to herbivore diet
20
21 261 at A Lanzada. Dental serial sections would be needed to test this hypothesis (see for
22
23 262 example Balasse et al. 2006).

24
25
26
27
28 263 The herbivore $\delta^{15}\text{N}$ also shows a wide range of variation, over 5.5‰ , and a mean
29
30 264 of $7.4 \pm 1.6\text{‰}$. In this context, it is remarkable that about half the samples have $\delta^{15}\text{N}$
31
32 265 between 8‰ and 10‰ which are elevated values in comparison with most other sites
33
34 266 from West and South-West Europe (e.g., Craig et al. 2009; Fuller et al. 2010; García-
35
36 267 Guixé et al. 2009; García et al. 2004; Müldner and Richards 2007; Stevens et al. 2010)
37
38 268 (see Fig. 4).

39
40
41
42 269 While a number of factors can lead to ^{15}N -enriched values in plants and
43
44 270 herbivore tissues, such an effect has been specifically observed for saline soils including
45
46 271 animals grazing in salt-marshes (Britton et al. 2008; Cloern et al. 2002; Heaton 1987;
47
48 272 Van Groenigen and Van Kessel 2002; Virginia and Delwiche 1982). It is therefore
49
50 273 reasonable to assume that the high herbivore $\delta^{15}\text{N}$ can be attributed to the environmental
51
52 274 conditions at A Lanzada. It is possible that traditional farming practices, which are
53
54 275 documented for the area in recent historical times and involve the fertilization of fields
55
56 276 with seaweed, seagrass or small crustaceans, were already in use in the more distant past
57
58
59
60

1
2
3 277 (Ferreiro García et al. 1993; Pérez García 1979; Villares et al. 2007), affecting the
4
5 278 nitrogen isotope composition of the crops (see Fraser et al. 2011). While direct evidence
6
7 279 for this is lacking for the Roman period, soil characteristics (i.e. acidic pH) could
8
9 280 suggest that these methods of fertilization may already have been practiced in antiquity.
10
11 281 Nevertheless, it is also likely that humans exploited the nearby O'Vao salt-marshes as
12
13 282 pasture grounds for at least part of the year. Salt-marsh grazing could also explain the
14
15 283 ^{13}C -enrichment in the bone collagen of several of the animals, either through the effect
16
17 284 of salinity on the carbon isotope composition of C_3 -plants (e.g. van Groenigen and van
18
19 285 Kessel 2002) or through direct input of C_4 -plants to the herbivore diet: one of the
20
21 286 common halophytes, or salt-loving plants, in the marshland near A Lanzada today is
22
23 287 *Spartina maritima* (Valdés-Bermejo and Silva Pando 1986), which follows the C_4
24
25 288 photosynthetic pathway (Sage and Monson 1999:221). Nevertheless, since millet, also a
26
27 289 C_4 -plant which is known to have been used for animal foddering in classical times (see
28
29 290 Spurr 1986), was cultivated in Roman Galicia (Dopazo Martínez et al. 1996), it also
30
31 291 may be responsible for the elevated $\delta^{13}\text{C}$ in the herbivores, as is the use of sea-weed as
32
33 292 supplementary fodder, for example during the winter months (see Balasse et al. 2006).
34
35
36
37
38

39 293 In summary, even though the method employed here does not allow us to
40
41 294 distinguish between the possible mechanisms and pathways that could explain the
42
43 295 herbivore isotope data, it appears that the coastal location and ecology of A Lanzada
44
45 296 had a significant impact on the isotopic composition of the foods available to animals
46
47 297 (and, by implication, humans) at the site, which will need to be taken into account when
48
49 298 interpreting the human data. Since it is likely that high herbivore $\delta^{15}\text{N}$ can be attributed
50
51 299 to the specific conditions at the coast, it can further be hypothesed that animals with
52
53 300 lower nitrogen isotope values ($<\sim 6\%$) were therefore not raised at A Lanzada but
54
55
56
57
58
59
60

1
2
3 301 brought there from further inland, demonstrating the wider connections of the site in
4
5 302 antiquity.
6
7

8 303 **Omnivores.** Although only three pigs were available for sampling, their results
9
10 304 are surprisingly varied, ranging from -21.2‰ to -16.4‰ for carbon and from 8.0‰ to
11
12 305 11.7‰ for nitrogen isotope values. This indicates very different diets or management
13
14 306 strategies, from exclusively C₃-terrestrial foods to a significant input of C₄-plants or,
15
16 307 perhaps more likely, giving the scavenging behaviour of pigs and the fact that fish-
17
18 308 waste must almost certainly have been freely accessible on site, marine-based protein.
19
20 309 Sample 906b (Fig. 4), the pig with the highest $\delta^{13}\text{C}$ is, again a young animal (less than
21
22 310 12 months) and its $\delta^{15}\text{N}$ is therefore likely elevated by the suckling effect.
23
24
25

26 311 **Marine Animals.** The fish bones were tentatively identified as belonging to six
27
28 312 different individuals of four species, (see Table 1). Even though the remains are not
29
30 313 from A Lanzada itself but from a site nearby, all fish, as well as one marine mammal
31
32 314 (dolphin), are native to the Galician coast and have evidently been frequently consumed
33
34 315 by past humans (the sampled dolphin vertebra also bore butchery marks). Therefore
35
36 316 they should provide suitable reference values for the local marine ecosystem.
37
38
39

40 317 The $\delta^{13}\text{C}$ average of the marine animals is $-12.0 \pm 0.6\text{‰}$ (see Table 2), which is
41
42 318 comparable to data from the Atlantic coast (Barrett et al. 2008). While the carbon
43
44 319 isotope range is relatively narrow (1.5‰), the $\delta^{15}\text{N}$ values spread over 5.9‰,
45
46 320 presumably reflecting differences in fish ecology, trophic level and size (Tables 1-2).
47
48
49

50 321 **Human diet at A Lanzada**

51
52 322 The isotopic data from the human bone collagen are remarkably variable.
53
54 323 Human samples have a $\delta^{13}\text{C}$ range of 5.9‰ (-18.7 to -12.8‰) with a mean of -
55
56 324 $16.0 \pm 1.5\text{‰}$ (1 s.d.); as well as $\delta^{15}\text{N}$ range of 3.9‰ (10.5‰ to 14.4‰) with a moderately
57
58
59
60

1
2
3 325 high average of $12.3 \pm 0.9\%$ (1 s.d.) (Fig.5). The carbon isotope ratios from A Lanzada
4
5 326 are among the most ^{13}C -enriched values observed in any Iberian population, including
6
7 327 Mesolithic hunter-gatherers (Fuller et al. 2010; García et al. 2004; García Guixé et al.
8
9 328 2006), but excluding other Galician coastal populations (López-Costas 2012). Although
10
11 329 some individuals from Islamic sites on Ibiza (Fuller et al. 2010) and Valencia
12
13 330 (Alexander et al. 2015) exhibit similar $\delta^{13}\text{C}$, the population averages are considerably
14
15 331 lower.

16
17
18
19 332 Even though the isotopic results suggest considerable heterogeneity in the
20
21 333 human diets at A Lanzada, no significant differences exist between males and females
22
23 334 (Student t-test $\delta^{13}\text{C}$: $t_{(51)} = -1.210$ $p=0.23$; $\delta^{15}\text{N}$: $t_{(51)} = -1.030$ $p=0.308$) or different ages-
24
25 335 at-death (Kruskal-Wallis test $\delta^{13}\text{C}$: $H_{(3)} = 0.022$ $p=0.99$; $\delta^{15}\text{N}$: $H_{(3)} = 5.119$ $p=0.16$)
26
27 336 regardless of the time period.

28
29
30
31 337 Burial rite, however, and in particular tomb typology shows an association with
32
33 338 the isotopic data. It is worth to remind that tomb typology is a clear reflection of the two
34
35 339 time periods they represent, Roman and post-Roman (see methods and material
36
37 340 section). Kruskal-Wallis combined with Dunn-Bonferroni post-hoc tests indicate that
38
39 341 the $\delta^{13}\text{C}$ of individuals buried in stone-slab graves are significantly different from those
40
41 342 in simple earth graves ($p=0.023$), *tegula* graves ($p=0.001$) and stone graves ($p=0.023$;
42
43 343 K-W test $H_{(3)} = 16.65$, $p=0.001$). Differences in $\delta^{15}\text{N}$ are not significant ($H_{(3)} = 6.69$,
44
45 344 $p=0.082$). This trend is even stronger when burials from the different time-periods are
46
47 345 compared.

48
49
50
51 346 Bones of individuals in post-Roman burials (which include all stone-slab graves
52
53 347 and a number of simple earth graves) are significantly enriched in ^{13}C and ^{15}N over
54
55 348 those of the Roman burials (which comprise stone graves, *tegula* graves as well as earth

1
2
3 349 graves) (Roman vs. post-Roman M-W test $\delta^{13}\text{C}$, $U=29,475$, $p<0.000$; $\delta^{15}\text{N}$, $U=9.905$,
4
5 350 $p=0.002$). Rather than dietary variation between individuals afforded different burial
6
7 351 rites, the observed isotopic differences between grave types therefore likely indicate a
8
9 352 significant shift in diet between the Roman and the post-Roman period. There are no
10
11 353 significant differences between males and females or age-groups within each of the time
12
13 354 periods (see Tables 6-7 and Fig. 6). The unique medieval skeleton from A Lanzada, the
14
15 355 245, plots most closely with the Roman samples (Fig. 5) and while a single sample is
16
17 356 insufficient to characterise the diet of a whole period, this result at least suggests a
18
19 357 continuation of dietary traditions at the site at least in broad terms.
20
21
22

23
24 358 Similarly high carbon isotope ratios observed in all the human samples from A
25
26 359 Lanzada have been previously related to marine fish or shellfish consumption in other
27
28 360 archaeological sites from different parts of Europe (e.g. Müldner and Richards 2007;
29
30 361 Richards and Hedges 1999; Schoeninger et al. 1983). The fact that A Lanzada site is
31
32 362 surrounded by the sea and deeply related to it, a variety of fish and shell species
33
34 363 (Vázquez Varela and García Quintela 1998), as well as fish hooks and net sinkers
35
36 364 (Suárez Otero and Fariña Busto 1990) and a fish-salting factory (Rodríguez Martínez et
37
38 365 al. 2011) were identified at the site, attesting to the importance of marine resources in
39
40 366 the life of the inhabitants. The $\delta^{15}\text{N}$ average indicates the presence of moderately
41
42 367 important sources of animal protein coming from the sea or inland. Even considering
43
44 368 the difficulties in interpreting the $\Delta^{15}\text{N}$ between human diet and collagen (O'Connell et
45
46 369 al. 2012), the observed values are in agreement with a high-moderate presence of
47
48 370 marine food in diet. An alternative explanation of ^{13}C enrichment is the direct or
49
50 371 indirect (animals fed with them) ingestion of C_4 plants, such as millet (e.g. Fuller et al.
51
52 372 2010; Reitsema et al. 2010; Tafuri et al. 2009). Since millet (*Setaria italica* and, more
53
54 373 commonly, *Panicum miliaceum*) was commonly cultivated at A Lanzada and
55
56
57
58
59
60

1
2
3 374 neighbouring areas during medieval and post-medieval times (Armas Castro 1992;
4
5 375 Seijas Montero 2001) and in other areas of Galicia at least from the Iron Age onwards
6
7 376 (Aira Rodríguez et al. 1990; Dopazo Martínez et al. 1996; Ramil-Rego 1993), its
8
9 377 inclusion in peoples' diet could be as common as marine resources probably were. More
10
11 378 so, if we consider that the herbivores also show elevated ^{13}C values. In summary, the
12
13 379 observed $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ data suggest that A Lanzada people may had have a diet with an
14
15 380 important input of marine resources and/or C_4 plants, the exploitation of which is
16
17 381 supported by the historical and archaeological data available for the area.
18
19

20
21 382 In accordance with the previous discussion on the global human averages, the
22
23 383 differences observed between Roman and post-Roman individuals must have also been
24
25 384 caused by differential intake of C_4/C_3 plants and/or marine/terrestrial resources in a
26
27 385 mixed diet. Both, the $\delta^{13}\text{C}$ and the $\delta^{15}\text{N}$, are significantly different between the two
28
29 386 groups, supporting the idea that post-Roman people may have consumed more marine
30
31 387 resources and C_4 plants than the Roman individuals. However, the isotopic differences
32
33 388 between the averages (post-Roman \bar{X} -Roman \bar{X}) are wider for carbon (2.4‰) than for
34
35 389 nitrogen (0.7‰), which suggests that C_4 plants intake must have been more influential
36
37 390 in the intra-population differences. The linear trend between $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values
38
39 391 within the Roman sample shows a higher correlation ($r=0.49$) than that for the post-
40
41 392 Roman one ($r=0.00$), which is totally flat (see fig. 7), a fact that also points in this
42
43 393 direction.
44
45
46
47

48
49 394 Unfortunately, there are no ichthyological and palaeobotanical studies on A
50
51 395 Lanzada for the periods studied by us, which prevents the comparison of the diachronic
52
53 396 trends in marine exploitation or millet cultivation from different methodological
54
55 397 perspectives. Similar analyses made on other Galician sites suggest that the
56
57
58
59
60

1
2
3 398 establishment of Roman rule did not have marked effect on local Iron Age subsistence
4
5 399 strategies based on terrestrial animals (Dopazo Martínez et al. 1996). However, the
6
7 400 exploitation of marine resources seems to have had a progressive increase in NW Spain
8
9 401 from the 2nd century BC onwards (although there is no data from post-Roman times)
10
11 402 (González Gómez de Agüero 2013). This enrichment parallels a seemingly marked
12
13 403 preference for more profitable species (e.g. sardine) and widespread access to marine
14
15 404 resources in coastal and inland regions from the 2nd century AD onwards (Fuertes Prieto
16
17 405 and Fernández Rodríguez 2010; González Gómez de Agüero 2013). In fact, millets,
18
19 406 mainly *Panicum miliaceum*, continued to be very important crops in Northwest Iberia
20
21 407 throughout the Roman period (Tereso 2012). In contrast, almost nothing is known about
22
23 408 subsistence change after the demise of the Roman Empire in this area, mostly due to a
24
25 409 scarcity of well-excavated post-Roman settlements. Nevertheless, pollen evidence
26
27 410 shows a relatively sudden increase in deforestation in 5th century AD (Martínez Cortizas
28
29 411 et al. 2005) which might suggest a change in agricultural preferences.
30
31
32
33

34 35 **Marine resources and millet vs. culture and environment**

36
37
38 413 Considering that the analytical data suggest an increase in the use of marine
39
40 414 resources and possibly also C₄ plants (probably millet) by the A Lanzada population in
41
42 415 the post-Roman compared to the Roman period, we proceed to contextualize the
43
44 416 possible causes. Since it is well documented that both foods were readily available
45
46 417 during both periods (see discussion in the previous section), it is necessary to
47
48 418 understand the motivations that may have led to an increase in either resource in
49
50 419 everyday diet.
51
52

53
54 420 Millet has a short vegetative cycle and a wide germination temperature range
55
56 421 (16° to 34° C) (James et al. 2011), which allows it to adapt well to different soils and
57
58
59
60

1
2
3 422 climates (Hunt and Jones 2008). That represented a clear advantage if a hard winter or
4
5 423 other event ruined crops with longer growing seasons. Similarly, fish and seafood were
6
7 424 abundant in A Lanzada coast and people could intensify their exploitation in case of
8
9 425 need.

10
11
12 426 Some important environmental changes also occurred across the Roman/post-
13
14 427 Roman transition. Modifications in marine currents from coastal areas near to A
15
16 428 Lanzada have been registered (Lebreiro et al. 2006; Muñoz Sobrino et al. 2014), which
17
18 429 may have led to an expansion of the salt-marshes and unpredictable changes in the
19
20 430 abundance of marine resources in estuaries. The climate was also affected (Martínez-
21
22 431 Cortizas et al. 1999; Mighall et al. 2006). The temperature and humidity conditions in
23
24 432 Galicia from the 2nd to the 4th century AD were not optimal for agriculture, but the
25
26 433 climate prevailing at this time would provide conditions necessary for a good growing
27
28 434 season for the majority of crops. In contrast, during post-Roman times there was a
29
30 435 considerable increase in humidity (pointing towards a two-fold increase in rainfall)
31
32 436 (Mighall et al. 2006) and temperatures fell in a similar way to those recorded in Central
33
34 437 Europe during the decline of the Roman Empire (Martínez-Cortizas et al. 1999).
35
36 438 Although rainfall seasonality has not been analyzed for this area, these latter weather
37
38 439 conditions could have been harmful for crops production especially for delicate cereals
39
40 440 such as wheat. The millets could have been used as a complement once other crops
41
42 441 failed, and even become a substantial part of human and animal diet. In this situation,
43
44 442 fish and shellfish could have also been used as alternative foods.

45
46
47
48
49
50
51 443 A probable increase in the consumption of millet between the Roman and post-
52
53 444 Roman periods has been reported for individuals from Central and Southeast European
54
55 445 populations, and has been convincingly explained by the post-Roman migrations
56
57 446 (Hakenbeck et al. 2010; Lightfoot et al. 2012). To our knowledge, there are no similar

1
2
3 447 reports for Galicia or Iberia, a fact that may be explained by the absence of specific
4
5 448 paleodiet studies or ancient texts. The post-Roman burials of A Lanzada, and more
6
7 449 specifically the stone slabs tombs (where individuals with the highest isotopic values
8
9 450 were found), represent a burial rite frequently associated with the Suevic or Germanic
10
11 451 rule in Northwest Spain (Fariña Busto and Suarez Otero 1997). Since the *Sueves* arrived
12
13 452 in Galicia from Central Europe, where there is evidence for an increase in millet
14
15 453 consumption in the post-Roman period (Hakenbeck et al. 2010; Rösch 1998), they
16
17 454 could have brought with them a preference for this cereal. Unfortunately, little is known
18
19 455 about the *Sueves*' way of life and food preferences and it is difficult to assess their
20
21 456 impact on the local population. On the other hand, the burial rite observed at A Lanzada
22
23 457 in the post-Roman period, the WE-orientation and the absence of grave goods in
24
25 458 particular, has been also associated with Christian burials (Blanco Freijeiro et al. 1967).
26
27 459 Although Christianity played a less important role in rural areas compared to urban ones
28
29 460 (Kulikowski 2004), at least part of the post-Roman individuals might have lived as
30
31 461 Christians. The Early Christian Iberian doctrines taught abstinence from meat (Ferreiro
32
33 462 2008) and ancient texts promoted the idea of a largely vegetarian diet with occasional
34
35 463 supplements of fish: "*the monks were encouraged to eat vegetables, greens, beans, and*
36
37 464 *on occasion fresh or salt fish* " (Campos Ruíz and Roca Melia 1971). An increase in
38
39 465 fish consumption at A Lanzada might therefore be explained by their conversion to
40
41 466 Christianity. Nevertheless, these early food rules were largely directed at a monastic
42
43 467 (rather than lay) audience and economic status should therefore have had a greater
44
45 468 influence than religion on dietary preferences.

51
52
53 469 Parallel to the arrival of new people and religions, important changes to the
54
55 470 economy of NW Iberian villages also took place. The Roman period (1st century AD
56
57 471 onwards) saw a high level of economic activity and coastal villages such as A Lanzada

1
2
3 472 were thriving trading settlements on a maritime route that connected the Mediterranean
4
5 473 with the North Sea (Naveiro López 1991). From the 4th century AD onwards, the
6
7 474 Roman Empire faced an economic crisis on the Iberian Peninsula which ultimately led
8
9 475 to its political break down (Kulikowski 2004) and resulted in a stark deterioration of the
10
11 476 Atlantic commerce (for NW Spain see Naveiro López 1991). The situation became
12
13 477 worse with the Germanic invasions and coastal settlements such as A Lanzada faced
14
15 478 isolation and political insecurity. The establishment of the Suevic Kingdom also appears
16
17 479 to have coincided with some dramatic environmental changes most likely resulting from
18
19 480 intensive economic activities, such as increasing deforestation, soil erosion,
20
21 481 metallurgical activity and pollution (Kylander et al. 2005; Martínez Cortizas et al. 2005;
22
23 482 Ramil-Rego et al. 1998). These factors could have caused a modification to the daily
24
25 483 subsistence strategies of the inhabitants of A Lanzada, such as in the way they exploited
26
27 484 their local resources, i.e. eating more fish if it was impossible to sell it.
28
29
30
31

32
33 485 In summary, the environmental and historical events that took place during the
34
35 486 occupation of the site provide observed number of plausible explanations to the
36
37 487 observed changes in human diet. Nevertheless, and especially since the factors are all
38
39 488 interrelated, it is difficult to distinguish whether these changes were primarily a
40
41 489 response to local or conditions or to the events that affected the Iberian Peninsula as a
42
43 490 whole at the time.
44
45
46
47
48

491

492 CONCLUSIONS

51
52 493 The analysis of this Roman and post-Roman community located on the very
53
54 494 fringes of Europe has revealed a very distinct food-web, which was highly dependent on
55
56 495 the surrounding marine environment. The connection with the sea and its resources was
57
58
59
60

1
2
3 496 presumably working in many ways. The results suggest that animals were managed
4
5 497 using different strategies, which includes possible grazing on saline pasture or the use of
6
7 498 seaweed foddering. A variable amount of C₄ plants, such as millet but also the
8
9 499 halophyte *Spartina sp.*, would also have been present in the diet of herbivores.
10
11 500 Humans' diet may have been also closely connected to marine resources along the time
12
13 501 increasing the consumed amount during specific periods (e.g. post-Roman time). The
14
15 502 observed strong relationship with the sea emphasizes the importance of considering a
16
17 503 local perspective to understand dietary preferences, even in the well-defined and known
18
19 504 Roman period.
20
21
22

23
24 505 Our study also suggests that the end of the Roman Empire and the Migration
25
26 506 Period had a profound impact on diet, as it presumably had in the European landscape.
27
28 507 Other authors' conclusions (Hakenbeck et al. 2010; Lightfoot et al. 2012) about the
29
30 508 increase in millet consumption during the post-Roman or the Early Medieval period
31
32 509 with respect to the Roman times are in agreement with our data. In the case of A
33
34 510 Lanzada, the rise of the use of millet may have been parallel to an increase in the
35
36 511 exploitation of local marine resources. We argue that the palaeoenvironmental
37
38 512 conditions, and more specifically the climate, could have played an important role in the
39
40 513 observed change in food preferences. However, based on the evidence available, it is
41
42 514 not possible to distinguish the effects of climate deterioration in NW Spain at the time
43
44 515 from the changes likely brought about by the arrival of new people, the *Sueves*, who
45
46 516 brought a new political and socioeconomic system with them. Understanding how the
47
48 517 establishment of Suevic rule in the area may have influenced the subsistence base and in
49
50 518 particular the reliance of marine resources and the cultivation of millet is still an open
51
52 519 question which will be addressed by further studies.
53
54
55
56
57
58
59
60

1
2
3 520 This project is the first stable isotope study of a Roman/post-Roman community
4
5 521 in the Iberian Peninsula (some isolated skeletons have been reported) and the largest on
6
7 522 the Southwestern fringe of the Roman Empire. For future palaeodietary studies, we
8
9 523 believe that well-defined local palaeoenvironmental reconstruction can be a highly
10
11 524 successful way to examine the causes of dietary change and to distinguish between
12
13 525 environmental factors and changes in cultural or economic preferences. This seems
14
15 526 particularly important in transitional periods such as after the end of the Roman Empire,
16
17 527 where profound environmental changes were coupled with significant political, social
18
19 528 and economic change, including migration. We consider that additional investigations
20
21 529 are necessary to understand the dietary adaptations in neighbouring areas and periods
22
23 530 will provide further insights into these transitions.
24
25
26
27

28 **ACKNOWLEDGEMENTS**

29
30
31 532 OLC is funded by Plan Galego I2C mod.A. The research was funded by projects
32
33 533 “Antropoloxía dos restos óseos humanos de Galicia” (Dirección Xeral de Patrimonio
34
35 534 Histórico), GPC2014/009 and R2014/001 of Xunta de Galicia. Many thanks to the
36
37 535 Museo Provincial de Pontevedra and Museo Provincial de Ourense for providing access
38
39 536 to the skeletal collections. Advice given by Miguel Botella has been a great help in
40
41 537 classifying skeletal features. An especial mention goes to Jose M. Rey Salgado and
42
43 538 Aleks Pluskowski for their help with the animal species identification, and to Eduardo
44
45 539 González and Víctor Bejega for the information about the fish strategies. Thanks to
46
47 540 Milton Núñez, Chus Casais, Antonio Martínez Cortizas, Francisco Fariña, Rafa
48
49 541 Rodríguez Martínez, David Varela, Antonio de la Peña, David Fernández Abella and
50
51 542 Ángel Costas for their ideas and advices to this study. We thank Tim Mighall for
52
53 543 reviewing and commenting a previous version of this manuscript.
54
55
56
57
58
59
60

LITERATURE CITED

- 544
- 545
- 546 Aira Rodríguez MJ, Ramil Rego P, and Alvarez Núñez A. 1990. Estudio paleocarpológico realizado en el
547 Castro de Penalba (Campolameiro, Pontevedra, España). *BotComplut* 16:81-89.
- 548 Alexander MM, Gerrard CM, Gutiérrez A, and Millard AR. 2015. Diet, society, and economy in late
549 medieval Spain: Stable isotope evidence from Muslims and Christians from Gandía, Valencia.
550 *Am. J. Phys. Anthropol.* 156(2):263-273.
- 551 Arias P, and Schulting RJ. 2010. Análisis de isótopos estables sobre los restos humanos de la Braña-
552 Arintero. Aproximación a la dieta de los grupos mesolíticos de la Cordillera Cantábrica. In:
553 Vidal J, and Prada ME, editors. *Los hombres mesolíticos de la cueva de La Braña-Arintero*
554 (Valdelugueros, León). Valladolid: Junta de Castilla y León. Estudios y Catálogos. p 130–137.
- 555 Armas Castro J. 1992. Pontevedra en los siglos XIII a XV, configuración y desarrollo de una villa
556 marinera en la Galicia Medieval. Pontevedra: Fundación Pedro Barrié de la Maza Conde de
557 Fenosa.
- 558 Balasse M, and Tresset A. 2002. Early weaning of Neolithic domestic cattle (Bercy, France) revealed by
559 intra-tooth variation in nitrogen isotope ratios. *J. Archaeol. Sci.* 29(8):853-859.
- 560 Balasse M, Tresset A, and Ambrose SH. 2006. Stable isotope evidence (delta C-13, delta O-18) for winter
561 feeding on seaweed by Neolithic sheep of Scotland. *J. Zool.*270(1):170-176.
- 562 Barrett J, Johnstone C, Harland J, Van Neer W, Eryvynck A, Makowiecki D, Heinrich D, Hufthammer
563 AK, Enghoff IB, Amundsen C et al. . 2008. Detecting the medieval cod trade: a new method and
564 first results. *J. Archaeol. Sci.* 35(4):850-861.
- 565 Blanco Freijeiro A, Fusté Ara ME, and García Alen A. 1961. La necrópolis galaico-romana de La
566 Lanzada (Noalla, Pontevedra), I. *Cuad Estud Gallegos* 16:141-158.
- 567 Blanco Freijeiro A, Fusté Ara ME, and García Alen A. 1967. La necrópolis galaico-romana de La
568 Lanzada (Noalla, Pontevedra), II. *Cuad Estud Gallegos* 22:5-23, 129-155.
- 569 Britton K, Muldner G, and Bell M. 2008. Stable isotope evidence for salt-marsh grazing in the Bronze
570 Age Severn Estuary, UK: implications for palaeodietary analysis at coastal sites. *J. Archaeol.*
571 *Sci.* 35(8):2111-2118.
- 572 Büntgen U, Tegel W, Nicolussi K, McCormick M, Frank D, Trouet V, Kaplan JO, Herzig F, Heussner K-
573 U, Wanner H et al. . 2011. 2500 years of European climate variability and human susceptibility.
574 *Science* 331(6017):578-582.
- 575 Campos Ruíz J, and Roca Melia I. 1971. San Leandro, San Isidoro, San Fructuoso, Reglas monásticas de
576 la España visigoda, Los tres libros de las “Sentencias”,. Madrid: Biblioteca de Autores
577 Cristianos. Santos Padres Españoles 2.
- 578 Carro Otero J, Masa Vázquez MC, and Varela Ogando ML. 1987. Un nuevo enterramiento de la
579 necrópolis galaico-romana de La Lanzada (Noalla-Pontevedra). *Pontevedra Arqueológica* 2:229-
580 252.
- 581 Castellanos S, and Viso IM. 2005. The local articulation of central power in the north of the Iberian
582 Peninsula (500–1000). *Early Medieval Europe* 13(1):1-42.
- 583 Cloern JE, Canuel EA, and Harris D. 2002. Stable carbon and nitrogen isotope composition of aquatic
584 and terrestrial plants of the San Francisco Bay estuarine system. *Limnol. Oceanogr.* 47(3):713-
585 729.
- 586 Collins MJ, and Galley P. 1998. Towards an optimal method of archaeological collagen extraction: the
587 influence of pH and grinding. *Anc. Biomol.* 2:209-222.
- 588 Craig OE, Biazzo M, O’Connell TC, Garnsey P, Martínez-Labarga C, Lelli R, Salvadei L, Tartaglia G,
589 Nava A, Reno L et al. . 2009. Stable Isotopic Evidence for Diet at the Imperial Roman Coastal
590 Site of Velia (1st and 2nd Centuries AD) in Southern Italy. *Am. J. Phys. Anthropol.* 139(4):572-
591 583.
- 592 Davis M. 2002. Putting meat on the bone: an investigation into palaeodiet in the Balearic Islands using
593 carbon and nitrogen stable isotope analysis. In: Waldren W, and Ensenyat J, editors. *World*
594 *islands in prehistory: international insular investigations V Deià International Conference of*
595 *Prehistory.* Oxford: BAR International Series 1095. p 198–216.
- 596 Díaz PC. 2011. *El reino suevo (411-585). Tres Cantos, Madrid: Akal, D L.*
- 597 Dopazo Martínez A, Fernández Rodríguez C, and Ramil Rego P. 1996. Arqueometría aplicada a
598 yacimientos galaico-romanos del NW peninsular, valoración de la actividad agrícola y ganadera.
599 In: Ramil Rego P, Fernández Rodríguez C, and Rodríguez Guitián M, editors. *Biogeografía*
600 *pleistocena -holocena de la Península Ibérica.* Santiago de Compostela: Consellería de Cultura. p
601 317-332.

- 1
2
3 602 Fariña Busto F. 1975. Excavación de A Lanzada (Sanxenxo-Pontevedra); Informe preliminar de la
4 603 campaña. Museo de Pontevedra 29:165-173.
5 604 Fariña Busto F, and Filgueira Valverde JE. 1974. Plan Nacional de Excavaciones 1973: A Lanzada
6 605 (Sanxenxo, Pontevedra). Museo de Pontevedra 28:83-86.
7 606 Fariña Busto F, and Suarez Otero J. 1997. As necrópoles xermano-suélicas. In: García Iglesias XM,
8 607 editor. Galicia Castrexa e Romana. Lugo: Consellería de Cultura e Comunicación Social.
9 608 Ferreiro A. 2008. De prohibitione carnis. Meat Abstention and the Priscillianists. Zeitschrift für Antikes
10 609 Christentum 11(3):464-478.
11 610 Ferreiro García IA, Acuña Castroviejo R, and Durán Neira C. 1993. As Algas en Galicia : alimentación e
12 611 outros usos: Dirección Xeral de Marisqueo e Acuicultura, D.L Galicia.
13 612 Ferro-Vázquez C, Martínez-Cortizas A, Nóvoa-Muñoz JC, Ballesteros-Arias P, and Criado-Boado F.
14 613 2014. 1500 years of soil use reconstructed from the chemical properties of a terraced soil
15 614 sequence. Quat Int (346):28-40.
16 615 Filgueira Valverde JE, and Blanco Freijeiro A. 1962. Excavaciones de La Lanzada. Noticiario
17 616 Arqueológico Hispánico V (1956-1961):137-152.
18 617 Fraser RA, Bogaard A, Heaton T, Charles M, Jones G, Christensen BT, Halstead P, Merbach I, Poulton
19 618 PR, Sparkes D et al. . 2011. Manuring and stable nitrogen isotope ratios in cereals and pulses:
20 619 towards a new archaeobotanical approach to the inference of land use and dietary practices. J.
21 620 Archaeol. Sci. 38(10):2790-2804.
22 621 Fuertes Prieto MN, and Fernández Rodríguez C. 2010. El comercio y consumo de moluscos en época
23 622 romana en Asturica Augusta (León). Férvedes 6:147-158.
24 623 Fuller B, Marquez-Grant N, and Richards MP. 2010. Investigation of Diachronic Dietary Patterns on the
25 624 Islands of Ibiza and Formentera, Spain: Evidence from Carbon and Nitrogen Stable Isotope
26 625 Ratio Analysis. Am. J. Phys. Anthropol. 143(4):512-522.
27 626 Fuller BT, Molleson TI, Harris DA, Gilmour LT, and Hedges REM. 2006. Isotopic evidence for
28 627 breastfeeding and possible adult dietary differences from Late/Sub-Roman Britain. Am. J. Phys.
29 628 Anthropol.129(1):45-54.
30 629 García-Guixé E, Martínez-Moreno J, Mora R, Nunez M, and Richards MP. 2009. Stable isotope analysis
31 630 of human and animal remains from the Late Upper Palaeolithic site of Balma Guilanya,
32 631 southeastern Pre-Pyrenees, Spain. J. Archaeol. Sci. 36(4):1018-1026.
33 632 García E, Subirà ME, and Richards MP. 2004. Régime et société d'après l'analyse des isotopes stable:
34 633 l'exemple de la population de "Can Reinés" (Mallorca, Espagne, 600 ap. J.C.). Anthropol 7:171-
35 634 176.
36 635 García Guixé E, Richards MP, and Subira ME. 2006. Palaeodiets of humans and fauna at the Spanish
37 636 Mesolithic site of El Collado. Curr. Anthropol. 47(3):549-556.
38 637 González Gómez de Agüero E. 2013. La ictiofauna de los yacimientos arqueológicos del noroeste de la
39 638 Península Ibérica. Doctoral thesis. León: University of León. 474 p.
40 639 González Ruibal A. 2004. Facing two seas: Mediterranean and Atlantic contacts in the North-West of
41 640 Iberia in the first millenium BC. Oxford Journal of Archaeology 23(3):287-317.
42 641 Hakenbeck S, McManus E, Geisler H, Grupe G, and O'Connell TC. 2010. Diet and Mobility in Early
43 642 Medieval Bavaria: A Study of Carbon and Nitrogen Stable Isotopes. Am. J. Phys.
44 643 Anthropol.143(2):235-249.
45 644 Heaton THE. 1987. The N-15/N-14 ratios of plants in South-Africa and Namibia- Relationship to climate
46 645 and coastal saline environments Oecologia 74(2):236-246.
47 646 Hedges REM, Clement JG, Thomas CDL, and O'Connell TC. 2007. Collagen turnover in the adult
48 647 femoral mid-shaft: Modeled from anthropogenic radiocarbon tracer measurements. Am. J. Phys.
49 648 Anthropol 133(2):808-816
50 649 Hunt H, and Jones M. 2008. Pathways across Asia : exploring the history of Panicum and Setaria in the
51 650 Indian subcontinent. Pragdhara 18:53-68.
52 651 James TK, Rahman A, McGill CR, and Trivedi PD. 2011. Biology and survival of broom corn millet
53 652 (Panicum miliaceum) seed. N. Z. Plant Prot. 64:142-148.
54 653 Jim S, Ambrose SH, and Evershed RP. 2004. Stable carbon isotopic evidence for differences in the
55 654 dietary origin of bone cholesterol, collagen and apatite: implications for their use in
56 655 palaeodietary reconstruction. Geochim Cosmochim Acta 68(1):61-72.
57 656 Kaal J, López-Costas O, and Martínez Cortizas A. 2016. Diagenetic effects on pyrolysis fingerprints of
58 657 extracted collagen in archaeological human bones from NW Spain, as determined by pyrolysis-
59 658 GC-MS. J. Archaeol. Sci. 65:1-10.
60 659 Kulikowski M. 2004. Late Roman Spain and its Cities. Baltimore: The Johns Hopkins University Press.

- 1
2
3 660 Kylander ME, Weiss DJ, Martínez Cortizas A, Spiro B, García-Sánchez R, and Coles BJ. 2005. Refining
4 661 the pre-industrial atmospheric Pb isotope evolution curve in Europe using an 8000 year old peat
5 662 core from NW Spain. *Earth Planet. Sci. Lett.* 240(2):467-485.
- 6 663 Lebreiro SM, Frances G, Abrantes FFG, Diz P, Bartels-Jonsdottir HB, Stroynowski ZN, Gil IM, Pena
7 664 LD, Rodrigues T, Jones PD et al. . 2006. Climate change and coastal hydrographic response
8 665 along the Atlantic Iberian margin (Tagus Prodelta and Muros Ria) during the last two millennia.
9 666 *Holocene* 16(7):1003-1015.
- 10 667 Lightfoot E, Šlaus M, and O'Connell TC. 2012. Changing cultures, changing cuisines: Cultural transitions
11 668 and dietary change in iron age, roman, and early medieval croatia. *Am. J. Phys. Anthropol.*
12 669 148(4):543-556.
- 13 670 Longin R. 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230(5291):241-242.
- 14 671 López-Costas O. 2015. Taphonomy and burial context of the Roman/post-Roman funerary areas (2nd to
15 672 6th centuries AD) of A Lanzada, NW Spain. *Estudos do Quaternário, APEQ* 12:55-67.
- 16 673 López-Costas O. 2012. Antropología de los restos óseos humanos de Galicia: estudio de la población
17 674 romano y medieval gallega. Doctoral thesis. Granada: University of Granada. 555 p.
- 18 675 López-Costas O, Lantes-Suárez O, and Martínez Cortizas A. 2016. Chemical compositional changes in
19 676 archaeological human bones due to diagenesis: type of bone vs soil environment. *J. Archaeol.*
20 677 *Sci.* in press.
- 21 678 López-Costas O, Müldner G, and Martínez Cortizas A. 2015. Diet and lifestyle in Bronze Age Northwest
22 679 Spain: the collective burial of Cova do Santo. *J. Archaeol. Sci.* 55(0):209-218.
- 23 680 Martínez-Cortizas A, Pontevedra-Pombal X, García-Rodeja E, Novoa-Munoz JC, and Shotyky W. 1999.
24 681 Mercury in a Spanish peat bog: Archive of climate change and atmospheric metal deposition.
25 682 *Science* 284(5416):939-942.
- 26 683 Martínez Cortizas A, Mighall T, Pontevedra Pombal X, Novoa Munoz JC, Peiteado Varela E, and Piñeiro
27 684 Rebolo R. 2005. Linking changes in atmospheric dust deposition, vegetation change and human
28 685 activities in northwest Spain during the last 5300 years. *Holocene* 15(5):698-706.
- 29 686 Mighall TM, Martínez Cortizas A, Biester H, and Turner SE. 2006. Proxy climate and vegetation changes
30 687 during the last five millennia in NW Iberia: Pollen and non-pollen palynomorph data from two
31 688 ombrotrophic peat bogs in the North Western Iberian Peninsula. *Rev. Palaeobot. Palynol.* 141(1-
32 689 2):203-223.
- 33 690 Mintz SW, and Du Bois CM. 2002. The anthropology of food and eating. *Annu Rev Anthropol* 31:99-
34 691 119.
- 35 692 Müldner G, and Richards MP. 2007. Diet and diversity at later medieval fishergate: The isotopic
36 693 evidence. *Am. J. Phys. Anthropol.* 134:162-174.
- 37 694 Munde M. 2009. An isotopic approach to diet in Medieval Spain. In: Baker S, Allen M, Middle S, and
38 695 Poole K, editors. *Food and Drink in Archaeology 2: University of Nottingham Conference 2009:*
39 696 *Prospect, Totnes.*
- 40 697 Muñoz Sobrino C, García-Moreiras I, Castro Y, Martínez Carreño N, de Blas E, Fernández Rodríguez C,
41 698 Judd A, and García-Gil S. 2014. Climate and anthropogenic factors influencing an estuarine
42 699 ecosystem from NW Iberia: new high resolution multiproxy analyses from San Simón Bay (Ría
43 700 de Vigo). *Quat Sci Rev* 93(0):11-33.
- 44 701 Musset L. 1975. *The Germanic invasions : The making of Europe Ad 400-600.* London: Elek.
- 45 702 Naveiro López J. 1991. *El comercio antiguo en el NW Peninsular. Lectura Histórica del Registro*
46 703 *Arqueológico.* Sada: do Castro.
- 47 704 O'Connell TC, Kneale CJ, Tasevska N, and Kuhnle GGC. 2012. The diet-body offset in human nitrogen
48 705 isotopic values: A controlled dietary study. *Am. J. Phys. Anthropol* 149(3):426-434.
- 49 706 Pérez García JM. 1979. *Un Modelo de sociedad rural de antiguo régimen en la Galicia costera : la*
50 707 *península del Salnés : (Jurisdicción de La Lanzada).* Santiago de Compostela: Universidade de
51 708 Santiago de Compostela. 416 p.
- 52 709 Pestle WJ, Crowley BE, and Weirauch MT. 2014. Quantifying Inter-Laboratory Variability in Stable
53 710 Isotope Analysis of Ancient Skeletal Remains. *PLoS ONE* 9(7):e102844.
- 54 711 Philpott R. 1991. *Burial Practices in Roman Britain. A Survey of Grave Treatment and Furnishing AD*
55 712 *43-410.* Oxford: Tempus Reparatum.
- 56 713 Ramil-Rego P. 1993. *Paleoethnobotánica de yacimientos arqueológicos holocenos de Galicia (N.O.*
57 714 *Cantábrico).* *Munibe* 45:165-174.
- 58 715 Ramil-Rego P, Munoz-Sobrino C, Rodriguez-Guitian M, and Gomez-Orellana L. 1998. Differences in the
59 716 vegetation of the North Iberian Peninsula during the last 16,000 years. *Plant Ecol* 138(1):41-62.
- 60 717 Reitsema LJ, Crews DE, and Polcyn M. 2010. Preliminary evidence for medieval Polish diet from carbon
718 and nitrogen stable isotopes. *J. Archaeol. Sci.* 37(7):1413-1423.

- 1
2
3 719 Richards MP, and Hedges REM. 1999. Stable isotope evidence for similarities in the types of marine
4 720 foods used by late mesolithic humans at sites along the Atlantic coast of Europe. *J. Archaeol.*
5 721 *Sci.* 26(6):717-722.
- 6 722 Rodríguez Martínez RM, Aboal Fernandez R, Castro Hierro V, Candela Cereijo C, and Ayán Vila JM.
7 723 2011. Una posible factoría prerromana en el noroeste. Primeras valoraciones de la intervención
8 724 en el Campo de A Lanzada (Sanxenxo, Pontevedra). *Férvedes* 7:159-168.
- 9 725 Rösch M. 1998. The history of crops and crop weeds in south-western Germany from the Neolithic period
10 726 to modern times, as shown by archaeobotanical evidence. *Veg Hist Archaeobot* 7(2):109-125.
- 11 727 Sage R, and Monson R, editors. 1999. *C4 plant biology*. San Diego: CA: Academic Press.
- 12 728 Salazar-García DC, Aura JE, Olària CR, Talamo S, Morales JV, and Richards MP. 2014. Isotope
13 729 evidence for the use of marine resources in the Eastern Iberian Mesolithic. *J. Archaeol. Sci.*
14 730 42(0):231-240.
- 15 731 Schoeninger M, DeNiro M, and Tauber H. 1983. Stable nitrogen isotope ratios of bone collagen reflect
16 732 marine and terrestrial components of prehistoric human diet. *Science* 220:1381-1383.
- 17 733 Schwarcz HP, and Schoeninger MJ. 1991. Stable isotope analyses in human nutritional ecology. *Yearb.*
18 734 *Phys. Anthropol.* 34:283-321.
- 19 735 Sealy J. 2001. Body tissue chemistry and Palaeodiet. In: Brothwell DR, and Pollard AM, editors.
20 736 *Handbook of Archaeological Sciences*. Chichester: John Wiley and Sons. p 269-279.
- 21 737 Seijas Montero M. 2001. Los prioratos de San Martín Pinario en el Sudoeste gallego: dominio territorial y
22 738 situación económica a fines del Antiguo Régimen. *Obradoiro Hist Mod* 10:121-140.
- 23 739 Spurr MS. 1986. *Arable Cultivation in Roman Italy, 200 B.C.- A.D.100*. London: Society for the
24 740 Promotion of Roman Studies. *J Roman Stud. Monographs*, 3.
- 25 741 Stevens RE, Lightfoot E, Hamilton J, Cunliffe B, and Hedges REM. 2010. Stable isotope investigations
26 742 of the Danebury. *Oxford Journal of Archaeology* 29(4):407-428.
- 27 743 Suarez Otero J, and Fariña Busto F. 1990. A Lanzada (Sanxenxo, Pontevedra), definición e interpretación
28 744 de un yacimiento castreño atípico. Apuntes para un estudio de los intercambios protohistóricos
29 745 en la costa atlántica peninsular. *Madrider Mitteilungen* 31:309-337.
- 30 746 Syme R. 1934. The Spanish War of Augustus (26-25 BC). *The American Journal of Philology* 55:193-
31 747 317.
- 32 748 Tafuri MA, Craig OE, and Canci A. 2009. Stable Isotope Evidence for the Consumption of Millet and
33 749 Other Plants in Bronze Age Italy. *Am. J. Phys. Anthropol.* 139(2):146-153.
- 34 750 Tereso JP. 2012. Environmental Change, agricultural development and social trends in NW Iberia from
35 751 the Late Prehistory to the Late Antiquity. Porto: Faculdade de Ciências do Porto. PhD thesis.
- 36 752 Toynbee JM. 1971. *Death and burial in the Roman World*. London: Thames and Hudson.
- 37 753 Valdés-Bermejo E, and Silva Pando J. 1986. *Vegetación del istmo de la Lanzada*. Pontevedra: Diputación
38 754 Provincial, Departamento de Cultura.
- 39 755 Van Groenigen J-W, and van Kessel C. 2002. Salinity-induced Patterns of Natural Abundance Carbon-13
40 756 and Nitrogen-15 in Plant and Soil. *Soil Sci Soc Am J* 66(2):489-498.
- 41 757 Van Strydonck M, Boudini M, and Ervynck A. 2002. Stable isotopes (¹³C and ¹⁵N) and diet: animal and
42 758 human bone collagen from prehistoric sites in Mallorca, Menorca and Formenter (Balearic
43 759 Islands, Spain). In: Waldren W, and Ensenyat J, editors. *World islands in prehistory:*
44 760 *international insular investigations V Deià International Conference of Prehistory*. Oxford: BAR
45 761 International Series. p 189-197.
- 46 762 Van Strydonck M, Boudini M, and Ervynck A. 2005. Spatial and temporal variation of dietary habits
47 763 during the prehistory of the Balearic Islands as reflected by ¹⁴C, ^{d15}N and ^{d13}C analyses on
48 764 human and animal bones. *Mayurga* 30:523-541.
- 49 765 Vázquez Varela JM, and García Quintela MV. 1998. *A vida cotiá na Galicia castrexa*. Santiago de
50 766 Compostela: Biblioteca de divulgación.
- 51 767 Villares R, Carral E, Lorenzana F, and Mosquera EL. 2007. Drift-seaweed evaluation for fertilizer use in
52 768 galiza (Northwest Spain): Tissue elemental characterization and site-sampling differences. *J*
53 769 *Sustain Agric* 31(1):45-60.
- 54 770 Virginia RA, and Delwiche CC. 1982. Natural N-15 abundance of presumed N₂-fixing and non N₂-fixing
55 771 plants from selected ecosystems. *Oecologia* 54(3):317-325.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Map of the site location including the main fresh water and marine resources near the site.
78x52mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Area of the site at present. The two phases of the necropolis and other archaeological sectors are emphasized, such as the Romanesque church (1), the medieval tower (2) and the Iron Age-Roman settlement area with the seafood salting installation (3). Number 4 indicates where the medieval skeleton was found. Modified from Google Earth ©2014 Google
338x177mm (96 x 96 DPI)




Figure 3. Pictures from two representative burials of the Roman (A) and post-Roman (B) phases of A Lanzada necropolis. The picture "A" corresponds to the earth grave of skeleton 202, which contained an imbrex placed under the head, ceramic vessels as gravegoods and hobnails (remains of nailed shoes). The picture "B" shows the stone slab tomb of skeleton 249. A second individual can be seen at the feet of 249, possibly representing an older burial that was moved aside (multiple non-consecutive burial).

63x34mm (300 x 300 DPI)




Figure 4. Animals bone collagen isotope results. Terrestrial samples are grouped by bone or tooth (dentin).
295x176mm (96 x 96 DPI)




Fig. 5. Bone collagen $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ values of animal and human samples from Lanzada grouped by time period.
270x174mm (96 x 96 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 6. Scatter plot of carbon and nitrogen isotope ratios for humans according to grave typology. Since the information about the burial typology of some skeletons was lost or not totally clear, the sample plot here is smaller than the represented according to the period of use.
240x170mm (96 x 96 DPI)




Figure 7. Scatter plot and linear regression of carbon and nitrogen isotope ratios for Roman and post-Roman humans
268x178mm (96 x 96 DPI)

TABLE 1. Carbon and nitrogen stable isotope ratios, collagen quality indicators, and tentatively species identification of the non human animals.

Sample number	Site	Animal group	Specie	Sample area	$\delta^{13}C$	$\delta^{15}N$	C/N	%C	%N	%Coll
904	A Lanzada	Herbivore	Cattle	horn	-18.8	8.7	3.2	35.5	12.8	11.3
907a	"	"	Cattle	calcaneus	-20.5	4.5	3.3	33.7	11.9	6.6
907d	"	"	Cattle	rib	-20.4	7.8	3.3	36.9	13.2	10.5
907f	"	"	Cattle	metapodial	-20.0	7.2	3.3	28.8	10.3	4.1
911	"	"	Cattle	tooth	-19.5	8.1	3.3	41.7	14.8	7.9
913a	"	"	Cattle	tooth	-20.5	8.1	3.3	42.7	15.0	10.2
913d	"	"	Cattle	talus	-21.5	5.8	3.4	40.3	13.7	3.1
914a	"	"	Cattle	tibia	-21.2	5.4	3.3	42.3	15.1	5.1
914b	"	"	Cattle	radio	-21.0	8.1	3.3	35.9	12.8	8.2
906c	"	"	Cattle/horse	tooth	-19.9	7.3	3.3	41.7	14.9	10.5
907e	"	"	Cattle/horse	humerus	-21.3	10.0	3.5	25.3	8.5	5.9
903	"	"	Ovicaprid	tibia	-19.5	5.2	3.4	29.2	10.1	5.4
905	"	"	Ovicaprid	metapodial	-19.1	9.3	3.2	33.7	12.2	6.9
906a	"	"	Ovicaprid	femur	-17.7	11.2	3.5	22.4	7.5	5.1
906d	"	"	Ovicaprid	cranium	-19.9	6.9	3.3	43.5	15.4	2.8
907b	"	"	Ovicaprid	cranium	-19.8	6.4	3.3	36.9	12.9	6.2
908	"	"	Ovicaprid	metapodial	-20.5	8.5	3.2	36.6	13.1	10.6
909	"	"	Ovicaprid	tibia	-19.8	5.2	3.2	41.0	14.8	15.6
910	"	"	Ovicaprid	tooth	-20.0	9.2	3.3	41.9	15.1	5.3
913b	"	"	Ovicaprid	tooth	-19.9	7.1	3.3	27.2	9.5	11.0
913c	"	"	Ovicaprid	humerus	-21.1	9.1	3.4	34.5	11.8	5.1
906b	"	Omnivore	Pig	femur	-16.4	11.7	3.3	40.5	14.3	2.6
907c	"	"	Pig	tibia	-19.2	8.9	3.4	33.8	11.6	2.5
912	"	"	Pig	metapodial	-21.2	8.0	3.3	34.2	12.0	8.6
936	Pontevedra	Marine fish	John Dory (<i>Zeus faber</i>)	jaw	-12.0	9.2	3.2	43.3	15.9	19.9
937	"	"	Hake	jaw	-11.8	15.1	3.1	38.8	14.6	15.8
938	"	"	Hake	jaw	-13.0	14.1	3.2	41.3	15.0	13.5
939	"	"	Red pogy (? <i>Pagruspagrus</i>)	jaw	-11.5	13.9	3.1	42.9	16.1	16.3
940	"	"	Tuna/Bonito	vertebra	-12.5	11.1	3.2	43.2	15.8	19.4
941	"	"	Tuna/Bonito	vertebra	-11.5	12.1	3.2	40.0	14.7	19.2
942	"	Marine mammal	Dolphin	vertebra	-11.7	11.9	3.2	40.5	14.6	21.6

TABLE 2. Statistical summary of the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ results for terrestrial animals from A Lanzada, NW Spain, and marine animals from a site 20 km away (Pontevedra).

	Cattle	Cattle/Horse	Sheep/goat	Suid	Marine fish	Dolphin
n	9	2	9	3	6	1
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\text{‰}$)	-20.4 \pm 0.9	-19.9, -21.3	-19.9 \pm 0.6	-18.9 \pm 2.4	-12.0 \pm 0.6	-11.7
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\text{‰}$)	7.1 \pm 1.5	7.3, 10.0	7.5 \pm 1.6	9.5 \pm 1.9	12.6 \pm 2.2	-11.9

TABLE 3. Carbon and nitrogen stable isotope ratios, collagen quality indicators, anthropological and archaeological information for A Lanzada humans.

Sample number	Period	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	%C	%N	%Coll	Sex	Age group	Sampled bone	Site area	Orientation	Grave typology	Notes
201	Roman	-16.7	11.6	3.4	40.4	13.8	3.0	M	MA	jaw	N area	SW-NE	eg	H
202	R.	-17.5	12.3	3.6	26.6	8.7	4.5	M	*	jaw	N area	S-N	tg	G; H
203	R.	-16.5	13.0	3.3	40.5	14.3	2.7	M	YA	jaw	N area	S-N	tg	G; H
204	R.	-17.2	12.6	3.3	35.8	12.5	14.6	F	*	jaw	N area	S-N	tg	G
205	R.	-17.2	11.6	3.2	42.9	15.4	16.5	F	*	jaw	N area	S-N	tg	G
206	R.	-17.0	11.7	3.3	37.7	13.3	7.3	F	*	jaw	N area	S-N	eg	G
207	R.	-14.9	13.9	3.4	26.9	9.2	6.3	M	YA	jaw	N area	S-N	eg	
208	R.	-17.0	12.1	3.4	42.3	14.5	3.1	M	YA	jaw	N area	*	eg	
209	R.	-15.8	14.3	3.3	43.1	15.3	14.4	F	MA	jaw	N area	S-N	tg	G
210	R.	-15.3	13.1	3.3	41.1	14.4	3.9	F	YA	jaw	N area	NE-SW	eg	
211	R.	-16.8	11.3	3.3	23.7	8.3	3.5	M	OA	jaw	N area	S-N	eg	
212	R.	-18.7	11.5	3.5	29.8	9.8	4.1	M	MA	jaw	*	*	*	
213	R.	-16.2	12.4	3.2	42.3	15.4	19.7	M	OA	jaw	N area	SE-NW	tg	
214	R.	-17.7	11.2	3.3	41.9	14.6	4.7	F	*	fibula	*	*	*	
215	R.	-17.4	11.8	3.2	38.1	13.8	21.6	M	YA	fibula	N area	SE-NW	tg	G; H
216	R.	-16.4	12.9	3.3	38.4	13.7	12.3	M	MA	rib	N area	SE-NW	tg	G; H
217	R.	-16.6	12.2	3.3	43.5	15.6	17.6	M	MA	occipital b.	N area	SE-NW	eg	G
218	R.	-15.3	12.6	3.5	25.0	8.2	4.8	F	YA	fibula	N area	SE-NW	eg	G; H
219	R.	-17.1	11.5	3.4	41.8	14.5	3.8	F	YA	ulna	N area	SW-NE	sg	G
220	R.	-16.6	12.0	3.3	42.1	15.0	16.3	M	*	radius	N area	SW-NE	tg	S
221	R.	-15.0	12.3	3.6	24.2	7.6	5.1	M	YA	fibula	N area	S-N	eg	
222	R.	-15.2	12.4	3.5	22.8	7.6	4.6	F	MA	fibula	N area	S-N	eg	
223	R.	-15.6	13.6	3.6	18.9	6.1	7.4	F	YA	fibula	N area	S-N	eg	
224	R.	-13.2	14.4	3.4	33.6	11.6	9.2	F	YA	fibula	N area	S-N	sg	
225	R.	-17.1	12.2	3.5	19.0	6.3	4.3	M	MA	ulna	N area	S-N	sg	G
226	R.	-16.7	11.6	3.2	40.5	14.6	19.5	F	MA	radius	N area	S-N	eg	G; H
227	R.	-17.2	11.3	3.2	43.1	15.7	22.8	F	MA	fibula	N area	S-N	sg	G; H
228	R.	-16.3	12.4	3.4	23.8	8.2	8.1	M	MA	radius	N area	SE-NW	sg	
229	R.	-16.1	12.7	3.3	41.1	14.6	12.2	F	MA	fibula	N area	S-N	eg	
230	R.	-17.2	11.0	3.3	34.8	12.5	12.9	M	OA	ulna	N area	S-N	eg	G; H
231	R.	-18.2	10.9	3.6	31.2	10.1	4.8	M	YA	fibula	N area	*	*	
232	R.	-18.0	12.3	3.4	32.8	11.2	5.6	M	MA	fibula	N area	*	*	
233	R.	-16.6	11.2	3.4	24.7	8.4	4.3	M	YA	radius	N area	*	*	
234	R.	-18.1	13.1	3.6	20.8	6.8	3.7	M	YA	radius	N area	*	*	
235	R.	-17.9	11.4	3.3	31.3	11.0	5.8	F	YA	humerus	N area	*	*	
236	R.	-16.9	11.9	3.5	40.6	13.7	2.7	F	*	humerus	N area	*	*	
237	R.	-16.6	11.1	3.3	37.1	13.2	7.2	*	SA	ulna	N area	*	*	
238	R.	-17.2	12.0	3.3	32.5	11.5	7.7	*	SA	tibia	N area	S-N	eg	G
239	R.	-15.6	10.5	3.2	44.7	16.4	22.7	*	SA	humerus	N area	*	*	
241	R.	-17.4	12.6	3.3	42.0	15.1	19.9	M	YA	rib	N area	*	*	
243	R.	-17.4	11.1	3.3	41.7	14.9	6.9	*	SA	phalange	N area	*	*	
247	R.	-17.0	11.8	3.4	39.6	13.8	5.0	F	MA	coxal b.	S area	*	*	
254	R.	-17.7	11.0	3.4	28.8	10.0	6.7	M	YA	rib	N area	*	*	

Sample number	Period	$\delta^{13}\text{C}$	$\delta^{15}\text{N}$	C/N	%C	%N	%Coll	Sex	Age group	Sampled bone	Site area	Orientation	Grave typology	Notes
240	Post-Roman	-12.8	12.6	3.3	41.6	14.6	7.4	M	YA	rib	S area	*	*	
242	P-R	-13.8	12.3	3.3	40.5	14.4	4.5	M	YA	rib	S area	*	eg	
244	P-R	-14.4	12.4	3.3	35.7	12.5	7.1	F	YA	rib	S area	W-E	sts	
246	P-R	-13.1	13.0	3.2	41.2	14.8	14.9	M	MA	rib	S area	W-E	sts	
248	P-R	-15.1	12.0	3.3	36.1	12.6	8.6	F	OA	rib	S area	*	*	
249	P-R	-14.8	12.7	3.4	26.1	8.8	4.3	M	OA	occipital b.	S area	W-E	sts	
250	P-R	-13.7	12.6	3.4	35.7	12.2	8.5	F	MA	rib	S area	*	*	
251	P-R	-13.3	12.7	3.4	29.1	10.1	6.0	M	MA	rib	S area	W-E	sts	
252	P-R	-14.4	13.0	3.3	32.7	11.6	6.8	F	YA	rib	S area	N-S	eg	
253	P-R	-13.6	12.7	3.3	35.2	12.6	8.6	F	YA	scapula	S area	*	*	
255	P-R	-15.1	13.8	3.4	39.6	13.6	6.3	F	MA	occipital b.	S area	W-E	sts	
256	P-R	-13.7	13.4	3.3	40.5	14.2	6.4	F	MA	rib	S area	W-E	sts	
257	P-R	-14.7	13.0	3.4	27.6	9.5	5.7	M	YA	occipital b.	S area	W-E	sts	
258	P-R	-13.7	13.3	3.3	31.7	11.1	6.3	*	SA	rib	S area	W-E	sts	
259	P-R	-14.2	12.3	3.3	36.4	12.8	10.7	*	SA	scapula	S area	*	*	
245	Medieval	-16.5	12.9	3.6	40.3	13.1	1.6	M	YA	tibia	W area	*	eg	

Anthropological and archaeological data was extracted from López Costas thesis (2012).

* information on burial type is lost or cannot be determined.

Key: M=male; F=female. SA Subadult (13-20 years), there were no individuals younger than 12 years old; YA=young adult (20-35 years); MA=middle adult (35-50 years); OA=old adult (50+ years); eg=earth grave; sg=stones grave; sts=stone slabs tomb; tg=tegulae grave; G=with grave goods; H=hobnails.

TABLE 4. Statistical summary of the $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ results for Roman, Post-Roman and Medieval periods.

	Roman period	Post-Roman Period	Medieval
n	43	15	1
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\text{‰}$)	-16.7 \pm 1.0	-14.3 \pm 0.7	-16.5
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\text{‰}$)	12.1 \pm 0.9	12.8 \pm 0.5	12.9

TABLE 5. Statistical summary of the human $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ according to grave typology

	Tegulae graves	Stone graves	Earth graves	Stone slabs tombs
n	9	5	18	8
$\bar{X} \pm \text{SD}$ ($\delta^{13}\text{C}\%$) (min, Max)	-16.8 \pm 0.6 (-17.5,-15.8)	-16.2 \pm 1.7 (-17.2,-13.2)	-16.0 \pm 1.0 (-17.2,-13.8)	-14.1 \pm 0.8 (-15.1,-13.1)
$\bar{X} \pm \text{SD}$ ($\delta^{15}\text{N}\%$) (min, Max)	12.5 \pm 0.8 (11.6, 14.3)	12.3 \pm 1.2 (11.4, 14.4)	12.4 \pm 0.8 (11.0, 13.9)	13.0 \pm 0.5 (12.4, 13.8)

TABLE 6. Statistical summary and inter-groups comparison analysis of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ arranged by estimated sex in Roman and Post-Roman phases.

	Male	Female	M-W-test
A Lanzada Roman			
n	22	17	
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\text{‰}$)	-16.9 \pm 0.9	-16.4 \pm 1.2	U 230.0 p=0.22
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\text{‰}$)	12.1 \pm 0.8	12.3 \pm 1.0	U 208.5 p=0.54
A LanzadaPost-Roman			
n	6	7	
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\text{‰}$)	-13.8 \pm 0.8	-14.3 \pm 0.6	U 13.0 p=0.29
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\text{‰}$)	12.7 \pm 0.3	12.8 \pm 0.6	U 22.5 p=0.83

TABLE 7. Statistical summary and inter-groups comparison analysis of $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ arranged by estimated age at death in Roman and Post-Roman phases.

	13-20 years	20-35 years	35-50 years	>50 years	K-W test
A Lanzada Roman					
n	4	16	13	3	
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\%$)	-16.7 \pm 0.8	-16.4 \pm 1.4	-16.7 \pm 0.9	-16.7 \pm 0.5	$H_{(3)}=0.08$, $p=0.99$
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\%$)	11.2 \pm 0.6	12.4 \pm 1.1	12.2 \pm 0.8	11.6 \pm 0.7	$H_{(3)}=7.06$, $p=0.07$
A Lanzada Post-Roman					
n	2	6	5	2	
$\bar{X} \pm SD$ ($\delta^{13}\text{C}\%$)	-14.2	-13.9 \pm 0.7	-13.8 \pm 0.8	-14.8 \pm 0.2	$H_{(3)}=4.13$, $p=0.25$
$\bar{X} \pm SD$ ($\delta^{15}\text{N}\%$)	12.8	12.7 \pm 0.3	13.1 \pm 0.5	12.3 \pm 0.5	$H_{(3)}=3.41$, $p=0.33$