

Predicting bee community responses to land-use changes: effects of geographic and taxonomic biases

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open access

De Palma, Adriana, Abrahamczyk, Stefan, Aizen, Marcelo A., Albrecht, Matthias, Basset, Yves, Bates, Adam, Blake, Robin J., Boutin, Céline, Bugter, Rob, Connop, Stuart, Cruz-López, Leopoldo, Cunningham, Saul A., Darvill, Ben, Diekötter, Tim, Dorn, Silvia, Downing, Nicola, Entling, Martin H., Farwig, Nina, Felicioli, Antonio, Fonte, Steven J., Fowler, Robert, Franzén, Markus, Goulson, Dave, Grass, Ingo, Hanley, Mick E., Hendrix, Stephen D., Herrmann, Farina, Herzog, Felix, Holzschuh, Andrea, Jauker, Birgit, Kessler, Michael, Knight, M. E., Kruess, Andreas, Lavelle, Patrick, Le Féon, Violette, Lentini, Pia, Malone, Louise A., Marshall, Jon, Pachón, Eliana Martínez, McFrederick, Quinn S., Morales, Carolina L., Mudri-Stojnic, Sonja, Nates-Parra, Guiomar, Nilsson, Sven G., Öckinger, Erik, Osgathorpe, Lynne, Parra-H, Alejandro, Peres, Carlos A., Persson, Anna S., Petanidou, Theodora, Poveda, Katja, Power, Eileen F., Quaranta, Marino, Quintero, Carolina, Rader, Romina, Richards, Miriam H., Roulston, T'ai, Rousseau, Laurent, Sadler, Jonathan P., Samnegård, Ulrika, Schellhorn, Nancy A., Schüepp, Christof, Schweiger, Oliver, Smith-Pardo, Allan H., Steffan-Dewenter, Ingolf, Stout, Jane C., Tonietto, Rebecca K., Tscharntke, Teja, Tylianakis, Jason M., Verboven, Hans A. F., Vergara, Carlos H., Verhulst, Jort, Westphal,

Catrin, Yoon, Hyung Joo and Purvis, Andy (2016) Predicting bee community responses to land-use changes: effects of geographic and taxonomic biases. *Scientific Reports*, 6. 31153. ISSN 2045-2322 doi: <https://doi.org/10.1038/srep31153> Available at <https://centaur.reading.ac.uk/66769/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/srep31153>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

SCIENTIFIC REPORTS

OPEN

Predicting bee community responses to land-use changes: Effects of geographic and taxonomic biases

Received: 29 February 2016

Accepted: 13 July 2016

Published: 11 August 2016

Adriana De Palma^{1,2}, Stefan Abrahamczyk³, Marcelo A. Aizen⁴, Matthias Albrecht⁵, Yves Basset⁶, Adam Bates⁷, Robin J. Blake⁸, Céline Boutin⁹, Rob Bugter¹⁰, Stuart Connop¹¹, Leopoldo Cruz-López¹², Saul A. Cunningham¹³, Ben Darvill¹⁴, Tim Diekötter^{15,16,17}, Silvia Dorn¹⁸, Nicola Downing¹⁹, Martin H. Entling²⁰, Nina Farwig²¹, Antonio Felicioli²², Steven J. Fonte²³, Robert Fowler²⁴, Markus Franzén²⁵, Dave Goulson²⁴, Ingo Grass²⁶, Mick E. Hanley²⁷, Stephen D. Hendrix²⁸, Farina Herrmann²⁶, Felix Herzog²⁹, Andrea Holzschuh³⁰, Birgit Jauker³¹, Michael Kessler³², M. E. Knight²⁷, Andreas Kruess³³, Patrick Lavelle^{34,35}, Violette Le Féon³⁶, Pia Lentini³⁷, Louise A. Malone³⁸, Jon Marshall³⁹, Eliana Martínez Pachón⁴⁰, Quinn S. McFrederick⁴¹, Carolina L. Morales⁴, Sonja Mudri-Stojnic⁴², Guimar Nates-Parra⁴⁰, Sven G. Nilsson⁴³, Erik Öckinger⁴⁴, Lynne Osgathorpe⁴⁵, Alejandro Parra-H^{46,47}, Carlos A. Peres⁴⁸, Anna S. Persson⁴³, Theodora Petanidou⁴⁹, Katja Poveda⁵⁰, Eileen F. Power⁵¹, Marino Quaranta⁵², Carolina Quintero⁴, Romina Rader⁵³, Miriam H. Richards⁵⁴, T'ai Roulston^{55,56}, Laurent Rousseau⁵⁷, Jonathan P. Sadler⁵⁸, Ulrika Samnegård⁵⁹, Nancy A. Schellhorn⁶⁰, Christof Schüepp⁶¹, Oliver Schweiger²⁵, Allan H. Smith-Pardo^{62,63}, Ingolf Steffan-Dewenter³⁰, Jane C. Stout⁵¹, Rebecca K. Tonietto^{64,65,66}, Teja Tschamntke²⁶, Jason M. Tylianakis^{1,67}, Hans A. F. Verboven⁶⁸, Carlos H. Vergara⁶⁹, Jort Verhulst⁷⁰, Catrin Westphal²⁶, Hyung Joo Yoon⁷¹ & Andy Purvis^{1,2}

Land-use change and intensification threaten bee populations worldwide, imperilling pollination services. Global models are needed to better characterise, project, and mitigate bees' responses to these human impacts. The available data are, however, geographically and taxonomically unrepresentative; most data are from North America and Western Europe, overrepresenting bumblebees and raising concerns that model results may not be generalizable to other regions and taxa. To assess whether the geographic and taxonomic biases of data could undermine effectiveness of models for conservation policy, we have collated from the published literature a global dataset of bee diversity at sites facing land-use change and intensification, and assess whether bee responses to these pressures vary across 11 regions (Western, Northern, Eastern and Southern Europe; North, Central and South America; Australia and New Zealand; South East Asia; Middle and Southern Africa) and between bumblebees and other bees. Our analyses highlight strong regionally-based responses of total abundance, species richness and Simpson's diversity to land use, caused by variation in the sensitivity of species and potentially in the nature of threats. These results suggest that global extrapolation of models based on geographically and taxonomically restricted data may underestimate the true uncertainty, increasing the risk of ecological surprises.

¹Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Rd, Ascot, Berkshire SL5 7PY, UK. ²Department of Life Sciences, Natural History Museum, Cromwell Road, London SW7 5BD, UK. ³Nees Institute for Plant Biodiversity, University of Bonn, Meckenheimer Allee 170, 53115 Bonn, Germany. ⁴Laboratorio Ecotono, INIBIOMA (CONICET - Universidad Nacional del Comahue), Quintral 1250, 8400 Bariloche, Río Negro, Argentina. ⁵Institute for Sustainability Sciences, Agroscope, Reckenholzstrasse 191, 8046 Zurich, Switzerland. ⁶Smithsonian Tropical Research Institute, Apartado 0843-03092, Balboa, Ancon, Panama City, Republic of Panama.

⁷Biosciences, Nottingham Trent University, Nottingham, NG11 8NS, UK. ⁸Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University of Reading, Earley Gate, Reading, RG6 6AR, UK. ⁹Science & Technology Branch, Environment and Climate Change Canada, 1125 Colonel By Drive, Carleton University, Ottawa, Ontario K1A 0H3, Canada. ¹⁰Alterra, Part of Wageningen University and Research, P.O. Box 47, 6700 AA Wageningen, Netherlands. ¹¹Sustainability Research Institute, University of East London, 4-6 University Way, Docklands, London E16 2RD, UK. ¹²Grupo de Ecología y Manejo de Artrópodos, El Colegio de la Frontera Sur (ECOSUR), Carretera Antigua Aeropuerto km 2.5. Tapachula, 30700 Chiapas, Mexico. ¹³CSIRO Land and Water, Canberra, ACT 2601, Australia. ¹⁴British Trust for Ornithology (Scotland), Biological and Environmental Sciences, University of Stirling, FK9 4LA, UK. ¹⁵Department of Landscape Ecology, Institute for Natural Resource Conservation, Kiel University, Olshausenstrasse 75, 24118 Kiel, Germany. ¹⁶Department of Biology, Nature Conservation, University Marburg, Marburg, Germany. ¹⁷Institute of Integrative Biology, ETH Zurich, Switzerland. ¹⁸Applied Entomology, ETH Zurich, Schmelzbergstr. 7/LFO, 8092 Zurich, Switzerland. ¹⁹RSPB, Scottish Headquarters 2 Lochside View, Edinburgh Park, Edinburgh, EH12 9DH, UK. ²⁰Institute for Environmental Sciences, University of Koblenz-Landau, Fortstr. 7, 76829 Landau, Germany. ²¹Conservation Ecology, Faculty of Biology, Philipps-Universität Marburg, Karl-von-Frisch-Str. 8, 35032 Marburg, Germany. ²²Dipartimento di Scienze Veterinarie, Viale delle Piagge 2, 56100, Pisa, Università di Pisa, Italia. ²³Department of Soil and Crop Sciences, Colorado State University, Fort Collins, CO 80523, USA. ²⁴School of Life Sciences, University of Sussex, BN19QG, UK. ²⁵Helmholtz Centre for Environmental Research - UFZ, Department of Community Ecology, Theodor-Lieser-Straße 4, 06120 Halle, Germany. ²⁶Agroecology, Department of Crop Sciences, Georg-August-University Göttingen, D-37077 Göttingen, Germany. ²⁷School of Biological Sciences, Plymouth University, Plymouth PL4 8AA, UK. ²⁸Department of Biology, University of Iowa, Iowa, USA. ²⁹Agroscope, Institut for Sustainability Sciences, CH-8046 Zurich, Switzerland. ³⁰Department of Animal Ecology and Tropical Biology, Biocenter, University of Würzburg, Am Hubland, 97074 Würzburg, Germany. ³¹Justus-Liebig University, Department of Animal Ecology, Heinrich-Buff-Ring 26-32, 35392 Giessen, Germany. ³²Institut für Systematische und Evolutionäre Botanik, Switzerland. ³³Dept. for Ecology and Conservation of Fauna and Flora, Federal Agency for Nature Conservation (Bundesamt für Naturschutz, BfN), Konstantinstrasse 110, D-53179 Bonn, Germany. ³⁴Institut de Recherche pour le Développement (IRD), 93143 Bondy Cedex, France. ³⁵Centro Internacional de Agricultura Tropical (CIAT), Tropical Soil Biology and Fertility Program, Latin American and Caribbean Region, Cali, Colombia. ³⁶INRA, UR 406 Abeilles et Environnement, CS 40509, F-84914 Avignon, France. ³⁷School of BioSciences, University of Melbourne, Parkville VIC 3010, Australia. ³⁸New Zealand Institute for Plant and Food Research Ltd, Private Bag 92169, Auckland Mail Centre, Auckland 1142, New Zealand. ³⁹Marshall Agroecology Ltd, 2 Nut Tree Cottages, Barton, Winscombe BS25 1DU, UK. ⁴⁰Departamento de Biología, Facultad de Ciencias, Universidad Nacional de Colombia, Sede Bogotá, Colombia. ⁴¹University of California, Riverside Department of Entomology, 900 University Avenue, Riverside, CA 92521, USA. ⁴²Department of Biology and Ecology, Faculty of Science, University of Novi Sad, 21000 Novi Sad, Serbia. ⁴³Department of Biology, Lund University, SE-223 62 Lund, Sweden. ⁴⁴Swedish University of Agricultural Sciences, Department of Ecology, Box 7044, SE-750 07 Uppsala, Sweden. ⁴⁵RSPB, UK Headquarters The Lodge, Sandy, Bedfordshire, UK. ⁴⁶Laboratorio de Investigaciones en Abejas, LABUN, Departamento de Biología, Facultad de Ciencias, Universidad Nacional de Colombia, Carrera 45 No. 26-85, Edif. Uriel Gutiérrez, Bogotá D.C., Colombia. ⁴⁷Corporación para la Gestión de Servicios Ecosistémicos, Polinización y Abejas - SEPyA, Bogotá D.C., Colombia. ⁴⁸School of Environmental Sciences, University of East Anglia, Norwich NR47TJ, UK. ⁴⁹Laboratory of Biogeography & Ecology, Department of Geography, University of the Aegean, 81100 Mytilene, Greece. ⁵⁰Entomology Department, Cornell University, Ithaca, NY 14850, USA. ⁵¹Botany, School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland. ⁵²CREA-ABP, Consiglio per la ricerca in agricoltura e l'analisi dell'economia agraria, Centro di ricerca per l'agrobiologia e la pedologia, Via di Lanciola 12/A, I-50125 - Cascine del Riccio, Firenze, Italy. ⁵³School of Environmental and Rural Science, University of New England, Armidale, New South Wales, Australia. ⁵⁴Department of Biological Sciences, Brock University, St. Catharines, Ontario, L2S 3A1, Canada. ⁵⁵Department of Environmental Sciences, University of Virginia, Charlottesville, Virginia 22904-4123, USA. ⁵⁶Blandy Experimental Farm, 400 Blandy Farm Lane, Boyce, Virginia 22620, USA. ⁵⁷Département des Sciences Biologiques, Université du Québec à Montréal, C.P. 8888, succursale Centre-ville, Montréal, Québec H3C 3P8, Canada. ⁵⁸GEES (School of Geography, Earth and Environmental Sciences, University of Birmingham, Birmingham B15 2TT, UK. ⁵⁹Department of Ecology, Environment and Plant Sciences, Stockholm University, SE-106 91 Stockholm, Sweden. ⁶⁰CSIRO, Dutton Park, QLD 4102, Australia. ⁶¹University of Bern, Institute of Ecology and Evolution, Community Ecology, Baltzerstrasse 6, 3012 Bern, Switzerland. ⁶²Animal and Plant Health Inspection Service, Plant Protection and Quarantine, United States Department of Agriculture (USDA), South San Francisco, CA 94080, USA. ⁶³Faculty of Sciences, National University of Colombia, Medellín (UNALMED), Colombia. ⁶⁴Plant Biology and Conservation, Northwestern University, 2205 Tech Drive, O.T. Hogan Hall Rm 2-1444, Evanston, IL 60208, USA. ⁶⁵Chicago Botanic Garden, 1000 Lake Cook Rd, Glencoe, IL 60011, USA. ⁶⁶Department of Biology, Saint Louis University, 3507 Laclede Avenue, Macelwane Hall, St. Louis, MO 63103-2010, USA. ⁶⁷Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand. ⁶⁸Division Forest, Nature, and Landscape, Department of Earth & Environmental Sciences, KU Leuven, Celestijnenlaan 200E, B-3001 Leuven, Belgium. ⁶⁹Departamento de Ciencias Químico-Biológicas, Universidad de las Américas Puebla, Mexico. ⁷⁰Spotvogellaan 68, 2566 PN, Den Haag, The Netherlands. ⁷¹Department of Agricultural Biology, National Institute of Agricultural Science, RDA, Wanju-gun, Jellabuk-do, 55365, Korea. Correspondence and requests for materials should be addressed to A.D.P. (email: adrid@nhm.ac.uk)

Bees are one of the most important groups of pollinators of economic crops^{1–3}, with both larvae and adults relying on floral products such as pollen and nectar³. Human impacts can reduce the diversity of pollinator assemblages^{4,5} and therefore can impact pollination efficiency and provision. This is a particular concern in agricultural settings, as over 35% of the volume of human food crops produced globally depend upon animal pollination to some extent⁶. Pollinator shortages can lead to reduced crop quality and yield^{7,8}, with potentially large economic

impacts⁹. There has therefore been much research into responses of bee communities to human impacts such as land-use change and intensification.

A number of syntheses have attempted to identify general trends in the response of bees to human impacts^{5,10}. However, their datasets have often been geographically limited, with the majority of data arising from North America and Western Europe¹¹. The geographic patterns of bee decline and diversity are not understood sufficiently well to ensure that such generalisations are valid^{11,12}. If species' responses to disturbance vary among regions, geographically-restricted models will be inadequate to support broad conclusions. The consequences of basing management strategies on extrapolations from such models could be severe, as many under-studied regions have a high economic dependency upon animal-pollinated crops^{11,13} and may generally have limited governmental capacity to adapt to environmental changes¹⁴.

Geographic variation in bee community responses could arise because differences in land-use history and practices mean that the threats facing assemblages differ across regions. Species subject to very recent disturbance may be more vulnerable, whereas extinction filters^{15–17} may have already removed many susceptible species from landscapes where the intensification of farming started already decades ago, such as in temperate European agricultural landscapes. Extinction debt may make matters worse still, if the full impact of land-use changes is not yet evident^{18,19}. In addition, differences in landscape context across regions can influence species' responses. For instance, Winfree *et al.*⁵ found that habitat loss and fragmentation significantly affected bee communities, but only in areas where little natural habitat still remained.

Bee community responses may also vary regionally because community composition varies geographically. Taxa can differ in their intrinsic susceptibility to land-use change and intensification, through having different functional response traits^{20–22}, the distribution of which within a community can affect resilience to pressures²³. A geographic bias towards North America and Western Europe has also resulted in a taxonomic bias; for instance, bumblebees (Apidae: *Bombus*) are particularly diverse in these areas, whereas large areas of the world have no native bumblebee species (e.g., most of Africa and Australasia). In addition, bumblebees are large, often abundant species with long flight seasons and relatively slow flight, making them fairly easy to sample and, in many cases, to identify. Bumblebees may be more or less sensitive than other bees due to their ecological traits and habitat requirements²⁴, which have been shown to influence responses to human impacts and vulnerability to decline^{25,26}. In addition, bumblebees have shown clearer declines than other bees in North America²⁵ and some European countries²⁷, so they may be atypical of broader bee diversity.

We compiled a global dataset of bee diversity from published sources of bee assemblages in sites differing in pressures such as land use, and used this to explore whether models of responses to human impacts are robust against geographic and taxonomic biases. Specifically, we hypothesized that bee responses to land-use pressures should vary significantly with region and with taxonomic group (i.e., bumblebees or other bees) and so models and projections will not be transferable across regions and taxa. Improved understanding in this area will help to clarify whether knowledge based on a few regions and taxa is sufficient to underpin policy decisions as well as highlight systems for future study.

Methods

Data Collation. Data were sought from the literature where bee species abundance and/or occurrence were reported for multiple sites. Suitable papers were identified by searching Web of Science at various times from 2011 to 2015, as well as searching journal alerts and assessing references cited in reviews. Papers were further considered if more than one site was sampled for bee diversity using the same sampling method in the same season and geographic coordinates of each site were available. Papers were prioritised if their data were collected from February 2000 onwards, so that biodiversity data could be matched with remote-sensed data from NASA's Moderate Resolution Imaging Spectroradiometer (MODIS). Data were supplemented with sources found through the PREDICTS project (www.predicts.org.uk), which aims to develop global statistical models of how local biodiversity responds to human impacts²⁸. The database presented here is not a comprehensive compilation of published sources on occurrence and abundance of bee species across sites differing in land use or intensity, because of regional differences in the ability to retrieve information about potential sources and because most researchers we contacted did not make their data available. The dataset will, however, still be useful for researchers wishing to study land-use impacts on this important taxonomic group.

Where possible we extracted site-level records of bee species (Hymenoptera: Apoidea) occurrence and abundance from suitable papers, along with data for other taxonomic groups if available. Raw data were usually not included within the papers or supplementary files, so the papers' corresponding authors were asked for these data. Relevant data were available from 69 papers, hereafter referred to as 'sources' (Table 1). Each source contains one or more studies, where a study is defined as the set of samples within the same country that were taken using the same methodology. By defining studies in this way, we reduce the impact of broad-scale biogeographic differences in diversity and avoid the confounding effects of methodological differences: within, but not between, studies, diversity data can be compared among sites in a straightforward fashion. Differences in sampling effort within a study were corrected for when necessary by dividing abundance by the sampling effort unit. This assumes a linear relationship between abundance and sampling effort; generalised additive models suggested that this assumption was appropriate (gamm4 package²⁹, see Supplementary Data S1 for details). Within each study, we recorded any blocked or split-plot design. The major land-use class and use intensity at each site were assessed based on information in the associated paper, using the scheme described in Hudson *et al.*²⁸ (reproduced in Supplementary Table S1). Briefly, land use was classified as primary vegetation (native vegetation not known to have ever been completely destroyed), secondary vegetation (where the primary vegetation has been completely destroyed; this can include naturally recovering, actively restored, or semi-natural sites), cropland (planted with herbaceous crops), plantation forest (planted with crop trees or shrubs), pasture (regularly or permanently grazed by livestock) or urban (areas with human habitation, where vegetation is predominantly managed for civic or

Reference	Country	Sampling years	Studies	Within-study sites	Bee taxa (% binomial)	Other taxa	mMLE
	Afrotropic		3	39	77	2304	
Basset <i>et al.</i> ^{67 +†}	Gabon	2001–2002	1	12	51 (19.61%)	1806	70
Gaigher & Samways ^{68 +†}	South Africa	2006	1	10	6 (0%)	383	nr
Grass <i>et al.</i> ^{69 +†‡}	South Africa	2011	1	17	21 (9.52%)	115	100
	Australasia		8	200	135	497	
Blanche <i>et al.</i> ^{70 +†}	Australia	2005	2	11	8 (89.36%)	17	nr
Cunningham <i>et al.</i> ^{71 +†}	Australia	2007–2008	1	24	69 (100%)	0	nr
Lentini <i>et al.</i> ^{72 +†}	Australia	2009–2010	1	104	36 (100%)	0	nr
Kessler <i>et al.</i> ^{73 +†}	Indonesia	2004–2005	1	15	9 (0%)	24	nr
Malone <i>et al.</i> ^{74 †‡}	New Zealand	2006–2007	1	2	9 (100%)	0	nr
Todd <i>et al.</i> ^{75 +†}	New Zealand	2007–2008	1	20	9 (100%)	442	27.3
Rader <i>et al.</i> ^{21 +†}	New Zealand	2008–2009	1	24	5 (100%)	20	nr
	Indo-Malay		4	16	1	0	
Liow <i>et al.</i> ^{76 +†‡}	Singapore, Malaysia	1999	4	16	1 (0%)	0	3000
	Nearctic		16	399	242	117	
Boutin <i>et al.</i> ^{77 +†}	Canada	2000	3	60	3 (0%)	116	nr
Richards <i>et al.</i> ^{78 +†}	Canada	2003	3	18	127 (95.04%)	0	nr
Hatfield & LeBuhn ^{79 †}	United States	2002–2003	1	120	13 (100%)	0	nr
McFrederick & LeBuhn ^{80 †‡}	United States	2003–2004	2	40	5 (100%)	0	nr
Shuler <i>et al.</i> ^{81 +†}	United States	2003	1	25	5 (60%)	0	nr
Winfree <i>et al.</i> ^{82 +†}	United States	2003	2	80	1 (0%)	0	nr
Kwaiser & Hendrix ^{83 +}	United States	2004	2	18	53 (97.22%)	1	nr
Julier & Roulston ^{84 +†}	United States	2006	1	20	3 (100%)	0	250
Tonietto <i>et al.</i> ^{85 +†}	United States	2006	1	18	67 (89.55%)	0	nr
	Neotropic		16	286	436	775	
Vázquez & Simberloff ^{86 +}	Argentina	1999, 2001	1	8	25 (52%)	104	nr
Quintero <i>et al.</i> ^{87 †}	Argentina	2000–2001	1	4	14 (35.71%)	38	1280
Schüpp <i>et al.</i> ^{88 +†}	Belize	2009–2010	1	15	43 (100%)	65	nr
Tonhasca <i>et al.</i> ^{89 +†‡}	Brazil	1997, 1999	1	9	21 (100%)	0	10
Barlow <i>et al.</i> ^{90 +†}	Brazil	2005	1	3	22 (75%)	0	3500
Smith-Pardo & Gonzalez ^{91 +†}	Colombia	1997	4	48	300 (46.2%)	0	nr
Parra-H & Nates-Parra ^{92 +†}	Colombia	2003	1	26	21 (100%)	0	nr
Poveda <i>et al.</i> ^{93 +†}	Colombia	2006–2007	2	34	4 (0%)	468	23
Tylianakis <i>et al.</i> ^{94 +†}	Ecuador	2003–2004	1	48	16 (0%)	16	71
Vergara & Badano ^{64 +†}	Mexico	2004	1	16	7 (71.43%)	8	nr
Fierro <i>et al.</i> ^{95 †‡}	Mexico	2009–2010	1	3	4 (100%)	0	346.41
Rousseau <i>et al.</i> ^{96 +†}	Nicaragua	2011	1	72	2 (100%)	81	30
	Palaearctic		64	2271	601	788	
Verboven <i>et al.</i> ^{97 †}	Belgium	2009	1	9	6 (66.67%)	0	11.34
Billeter <i>et al.</i> ^{98 +†} , Diekötter <i>et al.</i> ^{99 +†} and Le Féon <i>et al.</i> ^{100 +†}	Belgium, Czech Republic, Estonia, France, Germany, Netherlands, Switzerland	2001–2002	14	873	276 (98.46%)	7	nr
Kruess & Tschantke ^{101 +}	Germany	1996	2	34	17 (100%)	18	nr
Meyer <i>et al.</i> ^{102 +†}	Germany	2000, 2005	2	30	14 (75%)	8	34.51
Diekötter <i>et al.</i> ^{103 †}	Germany	2001	1	124	2 (100%)	0	353.55
Meyer <i>et al.</i> ^{104,105 +†}	Germany	2004	1	32	109 (100%)	75	nr
Herrmann <i>et al.</i> ^{106 †‡}	Germany	2005	2	26	1 (100%)	0	800
Holzschuh <i>et al.</i> ^{107 +}	Germany	2007	2	134	3 (33.33%)	1	100
Weiner <i>et al.</i> ^{108 +}	Germany	2007	1	29	59 (100%)	460	333
Nielsen <i>et al.</i> ^{109 +†‡}	Greece	2004	4	32	1 (0%)	0	nr
Power & Stout ^{110 +†}	Ireland	2009	1	20	9 (88.89%)	24	1200.24
Davis <i>et al.</i> ^{111 †‡}	Ireland, United Kingdom	2005, 2007, 2008, 2009	1	12	1 (100%)	0	nr
Quaranta <i>et al.</i> ^{112 +†}	Italy	2000	1	2	31 (100%)	0	200
Yoon <i>et al.</i> ¹¹³	Korea, Republic of	2000–2012	1	215	6 (100%)	1	nr
Kohler <i>et al.</i> ^{114 +†}	Netherlands	2004–2005	4	19	26 (95.48%)	56	1500
Continued							

Reference	Country	Sampling years	Studies	Within-study sites	Bee taxa (% binomial)	Other taxa	mMLE
Goulson <i>et al.</i> ^{115 †}	Poland	2006	1	32	22 (100%)	0	200
Mudri-Stojnic <i>et al.</i> ^{116 †‡}	Serbia	2011	1	16	55 (100%)	8	nr
Öckinger & Smith ¹¹⁷	Sweden	2004	1	36	11 (100%)	64	800
Franzén & Nilsson ^{118 †}	Sweden	2005	1	16	83 (100%)	43	nr
Samnegård <i>et al.</i> ^{119 †}	Sweden	2009	1	9	31 (100%)	0	90
Oertli <i>et al.</i> ^{120 †}	Switzerland	2001–2002	1	7	237 (100%)	0	2000
Albrecht <i>et al.</i> ^{121 †}	Switzerland	2003–2004	2	202	75 (100%)	0	nr
Farwig <i>et al.</i> ^{122 †}	Switzerland	2008	1	30	1 (0%)	0	nr
Schüepp <i>et al.</i> ^{123 †}	Switzerland	2008	1	30	11 (72.73%)	69	0.2
Darvill <i>et al.</i> ^{124 †}	United Kingdom	2001	1	17	3 (66.67%)	0	100
Marshall <i>et al.</i> ^{125 †}	United Kingdom	2003	2	84	25 (100%)	0	nr
Hanley (2005, unpublished data) [†]	United Kingdom	2004–2005	1	6	11 (100%)	0	1000
Knight <i>et al.</i> ^{126 †‡}	United Kingdom	2004	1	12	1 (100%)	0	3.16
Connop <i>et al.</i> ^{127 †‡}	United Kingdom	2005	1	5	2 (100%)	0	nr
Goulson <i>et al.</i> ^{128 †}	United Kingdom	2007	1	14	2 (100%)	0	200.25
Hanley <i>et al.</i> ^{129 †}	United Kingdom	2007–2010	1	34	6 (100%)	0	200.04
Blake <i>et al.</i> ^{130 †}	United Kingdom	2008–2010	2	6	8 (75%)	2	90
Redpath <i>et al.</i> ^{131 †}	United Kingdom	2008	1	11	7 (85.71%)	0	nr
Bates <i>et al.</i> ^{132 †}	United Kingdom	2009–2010	1	24	58 (100%)	50	56.6
Osgathorpe <i>et al.</i> ^{133 †}	United Kingdom	2009–2010	2	45	11 (90.91%)	1	nr
R. E. Fowler (PhD thesis, 2014) ^{†‡}	United Kingdom	2011–2012	1	36	75 (100%)	0	nr
Hanley (unpublished data, 2011) ^{†‡}	United Kingdom	2011	1	8	23 (82.61%)	110	nr

Table 1. Data sources and sample sizes. mMLE = largest Maximum Linear Extent (in meters) of any site in the source. MLE is the maximum distance between sampling points within a site, e.g. the length of a transect or the distance between pan traps. nr = not reported. Numbers of taxa are the numbers of unique taxa for which diversity measurements are given (so, if diversity measurements are available only for all bees combined, this would count as one taxon). The percentage of bee species with a known binomial name is also given (% binomial). Note that the figures here represent available data as curated by the PREDICTS team; these will not necessarily match figures in the original papers. †Data were used in the presented analysis. ‡Data will be incorporated into the PREDICTS database (which will be made openly available). *Data are available from the referenced paper. For all other datasets, please contact the corresponding author of that paper directly.

personal amenity). Use intensity was classified according to a three point scale: low, medium and high intensity. For instance, high-intensity cropland would be monocultures with many signs of intensification such as large fields with high levels of external inputs, irrigation and mechanisation; medium intensity cropland would only show some, but not all, features of higher intensity cropland; low-intensity would refer to small fields with mixed crops and little to no external inputs, irrigation or mechanisation. In one data source, information on the use intensity was unavailable at the site-level, so information at the landscape level was used.

The dataset contained 111 studies from 69 sources and 3211 within-study sites (Table 1). This amounted to 195,357 species diversity measurements (i.e., bee taxa and other taxa, Table 1), including 107,176 measurements of bee diversity (a single measurement being, for example, the abundance of a given species at a given site; see Supplementary Data S2 for species list).

Analysis. For this analysis, we did not include studies that recorded only particular target species (for instance, studies that were only interested in the abundance of a single species across sites), so that site-level diversity measures would be meaningful. The final dataset for the analysis included 101,524 diversity records from 837 bee species at 2421 sites from across the globe (North America: 239 sites; Central America: 103; South America: 176; Western Europe: 1211; Northern Europe: 325; Eastern Europe: 64; Southern Europe: 50; Middle and Southern Africa: 39; South Eastern Asia: 31; Australia and New Zealand: 183). In this reduced dataset, many combinations of land use and use intensity had too few sites to permit robust modelling. The data were therefore aggregated to give a variable of combined Land Use and Intensity (LUI) with the following levels: primary vegetation, secondary vegetation, low-intensity cropland, medium-intensity cropland, high-intensity cropland, pasture, plantation forest and urban. All LUI levels had at least 170 sites, except for plantation forest and urban areas, which were scarce in the dataset with only 105 and 94 sites respectively. Sites were also classified by region and subregion (according to United Nations classifications), with Middle and Southern Africa combined into a single category to increase the sample size.

For each site, we calculated three measures of bee community diversity as our response variables: total abundance, within-sample species richness and Simpson's diversity. Simpson's diversity was calculated as:

$$D = 1 - \sum_i P_i^2 \quad (1)$$

where P_i is the proportion of individuals belonging to species i . We use Simpson's diversity as it stabilises faster than species richness and other diversity measures as specimens accumulate³⁰.

As total abundance measurements are not necessarily integers (e.g. densities and effort-corrected measures), use of the Poisson error structure was not possible, so total abundance was $\ln + 1$ transformed before modelling to normalise residuals and equalise variance. Total abundance and Simpson's diversity were modelled using Gaussian errors (model-checking showed that these treatments were appropriate). Species richness was modelled with Poisson error distribution and log-link function; there was evidence of significant overdispersion in these models so an observation-level random effect was included to account for this (i.e., a Poisson-lognormal model)³¹.

All analyses were carried out using R 3.1.0³². We constructed models for each response variable, using mixed-effects models (lme4 package³³) to account for non-independence of data due to differences in collectors ('source'), sampling methodologies and biogeographic source pools ('study') and the spatial structure of sites ('block'); the initial random-effects structure was therefore block nested within study within source. The initial fixed-effects structure of models included LUI, subregion and their interaction. Subregion is treated as a fixed rather than random effect as we are interested in testing the effect, rather than simply estimating the variance associated with geographic subregion. We test differences in responses to LUI among subregions rather than assessing how responses vary with the latitude and longitude of sites, as subregions represent political differences in land-use patterns and data availability, as well as to some extent reflecting biogeographical differences in community composition.

The best random-effects structure was assessed using likelihood ratio tests³⁴, with models fit using Restricted Maximum Likelihood for total abundance and Simpson's diversity, and Maximum Likelihood for species richness. We then attempted to simplify the fixed-effects structure using backwards stepwise model simplification and likelihood ratio tests, with models fit using Maximum Likelihood^{34–36}. Significance of terms in the minimum adequate models were assessed using Type II Wald Chi Square Tests³⁷. However, to better appreciate the uncertainty in the models³⁸, if the interaction between LUI and subregion remained in the minimum adequate model, we also constructed the following models: additive model (with LUI and subregion included as additive effects); LUI only (univariate model); and subregion only (univariate). We then compared the explanatory power and predictive error of the interactive model with these simpler alternatives.

Explanatory power was calculated using the MuMIn package in R³⁹, as the marginal and conditional R^2_{GLMM} values: i.e., the variance explained by fixed effects alone and by fixed and random effects combined, respectively⁴⁰. Predictive error was calculated as the Mean Squared Error (MSE) from ten-fold cross validation, where the model was iteratively fit to nine-tenths of the data (training set), and validated on the final tenth (validation set); we did this by randomly assigning sites into ten approximately equal-sized groups⁴¹. As the data are structured, the training data may not be fully independent of the validation data⁴², but any bias in prediction error that this causes will apply equally to all models being compared as the random effect structures are identical. In addition, some combinations of explanatory variables only occur in few studies or sources; splitting the dataset by these higher-level strata would mean that some combinations would rarely appear in the training data, leading to overestimates of predictive error. MSE was decomposed into measures of bias and variance, which give an indication of the accuracy and precision of predictions respectively⁴³ (See Supplementary Methods for details).

The dataset was then subset to include only studies where both bumblebees (Apidae: *Bombus*) and other bees were sampled (bumblebees contributed over 19% of the bee abundance records); this resulted in 1636 sites from 47 studies. We calculated the site-level diversity measures separately for each group and fitted the initial model with a three-way interaction between LUI, subregion and taxonomic group (*Bombus* or otherwise). The initial random structure was as above, but included a site-level random effect to account for multiple samples (bumblebees and other bees) being taken from the same site. As above, we first attempted to simplify the initial model (both in terms of random effects and then fixed effects) and, if the initial three-way interaction remained in the model, compared the explanatory power and predictive error with simpler models, where responses to LUI were permitted to vary with subregion (LUI, subregion and their interaction) or with taxonomic group (LUI, taxonomic group and their interaction).

To further understand heterogeneity in community response to LUI, planned comparisons were performed (multcomp package⁴⁴). Within each subregion (and each taxonomic group, if assessed), we tested for differences between natural vegetation (primary vegetation) and all other land uses; between semi-natural vegetation (secondary vegetation) and all other land uses (except primary); whether low-intensity cropland differed from medium-intensity cropland; and whether medium-intensity cropland differed from high-intensity cropland. To avoid rank-deficiency, LUI and subregion were collapsed into a single factor in these models. Not all comparisons were possible in all subregions. Multiple comparisons were corrected for using the False Discovery Rate method to adjust significance values^{45,46}.

An alpha value of 0.05 was used in all tests for significance. Spatial autocorrelation was assessed in residuals of minimum adequate models using Moran's I, for each study in turn (spdep package^{47,48}). As multiple tests are carried out, we expect 5% of these to be significant by chance so we additionally test whether the proportion of studies showing autocorrelation exceeds this expected proportion (using a one-sided Chi squared test).

Results

For total abundance, Simpson's diversity and species richness, the minimum adequate models were those in which responses to LUI were free to vary among geographic subregions. These models also always had the greatest explanatory power and were always among the models having the lowest predictive error (Fig. 1). Overall,




Figure 1. The predictive error and explanatory power of models that include only the intercept (NULL), LUI alone, subregion alone, additive effects, or interactive effects. LUI = Land Use and Intensity. For explanatory power, solid bars show the marginal R^2_{GLMM} (the variance explained by fixed effects) and the hashed bars show the conditional R^2_{GLMM} (the variance explained by both random and fixed effects). Error bars show the standard error of the mean predictive error across 10 folds of cross validation. Note that the predictive error should only be compared among models assessing the same response variable, as absolute values depend on the measurement scale.




Figure 2. Predicted means of total (logged) abundance of bees for different land-use classes in each subregion, with 95% confidence intervals. Also shown are significant results of multiple comparisons, testing differences between natural (Primary vegetation) and semi-natural land uses (Secondary vegetation) to human-dominated land uses, and differences between low, medium and high intensity cropland (* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$).

however, explanatory power of fixed effects alone was fairly low, with most variation instead being attributed to random effects, which is not surprising given the variation in sampling methodology and effort among studies and sources.

For total abundance, the interaction between LUI and subregion explained a highly significant amount of variation ($\chi^2 = 133.15$, $df = 27$, $p < 0.0001$) (Fig. 2), which resulted in this model having the lowest predictive error compared to simpler models. The interaction between LUI and subregion was also significant for Simpson's diversity ($\chi^2 = 66.48$, $df = 27$, $p < 0.0001$) and species richness ($\chi^2 = 96.41$, $df = 27$, $p < 0.05$), although the predictive error was not much lower than for models based on subregion alone. In all cases, the interactive model had slightly higher bias than some simpler models, but the lowest variance (See Supplementary Table S2).

As Fig. 2 shows, the response of total bee abundance to land use differs significantly among regions. In Western Europe, agricultural land maintained higher bee abundances than secondary vegetation (low-intensity cropland: $z = 8.21$; medium-intensity cropland: $z = 5.33$; high-intensity cropland: $z = 9.19$; all $p < 0.0001$; pasture: $z = 4.18$, $p = 0.012$). Low-intensity cropland also maintained higher diversity than secondary vegetation (Simpson's diversity: $z = 4.22$, $p = 0.017$) and medium-intensity cropland (Simpson's diversity: $z = -5.68$, $p < 0.0001$; Species richness: $z = -4.82$, $p = 0.0015$).

In South America, bees were more sensitive to agricultural land uses: medium-intensity cropland maintained significantly lower Simpson's diversity than secondary vegetation ($z = -5.15$, $p = 0.00029$). Urbanization had differing effects between subregions, with increased species richness ($z = 5.29$, $p = 0.00022$) in Middle and Southern Africa, but no strong effect detected elsewhere.

When the dataset was split by taxon (*Bombus* vs. others), the best models for each response variable according to likelihood ratio tests included significant three way interactions between LUI, subregion and taxon (total abundance: $\chi^2 = 217.9$, $df = 13$, $p < 0.0001$; Simpson's diversity: $\chi^2 = 27.62$, $df = 13$, $p = 0.0102$; species richness: $\chi^2 = 76.08$, $df = 13$, $p < 0.0001$). These models also had the greatest explanatory power and lowest predictive error, compared with simpler models (Fig. 3); for total abundance, the lower predictive error was driven by both lower




Figure 3. The predictive error and explanatory power of models that include three way interactions between LUI, subregion and taxon (*Bombus* or not), and models with two way interactions between LUI and taxa, or LUI and Subregion. LUI = Land Use and Intensity. For explanatory power, solid bars show the marginal R^2_{GLMM} (the variance explained by fixed effects) and the hashed bars show the conditional R^2_{GLMM} (the variance explained by both random and fixed effects). Error bars show the standard error of the mean predictive error across 10 folds of cross validation. Note that the predictive error should only be compared among models assessing the same response variable, as absolute values depend on the measurement scale.

bias and variance, while for Simpson's diversity and species richness, the slightly higher bias was offset by lower variation (See Supplementary Table S2).

Planned comparisons provided more detail into these context- and taxon-specific differences in response to human impacts. In Western Europe, bumblebees and other species respond similarly in terms of total abundance, which was higher in cropland than in secondary vegetation (e.g. in low-intensity cropland, *Bombus*: $z = 6.85$, $p < 0.0001$; other species: $z = 9.33$, $p < 0.0001$). However, bumblebee species richness tended to be higher in low-intensity cropland than secondary vegetation ($z = 4.68$, $p = 0.004$), which was not true for other bees ($z = 1.96$, n.s.). Increasing agricultural intensity also resulted in a decline in species richness, but the response was consistent across taxa: low-intensity cropland maintained higher richness than medium-intensity cropland both in bumblebees ($z = -4.30$, $p = 0.016$) and other bees ($z = -3.75$, $p = 0.042$).

Bumblebees also responded differently from other bee species in North America. *Bombus* species richness was lower in secondary vegetation relative to primary vegetation ($z = -3.93$, $p = 0.027$), but this was not true for other bees ($z = 0.86$, n.s.); similarly, total abundance was slightly lower for bumblebees ($z = 3.64$, $p = 0.064$) but not other species ($z = 0.45$, n.s.). However, other genera in North America appeared to be relatively more sensitive to medium-intensity cropland, with reduced species richness relative to both primary vegetation ($z = -3.90$, $p = 0.027$) and secondary vegetation ($z = -3.33$, $p = 0.017$). Other genera were also more sensitive to urban areas, with reduced species richness in urban sites relative to primary vegetation ($z = -3.99$, $p = 0.027$), while bumblebees showed no significant response ($z = -2.5$, n.s.).

The number of studies showing significant autocorrelation was not significantly higher than the 5% expected by chance (See Supplementary Table S3 for details).

Discussion

Bees are facing declines across the globe as a result of changing and intensifying land use^{4,17,49}. Detailed statistical models that relate bee diversity to drivers of change have the potential to inform mitigation and conservation efforts and to help safeguard food security. However, the transferability of models based on restricted data to other regions and taxa is not guaranteed^{11,50}. If responses to threats are context dependent, extrapolation from well known study systems could carry significant risks for biodiversity and food security. The areas where food production is most highly dependent upon animal pollination are also those for which the fewest data are available^{11,13,51}, due to a lack of infrastructure and funding in many areas of the world¹¹. These same areas are often poorly buffered against the disruption of ecosystem service provision from whatever cause, meaning that effects of any ecological surprises on human well-being could be more severe here than elsewhere. We have shown that bee community responses to land-use change and agricultural intensification can indeed be highly context-dependent, but whether this impacts the transferability of models depends on the facet of diversity that is of interest.

The response of total abundance, Simpson's diversity and species richness of bee communities to land use and intensity (LUI) varied significantly with geographic region, in line with our hypotheses and with previous work in tropical regions⁵². For all response variables, the greatest predictive ability could only be achieved by allowing regional variation in responses to LUI; at the very least, it was necessary to allow regional variation in baseline diversity. This suggests that conclusions based on geographically restricted data cannot reliably be generalized to other regions. Indeed, bee community responses to agricultural intensification varied between regions; only in Western Europe was there an evident decline in diversity with increasing use-intensity of cropland, in line with previous suggestions that agricultural land-use intensity is more important in temperate than in tropical or subtropical systems¹⁰. The most negative impact of agriculture, however, was seen in South America, where Simpson's diversity was significantly lower than in secondary vegetation; this is congruent with a previous meta-analysis by Gibson *et al.*⁵² that focussed on tropical areas. The effects of urbanisation likewise depended on the subregion—with increased abundance in Africa but few effects seen elsewhere—but these inferences were based on relatively few data. More data from more regions are needed to better understand the impact of urbanisation on bee communities and associated ecosystem services⁵³.

It is likely that the geographic variation in responses is in part due to differences in community composition¹¹, as we found that taxonomic biases towards bumblebees, which frequently dominate datasets geographically limited to North America and Western Europe, can mask the responses of other species. In Western Europe, for instance, bumblebees had higher species diversity in low-intensity cropland than in secondary vegetation, while other bee species did not show the same effect. Bumblebees have longer flight distances than many smaller bees, so may be better able to persist in more human-dominated land-uses, where foraging resources tend to be further from nesting sites²², and can benefit from mass-flowering crops such as oilseed rape⁵⁴.

The effect of taxonomic group on responses to LUI also differed between subregions, suggesting that other factors may also affect generalities. For instance, geographic variation in the nature of threats may be important. Although our land-use intensity classification is applied in an equivalent fashion across regions, it remains extremely coarse. For example, high-intensity cropland may be more intensive in Western Europe than in South America, with regards to some pressures (e.g., pesticide load⁵⁵) but potentially not others (e.g. spatial extent of monocultures). Such variation in agricultural intensification among regions (even within the same land-use intensity class) could in part be driving observed regional differences in biodiversity responses. More detailed data on different aspects of land-use intensity, such as pesticide load and fertilizer application rates, as well as data on the landscape structure, would enable a more robust and precise analysis of how responses vary across regions. This limitation still highlights, however, that models mostly underpinned by data from regions with a long and intensive history of cultivation are unlikely to provide meaningful inferences for many other regions of the world.

Variation in response among regions could also be driven by differences in community composition and therefore in the distribution of traits that may confer resistance or resilience to human impacts. Previous work has shown that trait-based models of species distributions are only transferable—even within a subregion—when land cover is similar⁵⁶. Transferability across subregions is likely to be even more difficult: variability in the sensitivity

of bee communities will be influenced by a complex interaction between the trait distribution (and phylogeny) across communities and variation in the threats they face. For instance, in a global analysis, species that reproduce socially were more vulnerable to isolation and pesticide use than solitary species, but were less sensitive to tillage and agricultural intensity than solitary species²⁰; however, bees that reproduce solitarily are more common in temperate areas than the tropics⁵⁷, while the distribution of these pressures also vary regionally⁵⁵. While we only assessed how a single aspect of community composition may influence results (bumblebees vs other bees), further work into phylogenetic patterns of sensitivity may help to disentangle these two mechanisms that may be driving regional variation in responses. Another important extension to our work would be to explore the interaction of multiple threatening processes, as the pressures faced by bee communities can vary regionally⁵⁸. For instance, competition with introduced species and fragmentation are likely to be more important drivers of native bee diversity in the Neotropics than in temperate regions⁵⁹. While it was not the focus of this work, a spatially-explicit analysis of latitudinal gradients in vulnerability to land-use pressures may be an interesting avenue for further research, potentially highlighting other factors of the environment or community structure that could contribute to geographic variation in sensitivity. For example, species richness of bees peaks at approximately 35° latitude, in dry, Mediterranean climates⁶⁰, rather than in the tropics (as is the case for many other groups⁶¹) and this variation in baseline diversity may alter both actual and detected responses to human impacts.

Although our dataset includes over 2000 sites from five continents, it is not a comprehensive compilation of published sources and is still both geographically and taxonomically biased. Africa and Asia in particular are still poorly represented and as a consequence we may still be underestimating the uncertainty in bee responses to land use in these regions. Even biomes that have high bee diversity are underrepresented; for example, only six studies were in the Mediterranean biome although bee species richness tends to peak at this latitude⁶⁰. In addition, the explanatory power of fixed effects was fairly low, as most variation in diversity is explained by methodological differences between studies and sources in most models. Nonetheless, our analysis has important implications for pollinator research and conservation action. We show that results based on geographically and taxonomically restricted datasets may not be transferable to other regions. Responses vary across regions due to a combination of differences in the inherent vulnerability of species and variation in the nature of threats. The provision of pollination services can be influenced by the abundance^{62,63}, species diversity⁶⁴ and species richness^{64–66} of bee communities, although the relative importance of each facet of diversity appears to vary with study system^{62,64}. Therefore, if we are to safeguard pollinators and the services they provide, research effort to enhance the representativeness (if not the amount) of available data will be needed to make context-dependent recommendations and to better understand the state of pollination services worldwide.

References

- Garibaldi, L. A. *et al.* Services from Plant - Pollinator interactions in the Neotropics. In Rapidel, B., DeClerck, F., Le Coq, J. & Beer, J. (eds.) *Ecosystem services from agriculture and agroforestry: measurement and payment*, 119–139 (Earthscan, London, UK, 2011).
- Food and Agriculture Organization of the United Nations. *Pollination of cultivated plants in the Tropics, Issue 118* (Food & Agriculture Org., 1995).
- Kevan, P. G. & Baker, H. G. Insects as flower visitors and pollinators. *Annual Review of Entomology* **28**, 407–453 (1983).
- Potts, S. G. *et al.* Global pollinator declines: Trends, impacts and drivers. *Trends in Ecology & Evolution* **25**, 345–353 (2010).
- Winfree, R., Aguilar, R., Vázquez, D. P., Leubuh, G. & Aizen, M. A. A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology* **90**, 2068–2076 (2009).
- Klein, A.-M. *et al.* Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences* **274**, 303–313 (2007).
- Garratt, M. *et al.* Avoiding a bad apple: Insect pollination enhances fruit quality and economic value. *Agriculture, Ecosystems & Environment* **184**, 34–40 (2014).
- Richards, A. Does low biodiversity resulting from modern agricultural practice affect crop pollination and yield? *Annals of Botany* **88**, 165–172 (2001).
- Kevan, P. G. & Phillips, T. P. The economic impacts of pollinator declines: An approach to assessing the consequences. *Conservation Ecology* **5**, 8 (2001).
- Kennedy, C. M. *et al.* A global quantitative synthesis of local and landscape effects on wild bee pollinators in agroecosystems. *Ecology Letters* **16**, 584–599 (2013).
- Archer, C. R., Pirk, C. W. W., Carvalheiro, L. G. & Nicolson, S. W. Economic and ecological implications of geographic bias in pollinator ecology in the light of pollinator declines. *Oikos* **123**, 401–407 (2014).
- Mayer, C. *et al.* Pollination ecology in the 21st Century: Key questions for future research. *Journal of Pollination Ecology* **3**, 8–23 (2011).
- Gallai, N., Salles, J.-M., Settele, J. & Vaissière, B. E. Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics* **68**, 810–821 (2009).
- Mora, C. *et al.* The projected timing of climate departure from recent variability. *Nature* **502**, 183–187 (2013).
- Fritz, S. A., Bininda-Emonds, O. R. P. & Purvis, A. Geographical variation in predictors of mammalian extinction risk: big is bad, but only in the tropics. *Ecology Letters* **12**, 538–549 (2009).
- Balmford, A. Extinction filters and current resilience: the significance of past selection pressures for conservation biology. *Trends in Ecology & Evolution* **11**, 193–196 (1996).
- Ollerton, J., Erenler, H., Edwards, M. & Crockett, R. Extinctions of aculeate pollinators in Britain and the role of large-scale agricultural changes. *Science* **346**, 1360–1362 (2014).
- Kuussaari, M. *et al.* Extinction debt: a challenge for biodiversity conservation. *Trends in Ecology & Evolution* **24**, 564–571 (2009).
- Fritz, S. *et al.* Europe's other debt crisis caused by the long legacy of future extinctions. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 7342–7347 (2013).
- Williams, N. M. *et al.* Ecological and life-history traits predict bee species responses to environmental disturbances. *Biological Conservation* **143**, 2280–2291 (2010).
- Rader, R., Bartomeus, I., Tylianakis, J. M. & Laliberté, E. The winners and losers of land use intensification: pollinator community disassembly is non-random and alters functional diversity. *Diversity and Distributions* **20**, 908–917 (2014).
- De Palma, A. *et al.* Ecological traits affect the sensitivity of bees to land-use pressures in European agricultural landscapes. *Journal of Applied Ecology* **52**, 1567–1577 (2015).
- Moretti, M., De Bello, F., Roberts, S. P. M. & Potts, S. G. Taxonomical vs. functional responses of bee communities to fire in two contrasting climatic regions. *Journal of Animal Ecology* **78**, 98–108 (2009).

24. Colla, S. R. & Packer, L. Evidence for decline in eastern North American bumblebees (Hymenoptera: Apidae), with special focus on *Bombus affinis* Cresson. *Biodiversity and Conservation* **17**, 1379–1391 (2008).
25. Bartomeus, I. *et al.* Historical changes in northeastern US bee pollinators related to shared ecological traits. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 4656–60 (2013).
26. Williams, P., Colla, S. & Xie, Z. Bumblebee vulnerability: Common correlates of winners and losers across three continents. *Conservation Biology* **23**, 931–940 (2009).
27. Carvalheiro, L. G. *et al.* Species richness declines and biotic homogenisation have slowed down for NW-European pollinators and plants. *Ecology Letters* **16**, 870–878 (2013).
28. Hudson, L. N. *et al.* The PREDICTS database: a global database of how local terrestrial biodiversity responds to human impacts. *Ecology and Evolution* **4**, 4701–4735 (2014).
29. Wood, S. & Scheipl, F. gamm4: Generalized additive mixed models using mgcv and lme4. URL <http://CRAN.R-project.org/package=gamm4>. (2014).
30. Magurran, A. E. *Measuring biological diversity*. (Blackwell Science Ltd, Oxford, UK, 2004).
31. Harrison, X. A. Using observation-level random effects to model overdispersion in count data in ecology and evolution. *PeerJ* **2**, e616 (2014).
32. R Core Team R: *A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/> (2014).
33. Bates, D., Maechler, M., Bolker, B. & Walker, S. lme4: Linear mixed-effects models using Eigen and S4. URL <http://CRAN.R-project.org/package=lme4> (2014).
34. Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A. & Smith, G. M. *Mixed effects models and extensions in ecology with R*. No. iii in Statistics for Biology and Health (Springer New York, New York, NY, 2009).
35. Crawley, M. J. *The R book* (John Wiley & Sons Ltd, Chichester, 2007).
36. Murtaugh, P. A. Performance of several variable-selection methods applied to real ecological data. *Ecology Letters* **12**, 1061–1068 (2009).
37. Fox, J. & Weisberg, S. *An R Companion to Applied Regression*. (Sage, Thousand Oaks, CA, USA, 2011).
38. Whittingham, M. J., Stephens, P. A., Bradbury, R. B. & Freckleton, R. P. Why do we still use stepwise modelling in ecology and behaviour? *Journal of Animal Ecology* **75**, 1182–1189 (2006).
39. Barton, K. MuMIn: Multi-model inference. URL <http://CRAN.R-project.org/package=MuMIn> (2013).
40. Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. *Methods in Ecology and Evolution* **4**, 133–142 (2013).
41. Rodríguez, J. D., Pérez, A. & Lozano, J. A. Sensitivity analysis of kappa-fold cross validation in prediction error estimation. *IEEE transactions on pattern analysis and machine intelligence* **32**, 569–575 (2010).
42. Wang, W. & Gelman, A. Difficulty of selecting among multilevel models using predictive accuracy. *Statistics and Its Interface* **8**, 153–152 (1981).
43. Sheiner, L. B. & Beal, S. L. Some suggestions for measuring predictive performance. *Journal of Pharmacokinetics and Biopharmaceutics* **9**, 503–160 (1981).
44. Hothorn, T., Bretz, F. & Westfall, P. Simultaneous inference in general parametric models. *Biometrical Journal* **50**, 346–363 (2008).
45. Benjamin, Y. & Hochberg, Y. Controlling the False Discovery Rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B (Methodological)* **57**, 289–300 (1995).
46. Pike, N. Using false discovery rates for multiple comparisons in ecology and evolution. *Methods in Ecology and Evolution* **2**, 278–282 (2011).
47. Bivand, R. & Piras, G. Comparing implementations of estimation methods for spatial econometrics. *Journal of Statistical Software* **63**, 1–36 (2015).
48. Bivand, R. S., Hauke, J. & Kossowski, T. Computing the Jacobian in Gaussian spatial autoregressive models: An illustrated comparison of available methods. *Geographical Analysis* **45**, 150–179 (2013).
49. Roulston, T. H. & Goodell, K. The role of resources and risks in regulating wild bee populations. *Annual review of entomology* **56**, 293–312 (2011).
50. Petchey, O. L. *et al.* The ecological forecast horizon, and examples of its uses and determinants. *Ecology Letters* **18**, 597–611 (2015).
51. Lautenbach, S., Seppelt, R., Liebscher, J. & Dormann, C. F. Spatial and temporal trends of global pollination benefit. *PLoS ONE* **7**, e35954 (2012).
52. Gibson, L. *et al.* Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378–381 (2011).
53. Hernandez, J. L., Frankie, G. W. & Thorp, R. W. Ecology of urban bees: A review of current knowledge and directions for future study. *Cities and the Environment* **2**, 1–15 (2009).
54. Westphal, C., Steffan-Dewenter, I. & Tscharnkte, T. Mass flowering oilseed rape improves early colony growth but not sexual reproduction of bumblebees. *Journal of Applied Ecology* **46**, 187–193 (2009).
55. Schreinemachers, P. & Tipraqsa, P. Agricultural pesticides and land use intensification in high, middle and low income countries. *Food Policy* **37**, 616–626 (2012).
56. Powney, G. D., Preston, C. D., Purvis, A., Van Landuyt, W. & Roy, D. B. Can trait-based analyses of changes in species distribution be transferred to new geographic areas? *Global Ecology and Biogeography* **23**, 1009–1018 (2014).
57. Roubik, D. W. *Ecology and Natural History of tropical bees* (Cambridge University Press, Cambridge, 1992).
58. Brown, M. J. F. & Paxton, R. J. The conservation of bees: a global perspective. *Apidologie* **40**, 410–416 (2009).
59. Freitas, B. M. *et al.* Diversity, threats and conservation of native bees in the Neotropics. *Apidologie* **40**, 332–346 (2009).
60. Ollerton, J., Johnson, S. D. & Hingston, A. B. Geographic variation in diversity and specificity of pollination systems. 283–308 In Waser, N. M. & Ollerton, J. (eds.) *Plant-Pollinator Interactions: from Specialization to Generalization*. (University of Chicago Press, Chicago, USA, 2006).
61. Hillebrand, H. On the generality of the latitudinal diversity gradient. *The American Naturalist* **163**, 192–211 (2004).
62. Garibaldi, L. A. *et al.* Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science* **339**, 1608–1611 (2013).
63. Winfree, R., Fox, J. W., Williams, N. M., Reilly, J. R. & Cariveau, D. P. Abundance of common species, not species richness, drives delivery of a real-world ecosystem service. *Ecology Letters* **18**, 626–635 (2015).
64. Vergara, C. H. & Badano, E. I. Pollinator diversity increases fruit production in Mexican coffee plantations: The importance of rustic management systems. *Agriculture, Ecosystems & Environment* **129**, 117–123 (2009).
65. Hoehn, P., Tscharnkte, T., Tyliaakis, J. M. & Steffan-Dewenter, I. Functional group diversity of bee pollinators increases crop yield. *Proceedings of the Royal Society B: Biological Sciences* **275**, 2283–2291 (2008).
66. Rogers, S. R., Tarpy, D. R. & Burrack, H. J. Bee species diversity enhances productivity and stability in a perennial crop. *PLoS ONE* **9**, e97307 (2014).
67. Basset, Y. *et al.* Changes in arthropod assemblages along a wide gradient of disturbance in Gabon. *Conservation Biology* **22**, 1552–1563 (2008).
68. Gaigher, R. & Samways, M. J. Surface-active arthropods in organic vineyards, integrated vineyards and natural habitat in the Cape Floristic Region. *Journal of Insect Conservation* **14**, 595–605 (2010).
69. Grass, I., Berens, D. G., Peter, F. & Farwig, N. Additive effects of exotic plant abundance and land-use intensity on plant-pollinator interactions. *Oecologia* **173**, 913–923 (2013).

70. Blanche, K. R., Ludwig, J. A. & Cunningham, S. A. Proximity to rainforest enhances pollination and fruit set in orchards. *Journal of Applied Ecology* **43**, 1182–1187 (2006).
71. Cunningham, S. A., Schellhorn, N. A., Marcora, A. & Batley, M. Movement and phenology of bees in a subtropical Australian agricultural landscape. *Austral Ecology* **38**, 456–464 (2013).
72. Lentini, P. E., Martin, T. G., Gibbons, P., Fischer, J. & Cunningham, S. A. Supporting wild pollinators in a temperate agricultural landscape: Maintaining mosaics of natural features and production. *Biological Conservation* **149**, 84–92 (2012).
73. Kessler, M. *et al.* Alpha and beta diversity of plants and animals along a tropical land-use gradient. *Ecological Applications* **19**, 2142–2156 (2009).
74. Malone, L. *et al.* Observations on bee species visiting white clover in New Zealand pastures. *Journal of Apicultural Research* **49**, 284–286 (2010).
75. Todd, J. H. *et al.* Invertebrate community richness in New Zealand kiwifruit orchards under organic or integrated pest management. *Agriculture, Ecosystems & Environment* **141**, 32–38 (2011).
76. Liow, L. H., Sodhi, N. S. & Elmqvist, T. Bee diversity along a disturbance gradient in tropical lowland forests of south-east Asia. *Journal of Applied Ecology* **38**, 180–192 (2001).
77. Boutin, C., Martin, P. A. & Baril, A. Arthropod diversity as affected by agricultural management (organic and conventional farming), plant species, and landscape context. *Ecoscience* **16**, 492–501 (2009).
78. Richards, M. *et al.* Bee diversity in naturalizing patches of Carolinian grasslands in southern Ontario, Canada. *The Canadian Entomologist* **143**, 279–299 (2011).
79. Hatfield, R. & LeBuhn, G. Patch and landscape factors shape community assemblage of bumble bees, *Bombus* spp. (Hymenoptera: Apidae), in montane meadows. *Biological Conservation* **139**, 150–158 (2007).
80. McFrederick, Q. S. & LeBuhn, G. Are urban parks refuges for bumble bees *Bombus* spp. (Hymenoptera: Apidae)? *Biological Conservation* **129**, 372–382 (2006).
81. Shuler, R. E., Roulston, T. H. & Farris, G. E. Farming practices influence wild pollinator populations on squash and pumpkin. *Journal of Economic Entomology* **98**, 790–795 (2005).
82. Winfree, R., Griswold, T. & Kremen, C. Effect of human disturbance on bee communities in a forested ecosystem. *Conservation Biology* **21**, 213–223 (2007).
83. Kwaizer, K. S. & Hendrix, S. D. Diversity and abundance of bees (Hymenoptera: Apoidea) in native and ruderal grasslands of agriculturally dominated landscapes. *Agriculture, Ecosystems & Environment* **124**, 200–204 (2008).
84. Julier, H. E. & Roulston, T. H. Wild bee abundance and pollination service in cultivated pumpkins: Farm management, nesting behavior and landscape effects. *Journal of Economic Entomology* **102**, 563–573 (2009).
85. Tonietto, R., Fant, J., Ascher, J., Ellis, K. & Larkin, D. A comparison of bee communities of Chicago green roofs, parks and prairies. *Landscape and Urban Planning* **103**, 102–108 (2011).
86. Vázquez, D. P. & Simberloff, D. Ecological specialization and susceptibility to disturbance: conjectures and refutations. *The American naturalist* **159**, 606–623 (2002).
87. Quintero, C., Morales, C. L. & Aizen, M. A. Effects of anthropogenic habitat disturbance on local pollinator diversity and species turnover across a precipitation gradient. *Biodiversity and Conservation* **19**, 257–274 (2010).
88. Schüepp, C., Rittiner, S. & Entling, M. H. High bee and wasp diversity in a heterogeneous tropical farming system compared to protected forest. *PLoS ONE* **7**, e52109 (2012).
89. Tonhasca, A., Blackmer, J. L. & Albuquerque, G. S. Abundance and diversity of Euglossine bees in the fragmented landscape of the Brazilian Atlantic Forest. *Biotropica* **34**, 416–422 (2002).
90. Barlow, J. *et al.* Quantifying the biodiversity value of tropical primary, secondary, and plantation forests. *Proceedings of the National Academy of Sciences* **104**, 18555–18560 (2007).
91. Smith-Pardo, A. & Gonzalez, V. H. Diversidad de abejas (Hymenoptera: Apoidea) en estados sucesionales del bosque húmedo tropical TT - Bee diversity (Hymenoptera: Apoidea) in a tropical rainforest succession. *Acta Biológica Colombiana* **12**, 43–55 (2007).
92. Parra-H, A. & Nates-Parra, G. Variation of the orchid bees community (Hymenoptera: Apidae) in three altered habitats of the Colombian “llano” piedmont. *Revista de biología tropical* **55**, 931–941 (2007).
93. Poveda, K., Martínez, E., Kersch-Becker, M. F., Bonilla, M. A. & Tschardtke, T. Landscape simplification and altitude affect biodiversity, herbivory and Andean potato yield. *Journal of Applied Ecology* **49**, 513–522 (2012).
94. Tylianakis, J. M., Klein, A.-M. & Tschardtke, T. Spatiotemporal variation in the diversity of Hymenoptera across a tropical habitat gradient. *Ecology* **86**, 3296–3302 (2005).
95. Fierro, M., Cruz-López, L., Sánchez, D., Villanueva-Gutiérrez, R. & Vandame, R. Effect of biotic factors on the spatial distribution of stingless bees (Hymenoptera: Apidae, Meliponini) in fragmented Neotropical habitats. *Neotropical Entomology* **41**, 95–104 (2012).
96. Rousseau, L., Fonte, S. J., Téllez, O., van der Hoek, R. & Lavelle, P. Soil macrofauna as indicators of soil quality and land use impacts in smallholder agroecosystems of western Nicaragua. *Ecological Indicators* **27**, 71–82 (2013).
97. Verboven, H. A. F., Brys, R. & Hermy, M. Sex in the city: Reproductive success of *Digitalis purpurea* in a gradient from urban to rural sites. *Landscape and Urban Planning* **106**, 158–164 (2012).
98. Billeter, R. *et al.* Indicators for biodiversity in agricultural landscapes: a pan-European study. *Journal of Applied Ecology* **45**, 141–150 (2008).
99. Diekötter, T., Billeter, R. & Crist, T. O. Effects of landscape connectivity on the spatial distribution of insect diversity in agricultural mosaic landscapes. *Basic and Applied Ecology* **9**, 298–307 (2008).
100. Le Féon, V. *et al.* Intensification of agriculture, landscape composition and wild bee communities: A large scale study in four European countries. *Agriculture, Ecosystems & Environment* **137**, 143–150 (2010).
101. Kruess, A. & Tschardtke, T. Grazing intensity and the diversity of grasshoppers, butterflies, and trap-nesting bees and wasps. *Conservation Biology* **16**, 1570–1580 (2002).
102. Meyer, B., Gaebele, V. & Steffan-Dewenter, I. D. Patch size and landscape effects on pollinators and seed set of the Horseshoe Vetch, *Hippocrepis comosa*, in an agricultural landscape of Central Europe. *Entomologia Generalis* **30**, 173–185 (2007).
103. Diekötter, T., Walther-Hellwig, K., Conradi, M., Suter, M. & Frankl, R. Effects of landscape elements on the distribution of the rare bumblebee species *Bombus muscorum* in an agricultural landscape. *Biodiversity and Conservation* **15**, 57–68 (2006).
104. Meyer, B., Jauker, F. & Steffan-Dewenter, I. Contrasting resource-dependent responses of hoverfly richness and density to landscape structure. *Basic and Applied Ecology* **10**, 178–186 (2009).
105. Jauker, B., Krauss, J., Jauker, F. & Steffan-Dewenter, I. Linking life history traits to pollinator loss in fragmented calcareous grasslands. *Landscape Ecology* **28**, 107–120 (2013).
106. Herrmann, F., Westphal, C., Moritz, R. F. A. & Steffan-Dewenter, I. Genetic diversity and mass resources promote colony size and forager densities of a social bee (*Bombus pascuorum*) in agricultural landscapes. *Molecular Ecology* **16**, 1167–1178 (2007).
107. Holzschuh, A., Dormann, C. F., Tschardtke, T. & Steffan-Dewenter, I. Expansion of mass-flowering crops leads to transient pollinator dilution and reduced wild plant pollination. *Proceedings of the Royal Society B: Biological Sciences* **278**, 3444–3451 (2011).
108. Weiner, C. N., Werner, M., Linsenmair, K. E. & Blüthgen, N. Land use intensity in grasslands: Changes in biodiversity, species composition and specialisation in flower visitor networks. *Basic and Applied Ecology* **12**, 292–299 (2011).
109. Nielsen, A. *et al.* Assessing bee species richness in two Mediterranean communities: importance of habitat type and sampling techniques. *Ecological Research* **26**, 969–983 (2011).

110. Power, E. F. & Stout, J. C. Organic dairy farming: impacts on insect-flower interaction networks and pollination. *Journal of Applied Ecology* **48**, 561–569 (2011).
111. Davis, E. S., Murray, T. E., Fitzpatrick, Ü., Brown, M. J. F. & Paxton, R. J. Landscape effects on extremely fragmented populations of a rare solitary bee, *Colletes floralis*. *Molecular Ecology* **19**, 4922–4935 (2010).
112. Quaranta, M. *et al.* Wild bees in agroecosystems and semi-natural landscapes. 1997–2000 collection period in Italy. *Bulletin of Insectology* **57**, 11–61 (2004).
113. Yoon, H. J., Lee, K. Y., Kim, M. A. & Park, I. G. Local distribution and floral preferences of founder bumblebee queens in Korea. *Journal of Apiculture* **27**, 169–178 (2012).
114. Kohler, F., Verhulst, J., Van Klink, R. & Kleijn, D. At what spatial scale do high-quality habitats enhance the diversity of forbs and pollinators in intensively farmed landscapes? *Journal of Applied Ecology* **45**, 753–762 (2008).
115. Goulson, D., Lye, G. C. & Darvill, B. Diet breadth, coexistence and rarity in bumblebees. *Biodiversity and Conservation* **17**, 3269–3288 (2008).
116. Mudri-Stojnic, S., Andric, A., Józán, Z. & Vujic, A. Pollinator diversity (Hymenoptera and Diptera) in semi-natural habitats in Serbia during summer. *Archives of Biological Sciences* **64**, 777–786 (2012).
117. Öckinger, E. & Smith, H. G. Semi-natural grasslands as population sources for pollinating insects in agricultural landscapes. *Journal of Applied Ecology* **44**, 50–59 (2007).
118. Franzén, M. & Nilsson, S. G. How can we preserve and restore species richness of pollinating insects on agricultural land? *Ecography* **31**, 698–708 (2008).
119. Samnegård, U., Persson, A. S. & Smith, H. G. Gardens benefit bees and enhance pollination in intensively managed farmland. *Biological Conservation* **144**, 2602–2606 (2011).
120. Oertli, S., Mueller, A. & Dorn, S. Ecological and seasonal patterns in the diversity of a species-rich bee assemblage (Hymenoptera: Apoidea: Apiformes). *European Journal of Entomology* **102**, 53–63 (2005).
121. Albrecht, M. *et al.* Effects of ecological compensation meadows on arthropod diversity in adjacent intensively managed grassland. *Biological Conservation* **143**, 642–649 (2010).
122. Farwig, N. *et al.* Isolation from forest reduces pollination, seed predation and insect scavenging in Swiss farmland. *Landscape Ecology* **24**, 919–927 (2009).
123. Schüepp, C., Herrmann, J. D., Herzog, F. & Schmidt-Entling, M. H. Differential effects of habitat isolation and landscape composition on wasps, bees, and their enemies. *Oecologia* **165**, 713–721 (2011).
124. Darvill, B., Knight, M. E. & Goulson, D. Use of genetic markers to quantify bumblebee foraging range and nest density. *Oikos* **107**, 471–478 (2004).
125. Marshall, E. J. P., West, T. M. & Kleijn, D. Impacts of an agri-environment field margin prescription on the flora and fauna of arable farmland in different landscapes. *Agriculture, Ecosystems & Environment* **113**, 36–44 (2006).
126. Knight, M. E. *et al.* An interspecific comparison of foraging range and nest density of four bumblebee (*Bombus*) species. *Molecular Ecology* **14**, 1811–1820 (2005).
127. Connop, S., Hill, T., Steer, J. & Shaw, P. Microsatellite analysis reveals the spatial dynamics of *Bombus humilis* and *Bombus sylvarum*. *Insect Conservation and Diversity* **4**, 212–221 (2011).
128. Goulson, D. *et al.* Effects of land use at a landscape scale on bumblebee nest density and survival. *Journal of Applied Ecology* **47**, 1207–1215 (2010).
129. Hanley, M. E. *et al.* Increased bumblebee abundance along the margins of a mass flowering crop: evidence for pollinator spill-over. *Oikos* **120**, 1618–1624 (2011).
130. Blake, R. J., Westbury, D. B., Woodcock, B. A., Sutton, P. & Potts, S. G. Enhancing habitat to help the plight of the bumblebee. *Pest Management Science* **67**, 377–379 (2011).
131. Redpath, N., Osgathorpe, L. M., Park, K. & Goulson, D. Crofting and bumblebee conservation: The impact of land management practices on bumblebee populations in northwest Scotland. *Biological Conservation* **143**, 492–500 (2010).
132. Bates, A. J. *et al.* Changing bee and hoverfly pollinator assemblages along an urban-rural gradient. *PLoS ONE* **6**, e23459 (2011).
133. Osgathorpe, L. M., Park, K. & Goulson, D. The use of off-farm habitats by foraging bumblebees in agricultural landscapes: implications for conservation management. *Apidologie* **43**, 113–127 (2012).

Acknowledgements

We are grateful to all members of the PREDICTS team who aided in the collation and curation of data. We are especially grateful to all the many researchers who have made their data available to us. This paper is a contribution from the Imperial College Grand Challenges in Ecosystems and the Environment Initiative. The PREDICTS project is endorsed by the GEO BON. We thank BBSRC (grant BB/F017324/1 to ADP) and NERC (grant NE/J011193/1 and NE/M014533/1 to AP) for support.

Author Contributions

A.D.P. and A.P. designed the study, A.D.P. conducted the analyses and wrote the first draft of the manuscript. S.A., M.A.A., M.A., Y.B., A.B., R.J.B., C.B., R.B., S.C., L.C.-L., S.A.C., B.D., T.D., S.D., N.D., M.H.E., N.F., A.F., S.J.F., R.F., M.F., D.G., I.G., M.E.H., S.D.H., F.H., F.H., A.H., B.J., M.K., M.K., A.K., P.L., V.L.F., P.L., L.A.M., J.M., E.M.P., Q.S.M., C.L.M., S.M., G.N.P., S.G.N., E.Ö., L.O., A.P., C.P., A.S.P., T.P., K.P., E.F.P., M.Q., C.Q., R.R., M.H.R., T.R., L.R., J.P.S., U.S., N.A.S., C.S., O.S., A.H.S., I.S., J.C.S., R.K.T., T.T., J.M.T., H.A.V., C.H.V., J.V., C.W. and H.J.Y. collated ecological data. All authors contributed substantially to revisions of the manuscript.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: De Palma, A. *et al.* Predicting bee community responses to land-use changes: Effects of geographic and taxonomic biases. *Sci. Rep.* **6**, 31153; doi: 10.1038/srep31153 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>