

# *Nonlinear response of mid-latitude weather to the changing Arctic*

Article

Accepted Version

Overland, J. E., Dethloff, K., Francis, J. A., Hall, R. J., Hanna, E., Kim, S.-J., Screen, J. A., Shepherd, T. G. ORCID: <https://orcid.org/0000-0002-6631-9968> and Vihma, T. (2016) Nonlinear response of mid-latitude weather to the changing Arctic. *Nature Climate Change*, 6 (11). pp. 992-999. ISSN 1758-678X doi: 10.1038/nclimate3121 Available at <https://centaur.reading.ac.uk/67299/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/nclimate3121>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# Nonlinear Response of Midlatitude Weather to the Changing Arctic

James E. Overland<sup>1,\*</sup>, Klaus Dethloff<sup>2</sup>, Jennifer A. Francis<sup>3</sup>, Richard J. Hall<sup>4</sup>, Edward Hanna<sup>4</sup>,  
Seong-Joong Kim<sup>5</sup>, James A. Screen<sup>6</sup>, Theodore G. Shepherd<sup>7</sup>, and Timo Vihma<sup>8</sup>

<sup>1</sup>Pacific Marine Environmental Laboratory, NOAA, Seattle, Washington, USA

<sup>2</sup>Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Potsdam, Germany

<sup>3</sup>Department of Marine and Coastal Sciences, Rutgers University, New Jersey, USA

<sup>4</sup>Department of Geography, University of Sheffield, UK

<sup>5</sup>Korea Polar Research Institute, Korea

<sup>6</sup>College of Engineering, Mathematics and Physical Sciences, University of Exeter, UK

<sup>7</sup>Department of Meteorology, University of Reading, UK

<sup>8</sup>Finnish Meteorological Institute, Helsinki, Finland

\*Corresponding author: James Overland

NOAA/PMEL

7600 Sand Point Way NE

Seattle WA 98115

1-206-526-6795

[james.e.overland@noaa.gov](mailto:james.e.overland@noaa.gov)

Final Revision Submitted to Nature Climate Change: Perspectives

22 July 2016

Are continuing changes in the Arctic influencing wind patterns and the occurrence of extreme weather events in northern midlatitudes? The chaotic nature of atmospheric circulation precludes easy answers. Yet the topic is a major science challenge, as continued Arctic temperature increases are an inevitable aspect of anthropogenic global change. We propose a perspective that rejects simple cause-and-effect pathways, notes diagnostic challenges in interpreting atmospheric dynamics, and present a way forward based on understanding multiple processes that lead to uncertainties in Arctic/midlatitude weather and climate linkages. We emphasize community coordination for both scientific progress and communication to a broader public.

Various metrics indicate that the recent period of disproportionate Arctic warming relative to midlatitudes—referred to as Arctic Amplification (AA)—emerged from the noise of natural variability in the late 1990s<sup>1</sup>. This signal will strengthen as human activities continue to raise greenhouse gas concentrations<sup>2</sup>. The assessment of the potential for AA to influence broader hemispheric weather (referred to as linkages) is complex and controversial<sup>3-6</sup>. Yet with intensifying AA, we argue that the key question is not whether the melting Arctic will influence midlatitude weather patterns over the next decades, but rather what is the nature and magnitude of this influence relative to non-Arctic factors, and is it limited to specific regions, seasons, or types of weather events<sup>7</sup>?

Although studies arguing for linkages often highlight a single causal pathway, the complexity of atmospheric dynamics implies that such singular linkage pathways are unlikely. Nonlinearities in the climate system are particularly important in the Arctic and subarctic<sup>8,9,10</sup>. The climate change signal is larger than anywhere else in the Northern Hemisphere and the region possesses multiple

53    feedbacks. Coupling exists between the Arctic troposphere and the wintertime stratospheric polar  
54    vortex, which itself is highly nonlinear. A linkage pathway that may appear to be responsible for  
55    one series of events may not exist in another scenario with similar forcing. This is potentially  
56    reflected in observationally based studies that have struggled to find robust linkages<sup>11,12</sup>. Further,  
57    multiple runs of the same model with similar but slightly different initial conditions, termed  
58    ensemble members, show linkages in some subsets of ensemble runs but not in others<sup>13</sup>. This  
59    failure to detect direct connections is sometimes interpreted as evidence against linkages. Four  
60    properties (limitations) that contribute to the complexity of attribution of linkages are developed  
61    in this Perspective: *itinerancy* [seemingly random variations from state to state], *intermittency*  
62    [apparently different atmospheric responses under conditions of similar external forcing, such  
63    as sea-ice loss], *multiple influences* [simultaneous forcing by various factors, such as sea-  
64    surface temperature anomalies in the tropics, midlatitudes and Arctic], and *state dependence* [a  
65    response dependent on the prior state of the atmospheric circulation, e.g., the phase of the Arctic  
66    Oscillation (AO) atmospheric circulation index or the strength of the stratospheric vortex].

67  
68    We propose a system-level approach that recognizes multiple simultaneous processes, internal  
69    instabilities, and feedbacks. Progress in understanding Arctic/midlatitude linkages will require  
70    the use of probabilistic model forecasts that are based on case studies and high-resolution,  
71    ensemble solutions to the equations of motion and thermodynamics. Community coordinated  
72    model experiments and diagnostic studies of atmospheric dynamics are essential to resolve  
73    controversy and benefit efforts to communicate the impacts of linkages and uncertainties with a  
74    broad public.

**Arctic warming is unequivocal, substantial, and ongoing**

Changes in Arctic climate in the last two decades are substantial. Since 1980 Arctic temperature increases have exceeded those of the Northern Hemisphere average by at least a factor of two<sup>14</sup>. Over land north of 60°N, 12 of the past 15 years have exhibited the highest annual mean surface air temperatures since 1900. AA is also manifested in loss of sea ice, glaciers, snow and permafrost, a longer open-water season, and shifts in Arctic ecosystems. Sea ice has undergone an unprecedented decline over the past three decades with a two-thirds reduction in volume<sup>2</sup>. Comparable decreases in snow cover have occurred during May and June. AA is strongest in fall/winter with largest values over regions of sea ice loss<sup>15</sup>, while the areas of greatest warming in summer are located over high-latitude land where spring snow loss has occurred progressively earlier<sup>16</sup>.

This amplification of warming in the Arctic occurs for several reasons, all based on fundamental physical processes<sup>17,18</sup>. Among these are feedbacks related to albedo owing to a loss of snow and sea ice along with increases in heat-trapping water vapor and clouds. Increasing temperatures in the lower atmosphere elevate the height of mid-level pressure surfaces (geopotential height), leading to changes in poleward and regional gradients and, consequently, wind patterns<sup>19,20,21</sup>.

Based on over 30 climate model simulations presented in the most recent Intergovernmental Panel on Climate Change (IPCC) Assessment Report, future winter (November-March) surface temperatures in the Arctic (60-90°N) are projected to rise ~4°C by 2040, with a standard deviation of 1.6 °C, relative to the end of the previous century (1981-2000)<sup>2</sup>. This is roughly double the projected global increase and will likely be accompanied by sea ice free summers.

Past and near future emissions of anthropogenic CO<sub>2</sub> assure mid-century AA and global warming.

## **Living with an uncertain climate system**

The task of unraveling cause and effect of mechanisms linking changes in the large scale atmospheric circulation to AA is hampered by poor signal detection in a noisy system and complex climate dynamics, regardless of whether the approach is statistical analyses or targeted model simulations. Nonlinear relationships are widespread in the Arctic climate system, in which responses are not directly proportional to the change in forcing<sup>8,10,22</sup>. Further, when discussing anomalous weather or climate conditions, causation can have different meanings. Typically one factor is necessary but several supplementary factors may also be required. This can lead to confusion because only sufficient causes have deterministic predictive power<sup>23,24</sup>. Together these factors make linkage attribution challenging. Many previous data and modeling analyses start with straightforward Arctic changes using, for example, diminished sea ice, and at least implicitly assume quasi-linear, sufficient causal connections<sup>5,7,25-37</sup>. While this approach has been helpful in elucidating relevant linkage mechanisms, we provide a view at the system level that can mask simple cause and effect.

Thermodynamically (i.e., related to temperature gradients) forced wind systems on a rotating planet produce west-to-east flow at midlatitudes. This flow is dynamically unstable, creating north-south meanders that generate high- and low-pressure centers which can produce disruptive weather events. In addition to internal instability, variability in the wind pattern is forced by influences external to the midlatitude atmosphere that may themselves reflect internal variability

on longer timescales, such as sea-surface temperature anomalies in the tropics, midlatitudes, and ice-free parts of the Arctic. Remote forcings (i.e., changes outside the midlatitudes, remote in space and perhaps time) can influence the midlatitude circulation through linear and nonlinear atmospheric patterns, known as teleconnections. Extensive regions of positive temperature anomalies in the Arctic may increase the persistence of weather systems<sup>20,38</sup>. Further, troposphere-stratosphere connections can trigger changes in the regional wind patterns<sup>39</sup>.

Contributors to a lack of simple robust linkages include the four properties discussed as follows:

### *Itinerancy*

Itinerancy refers to the atmosphere spontaneously shifting from state to state based on instabilities in the wind field that can be amplified by internal and external variability. Such states can persist through nonlinear mechanisms<sup>10,22</sup>. Fig. 1(a, b) illustrates two configurations of the northern hemispheric wind pattern (tropospheric polar vortex) occurring at different times: the case shown in Fig. 1a is for a day in November 2013 that had a relatively circular flow pattern around the North Pole, and Fig. 1b shows another day two months later exhibiting a more north-south wavy flow pattern. Although the phrase *polar vortex* is formally reserved for the stratosphere, it is a useful term for discussing tropospheric geopotential height/wind configurations such as those shown in Fig. 1. The jet stream flows from west to east parallel to these geopotential height contours and is strongest where the contours are closest together. Shifts to and from a wavy pattern—known historically as the index cycle—and the varying longitudinal locations of ridges (northward peaks) and troughs (southward excursions) in the geopotential height pattern are part of the seemingly random, internal variability of atmospheric circulation. A wavier jet stream allows cold air from the Arctic to penetrate southward into midlatitudes, and



ridges transport warm air northward. Fig. 1(c, d) are corresponding temperature anomaly patterns for these two days. For the more circular jet stream, cold anomalies are mostly contained within the polar region along with warmer anomalies around midlatitudes (Fig. 1c). This particular pattern is not perfectly symmetric around the North Pole, as the center of the vortex is shifted into the western hemisphere. The wavier jet stream case has two warm and two cold anomaly regions in midlatitudes (Fig. 1d), to the west and east of the region of increased heights (ridges) over Alaska and Scandinavia. Many extreme weather events associated with wavy circulation patterns have occurred in the last decade<sup>40,41</sup>.

Multiple studies<sup>42,43,44</sup> illustrate the paradigm of itinerancy in describing the physical mechanisms driving shifts in atmospheric circulation. Atmospheric circulation can fluctuate between multiple states (referred to as local attractors) in irregular transitions, resulting in chaotic-like behavior on monthly, seasonal, and interannual time scales<sup>42</sup>. Chaos theory argues that the climate system can destabilize and suddenly shift into a new stable state<sup>45,46</sup>. On decadal timescales, increasing variability within a time series is a possible early-warning signal of a critical transition to a different state<sup>47</sup>.

Do observations indicate a recent increase in these types of sudden shifts in the atmospheric circulation? Although one might expect decreased sub-seasonal variability as the temperature contrast across the jet stream declines with AA<sup>48</sup>, recent observations suggest contrary evidence of stable or larger circulation variability and new extremes in several circulation indices. For example, an enhanced magnitude of both positive and negative excursions of the AO circulation index is evident in the last decade during Decembers based on data from 1950-2014<sup>49</sup>. Cohen<sup>50</sup>

notes an increase in midlatitude intraseasonal winter temperature variability from 1988/89 to 2014/15. Periods of relative persistence as well as increases in interannual variability have been noted in other related winter climate indices—such as the North Atlantic Oscillation (NAO), Greenland Blocking Index (GBI), and jet latitude metrics—although stability is more evident at other times of the year<sup>51,52,53</sup>. Observations from the next decade should reveal much about whether increasing variability and weather extremes are ongoing features of climate change or whether circulation-related extremes are damped by AA.

The ability of state-of-the-art climate models to correctly simulate the interplay between thermal and dynamical processes producing itinerancy on different spatial scales is limited. One manifestation of this is the continuing tendency for climate models to underestimate the frequency of blocking (a regional slowing of tropospheric winds)<sup>54</sup>. Also the signal to noise in models could be too weak, as appears to be the case for seasonal forecasts of the NAO<sup>55,56,57</sup>.

### *Intermittency*

*Intermittency* refers to necessary but insufficient causation and suggests an inconsistent response, evident at some times and not at others, or the same response arising from different combinations of Arctic conditions. In other words, the response is not a unique function of the forcing. If responses are intermittent, one will need a longer time series and/or a stronger signal to detect them. Often climate models and correlation analyses of observations produce differing estimates of how the climate will respond to the ongoing AA and loss of sea ice<sup>48,58</sup>. For example, climate model studies have reported shifts towards both the positive or negative phases of the AO and/or NAO, or no apparent shift, in response to AA<sup>13,19,34,39,59</sup>. Analyses that involve averaging over

large areas, long time periods, and/or many ensemble members may not reveal specific atmospheric responses to AA, such as enhanced jet-stream ridges and troughs that occur in specific locations. Despite some clear hypotheses for linkages, it remains difficult to prove that Arctic change has already had or not had an impact on midlatitude weather based on observations alone because of the short period since AA has become apparent<sup>5</sup>.

One approach to overcome the signal-to-noise problem is to use model simulations<sup>59</sup>. Large ensembles of climate simulations have been run with observed sea ice loss as the only forcing factor. In such large ensembles it is possible to answer the question: how many years of simulation are required for the impacts of sea ice loss to become detectable over the noise of internal climate variability? Depending on the metric used to detect changes, the spatial/temporal mean response to forcing often exceeds the length of observational records, suggesting that it may be a decade or more before the forced response to sea ice loss will clearly emerge from the noise of internal variability. Thermodynamic responses may be detected sooner than dynamical responses<sup>59,60</sup>. It may be that regional sea-ice loss will elicit robust signals in a shorter period.

The Arctic climate system is especially sensitive to external forces that can fundamentally alter climate and ecosystem functioning<sup>62</sup>. Nonlinear threshold behavior of the Arctic climate system to the loss of sea ice has been discussed<sup>63</sup>. There are qualitative hypotheses for the coupled Arctic/subarctic climate system<sup>64</sup> and new approaches such as nonlinear auto-regressive modeling for constructing linear and non-linear dynamical models (e.g. NARMAX)<sup>65,66</sup>. So far, NARMAX has been used to discern changing effects of glaciological, oceanographic and atmospheric conditions on Greenland iceberg numbers over the last century<sup>67</sup>. Novel methods to

distinguish between statistical and causal relationships<sup>68</sup>, the application of artificial intelligence such as evolutionary algorithms<sup>69</sup>, and a Bayesian Hierarchical Model approach may enable progress.

Evidence of systematic midlatitude responses to Arctic warming is beginning to emerge<sup>28-38</sup>. Linkage mechanisms vary with season, region, and system state, and they include both thermodynamic and dynamical processes. A complex web of pathways for linkages, as well as external forcing, is shown in Fig. 2, which summarizes selected recent references. Whilst these linkages shape the overall picture, considered individually they are subject to intermittency in cause and effect. To date, the most consistent regional linkage is supported by case studies and model simulations showing that reduced sea ice in the Barents/Kara Seas (northeast of Scandinavia) can lead to cold continental Asian temperatures<sup>33,70-74</sup>. A doubled probability of severe winters in central Eurasia with increased regional sea ice loss has been reported<sup>75</sup>. This singular linkage mechanism may be the exception rather than the rule<sup>7</sup>. Intermittency implies that frameworks allowing for multiple necessary causal factors may be required to accurately describe linkages in multiple locations.

### *Multiple influences*

Whilst a more consistent picture of linkages may emerge in future scenarios as AA strengthens, one needs to remember that sea ice loss is only one factor of many that influences, and is influenced by, climate change. For example, eastern North American weather is affected by sea-surface temperature patterns in the North Pacific and tropical Pacific<sup>76-79</sup> and also by sea ice loss in the Pacific sector of the Arctic<sup>32,33</sup>. The so-named Snowmageddon blizzard that hit eastern

North America in February 2010 was strengthened by the coincidence of moist, warm air associated with El Niño colliding with frigid air originating from Canada. Downstream influences on the Barents/Kara Sea region, noted for initiating sea ice linkages with eastern Asia, have been connected to the western North Atlantic<sup>80</sup>.

The Arctic can also be influenced by variability from midlatitudes. January through May 2016, for example, set new records for globally averaged temperatures along with the lowest recorded sea ice extent in those months since 1880. Extensive Arctic temperature anomalies of over 7° C were associated with strong southerly winds and warm air originating from the North Pacific, southwestern Russia and the northeastern Atlantic; anomalies for January 2016 are shown in Fig. 3. In contrast, the large scale wind pattern also resulted in a severe, week-long cold surge over eastern Asia during January 2016, evident as the blue region in Fig. 3.

On a hemispheric scale, the relative importance of Arctic versus non-Arctic forcing on atmospheric circulation patterns is uncertain. While models generally suggest that AA and sea ice loss favor a weakened and equatorward-shifted midlatitude storm track, warming of the tropical upper troposphere favors the opposite response<sup>81</sup>. Recent work suggests that Arctic influences may have started to exceed tropical influences in explaining subarctic variability<sup>50,82</sup>. In the long term, the direct warming effect of raised greenhouse gas concentrations favors warm anomalies over cold anomalies, leading to an overall hemispheric tendency for warmer winters<sup>4</sup>.

#### *State dependence*

Arctic thermodynamic influences (e.g., heat fluxes due to snow and sea ice loss, increased water

vapor, changes in clouds) can either reinforce or counteract the amplitude of regional geopotential height fields<sup>60,83</sup>. This response can depend on preexisting atmosphere-ocean conditions and the intensity of the index cycle<sup>49</sup> (state dependence), and can be considered a specific type of intermittency. For example, model simulations suggest that an amplification of the climatological ridge-trough pattern over North America, in response to Arctic sea ice loss, is conditional on the prevailing surface ocean state (Fig. 4). State dependence provides one explanation for why particular causal linkages may only constitute necessary but not sufficient causation.

Variability in the wintertime Arctic stratospheric is another mechanism for state dependence. In winter, planetary waves propagate between the troposphere and stratosphere, and the impacts of this propagation are sensitive to the state of the stratospheric polar vortex<sup>84</sup>. While a strong vortex is characterized by relatively fast-moving westerly winds and a cold core, sudden stratospheric warmings can occur, in which temperatures can increase by over 40° C in a matter of days<sup>85</sup>. These events can weaken, or even reverse, the stratospheric winds, leading to an eventual downward propagation of the circulation feature into the troposphere<sup>86</sup> and a tendency for a negative phase of the AO. This mechanism establishes memory in the system, as sea ice loss and snow cover in late fall can affect the tropospheric jet stream in late winter through lagged transfer of wave-induced disturbances involving the stratosphere<sup>39</sup>. Only models with realistic stratospheres are able to capture this mechanism.

## **Way Forward**

To summarize, the various linkages between AA, large scale midlatitude and tropical sea surface temperature fluctuations, and internal variability of atmospheric circulation are obscured by the four limitations discussed above. These limitations reflect the nonlinearity of climate system dynamics, and the study of linkages remains an unfinished puzzle. Handorf and Dethloff<sup>87</sup> report that current state-of-the-science climate models cannot yet reproduce observed changes in atmospheric teleconnection patterns because of shortcomings in capturing realistic natural variability as well as relationships between the most important teleconnections and patterns of temperature change. Until models are able to realistically reproduce these relationships, an understanding of subarctic climate variability and weather patterns in a warming world remains a challenge.

The complexities and limitations of the linkage issue work against the idea of parsimony in science, of direct causality, or of finding simple pathways. Given the complex web of linkages as illustrated in Fig. 2, an appropriate physics analogy is the effort to understand bulk thermodynamics for an ideal gas by examining only the mechanisms of individual molecular collisions without aggregating statistics. An approach is needed that recognizes multiple processes that act sometimes separately, sometimes interactively in a framework based on the equations of motion and thermodynamics. This is not an easy task but may be achieved through a combination of carefully designed, multi-investigator, coordinated, multi-model simulations, data analyses, and diagnostics.

Studies of linkages are motivated by the potential that a better understanding will benefit decision-makers in their efforts to prepare for impacts of climate change on multi-annual to

decadal timescales, as well as weather-prediction centers producing operational forecasts, particularly at the subseasonal to seasonal timescale. We offer the following recommendations:

- The climate science community needs to develop appropriate diagnostics to analyze model and reanalysis output to detect regional and intermittent responses. Here, major progress is achievable. Although internal variability is a principal characteristic of large scale atmospheric motions, there can be order in large scale atmospheric dynamics that should be further exploited, such as analyses based on potential vorticity (PV), progression of long waves, blocking persistence, and regional surface coupling.
- Nonlinearity and state dependence suggest that idealized and low-resolution climate models have limited explanatory power. Ultimately we need to use realistic models that are validated against observations. Improving the horizontal and vertical resolution is required to properly represent many regional dynamic processes such as jet stream meanders, blocks, polarity of the AO and NAO, teleconnections, surface-atmosphere interaction, stratosphere-troposphere interactions, atmospheric wave propagation, and shifts in planetary waviness<sup>88,89,90</sup>.
- Arctic and subarctic sub-regions are connected over large scales. System-wide studies can help in assessing polar versus tropical drivers on midlatitude jet stream variability.
- Model realism as well as improvements to weather forecasts would benefit from additional observations<sup>91</sup> in the Arctic and subarctic, and by improving global and Arctic meteorological reanalyses, particularly in their representation of surface fluxes<sup>92,93</sup>.
- Better coordination of the research community is needed for model experiments and data analyses, as the current controversy stems in part from uncoordinated efforts.



## Summary

Many recent studies of linkages have focused on direct effects attributed to specific changes in the Arctic, such as reductions in sea ice and snow cover. Disparate conclusions have been reached owing to the use of different data, models, approaches, metrics, and interpretations. Low signal-to-noise ratios and the regional, episodic, and state-dependent nature of linkages further complicate analyses and interpretations. Such efforts have rightly generated controversy.

Based on the large number of recent publications, progress is evident in understanding linkages and in uncovering their regional and seasonal nuances. However, basic limitations are inherent in these efforts. Fig. 5 offers a visualization of the current state of the science, presenting likely pathways for linkages between AA and midlatitude circulation at both the weather timescales (days) and for planetary waves (weeks), as noted on the left. Understanding such pathways can benefit from advanced atmospheric diagnostic and statistical methods. Limitations (center) in deciphering cause-and-effect derive from both itinerancy and multiple simultaneous sources of external forcing. A way forward (right) is through improved data, diagnostics, models, and international cooperation among scientists.

Wintertime cold spells, summer heatwaves, droughts and floods—and their connections to natural variability and forced change—will be topics of active research for years to come. We recommend that the meteorological community “embrace the chaos” as a dominant component of linkages between a rapidly warming Arctic and the midlatitude atmospheric circulation. Scientists should capitalize on and seek avenues to improve the realism and self-consistency of the physical processes in high-resolution numerical models that simultaneously incorporate multiple

processes and internal instabilities. Use of multiple ensembles is essential. Coordination efforts are necessary to move toward community consensus in the understanding of linkages and to better communicate knowns and unknowns to the public. Because of the potential impacts on billions of people living in northern midlatitudes, these priorities have been identified by national and international agencies, such as: the WMO/Polar Prediction Program (PPP), WCRP Climate and Cryosphere (CliC), WCRP Polar Climate Predictability Initiative (PCPI), the International Arctic Science Committee (IASC), the International Arctic Systems for Observing the Atmosphere (IASOA), the US National Science Foundation, NOAA, and the US CLIVAR Arctic-midlatitude working group.

## References

- 1 Serreze, M., Barrett, A., Stroeve, J., Kindig, D. & Holland, M. The emergence of surface-based Arctic amplification. *The Cryosphere* **3**, 11–19 (2009).
- 2 Overland, J. E., Wang, M., Walsh, J. E. & Stroeve, J. C. Future Arctic climate changes: Adaptation and mitigation timescales. *Earth's Future* **2**, 68–74, doi:10.1002/2013EF000162 (2014).
- 3 Francis, J. A. & Vavrus, S. J. Evidence for a wavier jet stream in response to rapid Arctic warming. *Environ. Res. Lett.* **10**, 014005, doi:10.1088/1748-9326/10/1/014005 (2015).
- 4 Wallace, J. M., Held, I. M., Thompson, D. W. J., Trenberth, K. E. & Walsh, J. E. Global warming and winter weather. *Science* **343**, 729–730, doi:10.1126/science.343.6172.729 (2014).
- 5 Barnes, E. A. & Screen, J. A. The impact of Arctic warming on the midlatitude jet-stream: Can it? Has it? Will it? *Clim. Change* **6**, 277–286, doi:10.1002/wcc.337 (2015).

- 374 6 Sun, L., Perlwitz, J., & Hoerling, M. What caused the recent “Warm Arctic, Cold  
375 Continents” trend pattern in winter temperatures? *Geophys. Res. Lett.* **43**,  
376 doi:10.1002/2016GL069024,(2016).
- 377 7 Overland, J. E. *et al.* The melting Arctic and mid-latitude weather patterns: Are they  
378 connected? *J. Clim.* **28**, 7917–7932, doi:10.1175/JCLI-D-14-00822.1 (2015).
- 379 8 Petoukhov, V. & Semenov, V. A. A link between reduced Barents-Kara sea ice and cold  
380 winter extremes over northern continents. *J. Geophys. Res.* **115**, D21111,  
381 doi:10.1029/2009JD013568 (2010).
- 382 9 Peings, Y. & Magnusdottir, G. Response of the wintertime Northern Hemisphere  
383 atmospheric circulation to current and projected Arctic sea ice decline. *J. Clim.* **27**, 244–  
384 264, doi:10.1175/JCLI-D-13-00272.1 (2014).
- 385 10 Semenov, V. A. & Latif, M. Nonlinear winter atmospheric circulation response to Arctic  
386 sea ice concentration anomalies for different periods during 1966–2012. *Environ. Res.*  
387 *Lett.* **10**, 054020, doi:10.1088/1748-9326/10/5/054020 (2015).
- 388 11 Screen, J. A. & Simmonds, I. Exploring links between Arctic amplification and mid-  
389 latitude weather. *Geophys. Res. Lett.* **40**, 959–964, doi:10.1002/grl.50174 (2013).
- 390 12 Barnes, E. A. Revisiting the evidence linking Arctic amplification to extreme weather in  
391 midlatitudes. *Geophys. Res. Lett.* **40**, 4734–4739, doi:10.1002/grl.50880 (2013).
- 392 13 Orsolini, Y. J., Senan, R., Benestad, R. E. & Melsom, A. Autumn atmospheric response  
393 to the 2007 low Arctic sea ice extent in coupled ocean–atmosphere hindcasts. *Clim.*  
394 *Dynam.* **38**, 2437–2448, doi:10.1007/s00382-011-1169-z (2012).

- 395 14 Overland, J. E. *et al.* Air temperature in *Arctic Report Card: Update for 2015* (2015);  
396 [http://www.arctic.noaa.gov/report15/air\\_temperature.html](http://www.arctic.noaa.gov/report15/air_temperature.html).
- 397 15 Screen, J. A. & Simmonds, I. The central role of diminishing sea ice in recent Arctic  
398 temperature amplification. *Nature* **464**, 1334–1337, doi:10.1038/nature09051 (2010).
- 399 16 Coumou, D., Lehmann, J. & Beckmann, J. The weakening summer circulation in the  
400 Northern Hemisphere mid-latitudes. *Science* **348**, 324–327, doi:10.1126/science.1261768  
401 (2015).
- 402 17 Pithan, F. & Mauritsen, T. Arctic amplification dominated by temperature feedbacks in  
403 contemporary climate models. *Nature Geosci.* **7**, 181–184, doi:10.1038/ngeo2071 (2014).
- 404 18 Taylor, P. C. *et al.* A decomposition of feedback contributions to polar warming  
405 amplification. *J. Clim.* **26**, 7023–7043, doi:10.1175/JCLI-D-12-00696.1 (2013).
- 406 19 Porter, D. F., Cassano, J. J. & Serreze, M. C. Local and large-scale atmospheric responses  
407 to reduced Arctic sea ice and ocean warming in the WRF model. *J. Geophys. Res.* **117**,  
408 D11115, doi:10.1029/2011JD016969 (2012).
- 409 20 Overland, J. E. & Wang, M. Y. Large-scale atmospheric circulation changes are  
410 associated with the recent loss of Arctic sea ice. *Tellus A* **62**, 1–9, doi:10.1111/j.1600-  
411 0870.2009.00421.x (2010).
- 412 21 Francis, J. A. & Vavrus, S. J. Evidence linking Arctic amplification to extreme weather in  
413 mid-latitudes. *Geophys. Res. Lett.* **39**, doi:10.1029/2012GL051000 (2012).
- 414 22 Palmer, T. N. A nonlinear dynamical perspective on climate prediction. *J. Clim.* **12**, 575–  
415 591 (1999).

- 416 23 Pearl, J. *Causality: Models, Reasoning and Inference* 2<sup>nd</sup> ed. (Cambridge University  
417 Press, 2009).
- 418 24 Hannart, A., Pearl, J., Otto, F. E. L., Naveau, P. & Ghil, M. Causal counterfactual theory  
419 for the attribution of weather and climate-related events. *Bull. Am. Meteorol. Soc.* **97**, 99–  
420 110, doi:10.1175/BAMS-D-14-00034.1 (2015).
- 421 25 Vihma, T. Effects of Arctic sea ice decline on weather and climate: A review. *Surv.*  
422 *Geophys.*, **35**, 1175–1214, doi:10.1007/s10712-014-9284-0 (2014).
- 423 26 Walsh, J. E. Intensified warming of the Arctic: Causes and impacts on middle latitudes.  
424 *Global Planet. Change* **117**, 52–63, doi:10.1016/j.gloplacha.2014.03.003 (2014).
- 425 27 Thomas, K. (ed.) National Academy of Sciences *Linkages between Arctic Warming and*  
426 *Mid-Latitude Weather Patterns* (The National Academies Press, 2014);  
427 [http://www.nap.edu/catalog/18727/linkages-between-arctic-warming-and-](http://www.nap.edu/catalog/18727/linkages-between-arctic-warming-and-midlatitudeweather-patterns)  
428 [midlatitudeweather-patterns](http://www.nap.edu/catalog/18727/linkages-between-arctic-warming-and-midlatitudeweather-patterns).
- 429 28 Cohen, J. *et al.* Recent Arctic amplification and extreme mid-latitude weather. *Nature*  
430 *Geosci.* **7**, 627–637, doi:10.1038/ngeo2234 (2014).
- 431 29 Jung, T. *et al.* Polar lower-latitude linkages and their role in weather and climate  
432 prediction. *Bull. Am. Meteorol. Soc.* **96**, ES197–ES200, doi:10.1175/BAMS-D-15-  
433 00121.1 (2015).
- 434 30 Hopsch, S., Cohen, J. & Dethloff, K. Analysis of a link between fall Arctic sea ice  
435 concentration and atmospheric patterns in the following winter. *Tellus A* **64**, 18624,  
436 doi:10.3402/tellusa.v64i0.18624 (2012).

- 437 31 Lee, M.-Y., Hong, C.-C. & Hsu, H.-H. Compounding effects of warm SST and reduced  
438 sea ice on the extreme circulation over the extratropical North Pacific and North America  
439 during the 2013–2014 boreal winter. *Geophys. Res. Lett.* **42**, 1612–1618,  
440 doi:10.1002/2014GL062956 (2015).
- 441 32 Kug, J.-S. *et al.* Two distinct influences of Arctic warming on cold winters over North  
442 America and East Asia. *Nature Geosci.* **8**, 759–762, doi:10.1038/ngeo2517 (2015).
- 443 33 King, M. P., Hell, M. & Keenlyside, N. Investigation of the atmospheric mechanisms  
444 related to the autumn sea ice and winter circulation link in the Northern Hemisphere.  
445 *Clim. Dynam.* **46**, 1185–1195, doi:10.1007/s00382-015-2639-5 (2015).
- 446 34 Pedersen, R., Cvijanovic, I., Langen, P. & Vinther, B. The impact of regional Arctic sea  
447 ice loss on atmospheric circulation and the NAO. *J. Clim.* **29**, 889–902,  
448 doi:10.1175/JCLI-D-15-0315.1 (2016).
- 449 35 Tang, Q., Zhang, X. Yang, X. & Francis, J. A. Cold winter extremes in northern  
450 continents linked to Arctic sea ice loss. *Environ. Res. Lett.* **8**, 014036, doi:10.1088/1748-  
451 9326/8/1/014036 (2013).
- 452 36 Furtado, J. C., Cohen, J. L. & Tziperman, E. The combined influences of autumnal snow  
453 and sea ice on Northern Hemisphere winters. *Geophys. Res. Lett.* **43**, 3478–3485,  
454 doi:10.1002/2016GL068108 (2016).
- 455 37 Dobricic, S., Vignati, E. & Russo, S. Large-scale atmospheric warming in winter and the  
456 Arctic sea ice retreat. *J. Clim.* **29**, 2869–2888, doi:10.1175/JCLI-D-15-0417.1 (2016).

457 38 Rinke, A., Dethloff, K., Dorn, W., Handorf, D. & Moore, J. C. Simulated Arctic  
458 atmospheric feedbacks associated with late summer sea ice anomalies. *J. Geophys. Res.*  
459 **118**, 7698–7714, doi:10.1002/jgrd.50584 (2013).

460 39 Nakamura, T. *et al.* A negative phase shift of the winter AO/NAO due to the recent  
461 Arctic sea-ice reduction in late autumn. *J. Geophys. Res. (Atmos.)* **120**, 3209–3227,  
462 doi:10.1002/2014JD022848 (2015).

463 40 Duarte, C., Lenton, T., Wadhams, P. & Wassmann, P. Abrupt climate change in the  
464 Arctic. *Nature Clim. Change* **2**, 60–62, doi:10.1038/nclimate1386 (2012).

465 41 Wu, B., Handorf, D., Dethloff, K., Rinke, A. & Hu, A. Winter weather patterns over  
466 northern Eurasia and Arctic sea ice loss. *Mon. Weather Rev.* **141**, 3786–3800,  
467 doi:10.1175/MWR-D-13-00046.1 (2013).

468 42 Corti, S., Molteni, F. & Palmer, T. N. Signature of recent climate change in frequencies  
469 of natural atmospheric circulation regimes. *Nature* **396**, 799–802 (1999).

470 43 Itoh, H. & Kimoto, M. Weather regimes, low-frequency oscillations, and principal  
471 patterns of variability: A perspective of extratropical low-frequency variability. *J. Atmos.*  
472 *Sci.* **56**, 2684–2705 (1999).

473 44 Sempf, M., Dethloff, K., Handorf, D. & Kurgansky, M. V. Toward understanding the  
474 dynamical origin of atmospheric regime behavior in a baroclinic model. *J. Atmos. Sci.* **64**,  
475 887–904, doi:10.1175/JAS3862.1 (2007).

476 45 Slingo, J. & Palmer, T. Uncertainty in weather and climate prediction. *Philos. Trans.*  
477 *Roy. Soc. A* **369**, 4751–4767 (2011).

478 46 Schmeits, M. J. & Dijkstra, H. A. Bimodal behavior of the Kuroshio and the Gulf Stream.  
479 *J. Phys. Oceanogr.* **31**, 3435–3456 (2001).

480 47 Davos, V. *et al.* Methods for detecting early warnings of critical transitions in time series  
481 illustrated using ecological data. *PLoS ONE* **7**, e41010,  
482 doi:10.1371/journal.pone.0041010 (2013).

483 48 Screen, J. A., Deser, C. & Sun, L. Projected changes in regional climate extremes arising  
484 from Arctic sea ice loss. *Environ. Res. Lett.* **10**, 084006 (2015).

485 49 Overland, J. E. & Wang, M. Increased variability in the early winter subarctic North  
486 American atmospheric circulation. *J. Clim.* **28**, 7297–7305, doi:10.1175/JCLI-D-15-  
487 0395.1 (2015).

488 50 Cohen, J. An observational analysis: Tropical relative to Arctic influence on midlatitude  
489 weather in the era of Arctic amplification. *Geophys. Res. Lett.* **43**,  
490 doi:10.1002/2016GL069102 (2016).

491 51 Hanna, E., Cropper, T. E., Jones, P. D., Scaife, A. A. & Allan, R. Recent seasonal  
492 asymmetric changes in the NAO (a marked summer decline and increased winter  
493 variability) and associated changes in the AO and Greenland Blocking Index. *Int. J.*  
494 *Climatol.* **35**, 2540–2554, doi:10.1002/joc.4157 (2015).

495 52 Woollings, T., Hannachi, A. & Hoskins, B. Variability of the North Atlantic eddy-driven  
496 jet stream. *Q. J. Roy. Meteorol. Soc.* **136**, 856–868, doi:10.1002/qj.625 (2010).

497 53 Hanna, E., Cropper, T. E., Hall, R. J. & Cappelen, J. Greenland Blocking Index 1851–  
498 2015: A regional climate change signal. *Int. J. Climatol.*, doi:10.1002/joc.4673 (2016).



499 54 Masato, G., Hoskins, B. J. & Woollings, T. Winter and summer Northern Hemisphere  
500 blocking in CMIP5 models. *J Clim.* **26**, 7044–7059, doi:10.1175/JCLI-D-12-00466.1  
501 (2013).

502 55 Scaife, A. A. *et al.* Skillful long-range prediction of European and North American  
503 winters. *Geophys. Res. Lett.* **41**, 2514–2519, doi:10.1002/2014GL059637 (2014).

504 56 Eade, R. *et al.* Do seasonal-to-decadal climate predictions underestimate the  
505 predictability of the real world? *Geophys. Res. Lett.* **41**, 5620–5628,  
506 doi:10.1002/2014GL061146 (2014).

507 57 Stockdale, T. N. *et al.* Atmospheric initial conditions and the predictability of the Arctic  
508 Oscillation. *Geophys. Res. Lett.* **42**, 1173–1179, doi:10.1002/2014GL062681 (2015).

509 58 Barnes, E. A. & Polvani, L. M. CMIP5 projections of Arctic amplification, of the North  
510 American/North Atlantic Circulation, and of their relationship. *J. Clim.* **28**, 5254–5271,  
511 doi:10.1175/JCLI-D-00589.1 (2015).

512 59 Screen, J. A., Deser, C., Simmonds, I. & Tomas, R. Atmospheric impacts of Arctic sea-  
513 ice loss, 1979–2009: Separating forced change from atmospheric internal variability.  
514 *Clim. Dyn.* **43**, 333–344, doi:10.1007/s00382-013-1830-9 (2014).

515 60 Shepherd, T. G. Atmospheric circulation as a source of uncertainty in climate change  
516 projections. *Nature Geosci.* **7**, 703–708, doi:10.1038/ngeo2253 (2014).

517 61 Hinzmann, L. *et al.* Trajectory of the Arctic as an integrated system. *Ecol. Appl.* **23**,  
518 1837–1868, doi:10.1890/11-1498.1 (2013).

519 62 Carstensen, J. & Weydmann, A. Tipping points in the Arctic: Eyeballing or statistical  
520 significance? *AMBIO* **41**, 34–43 (2012).

521 63 Eisenman, I. & Wettlaufer, J. S. Nonlinear threshold behavior during the loss of Arctic  
522 sea ice. *Proc. Natl. Acad. Sci. USA* **106**, 28–32, doi:10.1073/pnas.0806887106 (2009).

523 64 Mysak, L. A. & Venegas, S. A. Decadal climate oscillations in the Arctic: A new  
524 feedback loop for atmosphere-ice-ocean interactions. *Geophys. Res. Lett.* **25**, 3607–3610,  
525 (1998).

526 65 Billings, S. A., Chen, S. & Korenberg, M. J. Identification of MIMO non-linear systems  
527 using a forward-regression orthogonal estimator. *Int. J. Control* **49**, 2157–2189,  
528 doi:10.1080/00207178908559767 (1989).

529 66 Billings, S. A. *Nonlinear System Identification: NARMAX Methods in the Time,*  
530 *Frequency, and Spatio-Temporal Domains* (Wiley, 2013).

531 67 Bigg, G. R. *et al.* A century of variation in the dependence of Greenland iceberg calving  
532 on ice sheet surface mass balance and regional climate change. *Proc. R. Soc. A* **470**,  
533 20130662, doi:10.1098/rspa.2013.0662 (2014).

534 68 Kretschmer, M., Coumou, D., Donges, J. & Runge, J. Using causal effect networks to  
535 analyze different Arctic drivers of midlatitude winter circulation. *J. Clim.* **29**, 4069–4081  
536 doi:10.1175/JCLI-D-15-0654.1 (2016).

537 69 Stanislawska, K., Krawiec, K. & Kundzewicz, Z.W. Modeling global temperature  
538 changes with genetic programming. *Comput. Math. Appl.* **64**, 3717–3728 (2012).

539 70 Honda, M., Inoue, J. & Yamane, S. Influence of low Arctic sea-ice minima on  
540 anomalously cold Eurasian winters. *Geophys. Res. Lett.* **36**, L08707,  
541 doi:10.1029/2008GL037079 (2009).

542 71 Kim, B.-M. *et al.* Weakening of the stratospheric polar vortex by Arctic sea-ice loss.  
543 *Nature Commun.* **5**, 4646, doi:10.1038/ncomms5646 (2014).

544 72 Jaiser, R., Dethloff, K. & Handorf, D. Stratospheric response to Arctic sea ice retreat and  
545 associated planetary wave propagation changes. *Tellus A* **65**, 19375,  
546 doi:10.3402/tellusa.v65i0.19375 (2013).

547 73 Handorf, D., Jaiser, R., Dethloff, K., Rinke, A. & Cohen, J. Impacts of Arctic sea-ice and  
548 continental snow-cover changes on atmospheric winter teleconnections. *Geophys. Res.*  
549 *Lett.* **42**, 2367–2377 doi:10.1002/2015GL063203 (2015).

550 74 Luo, D. *et al.* Impact of Ural blocking on winter warm Arctic–cold Eurasian anomalies.  
551 Part I: Blocking-induced amplification. *J. Clim.* **29**, 3925–3947, doi:10.1175/JCLI-D-15-  
552 0611.1 (2016).

553 75 Mori, M. Watanabe, M., Shiogama, H., Inoue, J. & Kimoto, M. Robust Arctic sea-ice  
554 influence on the frequent Eurasian cold winters in past decades. *Nature Geosci.* **7**, 869–  
555 873, doi:10.1038/ngeo2277 (2014).

556 76 Ding, Q. *et al.* Tropical forcing of the recent rapid Arctic warming in northeastern  
557 Canada and Greenland. *Nature* **509**, 209–212, doi:10.1038/nature13260 (2014).

558 77 Perlwitz, J., Hoerling, M. & Dole, R. Arctic tropospheric warming: Causes and linkages  
559 to lower latitudes. *J. Clim.* **28**, 2154–2167 (2015).

560 78 Hartmann, D. L. Pacific sea surface temperature and the winter of 2014. *Geophys. Res.*  
561 *Lett.* **42**, 1894–1902, doi:10.1002/2015GL063083 (2015).

562 79 Screen J. & Francis, J. Contribution of sea-ice loss to Arctic amplification regulated by  
563 Pacific Ocean decadal variability. *Nature Clim. Change*, accepted (2016).

564 80 Sato, K., Inoue, J. & Watanabe, M. Influence of the Gulf Stream on the Barents Sea ice  
565 retreat and Eurasian coldness during early winter. *Environ. Res. Lett.* **9**, 084009 (2014).

566 81 Harvey, B. J., Shaffrey, L. C. & Woollings, T. Deconstructing the climate change  
567 response of the Northern Hemisphere wintertime storm tracks. *Clim. Dynam.* **45**, 2847–  
568 2860 (2015).

569 82 Feldstein, S. B. & Lee, S. Intraseasonal and interdecadal jet shifts in the Northern  
570 Hemisphere: The role of warm pool tropical convection and sea ice. *J. Clim.* **27**, 6497–  
571 6518, doi:10.1175/JCLI-D-14-00057.1 (2014).

572 83 Trenberth, K. E., Fasullo, J. T. & Shepherd, T. G. Attribution of climate extreme events.  
573 *Nature Clim. Change* **5**, 725–730, doi:10.1038/nclimate2657 (2015).

574 84 Sigmond, M. & Scinocca, J. F. The influence of the basic state on the Northern  
575 Hemisphere circulation response to climate change. *J. Clim.* **23**, 1434–1446 (2010).

576 85 Butler, A. H. *et al.* Defining sudden stratospheric warmings. *Bull. Am. Meteorol. Soc.* **96**,  
577 1913–1928, doi:10.1175/BAMS-D-13-00173.1 (2015).

578 86 Sigmond, M., Scinocca, J. F., Kharin, V. V. & Shepherd, T. G. Enhanced seasonal  
579 forecast skill following stratospheric sudden warmings. *Nature Geosci.* **6**, 98–102,  
580 (2013).

581 87 Handorf, D. & Dethloff, K. How well do state-of-the-art atmosphere-ocean general  
582 circulation models reproduce atmospheric teleconnection patterns? *Tellus A* **64**, 19777,  
583 doi:10.3402/tellusa.v64i0.19777 (2012).

584 88 Byrkjedal, Ø., Esau, I. N. & Kvamstø, N. G. Sensitivity of simulated wintertime Arctic  
585 atmosphere to vertical resolution in the ARPEGE/IFS model. *Clim. Dynam.* **30**, 687–701,  
586 doi:10.1007/s00382-007-0316-z (2008).

587 89 Wu, Y. & Smith, K. L. Response of Northern Hemisphere midlatitude circulation to  
588 Arctic amplification in a simple atmospheric general circulation model. *J. Clim.* **29**,  
589 2041–2058, doi:10.1175/JCLI-D-15-0602.1 (2016).

590 90 Anstey, J. A. *et al.* Multi-model analysis of Northern Hemisphere winter blocking: Model  
591 biases and the role of resolution. *J. Geophys. Res. Atmos.* **118**, 3956–3971,  
592 doi:10.1002/jgrd.50231 (2013).

593 91 Inoue, J. *et al.* Additional Arctic observations improve weather and sea-ice forecasts for  
594 the Northern Sea Route. *Sci. Rep.* **5**, 16868, doi:10.1038/srep16868, (2015).

595 92 Schlichtholz, P. Empirical relationships between summertime oceanic heat anomalies in  
596 the Nordic seas and large-scale atmospheric circulation in the following winter. *Clim.*  
597 *Dyn.*, doi:10.1007/s00382-015-2930-5 (2016).

598 93 Lindsay, R., Wensnahan, M., Schweiger, A. & Zhang, J. Evaluation of seven different  
599 atmospheric reanalysis products in the Arctic. *J. Clim.* **27**, 2588–2606, doi:10.1175/JCLI-  
600 D-13-00014.1 (2014).

601 94 Handorf, D., Dethloff, K., Marshall, A. G. & Lynch, A. Climate regime variability for  
602 past and present time slices simulated by the Fast Ocean Atmosphere Model. *J. Clim.* **22**,  
603 58–70, doi:10.1175/2008 JCLI2258.1 (2009).

- 95 Gervais, M., Atallah, E., Gyakum, J. R. & Tremblay, L. B., Arctic air masses in a  
warming world. *J. Clim.* **29**, 2359–2373, doi:10.1175/JCLI-D-15-0499.1. (2016).
- 96 Francis, J. & Skific, N. Evidence linking rapid Arctic warming to mid-latitude weather  
patterns. *Philos. Trans. R. Soc. Lond. Ser. A* **373**, 20140170, doi:10.1098/rsta.2014.0170  
(2015).
- 97 Screen, J. A. & Simmonds, I. Amplified mid-latitude planetary waves favour particular  
regional weather extremes. *Nature Clim. Change* **4**, 704–709, doi:10.1038/nclimate2271  
(2014).

To whom correspondence and requests for materials should be addressed:

James Overland [james.e.overland@noaa.gov](mailto:james.e.overland@noaa.gov)

Acknowledgements. JEO is supported by NOAA Arctic Research Project of the Climate  
Program Office. JAF is supported by NSF/ARCSS Grant 1304097. KD acknowledge support  
from the German DFG Transregional Collaborative Research Centre TR 172. RH and EH  
acknowledge support from the University of Sheffield’s Project Sunshine. S-JK was supported  
by the project of Korea Polar Research Institute (PE16010), and TV was supported by the  
Academy of Finland (Contract 259537). We appreciate the support of IASC, CliC and the  
University of Sheffield for hosting a productive workshop. PMEL Contribution Number 4429.

#### **Author Contributions & Competing Financial Interests statement**

JEO was the coordinating author and all other authors contributed ideas, analyses, and text.

There are no competing financial interests.

## Figure captions

Figure 1: (a, b) Geopotential height (units of meters) of the 500 hPa pressure surface, illustrating the northern hemisphere's tropospheric polar jet stream where height lines are closely spaced. Winds of the jet stream follow the direction parallel to contours, forming the persistent vortex that circulates counterclockwise around the North Pole. The primarily west-to-east wind flow can adopt a relatively circular pattern (a, for 15 November 2013) or a wavy one (b, for 5 January 2014). The lower panels (c, d) show the corresponding air temperature anomaly patterns (units of °C) for the same days at a lower atmospheric level (850 hPa).

Figure 2: A complex web of pathways that summarize examples of potential mechanisms that contribute to more frequent amplified flow and more persistent weather patterns in mid-latitudes. Numbers 1-11 refer to original literature listed below diagram, and [ ] refer to these citations in the current reference list. BK is Barents/Kara Seas area, EKE is eddy kinetic energy, and SLP is sea-level atmospheric pressure. For details on processes consult the original references.

Figure 3: Global air temperatures anomalies (°C) for January 2016 were the highest in the historical record for any January beginning in 1880. Southerly winds from midlatitudes contributed to the largest anomalies in the Arctic (+7° C). Note the cold anomaly (blue) over Asia. Source: NASA.

Figure 4: State dependence of the atmospheric response to Arctic sea-ice loss. Model simulated wintertime 500 hPa geopotential height responses to Arctic sea ice loss for two different surface ocean states. The responses are estimated from four 100-yr long atmospheric model simulations, with prescribed sea ice concentrations and sea surface temperatures. Experiments *A* and *C* have identical below-average sea ice conditions. Experiments *B* and *D* have identical above-average sea ice conditions. Experiments *A* and *B*, and *C* and *D*, have identical sea surface temperatures, but the two pairs have different sea surface temperatures from one another (i.e., *A* and *B* differ from *C* and *D*; see Supplementary Figure 1), capturing opposite phases of the Atlantic Multidecadal Oscillation (AMO). The response to sea-ice loss, under different surface ocean states, is estimated by contrasting experiments (a) *A* and *B*, and (b) *C* and *D*. The grey box highlights the midlatitude Pacific-American region, where a wave-train response to sea-ice loss is simulated for one SST state (a; negative AMO) but not the other (b; positive AMO), implying that the response to sea-ice loss is state dependent. Green hatching denotes responses that are statistically significant at the 95% ( $p=0.05$ ) confidence level.

Figure 5: Current state of the science for selected linkages. Arctic amplification and some pathways are known (left), but chaotic instabilities and multiple external forcing sources are noted under Limitations (center). (Right) A way forward is through improved data, models, and international cooperation of individual researchers.

## Figures






Figure 1. (a, b) Geopotential height (units of meters) of the 500 hPa pressure surface, illustrating the northern hemisphere's polar jet stream where height lines are closely spaced. Winds of the jet stream follow the direction parallel to contours, forming the persistent vortex that circulates counterclockwise around the North Pole. The primarily west-to-east wind flow can adopt a relatively circular pattern (a, for 15 November 2013) or a wavy one (b, for 5 January 2014). The lower panels (c, d) show the corresponding air temperature anomaly patterns (units of °C) for the same days at a lower atmospheric level (850 hPa).




Figure 2: A complex web of pathways that summarize examples of potential mechanisms that contribute to more frequent amplified flow and more persistent weather patterns in mid-latitudes. Numbers 1-11 refer to original literature listed below diagram, and [ ] refer to these citations in the current reference list. BK is Barents/Kara Seas area, EKE is eddy kinetic energy, and SLP is sea-level atmospheric pressure. For details on processes consult the original references.

January 2016

L-OTI(°C) Anomaly vs 1951-1980

1.13




Figure 3. Global air temperatures anomalies (°C) for January 2016 were the highest in the historical record for any January beginning in 1880. Southerly winds from northern midlatitudes contributed to the largest anomalies in the Arctic (+7° C). Note the cold anomaly (blue) over Asia. Source: NASA.




Figure 4:

State dependence of the atmospheric response to Arctic sea-ice loss. Model simulated wintertime 500 hPa geopotential height responses to Arctic sea ice loss for two different surface ocean states. The responses are estimated from four 100-yr long atmospheric model simulations, with prescribed sea ice concentrations and sea surface temperatures. Experiments *A* and *C* have identical below-average sea ice conditions. Experiments *B* and *D* have identical above-average sea ice conditions. Experiments *A* and *B*, and *C* and *D*, have identical sea surface temperatures, but the two pairs have different sea surface temperatures from one another (i.e., *A* and *B* differ from *C* and *D*; see Supplementary Figure 1), capturing opposite phases of the Atlantic Multidecadal Oscillation (AMO). The response to sea-ice loss, under different surface ocean states, is estimated by contrasting experiments (a) *A* and *B*, and (b) *C* and *D*. The grey box highlights the midlatitude Pacific-American region, where a wave-train response to sea-ice loss is simulated for one SST state (a; negative AMO) but not the other (b; positive AMO), implying that the response to sea-ice loss is state dependent. Green hatching denotes responses that are statistically significant at the 95% ( $p=0.05$ ) confidence level.




Figure 5. Current state of the science for selected linkages. Arctic amplification and some pathways are known (left), but chaotic instabilities and multiple external forcing sources are noted under Limitations (center). (Right) A way forward is through improved data, models, and international cooperation of individual researchers.



**a) Experiment A - B / C - D    b) Experiment A - C / B - D**




Figure S1: Prescribed surface boundary conditions. Differences in prescribed winter sea ice concentrations (a) and sea surface temperatures (b) between the experiments presented in Figure 4 of the main material. Experiments A and C have identical below-average sea ice conditions whilst experiments B and D have identical above-average sea ice conditions, and the difference between these is presented in (a). Experiments A and B, and C and D, have identical sea surface temperatures, but the two pairs have different sea surface temperatures from one another, with this difference shown in (b).