

*Dental caries as a measure of diet, health,
and difference in non-adults from urban
and rural Roman Britain*

Article

Accepted Version

Rohnbogner, A. and Lewis, M. ORCID: <https://orcid.org/0000-0001-6224-0278> (2016) Dental caries as a measure of diet, health, and difference in non-adults from urban and rural Roman Britain. *Dental Anthropology*, 29 (1). pp. 16-31. ISSN 1096-9411 Available at <https://centaur.reading.ac.uk/67761/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Publisher: Dental Anthropology Association

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Title

Dental caries as a measure of diet, health, and difference in non-adults from urban and rural Roman Britain.

Author name and affiliation (*corresponding author)

Anna Rohnbogner^{1*}

Mary Lewis¹

¹University of Reading, UK

Archaeology, School of Archaeology, Geography and Environmental Science, University of Reading, Reading, UK, RG6 6AB

*a.j.rohnbogner@reading.ac.uk; annarohnbogner@hotmail.com

Abstract

Dental disease in childhood has the potential to inform about food availability, social status, and feeding practices, in addition to contributing to a child's overall health status. This paper presents the first comprehensive overview of carious lesion frequencies in 433 non-adults (1-17 years), and 6283 erupted permanent and deciduous teeth from 15 urban and rural Romano-British settlements. Pooled deciduous and permanent caries rates were significantly higher in major urban sites (1.8%) compared to rural settlements (0.4%), with children from urban sites having significantly higher lesion rates in the deciduous dentition (3.0%), and in younger age groups with mixed dentitions. The differences in dental caries between urban and rural populations suggest disparities in maternal oral health, early childhood feeding practices, food preparation and access to refined carbohydrates. A richer, perhaps more 'Roman', cuisine was eaten in the urban settlements, as opposed to a more modest diet in the countryside. The effect of early childhood stress on caries frequency was explored using evidence for enamel hypoplasia. Co-occurrence of caries and enamel hypoplasia was highest in the major urban cohort (5.8%) and lowest in the rural sample (1.3%), suggesting that environmental stress was a contributing factor to carious lesion development in Romano-British urban children.

Keywords

Dental disease, carious lesion, Romanisation, dental enamel hypoplasia, co-occurrence

INTRODUCTION

Child health is an understudied subject in the archaeology of Roman Britain. Most of what we know about growing-up in the province is derived from the Classical literature. While archaeological research has mainly focussed on the architectural grandeur of the towns and high status *villa* settlements (Parkins, 1997; Millett, 2005; Mattingly, 2006; Pearce, 2008; Holbrook, 2015), rural settlements and their inhabitants have received scant attention preventing a more holistic view of life in *Britannia* (Taylor, 2007; McCarthy, 2013; Breeze, 2014; Fulford and Holbrook, 2014). Contrary to the long-held belief in the detrimental effects of the urban environment, recent bioarchaeological research has demonstrated that living in the Romano-British countryside also negatively affected health (Pitts and Griffin, 2012; Redfern et al., 2015). The late Romano-British *villa* economy may have provided health challenges for its workers, a subject yet to be fully explored. While non-adult skeletons are widely acknowledged to provide an intricate measure of population fitness in the past (Mensforth et al., 1978), comparisons of the morbidity and mortality of urban and rural Romano-British children have yet to be carried out. A child's diet would have reflected the cultural beliefs and social standing of their family. Carious lesion frequency can therefore provide valuable information on access to resources, eating habits and food preparation practices.

In children, calcified plaque (calculus), and pathological conditions of the dentition such as periapical lesions, periodontal disease and antemortem tooth loss are rare. Carious lesion frequencies in the deciduous and permanent teeth therefore provide the most effective measure for discussing past dental health childhood (Halcrow et al., 2013). The presence of caries in children not only indicates the levels of carbohydrates consumed, but also informs on cultural practices such as pre-mastication, and sharing of foods or utensils (Nield et al., 2008). Caries can cause considerable discomfort, resulting in intense pain, difficulty eating, reduced efficiency of the immune system (Baqain et al., 2004), and can affect speech development (Aligne et al., 2003). Dental studies that focus on children are rare in the palaeopathological literature and tend to focus on individual sites or have small sample sizes (e.g. Moore and Corbett, 1973; Whittaker et al., 1981; O'Sullivan et al., 1993; Clough and Boyle, 2010; Redfern et al., 2012). This study will provide an overview of carious lesion frequency in non-adults from 1st to early 5th century AD urban and rural settlements in Roman Britain. Due to the general absence of periodontal disease, pericapical lesions, antemortem tooth loss and calculus in the dentitions of children, the study relies on dental

caries as the sole measure of childhood dental health. The aim of the study is to assess whether an urban or rural living environment impacted on childhood dental health through the investigation of lesion frequencies of dental caries in the deciduous and permanent dentition.

Pathogenesis of dental caries in children

Caries is a progressive infectious disease of the deciduous and permanent dentition, with localised demineralisation of the dental hard tissues by organic acids (Larsen, 1997). *Streptococcus* and *Lactobacillus* bacteria in the oral cavity metabolise sugars and starches which create an acidic environment and demineralise tooth enamel (Byun et al. 2004). Teeth are remineralised as soon as pH levels are restored to neutral. If however, the pH is low for a prolonged period, the dynamic between de- and remineralisation is upset, resulting in a cavity (Gussy et al., 2006). *Streptococcus mutans*, the main cariogenic bacteria, are normally transmitted from the mother or other caregiver to the child through kissing or sharing implements (Nield et al., 2008). Colonisation with *S. mutans* can commence as early as six months old, although the highest risk for infection is at two years old. It takes around 13-16 months from colonisation for a lesion to develop (Kawashita et al., 2011).

‘Early Childhood Caries’ defined as caries in children aged 0-5 years, is a serious chronic condition in the modern world (Afroughi et al., 2010). Newly erupted deciduous teeth are particularly susceptible to caries due to incomplete maturation, large dental tubules and thinner enamel which is insufficient to prevent the progression of caries (Aligne et al., 2003; Schuurs, 2013). The area around the erupting tooth also provides favourable conditions for bacterial colonisation (Schuurs, 2013). Child saliva flow rate is slower than in adults, and has lower levels of secretory immunoglobulin A (IgA) concentrations. While IgA begins to be produced after one month of life, its formation can be impaired by high levels of cortisol in the blood through stress. Stress could therefore have a caries-promoting function. It may manifest as enamel defects, i.e. hypoplasia, which in turn may increase the risk of dental caries (Boyce et al., 2010). Hypoplasia may cause caries to progress quicker due to higher acid solubility of defective enamel, and greater adhesion of cariogenic bacteria at the site of a defect (Hong et al., 2009). The presence of enamel hypoplasia in individuals with caries may therefore be a valuable indicator of early childhood stress in relation to dental decay, and it is of interest to evaluate their co-occurrence.

Using the jaws of modern children, Afroughi and colleagues (2010) showed that once a tooth is infected, caries continues to spread to the teeth on either side or above, often in a symmetrical pattern. Risk of lesion development was greater for maxillary posterior teeth than for anterior or mandibular teeth. Early childhood caries may take on a rampant virulent form, known as ‘Severe Early Childhood Caries’ (SECC), ‘Nursing Caries’ or ‘Milk Bottle Syndrome’. The condition is characterised by rapid lesion development and dental decay primarily in the maxillary anterior dentition (Berkowitz, 2003; Azevedo et al., 2005). A probable case of SECC has recently been identified by Bonsall and colleagues (2015) in a 3-4 year old child from Roman Ancaster, Lincolnshire.

Diet is a crucial factor in the onset and proliferation of caries. Refined foods with a high carbohydrate and/or sugar content encourage the metabolic activity of oral bacteria and acid production, increasing the risk of lesion development (Powell, 1985; Prowse et al., 2008). A tough and fibrous diet has a cleaning effect, and vigorous mastication stimulates salivary flow which is alkaline, buffering against plaque acids (Duray, 1992; Moynihan, 2000). Soft sticky foods and prolonged snacking or sipping of sweetened fluids pose a greater risk for acid development (Hallet and O’Rourke, 2003). Dental disease is usually described by the different surfaces of the tooth that are affected: the occlusal surfaces, smooth surfaces of the crown including the interproximal areas, or the root (Moore and Corbett, 1973). Each of these surfaces has different cariogenic potential, therefore the location of carious lesions on the tooth may provide an insight into changes in diet (Ortner, 2003). With the intensification of cereal agriculture, caries at the root and cemento-enamel junction rises, and with the introduction of refined sugars, interproximal and fissure caries increase, especially in children (Hillson 1996, 283).

The Romano-British diet

There are a number of historical sources that make reference to ‘Roman’ food and drink, including Apicius’ collection of recipes, Pliny the Elder’s *Naturalis Historia* or Galen’s medical writings (Grocock and Grainger, 2006; Alcock, 2010). These sources reflect Mediterranean practices of the literate upper classes in the 1st and 2nd centuries AD, and it remains unknown how these are relevant for exploring diet in Roman Britain. When comparing literary evidence on diet to recent isotopic and osteological studies of populations from Rome (Rutgers et al. 2009; Killgrove and Tykot 2013), Velia (Craig et al. 2009), and

Portus Romae (Prowse et al. 2008), it transpires that a ‘typical’ Roman diet as described in Classical sources may not have been followed, or in fact existed.

Cereal products such as bread and porridge may have been the staple foods in Britain, although variation across the social and geographical strata is expected (Cool, 2006). Sugars would have been available as fructose from fruits, fruit juices, honey and syrups, or glucose in the carbohydrates consumed (Moore and Corbett, 1973; Bowman and Thomas, 1994; Cool, 2006; Bogdanov et al., 2008; Nassar et al., 2012). More advanced farming and plant cultivation techniques, alongside larger scale animal breeding, would have ensured a stable food supply for the army and non-producing urban population, whilst also putting increasing demands on the workers in the countryside (Mattingly, 1997; Dobney, 2001; Taylor, 2001; Pitts and Griffin, 2012; McCarthy, 2013; Redfern et al., 2015).

Studies by King (1984; 1999; 2001) and Cummings (2009) have demonstrated that access to meat and animal products was dependent on site type and status. Overall, marine and freshwater fish became increasingly fashionable, probably as a result of following Roman tastes (Locker, 2007). Isotopic studies on human bones (Richards et al., 1998; Redfern et al., 2010; Cheung et al., 2012; Müldner, 2013) and archaeobotany (van der Veen, 2008; van der Veen et al., 2007; 2008) have revealed temporal, cultural, social and sex differences in the consumption of terrestrial, plant and aquatic foods at an inter-site and intra-site level. Richard and colleagues’ (1998) study of Poundbury Camp in Dorset showed that animal protein and marine foods were only available to the few. Children are thought to have enjoyed more marine foods, and diet was more varied for the inhabitants of the towns (Redfern et al., 2010; Müldner, 2013). It also appears that marine and freshwater foods were primarily consumed in the urban environment rather than the surrounding villages, and that the male and female diet, at least within Roman Gloucester, differed (Cheung et al., 2012). To the benefit of Britons, a variety of new plant foods were introduced following the Roman conquest, improving the nutritional value of the Romano-British diet. Examples include cherries, carrots, or plum, also beet and cabbage, which are high in nutrients and vitamins (van der Veen et al., 2008). As a general pattern, plant foods were widely accessible and eaten in both town and country (van der Veen et al., 2007). It is of note that some areas of the province differed. Access to new plant foods was more restricted, or perhaps opposed, in the southwest (van der Veen et al., 2008).

In children, dietary experiences would have started at weaning with the gradual introduction of solid foods to supplement breastmilk. Galen’s writings and Soranus’ *Gynaecology*, both dating to the 2nd century AD, recommend the introduction of

supplementary foods from around six months (Fildes, 1986; Temkin, 1991). Weaning would have been complete by 2-4 years old, a practice that has been supported by isotopic analysis of non-adults from England and Italy (Fildes, 1986; Fuller et al., 2006; Prowse et al., 2008; Prowse, 2011; Nehlich et al., 2011; Redfern et al., 2012; Powell et al., 2014). The Roman weaning diet comprised mainly of cereal foods, such as porridge, or bread mixed with milk, wine or honey (Temkin, 1991; Garnsey, 1999). Pre-mastication of foods for weanlings was warned to be harmful, however the fact that the practice needed to be discouraged suggests it was a known method of infant feeding (Bradley, 1986; Temkin, 1991). Honey was also used as a popular medicinal aid. Soranus suggests rubbing honey on the gums of teething infants to soothe their pain (Temkin, 1991). Once weaned, the contents of the Roman early childhood diet are unknown. If diet was influenced by location and status, children would have been subjected to differential food allocation. Child dental disease rates may then also differ, depending on urban and rural site types, or high versus low status settlements. A large scale investigation of dental caries rates may therefore provide the first detailed evidence for cultural practices and dietary habits of children in Roman Britain.

MATERIALS AND METHODS

Site classification

The terms 'urban' and 'rural' are used to characterise the nature of the settlement rather than its geographic locale. In order to allow for a comparison of carious lesion frequencies between settlements under different levels of Roman influence, sites were divided into major urban (*coloniae, civitates*), minor urban (nucleated/small towns) and rural settlements (villages, farmsteads, *villa* estates). A common denominator for Romano-British towns, whether large or small, is the dependence on the hinterland for agricultural surplus to feed the non-producing urban population (Mattingly, 1997; Morley, 1997). 'Major urban' sites were defined as large legal and administrative settlements with a series of features including a grid layout for the street system and organised planning throughout, with public buildings, a forum and a spiritual focus (Wacher, 1974; Burnham and Wacher, 1990; Millett, 1990; Laurence et al., 2011). 'Minor urban' centres displayed some urban aspects, such as evidence for town planning and a market to facilitate local trade (Hingley, 1989; Burnham, 1993; 1995; Millett, 1995; Wilson, 2011). Rural sites were defined as undefended farmsteads, villages or *villa* estates with a predominantly agricultural focus (McCarthy, 2013). Depending on their location, rural sites would have exhibited varying economic and socio-cultural

dependence on nearby towns (Laurence et al., 2011; White, 2014). However, their agricultural focus still rendered them as rural in character and urbanisation of the countryside to the extent seen in Italy was not apparent in Roman Britain (Laurence, 2011). Current models on life in the Romano-British countryside consider *villa* economies as estates managed by landowners, with a peasant population that cultivates the land as tenants or freeholders, living either on the estate or surrounding villages (Taylor, 2001; Mattingly, 2006; McCarthy, 2013; Breeze, 2014).

There is a growing awareness of the difficulty in classifying Romano-British settlements (Mattingly, 1997; Millett, 1999; Burnham et al., 2001; Millett, 2001; Pearce, 2008; Rogers, 2011), and it is important to bear in mind that power in Roman Britain was not exclusively urban, as the elite often resided in the countryside or a town's immediate hinterland (Parkins, 1997; Pitts and Perring, 2006). In addition, not all urban cemeteries would have contained those living and dying exclusively within these large settlements, as many individuals may have been derived from the "urban periphery" (Goodman, 2007:1-2), or represent rural migrants (Griffin and Pitts, 2012; Redfern et al., 2015).

Materials

Dental caries was recorded in 433 non-adults (aged 1.1-17.0 years) and a total of 6283 erupted teeth (deciduous n=2910; permanent n=3373) from 15 Romano-British sites dating from the 1st-5th centuries AD (Figure 1, Table 1). Infants (birth to 1.0 year) were excluded from analysis due to the scarcity of erupted teeth within this age group. The following age categories defined by Lewis (2002) were used: 1.1-2.5 years, 2.6-6.5 years, 6.6-10.5 years, 10.6-14.5 years, 14.6-17.0 years. Age ranges serve to minimise age estimation error, and allow for cross-site comparison while corresponding with developmental milestones (Lampl and Johnston, 1996; Scheuer and Black, 2004; Lewis, 2010).

Methods

Individuals were aged based on their dental development either through macroscopic assessment or, wherever possible, using radiographs (Moorress et al., 1963a,b). As non-adults were not sexed, male and female tooth development stages were averaged. The cut-off point of 17.0 years was assigned when the roots of the third mandibular molar were complete but the apex remained open, giving an average age of 16.9 years old (Smith, 1991). When the apices of the available teeth were closed, an upper age estimate was derived using skeletal

maturation and diaphyseal lengths (Ubelaker, 1989; Scheuer and Black, 2000; 2004). Individuals were excluded once the femoral head was fully fused, with completion estimated between 14-17 years in females and 16-19 years for males (Scheuer and Black, 2000; 2004). It is recognised that the cut-off age of 17.0 years reflects a biological rather than chronological age, however it is based on the oldest, most accurate age estimate we currently have for non-adults (Lampl and Johnston, 1996). It is of course recognized that a biological age of 17.0 years would not necessarily have marked the end of childhood in Roman Britain. The lifecourse of Romano-British children is not fully explored, we therefore cannot use skeletal ages or age groups according to chronological ages that marked transitions in the lifecourse we are yet to define.

Only erupted teeth were included in the analysis, reflecting their susceptibility to caries. For teeth within the jaw, eruption was defined as those teeth in occlusion. The identification of loose teeth as erupted involved an assessment of root development stages (Moorrees et al., 1963a,b) and eruption patterns (Ubelaker, 1989), in addition to any evidence of wear or dental calculus (Buikstra and Ubelaker, 1994). As different teeth have varying degrees of susceptibility to caries, they were defined by type: deciduous or permanent, posterior (deciduous and permanent molars, permanent premolars) or anterior (deciduous and permanent incisors and canines) and location (i.e. root or crown). Carious lesion frequencies were reported accordingly. Caries was recorded macroscopically when the lesion perforated the tooth enamel. Statistical analysis was performed using a Pearson's chi-square test to evaluate the prevalence of lesions between the urban and rural sites, with a confidence limit set at 99.5% ($p=0.005$). Percent caries rates are presented as both crude prevalence rates (CPR) for the number of individuals observed and affected, or more precisely as true prevalence rates (TPR) determined by the number of teeth observed over those affected. The caries correction factor advocated by Lukacs (1995) was not applied as antemortem tooth loss is low in non-adult samples and loss of teeth through attrition was unlikely.

Enamel hypoplasia was recorded macroscopically when the tooth enamel was disrupted by circular or linear defects on two or more teeth on opposite sides of the dentition (Goodman and Armelagos, 1985; Goodman and Rose, 1990). This was done to avoid recording defects that may have resulted from localised trauma. The relationship between enamel hypoplasia and caries was quantified by Yule's Q.

RESULTS

Entire sample

The prevalence of caries for each site is listed in Table 1. For all sites and in the deciduous and permanent dentition combined, 11.3% (n=49/433) of individuals displayed caries, with a TPR of 1.4% (n=90/6283 teeth). Overall 2.1% of posterior teeth were affected (n=79/3678 teeth) compared to only 0.1% (n=2/2586 teeth) of the anterior teeth, a significant difference ($X^2=51.29$, $p<0.001$, d.f.=1). No lesions at the cemento-enamel junction (or 'root caries') were reported. Carious lesions were predominantly found on the occlusal surface of posterior teeth (TPR 1.1%, n=39/3678 teeth). Fissure caries accounted for 49.4% (n=39/79 teeth) of lesions in posterior teeth.

When the deciduous and permanent caries rates were compared, TPR was marginally higher in the deciduous dentition at 1.5% compared to 1.1%, but not significantly so ($X^2=1.58$, d.f.=2) (Table 2).

Inter-settlement comparisons

Nine carious teeth in the minor urban cohort were derived from the Ancaster toddler with SECC (Bonsall et al., 2015). To prevent skewing of the data, this individual was removed from analysis. The prevalence of caries was highest in the major urban sites (CPR 14.8%, TPR 1.8%), followed by minor urban (CPR 10.3%, TPR 1.1%) and rural settlements (CPR 3.8%, TPR 0.3%). The rural TPR is significantly lower than those reported from the urban settlements ($X^2=13.66$, $p<0.005$, d.f.=2).

Lesion frequency in posterior teeth was significantly lower in the rural sites at TPR 0.6% (n=4/659 teeth; $X^2=12.34$, $p<0.005$, d.f.=2) compared to both the major and minor urban sites. Lesions in the anterior dentition were only found in two non-adults, both from a major urban site, affecting the deciduous canine in the maxillary dentition of a 2.6-6.5-year old from Kingsholm, Gloucester, and the mandibular deciduous canine of a 6.6-10.5-year old from the northern cemetery at Winchester (Table 2).

In the deciduous teeth, significantly more lesions were recorded in the major urban cohort (n=32/1075 teeth) at TPR 3.0% ($X^2=27.03$, $p<0.001$, d.f.=2). Whereas no significant differences were observed for caries in the permanent teeth (Table 2).

Interproximal caries affecting the crown surface, either mesially or distally, was observed in 29 posterior teeth or 1.0% of the total sample (n=29/3019 teeth), but only in the major and minor urban sites with a revised prevalence rate of 38.7% (n=29/75 teeth). Again,

buccal caries was only observed in these urban cohorts with a prevalence of 10.7% (n=8/75 teeth).

Age differences

The prevalence of caries increased with age albeit with a slight decrease in the 10.6-14.5 year age group (CPR 16.7%, TPR 1.3%) probably as a result of the shedding of the primary dentition. Caries was not apparent until 10.6 years in the rural sample, whereas lesions were observed in the youngest cohort (1.1-2.5 years) in the major urban sample. In the 6.6-10.5-year cohort, the frequency of carious lesions was statistically higher in the major urban sample at TPR 3.7% (n=20/539 teeth; $X^2=11.36$, $p<0.005$, d.f.=2) compared to both the minor urban and rural sites. Out of the total of 20 carious teeth within this major urban group, 17 (85.0%) were deciduous molars (Table 3).

Stress and caries

In the major urban group, 11.4% of teeth (n=309/2700 teeth) had dental enamel defects, compared to 9.6% (n=236/2460 teeth) in the minor urban cohort, and 4.5% (n=49/1087 teeth) in the rural sample. The rural rate was significantly lower than those from both urban contexts ($X^2=43.66$, $p<0.001$, d.f.=2). The higher rates of enamel hypoplasia observed in major urban non-adults (TPR 11.4%) also matched the elevated rates of caries in these settlements. Overall, 39.6% (n=19/48) of the children with caries also had enamel hypoplasia. The Yule's Q statistic of 0.75 indicates that a moderate to strong relationship between the two conditions existed. This co-occurrence was highest in major urban sites at 5.8% (n=11/189) and lowest in rural children at 1.3% (n=1/78), although these differences were not significant ($X^2=3.26$, d.f.=2) (Table 4).

DISCUSSION

This study provides the first large scale examination of carious lesion frequencies in non-adults from urban and rural settlements in Roman Britain. Caries is age-progressive and multifactorial but can provide valuable insights into the cultural habits that affected non-adult diet, particularly where the characteristics of eating and drinking in childhood across Roman Britain are still largely unknown. This study is limited by the scarcity of skeletal and dental material from rural sites, a common problem within Romano-British bioarchaeology, and due to limited sample sizes it was also not possible to analyse variations in diet over time.

However, caries affected 11.3% of the Romano-British children, or 1.4% of all erupted teeth. Lesions were more frequent in children from major urban sites, where 14.8% of non-adults presented with caries. This is substantially higher than the non-adult rate reported at the Dorchester *civitas* (CPR 3.6%, Lewis pers. comm.) which is thought to represent a major urban settlement. Instead the Dorchester rates are more comparable to the rural pattern (CPR 3.8%) (Tables 2 and 5). The major urban TPR of 1.8% is however similar to that reported from the Lankhills cemetery for the Winchester *civitas* (TPR 1.7%) (Clough and Boyle, 2010).

When the deciduous teeth were examined, lesions were statistically more frequent in the major urban sample (TPR 3.0%), the only cohort to show individuals with caries in the anterior deciduous dentition. Higher rates of lesions in the deciduous compared to permanent dentition might be expected due to reduced enamel hardness and longer exposure to the oral environment (Hunter et al., 2000; Halcrow et al., 2013), although soft and carbohydrate-rich weaning foods further elevate early childhood caries (Temkin, 1991; Garnsey, 1999). Hence, the greater frequencies of deciduous caries in the major urban group suggest a prolonged exposure to cariogenic foods during early childhood (Veerkamp and Weerheijm, 1995; Berkowitz, 2003; Freeman and Stevens, 2008), while the presence of anterior deciduous caries may hint at the use of pacifiers (Azevedo et al., 2005). The significantly lower rate of deciduous caries in the rural settlements suggests a different early childhood diet or food processing techniques. Fewer carious lesions in the rural cohort may suggest that this part of the population were restricted in their consumption of agricultural products which are high in carbohydrates. Perhaps this reflects the primary role of rural inhabitants in this period who were expected to produce a surplus to provide for urban dwellers, the army and the elite (de la Bédoyère 1993, 86; Jones 1996, 208; MacMullen 1987; Whittaker and Garnsey 1997), where rural families would have been hesitant to consume their primary source of income. In turn, rural weaning foods may have been chosen that were less economically important and also less cariogenic. Differences in the archaeobotanical record between rural sites, particularly in the southwest where the sample sites of the current study are located, and urban settlements across Roman Britain were observed by van der Veen and colleagues (2008), and strengthen the argument for biased food allocation between producer and consumer. A diet high in grit, with hard and abrasive fibrous foods or contaminants may have also been consumed in rural areas, limiting caries development (Duray, 1992; Moynihan, 2000). Further research into dental wear in the urban and rural groups may help elucidate this

pattern.

While it is important to compare the results from this study with those that have gone before, this is challenging. Many of the studies that have discussed caries rates in Romano-British non-adults have very small sample sizes, and merged data without specifying which sites they included (Table 5). However, data for carious lesion frequency in non-adults is available for Rome itself. Killgrove (2010) reported true prevalence rates for 0-10 year olds, and 11-20 year olds. At Casal Bertone, a cemetery located close to the walls of the city of Rome, caries prevalence rates were 3.2% (0-10 year olds) and 2.9% (11-20 year olds), compared to 2.2% (n=33/1488 teeth in 1.1-10.5 year olds) and 1.5% (n=16/1096 teeth in 10.6-17.0 year olds) in the major urban cohort of the current study. Lesions were less frequent at the cemetery at Castelia Europarco which was located in an agricultural area of the suburbs of Rome. Just as in this study, no caries was reported in children under 10 years old, however the sample was small (n=2/121 teeth) (Killgrove 2010). Although we cannot make inferences on the specific foods eaten in Rome and the major urban towns of Roman Britain from this comparison, dental disease rates from both locations highlight the pertinent differences in oral health between residents of the towns and country at the centre and the fringes of the Empire. Prowse and colleagues (2008) reported a TPR of 3.8% for carious lesions in the deciduous teeth of 1-12 year olds from the necropolis at Isola Sacra at urban Portus. Although the deciduous caries rates for major urban settlements (3.0%) in the current study is significantly higher than elsewhere in Roman Britain, it is below that of children living in Italy, even outside of Rome. Additional data for child caries rates comes from Kellis II in Egypt, where Shukrum and Molto (2009) reported a TPR of 8.9% for deciduous teeth and TPR 3.8% for the permanent dentition, again substantially higher than seen in the British province. Overall, refined carbohydrates and a more fortified diet may have been more readily available to major urban residents in Roman Britain, compared to those living in the countryside. Yet, access to rich foods was more restricted than in Rome, the Italian urban centres and other provinces closer to the centre of the Empire. However, we also have to assume that genetic and geographic factors, such as immunity and fluoride exposure may have influenced caries susceptibility between these different populations.

Although the exact pathophysiology of a relationship between enamel hypoplasia and carious lesion development remains to be explored, a total of 39.6% of children with caries also displayed dental enamel hypoplasia. The result was anticipated, and is consistent with the caries-promoting function of stress. Stress may have elevated salivary cortisol levels

which in turn suppressed salivary IgA and enabled caries bacteria to spread (Boyce et al., 2010). Not only will these children have experienced a greater spread of cariogenic bacteria, but the quality of hypoplastic teeth would have exacerbated this effect. In the affected children, bacteria adhere to the site of the defect, and the thinned or defective enamel is more acid soluble, both allowing for caries to progress quicker (Hong et al., 2009). Co-occurrence of caries and enamel defects was more frequent in the urban cohorts. More bacteria in the oral cavity, and thinned enamel as a result of non-specific stress may have contributed to the progression of caries in those children. Since hypoplastic enamel defects were more frequent in the major urban cohort than in the rural children, both diet and non-specific stress may have contributed to the significant differences in carious lesion frequencies. This would suggest that higher status of the major urban children was not necessarily a given.

The youngest individual reported with carious lesions was a 1.1-2.5-year old from Butt Road, Colchester with occlusal caries on the deciduous maxillary second molar. The rise in caries TPRs from the youngest age group may reflect the timing of colonisation of the oral cavity with *S. mutans*. Carious lesions usually appear between 13-16 months after initial colonisation (Kawashita et al., 2011), suggesting the children in this study were affected at around 1 year of age. However, the administration of cariogenic foods and feeding habits may have occurred earlier, for instance between 6-12 months when infants start to sit up, develop better chewing and tasting mechanisms, and are eventually able to self-feed (Sheridan, 1991; Sellen, 2001; 2007; Carruth and Skinner, 2002; Delaney and Arvedson, 2008). Neither human breastmilk nor cow's milk are very cariogenic, but their caries-promoting properties are significantly increased when the infant is fed supplementary foods rich in carbohydrates such as fruits and honey (Moynihan, 2000; Azevedo et al., 2005; Kawashita et al., 2011). Today, *S. mutans* are mainly transmitted from the primary caregiver to the infant as soon as tooth surfaces have erupted (Gussy et al., 2006). In Rome, early child care was not only provided by the mother, but also by a *nutrix* or *paedagogus* among richer families (McWilliam, 2013), or by neighbours, friends and relatives for working women (Rawson, 2003a; Golden, 2011). It is likely similar practices were in place in Britain. For example, caries rates in Romano-British females are reported as 5.4% at Cannington (Brothwell et al., 2000), 3.9% at Butt Road, Colchester (Pinter-Bellows, 1993), and 5.4% at Bath Gate, Cirencester (Wells, 1982), higher than those observed in the non-adults from the same sites (TPRs of 0.4%, 1.8% and 1.0% respectively). The mechanism of *S. mutans* transmission may show that Romano-British care givers pre-chewed or tasted their infants'

food, and shared utensils with them when administering food or drink (Fildes, 1986). Historical evidence for pre-mastication refers to it as a practice to be avoided (Bradley, 1986; Temkin, 1991), which indicates that it might have indeed occurred, and probably had an influence on the transmission of oral infections from caregiver to child.

Lesion frequencies were higher in major urban and minor urban children from 10.6-17.0 years, at almost four times the rate reported for the rural sites. It is possible that this pattern is reflecting a richer diet of fortified foods for these adolescents. The fact that the majority of carious lesions in both the major urban and minor urban samples were interproximal and occlusal further attests to a softer and more refined diet (Vodanović et al., 2005). There is archaeological evidence for the fine milling of cereals into white flour (Cool, 2006; Alcock, 2010) and following a Roman diet, refined sugar may have been accessed in the form of honey and syrups (Moore and Corbett, 1973; Bowman and Thomas, 1994; Cool, 2006; Carreck, 2008; Crane, 2013). Honey has antibacterial properties which inhibit *S. mutans* growth, but whether honey is primarily cariogenic or cario-protective is debatable (Bogdanov et al., 2008; Nassar et al., 2012). However, it would have been the main sweetener in this time period and therefore frequently consumed. Grape syrup known as *defrutum*, *sapa* or *caroenum* was made in Spain or southern Gaul and was shipped to Britain in distinctive amphorae (Cool, 2006). Evidence for these amphorae in Britain is scarce and dates mainly to the 1st and 2nd centuries AD. These syrups were therefore either imported in different vessels in the later centuries, became even more difficult to get hold of or alternatively, ceased to be available in Roman Britain (Sealey and Davies, 1984; Sealey and Tyers, 1989; Monfort, 1998). However, *defrutum* could be made from boiling down any fruit juice in lead vessels, and would therefore have been widely available across Roman Britain (Farwell and Molleson, 1993; Roberts and Cox, 2003). In summary, carious lesion frequencies in children aged older than 10.6 years attest dietary differences between those living in the towns and country, probably linked with the availability of more refined and softer foods in the urban centres.

CONCLUSIONS

This research has demonstrated the value of reporting carious lesion frequency in non-adults. Its use is further increased by considering the location and type of carious lesions, and the presence of early childhood stress. Higher rates of dental disease in urban environments

suggest differences in diet between children in the countryside and the towns of *Britannia*. Elevated stress, measured as the prevalence of enamel hypoplasia, in the children growing up in urban settlements was accompanied by higher carious lesion frequency. However, we cannot currently estimate how exact the relationship between stress and enamel defects is to the incidence and severity of caries. Caries in the deciduous dentition allows for inferences to be made on weaning and feeding practices that promote dental decay, such as the sharing of utensils and food between caregiver and child. The lower frequency of carious lesions in the rural sample suggests a simpler diet consumed by these children, possibly as a result of biased food allocation in response to economic pressures. Urban populations may have had more access to processed and sweetened foodstuffs whereby refined carbohydrates were more readily available.

ACKNOWLEDGEMENTS

This research was funded by an AHRC Studentship. We thank the curators who provided access to the skeletal collections for this project: Robert Kruszynski, Natural History Museum London; Keith Fitzpatrick-Matthews, North Hertfordshire District Council Museums Service; Simon Mays, English Heritage; Christiane Jeuckens, Oxfordshire Museums Service; Jo Buckberry, Biological Anthropology Research Centre; Steve Minnitt, Somerset Heritage Centre; David Allen, Hampshire Arts and Museums Service; Sarah Wilson, Peterborough Museum; Helen Rees, Winchester Museum Service; Lorraine Cornwell, Rutland County Museum; Alison Brooks and Heather Dawson, Corinium Museum; Paul Sealey, Colchester and Ipswich Museums and Timothy Vickers, Luton Culture. The authors thanks the anonymous reviewers and editor for their constructive comments which have greatly improved an earlier draft of this paper.

LITERATURE CITED

Afroughi S, Faghihzadeh S, Khaledi MJ, Motlagh MG. 2010. Dental caries analysis in 3–5 years old children: a spatial modelling. *Arch Oral Biol* 55:374-378.

Aligne C, Moss M, Auinger P, Weitzman M. 2003. Association of pediatric dental caries with passive smoking. *JAMA* 289:1258-1264.

- Alcock JP. 2010. *Food in Roman Britain*. Stroud: The History Press.
- Azevedo TDPL, Bezerra ACB, de Toledo OA. 2005. Feeding habits and severe early childhood caries in Brazilian preschool children. *Pediatr Dent* 27:28-33.
- Bagain Z, Newman L, Hyde N. 2004. How serious are oral infections?. *J Laryngol Otol* 118:561-565.
- Bennion E. 1986. *Antique Dental Instruments*. London: Sotheby's.
- Berkowitz RB. 2003. Causes, treatment and prevention of early childhood caries: a microbiologic perspective. *J Can Dent Assoc* 69:304-307.
- Bogdanov S, Jurendic T, Sieber R, Gallmann P. 2008. Honey for nutrition and health: a review. *J Am Coll Nutr* 27:677-689.
- Bonfiglioli B, Brasili P and Belcastro MG. 2003. Dento-alveolar lesions and nutritional habits of a Roman Imperial age population (1st-4th c. AD): Quadrella (Molise, Italy). *HOMO* 54:36-56.
- Bonsall L, Ogden AR, Mays S. 2015. A case of early childhood caries (ECC) from late Roman Ancaster, England. *Int J Osteoarch*. DOI: 10.1002/oa.2441.
- Bowman AK, Thomas JD. 1994. *The Vindolanda Writing Tablets (Taulae Vindolandenses II)*. London: British Museum Press.
- Boyce WT, Den Besten PK, Stamperdahl J, Zhan L, Jiang Y, Adler NE, Featherstone JD. 2010. Social inequalities in childhood dental caries: the convergent roles of stress, bacteria and disadvantage. *Soc Sci Med* 71:1644-1652.
- Bradley KR. 1984. *Slaves and masters in the Roman Empire – a study in social control*. Oxford: Oxford University Press.
- Bradley KR. 1986. Wet-nursing at Rome: a study in social relations. In: Rawson B, editor. *The family in ancient Rome – new perspectives*. London: Routledge. p 201-229.
- Bradley KR. 1991. *Discovering the Roman family – studies in Roman social history*. Oxford: Oxford University Press.
- Bradley K. 1994. *Slavery and society at Rome – key themes in ancient history*. Cambridge: Cambridge University Press.

- Bradley K. 2005. The Roman child in sickness and in health. In: George M, editor. The Roman family in the Empire. Oxford: Oxford University Press. p 67-92.
- Breeze DJ (ed.). 2014. The Impact of Rome on the British Countryside. A Conference Organised by the Royal Archaeological Institute, Chester, 11-13 October 2013. London: Royal Archaeological Institute Supplementary Publication.
- Buikstra JE, Ubelaker DH (eds.). 1994. Standards for data collection from human skeletal remains. Fayetteville: Arkansas Archaeological Survey Research Series Number 44.
- Burleigh GR, Fitzpatrick-Matthews KJ. 2010. Excavations at Baldock, Hertfordshire, 1978-1994 – Volume 1 – An Iron Age and Romano-British cemetery at Wallington Road. Hitchin: North Hertfordshire District Council Museums Service and North Hertfordshire Archaeological Society.
- Burnham BC. 1993. The small towns of Roman Britain – the last fifty years. In Greep SJ, editor. Roman towns: The Wheeler inheritance. York: CBA Research Report 93. p 99-110.
- Burnham BC. 1995. Small towns: the British perspective. In: Brown AE, ed. Roman small towns in eastern England and beyond. Oxford: Oxbow Monograph 52. p 7-18.
- Burnham BC, Wachter J. 1990. The small towns of Roman Britain. London: Batsford.
- Burnham BC, Collis J, Dobinson C, Haselgrove C, Jones M. 2001. Themes for urban research, c 100 BC to AD 200. In: James S, Millett M, eds. Britons and Romans: advancing an archaeological agenda. York: CBA Research Report 125. p 67-76.
- Byun R, Nadkarni MA, Chhour K, Martin FE, Jacques NA, Hunter N. 2004. Quantitative analysis of diverse *Lactobacillus* species present in advanced dental caries. *J Clin Microbiol* 42:7, 3128-3136.
- Caffell A, Holst M. 2007. Osteological Analysis, Horncastle, East Lincolnshire. York: Unpublished Skeletal Report by York Osteoarchaeology. Commissioned by Lindsey Archaeological Services.
- Carreck NL. 2008. Are honey bees (*Apis mellifera* L.) native to the British Isles?. *J Apic Res* 47(4): 318-322.
- Carruth BR, Skinner JD. 2002. Feeding behaviours and other motor development in healthy children (2-24 months). *J Am Coll Nutr* 21: 88-96.

- Chambers RA. 1987. The late and sub-Roman cemetery at Queenford Farm, Dorchester-on-Thames, Oxon. *Oxoniensia* 52: 35-70.
- Chenery C, Müldner G, Evans J, Eckardt H, Lewis M. 2009. Strontium and stable isotope evidence for diet and mobility in Roman Gloucester, UK. *J Archaeol Sci* 37: 150-163.
- Cheung C, Schroeder H, Hedges REM. 2012. Diet, social differentiation and social change in Roman Britain: new isotopic evidence from Gloucestershire. *Archaeol Anthropol Sci* 4: 61-73.
- Clarke M, Locker D, Berall G, Pencharz P, Kenny DJ, Judd P. 2006. Malnourishment in a population of young children with severe early childhood caries. *Pedr Dent* 28: 2524-259.
- Clough S, Boyle A. 2010. Inhumations and disarticulated human bone. In: Booth P, Simmonds A, Boyle A, Clough S, Cool HEM, Poore D. *The late Roman cemetery at Lankhills, Winchester – excavations 2000-2005*. Oxford: Oxford Archaeology. p 339-399.
- Collis JR. 1968. Excavations at Owslebury, Hampshire: an interim report. *Ant J* 48: 18-31.
- Collis J. 1977. Owslebury (Hants) and the problem of burials on rural settlements. In: Reece R, ed. *Burial in the Roman World*. York: CBA Research Report 22. p 26-34.
- Cook DC, Buikstra JE. 1979. Health and differential survival in prehistoric populations: prenatal dental defects. *Am J Phys Anthropol* 51: 649-664.
- Cool HEM. 2006. *Eating and drinking in Roman Britain*. Cambridge: Cambridge University Press.
- Cox M. 1989. *The human bones from Ancaster*. London: Unpublished Ancient Monuments Laboratory Report 93/89.
- Craig OE, Biazzo M, O'Connell T, Garnsey P, Martinez-Labarga C, Lelli R, Salvadei L, Tartaglia G, Nava A, Reno L, Fiammenghi A, Rickards O, Bondioli L. 2009. Stable isotopic evidence for diet at the Imperial Roman coastal site of Velia (1st and 2nd centuries AD) in Southern Italy. *Am J Phys Anthropol* 139, 572-583.
- Crane EE. 2013. *The world history of beekeeping and honey hunting*. London: Routledge.
- Crummy N, Crossan C. 1993. Excavations at Butt Road Cemetery, 1976-9, 1986 and 1989. In: Crummy N, Crummy P, Crossan C. *Colchester Archaeological Report 9: Excavations of*

Roman and later cemeteries, churches and monastic sites in Colchester, 1971-88. Colchester: Colchester Archaeological Trust. p 4-163.

Cummings C. 2009. Meat consumption in Roman Britain: the evidence from stable isotopes. In: Driessen M, Heeren S, Hendriks J, Kemmers F, Visser R, eds. TRAC 2008: Proceedings of the eighteenth annual Theoretical Roman Archaeology Conference, Oxford. Oxford: Oxbow Books. p 73-83.

Dasen V. 2011. Childbirth and infancy in Greek and Roman Antiquity. In: Rawson B, ed. A companion to families in the Greek and Roman worlds. Chichester: Wiley-Blackwell. p 291-314.

de la Bédoyère G. 1993. English Heritage book of Roman villas and the countryside. London: B T Batsford/English Heritage.

Delaney AL, Arvedson JC. 2008. Development of swallowing and feeding: prenatal through first year of life. *Dev Dis Res Rev* 14: 105-117.

Dix B. 1983. Ashton Roman town. *SMidIA* 13: 18-20.

Dobney K. 2001. A place at the table: the role of vertebrate zooarchaeology within a Roman research agenda for Britain. In James S, Millett M, eds. *Britons and Romans: advancing an archaeological agenda*. York: CBA Research Report 125. p 36-45.

Duray SM. 1990. Deciduous enamel defects and caries susceptibility in a prehistoric Ohio population. *Am J Phys Anthropol* 81: 27-34.

Duray SM. 1992. Enamel defects and caries etiology: an historical perspective. In: Goodman AH, Capasso LL, eds. *Recent contributions to the study of enamel developmental defects*. *J Paleopathol Monographic Publications* 2. p 307-320.

Durham B, Rowley T. 1972. A cemetery site at Queensford Mill, Dorchester. *Oxoniensia* 37: 32-38.

Eckardt H. 1999. The Colchester 'child's grave'. *Britannia* 30: 57-90.

Faith R. 1997. *The English peasantry and the growth of lordship*. London: Leicester University Press.

Farwell DE, Molleson TI. 1993. *Poundbury Volume 2 – The Cemeteries*. Dorchester: Dorset Natural History and Archaeological Society Monograph Series 11.

- Fejerskov O, Bilde PG, Bizzarro M, Connelly JN, Thomsen JS, Nyvad B. 2012. Dental caries in Rome, 50-100 AD. *Caries Res* 46: 467-473.
- Fewtrell MS, Morgan JB, Duggan C, Gunnlaugsson G, Hibberd PL, Lucas A, Kleinman RE. 2007. Optimal duration of exclusive breastfeeding: what is the evidence to support current recommendations? *Am J Clin Nutr* 85(2): 635-638.
- Fildes V. 1981. The early history of the infant feeding bottle – I. *Nurs Times* 15: 128-129.
- Fildes VA. 1986. *Breasts, bottles and babies – a history of infant feeding*. Edinburgh: Edinburgh University Press.
- Fildes V. 1988. *Wet nursing – a history from Antiquity to the Present*. Oxford: Basil Blackwell.
- Fischer K. 1987. Nochmals ubuppa und tit(t)ina. *Philologus* 131(1-2): 156-157.
- Fischman SL. 1997. The history of oral hygiene products: how far have we come in 6000 years? *J Periodontol* 2000 15: 7-14.
- Freeman R, Stevens A. 2008. Nursing caries and buying time: an emerging theory of prolonged bottle feeding. *Community Dent Oral Epidemiol* 36: 425-433.
- Fuller BT, Molleson TI, Harris DA, Gilmour LT, Hedges REM. 2006. Isotopic evidence for breastfeeding and possible adult dietary differences from Late/Sub-Roman Britain. *A J Phys Anthropol* 129: 45-54.
- Garcin V, Velemínský P, Trefný P, Alduc-Le Bagousse A, Lefebvre A, Bruzek J. 2010. Dental health and lifestyle in four early mediaeval juvenile populations: comparisons between urban and rural individuals, and between coastal and inland settlements. *HOMO* 61(6): 421-439.
- Garnsey P. 1991. Child rearing in Ancient Italy. In: Kertzer DI and Saller RP, eds. *The family in Italy – from Antiquity to the Present*. New Haven: Yale University Press. p 48-65.
- Garnsey P. 1999. *Food and society in Classical Antiquity*. Cambridge: Cambridge University Press.
- Garnsey P, Saller R. 1987. *The Roman Empire – economy, society and culture*. Berkeley: University of California Press.

- Golden M. 2011. Other people's children. In: Rawson B, ed. *A companion to families in the Greek and Roman worlds*. Chichester: Wiley-Blackwell. p 262-275.
- Goodman PJ. 2007. *The Roman city and its periphery – From Rome to Gaul*. London: Routledge.
- Goodman AH, Armelagos GJ. 1985. Factors affecting the distribution of enamel hypoplasias within the human permanent dentition. *Am J Phys Anthropol* 68: 479-493.
- Goodman AH, Rose JC. 1990. Assessment of systemic physiological perturbations from dental enamel hypoplasias and associated histological structures. *Am J Phys Anthropol* 33 (S11), 59-110.
- Griffin R, Pitts M. 2011. Inequality at late Roman Baldock, UK – the impact of social factors on health and diet. *J Anthropol Res* 67: 533-556.
- Grocock C, Grainger S. 2006. *Apicius*. Totnes: Prospect Books.
- Gussy MG, Waters EG, Walsh O, Kilpatrick NM. 2006. Early childhood caries: current evidence for aetiology and prevention. *J Paediatr Child Health* 42: 37-43.
- Halcrow SE, Harris NJ, Tayles N, Ikehara-Quebral R, Pietrusewsky M. 2013. From the mouths of babes: dental caries in infants and children and the intensification of agriculture in mainland Southeast Asia. *Am J Phys Anthropol* 150: 409-420.
- Hallet KB, O'Rourke PK. 2003. Social and behavioural determinants of early childhood caries. *Aust Dent J* 48(1): 27-33.
- Harlow M, Laurence R. 2002. *Growing up and growing old in Ancient Rome – a life course approach*. London: Routledge.
- Harlow M, Laurence R. 2007. Age and ageing in the Roman Empire. In: Harlow M, Laurence R, eds. *Age and ageing in the Roman Empire*. Portsmouth: *J Roman Archaeol Supplementary Publications* 65. p 9-24.
- Heighway CM. 1980. Roman Cemeteries in Gloucester District. *Trans BGAS* 98: 57-72.
- Hemelrijk EA. 1999. *Matrona Docta – educated women in the Roman elite from Cornelia to Julia Domna*. London: Routledge.
- Hillson S. 1996. *Dental Anthropology*. Cambridge: Cambridge University Press.

- Hingley R. 1989. *Rural Settlement in Roman Britain*. London: Seaby.
- Hong L, Levy SM, Warren JJ, Broffitt B. 2009. Association between enamel hypoplasia and dental caries in primary second molars: a cohort study. *Caries Res* 43: 345-353.
- Humphrey LT. 2010. Weaning behaviour in human evolution. *Sem Cell Dev Biol* 21(4): 453-461.
- Hunter ML, West NX, Hughes JA, Newcombe RG, Addy M. 2000. Erosion of deciduous and permanent dental hard tissue in the oral environment. *J Dent* 28(4): 257-263.
- Hurst HR. 1985. *Kingsholm – Excavations at Kingsholm Close and Other Sites with a Discussion of the Archaeology of the Area*. Stroud: Gloucester Archaeological Publications.
- Hurst HR. 1986. *Gloucester – the Roman and later defences*. Stroud: Gloucester Archaeological Publications.
- Jackson R. 1988. *Doctors and diseases in the Roman Empire*. London: British Museum Publications.
- Jones ME. 1996. *The end of Roman Britain*. London: Cornell University Press.
- Katzenberg MA, Herring DA, Saunders SR. 1996. Weaning and infant mortality: evaluating the skeletal evidence. *Yearb Phys Anthropol* 39: 177-199.
- Katzenberg MA. 2012. The ecological approach: understanding past diet and the relationship between diet and disease. In: Grauer AL, ed. *A companion to paleopathology*. Chichester: Wiley-Blackwell. p 97-113.
- Kawashita Y, Kitamura M, Saito T. 2011. Early childhood caries. *Int J Dent* 2011: 1-7.
- Killgrove K and Tykot R H, 2013. Food for Rome: a stable isotope investigation of diet in the Imperial period (1st-3rd centuries AD). *J Anthropol Archaeol* 32 (1), 28-38
- King AC. 1984. Animal bones and the dietary identity of military and civilian groups in Roman Britain, Germany and Gaul. In: Blagg T, King A, eds. *Military and civilian in Roman Britain: cultural relationships in a frontier province*. Oxford: BAR British Series 136. p 187-217.
- King A. 1999. Diet in the Roman world: a regional inter-site comparison of the mammal bones. *J Roman Archaeol* 12: 168-202.

- King A. 2001. The Romanization of diet in the Western Empire: comparative archaeological studies. In: Keay S, Terrenato N, eds. *Italy and the West – comparative issues in Romanization*. Oxford: Oxbow Books. p 210-226.
- Laes C. 2007. Inscriptions from Rome and the history of childhood. In: Harlow M, Laurence R, eds. *Age and ageing in the Roman Empire*. Portsmouth: J Roman Archaeol Supplementary Publications 65. p 25-36.
- Laes C. 2011. *Children in the Roman Empire – outsiders within*. Cambridge: Cambridge University Press.
- Lampl M, Johnston FE. 1996. Problems in the ageing of skeletal juveniles: perspectives from maturation assessments of living children. *Am J Phys Anthropol* 101: 345-355.
- Larsen CS. 1997. *Bioarchaeology – Interpreting Behavior from the Human Skeleton*. Cambridge: Cambridge University Press.
- Laurence R, Cleary SE, Sears G. 2011. *The city in the Roman west c.250 BC – c.AD 250*. Cambridge: Cambridge University Press.
- Leech R. 1982. *Excavations at Catsgore 1970-1973 – a Romano-British village*. Taunton: Western Archaeological Trust Excavation Monograph 2.
- Leech R, Besly EM, Everton RF. 1981. The excavation of a Romano-British farmstead and cemetery on Bradley Hill, Somerton, Somerset. *Britannia* 12: 177-252.
- Lewis ME. 2002. Impact of Industrialization: comparative study of child health in four sites from Medieval and Postmedieval England (A.D. 850-1859). *Am J Phys Anthropol* 119: 211-223.
- Lewis ME. 2010. Life and death in a civitas capital: metabolic disease and trauma in the children from late Roman Dorchester, Dorset. *Am J Phys Anthropol* 142: 405-416.
- Locker A. 2007. In piscibus diversis, the bone evidence for fish consumption in Roman Britain. *Britannia* 38: 141-180.
- Lukacs JR. 1995. The ‘Caries Correction Factor’: a new method of calibrating dental caries rates to compensate for antemortem loss of teeth. *Int J Osteoarchaeol* 5: 151-156.
- Marter A, Agruss JC. 2007. Pacifiers: an update on use and misuse. *J Spec Pediatr Nurs* 12(4): 278-285.

- Matthews CL. 1981. A Romano-British inhumation cemetery at Dunstable. *Beds Arch J* 15.
- Mattingly DJ. 1997. Beyond belief? Drawing a line beneath the consumer city. In: Parkins HM, ed. *Roman urbanism – beyond the consumer city*. London: Routledge. p 210-218.
- Mattingly D. 2006. *An Imperial possession – Britain in the Roman Empire*. London: Penguin Group.
- McCarthy M. 2013. *The Romano-British peasant – towards a study of people, landscapes and work during the Roman occupation of Britain*. Oxford: Oxbow Books.
- MacMullen R, 1987. Late Roman slavery. *Historia* 36(3): 359-382.
- McConnell D, Grassam A, Mustchin A. 2012. Land adjacent to Great Casterton Primary School, Pickworth Road, Great Casterton, Rutland. Bury St. Edmunds: Archaeological Solutions Limited Unpublished Report.
- McDade TW, Worthman CM. 1998. The weanling's dilemma reconsidered: a biocultural analysis of breastfeeding ecology. *J Dev Behav Pediatr* 19(4): 286-299.
- McWilliam J. 2013. The socialization of Roman Children. In: Evans Grubbs J, Parkin T, eds. *The Oxford handbook of childhood and education in the Classical world*. Oxford: Oxford University Press. p 264-285.
- Mensforth RP, Lovejoy CO, Lallo JW, Armelagos GJ. 1978. The role of constitutional factors, diet and infectious disease in the aetiology of porotic hyperostosis and periosteal reactions in prehistoric infants and children. *Med Anthropol* 2: 1-59.
- Millett M. 1990. *The Romanization of Britain – an essay in archaeological interpretation*. Cambridge: Cambridge University Press.
- Millett M. 1995. Strategies for Roman small towns. In: Brown AE, ed. *Roman small towns in eastern England and beyond*. Oxford: Oxbow Monograph 52. p 29-38.
- Millett M. 1999. Coloniae and Romano-British studies. In: Hurst H, ed. *The Coloniae of Roman Britain: new Studies and a review*. Portsmouth: J Roman Archaeol Supplementary Series 36. p 191-196.
- Millet M. 2001. Approaches to urban societies. In: James S, Millett M, eds. *Britons and Romans: advancing an archaeological agenda*. York: CBA Research Report 125. p 60-66.

- Monfort CC. 1998. Britannia and the imports of Baetican and Lusitanian amphorae. *J Iber Archaeol*, 159-172.
- Moore WJ, Corbett E. 1973. The distribution of dental caries in ancient British populations – II. Iron Age, Romano-British and Mediaeval Periods. *Caries Res* 7: 139-153.
- Moorrees CFA, Fanning EA, Hunt EE. 1963a. Age variation of formation stages for ten permanent teeth. *J Dent Res* 42: 1490-1502.
- Moorrees CFA, Fanning EA, Hunt EE. 1963b. Formation and resorption of three deciduous teeth in children. *Am J Phys Anthropol* 21: 205-213.
- Morley N. 1997. Cities in context: urban systems in Roman Italy. In: Parkins HM, ed. *Roman urbanism: beyond the consumer city*. London: Routledge. p 42-58.
- Mouritsen H. 2011. The families of Roman slaves and freedmen. In Rawson B, ed. *A companion to families in the Greek and Roman worlds*. Chichester: Wiley-Blackwell. p 129-144.
- Moynihan P. 2000. Foods and factors that protect against dental caries. *Nutr Bull* 25: 281-286.
- Mullin D. 2006. The archaeological landscape of Frampton-on-Severn, Gloucestershire: updated project design. Stroud: Gloucester Archaeology Service.
- Müldner G. 2013. Stable isotopes and diet: their contribution to Romano-British research. *Antiquity* 87: 137-149.
- Nassar HM, Li M, Gregory RL. 2012. Effect of honey on *Streptococcus mutans* growth and biofilm formation. *Appl Environ Microbiol* 78(2): 536-540.
- Nehlich O, Fuller BT, Jay M, Mora A, Nicholson RA, Smith CI, Richards MP. 2011. Application of sulphur isotope ratios to examine weaning patterns and freshwater fish consumption in Roman Oxfordshire, UK. *Geochim Cosmochim Acta* 75: 4963-4977.
- Nield L, Stenger J, Kamat D. 2008. Common pediatric dental dilemmas. *Clin Pediatr* 47(2): 99-105.
- Ortner DJ. 2003. *Identification of Pathological Conditions in Human Skeletal Remains* (2nd edition). San Diego: Academic Press.

O'Sullivan EA, Williams SA, Wakefield RC, Cape JE and Curzon MEJ. 1993. Prevalence and site characteristics of dental caries in primary molar teeth from prehistoric times to the 18th century in England. *Caries Res* 27: 147-153.

Ottaway P. 2009. *Roman York*. Stroud: The History Press.

Ottaway PJ, Qualmann KE, Rees H, Scobie GD. 2012. *The Roman cemeteries and suburbs of Winchester – excavations 1971-86*. Winchester: Winchester Museums and English Heritage.

Parkins HM. 1997. The “consumer city” domesticated? The Roman city in elite economic strategies. In: Parkins HM, ed. *Roman urbanism – beyond the consumer city*. London: Routledge. p 83-111.

Pearce J. 2008. Burial evidence from Roman Britain – the un-numbered dead. *Pour une archaéologie du rite. Nouvelles perspectives de l'archaéologie funéraire, études réunies par John Scheid*. Paris: Collection de L'Ecole Française de Rome. p 29-42.

Pitts M, Griffin R. 2012. Exploring health and social well-being in Late Roman Britain: an intercemetary approach. *Am J Archaeol* 116(2): 253-276.

Pitts M, Perring D. 2006. The making of Britain's first urban landscapes: the case of late Iron Age and Roman Essex. *Britannia* 37: 189-212.

Powell ML. 1985. The analysis of dental wear and caries for dietary reconstruction. In: Gilbert RI, Mielke JH, eds. *The analysis of prehistoric diets*. Orlando: Academic Press. p 307-388.

Powell LA, Redfern RC, Millard AR, Gröcke DR. 2014. Infant feeding practices in Roman London: evidence from isotopic analyses. In: Carroll M, Graham E, eds. *Infant health and death in Roman Italy and beyond*. Portsmouth: J Roman Archaeol Supplementary Series 96. p 89-110.

Prowse TL. 2011. Diet and dental health through the life course in Roman Italy. In: Agarwal SC, Glencross BA, eds. *Social bioarchaeology*. Oxford: Blackwell Studies in Global Archaeology 15. p 410-437.

Prowse TL, Saunders SR, Schwarcz HP, Garnsey P, Macchiarelli R, Bondioli L. 2008. Isotopic and dental evidence for infant and young child feeding practices in an Imperial Roman skeletal sample. *Am J Phys Anthropol* 137: 294-308.

- Rahtz P, Hirst S, Wright SM. 2000. Cannington Cemetery – Excavations 1962-3 of Prehistoric, Roman, post-Roman, and later features at Cannington Park Quarry, near Bridgwater, Somerset. London: Society for the Promotion of Roman Studies, Britannia Monograph Series 17.
- Rawson B. 1966. Family life among the lower classes at Rome in the first two centuries of the Empire. *Class Philol* 61(2): 71-83.
- Rawson B. 1986a. Children in the Roman familia. In: Rawson B, ed. *The family in ancient Rome – new perspectives*. London: Routledge. p 170-200.
- Rawson B. 1986b. The Roman family. In: Rawson B, ed. *The family in ancient Rome – new perspectives*. London: Routledge. p 1-57.
- Rawson B. 1991a. Adult-child relationships in Roman society. In: Rawson B, ed. *Marriage, divorce and children in ancient Rome*. Oxford: Oxford University Press. p 7-30.
- Rawson B. 1991b. *Marriage, divorce and children in ancient Rome*. Oxford: Clarendon Press.
- Rawson B. 2003a. *Children and childhood in Roman Italy*. Oxford: Oxford University Press.
- Rawson B. 2003b. The roman family in recent research: state of the question. *Bib Int* 11(2): 119-138.
- Rawson B. 2005. Circulation of staff between Roman households. *Z Papyrologie Epigraphik* 151: 223-224.
- Redfern RC, Hamlin C, Beavan-Athfield N. 2010. Temporal changes in diet: a stable isotope analysis of late Iron Age and Roman Dorset, Britain. *J Archaeol Sci* 37: 1149-1160.
- Redfern RC, Millard AR, Hamlin C. 2012. A regional investigation of subadult dietary patterns and health in late Iron Age and Roman Dorset. *J Archaeol Sci* 39: 1249-1259.
- Redfern RC, DeWitte SN, Pearce J, Hamlin C, Dinwiddy KE. 2015. Urban-rural differences in Roman Dorset, England: a bioarchaeological perspective on Roman settlements. *Am J Phys Anthropol* 157: 107-120.
- Roberts C, Cox M. 2003. *Health and disease in Britain – from prehistory to the present day*. Stroud: Sutton Publishing.
- Rogers A. 2011. *Late Roman towns in Britain – rethinking change and decline*. Cambridge: Cambridge University Press.

- Rogers B, Arvedson J. 2005. Assessment of infant oral sensorimotor and swallowing function. *Ment Retard Dev D R* 11(1): 74-82.
- Rosenthal R. 1936. A short pictorial review of the evolution of infant-feeding vessels up to the beginning of the 19th century. *Bull Med Libr Assoc* 25(1-2): 89-94.
- Rutgers L V, van Strydonck M, Boudin M and van der Linde C, 2009. Stable isotope data from the early Christian catacombs of ancient Rome: new insights into the dietary habits of Rome's early Christians. *J Archaeol Sci* 36, 1127-1134
- Saller R. 1991. Corporal punishment, authority and obedience in the Roman household. In: Rawson B, ed. *Marriage, divorce and children in ancient Rome*. Oxford: Oxford University Press. p 144-165.
- Scheuer L, Black S. 2000. *Developmental juvenile osteology*. San Diego: Academic Press.
- Scheuer L, Black S. 2004. *The juvenile skeleton*. London: Elsevier Academic Press.
- Schuurs A. 2013. *Pathology of the hard dental tissues*. Chichester: Wiley-Blackwell.
- Schwartz JH. 2007. *Skeleton keys – an introduction to human skeletal morphology, development and analysis* (2nd edition). Oxford: Oxford University Press.
- Sealey PR, Davies GMR. 1984. Falernian wine at Roman Colchester. *Britannia* 15: 250-254.
- Sealey PR, Tyers PA. 1989. Olives from Roman Spain: a unique amphora find in British waters. *Antiq Journ* 69(1): 53-72.
- Sellen D W. 2001. Comparison of infant feeding patterns reported for nonindustrial populations with current recommendations. *J Nutr* 131: 2707-2715.
- Sellen DW. 2007. Evolution of infant and young child feeding: implications for contemporary public health. *Annu Rev Nutr* 27: 123-148.
- Sheridan MD. 1991. *From birth to five years*. Routledge: Routledge.
- Shukrum S, Molto J. 2009. Childhood oral health: dental pathology of Kellis II, Dakhleh, Egypt. In: Lewis M, Clegg M, eds. *Proceedings of the 9th Conference of the British Association for Biological Anthropology and Osteoarchaeology*. Oxford: Archaeopress. p 19-30.

- Sigismund-Nielsen H. 2007. Children for profit and pleasure. In: Harlow M, Laurence R, eds. *Age and ageing in the Roman Empire*. Portsmouth: J Roman Archaeol Supplementary Series 65. p 37-54.
- Smith BH. 1991. Standards of human tooth formation and dental age assessment. In: Kelley MA, Larsen CS, eds. *Advances in dental anthropology*. New York: Wiley Liss. p 143-168.
- Sreebny LM. 1983. Sugar and human dental caries. *World Rev Nutr Diet* 40: 19-65.
- Stead IM, Rigby V. 1986. Baldock: the excavation of a Roman and pre-Roman settlement, 1968-72. London: Society for the Promotion of Roman Studies.
- Taylor J. 2001. Rural society in Roman Britain. In: James S, Millett M, eds. *Britons and Romans: advancing an archaeological agenda*. York: CBA Research Report 125. p 46-59.
- Temkin O. 1991. *Soranus' Gynecology*. Baltimore: Johns Hopkins University Press.
- Todd M. 1975. Margidunum and Ancaster. In: Rodwell W, Rowley T, eds. *The 'Small Towns' of Roman Britain*. Oxford: BAR British Series 15. p 211-224.
- Touger-Decker R, van Loveren C. 2003. Sugars and dental caries. *Am J Clin Nutr* 78(4): 881-892.
- Ubelaker DH. 1989. *Human Skeletal Remains: Excavation, Analysis, Interpretation* (2nd edition). Washington: Taraxacum.
- Van der Veen M. 2008. Food as embodied material culture: diversity and change in plant food consumption in Roman Britain. *J Roman Archaeol* 21: 83-109.
- Van der Veen M, Livarda A, Hill A. 2007. The archaeobotany of Roman Britain: current state and identification of research priorities. *Britannia* 38: 181-210.
- Van der Veen M, Livarda A, Hill A. 2008. New plant foods in Roman Britain: dispersal and social access. *Environ Archaeol* 13: 11-36.
- Veerkamp JSJ, Weerheijm KL. 1995. Nursing-bottle caries: the importance of a developmental perspective. *J Dent Child* 62(6): 381-386.
- Viner L, Leech R. 1982. Bath Gate cemetery, 1969-1976. In: McWhirr, Viner L, Wells C. *Cirencester excavations II – Romano-British cemeteries at Cirencester*. Cirencester: Cirencester Excavation Committee. p 69-133.

Vodanović M, Brkić H, Šlaus M, Željko D, 2005. The frequency and distribution of caries in the mediaeval population of Bijelo Brdo in Croatia (10th-11th century). *Arch Oral Biol* 50: 669-680

Wacher J. 1974. *The Towns of Roman Britain*. London: Batsford.

Wenham LP. 1968. *The Romano-British cemetery at Trentholme Drive, York*. London: Ministry of Public Building and Works Archaeological Reports Number 5.

White R. 2014. *The Wroxeter Hinterland Project: exploring the relationship between country and town*. In: Breeze DJ, ed. *The impact of Rome on the British countryside. A conference organised by the Royal Archaeological Institute, Chester, 11-13 October 2013*, London: Royal Archaeological Institute Supplementary Publication. p 7-11.

Whitehead RG, Paul AA. 1984. Growth charts and the assessment of infant feeding practices in the western world and in developing countries. *Early Hum Dev* 9: 187-207.

Whittaker C R and Garnsey P, 1997. Rural life in the later Roman Empire. In: Cameron A, Garnsey P, eds. *The Cambridge Ancient History 13. The Late Empire, AD 337-425*. Cambridge: Cambridge University Press. p 277-311.

Whittaker DK, Molleson T, Bennett RB, Edwards A, Jenkins PR, Llewelyn JH. 1981. The prevalence and distribution of dental caries in a Romano-British population. *Arch Oral Biol* 26: 237-245.

Wiedemann T. 1989. *Adults and children in the Roman Empire*. London: Routledge.

Williams SA, Curzon MEJ. 1986. Observations of dental caries in primary teeth in some mediaeval British skull material. In: Cruwys E, Foley RA, eds. *Teeth and anthropology*. Oxford: BAR International Series 291. p 201-214.

Wilson P. 2011. *Introduction to heritage assets – Roman settlements*. London: English Heritage.

Woodward M and Walker ARP. 1994. Sugar consumption and dental caries: evidence from 90 countries. *Prev Dent* 176: 297-302

Table 1. Study sites by settlement date and type

Site	Date (AD)	Type	Total non-adults	TPR by tooth count a/p (%)	CPR by individual a/p (%)	Reference
Winchester (North, West, East)	1-4 th century	Major Urban	39	11/559 (2.0)	6/39 (15.4)	Ottaway et al. (2012)
Kingsholm, Gloucester	2-4 th century	Major Urban	9	9/135 (6.7)	4/9 (44.4)	Hurst (1985), (1986)
Gambier-Parry Lodge, Gloucester	2-4 th century	Major Urban	6	0/76 (0.0)	0/6 (0.0)	Heighway (1980); Mullin (2006)
Trentholme Drive, York	3-4 th century	Major Urban	17	2/262 (0.8)	2/17 (11.8)	Wenham (1968); Ottaway (2009)
Bath Gate, Cirencester	4 th century	Major Urban	38	5/483 (1.0)	4/38 (10.5)	Viner and Leech (1982)
Butt Road, Colchester	4-5 th century	Major Urban	80	22/1209 (1.8)	12/80 (15.0)	Crummy and Crossan (1993)
Baldock, Hertfordshire	2-4 th century	Minor Urban	25	2/414 (0.5)	2/25 (8.0)	Stead and Rigby (1986); Burleigh and Fitzpatrick-Matthews (2010)
Queenford Farm/Mill, Oxfordshire	3-4 th century	Minor Urban	51	8/729 (1.1)	3/51 (5.9)	Durham and Rowley (1972); Chambers (1987)
Ancaster, Lincolnshire	3-4 th century	Minor Urban	34	13/480 (2.7)	5/34 (14.7)	Todd (1975); Cox (1989)
Great Casterton, Rutland	3-4 th century	Minor Urban	27	7/383 (1.8)	4/27 (14.8)	McConnell et al. (2012)
Ashton, Northamptonshire	4 th century	Minor Urban	17	0/265 (0.0)	0/17 (0.0)	Dix (1983)
Dunstable, Bedfordshire	3-5 th century	Minor Urban	12	7/201 (3.5)	4/12 (33.3)	Matthews (1981)
Owslebury, Hampshire	1-4 th century	Rural	3	0/42 (0.0)	0/3 (0.0)	Collis (1968), (1977)
Cannington, Somerset	3-4 th century	Rural	69	4/977 (0.4)	3/69 (4.3)	Rahtz et al. (2000)
Catsgore, Somerset	2-5 th century	Rural	1	0/6 (0.0)	0/1 (0.0)	Leech (1982)
Bradley Hill, Somerset	4-5 th century	Rural	5	0/62 (0.0)	0/5 (0.0)	Leech et al. (1981)
Total non-adults			433	90/6283 (1.4)	49/433 (11.3)	

Table 2. Carious lesion frequencies by tooth type and individual

		Major Urban		Minor Urban		Rural		Total	
		Tth	Ind	Tth	Ind	Tth	Ind	Tth	Ind
Total	a/p	49/2724	28/189	28/2453*	17/165*	4/1087	3/78	81/6264*	48/432*
	%	1.8	14.8	1.5* (1.5)	10.3* (10.4)	0.3	3.8	1.3* (1.4)	11.1* (11.3)
Deciduous teeth	a/p	32/1075	14/125	10/1335	7/127	1/481	1/54	43/2891	22/306
	%	3.0	11.2	0.7	5.6	0.2	1.9	1.5	7.2
Permanent teeth	a/p	17/1649	14/124	18/1118	10/86	3/606	2/43	38/3373	26/253
	%	1.0	11.3	1.6	11.6	0.5	4.7	1.1	10.3
Anterior teeth	a/p	2/1100	2/171	0/1058	0/152	0/428	0/69	2/2586	2/392
	%	0.2	1.2	0	0	0	0	0.08	0.5
Posterior teeth	a/p	47/1624	28/185	28/1395	17/164	4/659	3/78	79/3678	48/427
	%	2.9	15.1	2.0	10.4	0.6	3.8	2.1	11.2

Tth = tooth count; Ind=individual count; a/p = number affected/number present; *corrected rates for the Ancaster child with SECC, uncorrected rates in brackets

Table 3. Carious lesion frequencies by age group and site type

Age (years)		Major Urban		Minor Urban		Rural		Total	
		Tth	Ind	Tth	Ind	Tth	Ind	Tth	Ind
1.1-2.5	a/p	1/394	1/38	0/571	0/49	0/193	0/21	1/1158	1/108
	%	0.3	2.6	0.0	0.0	0.0	0.0	0.1	0.9
2.6-6.5	a/p	12/555	7/50	7/774	4/56	0/254	0/22	19/1583	11/128
	%	2.2	14.0	0.9	7.1	0.0	0.0	1.2	8.6
6.6-10.5	a/p	20/539	8/35	4/393	4/27	0/112	0/8	24/1044	12/70
	%	3.7	22.9	1.0	14.8	0.0	0.0	2.3	17.1
10.6-14.5	a/p	11/883	8/44	11/520	5/25	1/315	1/15	23/1718	14/84
	%	1.3	18.2	2.1	20.0	0.3	6.7	1.3	16.7
14.6-17.0	a/p	5/213	4/20	6/195	4/8	3/351	2/12	14/759	10/40
	%	2.4	20.0	3.1	50.0	0.9	16.7	1.8	25.0

Tth = tooth count; Ind=individual count; a/p= number affected/number present

Table 4. Prevalence rates of enamel hypoplasia and co-morbidity

		Major Urban		Minor Urban		Rural		Total	
		Tth	Ind	Tth	Ind	Tth	Ind	Tth	Ind
Total	a/p	309/2700	54/214	236/2460	43/192	49/1087	12/105	594/6247	109/511
	%	11.4	25.2	9.6	22.4	4.5	11.4	9.5	21.3
Co-morbidity	a/p		11/189		7/165		1/78		19/432
	%		5.8		4.2		1.3		4.4

Tth = tooth count; Ind=individual count; a/p = number affected/number present

Table 5. Previously reported caries rates in Romano-British non-adults

Study	Site and context	Date (AD)	CPR (%)	TPR (%)	Comments
Moore and Corbett (1973)	Various unknown Major urban, military	1 st -5 th century		4.2	Deciduous teeth
O'Sullivan et al. (1993)	Various unknown Not stated	1 st -5 th century		16.0	Deciduous molars
Redfern et al. (2012)	Various (Dorset) Major urban, rural	1 st -5 th century		0.3/1.5	Deciduous/permanent teeth
Lewis (pers. comm.)	Poundbury Camp Dorchester Civitas	3 rd -5 th century	3.6	0.7/0.7	Deciduous/permanent teeth
Clough and Boyle (2010)	Lankhills Winchester Civitas	4 th century		1.7	Deciduous and permanent teeth
Caffell (2007)	Horncastle, Lincolnshire Fort/minor urban	4 th century		12.5/0.0	Deciduous/permanent teeth