

# *Cognitive bias modification of interpretation in children with social anxiety disorder*

Article

Accepted Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Orchard, F., Apetroaia, A., Clarke, K. and Creswell, C. (2017) Cognitive bias modification of interpretation in children with social anxiety disorder. *Journal of Anxiety Disorders*, 45. pp. 1-8. ISSN 0887-6185 doi: <https://doi.org/10.1016/j.janxdis.2016.10.012> Available at <http://centaur.reading.ac.uk/67936/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.janxdis.2016.10.012>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# Cognitive Bias Modification of Interpretation in Children with Social Anxiety Disorder

Faith Orchard<sup>1</sup>, Adela Apetroaia<sup>2</sup>, Kiri Clarke<sup>3</sup>, and Cathy Creswell<sup>1</sup>

<sup>1</sup>School of Psychology and Clinical Language Sciences, University of Reading,  
Whiteknights Road, Reading, RG6 6AL, United Kingdom

<sup>2</sup>Newham College University Centre, London, E15 4HT, United Kingdom

<sup>3</sup>Division of Psychology and Language Sciences, University College London, London,  
WC1E 6BT, United Kingdom

Corresponding author:

Faith Orchard,

School of Psychology and Clinical Language Sciences,

University of Reading, Reading, RG6 6AL, UK.

Telephone +44(0)118 378 8579

Email: f.orchard@reading.ac.uk

*Short Title:* Interpretation Bias and Child Social Anxiety

## Abstract

Negative (or a lack of positive) interpretation of ambiguous social situations has been hypothesised to maintain social anxiety disorder in children, yet there is currently limited evidence to support this. Cognitive Bias Modification of Interpretation (CBM-I) provides a means to explore the causal influence of interpretation bias on social anxiety disorder, and has been associated with a reduction in social anxiety symptoms in adults. Seven to twelve year old children with a diagnosis of social anxiety disorder completed CBM-I training, adapted from materials designed for socially anxious children in the community, or no training. Effects on interpretation bias and social anxiety were assessed. The adapted CBM-I training was not associated with significant changes in benign or negative interpretation. Unsurprisingly given the lack of successful interpretation training, there were no significant changes in child or parent reported social anxiety symptoms, clinician-rated severity or diagnoses and change in interpretation was not significantly associated with change in social anxiety. These findings contrast with some studies with community populations although it is possible that more intensive CBM-I training is required to fully test this hypothesis among clinical groups.

Keywords: social anxiety disorder, interpretation, cognitive bias modification, child

Social anxiety disorder is common in children (Beesdo et al., 2007), causes significant functional impairment (Erath, Flanagan, & Bierman, 2007; Van Ameringen, Mancini, & Farvolden, 2003) and is associated with long term risk of adult social anxiety as well as other mental health difficulties (Pine, Cohen, Gurley, Brook, & Ma, 1998; Zimmermann et al., 2003). Social information is often ambiguous, yet adults without social anxiety disorder often interpret this information in a positive manner. This helpful positive bias is lacking in adults with social anxiety disorder (Stopa & Clark, 2000; Hirsch & Mathews, 2000) and it is hypothesized that a lack of positive interpretation bias may play a fundamental role in the maintenance of social anxiety disorder (Hirsch & Clark, 2004).

Psychological treatments for social anxiety disorder among children typically include methods which aim to change interpretation bias (e.g. NICE, 2013). However it is far from clear that interpretation biases have a maintaining role in childhood social anxiety disorder as studies to date have typically examined cross-sectional associations, and where these have been conducted they have not provided consistent evidence for specific associations between social interpretation biases and social anxiety disorder (e.g. Muris et al., 2000). For example, Creswell, Murray & Cooper (2013) failed to find differences in either the frequency of threat interpretation in responses to ambiguous social scenarios, or in expectations of how a social challenge would turn out in children (7-12 years) with social anxiety disorder, other anxiety disorders and non-anxious children. However, although social anxiety disorder in adults is characterised by a lack of positive bias (e.g. Hirsch and Mathews, 2000; Garner, Mogg & Bradley, 2006), studies with children have typically failed to distinguish between increased positive and reduced negative interpretations, instead tending to treat these as on a continuum (e.g. Creswell, Murray, & Cooper, 2013). Furthermore, there has been limited examination of the prospective relationship

between cognitions and social anxiety in children (Muris, Huijding, Mayer, Remmerswaal, & Vreden, 2009).

One method that has the potential to determine causal influences of interpretation on social anxiety symptoms is Cognitive Bias Modification of Interpretation (CBM-I) which involves training participants to interpret ambiguous social stimuli in a more positive and/or less negative fashion. CBM-I has been associated with reduced social anxiety symptoms in both community and clinical adult populations (e.g. Murphy et al, 2007; Beard & Amir, 2008), and there is evidence that change in benign interpretation mediates the effect of training on social anxiety (Beard and Amir, 2008).

Recent applications of CBM-I with children have led to mixed results. On the basis of a meta-analysis of CBM (of attention and interpretation) interventions for mental health problems in children, Cristea, Mogoase, David and Cuijpers (2015) concluded that while CBM appeared to bring about significant changes in interpretation biases, these shifts did not translate to changes in child symptoms of anxiety, depression or general mental health (consistent with recent studies of Attention Bias Modification in the context of social anxiety; Carleton et al., 2015; Heeren, Mogoase, McNally, Schmitz & Philippot, 2015; Yao, Yu, Qian & Li; 2015). However, interpretation measures were collapsed to include both controlled in-lab experiments and real-life, ecologically valid measures, leaving the extent to which interpretation bias changed somewhat unclear. Where studies have focused on CBM-I and social anxiety specifically, findings have been mixed. One study reported a reduction in trait social anxiety among twenty two 10-11 year old children from a community population who scored highly on trait social anxiety, after 3 sessions of CBM-I with 45 trials (Vassilopoulos, Banerjee & Prantzalou, 2009). Yet, another study found no training effects on trait social anxiety among 77 10-13 year old

children, despite using similar methods (Vassilopoulos, Moberly, & Zisimatou, 2013). In both studies, interpretation bias training was associated with reduced negative interpretation (Vassilopoulos et al., 2009, 2013) and in Vassilopoulos et al. (2013) training was also associated with more benign interpretation of ambiguous social scenarios. One possible reason why CBM-I may have failed to translate to a change in social anxiety symptoms in Vassilopoulos et al. (2013) is that participants were an unselected sample who did not have elevated levels of social anxiety at the outset. Indeed, other studies with unselected populations have managed to successfully train interpretation biases but have failed to find an effect on general anxiety symptoms in children (Lester, Field, & Muris, 2011; Salemink & Wiers, 2011; though see Hirsch, Mathews & Clark, 2007, for contrasting evidence in adults); and it has been suggested that symptoms may be more likely to change following CBM-I for highly anxious child populations (e.g. Muris et al., 2008; Cristea et al., 2015). No studies to date have applied CBM-I with children who meet diagnostic criteria for social anxiety disorder, however, if successful in reducing social anxiety symptoms, this provides both valuable insights in to the relationship between interpretation and social anxiety and may lead to low-intensity treatment options (e.g. Beard et al., 2011).

We aimed to address whether CBM-I is associated with more benign interpretation and less negative interpretation of ambiguous social scenarios, a reduction in social anxiety symptoms and severity, and whether change in social anxiety was mediated by changes in interpretation. Some particular methodological features of the current study should be noted. We aimed to use an established method of CBM-I for social anxiety which had previously been used with high socially anxious children recruited from the community (Vassilopoulos et al., 2009). However, given previous CBM-I training programmes for social anxiety in children have relied

heavily on children's reading abilities and have not used highly systematised procedures (instead presenting training materials on cards which are read, circled and then turned over by participants to reveal 'correct' interpretations and therefore varying in timing of presentation, manner in which materials are read etc, e.g. Vassilopoulos et al., 2009, 2013) we adapted these previously used training materials for administration using experimental software with accompanying audio-materials. Furthermore, we asked participants to imagine themselves in the described scenarios since this has been shown to augment CBM-I in adults with depression (Holmes, Lang, & Shah, 2009).

The study hypotheses were as follows:

1. Children who receive the CBM-I training will have more benign interpretation and less negative interpretation post-training compared to children who do not receive CBM-I training.
2. Children who receive the CBM-I training will have lower scores on child, parent and clinician ratings of social anxiety post-training, compared to children who do not receive CBM-I training.
3. The association between group ('training', 'no training') and change in social anxiety will be mediated by the change in benign and negative interpretation biases, i.e., confirming a causal relationship between interpretation biases and anxiety.

### **Materials and Methods**

On the basis of the large effect found in Vassilopoulos et al. (2009), 30 participants were required to conduct repeated measures analyses of variance investigating within-between interactions (effect size  $F = 0.35$ ; power 95%; correlation among repeated measures, 0.48, as reported below). However, as an effect size has yet to be obtained with a clinical group, we



powered this study for a medium effect size ( $F = 0.25$ ), and so recruited  $n=56$  participants in order to provide at least 90% power to detect this more conservative effect.

## **Participants**

All participating children had been referred to the Berkshire Child Anxiety Clinic at the University of Reading by health or education professionals for assessment and treatment of an anxiety disorder. Children were invited to take part if they met the following inclusion criteria: (i) a primary anxiety disorder and a diagnosis of social anxiety disorder, (ii) aged 7-12 years, (iii) fluent English speakers, (iv) absence of learning difficulties (including autistic spectrum disorder), (v) absence of severe condition or risk that requires immediate treatment. A flow diagram showing recruitment and retention is shown in Figure 1. Fifty six clinically anxious children and their primary caregivers (all mothers) gave informed consent and took part in all stages of the study. Participants were randomized, using a random number generator, to either receive training (CBM) or not to receive training (NO CBM). The groups were well balanced on child age, gender, ethnicity, socioeconomic status, and symptoms of anxiety and low mood (see Table 1). No significant difference was found between groups for child anxiety disorder by comparing primary diagnosis ( $\chi^2(6) = 4.68, p = .59$ ) and frequency of social anxiety as the primary disorder ( $\chi^2(1) = 0.25, p = .62$ ). Groups did not differ on the presence of externalizing disorders ( $\chi^2(1) = 2.70, p = .10$ ) or mood disorders ( $\chi^2(1) = 1.46, p = .23$ ). (See Table 2 for frequencies of primary anxiety diagnoses and overall diagnoses). There was also no difference between the CBM group ( $M = 5.48, SD = .63$ ) and the NO CBM group ( $M = 5.37, SD = .74$ ) on ADIS-C/P clinical severity ratings of the primary diagnosis ( $t(54) = -.61, p = .54$ ). **Measures**

**Anxiety Disorders Interview Schedule for DSM IV for Children- Child and Parent Versions (ADIS-C/P; Silverman & Albano, 1996).** Children were assigned diagnoses on the

basis of the ADIS-C/P, a structured diagnostic interview with well-established psychometric properties (Silverman, Saavedra, & Pina, 2001). The interview covers anxiety disorders, mood disorders (depression and dysthymia) and behavioural disorders (conduct disorder, oppositional defiant disorder and attention deficit hyperactivity disorder). Where children met symptom criteria for a diagnosis (based on either child or parent report) they were assigned a clinical severity rating (CSR) ranging from 0 (*complete absence of psychopathology*) to 8 (*severe psychopathology*). As is conventional, only those children who met symptom criteria with a CSR of 4 or more (*moderate psychopathology*) were considered to meet diagnostic criteria. For the ADIS-C/P, as is standard, overall diagnoses and CSRs were assigned if the child met diagnostic criteria on the basis of either child or parent report, and the higher CSR of the two was taken. Assessors (psychology graduates) were trained on the standard administration and scoring of the ADIS-C/P through verbal instruction, listening to assessment audio-recordings and participating in diagnostic consensus discussions. The first 20 interviews conducted were then discussed with a consensus team, led by an experienced diagnostician (Consultant Clinical Psychologist). The assessor assigned diagnoses and CSRs prior to the consensus meeting and the consensus team independently allocated diagnoses and CSRs after the discussion. It is worth noting that the consensus discussions were limited by a reliance on how the team interpreted the interviewer's description, however audio recordings of ADIS-C/P interviews were available and were used to inform discussions / check decisions at times. Following the administration of 20 interviews, interrater reliability for each assessor was checked, and if assessors achieved reliability of at least .85, they were then required to discuss just one in six interviews with the consensus team (ongoing checks were conducted to prevent interrater drift). All assessors were reliable after 20 interviews. Overall reliability (on the first 20 and the subsequent interviews that were discussed)

was conducted for the assessment team and was found to be excellent (Diagnosis  $\alpha = .98$  (child report),  $\alpha = 1.00$  (mother report); CSR  $\alpha = .91$  (child report),  $\alpha = .97$  (mother report))<sup>1</sup>.

**Spence Children's Anxiety Scale (SCAS-C/P; Nauta et al., 2004; Spence, 1998).** The SCAS-c/p requires children/parents to rate how often they/their child experience/s each of 38 anxiety symptoms (presented alongside six positive filler items in the child report version) on a 4 point scale from 0 (*never*) to 3 (*always*). The social phobia scale of the SCAS-C/P was used as an indicator of self- and parent- reported social anxiety symptoms. The social phobia scale of the SCAS has been found to correlate highly with the Social Anxiety Scale for Children (SASC-R; Muris, Merckelbach, & Damsma, 2000). Internal consistency was acceptable to good (SCAS-c  $\alpha = 0.84$ ; SCAS-p  $\alpha = 0.84$ ; SCAS-SP-c  $\alpha = 0.69$ ; SCAS-SP-p  $\alpha = 0.80$ ).

**Short Mood and Feelings Questionnaire (SMFQ-C/P; Angold et al., 1995).** In order to assess the severity of common comorbid symptoms, and identify potential group differences, the SMFQ-c/p was administered to assess child and parent reported low mood. The SMFQ is a brief, 13 item measure which requires children/parents to report how often in the past two weeks the child has experienced a number of depressive symptoms on a 3 point scale from 0 (*not true*) to 2 (*certainly true*). Internal consistency was good (SMFQ-c  $\alpha = 0.80$ ; SMFQ-p  $\alpha = 0.87$ ).

**Ambiguous Scenarios Interview for Children (Vassilopoulos & Banerjee, 2009).** The ambiguous scenarios interview was used as a measure of both negative and benign interpretation. Interviews were conducted by a graduate research assistant and consisted of 16 ambiguous social scenarios which reflect events that commonly occur and are relevant for participants of this age, such as inviting classmates to your birthday party, approaching a group of peers, or going to a classmate's home to play together. Each scenario is followed by two thoughts which might occur

---

<sup>1</sup>As different assessors interviewed the mothers and child simultaneously, reliability figures for parent and child report were calculated separately.

to children in these situations reflecting a negative (for example, '[classmates] don't want to come [to party] because they don't like me'), or a benign judgement ('they don't know yet if they can come or not'). Children rated how likely they would be to endorse each explanation on a 5 point Likert scale. The first set of scenarios was presented at intake, the other set at the end of the training phase. Total ratings were calculated across the scenarios for each participant. The minimum score for benign or negative judgments was 8, the maximum was 40. The internal validity of the ASI overall was acceptable (benign  $\alpha = .76$ ; negative  $\alpha = .77$ ).

**CBM-I Training (adapted from Vassilopoulos and Banerjee, 2009).** The CBM-I training materials consisted of a total of 45 social scenarios presented during three sessions of 15 scenarios each, using translations of the materials developed by Vassilopoulos and Banerjee (2009). After being shown each ambiguous scenario participants were asked to select one of two alternative endings, a threatening and a non-threatening one, in counterbalanced order e.g. 'You enter the classroom and say hello to your schoolmate. However he/she doesn't say anything. Why do you think this happens?' (i) She has something else on her mind and she did not hear me or (ii) She doesn't like me anymore. Before selecting a response, participants were encouraged to reflect on the 'correct' interpretation. If the participants selected the non-threatening ending, they received the message 'This is correct', visually and aurally. If they selected the threatening interpretations, participants would see and hear the message 'This is the correct answer:' followed by the non-threatening ending. In both situations, participants were prompted by an audio message to think about how the non-threatening ending might explain the situation. A flow diagram representing the stages of the training procedure is shown in Figure 1.

The procedure was adapted from Vassilopoulos, Banerjee and Prantzalou's (2009) method in four ways. Firstly, materials were presented using E-Prime Version 2.0 rather than on

paper cards. To reduce the burden of reading, participants listened to each scenario read by a female actor in a friendly, neutral voice as scenarios were presented on the computer screen, in text, at the same time. Secondly, in order for children to understand the rationale of the study they were told that they would take part in a programme that would teach them some new ways of thinking in relation to their social worries. They were introduced to the connection between thoughts and emotions before the training programme began using two hand-outs from the 'Cool Kids' anxiety treatment programme (Lyneham, Abbott, Wignall & Rapee, 2003). A research assistant worked through the hand-outs with the child and encouraged them to identify the thoughts and feelings in the images. In the first one, 'The way I think and feel', children were given examples of four situations ('what happened?') followed by a thought ('what was I thinking?') and an emotion ('what was I feeling?'). Two of these situations were followed by a threatening interpretation, two by a non-threatening interpretation and the corresponding emotions. So as to not interfere with the training, the situations presented were non-social (for example, situation: 'a big dog comes near me'; thought: - 'the dog wants to play with me'; emotion - 'happy to play with the dog'). The second hand-out, 'How I feel depends on what I think', further explored the link between thoughts and emotions and children were encouraged to fill out examples of different thoughts and their associated emotions. As before, the situations presented were non-social. Thirdly, because accompanying imagery has been found to enhance the effects of CBM-I in studies with adult populations (Holmes, Lang & Shah, 2009), participants were given specific instructions to explain what an image was and to help them create a visual image of the scenarios. They were encouraged to practice imagining being at the beach, and seeing their favourite food, by concentrating on what they could see, hear, smell, feel and taste. They were then instructed to imagine that the training situation described is happening

to them, even if they find it unlikely (following Lothmann, Holmes, Chan & Lau, 2011). Finally, at the end of each training item an inference based on the non-threatening interpretation is presented and the child was asked to identify it as true or false. As in CBM with adult populations (Hayes, Hirsch, Krebs & Mathews, 2010), this was intended to reinforce the non-threatening interpretation and encourage active engagement in the generation of meaning. As we only presented positive interpretations at this stage, the ‘correct’ answer was always ‘True’ so the position of the ‘True’ and ‘False’ keys were varied to engage the child in thinking about the response. In keeping with some previous studies that have successfully trained altered interpretations, children did not receive feedback on these responses (e.g. Hayes et al., 2010, Beard & Amir, 2008, Micco, Henin & Hirshfeld-Becker; 2014).

### **Ethical considerations**

This study was reviewed by the Local Research Ethics Committee on behalf of the National Health Service and the University of Reading Research Ethics Committee. Parents and children were both provided with written and verbal information about the study. In order to participate in the study written parental consent and child assent were both required.

### **Procedure**

Children and their parents completed diagnostic interviews and symptom questionnaires as part of their routine clinical evaluation. All participants received a visit at home where they signed consent forms and completed the first set of the Ambiguous Scenarios Interview (Pre-Training) questions. Participants allocated to the CBM group were told that they would receive an experimental treatment which would teach them new ways of thinking. These participants attended a further three visits at the University to complete the CBM-I training. Each training session lasted approximately 30 minutes, with the first session lasting 45 minutes in order to also

complete the hand-outs first. The sessions were spaced as evenly as possible within two weeks. Children in the NO CBM group were assigned to the waitlist condition and were not required to attend any visits until the re-assessment. All families were informed that they were on a waiting list to receive treatment as usual immediately following the CBM study, and no families reported having started any additional treatment during this time.

Reassessments were scheduled for 4 weeks following the initial home visit. However, the reassessment was completed a mean of 6.11 weeks ( $SD = 2.84$ ) weeks after group allocation, and 1.10 weeks ( $SD = .34$ ) weeks after the final training session for the CBM group. This was due to rearrangements of sessions made by families. The time from allocation to reassessment did not differ between groups ( $t(54) = .10, p = .92$ ). At this reassessment all participants completed (i) the second set of the Ambiguous Scenarios Interview (Post-Training) questions, (ii) the SCAS-c/p, and (iii) the Social Phobia section of the ADIS-C/P. All post-training assessors were blind to participant group.

## **Results**

### **Preliminary analyses**

Continuous data were screened in relation to the assumptions of parametric tests (Tabachnick & Fidell, 2007). Where assumptions were violated, confirmatory analyses were conducted by running analyses with 1000 bootstrap samples or non-parametric alternatives. The majority of results were consistent, suggesting that the original analyses were robust to the violations of assumptions, so results based on the original (non-bootstrapped) analyses are presented for simplicity.

### **Change in interpretation bias**

To examine hypothesis one, two mixed design analyses of variance (ANOVA) were conducted with group (CBM vs. NOCBM) as the independent variable and measures of interpretation bias as the repeated dependent variables (see Table 3). This approach was taken rather than conducting a single multivariate analysis of variance due to concerns regarding collinearity given the high correlation between benign and negative interpretation scores at time 2 ( $r = -.51, p < .001$ ) (Field, 2009).

Significant, large main effects of time were found for both negative ( $V = .12, F(1,53) = 7.30, p = .01; d = .72$ ) and benign interpretation ( $V = .50, F(1,53) = 52.29, p < .001; d = 1.93$ ), reflecting the fact that participants had less negative and more benign interpretation post-compared to pre-training, regardless of group. The group x time interaction effect approached significance, reflecting a trend towards a greater increase in benign interpretation among the CBM group with a medium effect size ( $V = .07, F(1,53) = 3.84, p = .055; d = .52$ ). There was not a significant interaction between group and time for negative interpretation where the effect was small ( $V = .01, F(1,53) = .32, p = .58; d = .15$ ).

### **Change in social anxiety symptoms, severity and diagnoses**

To examine hypothesis two, three mixed design analyses of variance (ANOVA) were conducted with group (CBM vs. NOCBM) as the independent variable, and measures of self- and parent-reported social anxiety symptoms and clinical severity ratings of diagnoses as repeated dependent variables (see Table 3).

There was a main effect of time on child-reported social phobia symptoms on the SCAS ( $V = .09, F(1,54) = 5.09, p = .03; d = .60$ ) reflecting a decrease in symptoms from pre- to post-training regardless of group. A significant effect was not found for parent-reported symptoms ( $V = .01, F(1,54) = .77, p = .39; d = .23$ ). There was not a significant time x group interaction for



either parent ( $V = .01$ ,  $F(1,54) = .32$ ,  $p = .58$ ;  $d = .15$ ) or child-reported social phobia symptoms ( $V = .01$ ,  $F(1,54) = .28$ ,  $p = .60$ ;  $d = .14$ ).

There was not a significant main effect of time on social phobia clinical severity rating ( $V = .01$ ,  $F(1,54) = .35$ ,  $p = .56$ ;  $d = .16$ ), nor a significant interaction between time and group for clinical severity rating ( $V = .06$ ,  $F(1,54) = 3.27$ ,  $p = .08$ ;  $d = .48$ ) and notably the pattern of results was in the opposite direction to that predicted (see Table 3).

All participants in the CBM group maintained their social anxiety disorder diagnoses post-training, and only 2 participants no longer met criteria for social anxiety disorder in the NOCBM group, which did not reflect a significant difference between groups ( $\chi^2(1) = 2.23$ ,  $p = .14$ ).

Associations between change in social anxiety (child and parent report and clinician severity ratings) and change in benign or negative interpretations were not statistically significant after correcting for multiple tests (Bonferroni-corrected significance criterion level  $\alpha = .008$ ) (see Table 4). Unsurprisingly, given the lack of significant associations, there was no evidence of indirect effects (using the PROCESS macro; Hayes, 2013) of CBM on social anxiety symptoms or severity via change in benign or negative interpretation.

## **Discussion**

The current study is the first to investigate CBM-I in children with clinical levels of social anxiety. We administered an established method of modifying interpretation among children (with some adaptations aimed to standardise and enhance the procedure), however children with social anxiety disorder did not report significantly greater changes in benign or negative interpretation after receiving CBM-I training than those who did not (although differences in change in benign interpretations approached significance). While conclusions must

necessarily be tempered by the lack of successfully training interpretation bias, changes in interpretation were not significantly associated with changes in social anxiety symptoms, severity or diagnoses. It is worth noting that there was a trend towards significance, with a medium effect size, for a change in clinical severity ratings, however, this was not in the direction expected. Specifically the mean scores indicated that clinical severity increased in the CBM group and decreased in the NOCBM group.

These results differ from those of Vassilopoulos et al. (2009) who found a significant reduction in symptoms of anxiety following CBM-I among (non-clinical) children with elevated social anxiety symptoms. Notably, however, in that study children reported significant reductions in negative interpretation following the CBM-I procedure. It is possible that with a clinical population a greater intensity of training is required to bring about change in interpretation. For example, in the first trial of multisession CBM-I with adults diagnosed with social anxiety disorder the CBM-I training used a word sentence association paradigm, and the intensity of CBM-I training was substantially higher (12 x 20 minutes over 6 weeks, with 220 training trials in each session) than in the present study, and this training was associated with both a reduction in negative interpretation and an increase in positive interpretation of novel social situations (Amir & Taylor, 2012). Following Vassilopoulos and Banerjee (2009), there was a low dose of training in the present study, with only 45 training trials in total distributed over three sessions (i.e. 15 items per session). In comparison with most CBM-I multisession studies, this is a low dose both in terms of number of session and number of overall trials. The reason that we employed this method was based on its previous success and because we were concerned that children would not engage with CBM-I if too many sessions or trials were employed. However, it is notable that children of 10 years of age have been found to comply with large numbers of

(albeit far briefer) trials in Attention Bias Modification (ABM) procedures (768 trials per session; Bar-Haim, Morag & Glickman; 2011) so more extensive CBM-I training with children may well be feasible. Notably the CBM-I appeared to be an acceptable intervention for children referred for treatment for social anxiety disorder and their parents. Consistent with adult studies in which participants are paid to take part (Beard et al., 2011), retention to this study (where there was no participant payment) was high, with only two of the 29 CBM-I participants failing to complete the training. No adverse effects of the training were reported. This suggests that future studies with greater therapeutic dose or where CBM-I is used as an adjunct alongside, for example, CBT (e.g. Beard, 2011; although see Williams, O'Moore, Blackwell, Smith, Holmes & Andrews, 2015) may be feasible interventions for children with social anxiety disorder.

It is important to note that in the current study, both the CBM-I and control groups experienced an increase in benign interpretation and a reduction in negative interpretation over time. It is unclear whether this is an artefact of the test used, regression towards the mean, or a non-specific effect of being part of a study that investigates social anxiety where an assessor meets with the child and parent. Furthermore, it has not been formally established that the two sets of the ambiguous scenarios measures of interpretation are equivalent, however Vassilopoulos et al. (2009) found no significant differences on scores on the two sets administered before and after a no-training condition. Clearly, before firm conclusions regarding the role of interpretation bias in maintaining clinical levels of social anxiety in children, or the potential utility of CBM-I in this population can be determined, effects of higher training dose needs to be trialed and more effective training methods need to be developed, including the use of CBM training and assessment materials that are specifically tailored for individuals with social anxiety disorder. For example a key concern in the context of social anxiety disorder is

how the individual comes across to others, so materials which target this specifically may facilitate more effective training (e.g. Murphy et al., 2007).

The CBM-I training used here had previously been associated with a reduction in social anxiety following training among a high trait social anxiety community population (Vassilopoulos and Banerjee, 2009), however, in addition to the low dose, there are a number of reasons why the training may not have been optimal for application with clinically anxious children. Specifically, the CBM-I programme used here did not require participants to actively generate the meaning within trials, a factor which has been suggested to augment training effects (Mathews & Macintosh, 2000). CBM-I studies have typically either involved active generation (Hoppitt, Mathews, Yiend & Mackintosh, 2010) or the use of imagery, with imagery having similar beneficial effects to active generation on mood in adults (e.g. Holmes and Mathews, 2005). Consequently, we adapted Vassilopoulos and Banerjee's (2009) methods to include generation of self-imagery. However, on reflection requiring children to generate images of themselves in the social scenarios may have been counterproductive. Indeed, Vassilopoulos, Blackwell, Moberly and Karahaliou (2012) recently found that children who read verbal descriptions and thought about their meaning showed a greater reduction in negative interpretation than children who imagined the events. Furthermore, social anxiety is associated with the generation of stereotyped negative self-images in socially anxious individuals (e.g. Hackmann et al., 2000); it is therefore feasible that *negative* images were generated, that the children were unable to imagine themselves in the positive scenario, or even if they did so, that they did not believe that this is the way that the situation would actually go for them. Mathews, Ridgeway, Cook and Yiend (2007) used graded training where interpretation were initially benign and gradually moved on to more positive interpretation over time, minimising the

potential for the training materials to be rejected. This may be a useful approach in future research.

Other considerations in interpreting inconsistencies in findings in relation to previous studies include differences in the method of administration of CBM-I in the current study which presented training materials via computer rather than experimenter (reducing the potential for experimenter bias). We also made other adaptations which were intended to augment training effects, however it is possible that these may have had the opposite effect. For example, we only presented benign comprehension questions at the end of each training item and did not provide feedback on responses to this. Furthermore, we made the aims of training explicit, but recent research by Grafton, Mackintosh, Vujic and MacLeod (2014) found that explicit instructions given during CBM, designed to facilitate positive attentional bias, led to poorer outcomes under stress conditions, potentially in keeping with the lack of transfer to symptoms of social anxiety in the current study (though see Krebs et al., 2010, for contrasting results highlighting the potential influence of the manner in which aims are made explicit).

Finally, the lack of a significant association between change in interpretation and change in social anxiety may reflect the possibility that interpretation biases do not have an independent causal role in relation to social anxiety in children, although the failure to successfully train a change in bias means we must be extremely cautious in drawing conclusions. However this suggestion would be consistent with recent findings that have failed to establish a tendency towards greater threat interpretation of social scenarios among children with social anxiety disorder (Creswell et al., 2013). Notably, however, these studies have typically failed to make a distinction between negative and benign interpretations. Future studies would benefit from

consideration of developmental differences in the nature of the association between interpretation and social anxiety from childhood to adulthood.

Strengths of the current study were the inclusion of age and gender balanced groups from a referred population who all met diagnostic criteria for social anxiety disorder and consideration of potential confounding effects (behavioural disturbance and low mood). However, it is important to note certain limitations including the sample demographics (mostly high SES, Caucasian families) which limit the extent to which the findings can be generalised. We did not include a training control group, but as there were no differences in anxiety symptoms between the CBM and no intervention, this presents less of a problem in terms of interpretation. We also included children who met criteria for social anxiety disorder but this was not required to be their primary diagnosis. This may have meant that other interpretation biases were at play which may have accounted for the lack of translation of the effects of training to change in symptoms.

### **Conclusion**

CBM-I training, adapted from methods successfully used with community populations, was not associated with significant changes in benign or negative interpretation in response to ambiguous information or changes in self, parent or clinician-reported social anxiety post-training. As such this study is not able to provide evidence relating to the causal influence of interpretation on social anxiety in children. Higher doses of CBM-I training are likely to be required to fully test this hypothesis.

## Acknowledgements

The authors would like to thank the participating families and the staff at the Berkshire Child Anxiety Clinic, in particular Anna Alkozei, Debbie Andrews, Amy Corcoran, Jenny Crosby, Sue Cruddace, Kara Ellis-Jones, Katherine Fowler, Laura Hawkins, Kelsey Heard, Claire Hill, Zoe Hughes, Jessica Karalus, Ashton Kissoon, Hannah Maxted, Lara Payne, Emma Stephens, Helen Stubbs and Lucy Willetts for their help programming, collecting and coding data and to Colette Hirsch for her feedback on the manuscript. CC was funded by MRC Clinician Scientist Fellowship (G0601874) when this study was conducted and is now funded by an NIHR Research Professorship (RP\_2014-04-018). The views expressed are those of the authors and not necessarily those of the Medical Research Council, NHS, the NIHR or the Department of Health.

## References

- Amir, N., & Taylor, C. T. (2012). Interpretation training in individuals with generalized social anxiety disorder: a randomized controlled trial. *Journal of consulting and clinical psychology, 80*(3), 497.
- Angold, A., Costello, E. J., Messer, S. C., Pickles, A., Winder, F., & Silver, D. (1995). Development of a short questionnaire for use in epidemiological studies of depression in children and adolescents. *International Journal of Methods in Psychiatric Research, 5*, 237-249.
- Bar-Haim, Y., Morag, I., & Glickman, S. (2011). Training anxious children to disengage attention from threat: a randomized controlled trial. *Journal of Child Psychology and Psychiatry, 52*(8), 861-869.
- Beard, C., & Amir, N. (2008). A multi-session interpretation modification program: Changes in interpretation and social anxiety symptoms. *Behaviour Research and Therapy, 46*(10), 1135-1141.
- Beard, C., Weisberg, R. B. & Amir, N. (2011). Combined Cognitive Bias Modification Treatment for Social Anxiety Disorder: A Pilot Trial. *Depression & Anxiety, 28*(11), 981–988.
- Beesdo, K., Bittner, A., Pine, D. S., Stein, M. B., Höfler, M., Lieb, R., & Wittchen, H. U. (2007). Incidence of social anxiety disorder and the consistent risk for secondary depression in the first three decades of life. *Archives of General Psychiatry, 64*(8), 903-912.



- Carleton, R. N., Sapach, M. J. T., Oriet, C., Duranceau, S., Lix, L. M., Thibodeau, M. A., ... & Asmundson, G. J. (2015). A randomized controlled trial of attention modification for social anxiety disorder. *Journal of Anxiety Disorders*.
- Cristea, I. A., Mogoșe, C., David, D., & Cuijpers, P. (2015). Practitioner review: cognitive bias modification for mental health problems in children and adolescents: a meta-analysis. *Journal of Child Psychology and Psychiatry*.
- Creswell, C., Murray, L., & Cooper, P. (2013). Interpretation and expectation in childhood anxiety disorders: age effects and social specificity. *Journal of Abnormal Child Psychology*, 1-13.
- Erath, S. A., Flanagan, K. S., & Bierman, K. L. (2007). Social anxiety and peer relations in early adolescence: Behavioral and cognitive factors. *Journal of Abnormal Child Psychology*, 35(3), 405-416.
- Field, A. (2009). *Discovering Statistics Using SPSS*. London: Sage Publications.
- Garner M., Mogg K. & Bradley B. P. (2006). Fear-relevant selective associations and social anxiety: absence of a positive bias. *Behaviour Research & Therapy*, 44(2), 201-217.
- Grafton, B., Mackintosh, B., Vujic, T. & MacLeod, C. (2014). When Ignorance is Bliss: Explicit Instruction and the Efficacy of CBM-A for Anxiety. *Cognitive Therapy and Research*, 38, 172-188.
- Grey, S. J., & Mathews, A. M. (2009). Cognitive bias modification—Priming with an ambiguous homograph is necessary to detect an interpretation training effect. *Journal of behavior therapy and experimental psychiatry*, 40(2), 338-343.
- Hackmann, A., Clark, D. M. & McManus, F. (2000). Recurrent images and early memories in social phobia. *Behaviour Research & Therapy*. 38(6):601-10.

- Hayes, A. F. (2013). *Introduction to mediation, moderation, and conditional process analysis: A regression-based approach*. Guilford Press.
- Hayes, S., Hirsch, C. R., Krebs, G., & Mathews, A. (2010). The effects of modifying interpretation bias on worry in generalized anxiety disorder. *Behaviour research and therapy, 48*(3), 171-178.
- Heeren, A., Mogoșe, C., McNally, R. J., Schmitz, A., & Philippot, P. (2015). Does attention bias modification improve attentional control? A double-blind randomized experiment with individuals with social anxiety disorder. *Journal of anxiety disorders, 29*, 35-42.
- Hirsch, C. R., & Clark, D. M. (2004). Information-processing bias in social phobia. *Clinical psychology review, 24*(7), 799-825.
- Hirsch, C. R. & Mathews, A. (2000). Impaired positive inferential bias in social phobia. *Journal of Abnormal Psychology, 109*, 705–712.
- Hirsch, C. R., Mathews, A., & Clark, D. M. (2007). Inducing an interpretation bias changes self-imagery: A preliminary investigation. *Behaviour Research and Therapy, 45*(9), 2173-2181.
- Holmes, E. A., Lang, T. J. & Shah, D. M. (2009). Developing interpretation bias modification as a "cognitive vaccine" for depressed mood: Imagining positive events makes you feel better than thinking about them verbally. *Journal of Abnormal Psychology, 118*(1), 76-88.
- Holmes, E. A., & Mathews, A. (2010). Mental imagery in emotion and emotional disorders. *Clinical psychology review, 30*(3), 349-362.

- Hoppitt, L., Mathews, A., Yiend, J., & Mackintosh, B. (2010). Cognitive bias modification: The critical role of active training in modifying emotional responses. *Behavior Therapy*, *41*(1), 73-81.
- Krebs, G., Hirsch, C. R., & Mathews, A. (2010). The effect of attention modification with explicit vs. minimal instructions on worry. *Behaviour research and therapy*, *48*(3), 251-256.
- Lester, K. J., Field, A. P., & Muris, P. (2011). Experimental modification of interpretation bias about animal fear in young children: effects on cognition, avoidance behavior, anxiety vulnerability, and physiological responding. *Journal of Clinical Child & Adolescent Psychology*, *40*(6), 864-877.
- Lothmann, C., Holmes, E., Chan, S., & Lau, J. (2011). Cognitive bias training in adolescents: effects on interpretation bias and mood. *Journal of Child Psychology and Psychiatry*, *52*, 24-32.
- Lothmann, C., Viechtbauer, W., Höhn, P., Kasanova, Z., Haller, S. P., Drukker, M., ... & Lau, J. Y. (2014). How to boost positive interpretations? A meta-analysis of the effectiveness of cognitive bias modification for interpretation. *PloS one*, *9*(6), e100925.
- Lyneham, H. J., Abbott, M. J., Wignall, A., & Rapee, R. A. (2003). The cool kids anxiety treatment program. *Sydney: MUARU, Macquarie University. Misfud C & Rapee RM (2005) Early intervention for childhood anxiety in a school setting: outcomes for an economically disadvantaged population. Journal of the American Academy of Child and Adolescent Psychiatry*, *44*(10), 995.
- Mathews, A., & Mackintosh, B. (2000). Induced emotional interpretation bias and anxiety. *Journal of Abnormal Psychology*, *109*(4), 602-615.

- Mathews, A., Ridgeway, V., Cook, E., & Yiend, J. (2007). Inducing a benign interpretational bias reduces trait anxiety. *Journal of Behavior Therapy and Experimental Psychiatry*, 38(2), 225-236.
- Micco, J. A., Henin, A., & Hirshfeld-Becker, D. R. (2014). Efficacy of interpretation bias modification in depressed adolescents and young adults. *Cognitive therapy and research*, 38(2), 89-102.
- Muris, P., Huijding, J., Mayer, B., & Hameetman, M. (2008). A space odyssey: experimental manipulation of threat perception and anxiety-related interpretation bias in children. *Child Psychiatry and Human Development*, 39(4), 469-480.
- Muris, P., Huijding, J., Mayer, B., Remmerswaal, D., & Vreden, S. (2009). Ground control to Major Tom: experimental manipulation of anxiety-related interpretation bias by means of the “space odyssey” paradigm and effects on avoidance tendencies in children. *Journal of Anxiety Disorders*, 23(3), 333-340.
- Muris, P., Kindt, M., Bögels, S., Merckelbach, H., Gadet, B., & Moulart, V. (2000). Anxiety and threat perception abnormalities in normal children. *Journal of Psychopathology and Behavioural Assessment*, 22, 183–199.
- Muris, P., Merckelbach, H., & Damsma, E. (2000). Threat perception bias in nonreferred, socially anxious children. *Journal of Clinical Child Psychology*, 29(3), 348-359.
- Murphy, R., Hirsch, C. R., Mathews, A., Smith, K., & Clark, D. M. (2007). Facilitating a benign interpretation bias in a high socially anxious population. *Behaviour Research and Therapy*, 45(7), 1517-1529.
- Nauta, M. H., Scholing, A., Rapee, R. M., Abbott, M., Spence, S. H., & Waters, A. (2004). A parent-report measure of children's anxiety: psychometric properties and comparison with

- child-report in a clinic and normal sample. *Behaviour Research and Therapy*, 42(7), 813-839.
- National Institute for Health and Care Excellence (NICE). (2013). *Social anxiety disorder: recognition, assessment and treatment CG159*. London: National Institute for Health and Care Excellence.
- Pine, D. S., Cohen, P., Gurley, D., Brook, J., & Ma, Y. (1998). The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives of General Psychiatry*, 55(1), 56-64.
- Salemink, E., & Wiers, R. W. (2011). Modifying threat-related interpretive bias in adolescents. *Journal of Abnormal Child Psychology*, 39(7), 967-976.
- Silverman, W. K., & Albano, A. M. (1996). *The Anxiety Disorders Interview Schedule for DSM-IV: Child and Parent Versions*. The Psychological Corporation. San Antonio, TX.
- Silverman, W. K., Saavedra, L. M., & Pina, A. A. (2001). Test-retest reliability of anxiety symptoms and diagnoses with the Anxiety Disorders Interview Schedule for DSM-IV: Child and parent versions. *Journal of the American Academy of Child and Adolescent Psychiatry*, 40(8), 937-944.
- Spence, S. H. (1998). A measure of anxiety symptoms among children. *Behaviour Research and Therapy*, 36, 545-566.
- Stopa. L. & Clark, D.M. (2000). Social phobia and interpretation of social events. *Behaviour Research and Therapy*, 38, 759-777.
- Tabachnick, B. G., & Fidell, L. S. (2007). *Experimental designs using ANOVA*: Thomson/Brooks/Cole.

- Van Ameringen, M., Mancini, C., & Farvolden, P. (2003). The impact of anxiety disorders on educational achievement. *Journal of Anxiety Disorders, 17*(5), 561-571.
- Vassilopoulos, S. P., Banerjee, R., & Prantzalou, C. (2009). Experimental modification of interpretation bias in socially anxious children: Changes in interpretation, anticipated interpersonal anxiety, and social anxiety symptoms. *Behaviour Research and Therapy, 47*(12), 1085-1089.
- Vassilopoulos, S. P., Blackwell, S. E., Moberly, N. J., & Karahaliou, E. (2012). Comparing imagery and verbal instructions for the experimental modification of interpretation and judgmental bias in children. *Journal of behavior therapy and experimental psychiatry, 43*(1), 594-601.
- Vassilopoulos, S. P., Moberly, N. J., & Zisimatou, G. (2013). Experimentally Modifying Interpretations for Positive and Negative Social Scenarios in Children: A Preliminary Investigation. *Behavioural and Cognitive Psychotherapy, 41*(01), 103-116.
- Williams, A. D., O'Moore, K., Blackwell, S. E., Smith, J., Holmes, E. A., & Andrews, G. (2015). Positive imagery cognitive bias modification (CBM) and internet-based cognitive behavioral therapy (iCBT): A randomized controlled trial. *Journal of affective disorders, 178*, 131-141.
- Yao, N., Yu, H., Qian, M., & Li, S. (2015). Does attention redirection contribute to the effectiveness of attention bias modification on social anxiety?. *Journal of anxiety disorders, 36*, 52-62.
- Zimmermann, P., Wittchen, H.-U., Hofler, M., Pfister, H., Kessler, R. C., & Lieb, R. (2003). Primary anxiety disorders and the development of subsequent alcohol use disorders: a 4-

year community study of adolescents and young adults. *Psychological medicine*, 33(7), 1211-1222.