

*Infertility and ovarian follicle reserve depletion are associated with dysregulation of the FSH and LH receptor density in human antral follicles*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Regan, S. L. P., Knight, P. G. ORCID: <https://orcid.org/0000-0003-0300-1554>, Yovich, J. L., Stanger, J. D., Leung, Y., Arfuso, F., Dharmarajan, A. and Almahbobi, G. (2017) Infertility and ovarian follicle reserve depletion are associated with dysregulation of the FSH and LH receptor density in human antral follicles. *Molecular and Cellular Endocrinology*, 446. pp. 40-51. ISSN 0303-7207 doi: 10.1016/j.mce.2017.02.007 Available at <https://centaur.reading.ac.uk/69212/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.mce.2017.02.007>

Publisher: Elsevier

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Infertility and ovarian follicle reserve depletion are associated with  
dysregulation of the FSH and LH receptor density in human follicles**

Sheena L.P. Regan<sup>a\*</sup>, Phil G. Knight<sup>b</sup>, John L.Yovich<sup>c</sup>, James D. Stanger<sup>c</sup>, Yee Leung<sup>d</sup>, Frank Arfuso<sup>a</sup>,  
Arun Dharmarajan<sup>a</sup>, Ghanim Almahbobi<sup>e</sup>

<sup>a</sup>Stem Cell and Cancer Biology Laboratory, School of Biomedical Sciences, Curtin Health Innovation  
Research Institute, Curtin University, Perth, Australia. <sup>b</sup>School of Biological Sciences, Hopkins Building,  
University of Reading, Whiteknights, Reading RG6 6UB, UK. <sup>c</sup>PIVET Medical Centre, Perth, Australia.  
<sup>e</sup>Western Australian Gynaecologic Cancer Service, King Edward Memorial Hospital for Women, Perth,  
Australia. <sup>e</sup>School of Biomedical Sciences, Curtin Health Innovation Research Institute, Curtin University,  
Perth, Australia.

\* Dr Sheena LP Regan,

<sup>a</sup>Stem Cell and Cancer Biology Laboratory, School of Biomedical Sciences, Curtin Health Innovation  
Research Institute, Curtin University, Perth, Australia, GPO Box U1987, Perth, WA 6845, Australia  
Email: [sheenaregan@aapt.net.au](mailto:sheenaregan@aapt.net.au)

## Abstract

The low take-home baby rate in older women in Australia (5.8%) undergoing IVF is linked to the depletion of the ovarian reserve of primordial follicles. Oocyte depletion causes an irreversible change to ovarian function. We found that the young patient FSH receptor and LH receptor expression profile on the granulosa cells collected from different size follicles were similar to the expression profile reported in natural cycles in women and sheep. This was reversed in the older patients with poor ovarian reserve. The strong correlation of BMPR1B and FSH receptor density in the young was not present in the older women; whereas, the LH receptor and BMPR1B correlation was weak in the young but was strongly correlated in the older women. The reduced fertilisation and pregnancy rate was associated with a lower LH receptor density and a lack of essential down-regulation of the FSH and LH receptor. The mechanism regulating FSH and LH receptor expression appears to function independently, *in vivo*, from the dose of FSH gonadotrophin, rather than in response to it. Restoring an optimum receptor density may improve oocyte quality and the pregnancy rate in older women.



## 1. Introduction

As women age, the reserve of primordial follicles is depleted, and the quality of oocytes, fertilisation, and pregnancy rate are reduced. Following their initial recruitment from the ovarian reserve, activated primordial follicles grow and differentiate into pre-antral and small antral follicles (McGee and Hsueh, 2000). From the onset of puberty, cyclic fluctuations in follicle stimulating hormone (FSH) secretion from the anterior pituitary reach a threshold point sufficient to rescue a cohort of small antral follicles and initiate cyclic follicle recruitment (McGee and Hsueh, 2000). The number of antral follicles selected for dominance and ovulation is largely dependent on the regulatory action and the density of FSH receptors and LH receptors on the granulosa cell surface (Hillier, 2001, Baird, 1987, Baerwald, Adams and Pierson, 2012).

When the FSH level falls, the growth of the smaller follicles is reduced, and only the follicle with sufficient FSH and LH receptors continue to develop further because of their enhanced capacity to convert androstenedione to oestrogen for growth (Loumaye, Engrand, Shoham et al., 2003). As the ovarian primordial follicle reserve declines, the rate of cyclic recruitment of follicles diminishes (Baerwald et al., 2012, Almog, Shehata, Shalom-Paz et al., 2011). The number of these small antral follicles at the beginning of each cycle is representative of the ovarian reserve of primordial follicles that remain in the ovary.

Older patients, typically, have a slower follicle growth rate and a reduced number of granulosa cells per follicle (Santoro, Isaac, Neal-Perry et al., 2003). Other ovarian age related changes are associated with increased mitochondrial deletions in granulosa cells and reduced FSH receptor mRNA expression, which have been linked with infertility (González-Fernández, Peña, Hernández et al., 2010, Seifer, DeJesus and Hubbard, 2002, Cai, Lou, Dong et al., 2007). Reduced receptor density may directly contribute to poor oocyte quality by increasing the number of chromosomal errors (Maman, Yung, Kedem et al., 2012, Handyside, Montag, Magli et al., 2012).

In this study, the aim was to comprehensively profile the expression of granulosa FSH receptor and LH receptor protein in a range of patients of different ages and stages of ovarian primordial follicle depletion, who were receiving treatment for infertility. An average of ~8000 granulosa cells per follicle was collected from follicles ranging in size from 4 to 27 mm. Antibody labelling and flow cytometry were used to evaluate the receptor density. Previous studies of receptor expression have been confined to expression at the mRNA level, which may not be a reliable indicator of the level of translated 'mature' receptor protein expressed on the cell surface (Jeppesen, Kristensen, Nielsen et al., 2012, Pidoux, Gerbaud, Tsatsaris et al., 2007, Ascoli, Fanelli F and DL., 2002). The changes observed in receptor density may explain the adverse impact that ovarian reserve depletion has on fertility as women age.

## 2. Materials and Methods

### 2.1 Patients

A total of 415 follicles were collected from 56 patients undergoing standard fertility treatment with PIVET Medical Centre Perth, Western Australia, and are presented in Table 1. Patients were aged between 23 and 45 years, and follicles were collected irrespective of previous aetiology, but limited to exclude unusual medical conditions, hormonal dysfunction, and polycystic ovarian syndrome.

### 2.2 Human IVF: Ovarian stimulation, follicular fluid, and oocyte

Patient treatment consisted of two types of gonadotrophin releasing hormone-LH suppression (Puregon or Gonal F) in conjunction with commercially prepared recombinant human FSH, from cycle day 2 for ~10 days as previously described (Regan, Knight, Yovich et al., 2016). Ovulation was triggered with 10 000 IU HCG, and the collection of granulosa cells and oocyte retrieval was 36 hours later by transvaginal oocyte aspiration (Regan et al., 2016). *In vivo* human studies are more complex and therefore variables such as BMI have been minimised in this study (Table 1). We initially used the merino sheep model in a natural cycle (Regan, McFarlane, O'Shea et al., 2015) to establish a comparative data set with the human model during IVF treatment. The sheep and women both showed prerequisite down-regulation at the two critical follicle sizes that are equivalent to the size at the time of follicle selection and follicle maturation. Indeed,

the administration of the artificial LH surge to induce maturation appears not to alter the receptor expression in the young patient with a good ovarian reserve based on the similarities between the models and demonstrated in Fig.5.

Furthermore, analysis of the effect of rFSH dose on the receptor expression in a homogeneous group of patients with all variables controlled (ovarian reserve, age, follicle size, BMI, and AMH) was not significantly different (Fig. 6,  $p = 0.7$ ).

### **2.3 Ovarian reserve measured by the antral follicle count**

Patients received daily FSH according to a long established algorithm based on the patient's profile of age and ovarian reserve in order to predict the FSH dose required to stimulate multiple preovulatory follicles, as reported previously (Yovich, Stanger and Hinchliffe, 2012). Ovarian reserve was measured indirectly by the antral follicle count and was defined as the number of follicles between 2 - 10 mm in size that are present in total on ~day 2- 5 of a preliminary assessment cycle (Hansen, Hodnett, Knowlton et al., 2011).

The patients were divided into groups levels from A to E; good to poor ovarian reserve, respectively based on the algorithm, as described previously described (Regan et al., 2016), and a well-established clinical practice of patient treatment (Yovich et al., 2012): Poly cystic ovarian syndrome (PCOS) patients were excluded from the study group based on the Rotterdam criteria, initially prepared in 2003, and updated to reflect the advances in ultrasound technology(Lujan, Jarrett, Brooks et al., 2013): specifically; per ovary > 24 follicles, along with other criteria(Lujan et al., 2013). In the current study the combined ovary follicle total corresponded to Group A+ = 30-39 small follicles; group A = 20-29 small follicles; group B = 13-19 small follicles; group C = 9-12 small follicles; group D = 5-8 small follicles; group E =  $\leq 4$  small follicles.

### **2.4 Collection of granulosa cells**

The diameter of the follicle was calculated using ultrasonography as described previously (Regan et al., 2016). Flushing of the follicle by the clinician (Quinn's Advantage with Hepes, Sage Media, Pasadena, California) removed the loosely attached layers of granulosa cells. The cumulus ovarian complex was removed from the sample by the embryologist. The follicular fluid and flush was then layered onto a ficoll density gradient (555485; BD Biosciences, Perth, Australia) and centrifuged to isolate the granulosa cells

(Regan et al., 2016). Pure follicular fluid were analysed for oestrogen and progesterone using a random access immunoassay system (Siemens Medical Solutions, Bayswater, Victoria, Australia).

## **2.5 Immunolabelling of granulosa cells**

Aliquots of suspended granulosa cells ( $1 \times 10^6$  cells in 100  $\mu$ l) were immunolabelled using a double-indirect method as previously described (Regan et al., 2016). The antibodies have been used previously in human studies (Regan et al., 2016, Abir, Ben-Haroush, Melamed et al., 2008, Haÿ, Lemonnier, Fromigué et al., 2004), including flow cytometry analyses (Regan et al., 2016, Regan et al., 2015, Gao, De Geyter, Kossowska et al., 2007, Whiteman, Boldt J, Martinez J et al., 1991). Briefly, the cells were incubated with affinity purified goat polyclonal antibody for the FSH receptor (sc-7798), LH receptor (sc-26341), and BMPRII (sc-5679), (Santa Cruz Biotechnology, Santa Cruz, CA, USA), and then incubated with a donkey secondary antibody to goat IgG, conjugated to the fluorochrome Alexa 488 (Life Technologies Australia, Victoria, Australia), as described previously (Regan et al., 2016, Al-Samerria and Almahbobi, 2014 ). The routinely used monoclonal antibody CD45 was added to the LH receptor and BMPRII to enable the subtraction of the positive leukocyte common antigen to render a homogeneous population of granulosa cells (Fig. 1D).

Unstained samples or the substitution of a primary antibody with pre-immune goat IgG (Millennium Science, Surrey Hills, Victoria Australia) at the same concentration as the primary antibody served as a negative control for auto-fluorescence (Fig. 1A). A blocking peptide for the FSH receptor and BMPRII indicated nonspecific binding applied to human granulosa cells (sc-7798P, sc-5679P; Millennium Science, Surrey Hills, Victoria Australia), (Fig. 1B) as previously published (Cai et al., 2007, Gao et al., 2007). Pre-absorbed LH (Lutropin, Merck Serono, Frenchs Forest, NSW, Australia) also confirmed binding specificity (Fig. 1C, D). In the current study, the 'normal' goat IgG and unstained control cells emitted a similar average mean fluorescent intensity (MFI) and was subtracted from the receptor measurement. The auto-fluorescence and the nonspecific binding determined by the unstained control for each follicle was subtracted from each follicle as described previously (Regan et al., 2016). The data were analysed using FlowJo software (Tree Star Inc., Oregon, USA).

## 2.6 Microscopy

Re-suspended 10µl aliquots of FSH receptor immunolabelled, live granulosa cells were placed on slides and visualized using an Olympus DP 70 camera fitted to an Olympus BX-51 upright fluorescent microscope with a 40x UPlan N 0.4 N.A. objective (Olympus Imaging Australia, Macquarie Park, Australia) as described previously (Regan et al., 2016) (Fig.8B, D). Fluorescent microscopy revealed a positive staining of the cell membrane-bound FSH receptor as an intermittent, bright, ring-like pattern around the cells. All control samples showed negative staining. Granulosa cells ranged from 8 µm to 25 µm, with the average being 15 µm. We have validated the FSH receptor and LH receptor antibodies, and have published these findings (Al-Samerria and Almahbobi, 2014 ). 3D Image Analysis and immunofluorescent localisation and intensity quantification was performed on 10 um frozen sections in the sheep. In addition, this antibody has been used successfully by others in humans. Negative and positive controls were used and natural cell auto-fluorescence has been accounted for and subtracted for each sample, (Supplementary data Fig. S2) (Cai et al., 2007, Regan et al., 2015).

## 2.7 Flow cytometry

Selective gating of the whole sample to identify a pure granulosa cell population was achieved by graphing forward scatter to remove doublets (FSC-H verses FSC-A), presented in Fig. 1 and as previously described (Regan et al., 2016). The resulting population contained a uniform granulosa cell population that revealed positive staining for the FSH receptor, which is unique to granulosa cells (Hermann and Heckert, 2007). The data were analysed using FlowJo software (Tree Star Inc., Oregon, USA).

## 2.8 Statistics

Mean fluorescent intensity was obtained using ~8000 granulosa cells per individual follicle for the direct measurement of receptor protein expression. The data were subjected to statistical verification using one-way ANOVA with an uncorrected Fisher's LSD for follicular size using GraphPad Prism 6. Values in graphs are means ± S.E.M., and differences were considered significant if \*p<0.05, \*\*p<0.01, \*\*\*p<0.005, and \*\*\*\*p<0.001. The letter, such as 'a', signifies a statistical significant difference to the

matching letter (e.g. 'a\*'). The attached asterisk (a\*) indicates the significance level for the size follicle. A two tailed, student t-test was also used.

## **2.9 Human Ethics**

Informed consent was obtained from patients undergoing standard fertility treatment at PIVET Medical Centre, Perth, Australia, and from three patients undergoing risk reduction removal of the uterus and ovaries, who were recruited from King Edward Memorial Hospital (KEMH) Perth, Australia. Approval by the Human Research Ethics Committee of Curtin University of Technology and KEMH Women and Newborn Health Service ethics committee (WNHS) was obtained for this study (HR RD26-10:2010-2016), and all methods were performed in accordance with the relevant guidelines and regulations.

## **3. Results**

### **3.1 Follicle size and the extent of maturation of granulosa cells in the IVF patient**

During an IVF cycle, the granulosa cells from small follicles (8 mm) appear more compact and smaller in diameter (Fig. 2a). The granulosa cells from large follicles have a heterogeneous group of granulosa cells in different stages of maturation, which is referred to as luteinisation (Whiteman et al., 1991, Motta, 1969), (Fig. 2a, b, and c). The more mature granulosa cell has an expanded cytoplasm with prominent lipid droplets clustered around smooth endoplasmic reticulum, and are closely associated with numerous mitochondria as well as defined Golgi apparatus in the cytoplasm (Nottola, 1991). The extensive lipid droplets, observed in granulosa cells as grape-like spaces in the 23 mm follicles, indicate a greater steroid producing capacity and level of maturation. (Fig. 2c). The largest follicles produced a two fold increase in oestrogen and progesterone synthesis at the time of collection, Fig. 2d and 1e (Westergaard, Christensen and McNatty, 1986). High levels of progesterone synthesis by the granulosa cells are indicative of a greater extent of luteinisation even though all the follicles were exposed to an exogenous LH surge trigger injection.

The serum oestrogen peak was similar between the age groups; however, it is not until the ovarian reserve is used to distinguish the ageing process that more subtle differences are revealed. Peak serum oestrogen was approximately 6000 pmol/L for all age groups, but declined progressively as the ovarian reserve was depleted with increasing chronological age (Fig.3a). Age alone was not predictive of peak oestrogen levels, demonstrating no difference between age groups. There is a reduction in oestrogen output per follicle as the ovarian reserve is depleted, and is a reflection of the diminished number of follicles. Serum progesterone concentration was relatively uniform (2-4 nmol/L) but the lowest values were observed in the severely depleted patient groups of D and/or E, ( $p < 0.05$ , Fig. 3b).

### **3.2 The impact of age on FSH receptor and LH receptor density during follicle growth**

In the young (23-30 y) patient group with a good ovarian reserve (ovarian reserve group A+ & A), down-regulation of the FSH receptor was observed at follicle sizes corresponding to dominant follicle selection (10 mm,  $p = 0.0201$ ) and during the preovulatory maturation phase (24 mm,  $p = 0.0302$ ; Fig. 4a). The level of translated 'mature' LH receptor protein was expressed at a constant level on the granulosa cell surface of those follicles measuring 8 to 19 mm in the young patients, whereas the largest follicles (24 mm) expressed a significantly lower density of LH receptor ( $p = 0.0237$ , Fig. 4b).

In the older patients, a high level of granulosal FSH receptor was present in the very small antral follicles (4-8 mm, 35-40+ y, D & E), which was then reduced in a manner similar to that seen in younger patients with good ovarian reserve ( $p < 0.001$ , and  $p = 0.0259$ , respectively, Fig. 4a). The initial down-regulation was followed by a sequential up-regulation of FSH receptor in the largest follicles, in contrast to the decline observed in the young A+ & A patient group ( $p = 0.0013$ , and  $p = 0.0131$ , respectively, Fig. 4a). In the older patient group (35-40+ y group D & E) a significant increase in LH receptor was recorded in the 24-26 mm follicles ( $p = 0.0063$ , Fig. 4b). The LH receptor level after dominant follicle selection in the older patient group was significantly less than the level recorded in the young patient group (16 mm follicles:  $p = 0.0325$ , Fig. 4b).

### **3.3 The impact of ovarian reserve on receptor expression during follicular growth.**



In the 40+ y age group with a good ovarian reserve of B & C, FSH receptor down-regulation at the time of dominant follicle selection and maturation of the follicles ( $p < 0.01$  and  $p = 0.0182$ , respectively, Fig. 5a), was similar to the younger patients profile shown in Fig. 4. When the ovarian reserve was depleted within the same age group, both the FSH receptor and LH receptor expression at the time maturation was reversed (16 to 24 mm,  $p = 0.0286$  and 10 to 16 mm,  $p = 0.0411$ , respectively). The LH receptor was elevated in the largest follicles of the poor ovarian reserve 40+ y patient group D & E (19-23 mm,  $p = 0.0324$ , Fig. 5b).

The level of recombinant (r) FSH dose in the young was lower than the older patient group ( $p < 0.0001$ ) and consisted of ~100IU / day, whereas the older patients received an average of 450 IU / day (Fig. 5a). The effect of rFSH dose on FSH receptor or LH receptor expression was determined by administering either 300 or 600 IU / day for ~10 days during an IVF treatment cycle (Fig. 6b). The patients were matched for age (40-44y), ovarian reserve (AFC group E,  $< 5$  follicles, both ovaries combined), Anti-Mullerian hormone (AMH) ( $< 3.2$  p/mol/L), and size and range of follicles (10-22 mm). There was no significant difference in the effect of dose of rFSH on FSH receptor or LH receptor expression (Fig. 6b).

### **3.4 The relationship between BMPR1B and FSH receptor and LH receptor density as the ovarian reserve is depleted**

As the patient age increased, and the ovarian reserve declined, the correlation between BMPR1B and LH receptor density sequentially increased ( $R = 0.872$ ,  $p = 0.0063$ , Fig. 7a). The reversal of this correlation was observed between FSH receptor and BMPR1B. In the young patient group, the BMPR1B was aligned with the FSH receptor expression ( $R = 0.75$ ,  $p = 0.0044$ , Fig. 7b), followed by a complete dissociation in the relationship as ovarian reserve deteriorated ( $p = 0.4$ ). FSH receptor and LH receptor expression were not strongly correlated (Supplementary data Fig. S1).

## **4. Discussion**

Depletion of the ovarian reserve of primordial follicles has a considerable impact on young and older women. High achieving women, in particular, delay having children while they establish careers in business, elite sport, music, and other pursuits, only to find that the poor quality and reduced quantity of



their remaining oocytes prevents them from reproducing (Australian-Bureau-of-Statistics, 2001,Australian-Bureau-Statistics and 2006). In addition, the one child government policy in China has relaxed, creating the demand for a second child for older women. This novel, human, *in vivo* analysis reports the change in granulosa FSH receptor and LH receptor expression that occurs as the ovarian reserve is depleted in the young compared to the older women during IVF treatment. The present study is robust because of the in depth analysis of individual follicle sizes rather than pooled sample and the large number of granulosa cells analysed from each follicle via immunofluorescent flow cytometry.

The major findings of the study reveal that: 1. the young patient FSH receptor and LH receptor profile of expression on the granulosa cells collected from different size follicles were similar to the expression profile reported in natural cycles in women and sheep (Fig. 4) (Jeppesen et al., 2012,Regan et al., 2015). Conversely, in the older patients with poor ovarian reserve, down-regulation of the FSH receptor and LH receptor was not observed which may indicate reduced or delayed maturation of the granulosa cells in preparation for ovulation of the oocyte.

2. The overall effect of the disruption to receptor expression resulted in a shift in the strong correlation of BMPRII and FSH receptor density in the young as the ovarian reserve was depleted. Conversely, the BMPRII expression became strongly aligned with the density of LH receptor in the older women. This change indicates reduced or delayed granulosa cell luteinisation, and confirmed by the reduced progesterone in the poor ovarian reserve patients (Fig. 3 & 7). 3. In addition, impaired oocyte development was associated with a lower LH receptor density and a lack of essential down-regulation of the FSH and LH receptor; however additional contributing factors would also prevail. The poor oocyte quality was evident by the poor fertilisation rate, reduced pregnancy, and lower live birth rate (Table 1). 4. The gonadotrophin FSH has been strongly associated with the regulatory control of cyclic folliculogenesis (Yong, Baird, Yates et al., 1992). However, the mechanism regulating the expression of FSH receptor and LH receptor functions independently, *in vivo*, from the dose of gonadotrophin rFSH rather than in response to it in an *in vitro* scenario.

IVF stimulation of patients using rFSH extends the window of recruitment that promotes multiple dominant follicles. These follicles have been shown to grow at a similar rate compared to a natural cycle (Baird, 1987, Fauser and Van Heusden, 1997). We report that in patients matched for ovarian reserve, BMI, AMH, age, and follicle number and size, the impact of rFSH dose on receptor expression *in vivo* was not significant (Fig. 6b). Whereas, isolated granulosa cells in culture, responded differently to high concentrations of rFSH (Zhang and Roy, 2004). The time of dominant follicle selection and steroidogenic maturation would therefore progress in a similar manner to a natural cycle.

The high level of FSH receptors in the smallest follicles provides further support of a spatio-temporal range of differentiation of the follicles at the time of collection in an IVF cycle even though the LH surge has taken place. Indeed, the small follicles of 4-8 mm were morphologically granular in appearance with a relatively large nucleus compared to the cytoplasm, which indicates a follicular granulosa cell, and not a granulosa luteal cell (Fig. 2) (Nottola, Heyn, Camboni et al., 2006). It is therefore reasonable to relate the observed FSH receptor and LH receptor density profile to the underlying stages of antral follicle development, as indicated by the size of the follicle. Two critical stages of follicle development are dominant follicle selection (8 mm) and internalisation of receptors in an 'ovulatory follicle' (largest follicle).

The follicles were also shown to have a different follicular fluid steroid output based on size (Fig. 2d and e). The granulosa cells from a small follicle produced less oestrogen and progesterone compared to a larger 'ovulatory follicle', even though the follicles were exposed to the same gonadotrophin stimulation and the equivalent LH surge-ovulation-trigger, human chorionic gonadotrophin (HCG) prior to collection. Therefore, it remains that the mechanism regulating the expression of these receptors is independent, *in vivo*, from the gonadotrophin rFSH dose. In further support of this, post-transcriptional mechanisms have been identified in the regulation of these receptors (Menon and Menon, 2012).

In contrast, the clinical administration of HCG or the equivalent natural cycle LH-surge, induces many maturation changes referred to as luteinisation of the granulosa cell. These changes include degradation or internalisation of the BMPRII (Regan et al., 2016, Regan et al., 2015), LH receptor (Menon and Menon, 2012), and FSH receptor (Regan et al., 2015), cytoskeletal reorganisation of the granulosa cell, cessation of mitogenic proliferation, cumulus expansion, gap junction closure, resumption of meiosis, and general maturation of the oocyte (Izadyar, Zeinstra and Bevers, 1998, Fan, Liu, Shimada et al., 2009). Importantly, the lack of essential down-regulation of the receptors indicate disruption of this process, and may affect the LH surge-induced changes taking place at this time (Lyga, Volpe, Werthmann et al., 2016, Kash and Menon, 1998). However, the extent of luteinisation is dependent on the stage of development of the follicle at the time of the HCG/LH surge, as shown in Fig. 2, and also reported previously (Nottola et al., 2006). In particular, the level of LH receptor density during follicle development was consistent except for the largest ‘ovulatory’ follicles (Fig. 4). These large follicles are developmentally receptive to the LH surge-induced internalisation of the LH receptor (Menon and Menon, 2012) even though they were exposed to the same dose of HCG/LH surge trigger during IVF treatment (Fig. 2). This difference in response based on follicle size further supports our finding of differential maturation of the follicles in an IVF cycle. The ovulatory follicle internalisation of the LH receptor was also evident in natural cycles of women (Jeppesen et al., 2012) and in sheep studies conducted by our research group using the same analysis techniques (Regan et al., 2015).

The granulosa cells in the current study were collected from individual follicles and analysed immediately (fresh not frozen) to reduce any potential experiment-induced internalisation of receptors. Whereas, granulosa cells collected for culture were not responsive to rFSH stimulation (Breckwoldt, Selvaraj, Aharoni et al., 1996, Gutierrez, Campbell and Webb, 1997). The desensitised cells start to re-express receptors after 2-5 days of culture (Ophir, Yung, Maman et al., 2014). The lack of responsiveness of the FSH receptors to rFSH stimulation is consistent with our *in vivo* finding because typically, the largest, more prominent follicle will be collected for cell culture as it contains the greatest number of cells, and is

pooled with any smaller follicles. The pooling of the large and smaller follicles would mask the differences presented in the current study, and as previously documented (Regan et al., 2016).

The acquisition of LH receptors on the granulosa cells coincides with the first down-regulation of FSH receptors to promote dominant follicle selection (Maman et al., 2012, Jeppesen et al., 2012, Rice, Ojha, Whitehead et al., 2007, Sen, Prizant, Light et al., 2014). A high expression of the FSH receptor mRNA has been reported in granulosa cells from small antral follicles of ~6 mm, collected from a wide range of patients in natural cycles (7-38 years) (Jeppesen et al., 2012). The elevated FSH receptor levels were followed by a lower level of expression in ~9 and 15 mm follicles. The current study confirms and expands their data by reporting that at the time of dominant follicle selection (~8 mm follicles), a high level of granulosa FSH receptor density is observed in both the young and the older women regardless of ovarian reserve (Fig. 5). This is consistent with a metabolic change from high FSH dependency before dominant follicle selection and LH receptor expression induced by the high FSH receptor density (Rice et al., 2007, Sen et al., 2014), in both a natural cycle and a stimulated IVF cycle (Jeppesen et al., 2012). Therefore, the mechanism regulating early FSH receptor expression appears to be independent of either pituitary secreted FSH in a natural cycle or dose of rFSH received during IVF treatment.

This novel finding is consistent with the *in vivo*, insignificant effect of the dose of rFSH and receptor expression of either the FSH or LH receptor (Fig. 6b). Furthermore, it may be expected that if FSH receptors induce LH receptor expression (Rice et al., 2007, Sen et al., 2014), then a reduced ovarian reserve should not affect the expression of the LH receptors in the older women. However, it is apparent that the LH receptor density was reduced after dominant follicle selection when compared to the young patient group (16 mm, Fig. 4b). Therefore, other factors must influence the expression of LH receptors. Moreover, a reduced LH receptor density in the follicles in older women may be a contributing factor in the reduced quality of the oocyte and the resulting poor pregnancy rate (Table 1). The 35-38 year old group of women have a greater percentage of

women with the better B grade (45%) of ovarian reserve and a much lower rate of women with the poor ovarian reserve group of D (32.6%) and E group (5.9%) compared to the older women from 39-45y, (Fig. 8). Age has a direct impact on the frequency of chromosomal errors and fertility (Handyside et al., 2012). We demonstrate that the reduced fertility as we age is associated with a patient's ovarian reserve and the dysregulation of receptor signalling.

Expression of the type 1 transforming growth factor beta (TGF $\beta$ ) superfamily receptor, BMPR1B, has been shown to be down-regulated at the two critical stages of dominant follicle selection and prior to ovulation (Regan et al., 2016, Regan et al., 2015, Nakamura, Otsuka, Inagaki et al., 2012, Ogura Nose, Yoshino, Osuga et al., 2012, Miyoshi, Otsuka, Inagaki et al., 2007, Erickson and Shimasaki, 2003). The BMPR1B is the common receptor for several BMP ligands such as BMP 2, 4, 6, 7, and 15 (Knight and Glistler, 2003), and inhibits progesterone synthesis in favour of oestrogen synthesis during follicular growth. The BMP-induced inhibition prevents the onset of the LH surge-induced luteinisation (Pierre, Pisselet, Dupont et al., 2004, Tajima, Dantes, Yao et al., 2003, Val, Lefrançois-Martinez, Veyssi re et al., 2003, Abdo, Hisheh, Arfuso et al., 2008). In particular, at the time of the LH surge/ HCG trigger, the granulosa BMPR1B, FSH receptor, and LH receptor density of the largest follicles was shown to be down-regulated in sheep during a natural cycle (Regan et al., 2015), in women receiving IVF treatment (Regan et al., 2016), and in the current study (Fig. 4). The lack of down-regulation of the FSH receptor, LH receptor (Fig. 4), and BMPR1B (Regan et al., 2016) may delay or inhibit the production of progesterone and expression of progesterone receptors (Shimada, Nishibori, Isobe et al., 2003), (Fig. 3), which would directly impair germinal vesicle breakdown and meiotic resumption in the oocyte (Shimada and Terada, 2002). The inhibition of gonadotrophin-induced progesterone receptor formation has been linked to infertility and porcine oocyte quality (Park-Sarge and Mayo, 1994), and had a negative effect on the ovarian response to gonadotropin stimulation (Cai et al., 2007). Therefore, it is essential that the BMP inhibition is attenuated to promote maturation of the oocyte.

In the current study, a strong correlation between BMPR1B and FSH receptor expression was observed in the young patients. The FSH receptor is positively regulated by the BMP ligands 6 and 7 (Chen, Yu, Wang

et al., 2008, Shi, Yoshino, Osuga et al., 2010, Shi, Yoshino, Osuga et al., 2009), and as the ovarian reserve was depleted, the correlation was weakened (Fig. 7a). Significantly, the disruption to the FSH receptor expression was replaced with a strong correlation between BMPRII and the LH receptor expression in the older women (Fig. 7b). This shift further supports an association between dysregulated receptor expression and reduced maturation of the granulosa cells surrounding the oocyte in older women. Future research may involve restoring the optimum receptor density profile during the maturation phase to improve oocyte quality.

### **Acknowledgements**

The authors thank all the participants who generously donated their samples to this study, the clinical doctors, embryologists, and nursing staff.

### **Authors' roles**

SLPR conceived the study, experimental design, conducted all experiments, the analysis and interpretation of data, wrote the first draft of the manuscript and the final version of the paper. Obtained informed consent from patients and ethics approval. PK supervised, interpretation of data, contributed to the draft of the manuscript, interpretation of data, and critically revised the manuscript. JLY supervised, participated in the study design, participated in obtaining granulosa cells, interpretation of data, and critically revised the manuscript. JDS supervised, participated in obtaining granulosa cells, participated in the design of the study, and critically revised the manuscript. YL supervised, participated in the study design, obtained informed consents from patients and ethics approval, and critically revised the manuscript. FA supervised, contributed to the draft of the manuscript, interpretation of data, and critically revised the manuscript. AD supervised, participated in the study design, interpretation of data, contributed to the draft of the manuscript, and critically revised the manuscript. GA supervised, conceived the study, participated in the study design, interpretation of data, and critically revised the manuscript.

### **Funding**

SLPR was a recipient of an Australian Postgraduate Award and a Curtin University Postgraduate Scholarship. This work was supported by additional private external funding, which was gratefully accepted from Denby Macgregor. This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

# **Conflict of interest**

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

# **References**

- [1] McGee, E.A. and Hsueh, A.J.W., 2000. Initial and cyclic recruitment of ovarian follicles, *Endocrine Reviews*. 21, 200-214.
- [2] Hillier, S.G., 2001. Gonadotropic control of ovarian follicular growth and development, *Molecular and Cellular Endocrinology*. 179, 39-46.
- [3] Baird, D.T., 1987. A model for follicular selection and ovulation: lessons from superovulation, *The Journal of steroid biochemistry*. 27, 15-23.
- [4] Baerwald, A.R., Adams, G.P. and Pierson, R.A., 2012. Ovarian antral folliculogenesis during the human menstrual cycle: a review, *Human Reproduction Update*. 18, 73-91.
- [5] Loumaye, E., Engrand, P., Shoham, Z., Hillier, S.G. and Baird, D.T., 2003. Clinical evidence for an LH ceiling?, *Human Reproduction*. 18, 2719-2720.
- [6] Almog, B., Shehata, F., Shalom-Paz, E., Tan, S.L. and Tulandi, T., 2011. Age-related normogram for antral follicle count: McGill reference guide, *Fertility and Sterility*. 95, 663-666.
- [7] Santoro, N., Isaac, B., Neal-Perry, G., Adel, T., Weingart, L., Nussbaum, A., Thakur, S., Jinnai, H., Khosla, N. and Barad, D., 2003. Impaired Folliculogenesis and Ovulation in Older Reproductive Aged Women, *The Journal of Clinical Endocrinology & Metabolism*. 88, 5502-5509.



- 425 [8] González-Fernández, R., Peña, Ó., Hernández, J., Martín-Vasallo, P., Palumbo, A. and  
 426 Ávila, J., 2010. FSH receptor, KL1/2, P450, and PAPP genes in granulosa-lutein cells  
 427 from in vitro fertilization patients show a different expression pattern depending on the  
 428 infertility diagnosis, *Fertility and Sterility*. 94, 99-104.
- 429 [9] Seifer, D.B., DeJesus, V. and Hubbard, K., 2002. Mitochondrial deletions in luteinized  
 430 granulosa cells as a function of age in women undergoing in vitro fertilization, *Fertility  
 431 and Sterility*. 78, 1046-1048.
- 432 [10] Cai, J., Lou, H., Dong, M., Lu, X., Zhu, Y., Gao, H. and Huang, H., 2007. Poor ovarian  
 433 response to gonadotropin stimulation is associated with low expression of follicle-  
 434 stimulating hormone receptor in granulosa cells, *Fertility and Sterility*. 87, 1350-1356.
- 435 [11] Maman, E., Yung, Y., Kedem, A., Yerushalmi, G.M., Konopnicki, S., Cohen, B., Dor, J.  
 436 and Hourvitz, A., 2012. High expression of luteinizing hormone receptors messenger  
 437 RNA by human cumulus granulosa cells is in correlation with decreased fertilization,  
 438 *Fertility and Sterility*. 97, 592-598.
- 439 [12] Handyside, A.H., Montag, M., Magli, M.C., Repping, S., Harper, J., Schmutzler, A.,  
 440 Vesela, K., Gianaroli, L. and Geraedts, J., 2012. Multiple meiotic errors caused by  
 441 predivision of chromatids in women of advanced maternal age undergoing in vitro  
 442 fertilisation, *European Journal of Human Genetics*. 20, 742-747.
- 443 [13] Jeppesen, J.V., Kristensen, S.G., Nielsen, M.E., Humaidan, P., Dal Canto, M., Fadini, R.,  
 444 Schmidt, K.T., Ernst, E. and Yding Andersen, C., 2012. Lh-receptor gene expression in  
 445 human granulosa and cumulus cells from antral and preovulatory follicles, *The Journal of  
 446 Clinical Endocrinology & Metabolism*. 97, E1524-E1531.
- 447 [14] Pidoux, G., Gerbaud, P., Tsatsaris, V., Marpeau, O., Ferreira, F., Meduri, G.,  
 448 Guibourdenche, J., Badet, J., Evain-Brion, D. and Frendo, J.-L., 2007. Biochemical



characterization and modulation of LH/CG—receptor during human trophoblast differentiation, *Journal of cellular physiology*. 212, 26-35.

[15] Ascoli, M., Fanelli F and DL., S., 2002. The lutropin/choriogonadotropin receptor, a 2002 perspective, *Endocrine reviews*. Vol.23(2), 141-174

[16] Regan, S.L.P., Knight, P.G., Yovich, J., Stanger, J., Leung, Y., Arfuso, F., Dharmarajan, A. and Almahbobi, G., 2016. Dysregulation of granulosa bone morphogenetic protein receptor 1B density is associated with reduced ovarian reserve and the age-related decline in human fertility, *Molecular and Cellular Endocrinology*. 425, 84-93.

[17] Regan, S.L.P., McFarlane, J.R., O'Shea, T., Andronicos, N., Arfuso, F., Dharmarajan, A. and Almahbobi, G., 2015. Flow cytometric analysis of FSHR, BMRR1B, LHR and apoptosis in granulosa cells and ovulation rate in merino sheep, *Reproduction*. 150, 151-163.

[18] Yovich, J., Stanger, J. and Hinchliffe, P., 2012. Targeted gonadotrophin stimulation using the PIVET algorithm markedly reduces the risk of OHSS, *Reproductive BioMedicine Online*. 24, 281-292.

[19] Hansen, K.R., Hodnett, G.M., Knowlton, N. and Craig, L.B., 2011. Correlation of ovarian reserve tests with histologically determined primordial follicle number, *Fertility and Sterility*. 95, 170-175.

[20] Lujan, M.E., Jarrett, B.Y., Brooks, E.D., Reines, J.K., Peppin, A.K., Muhn, N., Haider, E., Pierson, R.A. and Chizen, D.R., 2013. Updated ultrasound criteria for polycystic ovary syndrome: reliable thresholds for elevated follicle population and ovarian volume, *Human Reproduction*. 28, 1361-1368.

[21] Abir, R., Ben-Haroush, A., Melamed, N., Felz, C., Krissi, H. and Fisch, B., 2008. Expression of bone morphogenetic proteins 4 and 7 and their receptors IA, IB, and II in human ovaries from fetuses and adults, *Fertility and Sterility*. 89, 1430-1440.

- 474 [22] Haÿ, E., Lemonnier, J., Fromigué, O., Guénou, H. and Marie, P.J., 2004. Bone  
475 morphogenetic protein receptor signaling mediates apoptosis independently of  
476 differentiation in osteoblastic cells, *Journal of Biological Chemistry*. 279, 1650-1658.
- 477 [23] Gao, S., De Geyter, C., Kossowska, K. and Zhang, H., 2007. FSH stimulates the  
478 expression of the ADAMTS-16 protease in mature human ovarian follicles, *Molecular*  
479 *Human Reproduction*. 13, 465-471.
- 480 [24] Whiteman, G., Boldt J, Martinez J and Pantazis C, 1991. Flow cytometric analysis of  
481 induced human graafian follicles.I. Demonstration and sorting of two luteinized cell  
482 populations, *Fertility and Sterility*. 56, 259-263.
- 483 [25] Al-Samerria, S. and Almahbobi, G., 2014 Three-dimensional image analysis to quantify  
484 the temporospatial expression of cellular receptors, *Journal of Medical and*  
485 *Bioengineering* 3, 179-182.
- 486 [26] Hermann, B.P. and Heckert, L.L., 2007. Transcriptional regulation of the FSH receptor:  
487 New perspectives, *Molecular and Cellular Endocrinology*. 260–262, 100-108.
- 488 [27] Motta, P.M., 1969. Electron microscope study on the human lutein cell with special  
489 reference to its secretory activity., *Z Zeltforsch Mikrosk Anat*. 98, 233-245.
- 490 [28] Nottola, S., Familiari G, Micara G, Aragona C, Motta PM, 1991 The ultrastructure of  
491 human cumulus-corona cells at the time of fertilization and early embryogenesis. A  
492 scanning and transmission electron microscopic study in an in vitro fertilization program,  
493 *Arch Histol Cytol*. . May;54(2):145-61.
- 494 [29] Westergaard, L., Christensen, I.J. and McNatty, K.P., 1986. Steroid levels in ovarian  
495 follicular fluid related to follicle size and health status during the normal menstrual cycle  
496 in women, *Human Reproduction*. 1, 227-232.
- 497 [30] Australian-Bureau-of-Statistics, 2001. Australian social trends 2001 Canberra.

- 498 [31] Australian-Bureau-Statistics and 2006. Births Australia 2005 ABS, Canberra., cat. no.  
499 33010.
- 500 [32] Yong, E.L., Baird, D.T., Yates, R., Reichert, L.E. and Hillier, S.G., 1992. Hormonal  
501 regulation of the growth and steroidogenic function of human granulosa cells, *The Journal*  
502 *of clinical endocrinology and metabolism*. 74, 842-849.
- 503 [33] Fauser, B.C. and Van Heusden, A.M., 1997. Manipulation of human ovarian function:  
504 physiological concepts and clinical consequences, *Endocrine reviews*. 18, 71-106.
- 505 [34] Zhang, Y.-M. and Roy, S.K., 2004. Downregulation of follicle-stimulating hormone (fsh)-  
506 receptor messenger rna levels in the hamster ovary: Effect of the endogenous and  
507 exogenous FSH, *Biology of Reproduction*. 70, 1580-1588.
- 508 [35] Nottola, S.A., Heyn, R., Camboni, A., Correr, S. and Macchiarelli, G., 2006.  
509 Ultrastructural characteristics of human granulosa cells in a coculture system for in vitro  
510 fertilization, *Microscopy Research and Technique*. 69, 508-516.
- 511 [36] Menon, K.M.J. and Menon, B., 2012. Structure, function and regulation of gonadotropin  
512 receptors- a perspective, *Molecular and Cellular Endocrinology*. 356, 88-97.
- 513 [37] Izadyar, F., Zeinstra, E. and Bevers, M.M., 1998. Follicle-stimulating hormone and  
514 growth hormone act differently on nuclear maturation while both enhance developmental  
515 competence of in vitro matured bovine oocytes, *Molecular reproduction and development*.  
516 51, 339-345.
- 517 [38] Fan, H., Liu, Z., Shimada, M., Sterneck, E., Johnson, P.F., Hedrick, S.M. and Richards,  
518 J.S., 2009. MAPK3/1 (ERK1/2) in ovarian granulosa cells are essential for female fertility,  
519 *Science*,. 324, 938-941.
- 520 [39] Lyga, S., Volpe, S., Werthmann, R.C., Götz, K., Sungkaworn, T., J., L.M. and Calebiro,  
521 D., 2016. Persistent cAMP signaling by internalized LH receptors in ovarian follicles,  
522 *Endocrinology*. 157(4), 1613-1621.

- 523 [40] Kash, J.C. and Menon, K.M.J., 1998. Identification of a hormonally regulated luteinizing  
524 hormone/human chorionic gonadotropin receptor mRNA binding protein: Increased mRNA  
525 binding during receptor down-regulation, *Journal of Biological Chemistry*. 273, 10658-  
526 10664.
- 527 [41] Breckwoldt, M., Selvaraj, N., Aharoni, D., Barash, A., Segal, I., Insler, V. and  
528 Amsterdam, A., 1996. Cell proliferation and apoptosis: Expression of Ad4-BP/cytochrome  
529 P450 side chain cleavage enzyme and induction of cell death in long-term cultures of  
530 human granulosa cells, *Molecular Human Reproduction*. 2, 391-400.
- 531 [42] Gutierrez, C., Campbell, B. and Webb, R., 1997. Development of a long-term bovine  
532 granulosa cell culture system: induction and maintenance of estradiol production, response  
533 to follicle-stimulating hormone, and morphological characteristics, *Biology of*  
534 *Reproduction*. 56, 608 - 616.
- 535 [43] Ophir, L., Yung, Y., Maman, E., Rubinstein, N., Yerushalmi, G.M., Haas, J., Barzilay, E.  
536 and Hourvitz, A., 2014. Establishment and validation of a model for non-luteinized human  
537 mural granulosa cell culture, *Molecular and Cellular Endocrinology*. 384, 165-174.
- 538 [44] Rice, S., Ojha, K., Whitehead, S. and Mason, H., 2007. Stage-specific expression of  
539 androgen receptor, follicle-stimulating hormone receptor, and anti-müllerian hormone  
540 type II receptor in single, isolated, human preantral follicles: Relevance to polycystic  
541 ovaries, *The Journal of Clinical Endocrinology & Metabolism*. 92, 1034-1040.
- 542 [45] Sen, A., Prizant, H., Light, A., Biswas, A., Hayes, E., Lee, H.-J., Barad, D., Gleicher, N.  
543 and Hammes, S.R., 2014. Androgens regulate ovarian follicular development by  
544 increasing follicle stimulating hormone receptor and microRNA-125b expression,  
545 *Proceedings of the National Academy of Sciences*. 111(8), 3008-3013.
- 546 [46] Nakamura, E., Otsuka, F., Inagaki, K., Miyoshi, T., Matsumoto, Y., Ogura, K.,  
547 Tsukamoto, N., Takeda, M. and Makino, H., 2012. Mutual regulation of growth hormone

- 548 and bone morphogenetic protein system in steroidogenesis by rat granulosa cells,  
549 Endocrinology. 153, 469-480.
- 550 [47] Ogura Nose, S., Yoshino, O., Osuga, Y., Shi, J., Hiroi, H., Yano, T. and Taketani, Y.,  
551 2012. Anti-Mullerian hormone (AMH) is induced by bone morphogenetic protein (BMP)  
552 cytokines in human granulosa cells, European journal of obstetrics & gynecology and  
553 reproductive biology. 164, 44-47.
- 554 [48] Miyoshi, T., Otsuka, F., Inagaki, K., Otani, H., Takeda, M., Suzuki, J., Goto, J., Ogura, T.  
555 and Makino, H., 2007. Differential regulation of steroidogenesis by bone morphogenetic  
556 proteins in granulosa cells: Involvement of extracellularly regulated kinase signaling and  
557 oocyte actions in follicle-stimulating hormone-induced estrogen production,  
558 Endocrinology. 148, 337-345.
- 559 [49] Erickson, G. and Shimasaki, S., 2003. The spatiotemporal expression pattern of the bone  
560 morphogenetic protein family in rat ovary cell types during the estrous cycle, Reprod Biol  
561 Endocrinol. 1(9), 1-20.
- 562 [50] Knight, P.G. and Glister, C., 2003. Local roles of TGF- $\beta$  superfamily members in the  
563 control of ovarian follicle development, Animal Reproduction Science. 78, 165-183.
- 564 [51] Pierre, A., Pisselet, C., Dupont, J., Mandon-Pepin, B., Monniaux, D., Monget, P. and  
565 Fabre, S., 2004. Molecular basis of bone morphogenetic protein-4 inhibitory action on  
566 progesterone secretion by ovine granulosa cells, J Mol Endocrinol. 33, 805 - 814.
- 567 [52] Tajima, K., Dantes, A., Yao, Z., Sorokina, K., Kotsuji, F., Seger, R. and Amsterdam, A.,  
568 2003. Down-regulation of steroidogenic response to gonadotropins in human and rat  
569 preovulatory granulosa cells involves mitogen-activated protein kinase activation and  
570 modulation of DAX-1 and steroidogenic factor-1, The Journal of clinical endocrinology  
571 and metabolism. 88, 2288-2299.

- 572 [53] Val, P., Lefrançois-Martinez, A.-M., Veyssi re, G. and Martinez, A., 2003. SF-1 a  
573 key player in the development and differentiation of steroidogenic tissues, Nuclear  
574 receptor. 1, 8-8.
- 575 [54] Abdo, M., Hisheh, S., Arfuso, F. and Dharmarajan, A., 2008. The expression of tumor  
576 necrosis factor-alpha, its receptors and steroidogenic acute regulatory protein during  
577 corpus luteum regression, Reproductive Biology and Endocrinology : RB&E. 6, 1-11.
- 578 [55] Shimada, M., Nishibori, M., Isobe, N., Kawano, N. and Terada, T., 2003. Luteinizing  
579 hormone receptor formation in cumulus cells surrounding porcine oocytes and its role  
580 during meiotic maturation of porcine oocytes, Biology of Reproduction. 68, 1142-1149.
- 581 [56] Shimada, M. and Terada, T., 2002. FSH and LH induce progesterone production and  
582 progesterone receptor synthesis in cumulus cells: a requirement for meiotic resumption in  
583 porcine oocytes, Molecular Human Reproduction. 8, 612-618.
- 584 [57] Park-Sarge, O.-K. and Mayo, K.E., 1994. Molecular biology of endocrine receptors in the  
585 ovary, in: Findlay, J.K. (Ed.), Molecular Biology of the Female Reproductive System.  
586 Academic Press, San Diego, pp. 153-205.
- 587 [58] Chen, A.Q., Yu, S., Wang, Z., Xu, Z. and Yang, Z., 2008. Stage-specific expression of  
588 bone morphogenetic protein type I and type II receptor genes: Effects of follicle-  
589 stimulating hormone on ovine antral follicles, Animal Reproduction Science. 111, 391-  
590 399.
- 591 [59] Shi, J., Yoshino, O., Osuga, Y., Nishii, O., Yano, T. and Taketani, Y., 2010. Bone  
592 morphogenetic protein 7 (BMP-7) increases the expression of follicle-stimulating  
593 hormone (FSH) receptor in human granulosa cells, Fertility and Sterility. 93, 1273-1279.
- 594 [60] Shi, J., Yoshino, O., Osuga, Y., Koga, K., Hirota, Y., Hirata, T., Yano, T., Nishii, O. and  
595 Taketani, Y., 2009. Bone morphogenetic protein-6 stimulates gene expression of follicle-

596 stimulating hormone receptor, inhibin/activin beta subunits, and anti-Müllerian hormone  
597 in human granulosa cells, Fertility and Sterility. 92, 1794-1798.

598

599

**\*Highlights (for review)**

- Reproductive ageing is linked to ovarian cellular function and infertility
- Granulosa FSHR and LHR density from 327 ovarian follicles from IVF patients
- Prerequisite FSHR and LHR down-regulation in older patients was not observed
- Ovarian reserve-impaired fertility was associated with lower granulosa cell LHR
- Ovarian reserve was linked to poor oocyte quality; fertilisation and pregnancy rate



**Table 1 Patient ovarian reserve, based on antral follicle count (AFC) and the number of follicles collected per group.**

AGE Year	IVF Patient	Total Follicle	BMI	Ovarian Reserve Group Follicles Collected						Fertility %			
				A+	A	B	C	D	E	Failed Fertilisation	Not Pregnant	Pregnant	Live Birth
23-30	11	95	24.1±4	31	64					0	36	**64	43
35-45	34	232	24.8±5			88	21	99	24	9	52	**39	18
*39-45	19	131	23.9±5			42	5	66	18	17	72	11	6


Ovarian reserve measured indirectly by the antral follicle count (AFC). Antral follicle count is the number of

follicles between 2-10 mm on day 2-5 of a cycle: A+ = 30-39 follicles; A = 20-29; B = 13-19; C = 9-12; D =


5-8; E = ≤4. Follicle count is based on the combined total from both ovaries. \*Subgroup of older patients.

\*\*1 Ectopic pregnancy. Frozen embryo transfers cycles included.


Figure



Figure




Figure



Figure


a

FSH receptor




b

LH receptor




OVARIAN  
RESERVE

GOOD	A+ = 30-39	A = 20-29	B = 13-19	C = 9-12	D = 5-8	E = ≤4	POOR
------	------------	-----------	-----------	----------	---------	--------	------

**a** FSH Receptor**b**


## LH Receptor




**OVARIAN  
RESERVE**

GOOD A+ = 30-39 A = 20-29 B = 13-19 C = 9-12 D = 5-8 E = ≤4 POOR

Figure




Figure





Figure



**Fig. 1 Validation of immunofluorescent labelling.**

**A.** Live granulosa cells, unstained control for FSH receptor auto-fluorescence (blue) compared to positive fluorescent signal (box). **B.** Live human granulosa cells with positive fluorescence for FSH receptor (a & b), and negative blocking agent for FSH receptor (c & d). Bar 10  $\mu$ m. **C.** Live granulosa cells, unstained control for LH receptor auto-fluorescence (red) compared to positive fluorescent signal (box), Gated and removed CD45 positive cells (circle). **D.** Live human granulosa cells with positive fluorescence for LH receptor (i & j), and negative blocking agent for LH receptor (k & l). Bar 10  $\mu$ m.

**Fig. 2 Comparison of granulosa cells from small, medium, and large human IVF follicles at different stages of maturation.**

a. Small antral follicle of 8 mm, compact morphology with large nucleus compared to cytoplasm. b. Medium size antral follicle of 14 mm, showing some larger granulosa cells, and many still with compact morphology similar to the small antral follicle. c. Granulosa cells from a large antral follicle of 24 mm, showing expanded cytoplasm with many lipid droplet spaces, arrows. Very few granulosa cells with a compact morphology (small dotted box), and the majority with an expanded cytoplasm (larger box). Air dried fresh samples were placed in a fume glass containing formalherhyde for 10 min and then stained with Oil red O, and Harris counterstained. Light microscope image at 40x magnification. Bar = 10  $\mu$ m. d and e. Follicular fluid oestrogen and progesterone concentration from a range of follicle sizes collected 36 hours after the LH ovulation surge induction trigger injection during IVF treatment . The data were subjected to statistical verification using one way ANOVA with an uncorrected Fisher's LSD for follicular size. Values in graphs are means  $\pm$  SEM. Differences were considered significant if  $p < 0.05$  indicated by an asterisk and \*;  $p < 0.01$  \*\*. The number within the column represents the number of follicles analysed of that size.

**Fig. 3 Peak serum oestrogen (a) and progesterone (b) levels from IVF patients during gonadotrophin stimulated cycles.**

Serum levels were taken at the time of peak oestrogen (just before ovulation and the LH surge) during a stimulated IVF cycle. Ovarian reserve was measured indirectly by the antral follicle count (AFC). Antral

follicle count is the number of follicles between 2-10 mm on day 2-5 of a cycle in both ovaries. The ovarian reserve is then classified into a group level from A to E, good to poor, respectively. The number of patients is indicated, and ranged in age from 23 to 45 years old. The data were subjected to statistical verification using one way ANOVA with an uncorrected Fisher's LSD for follicular size. Values in graphs are means  $\pm$  SEM. Differences were considered significant if  $p < 0.05$  indicated by an asterisk\*;  $p < 0.01$ \*\*;  $p < 0.005$ \*\*\* and  $p < 0.001$ \*\*\*\*. The letter 'a' was significantly different to the letter a\*\*, with an attached \* indicating the level of significance. The number within the column represents the number of follicles analysed for that group.

**Fig. 4 Granulosa FSH receptor and LH receptor density and ovarian reserve depletion in young compared to older patients.**

**a.** Follicle stimulating hormone (FSH) receptor. **b.** Luteinising hormone (LH) receptor. Patients were grouped according to ovarian reserve measured indirectly by the antral follicle count (AFC). Antral follicle count is the number of follicles from 2-10 mm on day 2-5 of a cycle. Follicle count is based on the combined total from both ovaries. Data were subjected to statistical verification using one-way ANOVA with an uncorrected Fisher's LSD. Values are means  $\pm$  S.E.M., and differences were considered significant if \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.005$ . The number within the column represents the number of follicles analysed for that group. The MFI is the average mean fluorescent intensity emitted by the granulosa cells surface receptors. The scale is set the same for all experiments but is arbitrary.

**Fig. 5 Granulosa FSH receptor and LH receptor density and ovarian reserve depletion within the same age group compared to young cohort.**

**a.** Follicle stimulating hormone (FSH) receptor and **b.** luteinising hormone (LH) receptor. Patients were grouped according to ovarian reserve measured indirectly by the antral follicle count (AFC). Antral follicle count is the number of follicles from 2-10 mm on day 2-5 of a cycle. Follicle count is based on the combined total from both ovaries. Data were subjected to statistical verification using one-way ANOVA with an uncorrected Fisher's LSD. Values are means  $\pm$  S.E.M., and differences were considered significant if \* $p < 0.05$ , \*\* $p < 0.01$  and \*\*\* $p < 0.005$ . The number within the column represents the number

of follicles analysed for that group. The MFI is the average mean fluorescent intensity emitted by the granulosa cells surface receptors. The scale is set the same for all experiments but is arbitrary.

**Fig. 6 The comparative effect of rFSH dose on FSH receptor and LH receptor expression**

a. The dose of gonadotrophins rFSH received by young (A+ & A, 23-30y) and older patients groups (D & E, 35-45y) based on ovarian reserve measured by AFC. The number within the column represents the number of follicles analysed for that group. Data were subjected to statistical verification using student t test,  $p = < 0.0001$ . b. The effect of dose of rFSH on granulosa receptor density in patients matched for aged, ovarian reserve, AMH, and size of follicles, 40+ y, with an ovarian reserve of E, an AMH  $< 3.2$ , and follicle size of 10-22 mm. Data were subjected to statistical verification using one-way ANOVA with an uncorrected Fisher's LSD. Values are means  $\pm$  S.E.M., and differences were considered significant if  $p < 0.05$ . The number within the column represents the number of follicles analysed for that group.

**Fig. 7 Correlation of BMPRII signalling with FSH receptor and LH receptor as the ovarian reserve is depleted.**

a. FSH receptor and b. LH receptor correlated with BMPRII signalling. Sequential graphs show increasing age and declining ovarian reserve. Linear regression analysis, R square indicated for each group. Ovarian reserve measured indirectly by the antral follicle count. Antral follicle count is the number of follicles between 2-10 mm on day 2-5 of a cycle. The data points are averages of the receptor expression for that follicle size (ie each dot represents a follicle size from 4 mm 8 mm, 10 mm, 14 mm, 16 mm, 19 mm and 24 mm) and the total number of follicles per age group graph corresponds to: 23-30 y,  $n = 95$  follicles; 31-34 y,  $n = 43$  follicles; 35-39 y,  $n = 67$  follicles; 40+y,  $n = 77$  follicles. Follicle count is based on the combined total from both ovaries. Data were subjected to statistical verification using linear regression Values are means  $\pm$  S.E.M., and differences were considered significant if  $p < 0.05$ .

**Fig. 8 The effect of ovarian reserve measured by AFC on fertility**

The percentage of patients from different age groups based on the remaining primordial reserve of follicles within the ovaries indirectly measured by AFC and grouped from B to E. The younger age range of 35-38 years has a higher percentage of good ovarian reserve (group B) and a smaller percentage of

poor ovarian reserve, and consequently an increased fertility rate (Table 1). The graph shows that the 35-38y group of women have a greater % of women with the best B grade (45%) of ovarian reserve, In addition, this group has a much lower rate of women with the very poor ovarian reserve D (32.6%) and E 5.9%.