

Weak product spaces of Dirichlet series

Article

Accepted Version

Brevig, O. F. and Perfekt, K.-M. (2016) Weak product spaces of Dirichlet series. *Integral Equations and Operator Theory*, 86 (4). pp. 453-473. ISSN 1420-8989 doi: <https://doi.org/10.1007/s00020-016-2320-3> Available at <http://centaur.reading.ac.uk/71300/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1007/s00020-016-2320-3>

To link to this article DOI: <http://dx.doi.org/10.1007/s00020-016-2320-3>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Weak product spaces of Dirichlet series

Ole Fredrik Brevig and Karl-Mikael Perfekt

Abstract. Let \mathcal{H}^2 denote the space of ordinary Dirichlet series with square summable coefficients, and let \mathcal{H}_0^2 denote its subspace consisting of series vanishing at $+\infty$. We investigate the weak product spaces $\mathcal{H}^2 \odot \mathcal{H}^2$ and $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$, finding that several pertinent problems are more tractable for the latter space. This surprising phenomenon is related to the fact that $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ does not contain the infinite-dimensional subspace of \mathcal{H}^2 of series which lift to linear functions on the infinite polydisc.

The problems considered stem from questions about the dual spaces of these weak product spaces, and are therefore naturally phrased in terms of multiplicative Hankel forms. We show that there are bounded, even Schatten class, multiplicative Hankel forms on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$ whose analytic symbols are not in \mathcal{H}^2 . Based on this result we examine Nehari's theorem for such Hankel forms. We define also the skew product spaces associated with $\mathcal{H}^2 \odot \mathcal{H}^2$ and $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$, with respect to both half-plane and polydisc differentiation, the latter arising from Bohr's point of view. In the process we supply square function characterizations of the Hardy spaces \mathcal{H}^p , for $0 < p < \infty$, from the viewpoints of both types of differentiation. Finally we compare the skew product spaces to the weak product spaces, leading naturally to an interesting Schur multiplier problem.

Mathematics Subject Classification (2010). Primary 47B35. Secondary 30B50.

Keywords. Dirichlet series, Hankel form, square function, weak product space.

1. Introduction

In this paper, we investigate certain properties of weak product spaces associated with the Hardy space of Dirichlet series,

$$\mathcal{H}^2 = \left\{ f(s) = \sum_{n=1}^{\infty} a_n n^{-s} : \|f\|_{\mathcal{H}^2} = \left(\sum_{n=1}^{\infty} |a_n|^2 \right)^{\frac{1}{2}} < \infty \right\},$$

and its subspace \mathcal{H}_0^2 , consisting of those $f \in \mathcal{H}^2$ with $a_1 = f(+\infty) = 0$. The main objects of study are the weak product spaces $\mathcal{H}^2 \odot \mathcal{H}^2$ and $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$. With $\mathcal{X} = \mathcal{H}^2$ or $\mathcal{X} = \mathcal{H}_0^2$, the weak product $\mathcal{X} \odot \mathcal{X}$ is defined as the Banach space completion of the finite sums $F = \sum_k f_k g_k$, where $f_k, g_k \in \mathcal{X}$, under the norm

$$\|F\|_{\mathcal{X} \odot \mathcal{X}} = \inf \sum_k \|f_k\|_{\mathcal{H}^2} \|g_k\|_{\mathcal{H}^2}.$$

The infimum is taken over all finite representations of F as a sum of products.

While a separate study of $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ may at first be thought unmotivated, we will find that the norm of this space is significantly larger for certain types of Dirichlet series (see Theorem 1 and its corollaries). The presence of such examples is related to the obstructions faced in producing monomials n^{-s} in a product $f_k g_k$, for $f_k, g_k \in \mathcal{H}_0^2$, when n is an integer with a low number of prime factors. In particular, elements of $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ contain no terms of the form p^{-s} , where p is a prime number. Hence there is an easily identifiable infinite-dimensional subspace of $\mathcal{H}^2 \odot \mathcal{H}^2$ which has trivial intersection with $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$.

The weak product space $\mathcal{H}^2 \odot \mathcal{H}^2$ was first investigated by Helson [21, 22] in an attempt to decide whether Nehari's theorem holds for multiplicative Hankel forms (see also Section 2). Helson's work was continued in [25], where it was demonstrated that Nehari's theorem does not hold in full generality. To explain his point of view, note that each sequence $\varrho \in \ell^2$ induces a (not necessarily bounded) multiplicative Hankel form on $\ell^2 \times \ell^2$,

$$\varrho(a, b) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_m b_n \varrho_{mn}, \quad a, b \in \ell^2. \quad (1)$$

The analytic symbol of (1) is the Dirichlet series

$$\varphi(s) = \sum_{n=1}^{\infty} \overline{\varrho_n} n^{-s}.$$

Indeed, if f and g are elements of \mathcal{H}^2 with coefficients a and b , respectively, we have that

$$H_\varphi(fg) = \langle fg, \varphi \rangle = \varrho(a, b). \quad (2)$$

Here, and throughout the rest of the paper, $\langle \cdot, \cdot \rangle$ denotes the inner product of \mathcal{H}^2 .

Now, from (2) it is clear that the multiplicative Hankel form (1) is bounded on $\ell^2 \times \ell^2$, or equivalently H_φ on $\mathcal{H}^2 \times \mathcal{H}^2$, if and only if φ induces

a bounded linear functional on $\mathcal{H}^2 \odot \mathcal{H}^2$ through the \mathcal{H}^2 -pairing, i.e. if and only if φ is in $(\mathcal{H}^2 \odot \mathcal{H}^2)^*$.

The first bona fide example of a multiplicative Hankel form was obtained in [11], by the following approach. Note first that the elements of \mathcal{H}^2 are analytic functions in the half-plane $\operatorname{Re} s > 1/2$, the reproducing kernel at each such s being given by $\zeta(w + \bar{s})$, where $\zeta(s) = \sum_{n \geq 1} n^{-s}$ is the Riemann zeta function. It is thus natural to consider the Carleman-type operator

$$\mathbf{H}f(s) = \int_{1/2}^{\infty} f(w) (\zeta(s+w) - 1) dw$$

acting on \mathcal{H}_0^2 , since $(\zeta(s+w) - 1)$ is the reproducing kernel of \mathcal{H}_0^2 at w , for $w > 1/2$. The matrix of the operator \mathbf{H} is that of the multiplicative Hankel form whose analytic symbol φ is the primitive of $(\zeta(s+1/2) - 1)$ in \mathcal{H}_0^2 . In [11] it was shown that the operator norm of \mathbf{H} on \mathcal{H}_0^2 is π , which in terms of its corresponding Hankel form means precisely that $|\langle fg, \varphi \rangle| \leq \pi \|f\|_{\mathcal{H}^2} \|g\|_{\mathcal{H}^2}$ for $f, g \in \mathcal{H}_0^2$. More explicitly written,

$$\left| \sum_{m=2}^{\infty} \sum_{n=2}^{\infty} \frac{a_m b_n}{\sqrt{mn} \log(mn)} \right| \leq \pi \left(\sum_{m=2}^{\infty} |a_m|^2 \right)^{\frac{1}{2}} \left(\sum_{n=2}^{\infty} |b_n|^2 \right)^{\frac{1}{2}}. \quad (3)$$

As explained more thoroughly in [11], inequality (3) is a multiplicative analogue of the classical Hilbert inequality

$$\left| \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{a_m b_n}{m+n} \right| \leq \pi \left(\sum_{m=1}^{\infty} |a_m|^2 \right)^{\frac{1}{2}} \left(\sum_{n=1}^{\infty} |b_n|^2 \right)^{\frac{1}{2}}. \quad (4)$$

There are several other versions of (4) which are also usually referred to as Hilbert's inequality — we direct the interested reader to [19, Ch. IX]. Let us extract a few facts. First, that by discretization of the continuous version of (4) and the Hermite–Hadamard inequality, the following improvement of (4) can be obtained.

$$\left| \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{a_m b_n}{m+n+1} \right| \leq \pi \left(\sum_{m=0}^{\infty} |a_m|^2 \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} |b_n|^2 \right)^{\frac{1}{2}}.$$

We mention without proof that the same procedure (with some additional straightforward estimates) yields in the multiplicative setting that

$$\left| \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} \frac{a_m b_n}{\sqrt{(m+1/2)(n+1/2)} \log((m+1/2)(n+1/2))} \right| \leq \pi \|a\|_{\ell^2} \|b\|_{\ell^2},$$

which of course no longer represents a multiplicative Hankel form.

The strongest version of Hilbert's inequality (4) is

$$\left| \sum_{\substack{m, n \geq 0 \\ m+n > 0}} \frac{a_m b_n}{m+n} \right| \leq \pi \left(\sum_{m=0}^{\infty} |a_m|^2 \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} |b_n|^2 \right)^{\frac{1}{2}}. \quad (5)$$

This last variant can also be stated for two-tailed sequences $\{a_m\}_{m \in \mathbb{Z}}$ and $\{b_n\}_{n \in \mathbb{Z}}$. The proof of (5) amounts to a concrete application of Nehari's

theorem on $H^2(\mathbb{T})$, since the associated Hankel form has the bounded symbol Φ of supremum norm π ,

$$\Phi(z) = -i \operatorname{Arg}(z) = i(\pi - \theta), \quad z = e^{i\theta}. \quad (6)$$

As far as the authors are aware, all proofs of (5) in the literature make use of (a reformulation of) (6).

Whether the multiplicative Hankel form (3) has a bounded symbol is an open problem that is related to a long standing embedding problem of \mathcal{H}^1 (see [11, Sec. 6]). It therefore natural to ask if we also have

$$|\langle fg, \varphi \rangle| \leq \pi \|f\|_{\mathcal{H}^2} \|g\|_{\mathcal{H}^2}, \quad f, g \in \mathcal{H}^2? \quad (7)$$

In light of the discussion above, this question actually turns out to be more subtle than what one might expect at first. We are unable to settle it, seemingly due to the lack of a Nehari theorem for multiplicative Hankel forms.

That (7) is significantly easier to settle for \mathcal{H}_0^2 than for \mathcal{H}^2 is not a peculiarity, but rather an ongoing theme for all the questions we will ask about product spaces in this paper. Note that inequality (3) is easily recast as a question about (the dual space of) $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$. More generally, elements of $(\mathcal{H}_0^2 \odot \mathcal{H}_0^2)^*$ correspond to multiplicative Hankel forms of the type (1), but with sums starting at $m, n = 2$. The remainder of this section is an overview of the problems that we will consider.

In Section 2, we investigate the difference between Hankel forms on $\mathcal{H}^2 \times \mathcal{H}^2$ and Hankel forms on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$. After some preliminaries, we obtain the main result of this section, Theorem 1, which allows us to embed any bounded operator $C: \ell^2 \rightarrow \ell^2$ into a Hankel form on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$. This result is the basis for our observation that $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ is significantly smaller than $\mathcal{H}^2 \odot \mathcal{H}^2$, and it is also an important tool in the proofs of our other main results.

Helson [21] proved that any Hankel form on $\mathcal{H}^2 \times \mathcal{H}^2$ which is of Hilbert-Schmidt class S_2 is induced by a bounded symbol on the infinite polytorus \mathbb{T}^∞ . In [10] it was shown that if $p > p_0 \approx 5.74$, then there is a Hankel form on $\mathcal{H}^2 \times \mathcal{H}^2$ of Schatten class S_p that does not have any bounded symbol, leading to the conjecture that the same might be true for all $p > 2$. In Theorem 6 we will prove that $p = 2$ is indeed critical in this sense for multiplicative Hankel forms acting on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$, leading us closer to optimality of Helson's result. In fact, for $p > 2$ we will even demonstrate the existence of forms in S_p that do not have square-integrable symbols on the polytorus.

The penultimate section is devoted to the study of the skew product space $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2)$. The motivation to study this space is twofold. Firstly, characterizations of the dual spaces of skew products are often significantly easier to obtain (see [1, 2]). Secondly, for the classical Hardy space H^2 , the comparison between $H^2 \odot H^2$ and $\partial^{-1}(H^2 \odot \partial H^2)$ leads naturally to a Schur multiplier problem for Hankel matrices. Much has been written about this problem, owing to the fact that it was closely related to Pisier's

construction of a polynomially bounded operator not similar to a contraction. We refer the reader to [9, 13, 16, 26].

We begin Section 3 by proving a square function characterization of \mathcal{H}^p , which is of independent interest for the study of Hardy spaces of Dirichlet series. Due to the notation involved, we defer a precise statement to Theorem 8. We first use this characterization to prove that

$$\mathcal{H}^2 \odot \mathcal{H}^2 \subseteq \partial^{-1} (\partial \mathcal{H}^2 \odot \mathcal{H}^2) \subsetneq \mathcal{H}^1. \quad (8)$$

We then study whether the first inclusion in (8) is strict. This appears to be a difficult question, but by Schur multiplier methods we are able to demonstrate that this is the case if every appearance of \mathcal{H}^2 in (8) is replaced by \mathcal{H}_0^2 .

Finally, in Section 4, we look at the material of Section 3 again, but with the Hardy spaces of the polydisc in mind. Noting that Dirichlet series differentiation gives rise to a rather unnatural differentiation operator on the polydisc, we prove instead a square function characterization of $H^p(\mathbb{T}^\infty)$ that is adapted to the radial differentiation operator

$$R = \sum_{j=1}^{\infty} z_j \partial_{z_j}. \quad (9)$$

This will allow us to conclude that on finite-dimensional tori, it holds that

$$H^2(\mathbb{T}^d) \odot H^2(\mathbb{T}^d) = R^{-1} (H^2(\mathbb{T}^d) \odot RH^2(\mathbb{T}^d)) = H^1(\mathbb{T}^d).$$

It also turns out that radial differentiation has a number theoretic interpretation when considered from the Dirichlet series point of view, something that too will be elaborated upon in Section 4.

Notation

As usual, $\{p_j\}_{j \geq 1}$ denotes the sequence of prime numbers in increasing order, and $\Omega(n)$ will denote the number of prime factors in n , counting multiplicities. We will write $f \ll g$ to indicate that there is some positive constant C so that $|f(x)| \leq C|g(x)|$. If both $f \ll g$ and $g \ll f$, we write $f \asymp g$.

When we speak of a Dirichlet series φ as an element of a dual space \mathcal{H}^* , where \mathcal{H} is a Banach space of Dirichlet series in which the space of Dirichlet polynomials \mathcal{P} is dense, we always mean that the functional induced by φ via the \mathcal{H}^2 -pairing is bounded. That is, $\varphi \in \mathcal{H}^*$ if and only if the functional

$$v_\varphi(f) = \langle f, \varphi \rangle, \quad f \in \mathcal{P},$$

extends to a bounded functional on \mathcal{H} . Similarly, when we write that $\mathcal{H}^* \subseteq \mathcal{X}$, where \mathcal{X} is a Banach space of Dirichlet series, we mean that for every functional $v \in \mathcal{H}^*$ there exists a $\varphi \in \mathcal{X}$ such that $v = v_\varphi$ (on \mathcal{P}) and $\|\varphi\|_{\mathcal{X}} \ll \|v\|_{\mathcal{H}^*}$.

2. Hankel forms and a matrix embedding

Much of the success in the theory of Hardy spaces of Dirichlet series is due to a simple observation of Bohr [7], which facilitates a link between Dirichlet series

and function theory in polydiscs. By identifying each prime number with an independent variable, $z_j = p_j^{-s}$, the Dirichlet series $f(s) = \sum_{n \geq 1} a_n n^{-s}$ is lifted to a function in the Hardy space of the countably infinite torus, $H^2(\mathbb{T}^\infty)$. More precisely, the prime factorization

$$n = \prod_{j=1}^{\infty} p_j^{\kappa_j}$$

associates to n the finite non-negative multi-index $\kappa(n) = (\kappa_1, \kappa_2, \kappa_3, \dots)$. This means that the Bohr lift of f is

$$\mathcal{B}f(z) = \sum_{n=1}^{\infty} a_n z^{\kappa(n)},$$

where $z = (z_1, z_2, z_3, \dots)$. The mapping $\mathcal{B}: \mathcal{H}^2 \rightarrow H^2(\mathbb{T}^\infty)$ is an isometric isomorphism that respects multiplication. \mathbb{T}^∞ is a compact abelian group, and its Haar measure is denoted by m_∞ . The measure m_∞ is equal to the product of the normalized Lebesgue measure on \mathbb{T} in each variable. In particular, $H^2(\mathbb{T}^d)$ is a natural subspace of $H^2(\mathbb{T}^\infty)$. We refer to [20, 27] for further properties of $H^2(\mathbb{T}^\infty)$.

In [4], Bayart introduced the spaces \mathcal{H}^p , for $1 \leq p < \infty$, as those Dirichlet series f such that $\mathcal{B}f \in H^p(\mathbb{T}^\infty)$, and we define the \mathcal{H}^p -norm as

$$\|f\|_{\mathcal{H}^p} = \left(\int_{\mathbb{T}^\infty} |\mathcal{B}f(z)|^p dm_\infty(z) \right)^{\frac{1}{p}}.$$

As above, $H^p(\mathbb{T}^d)$ is a natural subspace of $H^p(\mathbb{T}^\infty) \simeq \mathcal{H}^p$.

Returning to the multiplicative Hankel form H_φ defined in (2), the fact that \mathcal{B} respects multiplication implies that

$$H_\varphi(fg) = \langle \mathcal{B}f\mathcal{B}g, \mathcal{B}\varphi \rangle_{H^2(\mathbb{T}^\infty)}.$$

From this representation, it is clear that we may replace $\mathcal{B}\varphi$ with any $\psi \in L^2(\mathbb{T}^\infty)$ such that $P\psi = \mathcal{B}\varphi$, where P denotes the Riesz projection from $L^2(\mathbb{T}^\infty)$ to $H^2(\mathbb{T}^\infty)$. In this case, we also denote the Hankel form H_φ by H_ψ . If $\psi \in L^\infty(\mathbb{T}^\infty)$, then $\|H_\varphi\| \leq \|\psi\|_\infty$, where $\|H_\varphi\|$ denotes the norm of H_φ acting on $\mathcal{H}^2 \times \mathcal{H}^2$, and we say that H_φ has bounded symbol ψ . Note that if the functional

$$f \mapsto \langle f, \varphi \rangle, \quad f \in \mathcal{H}^1,$$

is bounded on $\mathcal{H}^1 \simeq H^1(\mathbb{T}^\infty) \subset L^1(\mathbb{T}^\infty)$, then H_φ has a bounded symbol by the Hahn-Banach theorem. Hence, H_φ has a bounded symbol if and only if $\varphi \in (\mathcal{H}^1)^*$.

The main result of [25] implies that there exist bounded multiplicative Hankel forms that do not have a bounded symbol. It should be pointed out that the proof is non-constructive, and no example of a bounded multiplicative Hankel form without a bounded symbol has been identified. On the other hand, if $d = 1$ then Nehari's theorem [24] states that every bounded Hankel form H_φ on $H^2(\mathbb{T}^d) \odot H^2(\mathbb{T}^d)$ has a bounded symbol $\psi \in L^\infty(\mathbb{T}^d)$.

Nehari's theorem has been extended to $d < \infty$ by Ferguson–Lacey [17] and Lacey–Terwilleger [23].

The matrix of the multiplicative Hankel form (2) is

$$M_\varrho = (\varrho_{mn})_{m,n \geq 1} = \begin{pmatrix} \varrho_1 & \varrho_2 & \varrho_3 & \cdots \\ \varrho_2 & \varrho_4 & \varrho_6 & \cdots \\ \varrho_3 & \varrho_6 & \varrho_9 & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}. \quad (10)$$

By isolating the first row and column in M_ϱ using the inner product representation of H_φ from (2), we obtain

$$H_\varphi(fg) = a_1 b_1 \varrho_1 + a_1 \langle g - b_1, \varphi \rangle + b_1 \langle f - a_1, \varphi \rangle + H_\varphi((f - a_1)(g - b_1)). \quad (11)$$

The left hand side is a bounded Hankel form if and only if $\varphi \in (\mathcal{H}^2 \odot \mathcal{H}^2)^*$, while the right hand side is bounded if and only if $\varphi \in (\mathcal{H}_0^2)^* = \mathcal{H}^2/\mathbb{C}$ and $\varphi \in (\mathcal{H}_0^2 \odot \mathcal{H}_0^2)^*$. While it is obvious that

$$(\mathcal{H}^2 \odot \mathcal{H}^2)^* \subseteq \mathcal{H}^2, \quad (12)$$

we shall now see that the corresponding statement for \mathcal{H}_0^2 is not true. This will follow immediately from our next result, which also is crucial in establishing the other main results of the paper.

Theorem 1 (Matrix embedding). *Let $C = (c_{j,k})_{j,k \geq 1}$ be an infinite matrix defining an operator on ℓ^2 . Consider the Dirichlet series*

$$\varphi(s) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} c_{j,k} (p_{2j-1} p_{2k})^{-s},$$

where $\{p_j\}_{j \geq 1}$ denotes the sequence of primes numbers in increasing order. Then

- (a) $\|H_\varphi\|_0 = \|C\|$,
- (b) $\|H_\varphi\| \asymp \|C\|_{S_2} = \|\varphi\|_{\mathcal{H}^2}$,

where $\|H_\varphi\|_0$ denotes the norm of H_φ acting on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$, and $\|C\|_{S_2}$ denotes the Hilbert–Schmidt matrix norm of C ,

$$\|C\|_{S_2} = \left(\sum_{j=1}^{\infty} \sum_{k=1}^{\infty} |c_{j,k}|^2 \right)^{\frac{1}{2}}.$$

Proof. Let $f, g \in \mathcal{H}_0^2$ with coefficients $\{a_j\}_{j \geq 1}$ and $\{b_k\}_{k \geq 1}$, respectively. Since there are no constant terms in \mathcal{H}_0^2 we have that

$$H_\varphi(fg) = \langle fg, \varphi \rangle = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} (a_{p_{2j-1}} b_{p_{2k}} + a_{p_{2k}} b_{p_{2j-1}}) c_{j,k}. \quad (13)$$

Note that for every prime p , a_p and b_p each only appear once in this sum. Let

$$\mathcal{H}_1 = \text{span}\{p_{2k-1}^{-s} : k \geq 1\}, \quad \mathcal{H}_2 = \text{span}\{p_{2k}^{-s} : k \geq 1\}, \quad \mathcal{H}_3 = \mathcal{H}_0^2 \ominus (\mathcal{H}_1 \oplus \mathcal{H}_2),$$

and let $P_{\mathcal{X}_j}$ denote the corresponding orthogonal projections. Let \mathbf{a}_j and \mathbf{b}_j denote the coefficient sequences, in the natural basis of \mathcal{X}_j , of $P_{\mathcal{X}_j}f$ and $P_{\mathcal{X}_j}g$, respectively. Then we may rewrite (13) as

$$H_\varphi(fg) = \langle C\mathbf{b}_2, \mathbf{a}_1 \rangle_{\ell^2} + \langle C\mathbf{a}_2, \mathbf{b}_1 \rangle_{\ell^2} = \langle \mathcal{J}(C^T \oplus C)(\mathbf{a}_1, \mathbf{a}_2), (\mathbf{b}_1, \mathbf{b}_2) \rangle_{\ell^2 \oplus \ell^2},$$

where \mathcal{J} is the involution on $\ell^2 \oplus \ell^2$ defined by $\mathcal{J}(\mathbf{a}_1, \mathbf{a}_2) = (\mathbf{a}_2, \mathbf{a}_1)$. We conclude that

$$H_\varphi|_{\mathcal{H}_0^2} \simeq \mathcal{J}(C^T \oplus C) \oplus 0,$$

completing the proof of (a). For (b), we first observe that setting $g = 1$ implies $\|H_\varphi\| \geq \|\varphi\|_{(\mathcal{H}^2)^*} = \|\varphi\|_{\mathcal{H}^2} = \|C\|_{S_2}$. Returning to the decomposition (11) we see that $\|H_\varphi\| \leq 4\|C\|_{S_2}$, by using (a). \square

As a corollary of Theorem 1, we obtain that a bounded Hankel form on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$ does not necessarily have a symbol in $L^2(\mathbb{T}^\infty)$, in stark contrast with the classical situation where bounded Hankel forms have bounded symbols. We also find that (12) does not hold for \mathcal{H}_0^2 .

Corollary 2. $(\mathcal{H}_0^2 \odot \mathcal{H}_0^2)^* \not\subseteq \mathcal{H}^2$. That is, there are bounded multiplicative Hankel forms H_φ on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$ with the property that there is no $\psi \in L^2(\mathbb{T}^\infty)$ such that $H_\varphi = H_\psi$.

Proof. Use Theorem 1 and let C be the matrix of the identity operator on ℓ^2 . \square

Actually, we have the following stronger version of Corollary 2, which can be proven by considering all diagonal operators C on ℓ^2 and using Theorem 1. It exemplifies concretely that $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ is in some ways significantly smaller than $\mathcal{H}^2 \odot \mathcal{H}^2$.

Corollary 3. *The Dirichlet series*

$$f(s) = \sum_{k=1}^{\infty} a_k (p_{2k-1} p_{2k})^{-s}$$

is in $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ if and only if $a \in \ell^1$, while it is in $\mathcal{H}^2 \odot \mathcal{H}^2$ if and only if $a \in \ell^2$.

Recall that $H^2(\mathbb{T}^d)$ is a natural subspace of $H^2(\mathbb{T}^\infty)$ and that if $f \in \mathcal{H}_0^2$, then $\mathcal{B}f(0) = 0$. We now observe that the inclusions behave as expected for the corresponding finite-dimensional subspaces of the weak product spaces.

Lemma 4. For $1 \leq d < \infty$, let $H_0^2(\mathbb{T}^d)$ denote the space of functions $F \in H^2(\mathbb{T}^d)$ for which $F(0, 0, \dots, 0) = 0$. Then

$$(H_0^2(\mathbb{T}^d) \odot H_0^2(\mathbb{T}^d))^* \subseteq H_0^2(\mathbb{T}^d) \oplus \text{Lin}(\mathbb{T}^d) \subseteq H_0^2(\mathbb{T}^d),$$

where $\text{Lin}(\mathbb{T}^d)$ denotes the subspace of $H_0^2(\mathbb{T}^d)$ consisting of linear functions,

$$\text{Lin}(\mathbb{T}^d) = \left\{ L(z) = \sum_{j=1}^d a_j z_j : a_j \in \mathbb{C} \right\}.$$

Proof. It is sufficient to show that

$$H_0^2(\mathbb{T}^d) \ominus \text{Lin}(\mathbb{T}^d) \subseteq H_0^2(\mathbb{T}^d) \odot H_0^2(\mathbb{T}^d),$$

since it follows that any functional in $(H_0^2(\mathbb{T}^d) \odot H_0^2(\mathbb{T}^d))^*$ must be represented by a unique element of $H_0^2(\mathbb{T}^d) \ominus \text{Lin}(\mathbb{T}^d)$. Every $F \in H_0^2(\mathbb{T}^d) \ominus \text{Lin}(\mathbb{T}^d)$ can be written

$$F(z) = \sum_{j=1}^d z_j F_j(z),$$

where $F_j \in H_0^2(\mathbb{T}^d)$. This representation of F is not unique, but we can always organize it so that $\sum_j \|F_j\|_{H^2(\mathbb{T}^d)}^2 = \|F\|_{H^2(\mathbb{T}^d)}^2$. By the computation

$$\|F\|_{H_0^2 \odot H_0^2} \leq \sum_{j=1}^d 1 \cdot \|F_j\|_{H^2(\mathbb{T}^d)} \leq \sqrt{d} \left(\sum_{j=1}^d \|F_j\|_{H^2(\mathbb{T}^d)}^2 \right)^{\frac{1}{2}} = \sqrt{d} \|F\|_{H^2(\mathbb{T}^d)},$$

we see that $F \in H_0^2 \odot H_0^2$. \square

It is clear that the final part of this argument breaks down for $d = \infty$; the key point being that the subspace $\text{Lin}(\mathbb{T}^\infty)$ of linear functions in $H_0^2(\mathbb{T}^\infty) \simeq \mathcal{H}_0^2$ is infinite-dimensional, which from the Dirichlet series point of view corresponds to the fact that there are infinitely many prime numbers. Even so, Corollary 2 is surprising. We stress that its conclusion is related to the additional arithmetical obstructions which appear when computing the norm of an element in $\mathcal{H}_0^2 \odot \mathcal{H}_0^2$ rather than in $\mathcal{H}^2 \odot \mathcal{H}^2$. The following result is intended to clarify this statement. In particular, it demonstrates that the subspace of linear functions actually is complemented in $\mathcal{H}^2 \odot \mathcal{H}^2$.

Theorem 5. *For a non-negative integer m , let P_m denote the projection on m -homogeneous Dirichlet series,*

$$P_m \sum_{n=1}^{\infty} a_n n^{-s} = \sum_{\Omega(n)=m} a_n n^{-s}.$$

Then P_m is a contraction on $\mathcal{H}^2 \odot \mathcal{H}^2$.

Proof. The case $m = 0$ is trivial. Let $m \geq 1$ and suppose that

$$F = \sum_k f_k g_k \tag{14}$$

is a finite sum. Then

$$P_m F(s) = \sum_k \sum_{j=0}^m P_j f_k(s) P_{m-j} g_k(s).$$

By applying the definition of the norm of $\mathcal{H}^2 \odot \mathcal{H}^2$ and the Cauchy–Schwarz inequality, we find that

$$\begin{aligned} \|P_m F\|_{\mathcal{H}^2 \odot \mathcal{H}^2} &\leq \sum_k \sum_{j=0}^m \|P_j f_k\|_{\mathcal{H}^2} \|P_{m-j} g_k\|_{\mathcal{H}^2} \\ &\leq \sum_k \left(\sum_{j=0}^m \|P_j f_k\|_{\mathcal{H}^2}^2 \right)^{\frac{1}{2}} \left(\sum_{j=0}^m \|P_{m-j} g_k\|_{\mathcal{H}^2}^2 \right)^{\frac{1}{2}} \\ &\leq \sum_k \|f_k\|_{\mathcal{H}^2} \|g_k\|_{\mathcal{H}^2}, \end{aligned}$$

the final inequality following from the fact that

$$\sum_{j=0}^{\infty} \|P_j f\|_{\mathcal{H}^2}^2 = \|f\|_{\mathcal{H}^2}^2, \quad f \in \mathcal{H}^2.$$

The proof is completed by taking the infimum over the representations (14). \square

We return to the matrix of H_φ acting on $\mathcal{H}^2 \times \mathcal{H}^2$ from (10). The matrix M_ρ^0 corresponding to the action of H_φ on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$ is obtained from M_ρ by deleting the first row and column. That is, $M_\rho^0 = (\rho_{mn})_{m,n \geq 2}$ in view of (10).

Now, suppose that H_φ is a compact form, i.e. that its matrix M defines a compact operator on ℓ^2 . Let

$$\Lambda = \{\lambda_1, \lambda_2, \dots\}$$

denote the singular value sequence of M . We say that H_φ is in the Schatten class S_p , $0 < p \leq \infty$, if $\Lambda \in \ell^p$, and we let $\|H_\varphi\|_{S_p} = \|\Lambda\|_{\ell^p}$. When speaking of a Hankel form H_φ we will write $S_p(\mathcal{H}^2)$ or $S_p(\mathcal{H}_0^2)$ to clarify which space is being considered; using Theorem 1 as in Corollary 3, it is easy to construct Hankel forms belonging to the latter Schatten class, but not to the former.

Helson [21] showed that if $H_\varphi \in S_p(\mathcal{H}^2)$ and $p = 2$, then H_φ has a bounded symbol. In [10], the authors showed that this is no longer the case when

$$p > p_0 \approx 5.738817179.$$

We will now investigate symbols for forms $H_\varphi \in S_p(\mathcal{H}_0^2)$. We start by verifying that Helson's result still holds for $S_2(\mathcal{H}_0^2)$.

As in Lemma 4, any bounded Hankel form on $\mathcal{H}_0^2 \times \mathcal{H}_0^2$ has a symbol φ in $(\mathcal{H}_0^2 \odot \mathcal{H}_0^2)^*$ of the form

$$\varphi(s) = \sum_{\Omega(n) \geq 2} \varrho_n n^{-s}.$$

From this fact, a computation shows that

$$\begin{aligned} \|H_\varphi\|_{S_2(\mathcal{H}_0^2)}^2 &= \sum_{m=2}^{\infty} \sum_{n=2}^{\infty} |H_\varphi(m^{-s}n^{-s})|^2 \\ &= \sum_{\Omega(n) \geq 2} (d(n) - 2) |\varrho_n|^2 \asymp \sum_{\Omega(n) \geq 2} d(n) |\varrho_n|^2. \end{aligned}$$

Here $d(n)$ denotes the number of divisors of n , and the final estimate follows from the fact that $d(n) - 2 \geq d(n)/3$ for n such that $\Omega(n) \geq 2$, seeing as $d(n) \geq \Omega(n) + 1$. Hence we can use Helson's inequality

$$\left(\sum_{n=1}^{\infty} \frac{|a_n|^2}{d(n)} \right)^{\frac{1}{2}} \leq \left\| \sum_{n=1}^{\infty} a_n n^{-s} \right\|_{\mathcal{H}^1}$$

to conclude that $\varphi \in (\mathcal{H}^1)^*$ whenever $H_\varphi \in S_2(\mathcal{H}_0^2)$. That is, H_φ has a bounded symbol whenever $H_\varphi \in S_2(\mathcal{H}_0^2)$. We now show that Helson's result is optimal for $S_p(\mathcal{H}_0^2)$.

Theorem 6. *For $p > 2$ there exist Hankel forms $H_\varphi \in S_p(\mathcal{H}_0^2)$ such that no ψ in $L^2(\mathbb{T}^\infty)$ satisfies $H_\varphi = H_\psi$. In particular, there exist Hankel forms $H_\varphi \in S_p(\mathcal{H}_0^2)$ for which there are no bounded symbols.*

Proof. Let $C = (c_{j,k})_{j,k \geq 1}$ be a matrix defining an operator on ℓ^2 which belongs to S_p but not to S_2 . In accordance with Theorem 1 let

$$\varphi(s) = \sum_{j=1}^{\infty} \sum_{k=1}^{\infty} c_{j,k} (p_{2j-1} p_{2k})^{-s}.$$

Since, as in the proof of Theorem 1, $H_\varphi|_{\mathcal{H}_0^2} \simeq \mathcal{J}(C^T \oplus C) \oplus 0$, we have that

$$\|H_\varphi\|_{S_p(\mathcal{H}_0^2)}^p = 2 \|C\|_{S_p}^p < \infty.$$

On the other hand, we have by assumption that

$$\|\varphi\|_{\mathcal{H}^2} = \|C\|_{S_2} = \infty. \quad \square$$

While Theorem 6 does not concern Hankel forms on $\mathcal{H}^2 \times \mathcal{H}^2$, we do consider it to give us an indication that $p = 2$ might be the critical value also in this case.

Conjecture 1. For every $p > 2$ there exists a multiplicative Hankel form H_φ in $S_p(\mathcal{H}^2)$ without a bounded symbol.

3. A square function characterization of \mathcal{H}^p and skew products

In the context of the classical Hardy spaces, it was Bourgain [9] who recalled the square function characterization of H^p due to Fefferman and Stein [15] and used it to the effect of showing that $H^2 \odot \partial H^2 \subseteq \partial H^1$, where ∂H^p

denotes the space consisting of the derivatives of all H^p -functions. In view of the fact that $\partial H^1 = \partial(H^2 \odot H^2) \subseteq H^2 \odot \partial H^2$ this immediately implies that

$$\partial^{-1}(H^2 \odot \partial H^2) = H^2 \odot H^2. \quad (15)$$

In terms of bilinear forms, we can naturally associate a Hankel-type form J_g to every element $g \in (\partial^{-1}(H^2 \odot \partial H^2))^*$. If an additive Hankel form H_g on $H^2 \times H^2$ corresponds to the matrix $(\hat{g}(j+k))_{j,k \geq 0}$, then J_g has matrix

$$\left(\frac{j+1}{j+k+1} \hat{g}(j+k) \right)_{j,k \geq 0}.$$

Hence Bourgain's lemma (15) can be equivalently rephrased to say that the map $H_g \mapsto J_g$ is bounded in operator norm. This statement actually carries greater interest than what its face value might suggest. The matrix

$$\left(\frac{j+1}{j+k+1} \right)_{j,k \geq 0}$$

is not a bounded Schur multiplier on all matrices, and hence the map $H_g \mapsto J_g$ is not completely bounded [13]. This observation is at the heart of Pisier's [26] construction of a polynomially bounded operator not similar to a contraction.

We define the skew product space $\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2)$ as the Banach space completion of the space of Dirichlet series F whose derivatives have a finite sum representation $F' = \sum_k f_k g_k'$, where $f_k, g_k \in \mathcal{H}^2$. The completion is taken under the norm

$$\|F\|_{\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2)} = |F(+\infty)| + \inf \sum_k \|f_k\|_{\mathcal{H}^2} \|g_k\|_{\mathcal{H}^2},$$

where the infimum is computed over all finite representations. From the product rule $(fg)' = f'g + fg'$ it is clear that

$$\mathcal{H}^2 \odot \mathcal{H}^2 \subseteq \partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2). \quad (16)$$

Our first goal is to establish a square function characterization of \mathcal{H}^p , for $0 < p < \infty$, and use it to show that $\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2) \subseteq \mathcal{H}^1$. We begin by recalling that the spaces \mathcal{H}^p are related to the Möbius invariant Hardy spaces in the right half-plane, \mathbb{C}_0 , defined as

$$H_1^p(\mathbb{C}_0) = \left\{ f \in \text{Hol}(\mathbb{C}_0) : \|f\|_{H_1^p(\mathbb{C}_0)} = \sup_{\sigma > 0} \left(\frac{1}{\pi} \int_{\mathbb{R}} |f(\sigma + it)|^p \frac{dt}{1+t^2} \right)^{\frac{1}{p}} < \infty \right\}.$$

Given a character $\chi \in \mathbb{T}^\infty$, we "twist" the Dirichlet series $f(s) = \sum_{n \geq 1} a_n n^{-s}$ to obtain

$$f_\chi(s) = \sum_{n=1}^{\infty} a_n \chi(n) n^{-s}, \quad \chi(n) = \chi^{\kappa(n)}.$$

We will require the following basic result, which can be extracted from Lemma 5 and Theorem 5 in [4].

Lemma 7. *Let $0 < p < \infty$, and suppose that $f \in \mathcal{H}^p$. For almost every $\chi \in \mathbb{T}^\infty$, $f_\chi \in H_1^p(\mathbb{C}_0)$. Moreover,*

$$\|f\|_{\mathcal{H}^p} = \left(\int_{\mathbb{T}^\infty} \|f_\chi\|_{H_1^p(\mathbb{C}_0)}^p dm_\infty(\chi) \right)^{\frac{1}{p}}.$$

Remark. The results in [4] are stated only for $p \geq 1$, but the same arguments lead to our statement of Lemma 7.

For $\tau \in \mathbb{R}$, let Γ_τ be the cone

$$\Gamma_\tau = \{\sigma + it : |t - \tau| < \sigma\}$$

in the right half-plane \mathbb{C}_0 , with vertex at $i\tau$. For a holomorphic function f in \mathbb{C}_0 , let Sf be the square function, or the Lusin area integral,

$$Sf(\tau) = \left(\int_{\Gamma_\tau} |f'(\sigma + it)|^2 d\sigma dt \right)^{1/2}, \quad \tau \in \mathbb{R},$$

and let f^* denote the non-tangential maximal function

$$f^*(\tau) = \sup_{s \in \Gamma_\tau} |f(s)|, \quad \tau \in \mathbb{R}.$$

Since $1/(1 + \tau^2)$ is a Muckenhoupt A_q -weight for all $q > 1$, it follows from Gundy and Wheeden [18] that $f \in H_1^p(\mathbb{C}_0)$ if and only if $f^* \in L_1^p(\mathbb{R}) = L^p((1 + \tau^2)^{-1} d\tau)$, for $0 < p < \infty$, with comparable norms. Furthermore, if $\lim_{\sigma \rightarrow \infty} f(\sigma + it) = 0$, then

$$\|f^*\|_{L_1^p(\mathbb{R})} \asymp \|Sf\|_{L_1^p(\mathbb{R})}. \quad (17)$$

This gives us a norm expression for functions in \mathcal{H}^p in terms of the square function.

Theorem 8. *Let $f(s) = \sum_{n \geq 1} a_n n^{-s}$. Then for any $0 < p < \infty$, we have*

$$\begin{aligned} \|f\|_{\mathcal{H}^p}^p &\asymp |a_1|^p + \int_{\mathbb{T}^\infty} \|S(f_\chi)\|_{L_1^p(\mathbb{R})}^p dm_\infty(\chi) \\ &= |a_1|^p + \int_{\mathbb{T}^\infty} \int_{\mathbb{R}} \left(\int_{\Gamma_\tau} |f'_\chi(\sigma + it)|^2 d\sigma dt \right)^{p/2} \frac{d\tau}{1 + \tau^2} dm_\infty(\chi). \end{aligned} \quad (18)$$

Proof. In view of (17) and Lemma 7 we obtain (18) for f with constant term $a_1 = 0$, that is, for $f \in \mathcal{H}_0^p$. Note that the linear functional $f \mapsto a_1$ is bounded on \mathcal{H}^p , corresponding to the functional $\mathcal{B}f \mapsto \mathcal{B}f(0)$ on $H^p(\mathbb{T}^\infty)$ [12]. Hence, the closed subspace \mathcal{H}_0^p is complemented in \mathcal{H}^p by \mathbb{C} , and (18) follows in general for $f \in \mathcal{H}^p$, with one side being finite if and only if the other is. \square

Corollary 9. $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2) \subseteq \mathcal{H}^1$.

Proof. Suppose that $f, g \in \mathcal{H}^2$, and that F is the Dirichlet series such that $F' = fg'$ with $F(+\infty) = 0$. Since $\|g - g(+\infty)\|_{\mathcal{H}^2} \leq \|g\|_{\mathcal{H}^2}$ it is for the

purpose of proving the statement justified to assume that $g(+\infty) = 0$. We then have that

$$\begin{aligned}
\|F\|_{\mathcal{H}^1} &\asymp \int_{\mathbb{T}^\infty} \int_{\mathbb{R}} \left(\int_{\Gamma_\tau} |f_\chi(\sigma + it)|^2 |g'_\chi(\sigma + it)|^2 d\sigma dt \right)^{1/2} \frac{d\tau}{1 + \tau^2} dm_\infty(\chi) \\
&\leq \int_{\mathbb{T}^\infty} \int_{\mathbb{R}} (f_\chi)^*(\tau) \left(\int_{\Gamma_\tau} |g'_\chi(\sigma + it)|^2 d\sigma dt \right)^{1/2} \frac{d\tau}{1 + \tau^2} dm_\infty(\chi) \\
&\leq \int_{\mathbb{T}^\infty} \|(f_\chi)^*\|_{L^2_1(\mathbb{R})} \left(\int_{\mathbb{R}} \int_{\Gamma_\tau} |g'_\chi(\sigma + it)|^2 d\sigma dt \frac{d\tau}{1 + \tau^2} \right)^{1/2} dm_\infty(\chi) \\
&\asymp \int_{\mathbb{T}^\infty} \|f_\chi\|_{H^2_1(\mathbb{C}_0)} \|g_\chi\|_{H^2_1(\mathbb{C}_0)} dm_\infty(\chi) \leq \|f\|_{\mathcal{H}^2} \|g\|_{\mathcal{H}^2}.
\end{aligned}$$

This proves that $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2) \subseteq \mathcal{H}^1$. \square

Before proceeding, we give a few remarks on the application of Theorem 8 to the Hardy space $H^p(\mathbb{T}^d)$ of a finite-dimensional polydisc, $d < \infty$. Let D denote the differentiation operator on Dirichlet series,

$$Df(s) = \partial f(s) = f'(s) = - \sum_{n=2}^{\infty} a_n \log(n) n^{-s}.$$

Consider a series f such that $\mathcal{B}f \in H^2(\mathbb{T}^d)$, i.e. such that $a_n = 0$ if $p_j | n$ for some $j > d$. Identifying p_j with the j th complex variable z_j , the differentiation operator D in the usual polydisc notation has the form

$$D\mathcal{B}f(z_1, \dots, z_d) = - \sum_{j=1}^d \log(p_j) z_j \partial_{z_j} \mathcal{B}f(z_1, \dots, z_d). \quad (19)$$

Hence Theorem 8 gives us a new type of square function characterization of $H^p(\mathbb{T}^d)$, in terms of the differentiation operator D . In analogy with Corollary 9 it can be used to prove that

$$D^{-1}(H^2(\mathbb{T}^d) \odot DH^2(\mathbb{T}^d)) \subseteq H^1(\mathbb{T}^d)$$

and by the characterization of $H^1(\mathbb{T}^d)$ due to Ferguson–Lacey [17] and Lacey–Terwilleger [23] we conclude that in the finite polydisc we have

$$D^{-1}(H^2(\mathbb{T}^d) \odot DH^2(\mathbb{T}^d)) = H^2(\mathbb{T}^d) \odot H^2(\mathbb{T}^d) = H^1(\mathbb{T}^d). \quad (20)$$

It should be objected, however, that the weighted differentiation operator D might not be natural in the setting of the polydisc. In Section 4 we shall consider the constructs of the present section for the infinite polydisc, using the radial differentiation operator instead of D , and in the process prove that (20) is valid also for radial differentiation and integration.

We return to the discussion of products of Dirichlet series spaces, and note that Corollary 9 in combination with (16) yields that

$$\mathcal{H}^2 \odot \mathcal{H}^2 \subseteq \partial^{-1}(\partial\mathcal{H}^2 \odot \mathcal{H}^2) \subseteq \mathcal{H}^1. \quad (21)$$

The remainder of this section is devoted to the investigation of whether these inclusions are strict. We begin with the following observation.

Lemma 10. Let $\varphi(s) = \sum_{k \geq 1} \bar{\rho}_k k^{-s}$ be a function in \mathcal{H}^2 . Then φ induces a bounded linear functional v_φ on $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2)$, via the \mathcal{H}^2 -pairing, if and only if the form

$$J_\varphi(a, b) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_m b_n \frac{\log n}{\log m + \log n} \rho_{mn} \quad (22)$$

is bounded on $\ell^2 \times \ell^2$, where the summand is understood to be 0 if $m = n = 1$. The corresponding norms are equivalent,

$$\|v_\varphi\| \asymp |\rho_1| + \|J_\varphi\|.$$

In particular, if $\rho_k \geq 0$ for all k , then $\varphi \in (\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2))^*$ if and only if $\varphi \in (\mathcal{H}^2 \odot \mathcal{H}^2)^*$, with equivalent norms.

Proof. Suppose that f and g are Dirichlet series with coefficient sequences a and b , respectively. Let $\partial^{-1}(f'g)$ denote the primitive of $f'g$ with constant term 0. Then

$$\langle \partial^{-1}(f'g), \varphi \rangle = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_m b_n \frac{\log n}{\log m + \log n} \rho_{mn},$$

proving the first part of the proposition. For the second part, note as per usual that the action of φ as an element in $(\mathcal{H}^2 \odot \mathcal{H}^2)^*$ corresponds to the multiplicative Hankel form

$$H_\varphi(a, b) = \sum_{m=1}^{\infty} \sum_{n=1}^{\infty} a_m b_n \rho_{mn}. \quad (23)$$

Hence, if $\rho_k \geq 0$ for all k , then

$$\|\varphi\|_{(\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2))^*} \ll \|\varphi\|_{(\mathcal{H}^2 \odot \mathcal{H}^2)^*}.$$

The converse inequality is a direct consequence of (16). \square

Ortega-Cerdá and Seip [25] showed that $\mathcal{H}^2 \odot \mathcal{H}^2 \subsetneq \mathcal{H}^1$. With Lemma 10, we are able to apply their technique to prove the corresponding statement for $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2)$.

Theorem 11. $\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2) \subsetneq \mathcal{H}^1$

Proof. Let d be a positive integer and consider the function

$$\varphi_d(s) = \prod_{j=1}^d (p_{2j-1}^{-s} + p_{2j}^{-s}),$$

where $\{p_j\}_{j \geq 1}$ again denotes the prime sequence. The norm of φ_d as an element of the dual of $\mathcal{H}^2 \odot \mathcal{H}^2$ is $2^{d/2}$ [25]. Since the coefficients of φ_d are non-negative, Lemma 10 hence shows that

$$\|\varphi_d\|_{(\partial^{-1}(\mathcal{H}^2 \odot \partial\mathcal{H}^2))^*} \asymp 2^{d/2}.$$

On the other hand, consider $f_d = \varphi_d$ as an element of \mathcal{H}^1 , $\|f_d\|_{\mathcal{H}^1} = (4/\pi)^d$ [25]. Since $\langle f_d, \varphi_d \rangle_2 = 2^d$, the functional induced by φ_d on \mathcal{H}^1 has norm at

least $(\pi/2)^d$. If it were the case that $\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2) = \mathcal{H}^1$, then the norm of φ_d as a functional on \mathcal{H}^1 and the norm as a functional on $\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2)$ would be equivalent, a contradiction as $d \rightarrow \infty$. \square

The remaining question of whether

$$(\partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2))^* = (\mathcal{H}^2 \odot \mathcal{H}^2)^* \quad (24)$$

or, equivalently, whether the first inclusion in (21) is strict, appears to be subtle. As we just saw in Lemma 10 it can be rephrased as to ask if the forms (22) and (23) are simultaneously bounded, which would mean precisely that

$$\left(\frac{\log n}{\log m + \log n} \right)_{m,n \geq 1}$$

is a Schur multiplier on the class of multiplicative Hankel forms. Specializing to the one-variable case by only considering integers of the form 2^k , we see that the analogue of (24) for the classical Hardy space $H^2(\mathbb{T})$ is equivalent to the statement that $(j+1)/(j+k+1)$ is a Schur multiplier on (additive) Hankel forms, as discussed in the introduction of this section.

However, by applying Theorem 1 in full force together with Schur multiplier techniques, we are able to show that the inclusion is strict when \mathcal{H}^2 is replaced by \mathcal{H}_0^2 . We define $\partial^{-1}(\mathcal{H}_0^2 \odot \partial \mathcal{H}_0^2)$ in exact analogy with our previous considerations, except that we impose all of its elements f to have constant term $f(+\infty) = 0$.

Theorem 12. $\mathcal{H}_0^2 \odot \mathcal{H}_0^2 \subsetneq \partial^{-1}(\mathcal{H}_0^2 \odot \partial \mathcal{H}_0^2)$.

Proof. Assume to the contrary that

$$\left(\frac{\log n}{\log m + \log n} \right)_{m,n \geq 2}$$

is a Schur multiplier on bounded multiplicative Hankel forms

$$\rho(a, b) = \sum_{m=2}^{\infty} \sum_{n=2}^{\infty} a_m b_n \rho_{mn}, \quad a, b \in \ell^2.$$

Applied to every symbol constructed by the procedure of Theorem 1, we conclude that

$$\left(\frac{\log p_{2j-1}}{\log p_{2k} + \log p_{2j-1}} \right)_{j,k \geq 1} \quad (25)$$

is a Schur multiplier on all matrices C defining bounded operators $C : \ell^2 \rightarrow \ell^2$. However, (25) cannot be a Schur multiplier, as this would defy Bennett's criterion [5], since

$$\lim_{j \rightarrow \infty} \lim_{k \rightarrow \infty} \frac{\log p_{2j-1}}{\log p_{2k} + \log p_{2j-1}} = 0,$$

while

$$\lim_{k \rightarrow \infty} \lim_{j \rightarrow \infty} \frac{\log p_{2j-1}}{\log p_{2k} + \log p_{2j-1}} = 1. \quad \square$$

It must be stressed that Theorem 12 *does not* imply that the inclusion in (16) is strict. If we attempt to apply the proof to $\mathcal{H}^2 \odot \mathcal{H}^2$, the matrices constructed by Theorem 1 are Hilbert–Schmidt. To be a Schur multiplier on Hilbert–Schmidt matrices means only to have bounded entries, so no contradiction is obtained. However, we do feel that Theorem 12 invokes the natural conjecture.

Conjecture 2. The inclusion between the standard weak product and its skew counterpart is strict, $\mathcal{H}^2 \odot \mathcal{H}^2 \subsetneq \partial^{-1}(\mathcal{H}^2 \odot \partial \mathcal{H}^2)$.

4. Radial differentiation

From the polydisc point of view, the constructs of the last section all arose from the weighted differentiation operator D of (19), obtained from the Dirichlet series formalism. In the present section we shall consider instead the more natural radial differentiation operator of equation (9). Before commencing, note that as in Theorem 5 every Dirichlet series may be decomposed into m -homogeneous subseries,

$$f(s) = \sum_{n=0}^{\infty} a_n n^{-s} = \sum_{m=0}^{\infty} \left(\sum_{\Omega(n)=m} a_n n^{-s} \right) = \sum_{m=0}^{\infty} P_m f(s).$$

Through the Bohr lift, this is equivalent to the corresponding decomposition of a power series in a countably infinite number of variables,

$$F(z) = \sum_{n=0}^{\infty} a_n z^{\kappa(n)} = \sum_{m=0}^{\infty} \left(\sum_{|\kappa(n)|=m} a_n z^{\kappa(n)} \right) = \sum_{m=0}^{\infty} P_m F(z), \quad z = (z_1, z_2, \dots).$$

We recall that $\kappa(n) = (\kappa_1, \kappa_2, \dots)$ is the finitely supported multi-index associated to every positive integer n through its prime decomposition, so that

$$|\kappa(n)| = \Omega(n) = \sum_j \kappa_j.$$

Consider now, for any $z \in \mathbb{T}^{\infty}$, the following power series in one variable w .

$$F_z(w) = F(zw) = \sum_{n=1}^{\infty} a_n z^{\kappa(n)} w^{\Omega(n)} = \sum_{m=0}^{\infty} P_m F(z) w^m.$$

Observe in particular that the m th coefficient of $F_z(w)$ is the m -homogeneous subseries of F . From here it is clear that differentiation in the auxiliary variable w allows us to capture the natural radial differentiation of the polydisc, since every monomial of order m is treated equally. This is further justified by the formal computation

$$w \frac{d}{dw} F_z(w) = w \sum_{j=1}^{\infty} z_j \partial_{z_j} F(zw) = (RF)_z(w).$$

We have the following analogue of Lemma 7. We also point out that through the Bohr lift a similar statement can be made for Dirichlet series.

Lemma 13. *Let $F \in H^p(\mathbb{T}^\infty)$, $0 < p < \infty$. Then $F_z \in H^p(\mathbb{T})$ for almost every $z \in \mathbb{T}^\infty$ and*

$$\|F\|_{H^p(\mathbb{T}^\infty)} = \left(\int_{\mathbb{T}^\infty} \|F_z\|_{H^p(\mathbb{T})}^p dm_\infty(z) \right)^{1/p}. \quad (26)$$

Proof. This follows from Fubini's theorem and the fact that $z \mapsto F(z)$ and $z \mapsto F(wz)$, for $w \in \mathbb{T}$, have equal $H^p(\mathbb{T}^\infty)$ -norm. \square

For $\theta \in [0, 2\pi)$, let $\Gamma_\alpha(\theta)$ denote the Stolz angle in \mathbb{D} with vertex at $e^{i\theta}$ and of some fixed aperture $\alpha < \pi/2$. The (slightly non-standard) square function Sg of a function g holomorphic in \mathbb{D} is given by

$$Sg(\theta) = \left(\int_{\Gamma_\alpha(\theta)} |wg'(w)|^2 dA(w) \right)^{1/2},$$

where dA denotes the normalized area element. If $g(0) = 0$ we have that $\|g\|_{H^p(\mathbb{T})} \asymp \|Sg\|_{L^p(\mathbb{T})}$. Since $F_z(0) = F(0)$ for every $z \in \mathbb{T}^\infty$, this immediately gives us the analogue of Theorem 8.

Theorem 14. *Let $F \in H^p(\mathbb{T}^\infty)$, $0 < p < \infty$. Then*

$$\|F\|_{H^p(\mathbb{T}^\infty)}^p \asymp |F(0)|^p + \int_{\mathbb{T}^\infty} \int_0^{2\pi} \left(\int_{\Gamma_\alpha(\theta)} |(RF)_z(w)|^2 dA(w) \right)^{p/2} \frac{d\theta}{2\pi} dm_\infty(z).$$

Now most of the arguments of the previous section can be repeated. We collect the results that follow without providing details. Note in particular the satisfying conclusion obtained for the finite-dimensional polydisc. Indeed, this result partly motivates the existence of this section.

Corollary 15. *We have that*

$$H^2(\mathbb{T}^\infty) \odot H^2(\mathbb{T}^\infty) \subseteq R^{-1} (H^2(\mathbb{T}^\infty) \odot RH^2(\mathbb{T}^\infty)) \subsetneq H^1(\mathbb{T}^\infty).$$

On the other hand, when $d < \infty$ it holds that

$$H^2(\mathbb{T}^d) \odot H^2(\mathbb{T}^d) = R^{-1} (H^2(\mathbb{T}^d) \odot RH^2(\mathbb{T}^d)) = H^1(\mathbb{T}^d).$$

We remark that it is not clear how to obtain Corollary 15 directly from the considerations in Section 3, due to the weights $\log p_j$ entering into Dirichlet series differentiation. In fact, suppose that $n = \prod_j p_j^{\kappa_j}$. Then

$$\log n = \sum_j \kappa_j \log p_j \quad \text{and} \quad \Omega(n) = \sum_j \kappa_j,$$

illustrating the fact that the R treats every prime equally, while the half-plane differentiation operator D does not. In particular, the proof of Theorem 12 does not yield any information when D is replaced by R , since the Schur multiplier vital to the proof has entries

$$\frac{\Omega(n)}{\Omega(m) + \Omega(n)} = \frac{\Omega(p_{2j-1})}{\Omega(p_{2k}) + \Omega(p_{2j-1})} = \frac{1}{2}.$$

It should also be pointed out that decomposing Dirichlet series (or power series on the infinite polydisc) into homogeneous subseries is not a new idea. It dates back at least to Bohnenblust–Hille [6], and has recently been applied to obtain results for composition operators on spaces of Dirichlet series [3] as well as L^1 -estimates for Dirichlet polynomials [8].

We conclude this paper by providing a charming inequality, which follows at once from Lemma 13 and the classical Hardy inequality

$$\sum_{m=0}^{\infty} \frac{|b_m|}{m+1} \leq \pi \left\| \sum_{m=0}^{\infty} b_m w^m \right\|_{H^1(\mathbb{T})}. \quad (27)$$

Corollary 16. *Let $f(s) = \sum_{n \geq 1} a_n n^{-s} \in \mathcal{H}^1$ and consider the m -homogeneous subseries $P_m f(s) = \sum_{\Omega(n)=m} a_n n^{-s}$. Then*

$$\sum_{m=0}^{\infty} \frac{\|P_m f\|_{\mathcal{H}^1}}{m+1} \leq \pi \|f\|_{\mathcal{H}^1}.$$

Corollary 16 can be compared to the estimate $\|P_m f\|_{\mathcal{H}^1} \leq \|f\|_{\mathcal{H}^1}$ appearing in [8, Lem. 3]. Returning to the beginnings of this paper, we mention that Hardy’s inequality (27) in turn can be obtained by viewing the bounded symbol for the sharpest version of Hilbert’s inequality (6) as an element in the dual of $H^1(\mathbb{T})$ (see [14, pp. 47–49]).

References

1. A. Aleman and K.-M. Perfekt, *Hankel forms and embedding theorems in weighted Dirichlet spaces*, Int. Math. Res. Not. IMRN (2012), no. 19, 4435–4448.
2. N. Arcozzi, R. Rochberg, E. T. Sawyer, and B. D. Wick, *The Dirichlet space: a survey*, New York J. Math. **17A** (2011), 45–86.
3. M. Bailleul and O. F. Brevig, *Composition operators on Bohr–Bergman spaces of Dirichlet series*, Ann. Acad. Sci. Fenn. Math. **41** (2016), no. 1, 129–142.
4. F. Bayart, *Hardy spaces of Dirichlet series and their composition operators*, Monatsh. Math. **136** (2002), no. 3, 203–236.
5. G. Bennett, *Schur multipliers*, Duke Math. J. **44** (1977), no. 3, 603–639.
6. H. F. Bohnenblust and E. Hille, *On the absolute convergence of Dirichlet series*, Ann. of Math. **32** (1931), no. 3, 600–622.
7. H. Bohr, *Ueber die Bedeutung der Potenzreihen unendlich vieler Variablen in der Theorie der Dirichletschen Reihen $\sum a_n/n^s$* , Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch–Physikalische Klasse **1913** (1913), 441–488.
8. A. Bondarenko and K. Seip, *Helson’s problem for sums of a random multiplicative function*, Mathematika **62** (2016), no. 1, 101–110.
9. J. Bourgain, *On the similarity problem for polynomially bounded operators on Hilbert space*, Israel J. Math. **54** (1986), no. 2, 227–241.
10. O. F. Brevig and K.-M. Perfekt, *Failure of Nehari’s theorem for multiplicative Hankel forms in Schatten classes*, Studia Math. **228** (2014), no. 2, 101–108.

11. O. F. Brevig, K.-M. Perfekt, K. Seip, A. G. Siskakis, and D. Vukotić, *The multiplicative Hilbert matrix*, Adv. Math. **302** (2016), 410–432.
12. B. J. Cole and T. W. Gamelin, *Representing measures and Hardy spaces for the infinite polydisk algebra*, Proc. London Math. Soc. (3) **53** (1986), no. 1, 112–142.
13. K. R. Davidson and V. I. Paulsen, *Polynomially bounded operators*, J. Reine Angew. Math. **487** (1997), 153–170.
14. P. L. Duren, *Theory of H^p spaces*, Pure and Applied Mathematics, Vol. 38, Academic Press, New York-London, 1970.
15. C. Fefferman and E. M. Stein, *H^p spaces of several variables*, Acta Math. **129** (1972), no. 3-4, 137–193.
16. S. H. Ferguson, *Polynomially bounded operators and Ext groups*, Proc. Amer. Math. Soc. **124** (1996), no. 9, 2779–2785.
17. S. H. Ferguson and M. T. Lacey, *A characterization of product BMO by commutators*, Acta Math. **189** (2002), no. 2, 143–160.
18. R. F. Gundy and R. L. Wheeden, *Weighted integral inequalities for the nontangential maximal function, Lusin area integral, and Walsh–Paley series*, Studia Math. **49** (1973/74), 107–124.
19. G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge University Press, 1988.
20. H. Hedenmalm, P. Lindqvist, and K. Seip, *A Hilbert space of Dirichlet series and systems of dilated functions in $L^2(0, 1)$* , Duke Math. J. **86** (1997), no. 1, 1–37.
21. H. Helson, *Hankel forms and sums of random variables*, Studia Math. **176** (2006), no. 1, 85–92.
22. ———, *Hankel forms*, Studia Math. **198** (2010), no. 1, 79–84.
23. M. T. Lacey and E. Terwilleger, *Hankel operators in several complex variables and product BMO*, Houston J. Math. **35** (2009), no. 1, 159–183.
24. Z. Nehari, *On bounded bilinear forms*, Ann. of Math. (2) **65** (1957), 153–162.
25. J. Ortega-Cerdà and K. Seip, *A lower bound in Nehari’s theorem on the polydisc*, J. Anal. Math. **118** (2012), no. 1, 339–342.
26. G. Pisier, *A polynomially bounded operator on Hilbert space which is not similar to a contraction*, J. Amer. Math. Soc. **10** (1997), no. 2, 351–369.
27. H. Queffélec and M. Queffélec, *Diophantine approximation and Dirichlet series*, Harish-Chandra Research Institute Lecture Notes, vol. 2, Hindustan Book Agency, New Delhi, 2013.

Ole Fredrik Brevig

Department of Mathematical Sciences, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway

e-mail: ole.brevig@math.ntnu.no

Karl-Mikael Perfekt

Department of Mathematical Sciences, Norwegian University of Science and Technology (NTNU), NO-7491 Trondheim, Norway

e-mail: karl-mikael.perfekt@math.ntnu.no