

## *Host-plant resistance*

Book or Report Section

Accepted Version

Van Emden, H. F. (2017) Host-plant resistance. In: Van Emden, H. F. and Harrington, R. (eds.) *Aphids as Crop Pests*, 2nd edition. CABI, Wallingford, pp. 515-532. ISBN 9781780647098 Available at <http://centaur.reading.ac.uk/72415/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Publisher: CABI

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# 22 Host-Plant Resistance

**Helmut F. van Emden**

*School of Agriculture, Policy and Development, The University of Reading, Earley Gate, Reading, Berkshire, RG6 6AR, UK.*

## Introduction

In this section, the word 'variety' will be used as an umbrella word for the many types of variation involved; varieties, cultivars, accessions, breeders' lines may all be included. The practical use of host-plant resistance (HPR) for aphid control in different crops is explored in the pest management section of this volume (Chapters 24-33).

It might be assumed that the role of symbionts in nitrogen metabolism (Chapter 6, this volume) should enable aphids to compensate for nutritional differences between crop varieties, and that the phloem-feeding habit would enable aphids to avoid many deterrent surface characters and allelochemicals in plants. However, the literature probably has more examples of HPR to aphids than to any other group of crop pests, including examples of nearly all the mechanisms of HPR known for insect pests in general. Moreover, HPR to aphids can be highly effective and dramatic; for example, the resistance of 'Avoncrisp' lettuce to *Pemphigus bursarius* (lettuce root aphid) (Fig. 22.1).

Resistance rankings between varieties can change with age and resistance to one aphid species may not give resistance to another. HPR to *Diuraphis noxia* (Russian wheat aphid) in wheat and barley seems especially specific, and has no effect on the several other species of cereal aphid (Robinson, 1992; Messina and Bloxham, 2004).

In recent years, molecular biology techniques have become routine, and much progress has been made in screening aphid populations for genetic variability. However, whether or not such genetic variability matches phenotypic variability (Thomas *et al.*, 2012), only differences in reaction to different varieties are relevant to HPR (Pompon *et al.*, 2011). Also, possibly because of the success against other pests of crops genetically engineered to express the *Bacillus*

*thuringiensis* (Bt) toxin, the emphasis in HPR studies on aphids has shifted to identifying genes (e.g. Shufran and Payton, 2009). This puts emphasis on single genes coding for proteins expressing a high level of toxins, a narrow and perhaps not the most desirable HPR mechanism (Chapter 23, this volume).

Before the advent of direct gene transfer, wild relatives within the natural crossing barriers were often the source of resistance genes (e.g. Hesler, 2013). Table 22.1 lists some of the wild relatives of crops often involved in resistance breeding for aphids, usually entailing the transfer of characteristics (e.g. hairiness, toughness, high toxin levels) lost in the process of domestication (van Emden, 1997).

Finally, HPR need not necessarily involve genetic change. It can sometimes be produced by some extrinsic influence on a susceptible variety ('pseudoresistance' or 'induced resistance'). Plant resistance to *Schizaphis graminum* (greenbug) in wheat after fertilization with silicon (Gomes *et al.*, 2005) is one example. HPR also is manifested when the aphid pest does not synchronize with a 'susceptibility window' in the plant's phenology. Plant resistance to aphids generally tends to increase with age ('age-related resistance'); Sarwar and Sattar (2013) reported that early-sown oilseed rape was effective in reducing incidence of and damage by *Myzus persicae* (peach–potato aphid).

Associated organisms (see 'endophytes' below) may also induce HPR.

## **Types of Host-Plant Resistance to Aphids**

The classification of Painter (1951), modified by Kogan and Ortman (1978), still has great value.

### **Antixenosis (close to Painter's 'non-preference')**

This is resistance to colonization by aphids, and shows in the high proportion of immigrating alatae which take off from a plant after superficial probes and within a few hours of landing. At the immigration stage, aphids seem to exhibit a high degree of non-preference, even on susceptible plants. Müller (1958) showed that the eventual four-fold greater population of *Aphis fabae* (black bean aphid) on the susceptible broad bean variety Schlandstedt than on the resistant Rastatt was based on respective re-take-off rates of 95% and an only slightly higher 99%. Antixenosis is usually assessed by liberating alatae over different varieties, both in choice and no-choice situations (Adams and van Emden, 1972), although van Emden *et al.* (1991) devised a test

(involving systemic insecticide with a correction for any detection thereof by the aphids) that identifies which aphids have ingested from the plants.

Tjallingii's (1988) electrical penetration graph (EPG) technique (Chapter 9, this volume) enables the continuous monitoring of aphid penetration and probing activity (Fig. 22.2) (Mentink *et al.*, 1984).

### **Antibiosis**

This negatively affects the multiplication of aphids that have colonized the plant, usually showing as reduced survival, growth and fecundity, and extended development time. These are measured easily on individuals confined on leaves in small clip cages, although care needs to be taken to ensure that any apparent antibiosis is no more than a first-generation expression of transfer to a novel plant. Survival, development time, and fecundity can be combined into a single statistic, the intrinsic rate of increase ( $r_m$ ), and another useful single statistic is the integral of the daily weight increase of individual aphids over a few days (mean relative growth rate or MRGR) (Chapter 5, this volume).  $r_m$  can readily be converted into 'population doubling time'.

### **Tolerance**

Tolerance of a variety shows as a better yield under the same aphid burden, both in numbers and duration. This is difficult to quantify, since any antibiosis defeats the 'equal burden' requirement, and intolerance to infestation will result in plant damage, which may cause a crash in the aphid population and defeat the 'equal duration' requirement. Havlickova (1997) has proposed a covariance model in which the main component is the slope of the relationship between the weights of infested and control seedlings. However, tolerance may perhaps be better quantified by the slopes of regressions of yield or plant biomass on initial aphid infestation.

### **What is the preferred type of resistance?**

Antixenosis is rarely effective in a no-choice situation, and puts pressure on the population for selection of genotypes not affected by the antibiosis (see 'biotypes' later), a danger not applicable to tolerance. Burd and Puterka (2009) challenge the prioritization of tolerance over antibiosis on

the grounds that crop selection was not a factor in the appearance of adapted greenbug genotypes. Moreover, a potential hazard of relying on tolerance is that farmers growing such varieties would not need to control aphids and so would breed populations to infest their neighbours' crops. Thus, there is general agreement (van Emden, 1997) that the preferred resistance is antibiosis coupled with some antixenosis, as has been found in some soybean lines (Hesler, 2013). Buschman and Ramaswamy (2012) have modelled HPR stability arising from two genes, one expressing an oviposition deterrent and the other a toxin. When the antibiosis was dominant, insect populations were managed for 150 years, but for much longer when it was recessive, presumably because the toxin was modelled as a stronger selection pressure than the deterrent. The combination of antibiosis and tolerance had also been recommended (Murugan *et al.*, 2010) as delaying biotype selection in *D. noxia*.

This chapter reviews HPR to aphids by the various anatomical, physiological, and biochemical mechanisms that have been correlated with HPR phenomena and Fig. 22.3 shows these with an indication of the order in which the aphid would encounter them. However, correlations are not the same as 'cause and effect'. Kazemi and van Emden (1992) found that some proposed chemical correlations with HPR in wheats to *Rhopalosiphum padi* (bird cherry–oat aphid) broke down when a wider range of wheats was tested. Similarly, Weibull (1994) found that correlations of chemistry with resistance to *R. padi* in parental barley lines broke down in the F1 generation. It seems it is not difficult to correlate resistance rankings with whatever chemical group represents one's own particular interest.

## **Mechanisms of Host-Plant Resistance to Aphids**

### **Mechanisms of antixenosis**

#### *Colour*

Colour can influence the preference of immigrating aphids. Alatae of *Brevicoryne brassicae* (cabbage aphid) do not settle well on red cabbage varieties, even though apterae caged on such varieties have a better growth rate than those on green varieties (Radcliffe and Chapman, 1965, 1966). Similarly, in a study of 50 mustard and oilseed rape varieties in India (Rana *et al.*, 2001), purple-leaved cultivars had fewer colonizing *Lipaphis erysimi* (mustard aphid) than dark green

ones, which in turn were less infested than those that were light green. Differences in the brightness of the yellow flowers had no effect on aphid colonization.

### *Palatability*

That so many common or Latin names of aphids reflect particular host plants (e.g. lupin aphid, *B. brassicae*, etc.) shows that aphids often accept or reject a plant on the basis of its secondary chemistry, even before reaching the phloem. Thus toxins (which would otherwise reflect antibiosis) become deterrents. Polyphagous aphids such as *M. persicae* will tolerate levels of secondary compounds such as glucosinolates in brassicas, but elevating levels still tolerable to the specialist *B. brassicae* will lead to antibiosis (van Emden, 1978; Cole, 1997b). In screening wild and cultivated brassicas for resistance to these two aphids, Cole (1997a) found that most of the variation in HPR was explained by four glucosinolates (2-OH-3-butenyl, 2-propenylglucosinolate, 3-methoxyindolyl and 4-pentylglucosinolate). High total phenol (especially ortho-dihydroxy phenol) content in oilseed brassicas deters infestation by *L. erysimi* (Sarwan Kumar Sangha, 2013) and the phenolic hydroxamic acids (Hx), often associated with antibiosis to aphids in cereals, are also a deterrent to *Metopolophium dirhodum* (rose-grain aphid), *Sitobion avenae* (English grain aphid) and *Schizaphis graminum*. *Rhopalosiphum maidis* (corn leaf aphid), however, is not deterred because it avoids Hx in high Hx plants by reducing puncturing of the cells by the stylets on their way to the phloem (Givovich and Niemeyer, 1995). In lucerne, other Hx compounds (the 3-amyl-4-hydroxycoumarins such as coumestrol) are deterrent to *Acyrtosiphon pisum* (pea aphid), as are saponins (Nicholas *et al.*, 2005). In tobacco, the antixenosis of some varieties to *M. persicae* is ascribed to sugar ester levels and alpha and beta monols on the leaf surface (Johnson *et al.*, 2002).

One mechanism of antixenosis in tomato to *Macrosiphum euphorbiae* (potato aphid) involves ethylene signalling. This has been shown to induce opposite effects in susceptible and resistant tomato varieties. It promotes aphid infestation on susceptible genotypes, but contributes to antixenosis in resistant ones (Wu *et al.*, 2015).

Recently a trial of GM wheat with a gene (resynthesized from peppermint to eliminate inhibitory compounds) expressing the alarm pheromone of aphids was carried out (Bruce *et al.*, 2015). Although cereal aphids were deterred in laboratory tests, no substantial reductions in the population of cereal aphids were observed in the field, but the idea of using behaviour-modifying compounds in this way is novel and has potential.

## *Waxiness*

Waxiness of the leaf surface (Fig. 22.4) has sometimes been positively linked with antixenosis to aphids, e.g. the epicuticle of some lucerne varieties resistant to *Therioaphis trifolii maculata* (spotted alfalfa aphid) had wax ester levels 50% higher than those in susceptible lucerne (Bergman *et al.*, 1991). Similarly Ren *et al.* (2014) found that the wax content on the leaf surface of the highly *Aphis gossypii*-resistant cucumber variety 'Axin' was 30 times higher than for the susceptible variety 'Pepino'. Wojcicka (2015) used the EPG technique (see earlier) to show that components in the wax of waxy genotypes of wheat completely stopped salivation and active ingestion by *R. padi*, yet it is often the less waxy (glossy) varieties that are resistant because of higher levels of deterrent chemicals in the reduced wax layer. Glossy wheats, antixenotic to *S. avenae*, contain dihydroketones as the deterrent compounds (Lowe *et al.*, 1985). With *A. pisum* studied on seven pea varieties, the glossy ones again were resistant (White and Eigenbrode, 2000), and glossy brassica varieties are also antixenotic to *B. brassicae* (Thompson, 1963; Ellis *et al.*, 1996), with reductions in aphid populations as high as 95% (Stoner, 1992).

Leaf surface wax components have also been implicated in the resistance of European red raspberries to *Amphorophora idaei* (large European raspberry aphid) conferred by gene  $A_{10}$  (Robertson *et al.*, 1991; Shepherd *et al.*, 1999a,b).

Ni *et al.* (1998) tested whether the ranking of resistance to *D. noxia* of wheat (most susceptible), oat (intermediate), and barley (most resistant) had a basis in surface wax by removing this with ethyl ether, but the resistance rankings remained as before.

## *Mechanical*

Hardness of the plant surface is a common resistance mechanism to chewing insects but not for aphids, where a more common cause of mechanical antixenotic resistance is difficulty in reaching the phloem. Pectin in the cellular middle lamella has a role in hindering access to the phloem (Dreyer and Campbell, 1984). Resistance to *S. graminum* in some sorghum varieties is due to increased methylation of the middle lamellar pectin (Dreyer and Campbell, 1984). A variety of melon resistant to *A. gossypii* deposits callose in attacked leaf veins.

A major qualitative chemical feature of two wild brassicas resistant to *B. brassicae* was the presence of gluconapin rather than the more usual glucobrassicin, but the former had no effect

on the aphid in artificial diet. Cole (1994a) proposed that the resistance mechanism was stylet blocking.

### *Local necrosis*

Local necrosis, i.e. the death of cells with the production of deterrent polyphenols wherever the aphid tries to insert its stylets, is a valuable resistance mechanism since it also gives protection against other piercing organisms such as fungal hyphae and nematodes (van Emden, 1987). The polyphenols are produced when aphid damage brings substrate and enzyme into contact. This is a 'hypersensitive response', perhaps an unusual concept of 'resistance'. It is the mechanism of resistance of apples to *Eriosoma lanigerum* (woolly apple aphid) (Wartenberg, 1953) and *Dysaphis plantaginea* (rosy apple aphid) (Alston and Briggs, 1970). Another example is the hypersensitive cell death response of some resistant barleys to attack by *D. noxia* (Belefantmiller *et al.*, 1994), though some aphid species may have reducing compounds in their saliva that counter the hypersensitive response of the plant (Miles, 1999).

### *Trichomes (non-glandular)*

Non-glandular trichomes are mechanical barriers to many small insects, including aphids. For example, high trichome density on wheat leaves deters *Sipha flava* (yellow sugar cane aphid), but any effect on *S. graminum* is questionable (Webster *et al.*, 1994). Pompon *et al.* (2010) attributed the resistance of two wild potato species to *M. euphorbiae*) and *Myzus persicae* to the trichomes on the leaf surface; high trichome density also deters *M. persicae* on crosses of tomato with wild potato (Simmons *et al.*, 2005). Pubescent broccoli is as resistant to *B. brassicae* as glossy varieties (see earlier) (Stoner, 1992). However, with *A. gossypii* on cotton, it is the glabrous varieties that have fewer aphids (Weathersbee *et al.*, 1994, 1995).

## **Mechanisms of antibiosis**

### *Glandular trichomes*

These are of two types: long slender ones that secrete drops of fluid from the tip, and short ones bearing a spherical glandular head (Fig. 22.5). Septa in the head separate substrate and enzyme

which are brought together to form polyphenols when the head is ruptured by contact with an aphid. These polyphenols are not only distasteful to the aphid (see earlier), but also harden on and disable the mouthparts and tarsi. Great interest has focused on breeding the presence of such trichomes on wild potato (especially *um berthaultii*) and tomato crossed with *Lycopersicon pennellii* (Simmons *et al.*, 2005) into varieties for resistance to aphids (Gibson, 1976), but none was ever commercialized. Any *M. euphorbiae* on the stems, however, multiply faster than on varieties without glandular trichomes (Ashouri *et al.*, 2001). The wild tomatoes *Solanum habrochaites* and *S. peruvianum* have a dense pubescence with both glandular and non-glandular trichomes (Kok-Yukomi, 1978), and Dreyer and Campbell (1987) observed that the resistance attributable to the glandular trichomes decreased as the density of non-glandular trichomes increased.

### *Toxins*

These are often the same chemicals that can also cause antixenosis, and they may only be induced following disruption of plant cell walls by aphid probing. The plant then produces signalling compounds such as jasmonic, salicylic, abscisic and gibberellic acids which lead to the mobilization of direct chemical defence responses (Smith and Boyko, 2007; Pompon *et al.*, 2010; Chapter 8, this volume). The resistance to *M. euphorbiae* of tomatoes carrying the Mi-1.2 gene results from jasmonic acid triggering a proteinase inhibitor (Cooper *et al.*, 2004).

Hydroxamic acids (Hx) are deterrent to many aphids (see earlier), but also decrease feeding in artificial diet of, for example, *S. avenae*. Also in diet, 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one (DIMBOA) decreases the survival of *M. dirhodum* (Cambier *et al.*, 2001), as do low concentrations of its decomposition product 6-methoxybenzoxazolin-2-one (MBOA) for *S. avenae* (Hansen, 2006). In whole plant studies, a wheat cultivar with high Hx was found to be antibiotic to *S. avenae* (Fuentes-Contreras and Niemeyer, 1998), and the growth rate of the aphid on three wheat and two oat cultivars was negatively correlated with Hx levels (Fuentes-Contreras *et al.*, 1996). Also, 26 Hungarian cultivars showed a negative correlation between Hx levels and populations of *R. padi* (Gianoli *et al.*, 1996).

Legumes are also well known to contain toxins. Ansari *et al.* (1989) discovered some varieties of cowpea (*Vigna unguiculata*) in which the leaves produced a powerful graft-transmissible but unidentified toxin that killed *Aphis craccivora* (cowpea aphid) very quickly, but did not seem to be detected by the aphid. Another example is lupin; the alkaloids in narrow-leaved lupins (*Lupinus angustifolius*) depress the fecundity of *Macrosiphum albifrons* (lupin

aphid) (Berlandier, 1996). *Myzus persicae* is less affected by these alkaloids (gramine, sparteine, lupanine, lupinine, 13-hydroxylupanine, and angustifoline) than *A. craccivora*, both in plants and in artificial diet (Ridsdill-Smith *et al.*, 2004).

On cotton, *A. gossypii* showed shorter longevity and lower fecundity on varieties bred to have high levels of the polyphenol gossypol than on two with lower levels (Du Li *et al.*, 2004).

Unique among the role played by alkaloids in plant defence against insects, including aphids, is the production of nitrogen-rich compounds by asexual *Epichloe/Neotyphodium* spp. facultative fungal endophytes of temperate grasses (Fig. 22.6). The bioactive compounds produced by these endophytes are ergot alkaloids (particularly ergovaline), indole diterpenes (lolitrems), pyrrolopyrazines (peramine), and pyrrolizidines (lolines). Ergot and lolitrem, but not peramine and lolines, are toxic to livestock (Young *et al.*, 2013).

Lolines alone protect infected tall fescue and perennial rye grass (*Lolium pratense*) from *R. padi* (Siegel *et al.*, 1990; Eichenseer *et al.*, 1991; Wilkinson *et al.*, 2000; Schardl *et al.*, 2007). Additionally, lolines alone in meadow fescue (Wilkinson *et al.*, 2000) and peramine only in other infected grasses (Siegel *et al.*, 1990) have been associated with resistance to *S. graminum*. Compared to endophyte-free plants, infected perennial ryegrass (Clement *et al.*, 1992), tall fescue (Clement *et al.*, 1996), and some wild perennial barley (*Hordeum* spp.) (Clement *et al.*, 1997), are strongly resistant to *D. noxia*. Endophyte infection of another wild temperate grass, *Phleum alpinum*, confers resistance to *R. padi* (Clement *et al.*, 2011). The expression of insect resistance (antixenosis, antibiosis) is affected by the host grass species/genotype, endophyte strain (including associated alkaloid profile) and insect species/biotype involved in a given interaction (Clement *et al.*, 1994; Popay, 2009; Crawford *et al.*, 2010) and soil nutrition (Lehtonen *et al.*, 2005).

Perennial ryegrass cultivars harbouring a specific endophyte strain (AR37) are used for protection against the root aphid *Aploneura lentisci*, and other pasture pests on farms in New Zealand (Hume *et al.*, 2007; Popay and Hume, 2013) and Australia (Moate *et al.*, 2012).

Although endophytic fungi are also frequently found in dicotyledenous plants (Saikkonen *et al.*, 1998), any role they may have in HPR to aphids (or indeed any other insect) has yet to be established.

*Bacillus thuringiensis* (Bt) toxins used for GM Lepidoptera resistance are not regarded as aphicides, but a GM potato cultivar with Bt against *Leptinotarsa decemlineata* (Colorado potato beetle) also depressed the growth and fecundity of *M. euphorbiae* (Ashouri *et al.*, 2001).

Other toxins studied specifically for their potential in GM crops against aphids include lectins, but these usually give only partial resistance to aphids. When 30 were tested in artificial diet against *A. pisum* (Rahbe *et al.*, 1995), most caused only low toxicity. However, those from

jack-pine (*Pinus banksiana*), love-lies-bleeding (*Amaranthus caudatus*), lentil (*Lens culinaris*), snowdrop (*Galanthus nivalis*) induced worthwhile mortality. Concanavilin A (the jack-pine lectin) was then tested on five other aphid species, but with very variable results. A chitin-binding lectin in two wild brassicas shows some insecticidal activity against *B. brassicae* (Cole, 1994b). There have also been studies with proteinase inhibitors. Oryzacystatin 1, effective against leaf-chewing insects in transgenic oilseed rape, inhibited growth of *A. gossypii*, *A. pisum*, and *M. persicae* (Rahbe *et al.*, 2003), and (in transgenic eggplant) of *M. persicae* and *Macrosiphum euphorbiae* (Ribeiro *et al.*, 2006). Tobacco genetically engineered to accumulate two ketosteroids (cholestan-3-one and cholest-4-en-3-one) in the phloem caused high mortality of *M. persicae* (Bouvaine *et al.*, 2014).

### *Nutritional factors*

Given the limiting levels of nitrogen dietary requirements for aphids in the phloem (Chapter 6, this volume), one might expect nutrition to be a valuable source of HPR. AucLair *et al.* (1957) studied several pea varieties and found a negative correlation between their resistance to *A. pisum* and the soluble and total amino nitrogen content of the leaves. Kazemi and van Emden (1992) analysed a wide provenance range of wheats (European, Iranian and the 'ancient' diploid wheat Einkorn (*Triticum monococcum*)) and found that the leaf concentration of three amino acids (alanine, histidine and threonine) was an excellent predictor for the performance of *R padi*, accounting for over 95% of the variation in the fecundity of the aphid. By contrast, Weibull (1994) analysed EDTA (ethylenediaminetetraacetic acid) leaf exudates as a better reflection of phloem content than whole leaf analyses, but found a negative correlation between several amino acids and the performance of *R. padi* on common barley (*Hordeum spontaneum*). However, since the segregating offspring of crosses showed no such relationship, no cause and effect was attached to the correlation.

Although nutritional factors are a mechanism of HPR to aphids, such resistance is particularly likely to fail under different environmental conditions such as different soils and fertilizer regimes.

### *Extrinsic factors*

Factors extrinsic to the plant may be responsible for greater mortality on the apparently 'resistant' than on the 'susceptible' variety. In the absence of such factors, true HPR may be much less, or even absent. Thus, Gowling and van Emden (1994) showed that the number of *M. dirhodum* falling from wheat plants was doubled when parasitoids were added, and doubled again if the wheat variety was partially aphid-resistant. The partially resistant variety 'Rapier' was only 25% resistant compared with the susceptible 'Armada', yet the increased falling of aphids in the presence of searching parasitoids caused it to appear 86% resistant.

Another example of extrinsic resistance is that of the 'leafless' pea varieties bred for low leaf area to be resistant to powdery mildew (*Erysiphe polygoni*) and, instead of leaves, it has a profusion of photosynthesizing tendrils (Fig. 22.7) (Kareiva and Sahakian, 1990). The apparent resistance to *A. pisum* in the field was considerably greater than the true intrinsic resistance of only some 8%, because the ladybirds *Coccinella septempunctata* and *Hippodamia convergens* could grip the tendrils, whereas a high proportion fell off the shiny leaves of normal peas. The proportion falling was from 47 to 26% for *C. septempunctata* and from 32 to 9% for *H. convergens*.

### **Mechanisms of tolerance**

Rather little is known about the mechanisms of tolerance of plants to aphids. Tolerance is often coupled with antibiosis, and the latter may explain the increased yield sufficiently for tolerance not to be identified. The examples of tolerance reported in the literature therefore tend to be dramatic, with crop growth and yield little affected despite high aphid infestation. For example, Hesler (2005) found that three triticale (x *Triticosecale* spp.) accessions showed difference in shoot length whether infested or uninfested with *R. padi*.

### *Compensation*

Some lines of wheat show much enhanced growth with increased tillering when attacked by *S. graminum* or *D. noxia* (Castro *et al.*, 2001). Such compensatory growth as a response to aphid attack may, however, lead to a delay in harvest.

### *Symptom expression*

The plant reaction to aphid saliva can vary with the particular genotypes of both aphid and plant involved. Thus, some genotypes of wheat are specifically tolerant to the E biotype of *S. graminum* with a lack of symptoms (Morgham *et al.*, 1994). A mechanism for such tolerance to *S. graminum* was proposed by Maxwell and Painter (1962), who compared two wheats, 'Dickson' (tolerant) and 'Pawnee' (intolerant). These two varieties differed as to the timing of the cessation of further expansion of the flag leaf, on the photosynthesis of which much of the grain yield depends. Since the aphids tend to feed in an aggregation on the underside of the leaf, growth there is inhibited, while the upper leaf surface continues growing. The leaf therefore curls strongly, reducing the effective leaf area exposed to solar radiation. In 'Dickson', the growth of the flag leaf stops earlier than with 'Pawnee', and feeding by the aphids on the underside of the leaf does therefore does not induce curling.

## **Further Considerations**

### **Yield drag or other fitness costs**

The majority of HPR mechanisms involve the production by the plant of more chemical compound or additional plant tissue, yet fitness costs of pest-resistance have rarely been reported. Thus, Assad *et al.* (2004) found no negative correlations between the height, grains per spike or total biomass of 14 wheat genotypes and their antixenosis to *D. noxia*, and in two Rag genes for resistance to *Aphis glycines* (soybean aphid), even when pyramided (i.e. combined on the same variety), incurred no yield penalty (Hesler *et al.*, 2013). Data of Gershenzon (1994) suggest that different secondary compounds can account for between 0.01 and 30% of the glucose in leaf tissue, but such static concentrations are negligible in comparison with the glucose production by photosynthesis over a period of time. Foyer *et al.* (2007) calculated that 30-40 min of photosynthesis (4-6 % of the total photosynthetic output over a 12 h day) would suffice to synthesize the DIMBOA in wheat and maize. Moreover photosynthesis is more often sink- than source-limited, and so any fitness costs would reveal themselves only under conditions of extreme stress.

### **Negative effects on natural enemies**

Crop varieties developed for resistance to aphids can have deleterious effects on natural enemies, especially where the resistance is based on toxins (see above). DIMBOA, a hydroxamic acid studied extensively in relation to HPR to cereal aphids (see earlier), is toxic to the ladybird *Eriopsis connexa*. Paradoxically, maize varieties with high DIMBOA cause less mortality of this predator than varieties with intermediate levels, since it can detect the compound in the aphids and avoid ingesting a toxic dose (Martos *et al.*, 1992). In contrast, the ladybird *Propylaea japonica* showed increased weight and faster development on a high gossypol aphid-resistant cotton variety (Du Li *et al.*, 2004).

The parasitoid *Aphidius rhopalosiphi*, reared on the smaller individuals of *M. dirhodum* on the partially aphid-resistant wheat 'Rapier', showed progressively smaller size and reduced egg load on emergence in consecutive generations. After ten generations on 'Rapier', size and egg load had decreased by 15 and over 50%, respectively. However, parasitoids returned to an aphid-susceptible variety ('Huntsman') regained their fitness in one generation (Salim Jan and H. F. van Emden, unpublished but summary data given by van Emden, 1995). Reductions in size and fecundity of *Aphidius nigripes* emerging from *M. euphorbiae* reared on Bt potato have also been attributed to the poorer quality of the host aphids (Ashouri *et al.*, 2001; Ashouri, 2004). *Aphis glycines* showed such reduced longevity on aphid-resistant soybeans that fewer *Binodoxys communis* could develop to the mummification stage (Ballman *et al.*, 2012).

## **Problem trading**

There is therefore always the possibility that the plant modification made in developing HPR to aphids will make the plant more susceptible to another damaging organism. An early example is found in the hairy South African cottons, resistant to leafhoppers but especially susceptible to *A. gossypii* (Dunnam and Clark, 1938). When lucerne varieties resistant to lucerne wilt (*Verticillium albo-atrum*) were released in California, they proved especially susceptible to *T. trifolii* (van den Bosch and Messenger, 1973).

Glossy brassica varieties resistant to *B. brassicae* suffer from increased *Phyllotreta cruciferae* (flea beetle) damage compared with waxy ones (Stoner, 1992; Eigenbrode *et al.*, 2000), and glossy peas resistant to *A. pisum* are more heavily attacked by *Sitona lineatus* (pea and bean weevil) (White and Eigenbrode, 2000).

Surprisingly, problem trading has also been reported in tall fescue plants resistant to *R. padi* because of alkaloids produced by the fungal endophyte *Neotyphodium coenophialum*. The

noctuid armyworm *Spodoptera frugiperda* showed enhanced performance on the endophyte-infected plants (Bultman and Bell, 2003).

In breeding F2 hybrid willows as biomass fuels, resistance to a *Chaitophorus* sp. aphid and the eriophyid mite *Aculus tetanothrix* were found to be inversely related (Czesak *et al.*, 2001).

## **Biotypes**

Species of aphid often show genotypic variation expressed through different behavioural traits, including sensitivity or tolerance to varieties bred for HPR. Tolerant variants are often referred to as 'resistance-breaking biotypes', but Blackman and Eastop (Chapter 1, this volume) express doubts about the use of the word because such genotypes may be just a single clone subject to recombination by sexual reproduction. However, the word 'biotype' is so well embedded in the literature of HPR that it would probably be misleading to the reader to use an alternative terminology here.

Some cowpea varieties producing a toxin strongly aphicidal to *A. craccivora* collected at the International Institute for Tropical Agriculture (IITA) in Nigeria proved susceptible to the first aphids collected from another location (Ansari, 1984), and the 'biotype problem' with pests indeed seems particularly common with aphids; 8% of 423 references between 1972 and 2015 concerning HPR and aphids are about this aspect, though many identify resistance-breaking biotypes in the glasshouse rather than reporting a problem in the field.

Biotypes in relation to HPR have proved particularly important with *S. graminum* on cereals, with the biotypes distinguished according to their responses to a clearly defined set of plant varieties. The resistance-breaking biotypes have proved a major headache in the field. For example, biotype E breaks the resistance incorporated in some sorghum varieties (Eisenbach and Mittler, 1987). No single resistance gene in either sorghum or wheat is effective against all biotypes of *S. graminum*. Anstead *et al.* (2003) located several biotypes of *S. graminum* on wild grasses, including one new biotype, and proposed that new biotypes are formed on wild grasses rather than on cereal crops.

The use of HPR against *A. glycines* in soybean has similarly identified at least three biotypes, though it is still possible to find soybean accessions resistant to all (Bansal *et al.*, 2013).

Attempts in the USA to manage *D. noxia* on cereals ran into biotype problems only five years after the appearance of the pest in Texas in 1986. Five biotypes have been distinguished (Shufran and Payton, 2009). The emergence of biotype 2 in Colorado made all resistance genes,

other than one in rye, ineffective (Murugan *et al.*, 2010). However, even the resistance-breaking biotypes vary in the chlorosis and stunting they cause (Shufran *et al.*, 2007). Some of the resistance genes no longer effective in the USA are still effective in South Africa (Tolmay *et al.*, 2013). Merrill *et al.* (2014) released two biotypes of *D. noxia* differing in their tolerance to resistant wheat in a known proportion at three sites along a gradient from northern Colorado to Florida, and found that the resistance-breaking biotype rapidly displaced the other, with an apparent more than 10-fold advantage.

Since the late 19th century, grapes in most parts of the world have been grafted on to rootstocks of old communion wine varieties resistant to *Daktulosphaira vitifoliae* (grape phylloxera) and brought from Europe to North America by the early settlers. This resistance lasted for a long time, but was finally broken by a biotype appearing in Germany in 1994 (Anonymous, 1994). Thus, in spite of the potential for 'biotypic breakdown' of resistance, HPR to aphids may be quite durable. Di Pietro and Caillaud (1998) reared *S. avenae* on resistant wheat (including highly resistant 'Einkorn ') for two years and could find no selection for breaking the HPR.

As can be seen from the above, the occurrence of biotypes is variable both between and within species (in relation to geographical distribution). Puterka *et al.* (1992) recorded seven biotypes of *D. noxia* worldwide, but as yet only a few are found in the USA. A detailed study of *Elatobium abietinum* (spruce aphid) compared 40 populations each from the UK and New Zealand (Nicol *et al.*, 1998). There were 28 genotypes (but of course not necessarily 28 biotypes) in the UK collections, but only one in New Zealand.

There is little information as to the mechanisms whereby aphid biotypes break HPR. It is, however, known that resistance to *S. graminum* in sorghums, based on increased methylation of the middle lamellar pectin, is defeated in resistance-breaking biotypes of the aphid by their saliva possessing enhanced pectin methyltransferase activity (Dreyer and Campbell, 1984). Biotypes of *Nasonovia ribisnigri* (currant-lettuce aphid) not affected by the Nr gene in resistant lettuce suppress the wound response of the sieve elements (ten Broeke, 2013). There is also the intriguing hypothesis that it is the symbionts that enable biotypes of *T. t. maculata* to break the resistance of some lucerne varieties (Ruggie and Gutierrez, 1995) and of some *M. euphorbiae*-resistant tomato varieties (Francis *et al.*, 2010).

Coping with the biotype problem involves the continual development of varieties with new resistance genes available if control begins to fail, as has been the history with *S. graminum* (Puterka and Peters, 1990). The breakdown of plant resistance is usually far more serious with fungal diseases, and plant pathologists have developed strategies for deploying resistance genes so as to delay such breakdown. These strategies are equally applicable to HPR against aphids.

One such strategy is combining more than one resistance gene in a variety ('pyramiding'). Porter *et al.* (2000) used a model for *S. graminum* to test whether pyramiding or sequential release of genes would maintain HPR for longer. The model predicted that pyramiding was the longer-lasting strategy, but Porter *et al.* claimed that experimental evidence actually pointed to the opposite conclusion. Bush *et al.* (1991) studied the value of variety mixtures against greenbug biotypes C and E. They used three varieties of wheat, one susceptible to the aphid (S) and one resistant to each of the biotypes C and E (CR and ER). The mixtures 3R:1S in separate experiments for each biotype, or 1CR:1ER:1S with both biotypes, managed the biotypes successfully, and the two-variety mix worked as well as the three-variety one.

### **Spread of viruses** (see also Chapter 15, this volume)

Some workers have argued that, because aphids on resistant varieties are more restless, the spread of virus in such varieties is likely to be accelerated, particularly with non-persistent virus diseases. Atiri *et al.* (1984) reported such greater spread of *Cowpea mosaic virus* in small cage experiments with *A. craccivora* on cowpeas. Other work in Africa (Roberts *et al.*, 1993) appears to confirm this in field experiments on chemical control of the same aphid with pyrethroids (which increase restlessness of aphids – van Emden and Service, 2004). In these experiments, the secondary spread of virus was greater on an aphid-resistant than on a susceptible cowpea variety. It is also possible that the phenomenon may occur with aphids in watermelon (Webb and Linda, (1993) – but see Kishaba *et al.* (1992) below) and with *M. persicae* in sugar beet, on which this aphid shows high mortality, especially as the plants get older. However, survival is improved on plants infected with either *Beet yellows virus* or *Beet mild yellowing virus* (Kift *et al.*, 1996). However, such reports are in the minority, perhaps partly because much of the spread of non-persistent viruses is by alatae of species for which the crop is not a host, and therefore totally resistant regardless of its resistance status to other aphid species. Even in respect of pest species, the overwhelming evidence is that HPR to aphids leads to less virus spread (van Emden, 1987). Thus Kishaba *et al.* (1992) compared *Watermelon mosaic virus 2* transmission by *A. gossypii* in one aphid-susceptible and three resistant water melon lines that were all equally good virus sources, and yet transmission by the aphid on the three resistant lines was reduced by 31, 71, and 74%. The number of aphids needed to give 59% infection was only 20 per plant on the susceptible line, but varied from 60-400 on the three aphid-resistant lines. Chen *et al.* (1997) also found that, for the water melon/*A. gossypii* combination, levels of non-persistent virus were negatively correlated with resistance to the aphid.

## **Interactions with other control measures**

The aphid literature provides many examples of HPR having a positive effect on the impact of natural enemies. HPR also affects the susceptibility of aphids to insecticides. Such interactions are not discussed here, but are reviewed in Chapter 23, this volume.

## **Conclusions**

Host-plant resistance has far greater potential for reducing populations of aphids than has as yet been exploited. This is partly because plant breeders have sought to use HPR as a single-component control measure, and have therefore screened for high resistance to aphids. Such levels of resistance are most likely to be based on toxic allelochemicals, often the expression of a single gene. This has certainly been the approach with the industrial production of insect-resistant transgenic crops, although as yet no transgenic event effective against aphids has been commercialized.

It will be evident from this account of HPR that such monogenetic resistance/ high expression of allelochemicals is likely to accentuate the potential disadvantages of HPR as a control method, including damaging side-effects on natural enemies and biotypic breakdown. The increased tolerance to insecticides that such HPR may cause is discussed on Chapter 23, this volume).

It is more broadly based, partial HPR to aphids that needs to be given greater research emphasis, especially given the worldwide move towards the integration of several control methods represented by IPM.

## **Acknowledgement**

I am very grateful to Professor Stephen Clement for up-dating the information on endophytes,

## **References**

Adams, J.B. and van Emden, H.F. (1972) The biological properties of aphids and their host plant relationships. In: van Emden, H.F. (ed.) *Aphid Technology*. Academic Press, London, pp. 6–104.

Alston, F.H. and Briggs, J.B. (1970) Inheritance of hypersensitivity to rosy apple aphid *Dysaphis plantaginea* in apples. *Canadian Journal of Genetics and Cytology* 12, 257–258.

Anonymous (1994) Die Rückkehr der Reblaus. *Profil, October* 1994, 11.

Ansari, A.K. (1984) Biology of *Aphis craccivora* (Koch) and varietal resistance of cowpeas. PhD thesis, The University of Reading, Reading, UK.

Ansari, A.K., van Emden, H.F. and Singh, S.R (1989) Graft-transmissibility of resistance to cowpea aphid, *Aphis craccivora* Koch (Hemiptera: Aphididae), in six highly antibiotic cowpea varieties. *Bulletin of Entomological Research* 79, 393–399.

Anstead, J.A., Burd, J.D. and Shufran, K.A. (2003) Over-summering and biotypic diversity of *Schizaphis graminum* (Homoptera: Aphididae) populations on noncultivated grass hosts. *Environmental Entomology* 32, 662–667.

Ashouri, A. (2004) Transgenic-Bt potato plant resistance to the Colorado potato beetle affect the aphid parasitoid *Aphidius nigripes*. *Mededelingen Faculteit Landbouwkundige en Toegepaste Biologische Wetenschappen Universiteit Gent* 69, 185–189.

Ashouri, A., Michaud, D. and Cloutier, C. (2001) Unexpected effects of different potato resistance factors to the Colorado potato beetle (Coleoptera: Chrysomelidae) on the potato aphid (Homoptera: Aphididae). *Environmental Entomology* 30, 524–532.

Assad, M.T., Mehrabi, A.M., Pakniyat, H. and Nematollahy, M.R. (2004). The effect of resistance components on reducing yield and its related characters in wheat as infected by *Diuraphis noxia* (Hemiptera: Aphididae). In: *Cereal Research Communications, No. 32*. Cereal Research Non-Profit Company, Zged, Hungary, pp. 69–73.

Atiri, G.I., Ekpo, E.J.A. and Thottappilly, G. (1984) The effect of aphid-resistance in cowpea on infestation and development of *Aphis craccivora* and the transmission of cowpea aphid-borne mosaic virus. *Annals of Applied Biology* 104, 339–346.

Auclair, J.L., Maltais, J.B. and Cartier, J.J. (1957) Factors in resistance of peas to the pea aphid, *Acyrtosiphon pisum* (Harr.) (Homoptera: Aphididae). II. Amino acids. *Canadian Entomologist* 89, 457–464.

Ballman, E.S., Ghising, K., Prischmann-Voldseth, D.A. and Harmon, J.P. (2012) Factors contributing to the poor performance of a soybean aphid parasitoid *Binodoxys communis* (Hymenoptera: Braconidae) on an herbivore resistant soybean cultivar. *Environmental Entomology* 41, 1417–1425.

Bansal, R., Mian, M.A.R. and Michel, A.P. (2013) Identification of novel sources of host plant resistance to known soybean aphid biotypes. *Journal of Economic Entomology* 106, 1479–1485.

Belefantmiller, H., Porter, D.R., Pierce, M.L. and Mort, A.J. (1994) An early indicator of resistance in barley to Russian wheat aphid. *Plant Physiology* 105, 1289–1294.

Bergman, D.K., Dillwith, J.W., Zarrabi, A.A., Caddel, J.L. and Berberet, R.C. (1991) Epicuticular lipids of alfalfa relative to its susceptibility to spotted alfalfa aphids (Homoptera, Aphididae). *Environmental Entomology* 20, 781–785.

Berlandier, F.A. (1996) Alkaloid level in narrow-leafed lupin, *Lupinus angustifolius*, influences green peach aphid reproductive performance. *Entomologia Experimentalis et Applicata* 79, 19–24.

van den Bosch, R. and Messenger, P.S. (1973) *Biological Control*. Intext, New York, 137 pp.

Bouvaine, S., Faure, M.L., Grebenok, R.J., Behmer, S T. and Douglas, A.E. (2014) A dietary test of putative deleterious sterols for the aphid *Myzus persicae*. *PLoS ONE* (on-line) 9(1), e86256.

ten Broeke, C.J.M. (2013) Unravelling the resistance mechanism of lettuce against *Nasonovia ribisnigri*. PhD thesis, Wageningen University, Wageningen, The Netherlands.

Bruce, T.J.A., Aradottir, G.I., Smart, L.E., Martin, J.L., Caulfield, J.C., Doherty, A., Sparks, C.A., Woodcock, C.M., Birkett, M.A., Napier, J.A., Jones, H.D. and Pickett, J.A. (2015) The first

crop plant genetically engineered to release an insect pheromone for defence. *Scientific Reports* (on-line) 5, 11183.

Bultman, T.L. and Bell, G.D. (2003) Interaction between fungal endophytes and environmental stressors influences plant resistance to insects. *Oikos* 103, 182–190.

Burd, J. and Puterka, G. (2009) Plant resistance management strategies for greenbug (*Schizaphis graminum* (Rondani)) (Homoptera: Aphididae) in wheat-sorghum cropping systems. *Redia* 92, 227–228.

Buschman, L.L. and Ramaswamy, S.B. (2012) How to build the non-host plant for stability in insect resistance management. *GM Crops* 3, 163–174.

Bush, L., Slosser, J.E., Worrall, W.D. and Horner, N.V. (1991) Potential of wheat cultivar mixtures for greenbug (Homoptera, Aphididae) management. *Journal of Economic Entomology* 84, 1619–1624.

Cambier, V., Hance, T. and De Hoffmann, E. (2001) Effects of 1,4-benzoxazin-3-one derivatives from maize on survival and fecundity of *Metopolophium dirhodum* (Walker) on artificial diet. *Journal of Chemical Ecology* 27, 359–370.

Castro, A.M., Ramos, S., Vasicek, A., Worland, A., Gimenez, D., Clua, A.A. and Suarez, E. (2001) Identification of wheat chromosomes involved with different types of resistance against greenbug (*Schizaphis graminum*, Rond.) and the Russian wheat aphid (*Diuraphis noxia*, Mordvilko). *Euphytica* 118, 321–330.

Chen, J.Q., Rahbe, Y., Delobel, B., Sauvion, N., Guillaud, J. and Febvay, G. (1997) Melon resistance to the aphid *Aphis gossypii*: behavioural analysis and chemical correlations with nitrogenous compounds. *Entomologia Experimentalis et Applicata* 85, 33–44.

Clement, S.L., Lester, D.G., Wilson, A.D. and Pike, K.S. (1992) Behavior and performance of *Diuraphis noxia* (Homoptera: Aphididae) on fungal endophyte-infected and uninfected perennial ryegrass. *Journal of Economic Entomology* 85, 583–588.

Clement, S.L., Kaiser, W.J. and Eichenseer, H. (1994) *Acremonium* endophytes in germplasms of major grasses and their utilization for insect resistance. In: Bacon, C.W. and White, J.M., Jr. (eds) *Biotechnology of Endophytic Fungi of Grasses*. CRC, Boca Raton, pp. 185–199.

Clement, S.L., Lester, D.G., Wilson, A.D., Johnson, R.C. and Bouton, J.H. (1996) Expression of Russian wheat aphid (Homoptera: Aphididae) resistance in genotypes of tall fescue harbouring different isolates of *Acremonium* endophyte. *Journal of Economic Entomology* 89, 766–770.

Clement, S.L., Wilson, A.D., Lester, D.G. and David, C.M. (1997) Fungal endophytes of wild barley and their effects on *Diuraphis noxia* population development. *Entomologia Experimentalis et Applicata* 82, 275–281.

Clement, S.L., Hu, J., Stewart, A.V., Wang, B. and Elberson, L.R. (2011) Detrimental and neutral effects of a wild grass-fungal endophyte symbiotum on insect preference and performance. *Journal of Insect Science* (on-line) 11, 77.

Cole, R.A. (1994a) Locating a resistance mechanism to the cabbage aphid in two wild brassicas. *Entomologia Experimentalis et Applicata* 71, 23–31.

Cole, R.A. (1994b) Isolation of a chitin-binding lectin, with insecticidal activity in chemically defined synthetic diets, from two wild brassica species with resistance to cabbage aphid *Brevicoryne brassicae*. *Entomologia Experimentalis et Applicata* 72, 181–187.

Cole, R.A. (1997a) The relative importance of glucosinolates and amino acids to the development of two aphid pests *Brevicoryne brassicae* and *Myzus persicae* on wild and cultivated *Brassica* species. *Entomologia Experimentalis et Applicata* 85, 121–133.

Cole, R.A. (1997b) Comparison of feeding behaviour of two *Brassica* pests *Brevicoryne brassicae* and *Myzus persicae* on wild and cultivated *Brassica* species. *Entomologia Experimentalis et Applicata* 85, 135–141.

Cooper, W.C., Jia, L. and Goggin, F.L. (2004). Acquired and R-gene-mediated resistance against the potato aphid in tomato. *Journal of Chemical Ecology* 30, 2527–2542.

- Crawford, K.M., Land, J.M. and Rudgers, J.A. (2010) Fungal endophytes of native grasses decrease insect herbivore preference and performance. *Oecologia* 164, 431–444.
- Czesak, M.E., Knee, M.J., Gale, R.G., Bodach, S.D. and Fritz, R.S. (2004) Genetic architecture of resistance to aphids and mites in a willow hybrid system. *Heredity* 93, 619–626.
- Deol, G.S., Wilde, G.E. and Gill, B.S. (1995) Host plant resistance in some wild wheats to the Russian wheat aphid, *Diuraphis noxia* (Mordvilko) (Homoptera: Aphididae). *Plant Breeding* 114, 545–546.
- Di Pietro, J.P. and Caillaud, C.M. (1998) Response to intracolonial selection for adaptation to resistant wheat in the English grain aphid (Homoptera: Aphididae). *Environmental Entomology* 27, 80–85.
- Di Pietro, J.P., Caillaud, C.M., Chaubet, P., Pierre, J.S. and Trottet, M. (1998) Variation in resistance to the grain aphid, *Sitobion avenae* (Sternorrhynca: Aphididae), among diploid wheat genotypes. *Plant Breeding* 117, 407–412.
- Dreyer, D.L. and Campbell, B.C. (1984) Association of the degree of methylation of intercellular pectin with plant resistance to aphids and with induction of aphid biotypes. *Experientia* 40, 224–226.
- Dreyer, D.L. and Campbell, B.C. (1987) Chemical basis of host-plant resistance to aphids. *Plant, Cell and Environment* 10, 353–361.
- Du Li, Ge Feng, Zhu SanRong and Parajulee, M.N. (2004) Effect of cotton cultivar on development and reproduction of *Aphis gossypii* (Homoptera: Aphididae) and its predator *Propylaea japonica* (Coleoptera: Coccinellidae). *Journal of Economic Entomology* 97, 1278–1283.
- Dunnam, E.W. and Clark, J.C. (1938) The cotton aphid in relation to the pilosity of cotton leaves. *Journal of Economic Entomology* 31, 663–666.

Eichenseer, H., Dahlman, D.L. and Bush, L.P. (1991) Influence of endophyte infection, plant-age and harvest interval on *Rhopalosiphum padi* survival and its relation to quantity of N-formyl and N-acetyl loline in tall fescue. *Entomologia Experimentalis et Applicata* 60, 29–38.

Eigenbrode, S.D., Kabalo, N.N. and Rutledge, C.E. (2000) Potential of reduced-waxbloom oilseed *Brassica* for insect pest resistance. *Journal of Agricultural and Urban Entomology* 17, 53–63.

Eisenbach, J. and Mittler, T.E. (1987) Extra-nuclear inheritance in a sexually produced aphid: the ability to overcome host plant resistance by biotype hybrids of the greenbug, *Schizaphis graminum*. *Experientia* 43, 332–334.

Ellis, P.R., Singh, R., Pink, D.A.C., Lynn, J.R. and Shaw, P.L. (1996) Resistance to *Brevicoryne brassicae* in horticultural brassicas. *Euphytica* 88, 85–96.

van Emden, H.F. (1978) Insects and secondary substances – an alternative viewpoint with special reference to aphids. In: Harborne, J.B. (ed.) *Phytochemical Aspects of Plant and Animal Co-evolution*. Academic Press, London, pp. 309–323.

van Emden, H.F. (1987) Cultural methods: the plant. In: Burn, A.J., Coaker, T.H. and Jepson, P.C. (eds) *Integrated Pest Management*. Academic Press, London, pp. 27–68.

van Emden, H.F. (1995). Host plant-Aphidophaga interactions. *Agriculture, Ecosystems and Environment* 52, 3–11.

van Emden, H.F. (1997) Host-plant resistance to insect pests. In: Pimentel, D. (ed.) *Techniques for Reducing Pesticide Use*. Wiley, Chichester, pp. 130–152.

van Emden, H.F. and Service, M.W. (2004) *Pest and Vector Control*. Cambridge University Press, Cambridge, 349 pp.

van Emden, H.F., Vidyasagar, P. and Kazemi, M.H. (1991) Use of systemic insecticide to measure antixenosis to aphids in plant choice experiments. *Entomologia Experimentalis et Applicata* 58, 69–74.

Foyer, C.H., Noctor, G. and van Emden, H.F. (2007) An evaluation of the costs of making specific secondary metabolites; is the yield penalty incurred by host plant resistance to insects due to competition for resources? *International Journal of Pest Management* 53, 175–182.

Francis, F., Guillonneau, F., Leprince, P., Pauw, E. de, Haubruge, E., Jia, L. and Goggin, F.L. (2010) Tritrophic interactions among *Macrosiphum euphorbiae* aphids, their host plants and endosymbionts: investigation by a proteomic approach. *Journal of Insect Physiology* 56, 575–585.

Fuentes-Contreras, E. and Niemeyer, H.M. (1998) DIMBOA glucoside, a wheat chemical defense, affects host acceptance and suitability of *Sitobion avenae* to the cereal aphid parasitoid *Aphidius rhopalosiphi*. *Journal of Chemical Ecology* 24, 371–381.

Fuentes-Contreras, J.E., Powell, W., Wadhams, L.J., Pickett, J.A. and Niemeyer, H.M. (1996) Influence of wheat and oat cultivars on the development of the cereal aphid parasitoid *Aphidius rhopalosiphi* and the generalist aphid parasitoid *Ephedrus plagiator*. *Annals of Applied Biology* 129, 181–187.

Gershenson, J. (1994) The cost of plant chemical defense against herbivory: a biochemical perspective. In: Bernays, E.A. (ed.) *Insect–Plant Interactions, Volume 5*. CRC, Boca Raton, pp. 173–205.

Gianoli, E., Papp, M. and Niemeyer, H.M. (1996) Costs and benefits of hydroxamic acids-related resistance on winter wheat against the bird cherry–oat aphid, *Rhopalosiphum padi* L. *Annals of Applied Biology* 129, 83–90.

Gibson, R.W. (1976) Glandular hairs are a possible means of limiting aphid damage to the potato crop. *Annals of Applied Biology* 82, 143–146.

Givovich, A. and Niemeyer, H.M. (1995) Comparison of the effect of hydroxamic acids from wheat on five species of cereal aphids. *Entomologia Experimentalis et Applicata* 74, 115–119.

Gomes, F.B., Moraes, J.C. de, Santos, C.D. dos and Goussain, M.M. (2005) Resistance induction in wheat plants by silicon and aphids. *Scientia Agricola* 62, 547–551.

- Gowling, G.R. and van Emden, H.F. (1994) Falling aphids enhance impact of biological control by parasitoids on partially aphid-resistant plant varieties. *Annals of Applied Biology* 125, 233–242.
- Hansen, L.M. (2006). Effect of 6-methoxybenzoxazolin-2-one (MBOA) on the reproductive rate of the grain aphid (*Sitobion avenae* F.). *Journal of Agriculture and Food Chemistry* 54, 1031–1035.
- Havlickova, H. (1997) Differences in level of tolerance to cereal aphids in five winter wheat cultivars. *Rostlinna Vyroba* 43, 593–596.
- Hesler, L.S. (2005) Resistance to *Rhopalosiphum padi* (Homoptera: Aphididae) in three triticales accessions. *Journal of Economic Entomology* 98, 603–610.
- Hesler, L.S. (2013) Resistance to soybean aphid among wild soybean lines under controlled conditions. *Crop Protection* 53, 139–146.
- Hesler, L.S., Chiozza, M.V., O'Neal, M.E., MacIntosh, G.C., Tilmon, K.J., Chandrasena, D.I., Tinsley, N.A., Cianzio, S.R., Costamagna, A.C., Cullen, E.M., DiFonzo, C.D., Potter, B.D., Ragsdale, D.W., Steffey, K. and Koehler, K.J. (2013) Performance and prospects of Rag genes for management of soybean aphid. *Entomologia Experimentalis et Applicata* 147, 201–216.
- Hume, D.E., Ryan, D.L., Cooper, B.M. and Popay, A.J. (2007) Agronomic performance of AR37-infected ryegrass in northern New Zealand. *Proceedings of the New Zealand Grassland Association* 69, 201–205.
- Johnson, A.W., Sisson, V.A., Snook, M.F., Fortnum, B.A. and Jackson, D.M. (2002) Aphid resistance and leaf surface chemistry of sugar ester producing tobaccos. *Journal of Entomological Research* 37, 154–165.
- Kareiva, P. and Sahakian, R. (1990) Tritrophic effects of a simple architectural mutation in pea plants. *Nature, London* 345, 433–434.
- Kazemi, M.H. and van Emden, H.F. (1992) Partial antibiosis to *Rhopalosiphum padi* in wheat and some phytochemical correlations. *Annals of Applied Biology* 121, 1–9.

- Kift, N.B., Dewar, A.M., Werker, A.R. and Dixon, A.F.G. (1996) The effect of plant age and infection with virus yellows on the survival of *Myzus persicae* on sugar beet. *Annals of Applied Biology* 129, 371–378.
- Kishaba, A.N., Castle, S.J., Coudriet, D.L., McCreight, J.D. and Bohn, G.W. (1992) Virus transmission by *Aphis gossypii* Glover to aphid-resistant and susceptible muskmelons. *Journal of the American Society for Horticultural Science* 117, 248–254.
- Kogan, M. and Ortman, E.F. (1978) Antixenosis – a new term proposed to define Painter’s ‘non-preference’ modality of resistance. *Bulletin of the Entomological Society of America* 24, 175–176.
- Kok-Yokomi, M.L. (1978) Mechanisms of host plant resistance in tomato to the potato aphid *Macrosiphum euphorbiae* (Thomas). *Dissertation Abstracts International B* 39, 1121.
- Lehtonen, P., Helander, M. and Saikkonen, K. (2005) Are endophyte-mediated effects on herbivores conditional on soil nutrients? *Oecologia* 142, 38–45.
- Lowe, H.J.B., Murphy, G.J.P. and Parker, M.I. (1985) Non-glaucousness, a probable aphid-resistance character of wheat. *Annals of Applied Biology* 106, 555–560.
- Martos, A., Givovich, A. and Niemeyer, H.M. (1992) Effect of DIMBOA, an aphid resistance factor in wheat, on the aphid predator *Eriopsis connexa* Germar (Coleoptera, Coccinellidae). *Journal of Chemical Ecology* 18, 469–479.
- Maxwell, F.G. and Painter, R.H. (1962) Auxin content of extracts of certain tolerant and susceptible host plants of *Toxoptera graminum*, *Macrosiphum pisi*, and *Therioaphis maculata*, and relation to host plant resistance. *Journal of Economic Entomology* 55, 46–56.
- Mentink, P.J.M., Kimmins, F.M., Harrewijn, P., Dieleman, F.L., Tjallingi, W.F., van Rheenen, B. and Ennink, A.H. (1984) Electrical penetration graphs combined with stylet cutting in the study of host plant resistance to aphids. *Entomologia Experimentalis et Applicata* 35, 210–213.

Merrill, S.C., Radolph, T.L., Peairs, F.B., Michels, G. J. and Walker, C.B. (2014) Examining the competitive advantage of *Diuraphis noxia* (Hemiptera: Aphididae) biotype 2 over biotype 1. *Journal of Economic Entomology* 107, 1471–1475.

Messina, F.J. and Bloxham, A.J. (2004) Plant resistance to the Russian wheat aphid: effects on a nontarget aphid and the role of induction. *Canadian Entomologist* 136, 129–137.

Migui, S.M. and Lamb, R.J. (2004) Seedling and adult plant resistance to *Sitobion avenae* (Hemiptera: Aphididae) in *Triticum monococcum* (Poaceae), an ancestor of wheat. *Bulletin of Entomological Research* 94, 35–46.

Miles, P.W. (1999) Aphid saliva. *Biological Reviews of the Cambridge Philosophical Society* 74, 41–85.

Moate, P.J., Williams, S.R.O, Grainger, C., Hannah, M.C., Mapleson, D., Auldish, M.J., Greenwood, J.S., Popay, A.J., Hume, D.E., Mace, W.J. and Wales, W.J. (2012) Effects of wild-type, AR1 and AR37 endophyte-infected perennial ryegrass on dairy production in Victoria, Australia. *Animal Production Science* 52, 1117–1130.

Morgham, A.T., Richardson, P.E., Campbell, R.K., Burd, J.D., Eikenbary, R.D. and Sumner, L.C. (1994) Ultrastructural responses of resistant and susceptible wheat to infestation by greenbug biotype E (Homoptera: Aphididae). *Annals of the Entomological Society of America* 87, 908–917.

Müller, H.J. (1958) The behaviour of *Aphis fabae* in selecting its host plants, especially different varieties of *Vicia faba*. *Entomologia Experimentalis et Applicata* 1, 66–72.

Murugan, M., Khan, S.A., Cardona, P.S., Orozco, G.V., Viswanathan, P., Reese, J., Starkey, S. and Smith, C.M. (2010) Variation of resistance in barley against biotypes 1 and 2 of the Russian wheat aphid (Hemiptera: Aphididae). *Journal of Economic Entomology* 103, 938–948.

Ni, X.Z., Quisenberry, S.S., Siegfried, B.D. and Lee, K.W. (1998) Influence of cereal leaf epicuticular wax on *Diuraphis noxia* probing behaviour and nymphoposition. *Entomologia Experimentalis et Applicata* 89, 111–118.

[Nicholas, A.](#), Birch, E. and [Leszczynski, B.](#) (2005) Saponins as a source of alfalfa resistance towards pea aphid, *Acyrtosiphon pisum* Harris. *Bulletin OILB/SROP* 28(10), 45–50.

Nicol, D., Armstrong, K.F., Wratten, S.D., Walsh, P.J., Straw, N.A., Cameron, C.M., Lahmann, C. and Frampton, C.M. (1998) Genetic diversity of an introduced pest, the green spruce aphid *Elatobium abietinum* (Hemiptera: Aphididae) in New Zealand and the United Kingdom. *Bulletin of Entomological Research* 88, 537–543.

Novy, R.G., Nasruddin, A., Ragsdale, D.W. and Radcliffe, E.B. (2002) Genetic resistance to potato leafroll virus, potato virus Y, and green peach aphid in progeny of *Solanum tuberosum*. *American Journal of Potato Research* 79, 9–18.

Painter, R.H. (1951) *Insect Resistance in Crop Plants*. Macmillan, New York, 520 pp.

[Pompon, J.](#), [Quiring, D.](#), [Giordanengo, P.](#) and [Pelletier, Y.](#) (2010) Role of host-plant selection in resistance of wild *Solanum* species to *Macrosiphum euphorbiae* and *Myzus persicae*. *Entomologia Experimentalis et Applicata*. 137, 73–85.

Pompon J., Li XiuQuing and Pelletier, Y. (2011) Resistance level to an aphid potato pest varies between genotypes from the same *Solanum* accession. *Journal of Economic Entomology* 104, 1075–1079.

Popay, A.J. (2009) Insect herbivory and defensive mutualisms between plants and fungi. In: White, J.F. Jr and Torres, M.S. (eds) *Defensive Mutualism in Microbial Symbiosis*, CRC Press, Boca Raton, pp. 347–366.

Popay, A.J. and Hume, D.E. (2013) Endophytes for improving ryegrass performance: Current status and future possibilities. *Proceedings of the 22nd International Grassland Congress, Sydney, September 2013*, 1625–1626.

Porter, D.R., Burd, J.D., Shufran, K.A. and Webster, J.A. (2000) Effect of pyramiding greenbug (Homoptera: Aphididae) resistance genes in wheat. *Journal of Economic Entomology* 93, 1315–1318.

Puterka, G.J. and Peters, D.C. (1990) Sexual reproduction and inheritance of virulence in the greenbug, *Schizaphis graminum* (Rondani). In: Campbell, R.K. and Eikenbary, R.D. (eds) *Aphid-Plant Genotype Interactions*. Elsevier, Amsterdam, pp. 289–318.

Puterka, G.J., Burd, J.D. and Burton, R.L. (1992) Biotypic variation in a worldwide collection of Russian wheat aphid (Homoptera: Aphididae). *Journal of Economic Entomology* 85, 1497–1506.

Radcliffe, E.B. and Chapman, R.K. (1965) The relative resistance to insect attack of three cabbage varieties at different stages of plant maturity. *Annals of the Entomological Society of America* 58, 897–902.

Radcliffe, E.B. and Chapman, R.K. (1966) Varietal resistance to insect attack in various cruciferous crops. *Journal of Economic Entomology* 59, 120–125.

Rahbe, Y., Sauvion, N., Febvay, G., Peumans, W.J. and Gatehouse, A.M.R. (1995) Toxicity of lectins and processing of ingested proteins in the pea aphid *Acyrtosiphon pisum*. *Entomologia Experimentalis et Applicata* 76, 143–155.

Rahbe, Y., Deraison, C., Bonade-Bottino, M., Girard, C., Nardon, C. and Jouanin, L. (2003) Effects of the cysteine protease inhibitor oryzacystatin (OC-1) on different aphids and reduced performance of *Myzus persicae* on OC-1 expressing transgenic oilseed rape. *Plant Science* 164, 441–450.

Rana, J.S., Khokhar, K.S. and Harvir Singh (2001) Influence of morphological traits of plants on the infestation by mustard aphid, *Lipaphis erysimi* (Kalt.) in rapeseed and mustard cultivars. *Indian Journal of Plant Protection* 29, 159–160.

Ren, J., Zhou, F., Chen, X., Hu, Q., Yang, A., Zhou, J., Shao, J. and Qian, S. (2014) Effect of physical characteristics of leaf on resistance of cucumber to aphid. *Chinese Journal of Eco-Agriculture* 23, 52–57 [in Chinese, with Abstract in English].

[Ribeiro, A.P.O.](#), [Pereira, E.J.G.](#), [Galvan, T.L.](#), [Picanco, M.C.](#), [Picoli, E.A.T.](#), [Silva, D.J.H. da](#), [Fari, M.G.](#) and [Otoni, W.C.](#) (2006) Effect of eggplant transformed with oryzacystatin gene on *Myzus persicae* and *Macrosiphum euphorbiae*. *Journal of Applied Entomology* 130, 84–90.

- Ridsdill-Smith, J., Edwards, O., Wang ShaoFang, Ghisalberti, E. and Reidy-Crofts, J. (2004) Aphid response to plant defensive compounds in plants. In: *Aphids in a New Millennium. Proceedings of the 6th International Symposium on Aphids, Rennes, September 2001*. INRA, Paris, pp. 491–497.
- Roberts, J.M.F., Hodgson, C.J., Jackai, L.E.N., Thottappilly, G. and Singh, S.R. (1993) Interaction between two synthetic pyrethroids and the spread of two nonpersistent viruses in cowpea. *Annals of Applied Biology* 122, 57–67.
- Robertson, G.W., Griffiths, D.W., Birch, A.N.E., Jones, A.T., McNicol, J.W. and Hall, J.E. (1991) Further evidence that resistance in raspberry to the virus vector aphid, *Amphorophora idaei*, is related to the chemical composition of the leaf surface. *Annals of Applied Biology* 119, 443–449.
- Robinson, J. (1992) Modes of resistance in barley seedlings to six aphid (Homoptera, Aphididae) species. *Journal of Economic Entomology* 85, 2510–2515.
- Ruggle, P. and Gutierrez, A.P. (1995) Use of life-tables to assess host-plant resistance in alfalfa to *Therioaphis trifolii f. maculata* (Homoptera: Aphididae) – hypothesis for maintenance of resistance. *Environmental Entomology* 24, 313–325.
- Saikkonen, K., Faeth, S.H., Helander, M.L. and Sullivan T.J. (1998) Fungal endophytes: a continuum of interactions with host plants. *Annual Review of Ecology and Systematics* 29, 319–343.
- Sarwan Kumar Sangha, M.K. (2013) Biochemical mechanism of resistance in some *Brassica* genotypes against *Lipaphis erysimi* (Kaltenbach) (Homoptera: Aphididae [sic]). *Vegetos* 26, 387–395.
- Sarwar, M. and Sattar, M. (2013). Varietals variability of winter rapeseed (*Brassica napus* L.) for their susceptibility to green aphid, *Myzus persicae* (Sulzer) (Homoptera: Aphididae). *Pakistan Journal of Zoology* 45, 883–888.
- Schardl, C.L., Gossman, R.B., Nagabhyru, P., Faulkner J.R. and Mallik, U.P. (2007) Loline alkaloids: currencies of mutualism. *Phytochemistry* 68, 980–996.

Shepherd, T., Robertson, G.W., Griffiths, D.W. and Birch, A.N.E. (1999a) Epicuticular wax composition in relation to aphid infestation and resistance in red raspberry (*Rubus idaeus* L.). *Phytochemistry* 52, 1239–1254.

Shepherd, T., Robertson, G.W., Griffiths, D.W. and Birch, A.N.E. (1999b) Epicuticular wax ester and triacylglycerol composition in relation to aphid infestation and resistance in red raspberry (*Rubus idaeus* L.). *Phytochemistry* 52, 1255–1267.

Shufran, K.A. and Payton, T.L. (2009) Limited genetic variation within and between Russian wheat aphid (Hemiptera: Aphididae) biotypes in the United States. *Journal of Economic Entomology* 102, 440–445.

Shufran, K.A., Mornhinweg, D.W., Baker, C.A. and Porter, D.R. (2007) Variation to cause host injury between Russian wheat aphid (Homoptera: Aphididae) clones virulent to Dn4 wheat. *Journal of Economic Entomology* 100, 1685–1691.

Siegel, M.R., Latch, G.C.M., Bush, L.P., Fannin, F.F., Rowan, D.D. and Tapper, B.A. (1990) Fungal endophyte-infected grasses: alkaloid accumulation and aphid response. *Journal of Chemical Ecology* 16, 3301–3315.

Simmons, A.T., McGrath, D. and Gurr, G.M. (2005). Trichome characteristics of F1 *Lycopersicon esculentum* × *L. pennellii* hybrids and effects on *Myzus persicae*. *Euphytica* 144, 313–320.

Smith, C.M. and Boyko, E.V. (2007) The molecular bases of plant resistance and defense responses to aphid feeding: current status. *Entomologia Experimentalis et Applicata* 122, 1–16.

Sotherton, N.W. and van Emden, H.F. (1982) Laboratory assessments of the resistance to the aphids, *Sitobion avenae* and *Metopolophium dirhodum* in three *Triticum* species and two modern wheat cultivars. *Annals of Applied Biology* 101, 99–107.

Stoner, K.A. (1992) Density of imported cabbageworms (Lepidoptera, Pieridae), cabbage aphids (Homoptera, Aphididae), and flea beetles (Coleoptera, Chrysomelidae) on glossy and trichome-bearing lines of *Brassica oleracea*. *Journal of Economic Entomology* 85, 1023–1030.

Thomas, S., Dogimont, C. and Boissot, N. (2012) Association between *Aphis gossypii* genotype and phenotype on melon accessions. *Arthropod - Plant Interactions* 6, 93–101.

Thompson, K.F. (1963) Resistance to the cabbage aphid (*Brevicoryne brassicae*) in *Brassica* plants. *Nature, London* 198, 209–210.

Tjallingii, W.F. (1988) Electrical recording of stylet penetration activities. In: Minks, A.K. and Harrewijn, P. (eds) *Aphids. Their Biology, Natural Enemies and Control, Volume 2B*. Elsevier, Amsterdam, pp. 95–108.

Tolmay, V.L. Jankielsohn, A. and Sydenham, S.L. (2013) Resistance evaluation of wheat germplasm containing Dn4 or Dny against Russian wheat aphid biotype RWASA3. *Journal of Applied Entomology* 137, 476–480.

Wartenberg, H. (1953) Über pflanzenphysiologische Ursachen des Massenwechsels der Apfelblattlaus (*Eriosoma lanigerum*) auf *Malus pumila*. *Mitteilungen der Biologischen Zentralanstalt Berlin-Dahlem* 75, 53–56.

Weathersbee, A.A., Hardee, D.D. and Meredith, W.R. (1994) Effects of cotton genotype on seasonal abundance of cotton aphid (Homoptera, Aphididae). *Journal of Agricultural Entomology* 11, 29–37.

Weathersbee, A.A., Hardee, D.D. and Meredith, W.R. (1995) Differences in yield response to cotton aphids (Homoptera, Aphididae) between smooth-leaf and hairy-leaf isogenic cotton lines. *Journal of Economic Entomology* 88, 749–754.

Webb, S.E. and Linda, S.B. (1993) Effect of oil and insecticide on epidemics of potyviruses in watermelon in Florida. *Plant Disease* 77, 869–874.

Webster, J.A., Inayatullah, C., Hamissou, M. and Mirkes, K.A. (1994) Leaf pubescence effects in wheat on yellow sugarcane aphids and greenbugs (Homoptera, Aphididae). *Journal of Economic Entomology* 87, 231–240.

Weibull, J. (1987) Work on plant resistance to *Rhopalosiphum padi* (L.) in oats and barley – present status. *Bulletin OILB/SROP* 10, 160–161.

Weibull, J. (1994) Glutamic acid content of phloem sap is not a good predictor of plant resistance to *Rhopalosiphum padi*. *Phytochemistry* 35, 601–602.

White, C. and Eigenbrode, S.D. (2000) Effects of surface wax variation in *Pisum sativum* on herbivorous and entomophagous insects in the field. *Environmental Entomology* 29, 773–780.

Wilkinson, H.H., Siegel, M.R., Blankenship, J.D., Mallory, A.C., Bush, L.P. and Schardl, C.L. (2000) Contribution of fungal loline alkaloids to protection from aphids in a grass-endophyte mutualism. *Molecular Plant-Microbe Interactions* 13, 1027–1033

Wojcicka, A. (2015) Aktywność mszycy zbożowej (*Sitobion avenae*) oraz czeremchowo-zbożowej (*Rhopalosiphum padi*) w procesie żerowania mszyc na sztucznych pożywkach z ekstraktami wosków powierzchniowych. *Progress in Plant Protection* 55, 14–19.

Wu, C.J., Avila, C.A. and Goggin, F.L. (2015) The ethylene response factor Pti5 contributes to potato aphid resistance in tomato independent of ethylene signalling. *Journal of Experimental Botany* 66, 559–570.

Young, C.A., Hume, D.E. and McCulley, R.L. (2013) Fungal endophytes of tall fescue and perennial ryegrass: pasture friend or foe? *Journal of Animal Science* 91, 2379–2394.

**Table 22.1.** Some examples of wild relatives as sources of genes in traditional breeding for host plant resistance to aphids.

---

<b>‘Wild’ relative</b>	<b>Crop</b>	<b>Resistance character (if known)</b>
	<b>Sample reference(s)</b>	
<i>Avena macrostachya</i> and <i>Avena barbata</i>	Oats Weibull (1987)	
<i>Solanum habrochaites</i> and <i>Lycopersicon pennellii</i>	Tomato Simmons <i>et al.</i> (2005)	

---

<i>Solanum peruvianum</i>	Tomato Kok-Yukom (1978)	Glandular hairs and high tomatine
<i>Solanum tuberosum</i>	Potato Novy <i>et al.</i> (2002)	
<i>Brassica fruticulosa</i> and <i>Brassica spinescens</i>	Cabbage Cole (1994a,b)	Bluconapin; lectin
<i>Triticum monococcum</i>	Wheat Sotherton and van Emden (1982);  Deol <i>et al.</i> (1995); Di Pietro <i>et al.</i> (1998)  Migui and Lamb (2004)	
<i>Triticum monococcum</i> subsp. <i>aegilopoides</i> <i>Triticum urartu</i>	Wheat Di Pietro <i>et al.</i> (1998)	
<i>Solanum berthaultii</i>	Potato Gibson (1976)	Glandular trichomes


---

Note: The same source of resistance may influence several aphid species attacking the same crop; hence, names of aphids affected are not included in the table. However, aphid names are usually given in the reference titles.


### Legends for figures




**Fig. 22.1.** Junction of experimental lettuce plant testing for host plant resistance to *Pemphigus bursarius*. The impact of variety as an aphid control measure is seen clearly in the contrast between the susceptible variety ‘Mildura’ in the foreground and the resistant ‘Avoncrisp’ behind (courtesy of J.A. Dunn).




**Fig 22.2.** Electrical penetration graph (EPG) recordings for *Nasonovia ribisnigri* (currant–lettuce aphid) on susceptible (a) and nearly isogenic resistant (b) lettuce varieties. On susceptible lettuce, beginning at about 170 min, one long continuous phloem phase (*p*) is shown. This starts with sieve element (watery) salivation (waveform *E1*) and continues for the rest of the 4 h recording (only the first 2 h are shown) with sieve element ingestion (waveform *E2*). In contrast, the graph from the resistant lettuce shows only two short phloem-feeding periods (*p*), mainly of *E1* waveform with some very brief switches to *E2*. The average start of the first phloem phase does not differ between the graphs from the two varieties; a phloem factor prevents sieve element ingestion (courtesy of W.F. Tjallingii).




**Fig.22.3.** Mechanisms of host plant resistance to aphids, from left to right approximately in the sequence in which they will affect colonizing aphids. Thickness of arrows indicates the likely outcome of the interaction between mechanism and aphid in terms of plant antixenosis, antibiosis, or tolerance.



**Fig. 22.4.** Electronmicrographs of wax quantity and its distribution on normal (a) and glossy (b) cabbage leaf surfaces (courtesy of S.G. Eigenbrode).



**Fig. 22.5.** Glandular trichomes. (a) Scanning electron micrograph of the leaf surface of *Solanum berthaultii* showing both trichomes with the four-celled head (magnified on insert) and those surmounted by a glandular vesicle. (b) Scanning electron micrograph of an aphid that was on *Solanum polyadenium*, showing the material from the glandular trichomes hardened on the feet (courtesy of R.W. Gibson).



**Fig. 22.6.** The endophytic fungus *Neotyphodium*. (a) Section through a tall fescue (*Lolium arundinaceum*) seed showing mycelium between the aleurone cells. (b) Electronmicrograph of hyphae, conidiophore and conidium (4  $\mu\text{m}$  long) in culture. Host is wild barley (*Hordeum* sp.) (courtesy of S.L. Clement).



**Fig. 22.7.** (a) Normal and (b) 'leafless' cultivars of peas ((b) courtesy of C. Coyne).