

A novel antibacterial peptide derived from Crocodylus siamensis haemoglobin hydrolysate induces membrane permeabilization causing iron dysregulation, oxidative stress and bacterial death

Article

Accepted Version

Lueangsakulthai, J., Jangpromma, N., Temsiripong, T., McKendrick, J. E. ORCID: <https://orcid.org/0000-0003-2275-0569>, Khunkitti, W., Maddocks, S.E. and Klaynongsruang, S. (2017) A novel antibacterial peptide derived from *Crocodylus siamensis* haemoglobin hydrolysate induces membrane permeabilization causing iron dysregulation, oxidative stress and bacterial death. *Journal of Applied Microbiology*, 123 (4). pp. 819-831. ISSN 1364-5072 doi: <https://doi.org/10.1111/jam.13539> Available at <https://centaur.reading.ac.uk/72721/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1111/jam.13539>

To link to this article DOI: <http://dx.doi.org/10.1111/jam.13539>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **A novel antibacterial peptide derived from *Crocodylus siamensis* hemoglobin hydrolysate induces membrane**
2 **permeabilisation causing iron dysregulation, oxidative stress and bacterial death**

3

4 J. Lueangsakulthai^{1,2}, N. Jangpromma^{1,3}, T. Temsiripong⁴, J.E. McKendrick⁵, W. Khunkitti⁶, S.E. Maddocks⁷ and S.
5 Klaynongsruang^{1,2}

6

7 ¹Protein and Proteomics Research Center for Commercial and Industrial Purposes (ProCCI)

8 ²Department of Biochemistry, Faculty of Science, Khon Kaen University, Khon Kaen, 40002, Thailand

9 ³Office of the Dean, Faculty of Science, Khon Kaen University, Khon Kaen, 40002, Thailand

10 ⁴Srirachamoda Co., Ltd. 383 Moo 4, Nongkham, Sriracha, Chonburi, 20230, Thailand

11 ⁵Department of Chemistry, The University of Reading, Reading, RG6 6UR, United Kingdom

12 ⁶Department of Pharmaceutical Technology, Faculty of Pharmaceutical Science, Khon Kaen University, Khon
13 Kaen, 40002, Thailand

14 ⁷Department of Biomedical Sciences, Cardiff School of Health Science, Cardiff Metropolitan University, Cardiff,
15 CF5 2YB, United Kingdom

16

17 **Correspondence**

18 Sarah E. Maddocks, Department of Biomedical Sciences, Cardiff School of Health Science, Cardiff Metropolitan
19 University, Cardiff, CF5 2YB, United Kingdom.

20 Email: smaddocks@cardiffmet.ac.uk

21 Sompong Klaynongsruang, Department of Biochemistry, Faculty of Science, Khon Kaen University, Khon Kaen,
22 40002, Thailand.

23 Email: somkly@kku.ac.th

24

25 **Abstract**

26 **Aims:** A novel antibacterial peptide from *Crocodylus siamensis* hemoglobin hydrolysate (CHHs) was characterised
27 for antimicrobial activity.

28 **Methods and Results:** CHHs was hydrolysed for 2 h (2h-CHH), 4 h (4h-CHH), 6 h (6h-CHH) and 8 h (8h-CHH).
29 8h-CHH showed antibacterial activity against *Escherichia coli*, *Staphylococcus aureus*, *Klebsiella pneumonia* and
30 *Pseudomonas aeruginosa* at concentrations of 20, 20, 20 and 10 mg ml⁻¹ (w/v), respectively. Fluorescent
31 microscopy revealed that 8h-CHH had bactericidal activity against *E. coli* and *P. aeruginosa*. β -galactosidase assay
32 supported by RT-qPCR demonstrated that 8h-CHH resulted in differential expression of genes involved in iron
33 homeostasis (*ftnA* and *bfd*) and oxidative stress (*sodA*, *soxR* and *oxyR*). Siderophore assay indicated that 8h-CHH
34 also impaired siderophore production with diminished expression of *pvdF*. This pattern of gene expression suggests
35 that 8h-CHH triggers the release of free ferric ions in the cytoplasm. However, decreased expression of genes
36 associated with the SOS response (*recA* and *lexA*) in combination with neutral comet revealed that no DNA damage
37 was caused by 8h-CHH. Membrane permeabilisation assay indicated that 8h-CHH caused membrane leakage
38 thought to mediate the antibacterial and iron-stress responses observed, due to loss of regulated iron transport. The
39 novel active peptide from 8h-CHH was determined as QAIHNEKVQAHGKKVL (QL17), with 41 %
40 hydrophobicity and +2 net charge.

41 **Conclusions:** The QAIHNEKVQAHGKKVL fragment of *Crocodylus siamensis* hemoglobin is antibacterial via a
42 mechanism that likely relies on iron dysregulation and oxidative stress which results in bacterial death.

43 **Significance and Impact of the Study:** We have described for the first time, a novel peptide derived from
44 *Crocodylus siamensis* hemoglobin hydrolysate, that has the potential to be developed as a novel antimicrobial
45 peptide.

46

47 **Keywords**

48 *Crocodylus siamensis*, hemoglobin hydrolysate, antibacterial, peptide, oxidative stress genes, iron homeostasis
49 genes.

50

51 **Running headline**

52 Antibacterial peptide from *C. siamensis* hemoglobin

53

54 **Introduction**

55 Bacterial infections account for a significant proportion of the global infectious disease burden, and morbidity and
56 mortality rates caused by infectious microbial agents pose serious public health concerns. This is exacerbated by
57 increasing resistance to antibiotics which is significant in an era where the development of new, synthetic
58 antibacterial drugs lags the emergence of antimicrobial resistance (Mbah *et al.* 2012). It is therefore paramount to
59 broaden the search for new antimicrobial substances, including the exploitation of novel sources where possible
60 (Song *et al.* 2012). A growing area of research has begun to focus on protein hydrolysates of animal origin, such as
61 goat whey protein hydrolyzed by treatment with alcalase, and which shows broad spectrum antibacterial activity
62 (Osman *et al.* 2016). Similarly, an antibacterial peptide derived from acid extract of chicken (*Gallus gallus*) blood
63 has efficacy against *E. coli* by a mechanism that results in toroidal pore formation and subsequent bacterial lysis
64 (Vasilchenko *et al.* 2016). Furthermore, a peptic hemoglobin hydrolysate from bovine hemoglobin has also been
65 shown to possess antibacterial activity (Froidevaux *et al.* 2001; Daoud *et al.* 2005; Arroume *et al.* 2006; Arroume *et*
66 *al.* 2008; Adje *et al.* 2011). Whilst these areas of research are largely in their infancy, they provide a diverse, yet
67 novel, means of informing the development of new peptide-based antimicrobial treatments.

68 *Crocodylus siamensis* is a small, freshwater crocodilian. In the wild these crocodiles experience many
69 traumatic wounds that might be expected to be rife with infection from endogenous environmental bacteria, but this
70 is not seen to be the case. *C. siamensis* (Siamese crocodile) hemoglobin constitutes the most abundant component in
71 crocodile blood and has been long associated with a broad spectrum of biological activity, including antimicrobial
72 (Srihongthong *et al.* 2012; Pakdeesuwan *et al.* 2017), antioxidant (Jandaruang *et al.* 2012; Srihongthong *et al.* 2012;
73 Phosri *et al.* 2014; Maijaroen *et al.* 2016; Pakdeesuwan *et al.* 2017; Phosri *et al.* 2017) and anti-inflammatory
74 activity (Phosri *et al.* 2014; Jangpromma *et al.* 2017; Phosri *et al.* 2017). Antibacterial activity has been attributed to
75 peptides derived from *C. siamensis* hemoglobin and they are currently thought to be targeted to the bacterial surface
76 (Srihongthong *et al.* 2012; Pakdeesuwan *et al.* 2017). However detailed studies of antimicrobial activity, mechanism
77 of action and critical peptide sequences that mediate the observed activity are currently lacking.

78 This study aimed to investigate the antibacterial activity of hemoglobin hydrolysate from *C. siamensis*
79 blood (CHH), hydrolysed for 2, 4, 6 and 8 h with pepsin, and to identify specific antimicrobial peptide fragments.
80 Minimum Inhibitory Concentration (MIC), Time-Killing kinetics and viability staining revealed both bactericidal
81 and bacteriostatic activity of the crude, hydrolysed peptide cocktail. To establish a mechanism of activity, the effect
82 of CHHs on bacterial iron homeostasis was investigated using *ftnA* and *bfd* linked reporter strains, and analysis of

83 siderophore production. Due to the close association of iron homeostasis and oxidative stress, the expression levels
84 of oxidative stress response genes (*oxyR*, *sodA* and *soxR*) were also investigated but ruled out the inclusion of the
85 SOS response genes or associated DNA damage as a mechanism of action. However, membrane permeability assay
86 indicated the hemoglobin fragments disrupted the cell envelope; the active peptide was found to be positively
87 charged and therefore likely functioned in a similar manner to cationic antimicrobial peptides.

88

89 **Materials and methods**

90 **Bacterial strains**

91 *Escherichia coli* (NCTC 10418), *Pseudomonas aeruginosa* (PAO1), *Staphylococcus aureus* (NCTC 13141),
92 *Klebsiella pneumonia* (ATCC 13883), *Bacillus subtilis* were maintained on nutrient agar (NA) or nutrient broth
93 (NB), aerobically at 37°C throughout the study, unless otherwise stated. *E. coli* MC4100, *E. coli* MC4100 *fnA-lacZ*,
94 *E. coli* MC4100 *bfd-lacZ*, *E. coli* H1914 Δ *fur-ftnA-lacZ*, *E. coli* H1914 Δ *fur-bfd-lacZ* carried stable chromosomal
95 mutations that allowed them to be maintained on NB or NA as described above; these strains were provided by Prof.
96 Simon Andrews (University of Reading).

97

98 **Hemoglobin extraction**

99 The extraction of hemoglobin from red blood cells (RBCs) was performed following the method of Srihongthong *et*
100 *al.* (2012). The RBCs were washed three times with phosphate buffered saline (PBS) pH 7.0 and centrifuged at 3000
101 $\times g$ for 5 min at 4°C. Ice-cold distilled water with five-fold volume was added to the RBCs pellet, vigorously mixed
102 and allowed to settle for 10 min. After centrifugation at 10000 $\times g$ for 20 min at 4°C the supernatant was collected,
103 lyophilized and stored at -70°C.

104

105 **Enzymatic hydrolysis**

106 Enzymatic hydrolysis was performed according to the method of Yu *et al.* (2006). Briefly, the hemoglobin solution
107 was digested with pepsin (at pH 2.0) using a ratio of enzyme to substrate (1:100 w/w) at 37°C for 2, 4, 6 and 8 h and
108 boiled at 95°C for 10 min to inactivate the enzyme. The insoluble material was removed by centrifugation at 7168 \times
109 g for 20 min. The supernatant was collected and adjusted to pH 7.0 by addition of 1 M HCl or 1 M NaOH. Finally
110 the supernatant (hydrolysate) was lyophilized and stored at -20°C.

111

112 **Minimum inhibitory concentration (MIC) and time –killing assay (TKA)**

113 The minimum inhibitory concentration (MIC) of CHHs was determined by microbroth dilution in a microtiter plate
114 assay system using a total volume of 110 μ l per well. Each well contained 10 μ l bacterial suspension (*E. coli* NCTC
115 10418, *S. aureus*, *K. pneumoniae* and *P. aeruginosa*; adjusted to OD 0.1 A₆₅₀) and 100 μ l of CHHs. Plates were
116 incubated for 24 h (MIC assay) or 9 h (TKA) at 37°C. An end point reading was taken for MIC, and hourly readings
117 were taken for TKA (A₆₅₀) (Spectrostar^{Nano}, BMG Labtech).

118

119 **Fluorescent microscopy using BacLight™ to assess bacterial viability**

120 Test microorganisms (*E. coli* NCTC 10418, *S. aureus*, *K. pneumoniae*, *B. subtilis* and *P. aeruginosa*) were incubated
121 for 16 h with 10, or 20 mg ml⁻¹ CHHs (determined from MIC). Cultures were centrifuged at 9000 \times g for 5 min and
122 the supernatant was discarded. The cell pellets were re-suspended in 100 μ l dH₂O containing SYTO9 and PI at a
123 ratio of 2:1. Cells were incubated in the dark at room temperature for 45 min and 10 μ l transferred to a glass slide.
124 Cells were visualized by fluorescent microscopy (Nikon eclipse 80i) using oil immersion and \times 100 lens. SYTO9
125 detection (viable cells) was used a 488 nm excitation and 520 nm emission filter. Propidium iodine (PI) detection
126 (non-viable cells) was used 543 nm excitation and 572 nm emission filter.

127

128 **β -galactosidase assay**

129 Test microorganisms (*E. coli* MC4100, *E. coli* MC4100 *ftnA-lacZ*, *E. coli* MC4100 *bfd-lacZ*, *E. coli* H1914 Δ *fur*-
130 *ftnA-lacZ*, *E. coli* H1914 Δ *fur-bfd-lacZ*) were cultured in 100 μ l NB in a 96-well microtiter plate until mid-log was
131 reached. Then 100 μ l of 2, 4, 6 and 8h-CHH (20 mg ml⁻¹) was added to each well followed by an additional
132 incubation for 1.5 h at 37°C. Bacteria were permeabilised with buffer containing 60 mM Na₂HPO₄·7 H₂O, 40 mM
133 NaH₂PO₄·H₂O, 10 mM KCl, 1 mM MgSO₄·7H₂O, 50 mM β -mercaptoethanol, 0.1 % SDS and 4 μ l of chloroform
134 for 5 min. Bacterial lysate was transferred to a fresh MTP and 20 μ l of ONPG was added to each sample. The plate
135 was incubated in the dark at 37°C for 10 min, and absorbance values were read at 420 nm and 550 nm
136 (Spectrostar^{Nano}, BMG Labtech). Miller units were calculated based on the following formula (Miller 1972):

137
$$\text{Miller Units} = 1,000 \times [\text{OD}_{420} - (1.75 \times \text{OD}_{550})] / [T \times V \times \text{OD}_{600}]$$

138 where T determines the reaction time in minute and V determines the volume of cultured assayed in ml.

139

140 **CAS agarose diffusion (CASD) assay**

141 CAS agarose diffusion assay followed the method of Schwyn and Neilands (1987). Pre-cultures of (*E. coli* MC4100,
142 *E. coli* NCTC 10418, *S. aureus*, *K. pneumoniae*, *B. subtilis* and *P. aeruginosa*) were equilibrated to OD 0.1 (A_{650})
143 and supplemented with 20 mg ml⁻¹ of 2, 4, 6 and 8h-CHH and 1 mM DTPA. Treated cultures were incubated for a
144 further 24 h at 37°C. After centrifuging at 9000 × *g* for 3 min (Rotina 380R centrifuge; Hettich, Germany) 50 μl
145 aliquots of the sample supernatant were added to wells bored into the center of Petri dishes filled with 25 ml CAS
146 agarose agar. After incubating in the dark at room temperature for 2 h, digital calipers were used to measure the
147 diameter (mm) of the yellow diffusion zone diameters which is an indicator of siderophore production and the
148 reduction of ferric iron.

149

150 **RNA extraction and cDNA synthesis**

151 Total RNA was extracted from bacterial cells per the manufacturer's instructions using the SV Total RNA extraction
152 kit (Promega, USA). Total RNA concentration and purity were determined using Nanodrop ND-1000
153 spectrophotometer (Nanodrop Technologies, USA). RNA samples with an absorbance ratio at OD 260/280 between
154 1.8-2.2 and OD 260/230 < 2.0 were used for further analysis. For each sample, cDNA was synthesized using the
155 High-Capacity cDNA Reverse Transcription kit according to the manufacturer's instructions (Applied Biosystem
156 Inc).

157

158 **Reverse transcription - real time PCR (RT-qPCR) assay**

159 Primers used in this study are shown in Table 1. All PCR reactions were performed using 7500 Fast Real-Time PCR
160 System machine under following conditions: 15 min at 95°C and 40 cycles of 3 s at 95°C, 30 s at 68°C in 96-well
161 clear optical reaction plates (Applied Biosystem, USA). The procedure ended by melt-curve ramping from 60 to
162 95°C for 20 min to check the PCR specificity. All RT-qPCR reactions were carried out in biological and technical
163 triplicate. A non-template control was also included in each run for each gene.

164

165 **Neutral comet assay**

166 Neutral comet assay was determined following the method of Solanky and Haydel (2012) with some modifications.
167 Frosted microscope glass slides with a clear window were pre-coated by dipping in 1% agarose solution prepared
168 with sterile water and were dried in an incubator at 40°C for 30 min. *E. coli* MC4100 cultures were incubated with
169 rotary agitation at 37°C until logarithmic phase of growth (OD ~0.1; A₆₀₀), and then diluted with NB broth to
170 achieve a concentration of ~10⁷ CFU ml⁻¹. The cells were collected by centrifugation (9000 × g, 15 min), washed
171 with 0.1× phosphate-buffered saline (PBS), and re-suspended in one of the following solutions: 1% TritonX-100; 5
172 mM H₂O₂; deionized H₂O and 8h-CHH 20 mg ml⁻¹. Exposed cells were then incubated for 30 min at 37°C.
173 After incubating the slide at 4°C for 10 min to allow the initial layer of agarose to cool, the coverslip was removed
174 and a subsequent agarose layer was added. The first layer consisted of 200 μl of 0.5% agarose prepared in 0.1× PBS
175 and maintained at 55 –60°C for 30 min. For the second layer, 2 μl of exposed cells was mixed thoroughly with 200
176 μl of 0.5% agarose solution and 100 μl of this mixture was transferred to the slide. A third layer was consisted of 5
177 μg ml⁻¹ RNaseA and 1 mg ml⁻¹ lysozyme in 0.5% agarose solution. Slides were refrigerated for 10 min at 4°C and
178 incubated for 30 min at 37°C. Embedded cells were then lysed by immersing slides in a solution containing 2.5 M
179 NaCl, 100 mM EDTA, 10 mM Tris pH 10 and 1% TritonX-100 for 1 h at room temperature. Following lysis, slides
180 were immersed in an enzyme digestion solution prepared with 2.5 M NaCl, 10 mM EDTA, 10 mM Tris pH 7.4, and
181 0.5 mg ml⁻¹ of proteinase K for 2 h at 37°C. Lysis and enzyme digestion steps were carried out in the dark to prevent
182 light exposure. Slides were immersed in buffer containing 300 mM sodium acetate and 100 mM Tris, pH 9 for 20
183 min. Slides were electrophoresed at 12 V for 50 min. Following electrophoresis, slides were sequentially immersed
184 in 1 M ammonium acetate prepared in ethanol for 20 min and 75% ethanol for another 30 min. Slides were then
185 allowed to dry. Prior to staining, slides were pretreated with a freshly prepared solution of 5% DMSO and 10 mM
186 NaH₂PO₄. While the slides were still wet, DNA was stained with 50 μl of 1 μM YOYO-1 in 5% DMSO and
187 visualized using a Nikon eclipse 80i fluorescent microscope at 100× magnification with the appropriate filter set for
188 YOYO-1 (excitation 491 nm and emission 509 nm). Comets were imaged and comet lengths were measured using
189 Volocity software version 5.5.

190

191 **Cytoplasmic membrane permeability assay**

192 Cytoplasmic membrane permeability assay was determined following the method of Chitemerere and
193 Mukanganyama (2014) with modifications. Briefly, bacteria (*E. coli* MC4100, *P. aeruginosa*, *S. aureus*) were grown

194 to mid-exponential phase (OD 0.2-0.4; A₆₅₀); 2 ml were mixed with an equivalent volume of 1 μM diSC3-5 dyes
195 and incubated 1 h, for maximal uptake of dye, then were collected by centrifugation (3000 × g for 5 min). Cells were
196 wash and re-suspended in 2 ml of buffer (5 mM HEPES, pH 7.2, 5 mM glucose) and the absorbance was measured
197 at an excitation wavelength of 622 nm and emission wavelength of 670 nm. Afterward, 2 ml of 100 mM KCl was
198 added to equilibrate the cytoplasmic and external K⁺ ion concentrations. Cells were mixed with and equal volume of
199 sample then the fluorescence was monitored at an excitation wavelength of 622 nm and emission wavelength of 670
200 nm at 1 h intervals. Dye released with de-ionised water was used as a negative control.

201

202 **Amino acid sequence analysis**

203 The active fraction of 8h-CHH was selected and the contained peptides identified using LTQ Orbitrap XL Mass
204 spectrometry employing the following search parameters: non-specified enzymatic cleavage with three possible
205 missed cleavages, +/-0.8 Da mass tolerances for MS and MS/MS, a peptide mass tolerance of +/-5 ppm,
206 methionine oxidation and Gln->pyro-Glu (N-term Q) variable modification and monoisotopic mass. Data were
207 additionally processed at the Mascot Server (<http://www.matrixscience.com/>) using MS/MS ion searches against
208 SwissProt (current release).

209

210 **Statistical analysis**

211 Statistical analysis was performed using ANOVA and followed by Dunnett's test. Data are presented as mean ±
212 SEM. A value of $P < 0.05$ was accepted to be significant ($*P < 0.05$, $**P < 0.01$, $***P < 0.001$).

213

214 **Results**

215 **Bacteriostatic and bactericidal effects of hemoglobin hydrolysate**

216 Bacteriostatic and bactericidal effects of hemoglobin hydrolysate were investigated by minimum inhibitory
217 concentration (MIC), time-killing assay (TKA) and fluorescent microscopy using BacLight™. The MIC was
218 determined using 2h-CHH, 4h-CHH, 6h-CHH and 8h-CHH, and results are presented in Table 2. Gram-negative
219 microorganisms generally required higher concentrations of hydrolysed CHHs to inhibit growth. The results showed
220 that all hydrolysed CHHs were antibacterial at concentrations of 20 mg ml⁻¹ (w/v) against *E. coli* and *S. aureus*.
221 Furthermore, there was antibacterial activity for all CHHs at concentrations of 10 mg ml⁻¹ (w/v) against *K.*

222 *pneumoniae* and *P. aeruginosa*. Significantly, the dose required to inhibit growth decreased as the length of time
223 that the hemoglobin was hydrolysed increased, suggesting that higher concentrations of shorter peptide fragments
224 mediated the inhibitory activity. This was verified by TKA which indicated that 8h-CHH resulted in the highest
225 percentage of bacterial death (Table 3). Samples analysed by BacLight™ viability staining and fluorescent
226 microscopy confirmed that 8h-CHH at a dose of 20 and 10 mg ml⁻¹ caused bacterial death (*E. coli* NCTC 10418 and
227 *P. aeruginosa*; Fig. 1a and 1b, respectively). At lower doses and shorter hydrolyze times, CHH treatments tended to
228 be bacteriostatic rather than bactericidal. Both mechanisms were investigated as described below.

229 230 **Hemoglobin hydrolysate alters bacterial iron homeostasis and causes oxidative stress without inducing** 231 **irreversible DNA-damage or the SOS response**

232 The β -galactosidase assay demonstrated that 2h-CHH, 4h-CHH, 6h-CHH and 8h-CHH at 20 mg ml⁻¹ decreased the
233 expression of *ftnA* while increasing the expression of *bfd*, under iron rich conditions (Fig. 2). Decreased expression
234 of *ftnA* is indicative of iron restriction whereas increased expression of *bfd* is affiliated with iron repletion and is
235 known to have a role in haem-iron handling. The expression profile was not altered in a Δfur background suggesting
236 that the dysregulated expression of *ftnA* and *bfd* was not the result of Fur-dependent iron-mediated regulation (Fig.
237 2). However, CASD assay indicated that CHHs decreased the production of siderophores (Fig. 3) and following
238 treatment with 8h-CHH, the expression of *pvdF* (associated with siderophore synthesis) also significantly decreased
239 ($P < 0.05$) (Fig. 4b) suggesting some degree of altered iron homeostasis.

240 Bacterial iron metabolism and oxidative stress are inextricably linked; RT-qPCR demonstrated that
241 following treatment with 8h-CHH, the expression of oxidative stress response genes *sodA* and *soxR* significantly
242 increased while the expression of *oxyR* decreased (Fig. 4a). Oxidative stress is correlated with DNA damage and
243 subsequent induction of the SOS system of repair. However, the expression of *recA* was diminished and *lexA*
244 unchanged following treatment with 8h-CHH suggesting that the SOS response was not initiated (Fig. 4c).
245 Therefore, the dysregulated iron homeostasis observed might be mediated by transcriptional responses associated
246 with oxidative stress, rather than in response to environmental iron availability.

247 248 **Hemoglobin hydrolysate does not cause DNA damage**

249 Neutral Comet Assay verified that CHH did not cause DNA damage. Treatment with 1% TritonX-100 and 5 mM
250 H₂O₂ (positive controls) produced an increase in comet length, and therefore DNA degradation, relative to the
251 negative control (DI sterilized water). Exposure of *E. coli* MC4100 to 1% TritonX-100 and 5 mM H₂O₂ for 30 min
252 increased the comet length values of 49.79 μ m and 21.12 μ m, respectively. Meanwhile the exposure of 8h-CHH
253 resulted in comet length values of 11.24 μ m and the negative control, 10.78 μ m as shown in Table 4. The 1%
254 TritonX-100 also yielded higher a comet length value, demonstrating higher DNA double strand break (DSB) levels
255 while DNA double strand break wasn't caused by 8h-CHH upon exposure in bacterial cells.

256

257 **Hemoglobin hydrolysate causes bacterial membrane leakage**

258 Many short peptides aggregate in bacterial membranes resulting in a loss of integrity, such loss of integrity allows
259 influx and efflux of various solutes and ions, including metal ions such as iron. It was hypothesized that membrane
260 disruption might underpin the iron homeostatic dysregulation and concurrent oxidative stress response described
261 above. Indeed, results from cytoplasmic membrane permeability assay, showed a gradual increase of diSC3-5 dye
262 release over time in the presence of 8h-CHH as well as in the presence of the 1% TritonX-100 positive control (Fig.
263 5a, 5b and 5c). Similarly, diSC3-5 was released from all test microorganisms following exposure to 8h-CHH for 60
264 min (Fig. 5). The effect was most marked for the Gram-negative test microorganisms (*E. coli* and *P. aeruginosa*)
265 indicating that 8h-CHH can permeabilise the cell envelope more effectively than that of Gram-positive
266 microorganisms (*S. aureus*).

267

268 **Active peptide was identified by LTQ Orbitrap XL mass spectrometry**

269 To establish the identity of the active CHH peptide, the primary sequence was determined using LTQ Orbitrap XL
270 mass spectrometry. As the most active CHH, the amino acid sequences of 8h-CHH was determined as
271 QAIHNEKVQAHGKKVL (QL17) corresponding to a molecular mass of 1895.07 Da. The obtained sequence was
272 uploaded to the antimicrobial database (<http://aps.unmc.edu/AP/main.php>) and the protein databank
273 (<https://www.ncbi.nlm.nih.gov>) for further characterization. As shown in Table 5, antibacterial peptide QL 17 had
274 hydrophobicity values of approximately 41%, with a net charge of +2. Alignment of the amino acid sequences of the
275 peptide fragments with *C. siamensis* hemoglobin indicated the antibacterial peptide originated from the β -subunit of

276 *C. siamensis* hemoglobin. The short length and positive charge of the peptide indicates that it might aggregate in the
277 bacterial cell envelope in much the same way as cationic antimicrobial peptides.

278

279 **Discussion**

280 Protein hydrolysates are gaining popularity for their potential therapeutic effects due to demonstrable efficacy and
281 low toxicity. This study used MIC, time-killing kinetics and viability staining to ascribe antimicrobial activity to
282 hemoglobin hydrolysate from *C. siamensis*. CHHs were either bacteriostatic or bactericidal depending on the species
283 of bacteria, dose and length of hydrolyzed time. Longer hydrolysis times correlated with higher inhibition of growth,
284 with 8h-CHH proving to be the most efficacious. The antibacterial mechanism was investigated with focus on iron
285 homeostasis and oxidative stress which in numerous pathogens is associated with host haem-iron availability.

286 The expression of *ftnA* was increased and that of both *bfd* and *pvdF* decreased following exposure to 8h-
287 CHH. Under iron repleted conditions expression of *ftnA* and *bfd* is ordinarily repressed via a process relying on the
288 ferric uptake regulator (Fur). The observed expression profile was no concordant with typical iron-Fur regulation
289 which was verified using a Δfur background, which showed the same expression profile as the wild-type
290 background. However, differential expression of *ftnA*, *bfd* and *pvdF* also occurs in response to oxidative stress,
291 which is closely allied to iron homeostasis (Zheng *et al.* 1999). Quantitative analysis of the expression of *soxR* and
292 *oxyR* indicated differential expression in response to CHH-treatment akin to that observed during oxidative stress,
293 indicating that CHH induced a state of oxidative stress in the microorganisms tested.

294 Uncontrolled oxidative stress results in widespread lipid, protein and DNA damage (Gault *et al.* 2016). If
295 DNA damage becomes too great, the bacterial SOS system is activated. *LexA* is the master regulator of the SOS
296 response; under normal conditions it represses the expression of genes encoding several DNA repair proteins,
297 including *recA*. *LexA* has a negative auto-regulatory function and when the SOS response is triggered, increased
298 levels of *lexA* are also produced which ultimately serves as a negative feedback mechanism to switch of the SOS
299 response once rescue is achieved (Michel 2005). In this study, the expression of *lexA* was unchanged and *recA*
300 decreased following treatment with CHH indicating that while oxidative stress occurred it was below the threshold
301 to necessary to induce the SOS system. This was verified by the absence of DNA damage (a key signal for SOS
302 induction) observed by neutral comet assay.

303 The bacterial response to outer membrane stress is distinct to the oxidative stress and SOS response, but
304 triggers several stress-associated, damage repair pathways to maintain membrane integrity and prevent influx/efflux
305 between the cytoplasm and external environment. Cationic antimicrobial peptides are well known to disrupt the
306 bacterial cell envelope by aggregating to form pores within the membrane. Ordinarily this causes catastrophic
307 damage resulting in bacterial death by lysis. Analysis of membrane integrity following treatment with CHH
308 indicated that 8h-CHH could permeabilise *E. coli*, *P. aeruginosa* and *S. aureus* (Chitemerere and Mukanganyama
309 2014). The active peptide from 8h-CHH was identified using LTQ-Orbitrap XL mass spectrometry and the sequence
310 determined as QAIHNEKVQAHGKKVL (QL17) corresponding to a molecular mass of 1895.07 Da. The peptide
311 QL17 had hydrophobicity values of about 41%, together with a net charge of +2. Given its positive charge, this
312 fragment is hypothesized to aggregate within the bacterial membrane, in the same way as cationic antimicrobial
313 peptides (Sato and Feix 2006). It is known that antimicrobial peptides positive charge combined with
314 hydrophobicity is critical for partitioning of the peptide into the bacterial cell membrane (Pata *et al.* 2011).

315 Alignment the amino acid sequences of the peptide fragment with *C. siamensis* hemoglobin indicated the
316 antibacterial peptide originated from the β -subunit of *C. siamensis* hemoglobin. This result agrees with Arroume *et*
317 *al.* (2008), who reported that antimicrobial products derived from hemoglobin hydrolysis are cleaved from the α -
318 and β -subunits. These products mostly consist of ~15–30 amino acid residues and have a molecular weight < 10
319 kDa. Of note, was the correlation between length of hydrolysis time and antimicrobial activity. As expected,
320 hemoglobin hydrolysed for longer periods of time comprised a higher concentration of small peptide fragments,
321 which in turn exhibited better antimicrobial efficacy, highlighting the importance of peptide fragment size for
322 activity.

323 Taken collectively, the data derived from this study indicates that the hemoglobin of *C. siamensis* can be
324 hydrolyzed to produce a novel antimicrobial peptide that at high concentrations mediates bacterial death by
325 aggregating in the cell envelope and damaging membrane integrity. The consequent influx of exogenous material
326 combined with an efflux of cytoplasmic content likely underpins the dysregulation of iron homeostasis and
327 concurrent oxidative stress. Whilst the potential to utilise antimicrobial peptides derived from hemoglobin warrants
328 further study, blood derived from *C. siamensis* is currently a waste-product of farming. Therefore, with larger
329 studies drawn from this exploratory research, it could instead be developed as a natural or synthetic antimicrobial
330 peptide, engineered to ensure maximum efficacy and minimal toxicity.

331

332 **Acknowledgements**

333 This research was financial supported by the Royal Golden Jubilee Ph.D. program (RGJ-PHD program), Grant No.
334 PhD 0258/2552 and the Newton Fund (British Council). We also grateful to the Protein and Proteomics Research
335 Center for Commercial and Industrial Purposes (ProCCI), Department of Biochemistry, Faculty of Science, Khon
336 Kaen University, Department of Pharmaceutical Technology, Faculty of Pharmaceutical Science, Khon Kaen
337 University, Department of chemistry, The University of Reading, United Kingdom and Department of Biomedical
338 Science, School of Health Science, Cardiff Metropolitan University, United Kingdom for essential support by
339 laboratory facilities. Additionally, we would like to thank Srirachamoda Co., Ltd., Chon Buri, Thailand for
340 supplying crocodile blood samples. We also thank Prof. Simon Andrews (University of Reading) for providing *E.*
341 *coli* strains.

342

343 **Conflict of Interest**

344 The authors confirm that this article content has no conflicts of interest.

345

346 **References**

- 347 Adje, E.Y., Balti, R., Kouach, M., Dhulster, P., Guillochon, D. and Arroume, N.N. (2011) Obtaining antimicrobial
348 peptides by controlled peptic hydrolysis of bovine hemoglobin. *Int J Biol Macromol* **49**, 143–153.
- 349 Ali, R., Lutfullah, G., Khan, A.A. and Rashid, M.I. (2013) Analysis of oxygen affinity in aquatic amphibian;
350 homology modelling of the major Haemoglobin component HbA1 from the African clawed frog (*Xenopus*
351 *laevis*, Anura). *Int J Bioinform Res Appl* **9**, 449–461.
- 352 Arroume, N.N., Delval, V.D., Miloudi, K., Daoud, R., Krier, F., Kouach, M., Briand, G. and Guillochon, D. (2006)
353 Isolation and characterization of four antibacterial peptides from bovine hemoglobin. *Peptides* **27**, 2082–
354 2089.
- 355 Arroume, N.N., Delval, V.D., Adje, E.Y., Traisnel, J., Krier, F., Mary, P., Kouach, M., Briand, G. and Guillochon,
356 D. (2008) Bovine hemoglobin: An attractive source of antibacterial peptides. *Peptides* **29**, 969–977.
- 357 Chitemerere, T.A. and Mukanganyama, S. (2014) Evaluation of cell membrane integrity as a potential antimicrobial
358 target for plant products. *BMC Complement Altern Med* **14**, 278.

359 Daoud, R., Dubois, V., Dodita, L.B., Arroume, N.N., Krier, F., Chihib, N.E., Mary, P., Kouach, M., Briand, G. and
360 Guillochon, D. (2005) New antibacterial peptide derived from bovine hemoglobin. *Peptides* **26**, 713–719.

361 Froidevaux, R., Kriera, F., Arroume, N.N., Marko, D.V., Kosciarz, E., Ruckebusch, C., Dhulster P. and Guillochon
362 D. (2001) Antibacterial activity of a pepsin-derived bovine hemoglobin fragment. *FEBS Letters* **491**, 159–
363 163.

364 Gault, M., Effantin, G. and Rodrigue, A. (2016) Ni exposure impacts the pool of free Fe and modifies DNA
365 supercoiling via metal induced oxidative stress in *Escherichia coli* K-12. *Free Radic Biol Med* **97**, 351–
366 361.

367 Jandaruang, J., Siritapetawee, J., Thumanu, K., Songsirithigul, C., Krittanai, C., Daduang, S., Dhiravisit, A. and
368 Thammasirirak, S. (2012) The effects of temperature and pH on secondary structure and antioxidant
369 activity of *Crocodylus siamensis* hemoglobin. *Protein J* **31**, 43–50.

370 Jangpromma, N., Poolperm, N., Pornsri, K., Anwised, P., Kabbua, T., Phosri, S., Daduang, S. and Klaynongsruang,
371 S. (2017) Proteomics profiling and inflammatory factor gene expression in LPS-stimulated RAW 264.7
372 cells treated with *Crocodylus siamensis* hemoglobin. *Chiang Mai J Sci* **44**, 1–16.

373 Maijaroen, S., Anwised, P., Klaynongsruang, S., Daduang, S. and Boonmee, A. (2016) Comparison of recombinant
374 α -hemoglobin from *Crocodylus siamensis* expressed in different cloning vectors and their biological
375 properties. *Protein Expr Purif* **118**, 55–63.

376 Mbah, J.A., Ngemenya, M.N., Abawah, A.L., Babiaka, S.B., Nubed, L.N., Nyongbela, K.D., Lemuh, N.D. and
377 Efange, S.M.N. (2012) Bioassay-guided discovery of antibacterial agents: *in vitro* screening of *Peperomia*
378 *vulcanica*, *Peperomia fernandopoioana* and *Scleria striatinux*. *Ann Clin Microbiol Antimicrob* **11**, 10.

379 Michel, B. (2005) After 30 years of study, the bacterial SOS response still surprises us. *PLoS Biol* **3**, 1174–1176.

380 Miller, J.H. Experiments in molecular genetics. Cold Spring Harbor Laboratory, NY; 1972. pp. 352–355.

381 Osman, A., Goda, H.A., Hamid, M.A., Badran, S.M. and Otte, J. (2016) Antibacterial peptides generated by alcalase
382 hydrolysis of goat whey. *LWT - Food Sci Technol* **65**, 480–486.

383 Pakdeesuan, A., Araki, T., Daduang, S., Payoungkiattikun, W., Jangpromma, N. and Klaynongsruang, S. (2017)
384 *In Vivo* wound healing activity of crocodile (*Crocodylus siamensis*) hemoglobin and evaluation of
385 antibacterial and antioxidant properties of hemoglobin and hemoglobin hydrolysate. *J Microbiol Biotechnol*
386 **27**, 26–35.

387 Pata, S., Yaraksa, N., Daduang, S., Temsiripong, Y., Svasti, J., Araki, T. and Thammasiriraka, S. (2011)
388 Characterization of the novel antibacterial peptide Leucrocicin from crocodile (*Crocodylus siamensis*) white
389 blood cell extracts. *Dev Comp Immunol* **35**, 545–553.

390 Phosri, S., Mahakunakorn, P., Lueangsakulthai, J., Jangpromma, N., Swatsitang, P., Daduang, S., Dhiravisit, A. and
391 Thammasirirak, S. (2014) An investigation of antioxidant and anti-inflammatory activities from blood
392 components of crocodile (*Crocodylus siamensis*). *Protein J* **33**, 484–492.

393 Phosri, S., Jangpromma, N., Patramanon, R., Kongyingyoes, B., Mahakunakorn, P. and Klaynongsruang, S. (2017).
394 Protective effect of crocodile hemoglobin and whole blood against hydrogen peroxide-induced oxidative
395 damage in Human Lung Fibroblasts (MRC-5) and inflammation in mice. *Inflammation* **40**, 205–220.

396 Sato, H. and Feix, J.B. (2006) Peptide–membrane interactions and mechanisms of membrane destruction by
397 amphipathic α -helical antimicrobial peptides. *Biochimica et Biophysica Acta* **1758**, 1245–1256.

398 Schwyn, B. and Neilands, J.B. (1987) Universal chemical assay for the detection and determination of siderophores.
399 *Anal Biochem* **160**, 47–56.

400 Solanky, D. and Haydel, S.E. (2012) Adaptation of the neutral bacterial comet assay to assess antimicrobial-
401 mediated DNA double-strand breaks in *Escherichia coli*. *J Microbiol Methods* **91**, 257–261.

402 Song, R., Wei, R.B., Luo, H.Y. and Wang, D.F. (2012) Isolation and characterization of an antibacterial peptide
403 fraction from the pepsin hydrolysate of Half-Fin Anchovy (*Setipinna taty*). *Molecules* **17**, 2980–2991.

404 Srihongthong, S., Pakdeesuwan, A., Daduang, S., Araki, T., Dhiravisit, A. and Thammasirirak, S. (2012) Complete
405 amino acid sequence of globin chains and biological activity of fragmented crocodile hemoglobin
406 (*Crocodylus siamensis*). *Protein J* **31**, 466–476.

407 Vasilchenko, A.S., Rogozhin, E.A., Vasilchenko, A.V., Kartashova, O.L. and Sycheva, M.V. (2016) Novel
408 haemoglobin-derived antimicrobial peptides from chicken (*Gallus gallus*) blood: purification, structural
409 aspects and biological activity. *J Appl Microbiol* **121**, 1546–1557.

410 Yu, Y., Hu, J., Miyaguchi, Y., Bai, X., Du, Y. and Lin, B. (2006) Isolation and characterization of angiotensin I-
411 converting enzyme inhibitory peptides derived from porcine hemoglobin. *Peptides* **27**, 2950–2956.

412 Zheng, M., Doan, B., Schneider, T.D. and Storz, G. (1999) OxyR and SoxRS regulation of fur. *J Bacteriol* **181**,
413 4639–4643.

414

415 **Table 1** Primer pairs for RT-qPCR

416

Species	Gene names	GeneBank accession no.	Region	Product size (bp)	Forward primer (5'→3')	Reverse primer (5'→3')
<i>E. coli</i>	<i>oxyR</i>	HG738867	2620691..2621608	197	CCCCGGCTTCAAAACAGAAA	GCTGGTGAAAGAGAGCGAAG
	<i>sodA</i>		2678668..2679288	187	CGAAGTCACGTTTCGATAGCC	CTGCCAGAATTTGCCAACCT
	<i>soxR</i>		4163425..4163889	172	CAGCGGCGATATAAACGTGA	CCAACTCTTCTCGCCATTGG
	<i>rec A</i>		3960012..3961073	173	GAAGAACAAAATCGTGTGCGC	CATTTCGCTTTACCTTGACCG
	<i>lex A</i>		4143071..4143679	158	GCAGGAAGAGGAAGAAGGGT	CTTTCATCGACATCCCCTG
<i>P. aeruginosa</i>	<i>rpoD</i>	BA000007	3952578..3954419	220	TTCGTACGCAAGAACGTCTG	AGGCCGGTTTCTTCTTCAAT
	<i>pvdF</i>	NC_002516	2652230..2653057	181	CGTACCAGCTCATCGAGGAT	AGACCCTGAACGACCTCTTG
	<i>rpoD</i>		634371..636224	186	GGGTCACATCGAACTGCTTG	TCATCGAGGACTCCACCATG

424

425 **Table 2** The MIC values of sample CHHs

426

Sample	MIC values (mg ml ⁻¹)				% Growth inhibition			
	<i>E.coli</i>	<i>K.pneumoniae</i>	<i>P.aeruginosa</i>	<i>S.aureus</i>	<i>E.coli</i>	<i>K.pneumoniae</i>	<i>P.aeruginosa</i>	<i>S.aureus</i>
2h-CHH	20	20	10	20	33.33***±0.003	57.14***±0.015	34.11***±0.004	38.87***±0.005
4h-CHH	20	10	10	10	21.74***±0.025	34.29***±0.002	25.10***±0.002	21.11***±0.031
6h-CHH	20	10	10	20	37.86***±0.007	45.14***±0.008	35.03***±0.008	25.53***±0.020
8h-CHH	20	20	10	20	24.64***±0.003	41.10***±0.113	44.61***±0.020	31.20***±0.195

431

432 *** denotes $P < 0.001$. Data expressed as a mean ± SEM of 3 independent experiments. Significance was measured using ANOVA followed by Dunnett's test.

433

434

435 **Table 3** Time-killing assay (TKA) of sample CHHs

436

437

438

439

440

441

442

443

Sample	%Growth inhibition at 9 h intervals			
	<i>E.coli</i>	<i>K.pneumoniae</i>	<i>P.aeruginosa</i>	<i>S.aureus</i>
2h-CHH	48.07***±0.001	3.21***±0.011	6.07***±0.004	31.23***±0.006
4h-CHH	55.64***±0.007	4.06***±0.002	4.98***±0.007	56.31***±0.007
6h-CHH	73.37***±0.001	0.06±0.002	10.65***±0.011	33.23***±0.013
8h-CHH	48.54***±0.020	12.00***±0.015	21.73***±0.004	17.88***±0.115

444 *** denotes $P < 0.001$. Data expressed as a mean ± SEM of 3 independent experiments. Significance was measured using ANOVA followed by Dunnett's test.

445

446 **Table 4** Neutral bacterial comet assay following *E.coli* exposure to 1 % TritonX-100, 5 mM H₂O₂, 8h-CHH and DI water.

447

448

449

450

451

Comet length values (μm)	<i>E.coli</i> treated with			
	1% (v/v) TritonX-100	5 mM H ₂ O ₂	8h-CHH	DI water
	49.79***±0.002	21.12***±0.010	11.24±0.014	10.78±0.009

452 *** denotes $P < 0.001$. Data expressed as a mean ± SEM of 3 independent experiments. Significance was measured using ANOVA followed by Dunnett's test.

453

454 **Table 5** Structural characteristics of the antibacterial peptide from 8h-CHH

Property	Peptide sequence	Hydrolysate	%Hydrophobicity	Net charge	Sequence alignment
Antibacterial	QAIHHNEKVQAHGKKVL (QL17)	8h-CHH	41%	+2	<i>C. siamensis</i> Hb β-subunit (position 53-69)

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477 **Figure captions**

478 **Figure 1** Representative images of Live and Dead staining assay (BacLight™ Fluorescent microscopy), red coloured
479 cells are dead cells and green coloured cells are live cells, showed the bactericidal effect of 8h-CHH at a dose of 20
480 and 10 mg ml⁻¹ against (a) *E.coli* and (b) *P.aeruginosa* PAO1, respectively.

481
482 **Figure 2** β- galactosidase assay. Expression of *ftnA* and *bfd* in response to treatment with CHHs does not exhibit
483 iron-dependent expression in a wild-type or Δfur background. Each bar represents the mean \pm SEM (n=3). ***
484 significant at $P < 0.001$.

485
486 **Figure 3** The effect of CHHs on siderophore production by CASD assay. Sample CHHs decreased the production of
487 siderophore from all of bacterial strains over a time frame of 2 h. Each bar represents the mean \pm SEM (n=3). ***
488 significant at $P < 0.001$.

489
490 **Figure 4** Effect of 8h-CHH on gene expression levels in *E.coli* and *P.aeruginosa* were analyzed by real time PCR
491 (RT-qPCR). (a) A fold change gene expression of *oxyR*, *sodA* and *soxR* were validated for oxidative stress response.
492 (b) A fold change gene expression of *pvdF* was validated for siderophore production and (c) a fold change gene
493 expression of *recA* and *lexA* were validated for DNA damage (SOS response). Each bar represents the mean \pm SEM
494 (n=3). *significant at $P < 0.05$, **significant at $P < 0.01$ and *** significant at $P < 0.001$.

495
496 **Figure 5** Measurement of 3'3 dipropylthiadicarbocyanine (diSC3-5) dye release overtime from (a) *E.coli*, (b)
497 *P.aeruginosa* PAO1 and (c) *S.aureus* membranes in the presence of 8h-CHH, permeabilizing agent 1% TritonX-100
498 and negative control Deionize water (DI).

499

500

501

502

503

504