

Evaluating soil extraction methods for chemical characterization of ultramafic soils in Kinabalu Park (Malaysia)

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

van der Ent, A., Nkrumah, P. N., Tibbett, M. ORCID: <https://orcid.org/0000-0003-0143-2190> and Echevarria, G. (2019) Evaluating soil extraction methods for chemical characterization of ultramafic soils in Kinabalu Park (Malaysia). *Journal of Geochemical Exploration*, 196. pp. 235-246. ISSN 0375-6742 doi: 10.1016/j.gexplo.2018.10.004 Available at <https://centaur.reading.ac.uk/80181/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.gexplo.2018.10.004>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Accepted Manuscript

Evaluating soil extraction methods for chemical characterization of ultramafic soils in Kinabalu Park (Malaysia)

Antony van der Ent, Philip Nti Nkrumah, Mark Tibbett, Guillaume Echevarria



PII: S0375-6742(18)30010-4
DOI: doi:[10.1016/j.gexplo.2018.10.004](https://doi.org/10.1016/j.gexplo.2018.10.004)
Reference: GEXPLO 6206

To appear in: *Journal of Geochemical Exploration*

Received date: 1 February 2018
Revised date: 14 August 2018
Accepted date: 15 October 2018

Please cite this article as: Antony van der Ent, Philip Nti Nkrumah, Mark Tibbett, Guillaume Echevarria , Evaluating soil extraction methods for chemical characterization of ultramafic soils in Kinabalu Park (Malaysia). *Gexplo* (2018), doi:[10.1016/j.gexplo.2018.10.004](https://doi.org/10.1016/j.gexplo.2018.10.004)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Evaluating soil extraction methods for chemical
characterization of ultramafic soils in Kinabalu Park
(Malaysia)**

Antony van der Ent^{1,2*}, Philip Nti Nkrumah¹, Mark Tibbett³ and Guillaume Echevarria^{1,2}

¹Centre for Mined Land Rehabilitation, Sustainable Minerals Institute, The University of Queensland, Australia.

²Université de Lorraine, INRA, Laboratoire Sols et Environnement, 54000 Nancy, France.

³Centre for Agri-Environmental Research & Soil Research Centre, School of Agriculture, Policy and Development, University of Reading, Reading, RG6 6AR, UK.

*Corresponding author: a.vanderent@uq.edu.au, Centre for Mined Land Rehabilitation, The University of Queensland, St Lucia QLD 4072, Australia.

ABSTRACT

Soils derived from ultramafic bedrock are known for hosting distinct vegetation types as a consequence of atypical soil chemistries consisting of high trace elements concentrations (Ni, Cr, Co) and exchangeable cation imbalances (high Mg:Ca quotients). Ecological studies use a range of single-stage extraction methods for chemical characterization of such soils in order to be able to interpret plant response, and ultimately to explain plant community composition. Few studies to date have compared different soil extraction methods in relation to tropical ultramafic soils. This study compares eight commonly used extraction methods on a large number of ultramafic soil samples collected from Kinabalu Park (Malaysia). The tested methods were: for trace elements: NH_4AC , DTPA, CaCl_2 , $\text{Sr}(\text{NO}_3)_2$ and Mehlich-3, for exchangeable cations: NH_4Ac and silverthioarea, and for plant-available phosphorus: Mehlich-3 and Olsen-P. These single-stage extraction methods were compared and evaluated for predictive power for chemically characterizing soils, interrelatedness and ecological application. The methods were also contrasted with a sequential extraction scheme. Finally, several operational parameters including molar ratio (0.01 and 0.1 M CaCl_2 , $\text{Sr}(\text{NO}_3)_2$) and pH buffering (DTPA-TEA) were also evaluated. The majority of single-stage extraction methods are highly inter-correlated and predictive power could be improved by including independent soil parameters (pH, CEC, pseudo-total element concentration) in the multivariate regression equation. Ecological interpretation remains difficult because of lack of experimental studies in relation to plant uptake response and potential phytotoxicity effects on tropical native plants from ultramafic soils.

Keywords: *carboxylic acid, DTPA, multivariate regression, single-stage extraction, sequential extraction scheme, strontium nitrate.*

INTRODUCTION

Ultramafic rocks are widespread on earth, particularly in tropical countries (Cuba, New Caledonia, Indonesia, Philippines and Malaysia). These rocks are parts of the upper mantle and consist largely of magnesium-iron-silicate minerals. Soils derived from such bedrock are relatively high in the trace elements nickel (Ni), cobalt (Co) and chromium (Cr), but concomitantly have cation imbalances as a result of high magnesium (Mg) but low calcium (Ca) (Proctor et al., 1981; Echevarria, 2018). This atypical soil chemistry has caused the occurrence of distinct vegetation types characterized by relatively low stature and high levels of endemism (Brooks, 1987; Proctor, 2003; van der Ent et al., 2018). The main soil-edaphic factors that are most often cited to be important in relation to the ecology of ultramafic soils are the (potential) phytotoxicity induced by Ni, and possibly by Co and Cr, and nutrient deficiency as a result of low Ca (and high Mg) and very low potassium (K) and phosphorus (P) (Vlams and Jenny, 1948; Walker, 1954; Proctor et al., 1981; Brooks, 1987; Proctor, 2003; Brady et al., 2005). Ecological studies have used single-stage soil extraction methods for: (i) Quantifying potential phytotoxic trace elements mainly Ni; (ii) Demonstrating (high) Mg:Ca quotients in the CEC; and (iii) Estimating potentially plant-available P (Vlams and Jenny, 1948; Proctor et al., 1981). Together these extractions aim to characterize the chemical properties of ultramafic soils under investigation with the ultimate objective of linking such information to ecological attributes, such as vegetation stunting, species-diversity per unit area or functional traits of individual species. However, none of the commonly used methods has been specifically developed for ultramafic soils, or for the use with regards to ecological parameters. Rather, most methods were originally developed for agricultural soils in relation to (trace element) nutrition and deficiency, or phytotoxicity risk assessments. Few studies have compared different methods in use in ecological studies on (tropical) ultramafic soils and evaluated the usefulness of such methods for the characterization of ultramafic soils.

Chemical extraction methods for estimating potential soil trace element phytotoxicity

Potentially phytoavailable trace element soil fractions can be estimated with a range of different methods, including: (a) Single-stage chemical extractants; (b) Sequential extractants schemes; (c) Ion exchange resin methods (IER); (d) Isotopic exchange kinetics (IEK); and (e) Diffusive Gradients in Thin-films (DGT) (Echevarria et al., 1998; Robinson et al., 1999). None of these methods, however, can exactly simulate trace elements availability to plants. Single-stage chemical extractants can be grouped in: (i) neutral salt-based extracts; (ii) chelator-based extracts; (iii) acid-based extracts; or (iv) synthetic root exudate-based extracts.

Important parameters for all extraction methods are the molar concentration of the extract, the liquid to solid ratio, pH, and the equilibration time (Meers et al., 2007).

Neutral salt-based extracts include CaCl_2 , $\text{Ca}(\text{NO}_3)_2$, KCl , NaNO_3 , and NH_4Ac in various molar concentrations. Initially, the aim for using such solutions was to assess the pool of exchangeable cations that are sorbed onto the Cation Exchange Capacity (CEC) (Ciesielski and Sterckeman, 1997). The ion replacement power in such extract solutions (with identical anions, such as NO_3^-) decreases according to the Z number of the element in the order: $\text{Ba}^{2+} > \text{Sr}^{2+} > \text{Ca}^{2+} > \text{Mg}^{2+} > \text{NH}_4^+ > \text{K}^+ > \text{Na}^+$ (Mengel and Kirkby, 2001). The specific chemical adsorption of divalent cations by the soil phases is mainly related to metal ion hydrolysis and increases with increasing pH with pK values of the metal ion hydrolysis, for example 9.9 (Ni) and 9.7 (Co) (Brümmer et al., 1986). However, ion exchange in soils is also indirectly influenced by soil pH because of the competition with H^+ ions for sorption onto the CEC (Tiller et al., 1984). Neutral salt extracts based on monovalent and divalent cations are essentially pH dependent (Anderson and Christensen, 1988; Echevarria et al., 2006), and have been reported to perform better than methods using chelators or methods that significantly acidify or buffer the soil, altering the pH at which the extraction actually occurs (Menzie et al., 2007). Frequently used are 0.01 or 0.1 M CaCl_2 solutions; and at the ionic strength of 0.01 M the extractant solution has a similar ionic strength to that of most soil solutions (Novozamsky et al., 1993; Houba et al., 2000). However, although in most 'normal' soils Ca^{2+} is the dominant cation, in ultramafic soils Mg^{2+} is generally the dominant cation, and hence an extractant based on Mg^{2+} might be more appropriate. Because Cl^- is a complexing anion to some metals as opposed to the NO_3^- counter ion (Gommy et al., 1998), and because the pH can be shifted by high Ca^{2+} levels, 0.1 M $\text{Sr}(\text{NO}_3)_2$ (Wang et al. 2003) and 0.01 M $\text{Sr}(\text{NO}_3)_2$ (Kukier and Chaney, 2004; Kukier et al., 2004) have also been proposed. In both the CaCl_2 and $\text{Sr}(\text{NO}_3)_2$ extracts, the divalent cations also promote coagulation of the colloidal material in the suspension, making higher molar concentrations that are required for monovalent cations (Na^+ , K^+ , NH_4^+) unnecessary (Meers et al., 2007). The 0.01 M $\text{Sr}(\text{NO}_3)_2$ extraction has proven to be effective in predicting extractable Ni and plant Ni uptake across a range of soils (Siebielec et al., 2007). In many studies pertaining ultramafic soils, 1 M NH_4Ac (pH 7.0) has been used for phytoavailable trace elements, and concomitantly for exchangeable cations (Proctor et al., 2000; Robinson et al., 2003; Brearley, 2005). Compared to divalent cations, NH_4^+ is less competitive for desorption of metals from the solid soil phases, but performs similar to K^+ on account of its nearly identical ion radius

(Gryschko et al., 2005; Gommy et al., 1998). The extractant is commonly buffered at pH 7 to prevent carbonate dissolution, but due to the increasing formation of NH_3 from NH_4^+ by dissociation with rising pH, metal ammine complexes can form (Gryschko et al., 2005). This might be important (but has not been studied) in the case of ultramafic soils, with high concentrations of Ni^{2+} , and the use of NH_4Ac has strong effect on soil extract pH and might also result in the formation of soluble Ni-hexammine complexes, which can potentially result in overestimation of exchangeable soil Ni. Also, in acidified ultramafic soils, buffering the extraction solution at pH 7 (2–3 units higher than soil pH) can generate CEC of Fe-oxides and change the retention of metal cations by these minerals, which dominate the soil composition (Becquer et al., 2001).

Chelator-based extractants based on synthetic amino-polycarboxylic acids include EDTA and DTPA. The DTPA (Diethylene triamine pentaacetic acid) method was originally developed to diagnose deficiency of micronutrients in soils (Lindsay and Norvell, 1978), but has been widely used for studies with ultramafic soils (L'Huillier and Edighoffer, 1996; Echevarria et al., 1998, 2006; Lazarus et al., 2011; Chardot-Jacques et al., 2013; Ünver et al., 2013). The DTPA-extract is made up of 0.005 M DTPA with 0.01 M CaCl_2 and is buffered at pH 7.3 with 0.01 M triethanolamine (TEA). The extraction of trace elements in this extract is promoted by the chelation action of DTPA and the Ca^{2+} exchange with other cations, as well as Cl^- complexation (Lindsay and Norvell, 1978; Hsiao et al., 2009). The buffer (TEA) was designed to prevent carbonate dissolution (Lindsay and Norvell, 1978), but carbonates are not likely to be important in ultramafic soils with pH 4–5.8, and Becquer et al. (1995) proposed unbuffered (excluding TEA) DTPA adjusted to pH 5.3. It is, however, important to keep the Ca:DTPA ratio intact because Ca:DTPA binding is necessary to control the exchangeability and chelation of DTPA (Lindsay and Norvell, 1978). The DTPA method was developed for soil deficient in trace elements, whereas ultramafic soils have a surplus of Ni, hence the DTPA-method risks over-saturation and the soil-extractant ratio needs to be adjusted accordingly (Kukier and Chaney, 2001; Li et al., 2000). Although some studies found (weak) correlation between soil Ni-DTPA and uptake in non-accumulating plants (L'Huillier and Edighoffer, 1996), such correlations are mostly restricted to comparing similar soils with a narrow pH range (Sukkariyah et al., 2005), and when applied to a variety of soils, poor prediction of phytoavailability has been reported (Menzies et al., 2007). This can be explained by the rather high chelation stability constants of Ni-DTPA and Co-DTPA at $\log K > 20.2$ and 19.3 respectively (Anderegg et al., 2005), which is unlikely representative for the chelation

capacities of carboxylic acids (for example citrate-Ni log K 5.4) in the plant rhizosphere. However, DTPA-extractable Ni can, in some cases, be strongly correlated to Ni uptake by hyperaccumulators in a limited range of pH (*i.e.* 4.2 to 5.6) in temperate ultramafic soils (Chardot et al., 2007), whereas it is not the case in *circum*-neutral Mediterranean ultramafic soils (Bani et al., 2009). However, in all these soils, the Ni-DTPA appears to be a useful extractant of the exact isotopically-exchangeable pools from which all plants take up Ni (Massoura et al., 2004; Chardot et al., 2007; Estrade et al., 2015): *i.e.* high-activity clays and hydrous Fe oxides (Massoura et al., 2006; Chardot et al., 2007; Bani et al., 2009, 2014).

Acid-based extracts include a digest, normally microwave-pressure-aided or in a hot block with mineral acids such as HCl and HNO₃. These extracts are used to provide 'pseudo-total' levels by dissolution of oxides, hydroxides, carbonates, organic matter and, if HF is also added to mix, to provide 'near-total' values by also breaking down silicate matrices. Such a digest gives a measure for virtually all trace elements present in the soil. In more dilute form, HNO₃ (0.1 M) can also be used to leach metals from the soils, and this has the benefit that the NO₃⁻ counter ion is not complexing. Alternatively (very) dilute organic acids (such as acetic, citric, formic, lactic and malic acid) can be used to mimic plant root exudates as these acids are the most abundant Low Molecular Weight Organic Acids (LMWOA) present in the rhizosphere of many plants (Meers et al., 2007). Wang et al. (2003) and Feng et al. (2005) proposed 10 mM LMWOA extraction solutions, consisting of formic acid, citric acid and malic acids, and reported good correlations with plant metal uptake. The acidity of these organic acids dissolves hydroxides and carbonates, and the citrate and malate counter ions complex Ni and other trace elements. However, the evidence for dilute solutions of carboxylic acids being suitable extractants for metals remains weak.

Selective sequential extractions (SSE) provide operationally defined solid-phase fractionations of metals over soil pools (Quantin et al., 2002). Various schemes have been proposed based on the (much simpler) BCR-protocol (Quevauviller et al., 1994), but for trace elements (Ni, Co, Cr) in ultramafic soils, the most reported is a 7-step programme that consists of the following stages: (1) Water soluble; (2) Exchangeable; (3) bound to Mn oxides; (4) Bound to amorphous Fe oxides; (5) Bound to crystalline Fe oxides; (6) Bound to organic matter; (7) and Residual (Quantin et al., 2002). Of these fractions, the water soluble and exchangeable are immediately phytoavailable, the fraction bound to Mn oxides, amorphous Fe oxides and bound to organic matter are potentially phytoavailable, whereas the

residual fraction is not phytoavailable. Together these labile and non-labile pools determine the mobility and phytoavailability of Ni, Cr and Co in ultramafic soils. Some of these labile fractions extracted individually (and not through a complete sequential procedure) from a wide range of ultramafic soils have been proved to be strongly correlated to isotopically-exchangeable Ni pools, which is the principal source of Ni available to plants in soils: *e.g.* amorphous Fe oxides (Massoura et al., 2004, 2006).

In this study, ultramafic soil samples were collected from a 700-km² area encompassing Kinabalu Park in Malaysia. These soil samples were analysed with the aims of characterizing soil chemical properties, in particular those of ecological relevance to plants, such as the cation exchange complex, availability of macro-nutrients, and concentrations of extractable trace elements. To that end, we compare and evaluate the results of a range of extraction methods commonly used in the field of tropical ecology with a focus on predictive power for characterizing soils, interrelatedness and ecological application.

MATERIALS AND METHODS

Study area and sample collection

Mount Kinabalu Park is located in Sabah (Malaysia) on the island of Borneo (6°5' N and 160°33' E), covering an area of 754 km² including two mountains: Kinabalu (4095 m) and Tambuyukon (2579 m). Although Mount Kinabalu itself is a granite pluton (Cottam et al., 2010), the lower slopes are covered with sedimentary rocks. Ultramafic rock outcrops appear like a collar around the massif on mid-elevation, and also outcrops on Mount Tambuyukon. In total, ultramafic outcrops cover 142 km² within the Park boundaries (Collenette, 1964). Kinabalu Park is covered with intact rainforest and has a humid tropical climate with a mean monthly air temperature of 20°C throughout the year at 1680 m, with a daily fluctuation of 7–9°C (Kitayama, 1991).

During 2010–2014, a large ecological research project was conducted in Kinabalu Park, and in the nearby Bidu-Bidu Hills and Trus Madi Forest Reserves, all in the Malaysian state of Sabah. The project was wide-ranging, but focused on the plant-soil relationships of the vegetation on ultramafic soils at these localities. Data from these soil samples has been previously reported (Van der Ent et al., 2015a,b; 2016a,b; 2018) and we refer to these publications for full details on the sample collection. Briefly, the soil samples were collected

from 14 different ultramafic localities (474–2950 m asl) in Kinabalu Park, including in total 95 discrete sample sites. At each site, 3 soil samples (1–2 kg) were collected in the A/B mineral horizon, and care was taken not to include organic surface layers. All soil samples were packed, brought to the local field station, air-dried at room temperature to constant weight (2–3 weeks), sieved to <2 mm, shipped to Australia, and gamma irradiated at Steritech Pty. Ltd. in Brisbane following Australian Quarantine Regulations.

Laboratory analyses

The chemical analysis of the soil samples took place in the laboratory of the Centre for Mined Land Rehabilitation (CMLR) at The University of Queensland in Australia. The soil samples (0.3 g) were digested using freshly prepared Aqua Regia (4 mL 70% nitric acid and 3 mL 37% hydrochloric acid per sample) in a digestion microwave for a 1-hour programme and diluted to 45 mL before analysis (Rayment and Higginson, 1992; method 17B1). Soil pH and electrical conductivity (EC) was obtained in a 1:2.5 soil:water mixture after 1-hour equilibrium time on an end-over-end shaker and 1-hour settling time. Plant-available phosphorus as Olsen-P (Olsen et al., 1954) was extracted with 1.0 g soil extracted with 20 mL 0.5 M NaHCO₃ (pH 8.5) for 30 minutes (Rayment and Higginson, 1992; method 9C1). Plant-available phosphorus ('ML-3') was also extracted with Mehlich-3 solution consisting of (0.2 M CH₃COOH + 0.25 M NH₄NO₃ + 0.015 M NH₄F + 0.013 M HNO₃ + 0.001 M EDTA at pH 2.50 ± 0.05), according to Mehlich (1984). This method is also used for phytoavailable trace elements, and as such provides a 'multi-functional' extract. Exchangeable trace elements (Ni, Co, Cr, Mn) were extracted in 0.1 M Sr(NO₃)₂ at a soil:solution ratio of 1:4 (10 g:40 mL) and 2-hour equilibrium time (Wang et al., 2003). This method was repeated on a selection of 25 samples using 0.01 M Sr(NO₃)₂ for a comparison. In addition, a second method for exchangeable trace elements was used with 0.01 M CaCl₂ separately with 2-hours equilibrium time (Houba et al. 2000; Meers et al., 2007). This method was also repeated on the same selection of 25 samples used in the Sr(NO₃)₂ method using 0.1 M CaCl₂.

Potentially phytoavailable trace elements (Ni, Co, Cr, Mn) were extracted with standard Diethylene triamine pentaacetic acid (DTPA) according to Lindsay and Norvell (1969), and also (separately) using an adaptation by Becquer et al. (1995), which excluding TEA, was adjusted to pH 5.3 and had an equilibrium time of 2-hours (instead of 1-hour). Another method for potentially phytoavailable trace elements was also used, by extraction with a mixture of carboxylic acids (acetic, malic and citrate acid in molar ratio of 1:2:2 at 0.01 M) at

a soil:solution ratio of 1:4 (10 g:40 mL) and 2-hour shaking equilibrium time (method loosely based on Feng et al., 2005). Exchangeable cations were extracted with silver-thioarea (Dohrmann, 2006) over a 16-hour equilibrium time on an end-over-end shaker in the dark (to prevent silver precipitation). In addition, a second method for exchangeable cations, the traditional 1 M pH 7.0 ammonium acetate (NH₄Ac) was used with 2-hours equilibrium time (Meers et al., 2007).

All soil extractions were undertaken in disposable 50 mL polypropylene (PP) centrifuge tubes. Soil samples were weighed using a 4-decimal balance and weights recorded for correction of the precise weights in the mass balance calculations. All samples were agitated ('equilibrated') for method-specific times using an end-over-end shaker at 60 rpm and subsequent centrifuged (10 minutes at 4000 rpm) and the supernatant was collected in 10 mL polyethylene tubes. The extraction methods and operational conditions are given in Table 1. Nickel, Co and Cr partitioning was evaluated with a 5-step selective sequential extraction scheme to provide operationally defined solid-phase trace element (Ni, Cr, Co, Mn) fractionation. This scheme is based on Quentin et al. (2002), which was in turn modified mainly from Leleyter and Probst (1999). Adaptations were made here by combining step 1 and step 2, and by using HNO₃/HF high-pressure microwave digests for the residual fraction (step 5) instead of an alkaline fusion as in Quentin et al. (2002). The step for the 'organic bound phase' was also omitted because the tested soils were extremely low in organic matter. As such the fractions were: water soluble and exchangeable (i), bound to Mn oxides (ii), bound to amorphous Fe oxides (iii), bound to crystalline Fe oxides (iv), and residual (v). After each extraction step, the tubes were centrifuged for 10 minutes at 4000 rpm and the supernatants were then filtered through 0.45 µm membranes. The residues were washed with 20 mL of TDI water, centrifuged again for 10 minutes at 4000 rpm, the water decanted, and the residue dried at 40°C prior to the next extraction step. The different extraction phases and operational conditions are presented in Table 2.

All soil extracts samples were analyzed with Inductively Coupled Plasma Atomic Emission Spectroscopy (ICP-AES) (Varian Vista Pro II) for Ni, Co, Cu, Zn, Mn, Fe, Mg, Ca, Na, K, S and P. Each method included 3 sample blanks, 2 NIST standards, 2 ASPAC reference soils, 3 random sample duplicates and 3 multi-element standards as part of the quality control. The ICP-AES instrument was calibrated using a 6-point multi-element standard (which included all measured elements) prepared in each extraction solution.

Statistical analysis

The soil chemistry data was analysed using the software package STATISTICA Version 9.0 (StatSoft), Excel for Mac version 2011 (Microsoft) and PRIMER Version 6 (PRIMER-E).

RESULTS

Extraction methods

In the context of ultramafic soils, the most frequently used extraction methods to estimate potentially phytoavailable Ni, Co and other trace elements are the DTPA-extract, NH_4Ac extract, CaCl_2 extract and the $\text{Sr}(\text{NO}_3)_2$ extract (McLaughlin et al. 2000; Kukier and Chaney, 2001; Wang et al. 2006). Of these DTPA-TEA, DTPA, 1 M NH_4Ac , 0.01 M and 0.1 M CaCl_2 , 0.1 M and 0.01 M $\text{Sr}(\text{NO}_3)_2$, were tested here, in addition to the Mehlich-3 extract and a 0.01 M mix of carboxylic acids. Figure 1 compares the different extraction methods for the amounts of Ni, Cr and Co extracted. Extractable amounts of Cr are extremely low for all extractants, but the acid-based extractants released the most Cr. The greatest amounts of Ni and Co were extracted with acid-based extracts, and relatively large amounts were also extracted with DTPA. Nickel is moderately extractable (3.5% of the mean pseudo-total soil Ni), but Cr was almost completely unavailable (0.009% of the mean pseudo-total soil Cr) in the DTPA extracts (Table 3). Figure 2 shows correlation for Co and Ni between DTPA extract and other extracts (0.1 M CaCl_2 , 0.1 M $\text{Sr}(\text{NO}_3)_2$ and Mehlich-3). For the Co and Ni, relatively high correlation exists between the DTPA and Mehlich-3 extracts, followed by CaCl_2 , but low correlation with the $\text{Sr}(\text{NO}_3)_2$ extract (Supplementary Fig. 1).

Buffered and unbuffered DTPA extractions

Unbuffered DTPA (excluding TEA, pH 5.3) was used for all soils ($n = 343$) and buffered DTPA (including TEA, pH 7.3) on a subset of soils ($n = 93$). Both methods correlate well ($r = 0.75$ for Co, 0.59 for Cr, 0.84 for Mn, and 0.80 for Ni; at $p < 0.01$), but the buffered DTPA extracts had considerably more Ni, Co and Mn compared to unbuffered DTPA, indicating the role of extraction solution pH in the extraction method.

$\text{Sr}(\text{NO}_3)_2$ and CaCl_2 (0.1 and 0.01M) extractions

The 0.1 M and 0.01 M $\text{Sr}(\text{NO}_3)_2$ and 0.1 M and 0.01 M CaCl_2 extractants were tested on a selection of 25 samples, as it was predicted that the higher molarity would result in higher extractable levels of Ni and Co (hence better detection precision during ICP-AES analysis),

but would also induce a stronger drop in pH due to displacement of H^+ . Comparing 0.01 M $CaCl_2$ with 0.1 M $Sr(NO_3)_2$ shows a very good correlation ($r = 0.99$), and a good correlation ($r = 0.82$ at $p < 0.01$) was obtained for 0.1 M $CaCl_2$ versus 0.1 M $Sr(NO_3)_2$ (Fig. 3). In both cases, the $Sr(NO_3)_2$ extracts had greater amounts of Ni (up to a factor 3), which can be explained by the greater cation displacement power for Sr^{2+} compared to Ca^{2+} at the same molar concentration. The change in pH in both extracts is somewhat erratic (in relation to the extracted amount of Ni) but vary little between the extractants. Calcium or Sr concentration does affect competition with sorption and pH of the extraction fluid which clearly affects extractable Ni (Fig. 3).

Multivariate regression of independent parameters

Soil extraction methods generally benefit from having factors for prediction of plant response such as soil pH and CEC incorporated in the multivariate regression model to increase accuracy (Haq et al., 1980; Meers et al., 2007; Siebielec et al., 2007; Römken et al., 2009). Here we use a multivariate regression expressing extractable Ni or Co contents as a function of independent factors using the following equation:

$$\log(M_{\text{extraction}}) = \alpha \log(M_{\text{total}}) + \beta \text{pH} + \gamma \log(\text{CEC}) + \delta$$

$M_{\text{extraction}}$	Metal extracted in $\mu\text{g g}^{-1}$
M_{total}	Pseudo-total metal concentration (HNO_3/HCl -digest) in $\mu\text{g g}^{-1}$
pH	Measured soil pH
CEC	Soil CEC in $\text{mmol}^{(+)}\text{kg}^{-1}$
α, β, γ and δ are constants	

All independent input variables were log-transformed (except for pH, which is already log-transformed) prior to regression analysis. The regression equation provides empirical information on relevant factors influencing Ni or Co extractability using various extractants, thus allowing for the identification of the most influencing of these factors. We tested various factors, including pH, EC, CEC, pseudo-total metal concentrations and extractable metal concentrations (other than Ni or Co), but after evaluating the fits only pH, CEC and pseudo-total metal concentrations provided improved predictability. Table 4 and 5 lists the calculated factors and constants for the multivariate analysis, and it is clear that in most cases pseudo-total Ni or Co is the single most important factor predicting extractability in the extracts.

Sequential extraction for trace element partitioning

Selective sequential extractions show that Cr is mainly associated with crystalline Fe-oxides, explaining its low extractability in the $\text{Sr}(\text{NO}_3)_2$ and DTPA extracts. However, Ni and Co are mainly bound in the Mn-oxides and most of all in amorphous Fe-oxides fractions and hence are more phytoavailable, as is evident from higher concentrations in the $\text{Sr}(\text{NO}_3)_2$ and DTPA extracts. Table 6 shows relative portions of Ni, Co and Cr over the 5 fractions of the selective extraction procedure.

Cation Exchange capacity (CEC) and component ions

Figure 4 shows K extractability in silverthiorea (AgTU) extract versus K in Mehlich-3 (ML-3), 0.1 M $\text{Sr}(\text{NO}_3)_2$ and carboxylic acid (CA) extracts. These extraction methods are highly intercorrelated for K, and for Ca and Mg (data not shown), and therefore all could be used for measuring cation exchange capacity and component ions. This has major benefits, as combining methods (for example the carboxylic acid extract) for both cations and trace elements reduces time and costs.

Plant available phosphorus

Olsen-P and Mehlich-3 extractions are relatively poorly correlated ($r = 0.41$ at $p < 0.01$), but given that the majority of the ultramafic soils are between pH 3.76–5.5, the Mehlich-3 method appears more appropriate. However, for near-neutral pH ultramafic soils, the Olsen-method would be more likely to be related to plant available P than the Mehlich-method.

DISCUSSION

Although ultramafic soils are enriched in the trace elements Ni, Co, Cr, and Mn, phytoavailable concentrations of these elements are only a fraction of the pseudo-total concentrations present in the soil. Nevertheless, these elements are potentially present at phytotoxic concentrations and are widely seen as important factors contributing to the adversity of strongly acidic ultramafic soils to plants (Crooke and Inkson, 1955; Kukier and Chaney, 2003). In addition, major cation imbalance towards Mg and the deficit in Ca, N, P and K are also thought to be major factors affecting plants growing in these soils (Walker et al., 1955; Brooks, 1987; Nagy and Proctor, 1997; Echevarria, 2018). The high geodiversity of Mount Kinabalu Park, a result of complex geology, paleo-history, topography and climate, has created ultramafic soils that are extremely diverse in their pedology and chemical properties (van der Ent et al., 2018). Appropriate chemical characterization of these soils is

therefore a challenge. Unfortunately, most chemical extractants used for characterizing soils (trace elements, nutrient status and cation exchange complex) were historically developed for agricultural soils with restricted ranges of main properties, or within ecotoxicological frameworks, and the appropriateness for using such methods for estimating ecological effects on plants growing in tropical ultramafic soils is questionable. Although there is only limited knowledge of the precise chemical interactions of these extractions in ultramafic soils, most extraction methods tested here perform well in characterizing different types of ultramafic soils. In ultramafic soils, Ni is mainly associated with low-charge (serpentine, talc) and high-charge (smectite) clays, with Fe-(Mn) oxides and with Layered Double Hydroxide (LDH) phase minerals (Siebecker and Sparks, 2010), but there is no extraction method specific for these phases, although Ni may be (partly) co-extracted in existing methods. However, exchangeable Al and Mg are more important in the case of the carboxylic and DTPA extractable Ni. This can be explained by the presence of Layered Double Hydroxide (LDH) phase minerals that can adsorb Ni, and from which the carboxylic acid and DTPA extractants can desorb (part of the) Ni. In the unbuffered salt extracts ($\text{Sr}(\text{NO}_3)_2$), soil pH is an important factor. This pH-dependence is not surprising because this extractant essentially represents solubility of Ni at soil pH. Regarding Cr, strong chelating agents (as anions) can desorb CrVI anions. In the case of DTPA, it is interesting to note that acid pH mobilizes less Cr than higher pH. This fits well with the fact that anions are more easily desorbed at higher pH. When Mn is highly available, Cr^{3+} can be oxidized by reducing MnIV into MnII. CrVI is then complexed onto the surface of Fe-oxides, and therefore desorbed by DTPA or carboxylates such as citrate, which in turn adsorb onto the surface of Fe-oxides such as other anionic extractants, *e.g.* phosphate (Raous et al., 2013).

Exchangeable cations are fundamental to buffering soil pH and hence directly and indirectly influence many soil processes. CEC soil extraction methods rely on attempting to displace all exchangeable cations. Exchangeable cations are often measured with the 1 M pH 7.0 ammonium acetate (NH_4Ac) method (DIN, 1995), with cobalt(III) hexamine trichloride method (Cisielski and Sterckeman, 1997; in the case of ultramafic soils: Raous et al., 2013) or the silverthioarea method (Pleysier and Juo, 1980; Searle, 1986; Dohrmann, 2006). The second method is not adapted to evaluate exchangeable Co in ultramafic soils. The last is, however, not frequently used for ultramafic soils, but has the benefit over NH_4Ac as it adopts the pH of the soil solution, has a greater cation displacement power, and exchange takes place at low ionic strength (0.01 M) (Pleysier and Juo, 1980; Proctor et al., 1981). As is typical for

ultramafic soils, the cation exchange complex is saturated with Mg^{2+} , but absolute concentrations of Ca^{2+} are not always low. The total cation exchangeable capacity (CEC) is lowest in the strongly leached Ferralsols, and high to extremely high in hypermagnesian Leptosols and serpentinitic Cambisols (Echevarria, 2018; van der Ent et al., 2018).

Apart from pseudo-total P (acid digest), P was extracted with the Olsen-P and Mehlich-3 methods, aimed at quantifying approximate plant available P concentrations. The Olsen-P method was initially developed for calcareous soils as the extraction solution is pH 8.5 (Sharpley et al., 2008), whereas the Mehlich-3 method was developed for non-calcareous soils with an extractant solution of pH 2.5 (Mehlich, 1984). The Mehlich-3 method is similar to the Bray-1 method as both employ a dilute NH_4F extraction medium (Lucero et al. 1998). Olsen P was proved to selectively extract isotopically-exchangeable P in a wide array of soils (Fardeau et al., 1988) which makes it a more universal method than initially thought.

A soil extraction method should be able to explain potential phytotoxicity effects as a result of trace elements, and should have predictive power for plant-uptake of these elements or at least a correlation with plant-uptake. However, plant uptake usually depends on the plant status and needs and may substantially vary between species. Even more, with high supply, plant uptake can respond without any correlation with the chemically mobile pool of an element (e.g. Ni for *Alyssum murale* in low pH soils: see Kukier et al., 2004; Bani et al., 2014). Ideally, the extractant provides simultaneous information about trace elements, exchangeable cations (major and trace) and macronutrients. Of the tested method, only Mehlich-3 method was specifically designed for such a 'multi-functional' purpose (Mehlich, 1984). This method uses dilute acetic acid combined with low concentration of EDTA for release and chelation of trace elements. Also present in the extraction solution are NH_4NO_3 to exchange cations from the exchange complex, and further NH_4F to extract P (hence essentially identical to the Bray-1 method of phytoavailable P). The 0.01 and 0.1 M $Sr(NO_3)_2$ extracts appear to achieve more consistent results compared to other dilute neutral salt extracts (0.01 and 0.1 MM $CaCl_2$), and have been shown to correlate with immediate Ni phytotoxicity (Kukier and Chaney, 2001; Siebielec et al., 2007). The buffered and unbuffered DTPA methods are highly correlated, despite the fact that the unbuffered version is more adaptable to the soil pH. The high correlation of the carboxylic acid method not only with DTPA, Mehlich-3, $CaCl_2$ and $Sr(NO_3)_2$ extractable trace elements (Ni, Cr, Co), but also with silverthiorexa-CEC and silverthiorexa exchangeable cations (Ca, Mg, K) means that this method is well-suited for

general characterization of tropical ultramafic soils. However, as with all methods tested here, ecological implications for tropical vegetation or elemental uptake in native plants growing in tropical ultramafic soils remains unstudied. Without plant uptake data, the predictive values of the soil extractions should be interpreted cautiously. Future research is required to assess suitable extraction methods for ultramafic soils by evaluating the correlation of extraction results with plant composition or deficiency, adequacy and toxicity of specific elements. There is the need to test whether the displaced soil solution extraction method (Proctor et al., 1981; Kukier et al., 2004; Coinchelin et al., 2012) could be useful in predicting short-term phytoavailability of trace elements in local ultramafic soils as it was shown that hyperaccumulators can concentrate Ni during active root uptake (up to five times) from the initial concentration in the displaced soil solution, that means, depletion of the labile pools is strongly active (Coinchelin et al., 2012). Furthermore, tropical ultramafic soils present suitable opportunity for use in Ni agromining (a technology that extracts strategic metals from the biomass of selected ‘metal crops’) (van der Ent et al., 2013b; 2015c). Hence, it is imperative to develop robust Ni phytoavailability assays to predict Ni yield in ‘metal crops’ as Ni accumulation by ‘hypernickelophores’ has little evident relationship with single soil extraction methods including displaced soil solution (Nkrumah et al., 2016). The fact that plant species native to (tropical) ultramafic soils have evolved while confronted with extreme soil chemistry means that they must be highly tolerant, and therefore the results from experimental trials using ‘normal’ plants are not immediately applicable. Finally, plants vary widely not only in their tolerance to soil chemistry, but also in their uptake characteristics of trace elements (Ni, Co, Cr, Mn) depending on genotypic and phenotypic controlled ecophysiologicals (Ernst, 2006; Ünver et al., 2013), therefore aiming for a soil extraction method that accurately predicts the response of all plant species on ultramafic soils is futile. Rather the method of choice needs to be inexpensive and simple, comparable to existing literature, as well as applicable to a wide range of different ultramafic soils. These conditions are met specifically with the carboxylic acid method. For more complex hyperaccumulator crops, both a mild extraction that is related to soil chemistry and a more intense DTPA extractions might be relevant to assess both the initial pool in the soil solution (*i.e.*, intensity) and the entire pool that replenishes the soil solution (*i.e.*, quantity) under significant depletion exerted by hyperaccumulators (Echevarria et al., 1998, 2006; Coinchelin et al., 2012).

ACKNOWLEDGEMENTS

We wish to thank Sabah Parks for their support and the SaBC for granting permission for conducting research in Sabah, and to extend our gratitude to Kalipin Al-Hafsan Bin Sampin and Hali bin Nassim in the field, and Vanessa Ullmann in the laboratory. The French National Research Agency through the national “Investissements d’avenir” program (ANR-10-LABX-21, LABEX RESSOURCES21) and through the ANR-14-CE04-0005 Project “Agromine” is acknowledged for funding support to A. van der Ent and P.N. Nkrumah. A. van der Ent is the recipient of a Discovery Early Career Researcher Award (DE160100429) from the Australian Research Council. P.N. Nkrumah is the recipient of an Australian Government Research Training Program Scholarship and UQ Centennial Scholarship at The University of Queensland, Australia.

REFERENCES

- Anderegg, G., Arnaud-Neu, F., Delgado, R., Felcman, J., Popov, K., 2005. Critical evaluation of stability constants of metal complexes of complexones for biomedical and environmental applications* (IUPAC Technical Report). *Pure and Applied Chemistry* 77(8), pp. 1445–1495.
- Anderson, P.R., Christensen, T.H., 1988. Distribution coefficients of Cd, Co, Ni, and Zn in soils. *Journal of Soil Science* 39, pp. 15–22.
- Bani, A., Echevarria, G., Mullaj, A., Reeves, R.D., Morel, J.L., Sulçe, S., 2009. Ni hyperaccumulation by *Brassicaceae* in serpentine soils of Albania and NW Greece. *Northeastern Naturalist* 16, pp. 385–404.
- Bani, A., Echevarria, G., Montargès-Pelletier, E., Gjoka, F., Sulce, S., Morel, J.L., 2014. Pedogenesis and nickel biogeochemistry in a typical Albanian ultramafic toposequence. *Environmental Monitoring and Assessment* 186, pp. 4431–4442.
- Becquer, T., Bourdon, E., Pétard, J., 1995. Disponibilité du nickel le long d'une toposéquence de sols développés sur roches ultramafiques de Nouvelle-Calédonie. *Comptes rendus de l'Académie des sciences. Série 2. Sciences de la terre et des planètes* 321(7), pp. 585–592.

Becquer, T., Pétard, J., Duwig, C., Bourdon, E., Moreau, R., Herbillon, A.J., 2001. Mineralogical, chemical and charge properties of Geric Ferralsols from New Caledonia. *Geoderma* 103(3), pp. 291–306.

Becquer, T., Quantin, C., Sicot, M., Boudot, J., 2003. Chromium availability in ultramafic soils from New Caledonia. *The Science of the Total Environment* 301(1-3), pp. 251–261.

Brady, K.U., Kruckeberg, A.R., Bradshaw, H.D.Jr., 2005. Evolutionary ecology of plant adaptation to serpentine soils. *Annual Review of Ecology, Evolution, and Systematics* 36, pp. 243–266.

Brearley, F., 2005. Nutrient limitation in a Malaysian ultramafic soil. *Journal of Tropical Forest Science* 17(4), pp. 596–609.

Brooks, R.R., 1987. *Serpentine and its vegetation: A multidisciplinary approach*. Dioscorides Press, Portland, Oregon, USA.

Bruemmer, G.W., Gerth, J., Herms, U. 1986. Heavy metal species, mobility and availability in soils. *Zeitschrift für Pflanzenernährung und Bodenkunde* 149, pp. 382–398.

Chardot, V., Echevarria, G., Gury, M., Massoura, S., Morel, J.L., 2007. Nickel bioavailability in an ultramafic toposequence in the Vosges Mountains (France). *Plant and Soil* 293(1-2), pp. 7–21.

Chardot-Jacques, V., Calvaruso, C., Simon, B., Turpault, M.-P., Echevarria, G., Morel, J.L., 2013. Chrysotile dissolution in the rhizosphere of the nickel hyperaccumulator *Leptoplax emarginata*. *Environmental Science and Technology* 47(6), pp. 2612–2620.

Coinchelin, D., Bartoli, F., Robin, C., Echevarria, G., 2012. Ecophysiology of nickel phytoaccumulation: a simplified biophysical approach. *Journal of Experimental Botany* 63(16), pp. 5815–5827.

Collenette, P., 1964. A short account of the geology and geological history of Mt Kinabalu. *Proceedings of the Royal Society of London, Series B* 161, pp. 56–63.

Cottam, M., Hall, R., Sperber, C., Armstrong, R., 2010. Pulsed emplacement of the Mount Kinabalu granite, Northern Borneo. *Journal of the Geological Society* 167(1), pp. 49–60.

Crooke, W.M., Inkson, R., 1955. The relationship between nickel toxicity and major nutrient supply. *Plant and Soil* 6(1), pp. 1–15.

Ciesielski, H., Sterckeman, T., 1997. Determination of cation exchange capacity and exchangeable cations in soils by means of cobalt hexamine trichloride. Effects of experimental conditions. *Agronomie* 17(1), pp. 1–7.

Dohrmann, R., 2006. Cation exchange capacity methodology II: A modified silver-thiourea method. *Applied Clay Science* 34(1-4), pp. 38–46.

Echevarria, G., 2018. Genesis and behaviour of ultramafic soils and consequences for nickel biogeochemistry. In: *Agromining: extracting unconventional resources from plants*, Mineral Resource Reviews series, Van der Ent A, Echevarria G, Baker AJM, Morel JL (Eds.), Springer, Cham, pp. 135–156.

Echevarria, G., Morel, J.L., Fardeau, J.C., Leclerc-Cessac, E., 1998. Assessment of phytoavailability of nickel in soils. *Journal of Environmental Quality* 27, pp. 1064–1070.

Echevarria, G., Massoura, S., Sterckeman, T., Becquer, T., Schwartz, C., Morel, J.L., 2006. Assessment and control of the bioavailability of Ni in soils. *Environmental Toxicology and Chemistry* 25, pp. 643–651.

Ernst, W., 2006. Evolution of metal tolerance in higher plants. *Forest Snow and Landscape Research* 80(3), pp. 251–274.

Estrade, N., Cloquet, C., Echevarria, G., Sterckeman, T., Deng, T-H-B., Tang, Y-T., Morel, J.L., 2015. Weathering and vegetation controls on nickel isotope fractionation in surface ultramafic environments (Albania). *Earth and Planetary Science Letters* 423, pp. 24–25.

Fandeur, D., Juillot, F., Morin, G., Olivi, L., Cognigni, A., Ambrosi, J.-P., *et al.* 2009. Synchrotron-based speciation of chromium in an Oxisol from New Caledonia: Importance of secondary Fe-oxyhydroxides. *American Mineralogist* 94(5-6), pp. 710–719.

Fardeau, J.C., Morel, C., Boniface, R., 1988. Why the Olsen method should be used to estimate available soil-phosphorus. *Agronomie* 8(7), pp. 577–584.

Feng, M., Shan, X., Zhang, S., Wen, B., 2005. A comparison of the rhizosphere-based method with DTPA, EDTA, CaCl_2 , and NaNO_3 extraction methods for prediction of bioavailability of metals in soil to barley. *Environmental Pollution* 137(2), pp. 231–240.

Garnier, J., Quantin, C., Martins, E.S., Becquer, T., 2006. Solid speciation and availability of chromium in ultramafic soils from Niquelândia, Brazil. *Journal of Geochemical Exploration* 88(1-3), pp. 206–209.

Gommy, C., Perdrix, E., Galloo, J.-C., Guillermo, R., 1998. Metal speciation in soil: extraction of exchangeable cations from a calcareous soil with a magnesium nitrate solution. *International Journal of Environmental Analytical Chemistry* 72(1), pp. 27–45.

Gryschko, R., Kuhnle, R., Terytze, K., Breuer, J., Stahr, K., 2004. Soil extraction of readily soluble heavy metals and As with 1 M NH_4NO_3 -Solution - Evaluation of DIN 19730 (6 pp). *Journal of Soils and Sediments* 5(2), pp. 101–106.

Hara, T., Sonoda, Y., 1979. Comparison of the toxicity of heavy metals to cabbage growth. *Plant and Soil* 51(1), pp. 127–133.

Haq, A.U., Bates, T.E., Soon, Y.K., 1980. Comparison of extractants for plant-available zinc, cadmium, nickel, and copper in contaminated soils. *Soil Science Society of America Journal*, 44(4), pp. 772–777.

Houba, V.J.G., Temminghof, E.J.M., Gaikhorst, G.A., van Vark, W., 2000. Soil analysis procedures using 0.01M calcium chloride as extraction reagent. *Communications in Soil Science and Plant Analysis*, 31 (2000), pp. 1299–1396.

Hsiao, K.H., Bao, K.H., Wang, S.H., Hseu, Z.Y., 2009. Extractable concentrations of cobalt from serpentine soils with several single-extraction procedures. *Communications in Soil Science and Plant Analysis* 40(13-14), pp. 2200–2224.

Kukier, U., Chaney, R.L., 2001. Amelioration of nickel phytotoxicity in muck and mineral soils. *Journal of Environmental Quality* 30(6), pp. 1949–1960.

Kukier, U., Chaney, R.L., 2003. In situ remediation of nickel phytotoxicity for different plant species. *Journal of Plant Nutrition* 27(3), pp. 465–495.

Lazarus, B.E., Richards, J.H., Claassen, V.P., O'Dell, R.E., Ferrell, M.A., 2011. Species specific plant-soil interactions influence plant distribution on serpentine soils. *Plant and Soil* 342(1-2), pp. 327–344.

Latham, M., 1975. Les sols d'un massif de roches ultrabasiques de Nouvelle-Caledonie: le Boulinda. Les sols à accumulation ferrugineuse. *Cahiers ORSTOM, Série Pédologie* 13, pp. 159–172.

Lee, B.D., Sears, S.K., Graham, R.C., Amrhein, C., Vali, H., 2003. Secondary mineral genesis from chlorite and serpentine in an ultramafic soil toposequence. *Soil Science Society of America Journal* 67(4), pp. 1309–1317.

Lewis, J.F., Draper, G., Espaillat, J., Proenza, J.A., Jiménez, J., 2006. Ophiolite-related ultramafic rocks (serpentinites) in the Caribbean region: A review of their occurrence, composition, origin, emplacement and Ni-laterite soil formation. *Geologica Acta* 4(1), pp. 237–264.

Leleyter, L., Probst, J.L., 1999. A new sequential extraction procedure for the speciation of particulate trace elements in river sediments. *International Journal of Environmental Analytical Chemistry* 73, pp. 109–128.

L'Huillier, L., Edighoffer, S., 1996. Extractability of nickel and its concentration in cultivated plants in Ni-rich ultramafic soils of New Caledonia. *Plant and Soil* 186(2), 255–264.

Lindsay, W.L., Norvell, W.A., 1978. Development of DTPA soil test for zinc, iron, manganese, and copper. *Soil Science Society of America Journal* 42, pp. 421–428

Lucero, D.W., Martens, D.C., McKenna, J.R., Starner, D.E., 1998. Comparison of Mehlich 3- and Bray 1-extractable phosphorus levels in a Starr clay loam amended with poultry litter. *Communications in Soil Science and Plant Analysis* 29(9-10), pp. 1133–1142.

Madden, M.S., 1988. Adapting the $\text{Sr}(\text{NO}_3)_2$ method for determining available cations to a routine soil testing procedure. M.Sc. thesis, University of Wisconsin, Madison.

Massoura, S. T., Echevarria, G., Leclerc-Cessac, E., Morel, J. L., 2004. Response of excluder, indicator and hyperaccumulator plants to nickel availability in soils. *Australian Journal of Soil Research* 42(8), pp. 933–938.

Massoura, S.T., Echevarria, G., Becquer, T., Ghanbaja, J., Leclerc-Cessac, E., Morel, J.L., 2006. Control of nickel availability by nickel bearing minerals in natural and anthropogenic soils. *Geoderma* 136(1-2), pp. 28–37.

McLaughlin, M.J., Zarcinas, B.A., Stevens, D.P., Cook, N., 2000. Soil testing for heavy metals. *Communications in Soil Science and Plant Analysis* 31(11-14), pp. 1661–1700.

Meers, E., Samson, R., Tack, F.M.G., Ruttens, A., Vandegheuchte, M., Vangronsveld, J., Verloo, M.G., 2007. Phytoavailability assessment of heavy metals in soils by single extractions and accumulation by *Phaseolus vulgaris*. *Environmental and Experimental Botany*, 60(3), pp. 385–396.

Mehlich, A., 1984. Mehlich-3 soil test extractant: a modification of Mehlich-2 extractant. *Communications in Soil Science and Plant Analysis* 15(12), pp. 1409–1416.

Mengel, K., Kirkby, E.A., 2001. *Principles of Plant Nutrition*. Springer (2001).

Menzies, N.W., Donn, M.J., Kopittke, P.M., 2007. Evaluation of extractants for estimation of the phytoavailable trace metals in soils. *Environmental Pollution*, 145(1), pp. 121–130.

Nagy, L., Proctor, J., 1997. Soil Mg and Ni as causal factors of plant occurrence and distribution at the Meikle Kilrannoch ultramafic site in Scotland. *New Phytologist* 135(3), pp. 561–566.

Nkrumah, P.N., Baker, A.J.M., Chaney, R.L., Erskine, P.D., Echevarria, G., Morel, J.L., van der Ent, A., 2016. Current status and challenges in developing nickel phytomining: an agronomic perspective. *Plant and Soil* 406, pp. 55–69.

Novozamsky, I., Lexmond, T.M., Houba, V.J.G., 1993. A single extraction procedure of soil for evaluation of uptake of some heavy metals by plants. *International Journal of Environmental Analytical Chemistry* 51(1), pp. 47–58.

Oze, C., Fendorf, S., Bird, D.K., Coleman, R.G., 2004. Chromium geochemistry in serpentinized ultramafic rocks and serpentine soils from the Franciscan complex of California. *American Journal of Science*, 304(1), pp. 67–101.

Pleysier, J.L., Juo, A., 1980. A single-extraction method using silver-thiourea for measuring exchangeable cations and effective CEC in soils with variable charges. *Soil Science*, 129(4), pp. 205–211.

Proctor, J., Baker, A.J.M., van Balgooy, M., Bruijnzeel, L., Jones, S., Madulid, D., 2000. Mount Bloomfield, Palawan, Philippines: forests on greywacke and serpentinized peridotite. *Edinburgh Journal of Botany*, 57(01), pp. 121–139.

Proctor, J., Johnston, W.R., Cottam, D.A., Wilson, A.B., 1981. Field-capacity water extracts from serpentine soils. *Nature* 294:245–246.

Proctor, J., 2003. Vegetation and soil and plant chemistry on ultramafic rocks in the tropical Far East. *Perspectives in Plant Ecology Evolution and Systematics*, 6(1-2), pp. 105–124.

Quantin, C., Becquer, T., Rouiller, J.H., Berthelin J., 2002. Redistribution of metals in a New Caledonia Ferralsol after microbial weathering. *Soil Science Society of America Journal*, 66(6), pp. 1797–1804.

Quantin, C., Ettler, V., Garnier, J., Šebek, O., 2008. Sources and extractibility of chromium and nickel in soil profiles developed on Czech serpentinites. *Comptes Rendus Geoscience* 340(12), pp. 872–882.

Quevauviller, P., Rauret, G., Muntau, H., 1994. Evaluation of a sequential extraction procedure for the determination of extractable trace metal contents in sediments. *Fresenius' Journal of Analytical Chemistry* 349, pp. 808–814.

Raous, S., Echevarria, G., Sterckeman, T., Hanna, K., Thomas, F., Martins, E.S., Becquer, T., 2013. Potentially toxic metals in ultramafic mining materials: identification of the main bearing and reactive phases. *Geoderma* 192, pp. 111–119.

Rayment, G.E., Higginson F.R., 1992. *Australian Laboratory Handbook of Soil and Water Chemical Methods*. Inkata Press, Melbourne.

Robinson, B.H., Brooks, R.R., Gregg, P.E.H., Kirkman, J.H., 1999. The nickel phytoextraction potential of some ultramafic soils as determined by sequential extraction. *Geoderma* 87, pp. 293–304.

Robinson, B., Fernandez, J., Madejon, P., Maranon, T., Murillo, J., Green, S., Clothier, B., 2003. Phytoextraction: an assessment of biogeochemical and economic viability. *Plant and Soil* 249(1), pp. 117–125.

Searle, P.L., 1986. The measurement of soil cation exchange properties using the single extraction, silver thiourea method-an evaluation using a range of New Zealand soils. *Soil Research* 24(2), pp. 193–200.

Römken, P.F., Guo, H.-Y., Chu, C.-L., Liu, T.-S., Chiang, C.-F., Koopmans, G.F., 2009. Characterization of soil heavy metal pools in paddy fields in Taiwan: chemical extraction and solid-solution partitioning. *Journal of Soils and Sediments* 9(3), pp. 216–228.

Sharpley, A.N., Kleinman, P.J., Weld, J.L., 2008. Environmental soil phosphorus indices. In: Carter, M.R., Gregorich, E.G., editors. *Soil Sampling and Methods of Analysis*, 2nd Edition. Canadian Society of Soil Science. CRC Press, Boca Raton, FL, pp. 141–159.

Siebecker, M., Sparks D.L., 2010. Nickel speciation in serpentine soils using synchrotron radiation techniques. 2010 19th World Congress of Soil Science, Soil Solutions for a Changing World. 1 – 6 August 2010, Brisbane, Australia.

Siebielec, G., Chaney, R.L., Kukier, U., 2007. Liming to remediate Ni contaminated soils with diverse properties and a wide range of Ni concentration. *Plant and Soil*, 299(1-2), pp. 117–130.

Sukkariyah, B.F., Evanylo, G., Zelazny, L., Chaney, R.L., 2005. Cadmium, copper, nickel, and zinc availability in a biosolids-amended piedmont soil years after application. *Journal of Environmental Quality* 34, pp. 2255–2262.

Tiller, K.G., Gehrt, J., Bruemmer, G., 1984. The sorption of Cd, Zn and Ni by soil clay fractions: Procedures for partition of bound forms and their interpretation. *Geoderma* 34, pp. 1–16.

Ünver, İ., Madenoğlu, S., Dilsiz, A., Namlı, A., 2013. Influence of rainfall and temperature on DTPA extractable nickel content of serpentine soils in Turkey. *Geoderma* 202-203(C), pp. 203–211.

Van der Ent, A., Baker, A.J.M., Reeves, R.D., Pollard, A.J., Schat, H., 2013a. Hyperaccumulators of metal and metalloid trace elements: Facts and fiction. *Plant and Soil* 362(1-2), pp. 1–16.

Van der Ent, A., Baker, A.J.M., van Balgooy, M.M.J., Tjoa, A. 2013b. Ultramafic nickel laterites in Indonesia (Sulawesi, Halmahera): mining, nickel hyperaccumulators and opportunities for phytomining. *Journal of Geochemical Exploration* 128: 72–79.

Van der Ent, A., Erskine, P.D., Sumail S., 2015a. Ecology of nickel hyperaccumulator plants from ultramafic soils in Sabah (Malaysia). *Chemoecology*, 25(5), pp. 243–259.

Van der Ent, A., Sumail, S., Clarke, C., 2015b. Habitat differentiation of obligate ultramafic *Nepenthes* endemic to Mount Kinabalu and Mount Tambuyukon (Sabah, Malaysia). *Plant Ecology* 216, pp. 789–807.

Van der Ent, A., Baker, A.J.M., Reeves, R.D., Chaney, R.L., Anderson, C.W.N., Meech, J.A., Erskine, P.D., Simonnot, M-O., Vaughan, J., Morel, J.L., Echevarria, G., Fogliani, B., Rongliang, Q., Mulligan, D.R., 2015c. Agromining: farming for metals in the future? *Environmental Science and Technology* 49, pp. 4773–4780.

Van der Ent, A., Erskine, P.D., Mulligan, D.R., Repin, R., Karim, R., 2016a. Vegetation on ultramafic edaphic islands in Kinabalu Park (Sabah, Malaysia) in relation to soil chemistry and altitude. *Plant and Soil*, 403(1), pp. 77–101.

Van der Ent, A., Echevarria, G., Tibbett, M., 2016b. Delimiting soil chemistry thresholds for nickel hyperaccumulator plants in Sabah (Malaysia). *Chemoecology* 26(2), pp. 67–82.

Van der Ent, A., Cardace, D., Tibbett, M., Echevarria, G., 2018. Ecological implications of pedogenesis and geochemistry of ultramafic soils in Kinabalu Park (Malaysia). *Catena* 160, pp. 154–169.

Vlams, J., Jenny, H., 1948. Calcium deficiency in serpentine soils as revealed by absorbent technique. *Science* 107, pp. 549.

Walker, R.B., 1954. The ecology of serpentine soils. II. Factors affecting plant growth on serpentine soils. *Ecology* 35, pp. 259–266.

Walker, R.B., Walker, H.M., Ashworth, P.R., 1955. Calcium-magnesium nutrition with special reference to serpentine soils. *Plant Physiology* 30(3), pp. 214–221.

Wang, W.-S., Shan, X.-Q., Wen, B., Zhang, S.-Z., 2003. Relationship between the extractable metals from soils and metals taken up by maize roots and shoots. *Chemosphere* 53(5), pp. 523–530.

TABLES

Table 1. Chemical soil extractants used in this study.

Table 2. Selective sequential extraction scheme.

Table 3. Elemental concentrations (ranges and means) in different extracts ($\mu\text{g g}^{-1}$), $n = 343$.

Table 4. Multivariate regression analysis of Ni-extractions and multivariate regression analysis of Co-extractions.

Table 5. Correlations of extractability of different elements in extracts (major cations), $n = 343$ ** denotes $p < 0.0001$.

Table 6. Sequential and non-sequential extraction of a) Ni, b) Co, c) Cr and d) Mn in 14 representative soils (as percentage of AR digest). The pH of the representative soils ranges from 5.11 to 9.19. Sequential extraction: exchangeable (Exch.), oxidisable Mn-oxides (Mn-OX), amorphous Fe-oxides (AM-Fe) or crystalline Fe-oxides (CR-FE); non-sequential extraction: carboxylic acid (Carbox.), 0.1 M strontium nitrate (SrNO_3)₂ or Melhich-3 (ML-3); and digest: digest with either $\text{HNO}_3 + \text{HCl}$ (AR) or $\text{HNO}_3 + \text{HCl} + \text{HF}$ (HF). <LOD denotes below detection limit.

FIGURES

Figure 1. Boxplots of Ni, Co, Cr, Mn in various soil extracts ($\mu\text{g g}^{-1}$). Key to symbols: open squares are the \pm mean, whiskers are \pm standard deviation, circles are outliers and asterisks are extreme outliers.

Figure 2. Plot of principal component analysis (PCA) of a) Ni extractability in DTPA, 0.1 M CaCl_2 , 0.1 M $\text{Sr}(\text{NO}_3)_2$ and Mehlich-3 extracts and b) Co extractability in DTPA, 0.1 M CaCl_2 , 0.1 M $\text{Sr}(\text{NO}_3)_2$ and Mehlich-3 extracts.

Figure 3. Principal component analysis of nickel extractability in extracts of 0.1 M $\text{Sr}(\text{NO}_3)_2$, 0.01 M and 0.1 M CaCl_2 , versus extraction solution pH.

Figure 4. Principal component analysis of extractable K in AgTU, carboxylic acid (CA), $\text{Sr}(\text{NO}_3)_2$ (0.1 M) and Mehlich-3 extracts.

TABLE 1

Extraction medium	Liquid:soil ratio	Equilibrium time	References
<i>Neutral-salt extracts</i>			
0.1 M Strontium nitrate	4:1	1 hr	Wang et al., 2003
0.01M Strontium nitrate	4:1	1 hr	Kukier et al., 2004
0.1 M Calcium chloride	4:1	1 hr	Meers et al., 2011
0.01 M Calcium chloride	4:1	1 hr	Meers et al., 2011
<i>Chelator extracts</i>			
DTPA-TEA pH 7.4	5:1	2 hrs	Lindsay and Norvell, 1978
DTPA pH 5.3	5:1	2 hrs	Becquer et al., 1995
<i>Acid based extracts</i>			
70% Nitric acid, 37% hydrochloric acid	12:0.3	1 hr	Rayment and Higginson, 1992
Carboxylic acid mix (Acetic, malic, citric acid)	4:1	2 hrs	Feng et al., 2005
<i>Exchangeable cations</i>			
1 M ammonium acetate	10:1	2 hrs	DIN, 1995
0.01M Silverthiorea	40:0.8	16 hrs	Pleysier and Juo, 1980
<i>Phosphorus extracts</i>			
Olsen-P (sodium hydrogen carbonate)	20:1	30 min.	Olsen et al., 1954
Mehlich-3 (ammonium fluoride based)	10:1	5 min.	Mehlich, 1985

TABLE 2

Fraction	Reagents	Conditions
Water soluble and exchangeable	0.01 M Sr(NO ₃) ₂	2 hours at 20°C
Bound to Mn-oxides	0.1 M hydroxylammonium chloride	30 minute at 20°C
Bound to amorphous Fe-oxides	0.2 M ammonium oxalate + 0.2 M oxalic acid	4 hours in the dark at 20°C
Bound to crystalline Fe-oxides	0.2M ammonium oxalate + 0.2M oxalic acid + 0.1 M ascorbic acid	30 minutes at 90°C in water bath
Residual	70% HNO ₃ + 32% HCl + 37% HF	1 hour at 115°C in digestion microwave

TABLE 3

	DTPA extract $\mu\text{g g}^{-1}$		Mehlich-3 extract $\mu\text{g g}^{-1}$		$\text{Sr}(\text{NO}_3)_2$ extract $\mu\text{g g}^{-1}$		Carboxylic extract $\mu\text{g g}^{-1}$		HNO_3/HCl digest $\mu\text{g g}^{-1}$	
Al	0.3–485	55	17–1750	415	0.1–220	15	5.8–4250	265	1200–118000	19600
Ca	–	–	3.1–2600	230	1.8–1770	210	2.1–1550	125	205–24500	2100
Co	0.1–95	20	0.5–70	15	0.1–25	1.8	0.2–275	40	0.5–1500	250
Cr	0.1–15	0.5	0.04–40	2.5	0.1–25	0.3	0.3–85	5.8	225–21800	4000
Cu	0.1–25	1.5	0.3–125	5.0	0.1–25	0.2	0.1–8.7	0.5	2.5–325	50
Fe	0.6–875	95	25–1750	210	0.1–35	1.1	12.1–2350	410	21500–535000	157000
K	1.0–110	30	<0.01–440	35	0.1–135	30	0.1–95	25	0.1–1060	90
Mg	2.9–1100	315	5.1–6850	980	5.0–3150	510	4.0–7850	620	270–235000	26700
Mn	0.4–780	190	0.7–1150	215	0.3–11800	445	0.4–3700	550	45–33500	3250
Na	1.9–25	10	6.6–345	25	0.5–30	7.1	0.1–65	5.8	0.1–360	105
Ni	0.2–275	50	0.4–620	55	0.1–65	7.9	0.2–405	55	15–7000	1440
P	0.1–15	0.7	<0.01–30	2.0	0.1–735	60	0.1–45	1.6	20–485	130
S	1.0–215	17	<0.01–115	16.5	0.1–20	1.2	0.8–80	9.6	85–755	350
Zn	0.1–4.8	0.7	0.1–5.0	1.0	0.1–15	3.0	0.1–3.1	0.8	15–375	115

TABLE 4

Extraction method	δ	α	β	γ
Carboxylic acid Ni	-3.45	0.70	0.40	0.30
DTPA Ni	-2.50	0.85	0.10	0.50
0.1 M Sr(NO₃)₂ Ni	-0.90	0.70	-0.25	0.45
Mehlich-3 Ni	-2.80	0.65	0.30	0.40
Carboxylic acid Co	-0.85	0.95	-0.07	0.19
DTPA Co	-0.025	0.90	-0.25	0.20
0.1 M Sr(NO₃)₂ Co	0.80	0.70	-0.45	0.05
Mehlich-3 Co	-0.40	0.65	-0.05	0.15

TABLE 5

	HNO ₃ /HCl	Sr(NO ₃) ₂	carboxylic acid	Mehlich-3	DTPA
Na					
Sr(NO ₃) ₂	-0.07				
carboxylic acid	-0.08	0.90**			
Mehlich-3	0.10	0.30**	0.35**		
DTPA	-0.10	0.90**	0.95**	0.30**	
silverthiorea	0.15	0.25**	0.25**	-0.10	0.25**
Mg					
Sr(NO ₃) ₂	0.50**				
carboxylic acid	0.70**	0.55**			
Mehlich-3	0.70**	0.80**	0.80**		
DTPA	0.60**	0.85**	0.65**	0.90**	
silverthiorea	0.40**	0.50**	0.65**	0.60**	0.60**
K					
Sr(NO ₃) ₂	0.20				
carboxylic acid	0.25**	0.90**			
Mehlich-3	0.25**	0.75**	0.80**		
DTPA	0.25**	0.90**	0.95**	0.75**	
silverthiorea	0.30**	0.85**	0.90**	0.75**	0.90**
Al					
Sr(NO ₃) ₂	0.50**				
carboxylic acid	0.70**	0.50**			
Mehlich-3	0.70**	0.55**	0.9**		
DTPA	0.50**	0.50**	0.70**	0.80**	
silverthiorea	0.50**	0.55**	0.65**	0.65**	0.75**
Ca					
Sr(NO ₃) ₂	0.55**				
carboxylic acid	0.55**	0.85**			
Mehlich-3	0.55**	0.85**	0.95**		
silverthiorea	0.55**	0.85**	0.95**	0.95**	

Table 5–continued.

	HNO ₃ /HCl	carboxylic acid	Sr(NO ₃) ₂	Mehlich-3
Co				
carboxylic acid	0.80**			
Sr(NO ₃) ₂	0.30**	0.20**		
Mehlich-3	0.70**	0.75**	0.60**	
DTPA	0.70**	0.80**	0.55**	0.95**
Cr				
carboxylic acid	0.09			
Sr(NO ₃) ₂	-0.04	0.35**		
Mehlich-3	-0.10	0.00	-0.05	
DTPA	0.10	0.50**	0.80**	-0.05
Fe				
carboxylic acid	-0.10			
Sr(NO ₃) ₂	-0.05	0.015		
Mehlich-3	-0.40**	0.60**	0.25**	
DTPA	-0.10	0.25**	0.60**	0.45**
Mn				
carboxylic acid	0.75**			
Sr(NO ₃) ₂	0.05	0.05		
Mehlich-3	0.55**	0.75*	0.005	
DTPA	0.55**	0.75**	0.025	0.90**
Ni				
carboxylic acid	0.5**			
Sr(NO ₃) ₂	0.25**	0.30**		
Mehlich-3	0.30**	0.65**	0.40**	
DTPA	0.50**	0.50**	0.55**	0.50**

TABLE 6a

Samples	pH	Sequential extraction				Non-sequential extraction			Digest	
		Exch.	Mn-OX %	AM-Fe	CR-Fe	Carbox.	DTPA %	ML-3	AR	HF
									$\mu\text{g g}^{-1}$	
1	5.66	3.83	0.35	12.2	21.1	5.6	10.0	11.9	420	630
2	5.67	2.76	0.12	3.2	9.2	1.3	3.0	3.9	1250	1400
3	5.82	2.09	0.39	4.3	42.7	3.9	3.9	5.5	595	855
4	5.11	1.69	0.13	2.8	15.6	1.9	2.6	3.5	305	615
5	6.48	0.31	0.22	34.2	11.4	7.9	10.0	7.0	1650	1650
6	7.21	0.02	1.02	15.6	4.1	10.6	3.3	4.2	1900	2100
7	6.94	0.81	0.23	6.9	4.9	7.0	6.2	4.5	2550	1950
8	5.59	0.10	0.01	0.2	6.0	0.1	0.2	0.2	2500	2950
9	9.19	0.00	0.66	13.0	2.6	5.2	0.4	26.8	2300	1250
10	9.19	0.05	1.09	19.6	5.2	15.2	0.8	43.1	1300	1300
11	6.86	0.17	0.06	16.1	9.1	3.6	2.0	4.1	3400	3400
12	6.44	0.28	1.02	12.6	7.2	6.2	10.9	4.0	1600	1600
13	7.15	0.11	0.55	16.0	8.4	1.9	2.0	1.9	2850	2350
14	6.82	0.44	0.88	29.7	11.5	9.2	17.5	8.7	2500	2300

TABLE 6b

Samples	Sequential extraction				Non-sequential extraction			Digest	
	Exch.	Mn-OX	AM-Fe	CR-Fe	Carbox.	DTPA	ML-3	AR	HF
	%				%			$\mu\text{g g}^{-1}$	
1	0.006	0.001	2.7	22.8	0.25	0.25	0.09	2390	2740
2	0.012	0.003	5.8	13.8	0.76	0.76	1.46	425	1290
3	0.007	0.002	1.1	33.5	0.13	0.13	0.12	625	3040
4	0.006	0.001	1.2	24.1	0.13	0.13	0.05	4050	4800
5	0.000	0.001	2.4	17.8	0.14	0.14	0.24	2030	3730
6	0.017	0.001	1.8	18.4	0.13	0.13	0.04	1520	2440
7	0.002	0.001	1.3	14.6	0.04	0.04	0.03	3000	5600
8	0.001	0.003	0.3	8.4	0.09	0.09	0.01	12600	10600
9	0.000	0.004	6.1	6.3	1.22	1.22	0.02	2960	630
10	0.004	0.014	14.7	14.8	6.16	6.16	0.06	1280	600
11	0.006	0.001	1.0	14.6	0.06	0.06	0.05	1980	2360
12	0.001	0.000	0.4	10.5	0.03	0.03	0.02	6520	8660
13	0.001	0.000	1.4	13.9	0.04	0.04	0.04	2490	2030
14	0.003	0.000	0.5	8.5	0.02	0.02	0.04	5420	4610

TABLE 6c

Samples	Sequential extraction				Non-sequential extraction			Digest	
	Exch.	Mn-OX	AM-Fe	CR-Fe	Carbox.	DTPA	ML-3	AR	HF
		%				%		$\mu\text{g g}^{-1}$	
1	1.57	1.3	13.6	19.0	7.5	8.8	9.02	25	7
2	2.35	1.0	2.9	7.0	1.8	3.3	3.29	100	9
3	2.01	5.6	13.2	6.8	36.1	21.7	26.80	75	8
4	7.74	2.7	7.4	4.1	17.1	16.9	14.80	160	10
5	0.00	1.2	34.5	2.2	17.0	3.4	3.85	280	16
6	0.04	2.3	22.7	0.3	31.9	3.1	7.44	330	18
7	1.17	3.5	25.6	0.4	20.4	4.8	7.11	480	25
8	25.70	9.4	66.7	0.5	78.2	64.3	90.60	4	0
9	0.02	0.1	0.4	9.3	1.0	0.4	5.64	140	7
10	0.23	0.2	0.2	21.9	1.8	1.2	13.10	60	7
11	0.02	0.6	29.1	2.0	12.2	0.7	6.61	360	17
12	0.01	3.6	15.6	1.1	15.7	2.9	3.97	560	27
13	0.10	2.0	24.3	1.2	3.8	5.3	2.44	260	14
14	0.02	2.6	23.4	1.9	13.6	5.9	2.55	410	18

TABLE 6d

Samples	Sequential extraction				Non-sequential extraction			Digest	
	Exch.	Mn-OX	AM-Fe	CR-Fe	Carbox.	DTPA	ML-3	AR	HF
	%				%			$\mu\text{g g}^{-1}$	
1	2.4	1.2	14.8	7.9	3.8	3.5	3.9	320	305
2	10.4	2.8	20.3	4.8	4.9	7.3	9.8	340	635
3	6.3	5.7	6.8	2.5	32.3	18.0	25.0	1330	1220
4	19.6	3.9	7.0	2.8	20.9	17.1	21.1	2000	1520
5	0.1	4.2	26.8	1.4	22.4	4.7	4.7	3120	3090
6	0.0	7.0	17.9	0.2	46.6	5.6	12.2	3300	4150
7	3.5	3.9	10.5	0.4	21.6	5.9	8.1	7540	4570
8	3.2	0.4	0.8	0.7	2.6	1.2	3.3	1560	1590
9	<LOD	0.5	0.4	0.4	1.3	<LOD	6.7	1950	575
10	0.0	1.4	1.1	1.1	6.1	0.6	18.3	655	540
11	0.1	2.3	22.4	1.9	14.3	1.5	9.2	4380	3030
12	<LOD	4.3	10.6	0.8	14.2	5.3	4.5	7330	5760
13	<LOD	3.1	21.2	0.5	4.1	4.4	2.8	3640	2230
14	<LOD	3.8	17.7	1.4	12.9	7.8	3.1	5460	3880

Highlights

1. Extraction solution pH affects metal extractability in tropical ultramafic soils.
2. Nickel bearing-phases influence its phytoavailability.
3. The carboxylic acid method is suited for characterizing tropical ultramafic soils.
4. Ecological interpretation remains difficult without plant uptake data.

ACCEPTED MANUSCRIPT







Figure 1



a)**b)**

Figure 2

Projection of the variables on the factor-plane (1 x 2)




Figure 3

Projection of the variables on the factor-plane (1 x 2)




Figure 4




Figure 5