Metal-tolerant fungal communities are delineated by high zinc, lead, and copper concentrations in metalliferous Gobi Desert Soils

Article

Accepted Version

Kerfah, D., Ogwu, M. C., Dagvatseren, D., Balt, A., Davaasuren, D., Enkhmandal, O., Purevsuren, T., Batbaatar, A., Tibbett, M., Undrakhbold, S., Boldgiv, B. and Adams, J. M. (2020) Metal-tolerant fungal communities are delineated by high zinc, lead, and copper concentrations in metalliferous Gobi Desert Soils. *Microbial Ecology*, 79. pp. 420-431. ISSN 1432-184X doi: <https://doi.org/10.1007/s00248-019-01405-8> Available at <https://centaur.reading.ac.uk/84864/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s00248-019-01405-8>

Publisher: Springer

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Metal tolerant fungal communities are delineated by high zinc, lead and copper concentrations in metalliferous Gobi desert soils

Dorsaf Kerfahi¹, Matthew Chidozie Ogwu², Dagvatseren Ariunzaya³, Altantsetseg Balt⁴, Dulamsuren Davaasuren⁴, Orsoo Enkhmandal³, Tsolmonjav Purevsuren⁴, Amgaa Batbaatar^{4,5}, Mark Tibbett⁶, Sainbileg Undrakhbold³, Bazartseren Boldgiv^{3*}, Jonathan M. Adams^{7*}

¹School of Applied Biosciences, College of Agriculture and Life Sciences, Kyungpook National University, Daegu, 41566, Republic of Korea.

²School of Bioscience and Veterinary Medicine, University of Camerino, 62032 Camerino, Marche – Floristic Research Center of the Appennines, Gran Sasso and Monti della Laga National Park, San Colombo, 67021 Barisciano, L'Aquila, Italy.

³Ecology Group, Department of Biology, School of Arts and Sciences, National University of Mongolia, Ulaanbaatar 14201, Mongolia.

⁴Environmental Department, Oyu Tolgoi LLC, Oyu Tolgoi Site, South Gobi Province, Mongolia.

⁵Department of Biological Sciences, Faculty of Science, University of Alberta, Edmonton, Alberta, T6G 2E9 Canada.

⁶Centre for Agri-Environmental Research, School of Agriculture, Policy and Development, University of Reading, Reading RG6 6AR, United Kingdom.

⁷School of Geographic and Oceanographic Sciences, Nanjing University, Nanjing 210008, China.

*** Correspondence:**

Jonathan M. Adams: School of Geographic and Oceanographic Sciences, Nanjing University, Nanjing 210008, China. Tel: (+44) 7504 853655

23 Email: foundinkualalumpur@yahoo.com

24

25 **Bazartseren Boldgiv:** Ecology Group, Department of Biology, School of Arts and Sciences,

26 National University of Mongolia, Ulaanbaatar 14201, Mongolia. Tel/Fax: (976-11) 32 54 35

27 Email: boldgiv@num.edu.mn

28

29 **Conflict of interest:**

30 The authors declare no conflict s of interest.

Abstract

The soil fungal ecology of the southern Gobi region of Mongolia has been little studied. We utilized the ITS1 region from soil DNA to study possible influences on soil fungal community variation. In the sample network, a distinctive fungal community was closely associated with high zinc (Zn), lead (Pb) and copper (Cu) concentrations. The pattern of occurrence suggests that high metal concentrations are natural and not a product of mining activities. The metal-associated fungal community differs little from the ‘normal’ community in its major OTUs, and in terms of major fungal guilds and taxa, and its distinctiveness depends on a combination of many less common OTUs. The fungal community in the sites with high metal concentrations is no less diverse than in areas with normal background levels. Overall, these findings raise interesting questions of the evolutionary origin and functional characteristics of this apparently ‘metal tolerant’ community, and of the associated soil biota in general. It is possible that rehabilitation of metal-contaminated mined soils from spoil heaps could benefit from the incorporation of fungi derived from these areas.

Keywords: Fungi, Gobi desert, Heavy metals, Metagenetics, Soil metal, ITS1 region

1 Introduction

Heavy metals occur widely in the parent materials of soils at concentrations that are broadly dependent on lithology. During pedogenesis metals become enriched or depleted in different soil horizons, depending on the soil-forming environment and the predominant chemical reactions for each element [1]. Soil may be naturally enriched in metals due to the influence of the underlying geology and the subsequent soil forming and ecological processes, for example, through pedogenic activities in ultramafic and serpentinised geologies (e.g. [2]) or enrichment in the surface horizons following plant uptake from depth (e.g. for Cu and Pb see [3, 4]).

Soils with high concentrations of metals are associated with a distinctive microbial communities. This association is not in itself novel: there is abundant evidence of distinctive communities of plants and bacteria associated with metal-rich areas in other parts of the world. Examples include lead-, zinc- and [5] copper-rich mine spoil heaps in the UK and natural serpentine rock areas such as California and Borneo [5, 6]. Soil bacteria have been found to develop metal tolerance under laboratory conditions, by enabling energy dependent efflux of metal ions [7-9]. While metal-rich soils have occasionally been sampled by culturing individual fungal strains from them [10], there are few studies on whole metal-tolerant communities of soil fungi from anywhere in the world. Fungi are known to tolerate and detoxify metals by several mechanisms including valence transformation, extra- and intracellular precipitation and active uptake [11, 12]. Biological mechanisms implicated in fungal survival include extracellular precipitation, transformation of metals, biosorption to cell wall and pigments, decreased transport or impermeability, efflux, intracellular compartmentalization and sequestration [12-14].

The behaviour and occurrence of naturally occurring trace elements in the ecology of the soil are complex and the key metals that might influence microbial communities are poorly understood.

69 The present study was conducted in a naturally metal-rich area of the Gobi, a large cold desert in
70 southern Mongolia and northern China. The desert basins of the Gobi are bounded by the Altai
71 Mountains and the grasslands and steppes of Mongolia on the north, by the Tibetan Plateau to the
72 southwest, and by the North China Plain to the southwest. It is the fifth largest desert in the world
73 and the largest in Asia [15]. The climate is continental, characterized by dry and cold winters and
74 a precipitation maximum in summer [16]. Our sampling was located in Oyu Tolgoi region, which
75 is situated in the South Gobi desert, Mongolia. This area has only recently been discovered to be
76 rich in metals, especially copper [17-20]. Large scale mining for copper started in 2013, with the
77 opening of the Oyu Tolgoi mine [21]. Five major copper deposits that extend over 6 km in a north-
78 northeast-oriented zone. These occur in a middle to late Paleozoic arc terrain and are related to
79 Late Devonian quartz monzodiorite intrusions. The Hugo Dummett deposits are the northernmost
80 and deepest, with up to 1,000 m of premineral sedimentary and volcanic cover rock remaining [17].
81 The area is characterized by sparse vegetation and large tracts of Quaternary sediments and loess.
82 Ephemeral streams cross the area and flow for a few short periods during an average summer.
83 Temperatures at Oyu Tolgoi range from +36°C to -25°C. Total precipitation is approximately 100
84 mm/year and occurs mainly in late spring and early summer [22]. Given the potential ecological
85 sensitivity of this area, several months after the opening of Oyu Tolgoi we undertook a baseline
86 study of the area within a 130 km radius of the mine. Given the known concentration of natural
87 copper deposits in the southern Gobi, we specifically examined whether metal content is a key
88 factor in the structure and composition of soil fungal communities in this area. This paper mainly
89 aims to critically appraise apparent associations between metal-rich soils and fungal communities
90 in SE Gobi to characterize these communities, and to consider their broader implications for
91 microbial ecology and the study of metal enriched environments.

92

93 **2 Methodology**

94 **2.1 Site description**

95 The sampling area (Supplementary Fig. S1) is located in south-eastern Mongolia, close to the
96 border with China, between the latitudes of 42°31 N to 43°36 N and longitudes of 106°34 E to
97 108°10 E. Samples were taken in mid-September 2013, towards the end of the vegetation growing
98 season.

99 This study commissioned by the Oyu Tolgoi mine company was initially intended to establish
100 reference data for the background state of the soils in this area for future monitoring of any effects
101 of dust contamination, ground water alteration, or displacement of grazing in the area surrounding
102 the mine. For this baseline study of soil chemistry and biota, we sampled a network of 34 sites
103 (Supplementary Table S2) chosen to represent the range of natural vegetation types of the south-
104 eastern Gobi. Thirty four separate 1 ha plot sample sites were assigned within a radius of 130 km
105 of the Oyu Tolgoi mine, their positions chosen by selecting representative examples of different
106 habitat types found in the Gobi, based on both local knowledge and on satellite imagery.
107 Vegetation coverage consisted of small shrubs (mostly *Chenopodiaceae* and *Asteraceae*) and
108 bunch grasses, with overall plant coverage on the hectare scale varying from 50% to 70%
109 (Supplementary Table S1). Common representative plant species are *Eurotia ceratoides*,
110 *Potania mongolica* and *Caragana korshinskii*. In each sampling site, we took one subsample at
111 each corner of a hectare square and another in the center. Each subsample consisted of a core, 5
112 cm deep and 2 cm in diameter. The 5 subsamples were combined into one composite sample,
113 mixed thoroughly, and brought back to the laboratory on the same day. Half of the mixed sample

was frozen at -20°C for later DNA extraction, while half was dried for soil chemical and physical analysis.

In most of the 1 ha plots, vegetation composition was recorded, with a species inventory of all vascular plants present. Due to time limitations, we were unable to make a species inventory of four of the plots.

2.2 Chemical Analysis and DNA extraction

Soil analyses were carried out in the Laboratory of Soil Science of the Institute of Geography, Mongolian Academy of Sciences, using standard protocols of the Soil Science Association of America (SSSA) [23]. Measured chemical parameters were pH, CaCO₃, total organic carbon (TOC), soil salinity (measured as electric conductivity, dS/m), nitrogen (N), soil texture (TX), chrome (Cr), lead (Pb), cadmium (Cd), copper (Cu) and zinc (Zn). All samples were transported to the laboratory at Oyu Tolgoi mine and frozen at -20°C within 3 hours of sampling. Within days frozen soils were processed for DNA extraction in the Laboratory of Ecological and Evolutionary Synthesis of the National University of Mongolia. Soil was sieved with a 3 mm sieve, and 0.35 g of the sieved soil was DNA extracted using the Power Soil DNA extraction kit (MoBio Laboratories, Carlsbad, CA, USA) according to the manufacturer's protocol.

2.3 PCR and sequencing

All the extracted DNA samples were amplified for ITS1 region by using the primer pair ITSIF (5'-CTTGGTCATTTAGAGGAAGTAA-3') and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3') [24, 25]. Polymerase chain reactions (PCR) were performed in 50 µl reactions using the following

temperature program: 95°C for 10 min; 30 cycles of 95°C for 30 s, 55°C for 30 s, 72°C for 30 s; and 72°C for 7 min. The resulting amplicons were purified using the QIAquick PCR purification kit (Qiagen, CA, USA), and sequenced using Illumina Miseq platform (paired end of 2 × 300) at Macrogen Incorporation (Seoul, Korea), following standard Illumina sequencing protocols [26].

2.4 Sequence processing

Miseq Illumina [24] sequence data were processed using Mothur platform [27]. In brief, the sequence data were pair assembled using make.contigs and the quality control was performed by identifying chimeric sequence via UCHIME [28] and by filtering chimera sequences. Operational Taxonomic Units (OTUs) were picked using UCLUST [29] with a threshold of 97% pairwise identity using the QIIME implementation [30]. This cut-off has often been used in fungal community analyses [31, 32]. After the process, singletons were screened in Mothur. Taxonomic classification was assigned using the UNITE database [33] using the classify command in Mothur. All of the ITS1 sequence data are available under the MG-RAST [34] project ID 17045 (<http://metagenomics.anl.gov/linkin.cgi?project=17045>). The FUNGuild were used to classify OTUs into trophic categories [35].

2.5 Statistical analysis

To perform the statistical analysis, all samples were standardized by random subsampling to 13,957 reads per sample. We used a *t-test* for normal data and *Wilcoxon rank-sum test* for non-normal data in R software package 2.15.2 to test whether the relative abundances of taxonomical groups were significantly different between samples in low metals sites and high metals sites.

OTUs richness and diversity indices were estimated using Mothur. To assess the correlation between richness/diversity and environmental variables, linear regression was performed in SigmaPlot v 10.0 (Systat Software, San Jose, CA). To test for spatial autocorrelation of the fungal community OTU composition, we used the Mantel test (Mantel Nonparametric Test Calculator 2.0) [36] to compare matrices of fungal community composition in relation to geographic distance between different samples.

An indicator species analysis [37-39] was performed using package labdsv [40] in R software to identify those OTUs that are characteristics of each high and low metal-enriched sites. We used the INDVAL analysis, which identifies indicator species based on OTU fidelity and relative abundance [39]. Only OTUs with significant ($P < 0.05$) and INDVAL values that were > 0.5 were considered as significant threshold for habitat specialization. The indicator species are defined as the most characteristic species of each group, found mostly in a single group and present in the majority of the samples belonging to that group. Bray-Curtis distance was calculated to analyze fungal community similarity. To reduce the contribution of highly abundant OTUs in relation to less abundant ones in the calculation of Bray-Curtis matrix, abundance data of OTUs were square root transformed. The pairwise differences in fungal community composition were calculated by analysis of similarity (ANOSIM) in relation to Bray-Curtis distance. A constrained analysis was conducted using CANOCO [41] to assess the effect of environmental variables on the fungal community. Forward selection was used in redundancy analysis (RDA) [42] to select significant explanatory variables with 999 permutations and only significant variables ($P < 0.05$) were included in the models.

Fungal functional guilds were assigned according to Nguyen *et al.* [35] and Tedersoo *et al.* [43] using an open annotation tool (FUNGuild). Here, we only accepted the guild assignment where

confidence ranking was “probable” or “highly probable”, as recommended by Nguyen *et al.* [35]. The functional guilds of fungi detected in this study principally included three major functional groups: pathogens, saprotrophs and symbionts

3 Results

The soils in all sites were alkaline, (mean pH 8.7; range pH 8.2 – 9.3), and contained on average 56% of sand, 34% of silt and 11% of clay. Results of soil physico-chemical analysis are shown in Supplementary Table S2. In several samples, Zn metal concentrations exceeded EPA, Eco-SSL regulation (<http://www.epa.gov/ecotox/ecossl/index.html>) levels of 50 mg/kg, reaching 233 mg/kg in some samples. Cu and Pb contents were also above Eco-SSL, EPA level of 50 mg/kg in some samples, the highest metal concentrations measured were around 2300 mg/kg for Cu and 130 mg/kg for Pb (Fig. 1). In combination, the combined Pb, Zn and Cu concentration averaged around 689.78 mg/kg (ranging from 54.7 mg/kg to 3061 mg/kg).

From the 34 soil DNA samples, we obtained a total of 2,459,044 quality reads, which were classified into 11,559 OTUs at the 97% similarity level. In presentation of results, we have distinguished ‘high metal concentration’ samples as those containing >50 ppm Zn, Cu and/or Pb. The fungal community averaged across all the samples was dominated by Ascomycota, with 79% of total reads (Fig. 2a). Basidiomycota was the next most abundant group with 7% of the total reads. Relative abundance of other phyla was under 1%. There was no significant difference at the phylum level when high metal samples and low metal samples were compared, with the exception of certain minor phyla representing less than 1% of total reads - including Blastocladiomycota (w=77, P=0.02), Chytridiomycota (w=198, P=0.007), Incertae sedis (w=62, P=0.01) and Streptophycophyta (w=172, P=0.01).

Among the detected fungal classes, *Dothideomycetes* and *Sordariomycetes* (Ascomycota) were the most abundant, with a relative abundance of 48% and 25% of reads respectively amongst all samples combined. *Tremellomycetes* (Basidiomycota) represented 5% of total reads, and *Eurotiomycetes* (Ascomycota) and *Agaricomycetes* (Basidiomycota) were each at around 1% of total reads. Dominant orders were *Pleosporales* (46%), *Hypocreales* (21%), *Filobasidiales* (5%) and *Sordariales* (4%) (Fig. 2b).

As Fig. 2c shows, 9 families (These families included *Nectriaceae* (18%), *Pleosporaceae* (17%), *Pleosporales* family *Incertae sedis* (15%), *Sporormiaceae* (10%), *Filobasidiaceae* (5%), *Chaetomiaceae* (2%), *Leptosphaeriaceae* (2%), *Lasiosphaeriaceae* (1.5%) and *Hypocreales* family *Incertae sedis* (1%)) made up less than 1% of the total community. Community composition at the family level did not vary in relation to metal content.

The most abundant genus across all the samples combined was *Gibberella* from the family *Nectriaceae*, representing 18% of total reads (Fig. 2d). Metal content had no statistically significant effect on genus level composition of the community (Supplementary Table S3).

Linear regression analysis showed that metal content had no significant effect on fungal diversity. Similarly, the other soil parameters (pH, soil texture, etc) did not influence the fungal diversity (Supplementary Fig. S2). The effect of metal content on fungal richness and diversity was further evaluated using multiple regression analyses. Metal content (i.e. Zn, Pb and Cu) did not show any correlation with OTU richness and with diversity indices (Supplementary Fig. S3).

Fungal diversity in the high metal samples was no lower than the normal metal samples (Supplementary Fig. S2). The heat map analysis of the 50 most abundant OTUs did not show any consistent difference between high metal and low metals sites, despite the difference of metal concentrations (Supplementary Fig. S4).

Indicator species analysis revealed fungal OTUs that sort between low and high metal sites. The OTUs classified as core community in the low metal samples were represented by 25 genera, and characterized by the genera *Phoma* (represented by the species *Phoma bulgarica*, *Phoma calidophila*, and *Phoma sp P31E4*), *Preussia*, *Giberella*, etc. There were 30 genera classified as core community in high metal samples represented most abundantly by an unclassified fungal genus previously detected in eastern US forest soils by [44] under the name species *fungal sp. CC 06 28*. Also members of the genera *Cochliobolus*, *Curvularia* and *Chaetomium* are abundant examples of the core community of the metal rich sites (Table 1).

Vegetation cover ($w=44.5$, $P=0.33$) and plant species composition ($t=-2.005$, $df=6.65$, $P=0.08$) did not differ in relation to any measured soil characteristic among the 30 quadrat samples which had vegetation data (Supplementary Fig. S5). Multiple regression analysis showed that vegetation cover was not correlated with either Zn soil content, or Cu soil content ($P>0.05$), whereas it significantly correlated with Pb soil content (Supplementary Fig. S6).

An RDA (Fig. 3) showed that metal concentration (Zn, Cu, Pb individually, or all three combined) was the strongest predictor of variation in fungal community composition among our samples. Higher metal content samples (defined in Fig 3 as >200 mg/Kg of Zn) tended to cluster separately in terms of fungal community composition (Fig 3). Together with two axes on the biplot, in an accumulative variance for the interaction between communities and variables, a total of 19.3% of variation was explained. Axis 1 explained 12.2% of the variation in the data, while axis 2 explained 15.8. Among the measured physico-chemical factors, Zn (pseudo- $F=3.7$, $P=0.001$), Pb (pseudo- $=1.3$, $P=0.05$) and silt (pseudo- $F=2.0$, $P=0.001$) were significant contributors to fungal community variability, and a forward test indicated that the most important factor was Zn.

ANOSIM performed on Bray-Curtis community matrix confirmed that samples from low metal content sites and high metal content sites varied significantly from each other (Global $R=0.48$; $P=0.001$). The Mantel test showed no effect of spatial distance on the composition of fungal communities amongst the sites (Mantel statistic $r=0.091$, $P=0.15$).

Fungal taxonomic functional analysis by FUNGuild categorized the fungal sequences into different trophic modes. 34% of all reads were identified as pathotroph, followed by pathotroph-symbiotroph (23%), saprotroph (19%), pathotroph-saprotroph (7%), and less than 1% for symbiotroph, saprotroph-symbiotroph and pathogen-symbiotroph (Fig. 4a). In this functional study, there was only significant difference in pathogen-symbiotroph (that represents less than 1% of total trophic modes) between the normal and high rich metal samples. There was no significant difference in trophic strategy composition between samples having normal metal content and samples having high metal content (Fig. 4b).

4 Discussion

4.1 *Community characteristics and comparison with other arid environments*

The soil fungal community across all our sampled sites in the Gobi was dominated by Ascomycota, with a much lower abundance of Basidiomycota and other phyla (Fig. 2a). This bias towards Ascomycota is typical of arid environments globally – whereas Basidiomycota normally dominate in forest soils [27, 45-48]. Within the phylum Ascomycota, the most abundant family across all samples was *Nectriaceae* (18% of total reads), which includes a number of common pathogens, but also saprobes [49]. Within this family, the genera *Gibberella* and *Fusarium* which contain both plant pathogens and saprobes [50-54] were at 18% and 1% relative abundance, respectively (Fig. 2d). This again is typical of arid environments: for example *Fusarium* is generally one of the

commonest fungi in desert environments globally, and members have often been found to be abundant in the presence of metals [55, 56].

The family *Pleosporaceae* (Ascomycota) was also very abundant in our samples (17% of total reads). Members of this group are typically necrotrophic pathogens and saprobes, especially associated with grasses [49]. Within the family, the genera *Mycocentrospora* (11% of total reads) and *Alternaria* (10% of total reads) were particularly abundant in these Gobi sites. *Alternaria* has also been isolated from the metal-rich soils elsewhere [56, 57]. *Mycocentrospora* forms chlamydospores which have thick walls for surviving in extreme environments, a feature that may allow it to live in the very variable water environment of the Gobi [57].

The most abundant family of Basidiomycota in our samples was *Filobasidiaceae*, belonging to the order *Filobasidiales* and the class *Tremellomycetes*. Their mean relative abundance was 4.8%, and they made up most of the Basidiomycota in these samples (the Basidiomycota averaging in total at 7% of reads). Genus *Cryptococcus* under family *Filobasidiaceae* is also known to exhibit tolerance to Cu and Zn [48]. *Filobasidiaceae* have been isolated from Antarctic ecosystems and have a very wide range of habitats. The relative abundance of Basidiomycota was much less than in typical samples in forested or damper environments, but typical of semi-arid and arid locations: generally Basidiomycota are less abundant in hot desert environments [49].

In terms of trophic guilds from FUNGuild, pathotrophism was the most abundant category with 34% of total fungal reads, although this may reflect the difficulties of guild assignment in very diverse genera such as *Giberella* and *Fusarium* known to contain saprobes [58]. Pathotrophism was the second abundant trophic mode with 23% of total reads. Saprotriphism was the third most abundant category overall, at 19% of total reads, which agrees with the family level results discussed above. The spore-forming habit of many saprotrophs may allow their

survival in mostly dry soils without any physiological activity [58, 59]. The least abundant trophic category was saprotroph-symbiotroph, accounting for only 0.2% of reads (Fig. 4).

4.2 Community patterns in relation to metal content

In the RDA for the Gobi plots, the community divides very clearly into two clusters (Fig. 3). There is clear tendency in the RDA for the metal rich samples (average Zn concentration in metal samples of 225.44 mg/kg, 1500 mg/kg Cu and 70.39 mg/kg Pb) to cluster on one side of the ordination diagram, with variation in fungal community composition mainly related to metal concentrations (Fig. 3), and with Zn and Pb as the strongest predictors.

The RDA analysis shows a strong consistent pattern in relation to metals, especially Zn and Pb. Despite the clear differentiation of the fungal community by soil metal content on the RDA ordinations, at the broad taxonomic level the composition of the metal-poor and the metal-rich communities at the phylum, class and family level is very similar (Fig. 2). The same dominant genera are also found in all samples, both metal-rich and normal (Fig. 2d). Comparing the heat maps of metal-poor and metal-rich samples, there is no clearly evident community difference in terms of the 50 most abundant OTUs shown (Supplementary Fig. S4). The same major OTUs are present across both sample sets. Whatever the differences that lead to the high metal samples clustering separately, they presumably involve either the overall effect of many rarer OTUs, or consistent but subtle differences in the abundance of both common and rare OTUs. These OTUs include the ‘indicator’ taxa mentioned above.

It is of course necessary to ask if the close relationship we observed between metal concentrations and fungal community composition is merely spurious, with these factors inter-correlating with some other soil parameters that actually play the important role in determining soil fungal

community composition. Since a broad range of soil parameters was measured (Supplementary Table S2) and Zn, Pb and Cu were by far the best predictors of fungal community variation (Fig. 3), it seems unlikely that metal concentrations are merely a proxy for other soil parameters. This agrees with a generally accepted view that high metal concentrations may lead to alteration of soil microbial community structure [61, 62].

The ‘high’ metal concentrations (Zn, Pb and Cu) seen here in some of our samples are not exceptionally high compared to some contaminated sites studied in Europe which have had around 10 times this concentration [63]. However, in their experimental studies Smolders et al. (2004) observed effects of Zn on soil microbial (mostly bacterial) activity starting at concentrations around 200 mg/kg of Zn, similar those we observed, which suggests that at the concentrations found in the Gobi sites, microbial ecology could be significantly affected by Zn and other metals. A background of previous work also suggests that soil fungi are especially susceptible to high Zn concentrations, compared to bacteria. A study by Speir *et al.* [65] found that increasing Zn concentrations between 0 and 400 mg/kg had a significant negative impact on enzymatic activities of soil fungi (in our study the higher metal sites had around 200 mg/kg of Zn). Soil fungal communities have been found to be more responsive to Zn than soil bacteria are [60, 66]. Most of the samples that cluster at the higher end in terms of Zn concentrations greatly exceed environmental tolerance guidelines for Zn (EPA, Eco-SSL database: <http://www.epa.gov/ecotox/ecossl/index.html>, Supplementary Table S2) and the concentration limits reported in Kabata-Pendias [67] who stated that the threshold upper limit value of Cu in surface soils should be around 100 ppm, whereas our most Cu rich soils have concentrations of an average of 1475 ppm. They also noted that upper limit for the Pb content is around 70 ppm whereas our most Pb rich soils have an average concentrations of 66 ppm, and the mean Zn for worldwide

soils is around 64 ppm, whereas our most of our Zn soils have concentrations of an average of 224 ppm. This is certainly suggestive that toxicity may have a role in selecting the distinctive community that is found in these places. In some previous studies, soil fungi were found to be more sensitive to Cu than Zn and tended to have tolerance of Zn [68, 69].

It is also necessary to consider the possibility that the summative effect of multiple metals in the soil may be having the observed effect on fungal communities. It may also be important that the metals in our sampled soils tend to co-occur at higher concentrations. In recent years, there has been growing awareness of the interactive effects that multiple metals may have on soil ecology. This phenomenon, known as mixture toxicity, takes place where synergies between metals arise [70, 71]. Synergistic effects occur when the combined effect of two metals is greater than the sum of the effect of each metal individually. For synergistic interactions to occur in the soil, interacting metals have both to co-occur and to be present at concentrations high enough to induce the synergy [72, 73]. It is plausible, given that Zn, Pb and Cu in our soils are strongly co-occurring, there is a synergistic effect at work here in affecting the soil community.

An important proviso on the case that metal ions are a major direct factor in the fungal ecology of this area, however, is that the soil pH in the metal-rich areas is high, and that this is predicted to limit bioavailability of metal ions [74, 75]. In general – based on idealized laboratory observations of the chemistry and solubility of metal salts - each unit increase in pH is forecast to result in halving of available concentrations [75-78]. As the average pH of the soils which contained high metals was 8.66, having a pH range between 8.43 to 8.87 (Supplementary Table S2), the actual bioavailability would be predicted to be lower than in neutral or acidic soils [79, 80]. Exactly how much soil pH affects metal ion availability in actual soils is uncertain: Smolders *et al.* [63] found that over a range of pH from 4.5 to 7.5, Zn toxicity effects on soil microbes were

unaffected by pH, suggesting that the theoretical limits are inaccurate. All that one can really state is that while correlation suggests that Zn, Cu and Pb are affecting the fungal communities in the Gobi, traditional inorganic chemistry predicts that their effects are weaker than they would be if soil pH were more acidic. Protection against toxic effects of metals in soils can also occur through various other mechanisms: for example organic matter content, clay content and iron oxide content [78, 81-83], and it is possible that these too affect the biological availability of Zn and other metals in the soil.

4.3 Community patterns in relation to vegetation

Vegetation composition and percentage coverage did not correlate with fungal community composition in the sampled areas. Despite the apparent effects of these metals on the fungal community, the visual appearance of the Gobi ecosystem in the high metal areas does not suggest intense toxicity by metals. In the 30 samples which had vegetation data, the plant diversity and vegetation coverage of the most metal-rich 1 ha plots we obtained from the Gobi was no less than the 1 ha plots with normal metal concentrations, suggesting that: 1) the plants themselves have evolved local metal-tolerant ecotypes, 2) there might be evolutionary adaptation by the soil biota, perhaps in sequestering metal ions (and possibly making these metal ions unavailable to the plants in the process) or 3) due to high pH in limiting availability, the effects of high Zn or Cu or Pb concentrations are marginal in terms of plant growth and ecosystem function, despite a subtle effect that can be detected in ordination of the fungal community.

4.4 High diversity of the metal-rich soil fungal communities

386 Despite the high concentrations of metals in some of the soils, and the apparently dominant role
387 of metal concentrations in determining variation in fungal community structure, the diversity of
388 soil fungi in the metal-rich 1 ha plot samples was no less than in the local soils with normal
389 background levels of metals (Supplementary Fig. S2).

390 Extreme environments are usually seen as being associated with lower diversity [84]. This is the
391 case for example with soil bacterial communities of metal-polluted soils, and of extreme high and
392 low pH conditions [85-88]. However, in a previous study of the effects of application of sewage
393 sludge rich in metals, similarly high metal concentrations in soils still only showed minor effects
394 on microbial diversity [60].

395 In the system we are studying here, there seems to be no association between diversity and
396 'extreme' conditions (high Zn, Cu and Pb concentrations), for soil fungi at least. It is possible that
397 the high soil pH acts as a protectant against the worst effects of the metal ions on cell physiology
398 preventing the diversity-suppressing effects normally associated with an extreme environment.
399 Relatively low bioavailability could explain why fungal diversity is comparable with the other soil
400 samples with normal background concentrations of these metals, even though the metal
401 concentration is apparently having enough biological effect to make it a strong predictor of
402 community variation. An alternative, or additional, explanation for why these metal-rich soils
403 remain high in fungal diversity is that high metal concentrations in this area have existed for long
404 geological periods, allowing a very diverse soil fungal community to build up.

406 **4.5** *Are the high metal concentrations natural or influenced by anthropogenic activity?*

407 The source of the Zn and also the Cu and Pb found in many of the sites we sampled is unclear.
408 Most of our more metal-rich sites are found across a broad radius south-east of the Oyu Tolgoi

mine site, and the predominant wind direction in the area is north-west to south-east [89], which could imply that dust from the mine is the source. However, this seems unlikely, as the concentration of Zn, Cu and Pb does not show any relationship with distance from the mine, with some of the most metal rich samples being 20-80 km or more away from the mine, in areas never before mined. Scattered amongst the metal rich sites were also other sites that have normal background levels of metals, which would not be expected if an extensive metal-rich dust plume was spreading out across the desert. Also, the mine had only recently opened, producing its first ore in 2013 which was the year of sampling [21], which further implies that it is unlikely to have provided such extensive contamination of Zn, Cu and Pb over such a large area, and to have produced any noticeable changes in a fungal community averaged through the sampling depth of 0-10 cm . Additionally, a further set of very metal rich samples are found 30-40 km north-east of the mine, well upwind from the mine.

The most plausible explanation is that the high soil metal concentrations we observe here in the south eastern Gobi are natural, a consequence of the geological enrichment of rocks in this area with metal ores [90, 91], followed by weathering to form metal-rich soils.

5 Conclusions

The area around Oyu Tolgoi appears to be an unusual system, with naturally high soil concentrations of Zn, Cu and Pb in many areas [90, 91]. The strength of correlation suggests that Zn in particular dominates the community composition of soil fungi in this area, although Cu and Pb might also play an important part since they also tend to occur at high concentrations in the same Zn-rich areas. Despite the high metal concentrations, diversity of fungi in the metal-rich

431 areas is as high as in areas with normal background metal concentrations, which suggests the
432 possibility of a long history of specialized adaptation by the soil biota.

433 As a naturally metal-rich system, the soil ecology of Oyu Tolgoi area deserves further study. It
434 appears to offer a natural analog to anthropogenic metal-contaminated sites associated with
435 industrial activity around the world. Globally, there are very few known examples of naturally
436 metalliferous soil, the few exceptions being serpentine rock outcrops (rich in Ni), and Zn-rich sites
437 in central Europe [5, 92] – but even these are much more localized in extent than the Gobi area
438 that we studied. To our knowledge, no naturally metalliferous soil has ever been thoroughly
439 investigated from a microbial viewpoint. It would be interesting to know what (if any) special
440 adaptations the fungi in these metal-enriched soils have to the presence of metal ions, and whether
441 the distinctive community composition also extends to other soil organisms such as bacteria, soil
442 metazoans and archaea. Further studies should also include the soil metagenome, its
443 metatranscriptome, and soil properties such as potential soil respiration rate. Such aspects could
444 then be compared to those of anthropogenically contaminated sites, for potential lessons in terms
445 of the processes of community adaptation over time, and practical guidance for rehabilitation of
446 contaminated land.

447 **Acknowledgments**

449 This work was supported by the National Research Foundation (NRF) grant funded by the Korean
450 government, Ministry of Education, Science and Technology (MEST) (NRF 2013-031400). We
451 also received logistical support for field work and soil analytical services from the Oyu Tolgoi
452 mining company. DA, OE, SU, BB were also supported by the Taylor Family-Asia Foundation
453 Endowed Chair in Ecology and Conservation Biology.

455 **References**

- 456 1. Adriano DC (2001) ArsenicTrace elements in terrestrial environments. Springer, pp. 219-
457 261
- 458 2. Van Der Ent A, Cardace D, Tibbett M, Echevarria G (2018) Ecological implications of
459 pedogenesis and geochemistry of ultramafic soils in Kinabalu Park (Malaysia). *Catena* 160:
460 154-169.
- 461 3. Baker D, Senft J (1995) Copper (179-205) i Alloway. BJ (red) Heavy metals in.
- 462 4. Alloway B (1995) Cadmium, Ch. 6. Heavy Metals in Soils: London, Blackie: 100-124.
- 463 5. Hildebrandt U, Hoef-Emden K, Backhausen S, Bothe H, Božek M, Siuta A, Kuta E (2006)
464 The rare, endemic zinc violets of Central Europe originate from *Viola lutea* Huds. *Plant*
465 *Syst Evol* 257: 205-222.
- 466 6. Ernst WH (2006) Evolution of metal tolerance in higher plants. *For Snow Landsc Res* 80:
467 251-274.
- 468 7. Diaz-Ravina M, Baath E (1996) Development of metal tolerance in soil bacterial
469 communities exposed to experimentally increased metal levels. *Appl Environ Microbiol*
470 62: 2970-2977.
- 471 8. Hassen A, Saidi N, Cherif M, Boudabous A (1998) Resistance of environmental bacteria
472 to heavy metals. *Bioresour Technol* 64: 7-15.
- 473 9. Piotrowska-Seget Z, Cycoń M, Kozdroj J (2005) Metal-tolerant bacteria occurring in
474 heavily polluted soil and mine spoil. *Appl Soil Ecol* 28: 237-246.
- 475 10. Zarei M, Hempel S, Wubet T, Schäfer T, Savaghebi G, Jouzani GS, Nekouei MK, Buscot
476 F (2010) Molecular diversity of arbuscular mycorrhizal fungi in relation to soil chemical
477 properties and heavy metal contamination. *Environ Pollut* 158: 2757-2765.
- 478 11. Ashida J (1965) Adaptation of fungi to metal toxicants. *Annu Rev Phytopathol* 3: 153-174.
- 479 12. Gadd GM (1993) Interactions of fungip with toxic metals. *New Phytologist* 124: 25-60.
- 480 13. Ross I (1975) Some effects of heavy metals on fungal cells. *Trans Br Mycol Soc* 64: 175-
481 193.
- 482 14. Mehra RK, Winge DR (1991) Metal ion resistance in fungi: molecular mechanisms and
483 their regulated expression. *J Cell Biochem* 45: 30-40.
- 484 15. Magsar U, Baasansuren E, Tovvudorj M-E, Shijirbaatar O, Chinbaatar Z, Lkhagvadorj K,
485 Kwon O (2018) Medicinal plant diversity in the southern and eastern Gobi Desert region,
486 Mongolia. *J Ecol Environ* 42: 4.
- 487 16. An C-B, Chen F-H, Barton L (2008) Holocene environmental changes in Mongolia: a
488 review. *Global Planet Change* 63: 283-289.
- 489 17. Khashgerel B-E, Rye RO, Hedenquist JW, Kavalieris I (2006) Geology and reconnaissance
490 stable isotope study of the Oyu Tolgoi porphyry Cu-Au system, South Gobi, Mongolia.
491 *Econ Geol* 101: 503-522.
- 492 18. Panteleyev A, Eng P (2002) Report on the Geology and Mineral Potential of ‘Shivee
493 Tolgoi’Property, Southern Gobi Gold-Copper Belt, Mongolia. Private Report for Entrée
494 Resources Inc.
- 495 19. Wainwright AJ, Tosdal RM, Forster CN, Kirwin DJ, Lewis PD, Wooden JL (2011)
496 Devonian and carboniferous arcs of the oyu tolgoi porphyry Cu-Au district, South Gobi
497 region, Mongolia. *Geol Soc Am Bull* 123: 306-328.
- 498 20. Suzuki Y (2013) Conflict between mining development and nomadism in MongoliaThe
499 Mongolian Ecosystem Network. Springer, pp. 269-294

21. Miller J (2013) Oyu Tolgoi: A Nation Changer. *Eng Min J* 214: 68.
22. Kirwin D, Forster C, Kavalieris I, Crane D, Orssich C, Panther C, Garamjav D, Munkhbat T, Niislelkhoo G (2005) The Oyu Tolgoi copper-gold porphyry deposits, south Gobi, Mongolia. *Geodynamics and metallogeny of Mongolia with a special emphasis on copper and gold deposits SEG-IGOD field trip*: 14-16.
23. Topp GC, Reynolds WD, Green RE (1993) Advances in measurement of soil physical properties: Bringing theory into practice. *Soil Sci* 156: 429-432.
24. Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for basidiomycetes-application to the identification of mycorrhizae and rusts. *Mol Ecol* 2: 113-118.
25. White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR protocols: a guide to methods and applications* 18: 315-322.
26. Grada A, Weinbrecht K (2013) Next-generation sequencing: methodology and application. *J Invest Dermatol* 133: 1-4.
27. Schoch CL, Sung G-H, López-Giráldez F, Townsend JP, Miadlikowska J, Hofstetter V, Robbertse B, Matheny PB, Kauff F, Wang Z (2009) The Ascomycota tree of life: a phylum-wide phylogeny clarifies the origin and evolution of fundamental reproductive and ecological traits. *Syst Biol*: syp020.
28. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME improves sensitivity and speed of chimera detection. *Bioinformatics* 27: 2194-2200.
29. Edgar RC (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26: 2460-2461.
30. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N, Pena AG, Goodrich JK, Gordon JI (2010) QIIME allows analysis of high-throughput community sequencing data. *Nat Methods* 7: 335.
31. Nilsson RH, Kristiansson E, Ryberg M, Hallenberg N, Larsson KH (2008) Intraspecific ITS variability in the kingdom Fungi as expressed in the international sequence databases and its implications for molecular species identification. *Evol Bioinformatics* 4.
32. Tedersoo L, Nilsson RH, Abarenkov K, Jairus T, Sadam A, Saar I, Bahram M, Bechem E, Chuyong G, Kõljalg U (2010) 454 Pyrosequencing and Sanger sequencing of tropical mycorrhizal fungi provide similar results but reveal substantial methodological biases. *New Phytol* 188: 291-301.
33. Abarenkov K, Henrik Nilsson R, Larsson KH, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjoller R, Larsson E, Pennanen T (2010) The UNITE database for molecular identification of fungi—recent updates and future perspectives. *New Phytol* 186: 281-285.
34. Meyer F, Paarmann D, D'Souza M, Olson R, Glass EM, Kubal M, Paczian T, Rodriguez A, Stevens R, Wilke A (2008) The metagenomics RAST server—a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC bioinformatics* 9: 386.
35. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG (2016) FUNGuild: an open annotation tool for parsing fungal community datasets by ecological guild. *Fungal Ecol* 20: 241-248.
36. Liedloff A (1999) Mantel nonparametric test calculator. School of Natural Resource Sciences, Queensland University of Technology, Queensland, Australia.
37. Bakker JD (2008) Increasing the utility of indicator species analysis. *J Appl Ecol* 45: 1829-1835.

- 546 38. De Cáceres M, Legendre P, Moretti M (2010) Improving indicator species analysis by
547 combining groups of sites. *Oikos* 119: 1674-1684.
- 548 39. Dufrêne M, Legendre P (1997) Species assemblages and indicator species: the need for a
549 flexible asymmetrical approach. *Ecol Monogr* 67: 345-366.
- 550 40. Roberts DW (2007) labdsv: Ordination and multivariate analysis for ecology. R package
551 version 1.
- 552 41. Šmilauer P, Lepš J (2014) Multivariate analysis of ecological data using CANOCO 5.
553 Cambridge university press
- 554 42. Ter Braak CJ (1986) Canonical correspondence analysis: a new eigenvector technique for
555 multivariate direct gradient analysis. *Ecology* 67: 1167-1179.
- 556 43. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-
557 Palacios AM, Thu PQ, Suija A (2014) Global diversity and geography of soil fungi.
558 *Science* 346: 1256688.
- 559 44. Baird R, Stokes CE, Wood-Jones A, Alexander M, Watson C, Taylor G, Johnson K,
560 Remaley T, Diehl S (2014) Fleshy Saprobic and Ectomycorrhizal Fungal Communities
561 Associated with Healthy and Declining Eastern Hemlock Stands in Great Smoky
562 Mountains National Park. *Southeast Nat* 13: 192-219.
- 563 45. Buee M, Reich M, Murat C, Morin E, Nilsson RH, Uroz S, Martin F (2009) 454
564 Pyrosequencing analyses of forest soils reveal an unexpectedly high fungal diversity. *New*
565 *Phytol* 184: 449-456.
- 566 46. Matsushita Y, Bao Z, Kurose D, Okada H, Takemoto S, Sawada A, Nagase H, Takano M,
567 Murakami H, Koitabashi M (2015) Community structure, diversity, and species dominance
568 of bacteria, fungi, and nematodes from naturally and conventionally farmed soil: a case
569 study on Japanese apple orchards. *Organic Agriculture* 5: 11-28.
- 570 47. Wang H, Guo S, Huang M, Thorsten LH, Wei J (2010) Ascomycota has a faster
571 evolutionary rate and higher species diversity than Basidiomycota. *Science China Life*
572 *Sciences* 53: 1163-1169.
- 573 48. Vadkertiová R, Sláviková E (2006) Metal tolerance of yeasts isolated from water, soil and
574 plant environments. *J Basic Microbiol* 46: 145-152.
- 575 49. Cannon PF, Kirk PM (2007) Fungal families of the world. Cabi
- 576 50. Yoshida N, Sakai Y, Serata M, Tani Y, Kato N (1995) Distribution and properties of
577 fructosyl amino acid oxidase in fungi. *Appl Environ Microbiol* 61: 4487-4489.
- 578 51. Tudzynski B, Hedden P, Carrera E, Gaskin P (2001) The P450-4 gene of *Gibberella*
579 *fujikuroi* encodes ent-kaurene oxidase in the gibberellin biosynthesis pathway. *Appl*
580 *Environ Microbiol* 67: 3514-3522.
- 581 52. Tani Y, Mori N, Ogata K, Yamada H (1979) Production and Purification of Choline
582 Oxidase from *Cylindrocarpon didymum* M-1. *Agric Biol Chem* 43: 815-820.
- 583 53. Malonek S, Rojas M, Hedden P, Gaskin P, Hopkins P, Tudzynski B (2005) Functional
584 characterization of two cytochrome P450 monooxygenase genes, P450-1 and P450-4, of
585 the gibberellic acid gene cluster in *Fusarium proliferatum* (*Gibberella fujikuroi* MP-D).
586 *Appl Environ Microbiol* 71: 1462-1472.
- 587 54. Troncoso C, González X, Bömke C, Tudzynski B, Gong F, Hedden P, Rojas MC (2010)
588 Gibberellin biosynthesis and gibberellin oxidase activities in *Fusarium sacchari*, *Fusarium*
589 *konzum* and *Fusarium subglutinans* strains. *Phytochemistry* 71: 1322-1331.
- 590 55. Bhatnagar A, Bhatnagar M (2005) Microbial diversity in desert ecosystems. *Curr Sci* 89:
591 91-100.

56. Zafar S, Aqil F, Ahmad I (2007) Metal tolerance and biosorption potential of filamentous fungi isolated from metal contaminated agricultural soil. *Bioresour Technol* 98: 2557-2561.
57. Dix NJ (2012) *Fungal ecology*. Springer Science & Business Media
58. Cicccarone C, Rambelli A (2000) A study on micro-fungi in arid areas: notes on soil saprotrophs and animal opportunistic pathogens. *Plant Biosyst* 134: 25-29.
59. Sterflinger K, Tesei D, Zakharova K (2012) Fungi in hot and cold deserts with particular reference to microcolonial fungi. *Fungal Ecol* 5: 453-462.
60. Anderson IC, Parkin PI, Campbell CD (2008) DNA-and RNA-derived assessments of fungal community composition in soil amended with sewage sludge rich in cadmium, copper and zinc. *Soil Biol Biochem* 40: 2358-2365.
61. Frostegård Å, Tunlid A, Bååth E (1996) Changes in microbial community structure during long-term incubation in two soils experimentally contaminated with metals. *Soil Biol Biochem* 28: 55-63.
62. Bååth E, Díaz-Raviña M, Frostegård Å, Campbell CD (1998) Effect of metal-rich sludge amendments on the soil microbial community. *Appl Environ Microbiol* 64: 238-245.
63. Smolders E, Buekers J, Oliver I, McLaughlin MJ (2004) Soil properties affecting toxicity of zinc to soil microbial properties in laboratory-spiked and field-contaminated soils. *Environ Toxicol Chem: An Inter J* 23: 2633-2640.
64. Kim M, Boldgiv B, Singh D, Chun J, Lkhagva A, Adams JM (2013) Structure of soil bacterial communities in relation to environmental variables in a semi-arid region of Mongolia. *J Arid Environ* 89: 38-44.
65. Speir TW, Van Schaik AP, Hunter LC, Ryburn JL, Percival HJ (2007) Attempts to derive EC 50 values for heavy metals from land-applied Cu-, Ni-, and Zn-spiked sewage sludge. *Soil Biol Biochem* 39: 539-549.
66. Macdonald C, Singh B, Peck J, Van Schaik A, Hunter L, Horswell J, Campbell C, Speir T (2007) Long-term exposure to Zn-spiked sewage sludge alters soil community structure. *Soil Biol Biochem* 39: 2576-2586.
67. Kabata-Pendias A (2010) *Trace elements in soils and plants*. CRC press
68. Gadd GM (1994) Interactions of fungi with toxic metals. *The Genus Aspergillus*. Springer, pp. 361-374
69. Blaudez D, Jacob C, Turnau K, Colpaert J, Ahonen-Jonnarth U, Finlay R, Botton B, Chalot M (2000) Differential responses of ectomycorrhizal fungi to heavy metals in vitro. *Mycol Res* 104: 1366-1371.
70. Yang Z.X, Liu S.Q, Zheng D.W, Feng S.D (2006) Effects of cadmium, zinc and lead on soil enzyme activities. *J Environ Sci* 18: 1135-1141.
71. Wang Y-P, SHI J-y, Qi L, Chen X-c, Chen Y-x (2007) Heavy metal availability and impact on activity of soil microorganisms along a Cu/Zn contamination gradient. *J Environ Sci* 19: 848-853.
72. Cedergreen N (2014) Quantifying synergy: a systematic review of mixture toxicity studies within environmental toxicology. *PloS one* 9: e96580.
73. Wu X, Cobbina SJ, Mao G, Xu H, Zhang Z, Yang L (2016) A review of toxicity and mechanisms of individual and mixtures of heavy metals in the environment. *Environ Sci Pollut Res* 23: 8244-8259.
74. Rajapaksha R, Tobor-Kapłon M, Bååth E (2004) Metal toxicity affects fungal and bacterial activities in soil differently. *Appl Environ Microbiol* 70: 2966-2973.

75. Wang X, Wei D, Ma Y, McLaughlin MJ (2015) Derivation of soil ecological criteria for copper in Chinese soils. *PloS one* 10: e0133941.
76. Christensen TH (1984) Cadmium soil sorption at low concentrations: I. Effect of time, cadmium load, pH, and calcium. *Water Air Soil Pollut* 21: 105-114.
77. Appel C, Ma L (2002) Concentration, pH, and surface charge effects on cadmium and lead sorption in three tropical soils. *J Environ Quality* 31: 581-589.
78. Bradl HB (2004) Adsorption of heavy metal ions on soils and soils constituents. *J Colloid Inter Sci* 277: 1-18.
79. Munoz-Melendez G, Korre A, Parry S (2000) Influence of soil pH on the fractionation of Cr, Cu and Zn in solid phases from a landfill site. *Environ Pollut* 110: 497-504.
80. Wang Y, Shi J, Wang H, Lin Q, Chen X, Chen Y (2007) The influence of soil heavy metals pollution on soil microbial biomass, enzyme activity, and community composition near a copper smelter. *Ecotoxicol Environ Safety* 67: 75-81.
81. Smolders E, Oorts K, Van Sprang P, Schoeters I, Janssen CR, McGrath SP, McLaughlin MJ (2009) Toxicity of trace metals in soil as affected by soil type and aging after contamination: using calibrated bioavailability models to set ecological soil standards. *Environ Toxicol Chem* 28: 1633-1642.
82. Babich H, Stotzky G, Ehrlich H (1980) Environmental factors that influence the toxicity of heavy metal and gaseous pollutants to microorganisms. *CRC Crit Rev Microbiol* 8: 99-145.
83. Giller KE, Witter E, Mcgrath SP (1998) Toxicity of heavy metals to microorganisms and microbial processes in agricultural soils: a review. *Soil Biol Biochem* 30: 1389-1414.
84. Paul EA (2014) *Soil microbiology, ecology and biochemistry*. Academic press
85. Gans J, Wolinsky M, Dunbar J (2005) Computational improvements reveal great bacterial diversity and high metal toxicity in soil. *Science* 309: 1387-1390.
86. Shen C, Xiong J, Zhang H, Feng Y, Lin X, Li X, Liang W, Chu H (2013) Soil pH drives the spatial distribution of bacterial communities along elevation on Changbai Mountain. *Soil Biol Biochem* 57: 204-211.
87. Yanagawa K, Morono Y, de Beer D, Haeckel M, Sunamura M, Futagami T, Hoshino T, Terada T, Nakamura K-i, Urabe T (2013) Metabolically active microbial communities in marine sediment under high-CO₂ and low-pH extremes. *The ISME J* 7: 555-567.
88. Sorokin DY, Berben T, Melton ED, Overmars L, Vavourakis CD, Muyzer G (2014) Microbial diversity and biogeochemical cycling in soda lakes. *Extremophiles* 18: 791-809.
89. Zhibao D, Xunming W, Lianyou L (2000) Wind erosion in arid and semiarid China: an overview. *J Soil Water Conserv* 55: 439-444.
90. Khashgerel B-E, Kavalieris I, Hayashi K-i (2008) Mineralogy, textures, and whole-rock geochemistry of advanced argillic alteration: Hugo Dummett porphyry Cu–Au deposit, Oyu Tolgoi mineral district, Mongolia. *Mineralium Deposita* 43: 913-932.
91. Perelló J, Cox D, Garamjav D, Sanjdorj S, Diakov S, Schissel D, Munkhbat TO, Oyun G (2001) Oyu Tolgoi, Mongolia: siluro-devonian porphyry Cu-Au-(Mo) and high-sulfidation Cu mineralization with a cretaceous chalcocite blanket. *Econ Geol* 96: 1407-1428.
92. Brady KU, Kruckeberg AR, Bradshaw Jr H (2005) Evolutionary ecology of plant adaptation to serpentine soils. *Annu Rev Ecol Evol Syst*: 243-266.

681 **Figure legends**

682 **Fig. 1** Metal (Zn, Cu and Pb) contents in low metal samples and high metal samples. High metal
683 is defined by cutoff concentrations of 50 mg/kg in Zn, Pb and Cu.

684 **Fig. 2** Relative abundance of the detected fungal taxa observed in the Gobi samples (a) at the
685 phylum level, (b) class level, (c) Family level and (d) genus level.

686 **Fig. 3** Redundancy Analysis (RDA) ordination plot of fungal community composition based on
687 ITS1 gene OTUs and a vector overlay of the environmental variables. The significant
688 environmental variables were shown in red arrows. Red dots denote samples having low heavy
689 metal contents and blue dots denote samples having high heavy metal contents.

690 **Fig. 4** Detected fungal trophic mode by FUNGuild. (a) Classified trophic modes and (b) Relative
691 abundance of each trophic mode.

692 **Table legend**

693 **Table 1** Results of indicator species analysis.