Egg yolk phospholipids: a functional food material to generate deep-fat frying odorants

Article

Accepted Version

Chen, D.-W., Balagiannis, D. P. and Parker, J. K. (2019) Egg yolk phospholipids: a functional food material to generate deep-fat frying odorants. *Journal of the Science of Food and Agriculture*, 99 (14). pp. 6638-6643. ISSN 1097-0010 doi: <https://doi.org/10.1002/jsfa.9939> Available at <https://centaur.reading.ac.uk/85111/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1002/jsfa.9939>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Egg yolk phospholipids: a functional food material to generate**
2 **deep-fat frying odorants**

De-Wei Chen^{a,b,*}, Dimitrios P. Balagiannis^b, Jane K. Parker^b

^a Department of Food Science, Guangxi University, 530004, Nanning, Guangxi,
China

^b Department of Food and Nutritional Sciences, University of Reading, Whiteknights,
Reading RG6 6AP, UK

*Corresponding author: Dr De-Wei Chen, Department of Food Science, Guangxi
University, 530004, Nanning, Guangxi, China

Tel: 0086-(0)771-3237305; E-mail: chendw@gxu.edu.cn

4 **Abstract**

5 BACKGROUND

6 Phospholipids are an important precursor for the generation of carbonyl compounds
7 which play a significant role in the characteristic aroma of deep-fat fried foods.

8 RESULTS

9 Phospholipids extracted from hen egg yolks were added into sunflower oil (2.0 g/kg)
10 and heated with or without chicken meat at 160 °C for 10 min, and then dynamic
11 headspace extraction and GC-MS were used to extract and analyze the volatiles. The
12 results showed that the characteristic deep-fat frying odorants, such as
13 (*E,E*)-2,4-decadienal, (*E,Z*)-2,4-decadienal, as well as 1-octen-3-one, (*E*)-2-nonenal,
14 octanal, methional, dimethyl disulfide and alkylpyrazines had increased by 3-65
15 times in the sunflower oil with added phospholipids, and increased up to 6 times in
16 chicken meat which had been treated with phospholipids prior to heating.

17 CONCLUSION

18 There is potential for the food industry to use low levels of phospholipids,
19 particularly egg yolk phospholipids, to increase deep-fat frying odorants in a wide
20 range of deep fried products.

21 **Keywords:** deep-fat frying; odorants; phospholipids; egg yolk; chicken meat;
22 2,4-decadienal.

Introduction

Deep-fat frying is a popular cooking method, and the deep-fat frying aroma is very attractive for the consumer. Deep-fat frying aroma comes from the oxidation and decomposition of the lipids, as well as Maillard reaction, and it always occurs at high cooking temperatures.¹ (*E,E*)-2,4-Decadienal (deep-fried), (*E,Z*)-2,4-decadienal (deep-fried), pyrazines (roasty), 2- and 3-methylbutanal (malty), and methional (cooked potato-like odor) are the key odorants in fried potato chips.¹⁻³

Thuerer et al showed that the fatty acid [profile](#) of the frying oil significantly influenced the concentration of 2,4-decadienals.³ They found that the concentration of 2,4-decadienals in safflower oil was about 12 times higher than in coconut oil, as safflower oil is richer in linoleic acid, the main precursor of 2,4-decadienals. The concentration of the Strecker aldehydes and alkylpyrazines were not clearly influenced by the frying oils, but frying temperature is an important parameter as it has a great influence on their formation.⁴ Compared with conventional fried (at 165 °C) crisps, vacuum frying (at 120 °C) crisps reduced the formation of lipid-derived compounds and Maillard reaction products, especially alkylpyrazines.⁴

[Egg yolk phospholipids are rich in linoleic acid and arachidonic acid,](#)⁵ and it can be used as a source of important precursors for the generation of lipid-derived odorants.^{6,7} For example, hexanal, (*E,E*)-2,4-decadienal, 1-octen-3-one, (*E*)-2-decenal and (*E*)-2-undecenal were produced when egg yolk phospholipids were thermally treated at 145 °C for 20 min.⁶ In our previous work,⁷ the lipid-derived compounds contributing to the characteristic chicken meat aroma

increased sharply when egg yolk phospholipids were heated in water at 100 °C for 20 min, and chicken meat aroma was improved when egg yolk phospholipids were added to minced chicken meat prior to heating under the same conditions.

The hypothesis of this work is that egg yolk phospholipids can be the primary precursors of deep-fat frying odorants, which could be used specifically to increase the key odorants of deep-fried food. Although deep-fat frying odorants have been widely studied and produced by different methods, to the best of our knowledge, no research has been published where egg yolk phospholipids have been used in the deep-fried food. The objective of this work is to study the impact on deep-fried odorants when egg yolk phospholipids are added into sunflower oil and heated at frying temperatures with and without chicken meat.

Materials and methods

Reagents and Chemicals

Authentic chemicals of reference aroma compounds were purchased from a range of laboratory chemical suppliers in U.K.. 1,2-Dichlorobenzene in methanol (130.6 ng/ μ L) and alkane standard C5–C25 (100 ng/ μ L in diethyl ether), used as GC-MS standards, were obtained from Sigma-Aldrich Co. Ltd. (Gillingham, U.K.).

Phospholipids extraction

Commercial fresh hen eggs were bought from a local supermarket for this study. The extraction method employed was that reported by Chen et al.⁷ Briefly, fresh egg yolk (40 g, obtained from six hen eggs) lipids were extracted with ethanol; then the crude phospholipids were dissolved in hexane; next, cold acetone was used to

precipitate phospholipids; finally the high purity phospholipids were obtained. The minor residual solvents in the phospholipids were removed under high vacuum at room temperature for 10 h.

Phospholipids purity

This method was according to the AOCS Official Method.⁸ Firstly, the phospholipids sample was ashed in the presence of zinc oxide, followed by the spectrophotometric measurement of phosphorus as a blue phosphomolybdic acid complex, and finally a conversion factor of 30 was used to convert phosphorus to phospholipids.

Sample preparation

To disperse the phospholipids, 0.50 g phospholipids and 9.50 g salt were ground and thoroughly mixed. Commercial sunflower oil and fresh chicken breast fillets without skin or bone were bought from a local supermarket for this study. The chicken meat (~200 g) was ground in a domestic meat mincer (Kenwood, Havant, UK) and thoroughly mixed. The samples were prepared as follows:

- 1) Sunflower oil sample: 0.57 g salt, 15.0 g sunflower oil.
- 2) Sunflower oil & phospholipids sample: 0.60 g phospholipids and salt mixture, 15.0 g sunflower oil.
- 3) Chicken meat sample: 1.50 g chicken meat, 0.57 g salt, 15.0 g sunflower oil.
- 4) Chicken meat & phospholipids sample: 1.50 g chicken meat, 0.60 g phospholipids and salt mixture, 15.0 g sunflower oil.

Finally the samples were sealed in 100 mL glass Duran bottles and heated in an oil

89 bath (160 °C) for 10 min and then cooled in an ice-bath. Each treatment was carried
90 out in quadruplicate and all samples were prepared from the same batch of egg,
91 sunflower oil and chicken mince.

92 **Dynamic Headspace Extraction (DHE)**

93 DHE was used for the extraction of the volatiles, following the method described
94 by Chen et al with minor modifications.⁷ The entire contents of each Duran bottle
95 were transferred into a 250 mL conical flask fitted with a Dreschel head. The flask
96 was incubated in a water bath at 50 °C, and the volatiles in the headspace were swept
97 onto Tenax absorbent using a flow of nitrogen (40 mL/min) for 60 min. After
98 sweeping, 1.0 µL 1,2-dichlorobenzene in methanol (130.6 ng/µL) was added as an
99 internal standard to the trap, followed by a purge of 100 mL/min for 10 min to
100 remove excess solvent and moisture.

101 **GC-MS Analysis of Volatile Compounds**

102 The DHE samples were analysed using an Agilent 7890A-5975 GC-MS system
103 (Agilent Technologies Co. Ltd., Palo Alto, CA, USA) equipped with an automated
104 thermal desorber (Turbomatrix ATD), using a DB 5 column (60 m × 0.25 mm i.d., 1
105 µm film thickness from J&W Scientific, Agilent, Palo Alto, CA, USA) under
106 instrumental conditions described by Chen et al.⁷ The identification of the
107 compounds was based on the comparison of their mass spectra with spectra from the
108 NIST 11 Mass Spectral Database.⁹ The linear retention index (LRI) was calculated
109 for each volatile using the retention times of a series of C5–C25 n-alkanes. The
110 identities of most of the volatiles were confirmed if their mass spectra and LRI

matched those of authentic compounds run under the same analytical conditions in our laboratory. Volatiles were considered as tentatively identified by matching their mass spectra with the reference mass spectra in the NIST mass spectral library, and by comparison of their LRI to the NIST Chemistry WebBook database.¹⁰ Volatiles were semi-quantitatively determined by comparison of the peak areas against that of the internal standard using a response factor of 1 for each compound.

Statistical Analysis

The GC–MS data were analyzed using Independent-Samples T Tests (SPSS, 23, IBM) comparing the sunflower oil sample with the sunflower oil & phospholipids sample, and comparing the chicken meat sample with the chicken meat & phospholipids sample. Significant differences are shown by **, $p \leq 0.01$; *, $p \leq 0.05$; ns, $p > 0.05$.

Results and Discussion

The purity of the extracted phospholipids

The purity of the extracted egg yolk phospholipids was $95.2 \pm 3.0\%$, which was coincided with the report of Gladkowski et al.¹¹ According to their report, the purity of the phospholipids extracted by this method was very high (95.9%), and the phospholipids contained phosphatidylcholine (78%) and phosphatidylethanolamine (21%).

Characteristic deep-fat frying odorants derived from lipids

The results are shown in Table 1. The most abundant odorants detected were mainly aldehydes, such as (*E*)-2-heptenal, hexanal and (*E,E*)-2,4-decadienal.

133 (*E,E*)-2,4-Decadienal (fatty, fried), (*E,Z*)-2,4-decadienal (fatty, fried), 1-octen-3-one
 134 (mouldy, mushroom-like), (*E,E*)-2,4-nonadienal (fatty, fried, green), are the
 135 characteristic deep-fat frying odorants.¹ These compounds are formed from the
 136 thermal oxidation of unsaturated fatty acids, especially from the ω -6 fatty acids
 137 (linoleate and arachidonate).^{12, 13} Both (*E,E*)-2,4-decadienal and hexanal are often
 138 used as primary marker compounds of the oxidation of ω -6 fatty acids.¹⁴

139 Sunflower oil is widely used for deep-fat frying process,¹⁵ and it is rich in linoleic
 140 acid (about 62% of the total fatty acids),¹⁶ and the linoleate residues are present as
 141 triglycerides (as there are only trace phospholipids presented in the refined sunflower
 142 oil). Both the C9 and C13 position of the linoleate residues in triglycerides are
 143 favored positions to form the hydroperoxides during autoxidation,^{14, 17} and the C9
 144 and C13 hydroperoxide are the precursors for 2,4-decadienal and hexanal,
 145 respectively. So linoleate residues present in triglycerides can produce both
 146 2,4-decadienal and hexanal.

147 **Characteristic odorants derived from the Maillard reaction**

148 Alkylpyrazines, originating via the Maillard reaction, are important contributors to
 149 typical roasted, nutty, meaty, earthy, and popcorn-like note of many heated food
 150 products. The dominant alkylpyrazines in the present study were methylpyrazine and
 151 2,5-dimethylpyrazine. The Strecker aldehydes, for example, 2- and 3-methylbutanal
 152 (malty), methional (cooked potato-like odor), as well as phenylacetaldehyde (green,
 153 sweet), are the key odorants in fried potato chips, which are originated from
 154 isoleucine, leucine, methionine and phenylalanine, respectively. 2-Methylpropanal

(pungent) is formed from valine.¹⁸ Benzaldehyde (almond) is formed from phenylacetaldehyde,¹⁹ and methional may degrade to methanethiol, which in turn could produce dimethyl disulfide and dimethyl trisulfide.²⁰

Sunflower oil samples

The contents of hexanal and (*E,E*)-2,4-decadienal in the headspace collected from the heated sunflower oil sample were approximately 800 ng and 2 ng respectively. When the phospholipids were added to the sunflower oil prior to heating, hexanal increased by about 20% whereas the concentration of (*E,E*)-2,4-decadienal was 65 times higher than in the heated sunflower oil alone. As both (*E,E*)-2,4-decadienal and hexanal are the markers of oxidation, this shows that the egg yolk phospholipids under the present frying condition (160 °C, 10 min) were more prone to oxidation. It also demonstrates how the chemical environment of the linoleate residue influences the relative rates of formation of hexanal and (*E,E*)-2,4-decadienal.

The presence of phospholipids in the oil has two effects. Firstly it is clear that incorporation of the phospholipids increased the rate of oxidation. Owing to the amphiphilic nature of phospholipids, they tend to be present in a dispersed form, increasing their surface area, and increasing their exposure to water soluble pro-oxidants. In meat, these are likely to be pro-oxidant metals, rendering the phospholipids highly susceptible to oxidation and to the generation of large quantities of lipid-derived volatiles.²¹ Linoleic acid is the predominant PUFA (about 14% of the total fatty acids) in egg yolk phospholipids.⁵ Secondly, phospholipids direct the position of oxidation, promoting formation of the C9

hydroperoxide-derived volatiles in linolate residues over those derived from C13 hydroperoxides.²² It was reported that for linoleate residues bound in a phospholipid, autoxidation at the C9 position is the most favored position for formation of hydroperoxides.²² The C9 hydroperoxide is the precursor for 2,4-decadienals, so 2,4-decadienals are the dominant product for linoleate assembled in phospholipids. When the phospholipids were added to the sunflower oil sample, there was a sharp increase in (*E,E*)-2,4-decadienal (130 ng vs. 2 ng) and only a relatively small increase in hexanal (988 ng vs. 804 ng) which corroborates the finding of Reis and Spickett.²² In our previous study,⁷ we also observed a much greater increase in (*E,E*)-2,4-decadienal than in hexanal when phospholipids were added to water prior to a mild heating procedure.

Once the lipid oxidation process has been initiated by the more reactive fatty acids, this can promote the oxidation of the less reactive fatty acids.²³ The oxidised phospholipids not only produced lipid-derived aldehydes, but also promoted the oxidation of sunflower oil, explaining the increase in ω -3 and ω -9 derived volatiles in the sunflower oil & phospholipids sample.

The presence of some Maillard reaction products in the sunflower oil & phospholipids sample was not surprising, as low levels of Maillard reaction precursors are co-extracted with the phospholipids when polar solvents (ethanol and acetone) are used. In addition, both the primary amine group present in phosphatidylethanolamine and aldehydes produced from lipid oxidation can serve as Maillard reaction substrates.²¹ It is also consistent with the previous reports^{24, 25} that

the products of phospholipids oxidation can contribute to the Maillard reaction.

Sunflower oil samples with chicken meat

The trends in odorants in the heated chicken meat samples were consistent with those in the heated oil samples, but the magnitude of the increases were significantly diminished. Some lipid-derived aldehydes and most of the Maillard reaction products were significantly higher in the chicken meat & phospholipids sample compared to the chicken meat sample. The presence of meat provides a rich source of precursors and reactive intermediates which can influence the formation of volatiles and explain the changes in both lipid- and Maillard-derived compounds. In the chicken meat systems, the smaller increase in lipid-derived compounds with added phospholipids is consistent with previous findings.⁷ Examination of Table 1 shows that some odorants, such as 2,4-alkadienals, had much smaller increases when chicken meat was added to the samples. This apparent “loss” of 2,4-decadienals in the presence of meat could be attributed to the following. Firstly, the temperature was lower in the frying meat system than in the frying oil system (because of water evaporation), and the formation of those odorants were highly associated with the temperature.⁴ Secondly, those odorants may get trapped within the meat.²⁶ Thirdly, there may be interaction with other components of the meat (such as NH₃, H₂S, free amino groups and reactive dicarbonyls). For example 2,4-decadienal reacts with ammonia to form 2-pentylpyridine via Maillard reaction.²⁷ Finally, the degradation of this highly reactive alkadienal, such as 2,4-decadienal can undergo retroaldolisation to form hexanal and 2-octenal.²⁸ The higher productions of

2-pentylpyridine, 2-octenal and hexanal in the chicken meat & phospholipids sample could indicate more extensive degradation of 2,4-decadienal.

The small increase in 3-methylbutanal, and a similar trend for the other Strecker aldehydes, is due to the participation of unsaturated lipid-derived compounds in the Strecker degradation.²⁹ The parallel formation of α -aminoketones may also promote the formation of pyrazines. This increase is observed for 4 of the 5 pyrazines detected.

The key odorants in chicken & phospholipids sample

Both isomers of 2,4-decadienal not only impart characteristic deep-fried odor,¹ but were also the odorants with highest flavour dilution factor in chicken meat broths.³⁰ ³¹ (*E,E*)-2,4-Decadienal contributes a fatty chicken note with a very low odour detection threshold (0.07 $\mu\text{g/kg}$).³² In the present study, the contents of both isomers of 2,4-decadienal had a 3-4 times increase in the chicken meat & phospholipids samples, which indicated that the characteristic fatty fried chicken notes could arise from the increases in 2,4-decadienals. Besides (*E,E*)-2,4-decadienal, 1-octen-3-one, alkylpyrazines and the Strecker aldehydes are also the most significant contributors to deep-fried chicken meat flavor. Under the present processing conditions, the content of most of alkylpyrazines and the Strecker aldehydes had significantly increased, especially methional had a about 4 times increase in the phospholipids added chicken meat samples, which means that the fried meat note was enhanced by egg yolk phospholipids.

Conclusion

It has been demonstrated that the characteristic deep-fat frying odorants and fried chicken meat odorants, such as (*E,E*)- and (*E,Z*)- 2,4-decadienal, 1-octen-3-one, (*E*)-2-nonenal, octanal, methional, disulfide dimethyl, methylpyrazine and trimethylpyrazine significantly increased when a small amount (2.0 g/kg) of egg yolk phospholipids was added into sunflower oil samples with or without chicken present prior to heating at frying temperatures. This is of potential interest to the food industry and catering industry, however, its usage and effects on the deep-fat frying aroma need further sensory research, as food aroma is a delicate balance of the odorants.

Acknowledgements

This study was supported by the National Key R&D Program of China (2018YFD0901003) and China Scholarship Council (No. 201706665009).

Conflict of interest

There is no conflict of interest about this article.

References

1. Wagner R K and Grosch W, Key odorants of French fries. *J AM OIL CHEM SOC* 75:1385-1392 (1998).
2. Van Loon W, Linssen J, Legger A, Posthumus MA and Voragen A, Identification and olfactometry of French fries flavour extracted at mouth conditions. *FOOD CHEM* 90: 417-425 (2005).
3. Thuerer A and Granvogl M, Generation of Desired Aroma-Active as Well as

- 264 Undesired Toxicologically Relevant Compounds during Deep-Frying of Potatoes
265 with Different Edible Vegetable Fats and Oils. *J AGR FOOD CHEM* 64:
266 9107-9115 (2016).
- 267 4. Belkova B, Hradecky J, Hurkova K, Forstova V, Vaclavik L and Hajslova J, Impact
268 of vacuum frying on quality of potato crisps and frying oil. *FOOD CHEM* 241:
269 51-59 (2018).
- 270 5. Katz MA, Dugan LR and Dawson LE, Fatty acids in neutral lipids and
271 phospholipids from chicken tissues, *J FOOD SCI* 5: 717-720 (1966).
- 272 6. Lin JM and Blank I, Odorants generated by thermally induced degradation of
273 phospholipids. *J AGR FOOD CHEM* 51: 4364-4369 (2003).
- 274 7. Chen D, Balagiannis DP and Parker JK, Use of egg yolk phospholipids to generate
275 chicken meat odorants. *FOOD CHEM* 286: 71-77 (2019).
- 276 8. AOCS Official Method Ca 12-55. Phosphorus. Reapproved 2009. (2009).
- 277 9. NIST/EPA/MSDC. Mass Spectral Database (versions for PC and for Mass
278 Spectrometer Database Systems). National Institute of Standards and
279 Technology, Gaithersburg. (1992).
- 280 10. NIST Chemistry WebBook. NIST Standard Reference Database, SRD Number
281 69. <https://webbook.nist.gov/chemistry/>. (2017).
- 282 11. Gładkowski W, Chojnacka A, Kielbowicz G, Trziszka T, Wawrzenczyk C,
283 Isolation of Pure Phospholipid Fraction from Egg Yolk. *J AM OIL CHEM SOC*
284 89:179-182 (2012).
- 285 12. Zhou L, Zhao M, Bindler F and Marchioni E, Comparison of the Volatiles Formed

- 286 by Oxidation of Phosphatidylcholine to Triglyceride in Model Systems. *J AGR*
287 *FOOD CHEM* 62: 8295-8301 (2014).
- 288 13. Ladikos D and Lougovois V, Lipid oxidation in muscle foods: A review. *FOOD*
289 *CHEM* 35: 295-314 (1990).
- 290 14. Choe E and Min DB, Mechanisms and factors for edible oil oxidation, *COMPR*
291 *REV FOOD SCI* 5: 169-186 (2006).
- 292 15. Ekiz E and Oz F, The effects of different frying oils on the formation of
293 heterocyclic aromatic amines in meatballs and the changes in fatty acid
294 compositions of meatballs and frying oils. *J SCI FOOD AGR* 99: 1509-1518
295 (2019).
- 296 16. Dorni C, Sharma P, Saikia Gand Longvah T, Fatty acid profile of edible oils and
297 fats consumed in India. *FOOD CHEM* 238: 9-15 (2018).
- 298 17. Ho CT and Chen QY, LIPIDS IN FOOD FLAVORS - AN OVERVIEW, in *ACS*
299 *SYMPOSIUM SERIES*, eds. C. T. Ho and T. G. Hartman, AMER CHEMICAL
300 SOC, WASHINGTON pp. 2-14 (1994).
- 301 18. Whitfield FB and Mottram DS, Volatiles from interactions of Maillard reactions
302 and lipids. *CRIT REV FOOD SCI Nutr* 31: 1-58 (1992).
- 303 19. Chu FL and Yaylayan VA, Model Studies on the Oxygen-Induced Formation of
304 Benzaldehyde from Phenylacetaldehyde Using Pyrolysis GC-MS and FTIR. *J*
305 *AGR FOOD CHEM* 56: 10697-10704 (2008).
- 306 20. Mandin O, Duckham SC and Ames JM, Volatile compounds from potato-like
307 model systems. *J AGR FOOD CHEM* 47: 2355-2359 (1999).

- 308 21. Cui L and Decker EA, Phospholipids in foods: prooxidants or antioxidants?. *J SCI*
 309 *FOOD AGR* 96: 18-31 (2016).
- 310 22. Reis A and Spickett CM, Chemistry of phospholipid oxidation.
 311 *BBA-BIOMEMBRANES* 1818: 2374-2387 (2012).
- 312 23. Elmore JS, Mottram DS, Enser M and Wood JD, Effect of the polyunsaturated
 313 fatty acid composition of beef muscle on the profile of aroma volatiles. *J AGR*
 314 *FOOD CHEM* 47: 1619-1625 (1999).
- 315 24. Hidalgo FJ and Zamora R, Strecker-type degradation produced by the lipid
 316 oxidation products 4,5-epoxy-2-alkenals. *J AGR FOOD CHEM* 52: 7126-7131
 317 (2004).
- 318 25. Hidalgo FJ and Zamora R, Amino Acid Degradations Produced by Lipid
 319 Oxidation Products. *CRIT REV FOOD SCI Nutr* 56: 1242-1252 (2016).
- 320 26. Perez-Juan M, Flores M and Toldra F, Effect of pork meat proteins on the binding
 321 of volatile compounds. *FOOD CHEM* 108: 1226-1233 (2008).
- 322 27. Mottram DS, Flavour formation in meat and meat products: a review. *FOOD*
 323 *CHEM* 62: 415-424 (1998).
- 324 28. Frankel EN, Recent advances in lipid oxidation. *J SCI FOOD AGR* 54: 495-511
 325 (1991).
- 326 29. Gallardo E, De Schutter DP, Zamora R, Derdelinckx G, Delvaux FR and Hidalgo
 327 FJ, Influence of lipids in the generation of phenylacetaldehyde in wort-related
 328 model systems. *J AGR FOOD CHEM* 56: 3155-3159 (2008).
- 329 30. Feng Y, Cai Y, Fu X, Zheng L, Xiao Z and Zhao M, Comparison of aroma-active

- 330 compounds in broiler broth and native chicken broth by aroma extract dilution
331 analysis (AEDA), odor activity value (OAV) and omission experiment. *FOOD*
332 *CHEM* 265: 274-280 (2018).
- 333 31. Fan M, Xiao Q, Xie J, Cheng J, Sun B, Du W, Wang Y and Wang T, Aroma
334 Compounds in Chicken Broths of Beijing Youji and Commercial Broilers. *J AGR*
335 *FOOD CHEM* 66: 10242-10251 (2018).
- 336 32. Shi H and Ho CT, *The flavour of poultry meat. In Flavor of meat and meat*
337 *products (pp. 52-70)*, Springer, Boston, MA (1994).

338 Table 1. Mean Values (approx ng/sample extraction) (n=4) of the Volatile Compounds Identified in Headspace of the Samples.

			Sunflower oil	Sunflower oil & Phospholipids		Chicken meat	Chicken meat & Phospholipids	
Compound Name	LRI DB5 ¹	ID ²	Mean±SD ³	Mean±SD ³	Sig ⁴	Mean±SD ³	Mean±SD ³	Sig ⁵
ω-3 derivatives								
2-Hexenal, (<i>E</i>)-	857	A	112±19	310±57	**	106±48	116±51	ns
2,4-Heptadienal, (<i>E,Z</i>)-	1000	B	1.73±0.30	3.82±1.09	*	0.79±0.52	0.91±0.26	ns
2,4-Heptadienal, (<i>E,E</i>)-	1015	A	5.52±0.79	8.89±2.39	*	2.33±1.32	2.77±0.91	ns
5-Ethyl-1-cyclopentene-1-carboxaldehyde	1036	A	0.80±0.18	81.5±15.5	**	17.7±5.2	28.5±13.5	ns
ω-6 derivatives								
Pentane	500	A	59.9±35	282±99	*	53.3±12.2	69.2±32.9	ns
Pentanal	706	A	156±43	257±39	**	201±100	150±48	ns
1-Pentanol	765	A	23.7±2.2	47.6±7.2	**	191±60	214±86	ns
Hexanal	813	A	804±121	988±158	*	352±110	423±278	ns
1-Hexanol	869	A	8.07±1.14	10.5±2.6	ns	10.6±1.8	22.1±4.3	**
2-Heptanone	889	A	9.43±1.01	25.9±5.4	**	8.12±3.94	13.6±3.6	ns
Heptanal	904	A	35.6±3.5	135±36	**	36.2±21.6	53.5±14.9	ns
2-Heptenal, (<i>E</i>)-	966	A	655±260	1456±286	**	603±297	608±181	ns
1-Octen-3-one	978	A	10.2±3.7	55±22	**	11.8±4.5	22.9±4.6	*
1-Octen-3-ol	983	A	101±16	174±26	**	72.8±45.4	63.7±37.0	ns
2,3-Octanedione	984	A	1.42±0.28	33.9±8.6	**	4.48±2.25	9.54±4.11	*
2-Pentylfuran	990	A	16.7±3.6	33.2±13.5	*	9.71±7.35	13.1±6.9	ns

2-Octenal, (<i>E</i>)-	1060	A	25.8±3.2	145±48	**	40.1±27.9	57.2±18.5	ns
2-Octen-1-ol, (<i>E</i>)-	1067	A	1.49±0.50	7.59±1.44	**	3.15±0.89	4.37±1.61	ns
3-Nonen-2-one	1142	A	1.05±0.21	7.30±1.65	**	1.34±0.39	2.09±0.62	ns
2,4-Nonadienal, (<i>E,Z</i>)-	1200	B	0.27±0.05	1.15±0.36	**	0.35±0.20	0.25±0.11	ns
2,4-Nonadienal, (<i>E,E</i>)-	1222	A	0.79±0.12	4.22±1.31	**	1.11±0.59	1.33±0.68	ns
2,4-Decadienal, (<i>E,Z</i>)-	1299	B	1.20±0.20	47.92±26.53	*	3.29±1.36	10.4±3.6	*
2,4-Decadienal, (<i>E,E</i>)-	1325	A	2.01±0.83	130±70	*	5.68±1.91	22.4±7.6	*
ω-9 derivatives								
1-Octene	791	A	20.7±3.8	184±50	**	15.4±12.4	40.88±28.67	ns
Octane	800	A	43.2±6.9	147±32	**	73.0±23.1	70.6±20.6	ns
2-Octene, (<i>Z</i>)-	815	A	22.4±3.8	160±45	**	28.3±22.4	93.4±57.0	ns
Octanal	1004	A	16.4±1.8	59.0±19.0	**	18.5±6.2	28.5±6.7	ns
1-Octanol	1070	A	2.21±0.46	6.86±2.55	**	3.70±0.94	3.79±0.91	ns
Nonanal	1108	A	51.8±8.2	113±28	**	41.5±14.7	57.4±18.1	ns
2-Nonenal, (<i>E</i>)-	1164	A	2.79±0.39	10.9±2.1	**	2.66±0.64	5.14±1.29	*
1-Nonanol	1173	A	3.50±0.62	2.88±0.82	ns	2.94±0.67	2.73±0.52	ns
Decanal	1208	A	25.2±2.9	17.9±7.20	ns	22.8±6.5	22.3±1.4	ns
2-Decenal, (<i>E</i>)-	1266	A	3.60±0.81	17.4±7.3	**	5.42±1.59	8.21±2.79	ns
2-Undecenal	1368	A	3.55±1.35	3.11±1.5	ns	2.15±0.60	0.96±0.64	*
Maillard reaction products								
2-Methylpropanal	555	A	65.5±44.6	118±18	ns	45.8±11	74.8±20.0	ns
3-Methylbutanal	665	A	3.55±1.39	660.±90	**	420±65	664±138	**
2-Methylbutanal	674	A	2.50±0.75	275±50	**	144±9.8	170±29	ns
Methylpyrazine	829	A	0.17±0.11	7.78±2.38	**	8.41±0.64	17.4±5.6	**

Methional	912	A	nd	33.0±4.5	**	24.0±5.7	90.8±26.1	**
2,5-Dimethylpyrazine	917	A	nd	7.05±1.52	**	4.03±1.1	12.3±0.8	**
2-Ethylpyrazine	969	A	0.07±0.02	0.95±1.41	ns	0.12±0.23	0.18±0.35	ns
2-Ethy-6-methylpyrazine	1001	A	nd	0.35±0.14	**	0.34±0.16	0.69±0.01	*
Trimethylpyrazine	1006	A	1.60±1.09	6.79±4.12	*	0.19±0.11	1.05±0.26	**
Benzeneacetaldehyde	1049	A	3.72±1.86	17.3±4.8	**	36.7±4.1	102±25	**
2-Pentylpyridine	1204	B	0.04±0.02	2.20±0.63	**	0.05±0.01	0.44±0.24	*
Miscellaneous								
Dimethyl disulfide	745	A	0.05±0.00	14.3±10.9	*	12.1±1.7	30.9±9.9	*
Toluene	767	A	9.66±8.01	11.8±0.87	ns	10.1±2.7	25.6±28.5	ns
Acetylacetone	788	B	5.85±0.85	25.3±41.1	ns	3.35±1.58	16.±33	ns
3-Cyclohepten-1-one	826	C	4.52±0.51	8.47±16.8	ns	5.14±2.38	9.7±3.9	ns
2-Methylcyclopentanone	844	A	3.46±0.57	25.7±14.6	*	5.43±2.61	6.35±2.94	ns
α -Pinene	934	A	11.0±1.2	9.97±0.27	ns	8.26±2.68	5.02±1.51	ns
Benzaldehyde	971	A	42.4±18.8	25.0±4.0	ns	61.7±11.9	21.3±9.0	**
Dimethyl trisulfide	976	A	0.04±0.01	1.28±1.48	ns	14.2±6.23	14.1±2.7	ns
D-Limonene	1031	A	0.38±0.11	0.32±0.09	ns	10.1±4.3	14.0±7.8	ns
Undecane	1100	A	2.35±0.33	4.31±1.23	*	2.84±0.27	2.29±0.19	*
Dodecane	1199	A	1.11±0.38	1.02±0.17	ns	0.92±0.17	0.64±0.12	*
Undecanal	1309	A	2.26±0.34	1.02±0.41	**	2.07±0.64	1.47±0.19	ns
Tetradecane	1399	A	1.23±0.39	0.53±0.29	*	0.79±0.39	0.55±0.21	ns
Dodecanal	1411	A	2.24±0.40	0.75±0.40	**	2.38±1.14	1.06±0.04	*
Pentadecane	1499	A	1.05±0.27	0.44±0.32	*	1.07±0.73	0.36±0.03	ns

339 ¹Linear retention indices determined on a DB 5 column.

340 ²Confirmation of identity where A = mass spectrum and LRI agree with those of an authentic compound; B = mass spectrum agrees with
341 reference spectrum in the NIST mass spectral database and the LRI value agrees with that in the database (NIST Chemistry WebBook, 2017); C
342 = mass spectrum agrees with reference spectrum in the NIST mass spectral database (NIST/EPA/MSDC, 1992).

343 ³Approximate amount (mean, n=4) collected from the headspace, calculated by comparison of peak area with that of 1,2-dichlorobenzene
344 (130.6 ng) with a response factor of 1, nd = not detected.

345 ⁴Independent-Samples T Tests was used to compare group samples (sunflower oils sample vs. sunflower oils & phospholipids sample), significant
346 differences are shown by **, $p \leq 0.01$; *, $p \leq 0.05$; ns, $p > 0.05$.

347 ⁵Independent-Samples T Tests was used to compare group samples (chicken meat sample vs. chicken meat & phospholipids sample), significant
348 differences are shown by **, $p \leq 0.01$; *, $p \leq 0.05$; ns, $p > 0.05$.