

# *On certain sums concerning the gcd's and lcm's of k positive integers*

Article

Accepted Version

Hilberdink, T., Luca, F. and Toth, L. (2020) On certain sums concerning the gcd's and lcm's of k positive integers. *International Journal of Number Theory*, 16 (1). pp. 77-90. ISSN 1793-7310 doi: <https://doi.org/10.1142/S1793042120500049> Available at <http://centaur.reading.ac.uk/85149/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1142/S1793042120500049>

Publisher: World Scientific

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

International Journal of Number Theory  
© World Scientific Publishing Company

## On certain sums concerning the gcd's and lcm's of $k$ positive integers

Titus Hilberdink

*Department of Mathematics, University of Reading, Whiteknights  
PO Box 220, Reading RG6 6AX, UK  
t.w.hilberdink@reading.ac.uk*

Florian Luca

*School of Mathematics, University of the Witwatersrand  
Private Bag X3, WITS 2050, Johannesburg, South Africa  
and  
Research Group in Algebraic Structures and Applications  
King Abdulaziz University, Jeddah, Saudi Arabia  
and  
Department of Mathematics, Faculty of Sciences, University of Ostrava  
30 dubna 22, 701 03 Ostrava 1, Czech Republic  
florian.luca@wits.ac.za*

László Tóth

*Department of Mathematics, University of Pécs  
Ifjúság útja 6, H-7624 Pécs, Hungary  
ltoth@gamma.ttk.pte.hu*

Received (Day Month Year)  
Accepted (Day Month Year)

We use elementary arguments to prove results on the order of magnitude of certain sums concerning the gcd's and lcm's of  $k$  positive integers, where  $k \geq 2$  is fixed. We refine and generalize an asymptotic formula of Bordellès (2007), and extend certain related results of Hilberdink and Tóth (2016). We also formulate some conjectures and open problems.

*Keywords:* greatest common divisor; least common multiple; gcd-sum function; lcm-sum function; asymptotic formula; order of magnitude

Mathematics Subject Classification 2010: 11A25, 11N37

### 1. Introduction

Consider the gcd-sum function

$$G(n) := \sum_{k=1}^n (k, n) = \sum_{d|n} d\varphi(n/d) \quad (n \in \mathbb{N}),$$

where  $\varphi(n)$  is Euler's totient function. The function  $G(n)$  is multiplicative and the asymptotic formula

$$\sum_{n \leq x} G(n) = \frac{x^2}{2\zeta(2)} \left( \log x + 2\gamma - \frac{1}{2} - \frac{\zeta'(2)}{\zeta(2)} \right) + O(x^{1+\theta+\varepsilon}), \quad (1.1)$$

holds for every  $\varepsilon > 0$ , where  $\gamma$  is Euler's constant, and  $\theta$  is the exponent appearing in Dirichlet's divisor problem. See the survey paper [8] by the third author.

The function

$$G^{(-1)}(n) := \sum_{k=1}^n \frac{1}{(k, n)} = \sum_{d|n} \frac{\varphi(n/d)}{d} \quad (n \in \mathbb{N}),$$

is also multiplicative. Bordellès [1, Th. 5.1] deduced that

$$\sum_{n \leq x} G^{(-1)}(n) = \frac{\zeta(3)}{2\zeta(2)} x^2 + O\left(x(\log x)^{2/3}(\log \log x)^{4/3}\right). \quad (1.2)$$

The error term of estimate (1.2) comes from the classical result of Walfisz [9, Satz 1, p. 144],

$$R(x) := \sum_{n \leq x} \varphi(n) - \frac{1}{2\zeta(2)} x^2 = O\left(x(\log x)^{2/3}(\log \log x)^{4/3}\right). \quad (1.3)$$

We remark that recently (1.3) was improved by Liu [4] into

$$R(x) = O\left(x(\log x)^{2/3}(\log \log x)^{1/3}\right), \quad (1.4)$$

therefore, this serves as the remainder of (1.2). Also see the preprint by Suzuki [7].

The lcm-sum function

$$L(n) := \sum_{k=1}^n [k, n] = \frac{n}{2} \left( 1 + \sum_{d|n} d\varphi(d) \right) \quad (n \in \mathbb{N}).$$

was investigated by Bordellès [1], Ikeda and Matsuoka [3], and others. The function  $L(n)$  is not multiplicative and one has, see [1, Th. 6.3],

$$\sum_{n \leq x} L(n) = \frac{\zeta(3)}{8\zeta(2)} x^4 + O\left(x^3(\log x)^{2/3}(\log \log x)^{4/3}\right). \quad (1.5)$$

By using (1.4), the exponent of the  $\log \log x$  factor in the error of (1.5) can be improved into  $1/3$ .

Now let

$$L^{(-1)}(n) := \sum_{k=1}^n \frac{1}{[k, n]} \quad (n \in \mathbb{N}).$$

Bordellès [1, Th. 7.1] proved that

$$\sum_{n \leq x} L^{(-1)}(n) = \frac{1}{\pi^2} (\log x)^3 + A(\log x)^2 + O(\log x), \quad (1.6)$$

On certain sums concerning the gcd's and lcm's of  $k$  positive integers 3

with an explicitly given constant  $A$ .

By the general identity

$$\sum_{m,n \leq x} \psi(m, n) = 2 \sum_{n \leq x} \sum_{m=1}^n \psi(m, n) - \sum_{n \leq x} \psi(n, n),$$

valid for any function  $\psi : \mathbb{N}^2 \rightarrow \mathbb{C}$ , which is symmetric in the variables, (1.1), (1.2), (1.5) and (1.6), together with the remark on (1.4) lead to the asymptotic formulas

$$\sum_{m,n \leq x} (m, n) = \frac{x^2}{\zeta(2)} \left( \log x + 2\gamma - \frac{1}{2} - \frac{\zeta(2)}{2} - \frac{\zeta'(2)}{\zeta(2)} \right) + O(x^{1+\theta+\varepsilon}), \quad (1.7)$$

$$\sum_{m,n \leq x} \frac{1}{(m, n)} = \frac{\zeta(3)}{\zeta(2)} x^2 + O(x(\log x)^{2/3}(\log \log x)^{1/3}), \quad (1.8)$$

$$\sum_{m,n \leq x} [m, n] = \frac{\zeta(3)}{4\zeta(2)} x^4 + O(x^3(\log x)^{2/3}(\log \log x)^{1/3}), \quad (1.9)$$

and

$$\sum_{m,n \leq x} \frac{1}{[m, n]} = \frac{2}{\pi^2} (\log x)^3 + A_1 (\log x)^2 + O(\log x), \quad (1.10)$$

respectively, where  $A_1 = 2A$ .

It is easy to generalize (1.7) and (1.8) for sums with  $k$  variables by using the general identity

$$\sum_{n_1, \dots, n_k \leq x} f((n_1, \dots, n_k)) = \sum_{d \leq x} (\mu * f)(d) \lfloor x/d \rfloor^k,$$

where  $f$  is an arbitrary arithmetic function,  $\mu$  is the Möbius function and  $*$  stands for the Dirichlet convolution of arithmetic functions. For example, we have the next result: For any  $k \geq 3$ ,

$$\sum_{n_1, \dots, n_k \leq x} \frac{1}{(n_1, \dots, n_k)} = \frac{\zeta(k+1)}{\zeta(k)} x^k + O(x^{k-1}).$$

However, it is more difficult to derive asymptotic formulas for similar sums involving the lcm  $[n_1, \dots, n_k]$ . As corollaries of more general results concerning a large class of functions  $f$ , the first and third authors [2, Cor 1] proved that for any  $k \geq 3$  and any real number  $r > -1$ ,

$$\sum_{n_1, \dots, n_k \leq x} [n_1, \dots, n_k]^r = A_{r,k} x^{k(r+1)} + O(x^{k(r+1) - \frac{1}{2} \min(r+1, 1) + \varepsilon}) \quad (1.11)$$

and

$$\sum_{n_1, \dots, n_k \leq x} \left( \frac{[n_1, \dots, n_k]}{n_1 \cdots n_k} \right)^r = A_{r,k} x^k + O(x^{k - \frac{1}{2} \min(r+1, 1) + \varepsilon}),$$

where  $A_{k,r}$  are explicitly given constants. Here, (1.11) is the  $k$  dimensional generalization of (1.9). Furthermore, [2, Cor 2] shows that for any  $k \geq 3$  and any real number  $r > 0$ ,

$$\sum_{n_1, \dots, n_k \leq x} \left( \frac{[n_1, \dots, n_k]}{(n_1, \dots, n_k)} \right)^r = B_{r,k} x^{k(r+1)} + O\left(x^{k(r+1) - \frac{1}{2} + \varepsilon}\right),$$

with explicitly given constants  $B_{k,r}$ . The proofs use the fact that  $(n_1, \dots, n_k)$  and  $[n_1, \dots, n_k]$  are multiplicative functions of  $k$  variables and the associated multiple Dirichlet series factor over the primes into Euler products. The proofs given in [2] cannot be applied in the case  $r = -1$ .

It is the goal of the present paper to investigate the order of magnitude of the sums

$$S_k(x) := \sum_{n_1, \dots, n_k \leq x} \frac{1}{[n_1, \dots, n_k]}, \tag{1.12}$$

$$T_k(x) := \sum_{n_1, \dots, n_k \leq x} \frac{(n_1, \dots, n_k)}{[n_1, \dots, n_k]}, \tag{1.13}$$

$$U_k(x) := \sum_{\substack{n_1, \dots, n_k \leq x \\ (n_1, \dots, n_k) = 1}} \frac{1}{[n_1, \dots, n_k]}, \tag{1.14}$$

$$V_k(x) := \sum_{n_1, \dots, n_k \leq x} \frac{n_1 \cdots n_k}{[n_1, \dots, n_k]}, \tag{1.15}$$

where  $k \geq 2$  is fixed, by using elementary arguments. Theorem 2.1, concerning the sum  $S_2(x)$ , refines formulas (1.6) and (1.10) of Bordellès [1]. Theorems 2.3 and 3.1 give the exact order of magnitude of the sums  $S_k(x)$  and  $U_k(x)$ , respectively, for  $k \geq 3$ . Theorem 4.1 concerns the sums  $V_k(x)$ , while Theorem 5.2 provides an asymptotic formula with remainder term for  $T_k(x)$ , for any fixed  $k \geq 2$ . Some conjectures and open problems are formulated as well.

## 2. The sums $S_k(x)$

First consider the sums  $S_k(x)$  defined by (1.12). In the case  $k = 2$  we use Dirichlet's hyperbola method to prove the next result, which improves formulas (1.6) and (1.10).

### Theorem 2.1.

$$\sum_{n \leq x} L^{(-1)}(n) = \frac{1}{\pi^2} (\log x)^3 + A (\log x)^2 + B \log x + C + O\left(x^{-1/2} (\log x)^2\right), \tag{2.1}$$

that is,

$$\sum_{m, n \leq x} \frac{1}{[m, n]} = \frac{2}{\pi^2} (\log x)^3 + A_1 (\log x)^2 + B_1 \log x + C_1 + O\left(x^{-1/2} (\log x)^2\right),$$

On certain sums concerning the gcd's and lcm's of  $k$  positive integers 5

where the constants  $A, B, C$  can be explicitly computed, and  $A_1 = 2A$ ,  $B_1 = 2B - 1$ ,  $C_1 = C - \gamma$ .

**Proof.** We have

$$L^{(-1)}(n) = \sum_{k=1}^n \frac{(k, n)}{kn} = \frac{1}{n} \sum_{d|n} d \sum_{\substack{k=1 \\ (k, n)=d}}^n \frac{1}{k} = \frac{1}{n} \sum_{d|n} \sum_{\substack{t=1 \\ (t, n/d)=1}}^{n/d} \frac{1}{t} = \frac{1}{n} \sum_{d|n} h(d), \quad (2.2)$$

where

$$\begin{aligned} h(n) &:= \sum_{\substack{m=1 \\ (m, n)=1}}^n \frac{1}{m} = \sum_{m=1}^n \frac{1}{m} \sum_{d|(m, n)} \mu(d) = \sum_{d|n} \frac{\mu(d)}{d} \sum_{j=1}^{n/d} \frac{1}{j} \\ &= \sum_{d|n} \frac{\mu(d)}{d} \left( \log \frac{n}{d} + \gamma + O\left(\frac{d}{n}\right) \right) = \sum_{d|n} \frac{\mu(d)}{d} \log \frac{n}{d} + \gamma \frac{\varphi(n)}{n} + O\left(\frac{2^{\omega(n)}}{n}\right). \end{aligned}$$

Hence,

$$H(x) := \sum_{n \leq x} h(n) = \sum_{d \leq x} \frac{\mu(d)}{d} \sum_{m \leq x/d} \log m + \gamma \sum_{n \leq x} \frac{\varphi(n)}{n} + O\left(\sum_{n \leq x} \frac{2^{\omega(n)}}{n}\right).$$

By using the known estimates

$$\begin{aligned} \sum_{n \leq x} \log n &= x \log x - x + O(\log x), \\ \sum_{n \leq x} \frac{\varphi(n)}{n} &= \frac{6}{\pi^2} x + O(\log x), \\ \sum_{n \leq x} \frac{2^{\omega(n)}}{n} &= O((\log x)^2), \end{aligned}$$

we deduce that

$$\begin{aligned} H(x) &= (x \log x - x) \sum_{d \leq x} \frac{\mu(d)}{d^2} - x \sum_{d \leq x} \frac{\mu(d) \log d}{d^2} + \frac{6}{\pi^2} \gamma x + O((\log x)^2) \\ &= \frac{6}{\pi^2} (x \log x + cx) + O((\log x)^2), \end{aligned} \quad (2.3)$$

with a certain constant  $c$ . Let  $\mathbf{1}(n) = 1$  ( $n \in \mathbb{N}$ ), and let  $*$  denote the Dirichlet convolution. By Dirichlet's hyperbola method,

$$\begin{aligned} \sum_{n \leq x} (\mathbf{1} * h)(n) &= \sum_{n \leq \sqrt{x}} (H(x/n) + h(n)[x/n]) - [\sqrt{x}]H(\sqrt{x}) \\ &= \sum_{n \leq \sqrt{x}} H(x/n) + x \sum_{n \leq \sqrt{x}} \frac{h(n)}{n} - \sqrt{x}H(\sqrt{x}) + O(H(\sqrt{x})). \end{aligned}$$

6 *T. Hilberdink, F. Luca, L. Tóth*

By partial summation,

$$x \sum_{n \leq \sqrt{x}} \frac{h(n)}{n} = \sqrt{x}H(\sqrt{x}) + x \int_1^{\sqrt{x}} \frac{H(t)}{t^2} dt,$$

and using (2.3) we deduce

$$\begin{aligned} \sum_{n \leq x} (\mathbf{1} * h)(n) &= \frac{6}{\pi^2} \sum_{n \leq \sqrt{x}} \left( \frac{x}{n} \log \left( \frac{x}{n} \right) + c \left( \frac{x}{n} \right) \right) + \frac{6x}{\pi^2} \int_1^{\sqrt{x}} \left( \frac{\log t}{t} + c \right) \frac{dt}{t} + O(\sqrt{x}(\log x)^2) \\ &= x \left( \frac{3}{\pi^2} (\log x)^2 + a \log x + b \right) + O(\sqrt{x}(\log x)^2), \end{aligned}$$

for some constants  $a, b$ , which can be explicitly calculated.

Here  $(\mathbf{1} * h)(n) = nL^{(-1)}(n)$ , according to (2.2), and we obtain (2.1) by partial summation.  $\square$

It is more difficult to handle the sums  $S_k(x)$  in the case  $k \geq 3$ . We will apply the following general result proved by the second and third authors [5], using elementary arguments.

**Theorem 2.2.** ([5]) *Let  $k$  be a positive integer and let  $f : \mathbb{N} \rightarrow \mathbb{C}$  be a multiplicative function satisfying the following properties:*

- (i)  $f(p) = k$  for every prime  $p$ ,
- (ii)  $f(p^\nu) = \nu^{O(1)}$  for every prime  $p$  and every integer  $\nu \geq 2$ , where the constant implied by the  $O$  symbol is uniform in  $p$ .

Then

$$\sum_{n \leq x} \frac{f(n)}{n} = \frac{1}{k!} C_f (\log x)^k + D_f (\log x)^{k-1} + O((\log x)^{k-2}),$$

where  $C_f$  and  $D_f$  are constants,

$$C_f = \prod_p \left( 1 - \frac{1}{p} \right)^k \left( \sum_{\nu=0}^{\infty} \frac{f(p^\nu)}{p^\nu} \right).$$

We have the following result.

**Theorem 2.3.** *Let  $k \geq 3$  be a fixed integer. Then*

$$S_k(x) \asymp (\log x)^{2k-1} \quad \text{as } x \rightarrow \infty.$$

**Proof.** Since  $[n_1, \dots, n_k] \leq n_1 \cdots n_k \leq x^k$ , we can write

$$S_k(x) = \sum_{n \leq x^k} \frac{1}{n} \sum_{\substack{n_1, \dots, n_k \leq x \\ [n_1, \dots, n_k] = n}} 1 \tag{2.4}$$



On certain sums concerning the gcd's and lcm's of  $k$  positive integers 7

Let

$$a_k(n) := \sum_{\substack{n_1, \dots, n_k \in \mathbb{N} \\ [n_1, \dots, n_k] = n}} 1.$$

Now if  $n \leq x$ , then the inner sum in (2.4) is just  $a_k(n)$  (since  $n \leq x$  forces  $n_1, \dots, n_k \leq x$ ), while in any case it is at most  $a_k(n)$ . Thus

$$\sum_{n \leq x} \frac{a_k(n)}{n} \leq S_k(x) \leq \sum_{n \leq x^k} \frac{a_k(n)}{n}. \quad (2.5)$$

To see the properties of the function  $a_k(n)$  write

$$\sum_{d|n} a_k(d) = \sum_{d|n} \sum_{[n_1, \dots, n_k] = d} 1 = \sum_{[n_1, \dots, n_k] | n} 1 = \sum_{n_1 | n, \dots, n_k | n} 1 = \tau(n)^k.$$

Therefore, by Möbius inversion, we have  $a_k = \mu * \tau^k$ . This shows that  $a_k(n)$  is multiplicative and its values at the prime powers  $p^\nu$  are given by  $a_k(p^\nu) = (\nu + 1)^k - \nu^k$  ( $\nu \geq 1$ ). In particular,  $a_k(p) = 2^k - 1$ .

Applying Theorem 2.2 for the function  $f(n) = a_k(n)$ , with  $2^k - 1$  instead of  $k$ , we get that

$$\sum_{n \leq x} \frac{a_k(n)}{n} \sim \alpha_k (\log x)^{2^k - 1} \quad \text{as } x \rightarrow \infty, \quad (2.6)$$

for some constant  $\alpha_k$ . Now, from (2.5) and (2.6) the result follows.  $\square$

**Remark 2.4.** It is natural to expect that  $S_k(x) \sim c_k (\log x)^{2^k - 1}$  as  $x \rightarrow \infty$ , with a certain constant  $c_k$ . In fact, in view of Theorem 2.1, the plausible conjecture is that

$$S_k(x) = P_{2^k - 1}(\log x) + O(x^{-r}), \quad (2.7)$$

where  $P_{2^k - 1}(t)$  is a polynomial in  $t$  of degree  $2^k - 1$  and  $r$  is a positive real number. We pose as an open problem to find the constants  $c_k$  and to prove (2.7).

### 3. The sums $U_k(x)$

Next consider the sums  $U_k(x)$  defined by (1.14). In the case  $k = 2$ ,

$$U_2(x) \sim \frac{6}{\pi^2} (\log x)^2 \quad \text{as } x \rightarrow \infty,$$

and it is not difficult to deduce a more precise asymptotic formula.

We have the following general result.

**Theorem 3.1.** *Let  $k \geq 3$  be a fixed integer. Then*

$$U_k(x) \asymp (\log x)^{2^k - 2} \quad \text{as } x \rightarrow \infty.$$

8 *T. Hilberdink, F. Luca, L. Tóth*

**Proof.** Similar to the proof of Theorem 2.3. We have

$$U_k(x) = \sum_{\substack{n_1, \dots, n_k \leq x \\ (n_1, \dots, n_k) = 1}} \frac{1}{[n_1, \dots, n_k]} = \sum_{n \leq x^k} \frac{1}{n} \sum_{\substack{n_1, \dots, n_k \leq x \\ [n_1, \dots, n_k] = n \\ (n_1, \dots, n_k) = 1}} 1. \quad (3.1)$$

Let

$$b_k(n) = \sum_{\substack{n_1, \dots, n_k \in \mathbb{N} \\ [n_1, \dots, n_k] = n \\ (n_1, \dots, n_k) = 1}} 1.$$

Now if  $n \leq x$ , then the inner sum in (3.1) is exactly  $b_k(n)$ , while in any case it is at most  $b_k(n)$ . Thus

$$\sum_{n \leq x} \frac{b_k(n)}{n} \leq U_k(x) \leq \sum_{n \leq x^k} \frac{b_k(n)}{n}. \quad (3.2)$$

Write

$$\begin{aligned} \sum_{d|n} b_k(d) &= \sum_{d|n} \sum_{\substack{[n_1, \dots, n_k] = d \\ (n_1, \dots, n_k) = 1}} 1 = \sum_{\substack{[n_1, \dots, n_k] | n \\ (n_1, \dots, n_k) = 1}} 1 \\ &= \sum_{n_1 | n, \dots, n_k | n} \sum_{\delta | (n_1, \dots, n_k)} \mu(\delta) = \sum_{\delta a_1 b_1 = n, \dots, \delta a_k b_k = n} \mu(\delta) \\ &= \sum_{\delta t = n} \mu(\delta) \sum_{a_1 b_1 = t} 1 \cdots \sum_{a_k b_k = t} 1 = \sum_{\delta t = n} \mu(\delta) \tau(t)^k. \end{aligned}$$

Therefore, by Möbius inversion  $b_k = \mu * \mu * \tau^k$ . This shows that  $b_k(n)$  is multiplicative and its values at the prime powers  $p^\nu$  are given by  $b_k(p^\nu) = (\nu + 1)^k - 2\nu^k + (\nu - 1)^k$  ( $\nu \geq 1$ ). In particular,  $b_k(p) = 2^k - 2$ .

Applying now Theorem 2.2 for the function  $f(n) = b_k(n)$ , with  $2^k - 2$  instead of  $k$ , we deduce that

$$\sum_{n \leq x} \frac{b_k(n)}{n} \sim \alpha'_k (\log x)^{2^k - 2} \quad \text{as } x \rightarrow \infty \quad (3.3)$$

for some constant  $\alpha'_k$ . Now, from (3.2) and (3.3) we have  $U_k(x) \asymp (\log x)^{2^k - 2}$ .  $\square$

**Remark 3.2.** We conjecture that  $U_k(x) \sim d_k (\log x)^{2^k - 2}$  as  $x \rightarrow \infty$ , with a certain constant  $d_k$ . The sums  $S_k(x)$  and  $U_k(x)$  are strongly related. Namely, by grouping the terms according to the values  $(n_1, \dots, n_k) = d$  one obtains

$$S_k(x) = \sum_{d \leq x} \frac{1}{d} U_k(x/d), \quad (3.4)$$

and conversely,

$$U_k(x) = \sum_{d \leq x} \frac{\mu(d)}{d} S_k(x/d). \quad (3.5)$$

If  $U_k(x) \sim d_k(\log x)^{2^k-2}$  holds, then by (3.4) it follows that  $S_k(x) \sim \frac{d_k}{2^k-1}(\log x)^{2^k-1}$ . Conversely, assume that the asymptotic formula (2.7) is true, where  $c_k$  is the leading coefficient of the polynomial  $P_{2^k-1}(t)$ . Then (3.5), together with the well known results

$$\sum_{n \leq x} \frac{\mu(n)}{n} = O((\log x)^{-1}), \quad \sum_{n=1}^{\infty} \frac{\mu(n) \log n}{n} = -1,$$

and Shapiro's estimates [6, Th. 4.1]

$$\sum_{n \leq x} \frac{\mu(n)}{n} \left( \log \left( \frac{x}{n} \right) \right)^m = m(\log x)^{m-1} + \sum_{i=1}^{m-2} c_i^{(m)} (\log x)^i + O(1),$$

valid for any integer  $m \geq 2$ , where  $c_i^{(m)}$  are constants, imply that

$$U_k(x) = (2^k - 1)c_k(\log x)^{2^k-2} + b_{2^k-3}(\log x)^{2^k-3} + \dots + b_1 \log x + O(1),$$

with some constants  $b_i$ .

#### 4. The sums $V_k(x)$

The sums  $V_k(x)$  defined by (1.15) are sums of integers. In the case  $k = 2$  we have, according to (1.7),

$$V_2(x) = \sum_{m, n \leq x} (m, n) \sim \frac{6}{\pi^2} x^2 \log x. \quad (4.1)$$

**Theorem 4.1.** *Let  $k \geq 3$  be a fixed integer. Then*

$$x^k \ll V_k(x) \ll x^k (\log x)^{2^k-2} \quad \text{as } x \rightarrow \infty.$$

**Proof.** The lower bound is trivial by  $n_1 \cdots n_k \geq [n_1, \dots, n_k]$ . Also, by grouping the terms according to the values  $(n_1, \dots, n_k) = d$ , and by denoting  $M = \max(m_1, \dots, m_k)$  we have

$$\begin{aligned} V_k(x) &= \sum_{\substack{dm_1, \dots, dm_k \leq x \\ (m_1, \dots, m_k) = 1}} \frac{dm_1 \cdots dm_k}{[dm_1, \dots, dm_k]} = \sum_{\substack{m_1, \dots, m_k \leq x \\ (m_1, \dots, m_k) = 1}} \frac{m_1 \cdots m_k}{[m_1, \dots, m_k]} \sum_{d \leq x/M} d^{k-1} \\ &\ll x^k \sum_{\substack{m_1, \dots, m_k \leq x \\ (m_1, \dots, m_k) = 1}} \frac{m_1 \cdots m_k}{[m_1, \dots, m_k] M^k} \leq x^k \sum_{\substack{m_1, \dots, m_k \leq x \\ (m_1, \dots, m_k) = 1}} \frac{1}{[m_1, \dots, m_k]} = x^k U_k(x), \end{aligned}$$

and the upper bound follows from Theorem 3.1.  $\square$

10 T. Hilberdink, F. Luca, L. Tóth

**Remark 4.2.** We conjecture that  $V_k(x) \sim \lambda_k x^k (\log x)^{2^k - k - 1}$  as  $x \rightarrow \infty$ , with a certain constant  $\lambda_k$ , in accordance with (4.1) for the case  $k = 2$ . We pose as another open problem to prove this and to find the constants  $\lambda_k$ .

### 5. The sums $T_k(x)$

Finally, we investigate the sums  $T_k(x)$  defined by (1.13) and establish an asymptotic formula with remainder term for it. We give a short direct proof in the case  $k = 2$ . Then for any fixed  $k \geq 2$  we use multiple Dirichlet series to get the result.

Let

$$F(n) := \sum_{k=1}^n \frac{(k, n)}{[k, n]} \quad (n \in \mathbb{N}). \quad (5.1)$$

**Theorem 5.1.**

$$\sum_{n \leq x} F(n) = 2x + O((\log x)^2), \quad (5.2)$$

that is,

$$\sum_{m, n \leq x} \frac{(m, n)}{[m, n]} = 3x + O((\log x)^2).$$

**Proof.** Let  $\phi_2(n) = \sum_{d|n} d^2 \mu(n/d)$  be the Jordan function of order 2. We have

$$\begin{aligned} F(n) &= \sum_{k=1}^n \frac{(k, n)^2}{kn} = \frac{1}{n} \sum_{k=1}^n \frac{1}{k} \sum_{d|(k, n)} \phi_2(d) = \frac{1}{n} \sum_{d|n} \phi_2(d) \sum_{\substack{k=1 \\ d|k}}^n \frac{1}{k} \\ &= \frac{1}{n} \sum_{d|n} \frac{\phi_2(d)}{d} \sum_{j=1}^{n/d} \frac{1}{j} = \frac{1}{n} \sum_{d|n} \frac{\phi_2(d)}{d} H_{n/d}, \end{aligned}$$

where  $H_m = \sum_{j=1}^m 1/j$  is the harmonic sum. Therefore, using that

$$\sum_{n \leq x} \frac{\phi_2(n)}{n^2} = \frac{x}{\zeta(3)} + O(1),$$

we deduce

$$\begin{aligned} \sum_{n \leq x} F(n) &= \sum_{dm \leq x} \frac{\phi_2(d)}{d^2 m} H_m = \sum_{m \leq x} \frac{H_m}{m} \sum_{d \leq x/m} \frac{\phi_2(d)}{d^2} \\ &= \sum_{m \leq x} \frac{H_m}{m} \left( \frac{x}{\zeta(3)m} + O(1) \right) = \frac{x}{\zeta(3)} \sum_{m \leq x} \frac{H_m}{m^2} + O\left( \sum_{m \leq x} \frac{H_m}{m} \right) \end{aligned}$$

On certain sums concerning the gcd's and lcm's of  $k$  positive integers 11

$$\begin{aligned} &= \frac{x}{\zeta(3)} \sum_{m=1}^{\infty} \frac{H_m}{m^2} + O\left(x \sum_{m>x} \frac{H_m}{m^2}\right) + O\left(\sum_{m\leq x} \frac{H_m}{m}\right) \\ &= \frac{x}{\zeta(3)} \cdot 2\zeta(3) + O\left(x \sum_{m>x} \frac{\log m}{m^2}\right) + O\left(\sum_{m\leq x} \frac{\log m}{m}\right) = 2x + O((\log x)^2), \end{aligned}$$

by using that

$$\sum_{n=1}^{\infty} \frac{H_n}{n^2} = 2\zeta(3), \quad (5.3)$$

which is Euler's result.  $\square$

**Theorem 5.2.** *If  $k \geq 2$ , then*

$$T_k(x) = \beta_k x + O\left((\log x)^{2^k-2}\right),$$

where

$$\beta_k := \sum_{\substack{n_1, \dots, n_k=1 \\ (n_1, \dots, n_k)=1}}^{\infty} \frac{1}{[n_1, \dots, n_k] \max(n_1, \dots, n_k)} = \frac{1}{\zeta(2)} \sum_{n_1, \dots, n_k=1}^{\infty} \frac{1}{[n_1, \dots, n_k] \max(n_1, \dots, n_k)}.$$

**Proof.** By grouping the terms according to  $(n_1, \dots, n_k) = d$ , where  $n_j = dm_j$  ( $1 \leq j \leq k$ ),  $(m_1, \dots, m_k) = 1$ , we have

$$\begin{aligned} T_k(x) &= \sum_{\substack{dm_1, \dots, dm_k \leq x \\ (m_1, \dots, m_k)=1}} \frac{d}{[dm_1, \dots, dm_k]} = \sum_{\substack{dm_1, \dots, dm_k \leq x \\ (m_1, \dots, m_k)=1}} \frac{1}{[m_1, \dots, m_k]} \\ &= \sum_{\substack{m_1, \dots, m_k \leq x \\ (m_1, \dots, m_k)=1}} \frac{1}{[m_1, \dots, m_k]} \sum_{d \leq x/M} 1 = \sum_{\substack{m_1, \dots, m_k \leq x \\ (m_1, \dots, m_k)=1}} \frac{\lfloor x/M \rfloor}{[m_1, \dots, m_k]}, \end{aligned}$$

where  $M = \max(m_1, \dots, m_k)$ . Let

$$h(n_1, \dots, n_k) := \begin{cases} \frac{1}{[n_1, \dots, n_k]}, & \text{if } (n_1, \dots, n_k) = 1, \\ 0, & \text{otherwise.} \end{cases}$$

Hence,

$$T_k(x) = x \sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{\max(n_1, \dots, n_k)} + O\left(\sum_{n_1, \dots, n_k \leq x} h(n_1, \dots, n_k)\right) \quad (5.4)$$

and we estimate the right-hand sums in turn. Here  $h(n_1, \dots, n_k)$  is a symmetric and multiplicative function of  $k$  variables and for prime powers  $p^{\nu_1}, \dots, p^{\nu_k}$  ( $\nu_1, \dots, \nu_k \geq 0$ ) one has

$$h(p^{\nu_1}, \dots, p^{\nu_k}) = \begin{cases} \frac{1}{p^{\max(\nu_1, \dots, \nu_k)}}, & \text{if } \min(\nu_1, \dots, \nu_k) = 0, \\ 0, & \text{otherwise.} \end{cases}$$

12 *T. Hilberdink, F. Luca, L. Tóth*

Consider its Dirichlet series

$$H(s_1, \dots, s_k) := \sum_{n_1, \dots, n_k=1}^{\infty} \frac{h(n_1, \dots, n_k)}{n_1^{s_1} \cdots n_k^{s_k}} = \prod_p \sum_{\substack{\nu_1, \dots, \nu_k=0 \\ \min(\nu_1, \dots, \nu_k)=0}}^{\infty} \frac{1}{p^{\max(\nu_1, \dots, \nu_k) + \nu_1 s_1 + \cdots + \nu_k s_k}}.$$

By grouping the terms according to the values of  $r = \max(\nu_1, \dots, \nu_k)$  we deduce

$$H(s_1, \dots, s_k) = \prod_p \frac{1}{p^r} \sum_{r=0}^{\infty} \sum_{\substack{\nu_1, \dots, \nu_k=0 \\ \max(\nu_1, \dots, \nu_k)=r \\ \min(\nu_1, \dots, \nu_k)=0}} \frac{1}{p^{\nu_1 s_1 + \cdots + \nu_k s_k}},$$

which converges absolutely for  $\Re s_j > 0$  ( $1 \leq j \leq k$ ).

We shall need an estimate for  $H_k(\varepsilon, \dots, \varepsilon)$  for  $\varepsilon > 0$  (small). We have

$$H(\varepsilon, \dots, \varepsilon) = \prod_p \left( 1 + \frac{1}{p} \sum_{j=1}^{k-1} \binom{k}{j} \frac{1}{p^{j\varepsilon}} + O\left(\frac{1}{p^2}\right) \right).$$

Therefore,

$$\log H(\varepsilon, \dots, \varepsilon) = \sum_p \frac{1}{p} \sum_{j=1}^{k-1} \binom{k}{j} \frac{1}{p^{j\varepsilon}} + O(1) = \sum_{j=1}^{k-1} \binom{k}{j} \sum_p \frac{1}{p^{1+j\varepsilon}} + O(1).$$

But  $\sum_p p^{-1-\varepsilon} = \log \frac{1}{\varepsilon} + O(1)$  as  $\varepsilon \rightarrow 0$ . Thus,

$$H(\varepsilon, \dots, \varepsilon) = \exp \left( \sum_{j=1}^{k-1} \binom{k}{j} \log \frac{1}{\varepsilon} + O(1) \right) \asymp \left( \frac{1}{\varepsilon} \right)^{2^k - 2}. \quad (5.5)$$

Furthermore, for any  $\varepsilon > 0$ , we have

$$\begin{aligned} \sum_{n_1, \dots, n_k \leq x} h(n_1, \dots, n_k) &= \sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{(n_1 \cdots n_k)^{\varepsilon/k}} (n_1 \cdots n_k)^{\varepsilon/k} \\ &\leq x^\varepsilon \sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{(n_1 \cdots n_k)^{\varepsilon/k}} \leq x^\varepsilon H(\varepsilon/k, \dots, \varepsilon/k). \end{aligned} \quad (5.6)$$

Next, note that  $\max(n_1, \dots, n_k) \geq (n_1 \cdots n_k)^{1/k}$ , so that

$$\sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{\max(n_1, \dots, n_k)} \leq \sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{(n_1 \cdots n_k)^{1/k}} \leq H(\varepsilon/k, \dots, \varepsilon/k),$$

which converges. Hence,

$$\beta_k = \sum_{n_1, \dots, n_k=1}^{\infty} \frac{h(n_1, \dots, n_k)}{\max(n_1, \dots, n_k)}$$

is finite and  $\beta_k \leq H(\varepsilon/k, \dots, \varepsilon/k)$ . Also,

$$\beta_k - \sum_{n_1, \dots, n_k \leq x} \frac{h(n_1, \dots, n_k)}{\max(n_1, \dots, n_k)} = \sum_{\substack{n_1, \dots, n_k \in \mathbb{N} \\ \text{some } n_i > x}} \frac{h(n_1, \dots, n_k)}{\max(n_1, \dots, n_k)}$$

On certain sums concerning the gcd's and lcm's of  $k$  positive integers 13

$$\begin{aligned} &\leq k \sum_{\substack{n_1 \geq n_2, \dots, n_k \\ n_1 > x}} \frac{h(n_1, \dots, n_k)}{n_1} \leq k \sum_{\substack{n_1 \geq n_2, \dots, n_k \\ n_1 > x}} \frac{h(n_1, \dots, n_k)}{n_1^{1-\varepsilon} (n_1 n_2 \cdots n_k)^{\varepsilon/k}} \\ &\leq \frac{k}{x^{1-\varepsilon}} \sum_{n_1, \dots, n_k=1}^{\infty} \frac{h(n_1, \dots, n_k)}{(n_1 \cdots n_k)^{\varepsilon/k}} = kx^{\varepsilon-1} H(\varepsilon/k, \dots, \varepsilon/k). \end{aligned} \quad (5.7)$$

Hence, (5.4) and the estimates (5.6), (5.7) give

$$T_k(x) = \beta_k x + O(x^\varepsilon H(\varepsilon/k, \dots, \varepsilon/k)).$$

Now we choose  $\varepsilon = 1/\log x$  and use the bound (5.5). The proof is complete.  $\square$

**Remark 5.3.** For  $k = 2$ , Theorem 5.2 recovers Theorem 5.1. Note that

$$\beta_2 = \frac{1}{\zeta(3)} \sum_{m,n=1}^{\infty} \frac{1}{mn \max(m,n)} = \frac{2}{\zeta(3)} \sum_{m=1}^{\infty} \frac{1}{m^2} \sum_{n=1}^m \frac{1}{n} - 1 = 3,$$

by Euler's result (5.3). Is it possible to evaluate the constants  $\beta_k$  for any  $k \geq 2$ ?

The sums  $T_k(x)$  and  $U_k(x)$  are related by the formulas

$$T_k(x) = \sum_{d \leq x} U_k(x/d), \quad U_k(x) = \sum_{d \leq x} \mu(d) T_k(x/d).$$

## Acknowledgments

Part of the work in this paper was done when both the second and third authors visited the Max Plank Institute for Mathematics Bonn, in February 2017. They thank this institution for hospitality and a fruitful working environment. F. L. was also supported by grant CPRR160325161141 and an A-rated scientist award both from the NRF of South Africa and by grant no. 17-02804S of the Czech Granting Agency. L. T. was also supported by the European Union, co-financed by the European Social Fund EFOP-3.6.1.-16-2016-00004.

## References

- [1] O. Bordellès, Mean values of generalized gcd-sum and lcm-sum functions, *J. Integer Seq.* **10** (2007), Article 07.9.2, 13 pp.
- [2] T. Hilberdink and L. Tóth, On the average value of the least common multiple of  $k$  positive integers, *J. Number Theory* **169** (2016), 327–341.
- [3] S. Ikeda and K. Matsuoka, On the lcm-sum function, *J. Integer Seq.* **17** (2014), Article 14.1.7, 11 pp.
- [4] H.-Q. Liu, On Euler's function, *Proc. Roy. Soc. Edinburgh Sect. A* **146** (2016), 769–775.
- [5] F. Luca and L. Tóth, The  $r$ th moment of the divisor function: an elementary approach, *J. Integer Seq.* **20** (2017), Article 17.7.4, 8 pp.
- [6] H. N. Shapiro, On a theorem of Selberg and generalizations, *Ann. Math.* **51** (1950), 485–497.

14 *T. Hilberdink, F. Luca, L. Tóth*

- [7] Y. Suzuki, On error term estimates á la Walfisz for mean values of arithmetic functions, Preprint, 2018, 32 pp., arXiv:1811.02556 [mathNT].
- [8] L. Tóth, A survey of gcd-sum functions, *J. Integer Seq.* **13** (2010), Article 10.8.1, 23 pp.
- [9] A. Walfisz, *Weylsche Exponentialsummen in der neueren Zahlentheorie*, Mathematische Forschungsberichte, XV, VEB Deutscher Verlag der Wissenschaften, 1963.