

*Prolonged exposure to manure from
livestock administered antibiotics
decreases ecosystem carbon-use
efficiency and alters nitrogen cycling*

Article

Accepted Version

Wepking, C., Badgley, B., Barrett, J. E., Knowlton, K. F., Lucas, J. M., Minick, K. J., Ray, P. P. ORCID: <https://orcid.org/0000-0001-8375-8279>, Shawver, S. E. and Strickland, M. S. (2019) Prolonged exposure to manure from livestock administered antibiotics decreases ecosystem carbon-use efficiency and alters nitrogen cycling. *Ecology Letters*, 22 (12). pp. 2067-2076. ISSN 1461-0248 doi: 10.1111/ele.13390 Available at <https://centaur.reading.ac.uk/85786/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/ele.13390>

Publisher: Wiley

copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 Title: Prolonged exposure to manure from livestock administered antibiotics
 2 decreases ecosystem carbon-use efficiency and alters nitrogen cycling

3
 4 Authors: Carl Wepking^{a,b,*}, Brian Badgley^c, John E. Barrett^b, Katharine F.
 5 Knowlton^d, Jane M. Lucas^e, Kevan Minick^f, Partha P. Ray^g, Sarah Shawver^c,
 6 Michael S. Strickland^{e*}

7
 8 ^a School of Global Environmental Sustainability, Colorado State University, Fort
 9 Collins, CO, 80523; ^b Department of Biological, Virginia Tech, Blacksburg, VA,
 10 24061; ^c Sciences Department of Crop and Soil Environmental Sciences, Virginia
 11 Tech, Blacksburg, VA, 24061; ^d Department of Dairy Science, Virginia Tech,
 12 Blacksburg, VA, 24060; ^e Department of Soil and Water Systems, University of
 13 Idaho, Moscow, ID, 83844; ^f Department of Forestry and Environmental
 14 Resources, NC State University, Raleigh, NC, 27695; ^g Animal, Dairy and Food
 15 Chain Sciences, School of Agriculture, Policy and Development, University of
 16 Reading, Reading RG6 6AR

17
 18 ORCID: CW: 0000-0003-1910-5999; MSS: 0000-0001-5349-0363

19
 20 Keywords: elemental cycles, microbial ecology, stable isotopes, ecosystem
 21 function, antibiotics, agroecology

22
 23 Authorship Statement: C.W. and M.S.S. designed research. C.W., B.B., J.E.B.,
 24 K.F.K., K.M., P.P.R., S.S., and M.S.S. performed research. C.W., J.M.L and
 25 M.S.S. analyzed the data; C.W., and M.S.S. wrote the manuscript; all authors
 26 participated in editing and revision of the manuscript.

27
 28 Data Statement: The data that support the findings of this study are available at
 29 <https://datadryad.org/resource/doi:xxxxxx> (placeholder until this Dryad dataset is
 30 official).

31
 32 Number of words in abstract: 147
 33 Number of words in main text: 4954
 34 Number of figures: 4 (additionally, 3 supplementary figures
 35 and 5 supplementary tables)
 36 Number of references: 75 (plus 17 in supplementary materials)

37
 38 *Corresponding authors and contact information: 48
 39 49 Michael S. Strickland
 40 Carl Wepking 50 Department of Soil and Water Systems
 41 School of Global Environmental Sustainability – 51 University of Idaho
 42 Colorado State University 52 875 Perimeter Dr. MS 2340
 43 108 Johnson Hall 53 Moscow, ID 83844-2340
 44 Fort Collins, CO 80523 54 mstrickland@uidaho.edu
 45 carl.wepking@colostate.edu 55 phone: 208-885-0960
 46 phone: 970-491-7647 / fax: 970-492-4130

57 **Abstract:**

58

59 Microbial communities drive soil ecosystem function but are also susceptible to
60 environmental disturbances. We investigated whether exposure to manure
61 sourced from cattle either administered or not administered antibiotics affected
62 microbially-mediated terrestrial ecosystem function. We quantified changes in
63 microbial community composition, and terrestrial elemental cycling via a stable
64 isotope pulse-chase. Exposure to manure from antibiotic-treated cattle caused: *i)*
65 changes in microbial community structure; and *ii)* alterations in elemental cycling
66 throughout the terrestrial system. This exposure caused changes in
67 fungal:bacterial, as well as changes in bacterial community structure.
68 Additionally, exposure to manure from cattle treated with pirlimycin resulted in an
69 approximate two-fold increase in ecosystem respiration of recently fixed-carbon,
70 and a greater proportion of recently-added nitrogen in plant and soil pools
71 compared to the control manure. Manure from antibiotic-treated cattle therefore
72 affects terrestrial ecosystem function via the soil microbiome, causing decreased
73 ecosystem carbon use efficiency, and altered nitrogen cycling.

74

75 Introduction:

76

77 Use of antibiotics is under heightened scrutiny due to the increased prevalence
78 of antibiotic resistant pathogens (1–3). Antibiotic resistance is a multifaceted
79 problem and although the primary focus is on more stringent use of antibiotics in
80 medical settings, the use of antibiotics in the livestock sector is gaining increased
81 attention (4–13). In the United States, 80% of antibiotics are used in livestock
82 production, representing approximately 15-million kg of antibiotics annually (14,
83 15); globally livestock antibiotic use is projected to increase by 67% between
84 2010 and 2030 (4). After dosing, 40-90% of antibiotics are excreted by livestock
85 either intact or as a biologically active metabolite (16–18). Livestock manure
86 either collects in pastures or is applied to cultivated fields as fertilizer, therefore
87 potentially contributing up to 13-million kg of antibiotics to the environment
88 annually (14, 18). This widespread antibiotic exposure can affect human health
89 through the spread of antibiotic resistance, and also has the potential to directly
90 affect soil microbial communities and the ecosystem processes they regulate
91 (19–21).

92

93 The effect of antibiotics is an important consideration because microbial
94 communities are key drivers of ecosystem function. Soil microbial communities
95 play an important role in decomposition and elemental cycling in soils (22–25),
96 and impact the composition and productivity of plant communities (26) often
97 through beneficial and detrimental symbioses, and plant-microbe competition for
98 nutrients (27–31). While it is well known that soil microbes compete and signal
99 via antibiotics (32–34), the type and amount of antibiotics that soil microbial
100 communities are exposed to in agroecosystems are often novel and certainly
101 present in amounts far surpassing those found in soils naturally (8, 35).

102

103 Evidence is mounting that antibiotics can alter both soil microbial composition
104 through selection by antibiotic pressure, and physiology (21) through a stress
105 response (36) with the potential to affect ecosystem function. For instance, in
106 settings with known exposure to antibiotics microbial efficiency has been shown
107 to decrease, as evidenced by increased microbial mass-specific respiration with
108 a subsequent increase in the abundance of antibiotic resistance genes (21),
109 indicating that the metabolic costs associated with maintaining active antibiotic
110 resistance may reduce microbial efficiency (37). Antibiotic exposure has also
111 been shown to increase methane fluxes from manure (38). In addition to these
112 carbon (C) cycling effects, antibiotic exposure may also affect nutrient cycling.
113 Because production of microbial biomass is more demanding for nutrients (*e.g.*
114 nitrogen; N), the shift away from biomass production towards metabolic pathways
115 associated with a stress response could reduce microbial nutrient immobilization,
116 potentially increasing nutrient losses from ecosystems (36).

117 To investigate the potential effects of prolonged exposure to manure from
118 livestock treated with antibiotics (hereon these effects are referred to as antibiotic
119 effects) on microbial communities and ecosystem functioning, we applied manure
120 from three groups of cattle (those that received the bactericidal antibiotic

121 cephalosporin, those that received the bacteriostatic antibiotic pirlimycin, and control
122 cattle receiving no antibiotics) to grassland plots in a common-garden
123 experiment, along with a no-manure control. The relative impacts of antibiotics on
124 soil microbial communities were examined via determination of fungal:bacterial
125 ratio (hereon F:B) and 16S and ITS metabarcoding (to assess bacterial and
126 fungal community composition, respectively), and on ecosystem processes via a
127 ^{13}C and ^{15}N stable isotope pulse-chase. We expected that manure itself would
128 positively affect plant growth and lead to an increase in soil C pools. However,
129 when manure was sourced from cattle administered antibiotics, we expected a
130 greater loss of C via respiration, as well as, an overall decrease in ecosystem C-
131 use efficiency due to decreased microbial efficiency (specifically bacterial; 21).
132 Antibiotics are likely to lead to an increase in F:B (39, 40). The implications of this
133 on an ecosystem-scale are subject to debate: given the classical understanding
134 of fungal versus bacterial contribution to biogeochemical processes, we would
135 expect that systems with a higher F:B would retain more C and N (41, 42).
136 Alternatively, recent work has shown that C and N mineralization are unrelated to
137 the relative dominance of bacteria and fungi (43). Therefore, outcomes from this
138 experiment could lend support to either theory in light of recent challenges to the
139 classical understanding of fungal versus bacterial contribution to biogeochemical
140 processes.

141

142 **Materials and Methods:**

143

144 *Experimental Design:*

145

146 A common garden experiment with a randomized block design (four treatments,
147 $n=6$) was conducted at Kentland Farm, Blacksburg, VA, USA (37.199490, -
148 80.584659; 547-m elevation; Unison and Braddock cobbly soils; dominant plant
149 cover is grasses, mostly tall fescue, as well as some herbaceous cover including
150 members of the Lamiaceae and Plantaginaceae families). Treatments included
151 three manure additions (manure from cattle given no antibiotics, or manure from
152 cattle given either cephalosporin benzathine or pirlimycin hydrochloride) and one
153 control treatment that received no manure. Both antibiotics are commonly used in
154 the prevention of mastitis in dairy cattle, however they vary in a number of ways
155 including their fate in the environment (44) and mode of action. Cephalosporin
156 benzathine (Molecular weight = 365.4 g mol⁻¹; pKa = 2.2; water solubility = 3,430
157 mg L⁻¹) is bactericidal, damaging the structural integrity of bacterial cell
158 membranes, whereas pirlimycin hydrochloride (Molecular weight = 447.4 g mol⁻¹;
159 pKa = 8.4; water solubility = 64,900 mg L⁻¹) is bacteriostatic, inhibiting protein
160 synthesis. Hereon we refer to these four treatments as no-manure control (NMC),
161 control manure (Con), cephalosporin manure (Ceph), and pirlimycin manure (Pir).

162

163 Manure was applied to appropriate treatments at a monthly rate of 648-g-m⁻² of
164 wet-weight manure starting in October, 2014 until May, 2015 (213 days) –
165 totaling 4,536-g of manure m⁻². This amount of manure corresponds with the
166 amount of manure expected given a typical dairy cattle stocking density.

167
168 For information regarding manure properties, sourcing and within-manure
169 antibiotic quantification see supplementary materials.

170
171 *Pulse-chase experiment:*

172
173 Field sampling was conducted in May, 2015. In order to determine whether
174 antibiotic use in dairy cattle affects system-wide elemental cycling, a ^{13}C and ^{15}N
175 stable isotope pulse-chase experiment was conducted. The use of ^{13}C allowed
176 for the tracking of recently photosynthesized C through both above- and
177 belowground C pools. To accomplish ^{13}C -labeling, a $\sim 1\text{-m}^2$ subplot within each
178 treatment plot was covered with a 0.6-m^3 ($0.99\text{-m} \times 0.99\text{-m} \times 0.61\text{-m}$) transparent
179 acrylic chamber (Figure S1). To prevent gas exchange from outside the
180 chamber, the chamber was fitted into a rubber lined wooden base that was
181 trenched 10-cm into the soil. The rubber liner was then adhered to the acrylic
182 glass chamber using silicon grease. $^{13}\text{CO}_2$ was introduced into each chamber via
183 gas-tight ports by reacting 1-g sodium carbonate ($\text{Na}^{13}\text{CO}_3$, 99 atom% ^{13}C ,
184 Sigma-Aldrich; CAS number: 9367-48-4; 113-mg of ^{13}C equivalent) with excess
185 hydrochloric acid. Air was circulated within the chambers using a centrally
186 located internal battery-operated fan. Chamber temperature was monitored using
187 an internal thermometer. CO_2 concentrations within the chamber were monitored
188 via a LI-8100 infrared gas analyzer (Li-Cor Biosciences, Lincoln, NE). Chambers
189 were removed after CO_2 levels returned to pre-pulse levels. As temperatures in
190 the chambers can be high during mid-day, pulsing was limited to early morning
191 and late afternoon. The amount of ^{13}C fixed by the plant communities was
192 determined by taking foliar clip samples immediately post-pulse.

193
194 Following the ^{13}C pulse-labeling, each plot was also labeled with ^{15}N ammonium
195 nitrate ($^{15}\text{NH}_4^{15}\text{NO}_3$; 98 atom%; Sigma-Aldrich; CAS Number: 31432-46-9; 67-mg
196 of ^{15}N equivalent) in order to examine N-dynamics in response to manure and
197 antibiotic treatments. Ammonium nitrate (300-mg in 1-L of DI water) was added
198 evenly to the soil surface of each 1-m^2 plot. The amount of ^{15}N , similar to
199 Fraterrigo *et al.* (45), was kept low to avoid a fertilization effect.

200
201 Upon completion of pulse-labeling, we destructively harvested 0.05-m^2 sub-plots
202 within each 1-m^2 plot at 1, 2, and 7-days post-labeling. An additional sub-plot
203 was harvested from each 1-m^2 experimental plot prior to pulse-labeling in order
204 to determine natural abundance of ^{13}C and ^{15}N . Aboveground plant material from
205 each sub-plot was harvested by clipping it at the soil surface. Aboveground plant
206 biomass samples were air-dried, weighed, and milled for elemental and isotope
207 analyses. The belowground portion of the sub-plot was sampled to 10-cm depth;
208 roots and soil were then separated. Root material was initially air-dried and then
209 later washed, air-dried, weighed, and milled for elemental and isotopic analyses.
210 Soils were sieved (4.75-mm), homogenized, and stored at either -80°C , 4°C , or
211 air-dried depending on future analyses (see below).

212

213 For soils, we determined POM and mineral-associated soil C and N, and soil
214 microbial biomass C and N. POM and mineral-associated C and N was
215 determined on air dried soil samples (46). Microbial biomass C and N were
216 determined following the chloroform fumigation extraction (CFE) procedure
217 outlined in Fierer and Schimel (47). Briefly, 40 mls of 0.5M K₂SO₄ was added to
218 one of each 7-g dry mass equivalent soil pair. One of each pair is then exposed
219 to 1-ml of ethanol-free chloroform to lyse microbial cells and accumulate
220 microbial C and N. Samples are capped, and shaken for 4-h. Samples were then
221 allowed to settle before filtration. Microbial biomass was estimated as the
222 difference between the quantity of C and N between the fumigated and un-
223 fumigated samples. Total organic C and N were then calculated for both the
224 fumigated and un-fumigated samples using a Vario TOC Cube (Elementar,
225 Langensfeld, Germany).

226

227 $\delta^{13}\text{C}$ and $\delta^{15}\text{N}$ in above- and belowground plant biomass, POM and mineral-
228 associated pools were determined using a Costech ECS 4010 Elemental
229 Analyzer (Costech Analytical, Valencia, CA, USA) paired with an Thermo Delta
230 Plus Advantage Isotope Ratio Mass Spectrometer (IRMS; Thermo Fisher
231 Scientific™, Waltham, MA, USA).

232

233 Prior to each destructive harvest event ecosystem respiration was measured
234 using a LI-8100 Infrared Gas Analyzer (Li-Cor Biosciences, Lincoln, NE, USA).
235 Additionally, two 15-ml subsamples of respired air were captured using a gas
236 syringe and air-evacuated exetainers in order to determine the $\delta^{13}\text{C}$ of respired
237 CO₂. The first of two subsamples were collected within the first 15-seconds of a
238 2-minute respiration measurement period, and the second subsample was
239 collected in the final 15-seconds. Both subsamples were then analyzed for $\delta^{13}\text{C}$
240 using a GasBench II IRMS (Thermo Fisher Scientific™, Waltham, MA, USA).
241 Data were then recalculated to account for varying heights of soil collars and
242 adjusted to optimize the r^2 of the respiration trend-line from each 2-minute
243 measurement.

244

245 The amount of ¹³C fixed, respired, and the amount of both ¹³C and ¹⁵N contained
246 in above- and belowground pools was derived using standard isotopic mixing
247 models (48). The amount of C and N derived from ¹³C and ¹⁵N additions was
248 calculated as atom excess in a given C or N pool. The atom% excess of a given
249 pool was then multiplied by the total C or N in that pool, giving the mass of ¹³C or
250 ¹⁵N label. The proportion of label in a given pool was calculated as the mass of
251 ¹³C or ¹⁵N label divided by the total amount of C or N of that specific pool.

252 Cumulative ecosystem respiration was calculated via integration. See
253 Supplemental Methods for details related to additional soil parameters, microbial
254 catabolic response profiles, and microbial community composition measured in
255 conjunction with the pulse-chase experiment.

256

257 *Statistics and analysis:*

258

259 Data was analyzed using linear mixed models (LMM; 'lme4' package; 63) with
260 treatment as a fixed effect and plot nested within block as a random effect to
261 account for sampling of plots across time. Model selection (additive vs.
262 interactive) was determined by lowest Akaike information criterion (AIC) score
263 (50). Normality of variance was tested using a Wilk-Shapiro test. Data with non-
264 normal variance was either log or square-root transformed. If normality
265 assumptions were still not met, generalized linear models (GLM; 'car' package;
266 64) were used using the Gamma family and either the inverse or log link function
267 as all data was continuous and positive (variables containing negative values
268 were standardized). Wald χ^2 tests were used to assess model significance for
269 GLMs. Data was analyzed using the R statistical platform (52). Degrees of
270 freedom for linear mixed models were calculated using Satterthwaite
271 approximations.

272

273 For all analyses, we consider statistical significance at $P < 0.05$, and marginal
274 significance at $P < 0.10$. However, it should be noted that it has typically been
275 deemed acceptable to consider changes in soil C pools at $P < 0.10$ (53, 54),
276 given that soil C is inherently heterogeneous.

277

278 **Results/Discussion:**

279

280 *Antibiotic effects on active and total microbial biomass:*

281

282 Prolonged manure additions, regardless of antibiotic involvement, increased C
283 mineralization – an estimate of bioavailable soil C – when compared to NMC
284 ($F_{3,15} = 11.8$, $P < 0.001$; Table S1). More surprising was the observation that
285 active microbial biomass – determined via SIR – was differentially affected by the
286 antibiotic status of the manure (again, we are referring to the effect of exposure
287 to manure from cows given an antibiotic, as an antibiotic effect; Figure S2a; $P <$
288 0.01). Specifically, the Ceph treatment exhibited greater active microbial biomass
289 in comparison to the other treatments. Similar to the increase in respiration
290 observed for SIR, increased microbial activity was observed across a range of C-
291 substrates for the Ceph treatment in a catabolic response profile (CRP; Figure
292 S2b). In contrast, we observed a marginally significant treatment effect on total
293 microbial biomass C ($P = 0.07$) and N ($P = 0.08$), primarily driven by a trend
294 towards increased microbial C and N in the Con treatment (Table S2). This
295 contrast between active and total microbial biomass may suggest physiological
296 changes, specifically greater mass-specific activity for the Ceph treatment,
297 consistent with findings from previous investigations (21). As discussed in the
298 corresponding section, the Ceph treatment did not differ from the other antibiotic
299 manure treatment, Pir, in terms of microbial community composition. Therefore,
300 elevated microbial activity could be due to two, non-mutually exclusive, factors: *i*)
301 the increased presence of lysed cellular material from the action of cephalosporin, a
302 bactericidal antibiotic, or *ii*) from a stress response of the microbial community,
303 due to the added metabolic cost of maintaining antibiotic resistance (36). This
304 stress response is consistent with previous research on cephalosporin use on dairy

305 cattle in pasture systems, and has been suggested as a possible cause of
306 altered ecosystem C cycling, through reduced microbial efficiency (21).

307

308 *Antibiotic effects on fungal:bacterial dominance:*

309

310 As antibiotics detrimentally effect bacteria, we assessed F:B via qPCR to
311 determine shifts in fungi and bacteria. Overall, we observed a significant
312 treatment effect for F:B (Figure 1; $F_{3,63} = 6.497$, $P < 0.001$) as has been
313 previously observed (55). Fungal counts increased in the soils receiving manure
314 compared to NMC, as well as in Pir compared to Con. NMC had the lowest ratio,
315 indicating that without manure and associated antibiotics the system is relatively
316 more dominated by bacteria. Notably, within the Pir and Ceph manure
317 treatments, the differences in F:B were due to declines in bacteria (*i.e.* 16S gene
318 abundance), whereas little change in fungi (*i.e.* ITS abundance) was observed.
319 This increase in F:B in the Pir treatment may be driven by pirlimycin's
320 bacteriostatic mode of action: pirlimycin typically reduces bacterial growth but
321 does not induce cell lysis. Conversely, cephalosporin – a bactericidal antibiotic –
322 causes cell lysis. Lysed cells, as suggested above, cause an increase in labile
323 resources that potentially favor bacteria in spite of the direct negative effects of
324 the antibiotic (56, 57). This potential net positive effect for bacteria under the
325 Ceph compared to the Pir treatment is supported by a pairwise marginally
326 significant increase in 16S copies (Figure 1; $P = 0.08$). It is also possible that
327 decreased inhibition of bacteria in the Ceph treatment can be attributed to
328 cephalosporin being relatively more easily degraded than pirlimycin. However,
329 evidence of cephalosporin's effect on microbial functional properties was in fact
330 observed, therefore degradability is unlikely to be the explanation for the
331 difference in bacterial effects between antibiotic treatments. Compared to NMC,
332 the addition of manure, regardless of antibiotic involvement, increased the
333 abundance of fungi (*i.e.* ITS copies) in soil ($F_{3,68} = 5.868$, $P < 0.01$), with Con,
334 Ceph, and Pir treatments having greater numbers of ITS copies than NMC.
335 Within the manure-addition treatments, Ceph had a marginally lower abundance
336 of fungi compared to Con ($P < 0.10$), this too could be driven by the mode of
337 action related to this antibiotic.

338

339 The primary fungal effect appeared to be driven by manure itself, given that all
340 manure additions increased ITS copies compared to NMC. This could be
341 attributed in part to coprophilous fungi, which specialize in the decomposition of
342 fecal matter, previously shown to be elevated in conjunction with manure (21).
343 Additionally, the highest counts of 16S and ITS copies were measured in the Con
344 treatment. This is likely attributed to the influx of manure-derived resources in the
345 absence of antibiotics. Together these results suggest that while manure
346 additions increase F:B, manure from cattle administered antibiotics tends to lead
347 to even greater increases, primarily driven by decreased bacterial abundance.

348

349 *Antibiotic effects on microbial community composition:*

350

351 The results of our community composition assessment largely mirrored the
352 results of the F:B analysis. Bacterial communities changed across our treatments
353 (Figure 2: $pseudo-F_{3,23} = 1.15$; $P < 0.05$; note, the random effect 'block' was
354 dropped from this model because it was non-significant), but fungal communities
355 did not (Figure S3: $pseudo-F_{3,23} = 1.01$; $P = 0.18$). The treatment effect on
356 bacteria was largely driven by differences between Con, and both Pir and Ceph.
357 Notably, a marginally significant pairwise difference was observed between Pir
358 and Ceph ($P = 0.064$). NMC did not differ from the other three treatments, in fact,
359 as NMC can be viewed as a baseline, shifts in bacterial community composition
360 from manure exposure were dependent on the antibiotic status of the manure. If
361 the manure was sourced from cattle administered an antibiotic, the community
362 shifted to the lower right in ordination space, while control manure caused the
363 community to shift in the opposite direction, with NMC situated between (Figure
364 2a). Additionally, bacterial beta diversity did not differ between treatments (Figure
365 2a; $pseudo-F_{3,20} = 1.12$; $p = 0.31$), suggesting that the microbial communities in
366 our antibiotic treatments are distinct from control environments, and not just more
367 variable in composition. While we did not seek to document the impact of the
368 fecal microbiome on the soil microbiome, previous studies have shown that the
369 fecal microbiome can be impacted by antibiotic exposure (38). Therefore, further
370 research into the quantification of this effect would be beneficial, especially
371 investigations into interactions between the fecal and soil microbiomes.

372
373 To further investigate OTUs that possibly drove treatment differences, we
374 identified potential OTUs of interest via SIMPER that were common across all
375 pairwise treatment comparisons. This resulted in 32 common OTUs of which only
376 6 exhibited significant differences between treatments (Figure 2b). Of these 6
377 OTUs, 4 were associated with Phyla Acidobacteria and γ -Proteobacteria (2 in
378 each), and 2 were associated with the Phyla Bacteroidetes and Verrucomicrobia
379 (1 in each). Interestingly, the two γ -Proteobacteria were associated with families
380 Acinetobacter and Xanthomonadaceae, which are typically associated with the
381 environment but also include members of human health concern (58). In fact,
382 Wepking *et al.* (21) observed a similar increase in the genus Acinetobacter in
383 response to cattle administered a cephalosporin. Our results add further support
384 to the likely influence of antibiotics on soil community structure, and further
385 support the proposition that inputs of manure from cattle given antibiotics can
386 shift soil microbial communities towards organisms that are related to those of
387 human health concern (59, 60).

388
389 Interestingly, though, several OTUs associated with phyla that we expect to be
390 more oligotrophic in nature (*i.e.* Acidobacteria, Verrucomicrobia; 49, 50) also
391 exhibited greater relative abundance in Pir and Ceph. Even some taxa in the
392 family Cytophagaceae could be classed as oligotrophs, especially those involved
393 in cellulose degradation (63). This greater relative abundance of oligotrophic
394 taxa, primarily in Pir, may be due *not* to an increase in these groups but to a
395 decrease in other potentially more copiotrophic groups. That is in the Pir
396 treatment there was an observed decrease in 16S abundance, suggesting a

397 decline in overall bacterial abundance. Such a decrease, if driven by the
398 antibiotic pirlimycin may have been disproportionate because the antibiotic is
399 bacteriostatic and, as such is likely to have a more detrimental effect on active
400 bacteria (64, 65). Overall, these results suggest that manure from cattle given
401 antibiotics versus those not, has the potential to lead to shifts in soil bacterial
402 community composition and F:B dominance in a relatively short time (*i.e.* ~8
403 months) with implications on microbially-mediated ecosystem function.

404

405 *Antibiotic effects on carbon and nitrogen dynamics:*

406

407 Few differences were observed in most pools of recently fixed C (Table S4), and
408 in the amount of ^{13}C fixed across manure-amended treatments (NMC fixed more
409 ^{13}C relative to total plant C, likely due to the lower plant biomass and identical
410 amount of labeled C added to the chamber; $F_{3,20} = 3.07$, $P = 0.05$). However, we
411 did observe a significant effect of both treatment ($\chi^2 = 18.52$, $df = 3$, $P < 0.001$),
412 and time ($\chi^2 = 87.18$, $df = 2$, $P < 0.001$), as well as a treatment \times time interaction
413 ($\chi^2 = 41.13$, $df = 6$, $P < 0.001$), when examining the ecosystem respiration of
414 recently fixed C (Figure 2a). Specifically, the Pir treatment exhibited greater initial
415 respiration of ^{13}C compared to the other treatments, but by day 7 of the
416 experiment, respiration of ^{13}C for this treatment was nearly zero (Figure 3a). The
417 NMC, Pir, and Con treatments exhibited similar respiration dynamics (Figure 3a).
418 Ecosystem respiration dynamics for the Ceph treatment were more constant
419 during the sampling period compared to the other three treatments (Figure 3a). A
420 marginally significant difference in cumulative ^{13}C respired across the entire
421 sampling period was explained by the greatest amount of ^{13}C being respired in
422 the Pir treatment with the NMC and Ceph treatments intermediate, and the Con
423 treatment the lowest ($F_{3,15} = 2.72$; $P = 0.08$; Figure 3a). In fact, nearly twice the
424 amount of newly photosynthesized C was respired – not retained in the soil – in
425 the Pir treatment compared to the Con treatment (Figure 3a).

426

427 These results suggest that manure from cattle administered antibiotics can alter
428 both ecosystem respiration dynamics of recently fixed ^{13}C (*i.e.* Ceph) as well as
429 the total amount of C respired (*i.e.* Pir) compared to manure from antibiotic-free
430 cattle. Manure additions from cattle not administered antibiotics may initially
431 suppress respiration slightly compared to sites receiving no manure, possibly
432 driven by decreased plant demand for nutrients. Additionally, if less recently fixed
433 C is lost from a system via respiration it is likely that more C will be sequestered
434 in that system. This supposition is supported by the significant treatment effect
435 on the proportion of ^{13}C recovered in the mineral pool during the entire
436 experiment (Table S4) with the most ^{13}C recovered in the NMC treatment
437 followed by the Con treatment. Further, at the conclusion of the pulse-chase we
438 observed a marginally significant treatment effect for ^{13}C found in the mineral-
439 associated soil C pool with the most ^{13}C recovered in the Con treatment among
440 the treatments containing manure (Figure 3b; $F_{3,15} = 3.04$, $P = 0.06$). Given the
441 slow turnover of the mineral-associated soil C pool (66), our results suggest that
442 inputs of manure from cattle administered antibiotics may decrease C-

443 sequestration potential. Direct evidence for this potential is the observation of a
 444 significant treatment effect for the ratio of fixed ^{13}C to respired ^{13}C (Figure 3c;
 445 $F_{3,15} = 3.65$, $P < 0.05$), an indicator of ecosystem-scale C-use efficiency (37).
 446 Specifically, we observed that the Pir treatment had the lowest overall C-use
 447 efficiency, Con had the greatest, and both NMC and Ceph were intermediate.
 448 Soils receiving the Con treatment fixed 2.5-fold more C for every unit of C
 449 respired than did the Pir treatment. Together these results indicate that manure
 450 inputs from animals administered antibiotics have the potential to increase C
 451 losses from ecosystems compared to manure inputs from animals not
 452 administered antibiotics. However, our results also indicate that this effect on C-
 453 cycling may be influenced by the specific choice of antibiotics. Further
 454 investigation to examine the ecosystem effects of administering an array of
 455 antibiotics is merited, especially as agricultural management practices are
 456 increasingly seen as opportunities to mediate global climate change (67).

457
 458 A greater proportion of ^{15}N relative to the total N pool was observed with the Pir
 459 treatment compared to the other manure-amended plots across all pools
 460 sampled (Figure 4, Table S4), but not necessarily in comparison to NMC.
 461 Measuring ^{15}N as a proportion of the total N pool addresses potential difference
 462 in plant biomass between treatments (68). Within the aboveground biomass ($F_{3,61}$
 463 $= 8.08$, $P < 0.001$; Figure 4a, Table S4; analyzed as additive model based on
 464 quality of model using AIC score) and belowground biomass ($F_{3,55} = 4.53$, $P <$
 465 0.01 ; Figure 4b, Table S4) significantly more ^{15}N was found in the Pir treatment.
 466 In addition, a significant and marginally significant time effect was observed in
 467 the proportion of ^{15}N in the aboveground and the belowground biomass,
 468 respectively ($F_{2,61} = 6.42$, $P < 0.005$; $F_{2,55} = 2.74$, $P < 0.10$, respectively; Table
 469 S4; pooled across treatment). This was characterized by an increased in the
 470 proportion of ^{15}N in plant biomass across time. We also observed a significant
 471 treatment effect for total N in aboveground plant biomass ($F_{3,61} = 8.48$, $P < 0.001$;
 472 Table S2) and a marginally significant treatment effect for total N in belowground
 473 plant biomass ($F_{3,55} = 2.55$, $P = 0.06$; Table S2; the former was analyzed as
 474 additive model, and the latter as an interactive model based on AIC model
 475 score). This effect was likely due to greater biomass in the treatments receiving
 476 manure versus NMC (Table S2).

477
 478 As observed in plant biomass, a greater proportion of ^{15}N in POM (Treatment:
 479 $F_{3,61} = 4.77$, $P < 0.005$; Figure 3c, Table S4) and mineral-associated (Treatment:
 480 $F_{3,61} = 2.49$, $P < 0.10$; Figure 3d, Table S4) soil N pools was observed in the Pir
 481 treatment at the conclusion of the experiment compared to the other treatments.
 482 The effect of Pir on ^{15}N in the POM N fraction was likely attributable to the root
 483 biomass, due to the contribution of plant derived constituents to this pool (69).
 484 Abundance of ^{15}N in the mineral pool was also increased with Pir, possibly due to
 485 decreased plant-microbe competition for N due to an increased F:B in the
 486 microbial community. Microbial communities with a higher F:B typically have a
 487 higher C:N due to reduced N demand of fungi (70). With the Pir system being

488 more fungally dominated, the overall microbial demand for N is likely lower than
489 for the other three treatments.

490

491 We see an increased loss of recently fixed C in the Pir treatment but also
492 increased uptake of recently added N in this treatment compared to the other
493 treatments. For C, this is particularly unexpected given the increased F:B
494 associated with the Pir treatment. However, the notion that increased F:B leads
495 to a less leaky C-cycle has been called into question (43, 71). Rousk and Frey
496 (43) found that bacterial dominance is linked to a less leaky C-cycle and a less
497 leaky N-cycle, while we observed the potential for greater plant uptake of N. This
498 disparity between our results and those of Rousk and Frey may be because our
499 research was conducted in a grassland system and theirs in a forest. Of
500 particular relevance – among the many differences between these systems – the
501 uptake of available N is likely greater in grasses and forbs during peak growth,
502 over a short period of time, compared to N uptake in trees. When only
503 considering the microbial community, the Pir treatment (*i.e.* higher F:B) appears
504 to have a leakier N cycle – but when the plant community is also included, this
505 effect is diminished. This highlights the potential for antibiotics to alter plant-
506 microbe interactions and lead to shifts in ecosystem processes.

507

508 Mechanistically, these effects on N dynamics could be due to altered microbe-
509 plant competition for N. In this instance a bacteriostatic antibiotic – pirlimycin –
510 increased F:B leading to a leakier C cycle, but also a decrease in plant-microbe
511 competition for N (evidenced by increased plant N uptake; 52). Another potential
512 mechanism is reduced competition with mycorrhizal fungi for N with the Pir
513 treatment. Given recent evidence suggesting a C cost associated with N uptake
514 (73) for mycorrhizal symbionts, if mycorrhizal N uptake increases with a reduction
515 in bacteria then plant N uptake may increase but more C may be lost from the
516 system. Finally, as this experiment was conducted during peak plant growth and
517 N demand, more N may in fact be lost from the system with decreased plant N
518 demand.

519

520 **Conclusion:**

521

522 Antibiotics affect not just the soil microbiome but the entire ecosystem; how the
523 ecosystem is affected depends on the antibiotic's mode of action (*i.e.* bactericidal
524 vs. bacteriostatic). Of the two antibiotics investigated, one in particular –
525 pirlimycin – alters both C and N cycling. This is likely due to changes in microbial
526 composition – as demonstrated by increased F:B, and shifts in bacterial
527 community composition. Increased availability of N appears to occur with
528 decreased C retention in the system – subsequently decreased whole ecosystem
529 C-use efficiency. In contrast, cephalosporin increases microbial activity as a stress
530 response – in keeping with previously published research which showed
531 decreased microbial efficiency and increased soil C loss (21). While the majority
532 of attention is paid to livestock antibiotic use from the perspective of the
533 proliferation of antibiotic resistant pathogens and antibiotic resistance genes (74,

534 75), the impacts on biogeochemical cycling have been overlooked. With global
535 livestock antibiotic use projected to increase by 67% between 2010 and 2030 (4)
536 combined with increasing atmospheric CO₂ concentrations, understanding and
537 accounting for the effects antibiotics have on soil microbial communities and
538 whole ecosystem function is imperative.

539

540 ***Acknowledgements:***

541

542 We thank the University of Georgia Stable Isotope Ecology Laboratory, and the
543 Yale Analytical and Stable Isotope Center for analysis of stable isotopic samples.

544 Thanks as well to the many graduate and undergraduate researchers who
545 helped with the field and lab work associated with this study.

546

547 *Funding:*

548

549 USDA-NIFA: Award number: 2013-67019-21363

550 NSF: Award number: 1832888

551 **Works Cited:**

552

- 553 1. O'Neill J (2016) *Tackling drug-resistant infections globally: final report and*
554 *recommendations. The review on antimicrobial resistance.*
- 555 2. Rubin RJ, et al. (1999) The economic impact of Staphylococcus aureus
556 infection in New York City hospitals. *Emerg Infect Dis* 5(1):9–17.
- 557 3. Klein E, Smith DL, Laxminarayan R (2007) Hospitalizations and deaths
558 caused by methicillin-resistant Staphylococcus aureus, United States,
559 1999-2005. *Emerg Infect Dis* 13(12):1840–1846.
- 560 4. Van Boeckel TP, et al. (2015) Global trends in antimicrobial use in food
561 animals. *Proc Natl Acad Sci U S A* 112(18):5649–5654.
- 562 5. Koch BJ, Hungate BA, Price LB (2017) Food-animal production and the
563 spread of antibiotic resistance: the role of ecology. *Front Ecol Environ*
564 15(6):309–318.
- 565 6. Van Boeckel TP, et al. (2017) Reducing antimicrobial use in food animals.
566 *Science (80-)* 357(6358):1350–1352.
- 567 7. Kemper N (2008) Veterinary antibiotics in the aquatic and terrestrial
568 environment. *Ecol Indic* 8(1):1–13.
- 569 8. Chee-Sanford JC, et al. (2009) Fate and transport of antibiotic residues
570 and antibiotic resistance genes following land application of manure waste.
571 *J Environ Qual* 38(3):1086.
- 572 9. Zhu Y-G, et al. (2013) Diverse and abundant antibiotic resistance genes in
573 Chinese swine farms. *Proc Natl Acad Sci* 110(9):3435–3440.
- 574 10. Jechalke S, Heuer H, Siemens J, Amelung W, Smalla K (2014) Fate and
575 effects of veterinary antibiotics in soil. *Trends Microbiol* 22(9):536–545.
- 576 11. Ghosh S, LaPara TM (2007) The effects of subtherapeutic antibiotic use in
577 farm animals on the proliferation and persistence of antibiotic resistance
578 among soil bacteria. *ISME J* 1(3):191–203.
- 579 12. Forsberg KJ, et al. (2012) The shared antibiotic resistome of soil bacteria
580 and human pathogens. *Science (80-)* 337(6098):1107–1111.
- 581 13. Popowska M, et al. (2012) Influence of soil use on prevalence of
582 tetracycline, streptomycin, and erythromycin resistance and associated
583 resistance genes. *Antimicrob Agents Chemother* 56(3):1434–1443.
- 584 14. FDA (2013) *2013 summary report on antimicrobials sold or distributed for*
585 *use in food-producing animals.*
- 586 15. Lipsitch M, Singer RS, Levin BR (2002) Antibiotics in agriculture: When is it
587 time to close the barn door? *Proc Natl Acad Sci* 99(9):5752–5754.
- 588 16. Toth JD, Feng Y, Dou Z (2011) Veterinary antibiotics at environmentally
589 relevant concentrations inhibit soil iron reduction and nitrification. *Soil Biol*
590 *Biochem* 43(12):2470–2472.
- 591 17. Gutiérrez IR, Watanabe N, Harter T, Glaser B, Radke M (2010) Effect of
592 sulfonamide antibiotics on microbial diversity and activity in a Californian
593 Mollic Haploxeralf. *J Soils Sediments* 10(3):537–544.
- 594 18. Sarmah AK, Meyer MT, Boxall ABA (2006) A global perspective on the
595 use, sales, exposure pathways, occurrence, fate and effects of veterinary
596 antibiotics (VAs) in the environment. *Chemosphere* 65(5):725–759.


- 597 19. Roose-Amsaleg C, Laverman AM (2016) Do antibiotics have
598 environmental side-effects? Impact of synthetic antibiotics on
599 biogeochemical processes. *Environ Sci Pollut Res* 23(5):4000–4012.
- 600 20. Grenni P, Ancona V, Barra Caracciolo A (2018) Ecological effects of
601 antibiotics on natural ecosystems: A review. *Microchem J* 136:25–39.
- 602 21. Wepking C, et al. (2017) Exposure to dairy manure leads to greater
603 antibiotic resistance and increased mass-specific respiration in soil
604 microbial communities. *Proc R Soc B Biol Sci* 284:1–9.
- 605 22. Zak DR, Blackwood CB, Waldrop MP (2006) A molecular dawn for
606 biogeochemistry. *Trends Ecol Evol* 21(6):288–295.
- 607 23. Schimel JP, Schaeffer SM (2012) Microbial control over carbon cycling in
608 soil. *Front Microbiol* 3(348):1–11.
- 609 24. Philippot L, et al. (2013) Loss in microbial diversity affects nitrogen cycling
610 in soil. *ISME J* 7(8):1609–1619.
- 611 25. Graham EB, et al. (2014) Do we need to understand microbial communities
612 to predict ecosystem function? A comparison of statistical models of
613 nitrogen cycling processes. *Soil Biol Biochem* 68:279–282.
- 614 26. van der Heijden MGA, Bardgett RD, van Straalen NM (2008) The unseen
615 majority: soil microbes as drivers of plant diversity and productivity in
616 terrestrial ecosystems. *Ecol Lett* 11(3):296–310.
- 617 27. Bonfante P, Anca I-A (2009) Plants, mycorrhizal fungi, and bacteria: a
618 network of interactions. *Annu Rev Microbiol* 63(1):363–383.
- 619 28. Raaijmakers JM, Paulitz TC, Steinberg C, Alabouvette C, Moënne-Loccoz
620 Y (2009) The rhizosphere: a playground and battlefield for soilborne
621 pathogens and beneficial microorganisms. *Plant Soil* 321(1–2):341–361.
- 622 29. van der Heijden MGA, Wagg C (2013) Soil microbial diversity and agro-
623 ecosystem functioning. *Plant Soil* 363(1–2):1–5.
- 624 30. Dessaux Y, Grandclément C, Faure D (2016) Engineering the rhizosphere.
625 *Trends Plant Sci* 21(3):266–278.
- 626 31. Raaijmakers JM, Mazzola M (2016) Soil immune responses. *Science (80-)*
627 352(6292):1392–1393.
- 628 32. Waksman S, Woodruff HB (1940) The soil as a source of microorganisms
629 antagonistic to disease-producing bacteria. *J Bacteriol* 40(4):581–600.
- 630 33. Martínez JL (2008) Antibiotics and antibiotic resistance genes in natural
631 environments. *Science (80-)* 321(5887):365–367.
- 632 34. Fajardo A, Martínez JL (2008) Antibiotics as signals that trigger specific
633 bacterial responses. *Curr Opin Microbiol* 11(2):161–167.
- 634 35. Aust MO, et al. (2008) Distribution of sulfamethazine, chlortetracycline and
635 tylosin in manure and soil of Canadian feedlots after subtherapeutic use in
636 cattle. *Environ Pollut* 156(3):1243–1251.
- 637 36. Schimel J, Balsler TC, Wallenstein M (2007) Microbial stress-response
638 physiology and its implications for ecosystem function. *Ecology*
639 88(6):1386–1394.
- 640 37. Geyer KM, Kyker-Snowman E, Grandy AS, Frey SD (2016) Microbial
641 carbon use efficiency: accounting for population, community, and
642 ecosystem-scale controls over the fate of metabolized organic matter.

- 643 *Biogeochemistry* 127(2–3):173–188.
- 644 38. Hammer TJ, et al. (2016) Treating cattle with antibiotics affects greenhouse
645 gas emissions, and microbiota in dung and dung beetles. *Proc R Soc B*
646 *Biol Sci* 283(1831):20160150.
- 647 39. Thiele-Bruhn S, Beck IC (2005) Effects of sulfonamide and tetracycline
648 antibiotics on soil microbial activity and microbial biomass. *Chemosphere*
649 59(4):457–465.
- 650 40. Demoling LA, Bååth E, Greve G, Wouterse M, Schmitt H (2009) Effects of
651 sulfamethoxazole on soil microbial communities after adding substrate. *Soil*
652 *Biol Biochem* 41(4):840–848.
- 653 41. Bardgett RD, McAlister E (1999) The measurement of soil fungal:bacterial
654 biomass ratios as an indicator of ecosystem self-regulation in temperate
655 meadow grasslands. *Biol Fertil Soils* 29(3):282–290.
- 656 42. Malik AA, et al. (2016) Soil fungal:bacterial ratios are linked to altered
657 carbon cycling. *Front Microbiol* 7(AUG):1–11.
- 658 43. Rousk J, Frey SD (2015) Revisiting the hypothesis that fungal-to-bacterial
659 dominance characterizes turnover of soil organic matter and nutrients. *Ecol*
660 *Monogr* 85(3):457–472.
- 661 44. Wind L, et al. (2018) Fate of pirlimycin and antibiotic-resistant fecal
662 coliforms in field plots amended with dairy manure or compost during
663 vegetable cultivation. *J Environ Qual* 47(3):436–444.
- 664 45. Fraterrigo JM, Strickland MS, Keiser AD, Bradford MA (2011) Nitrogen
665 uptake and preference in a forest understory following invasion by an
666 exotic grass. *Oecologia* 167(3):781–791.
- 667 46. Lal R (2001) *Assessment methods for soil carbon* (Lewis Publishers, Boca
668 Raton).
- 669 47. Fierer N, Schimel JP (2002) Effects of drying–rewetting frequency on soil
670 carbon and nitrogen transformations. *Soil Biol Biochem* 34(6):777–787.
- 671 48. Fry B (2006) *Stable Isotope Ecology* (Springer New York, New York, NY)
672 doi:10.1007/0-387-33745-8.
- 673 49. Bates D, Maechler Martin, Walker S (2016) Package “lme4.” *CRAN*
674 *Repos*:113.
- 675 50. Akaike H (1973) Information theory and an extension of the maximum
676 likelihood principle. *Second International Symposium on Information*
677 *Theory* doi:10.1007/978-1-4612-1694-0_15.
- 678 51. Fox J, et al. (2016) Package ‘car.’ *CRAN Repos*.
679 doi:10.1177/0049124105277200.
- 680 52. R Core Team (2017) R: A language and environment for statistical
681 computing. *R Found Stat Comput Vienna, Austria*:R Foundation for
682 Statistical Computing.
- 683 53. Carney KM, Hungate BA, Drake BG, Megonigal JP (2007) Altered soil
684 microbial community at elevated CO₂ leads to loss of soil carbon. *Proc*
685 *Natl Acad Sci* 104(12):4990–4995.
- 686 54. Strickland MS, Devore JL, Maerz JC, Bradford MA (2010) Grass invasion
687 of a hardwood forest is associated with declines in belowground carbon
688 pools. *Glob Chang Biol* 16(4):1338–1350.

- 689 55. Thiele-Bruhn S, Beck I-C (2005) Effects of sulfonamide and tetracycline
690 antibiotics on soil microbial activity and microbial biomass. *Chemosphere*
691 59(4):457–465.
- 692 56. Hendrix PF, et al. (1986) Detritus food webs in conventional and no-tillage
693 agroecosystems. *Bioscience* 36(6):374–380.
- 694 57. Rousk J, Bååth E (2007) Fungal and bacterial growth in soil with plant
695 materials of different C/N ratios. *FEMS Microbiol Ecol* 62(3):258–267.
- 696 58. Berendonk TU, et al. (2015) Tackling antibiotic resistance: the
697 environmental framework. *Nat Rev Microbiol* 13(5):310–317.
- 698 59. Marti R, et al. (2013) Impact of manure fertilization on the abundance of
699 antibiotic-resistant bacteria and frequency of detection of antibiotic
700 resistance genes in soil and on vegetables at harvest. *Appl Environ*
701 *Microbiol* 79(18):5701–5709.
- 702 60. Wind L, et al. (2018) Fate of pirlimycin and antibiotic-resistant fecal
703 coliforms in field plots amended with dairy manure or compost during
704 vegetable cultivation. *J Environ Qual* 47(3). doi:10.2134/jeq2017.12.0491.
- 705 61. Fierer N, et al. (2012) Comparative metagenomic, phylogenetic and
706 physiological analyses of soil microbial communities across nitrogen
707 gradients. *ISME J* 6(5):1007–1017.
- 708 62. Bergmann GT, et al. (2011) The under-recognized dominance of
709 Verrucomicrobia in soil bacterial communities. *Soil Biol Biochem*
710 43(7):1450–1455.
- 711 63. McBride MJ, Liu W, Lu X, Zhu Y, Zhang W (2014) The family
712 cytophagaceae. *The Prokaryotes: Other Major Lineages of Bacteria and*
713 *The Archaea* doi:10.1007/978-3-642-38954-2_382.
- 714 64. Ding C, He J (2010) Effect of antibiotics in the environment on microbial
715 populations. *Appl Microbiol Biotechnol* 87(3):925–941.
- 716 65. Lobritz MA, et al. (2015) Antibiotic efficacy is linked to bacterial cellular
717 respiration. *Proc Natl Acad Sci* 112(27):8173–8180.
- 718 66. Schlesinger WH, Lichter J (2001) Limited carbon storage in soil and litter of
719 experimental forest plots under increased atmospheric CO₂. *Nature*
720 411:466–469.
- 721 67. Griscom BW, et al. (2017) Natural climate solutions. *Proc Natl Acad Sci*
722 114(44):11645–11650.
- 723 68. Fraterrigo JM, Strickland MS, Keiser AD, Bradford MA (2011) Nitrogen
724 uptake and preference in a forest understory following invasion by an
725 exotic grass. *Oecologia*. doi:10.1007/s00442-011-2030-0.
- 726 69. Grandy AS, Neff JC (2008) Molecular C dynamics downstream: The
727 biochemical decomposition sequence and its impact on soil organic matter
728 structure and function. *Sci Total Environ* 404(2–3):297–307.
- 729 70. Mouginot C, et al. (2014) Elemental stoichiometry of fungi and bacteria
730 strains from grassland leaf litter. *Soil Biol Biochem* 76:278–285.
- 731 71. Strickland MS, Rousk J (2010) Considering fungal:bacterial dominance in
732 soils – Methods, controls, and ecosystem implications. *Soil Biol Biochem*
733 42(9):1385–1395.
- 734 72. Schimel JP, Bennett J (2004) Nitrogen mineralization: Challenges of a



- 735 changing paradigm. *Ecology*. doi:10.1890/03-8002.
- 736 73. Shi M, Fisher JB, Brzostek ER, Phillips RP (2016) Carbon cost of plant
737 nitrogen acquisition: global carbon cycle impact from an improved plant
738 nitrogen cycle in the Community Land Model. *Glob Chang Biol* 22(3):1299–
739 1314.
- 740 74. Udikovic-Kolic N, Wichmann F, Broderick NA, Handelsman J (2014) Bloom
741 of resident antibiotic-resistant bacteria in soil following manure fertilization.
742 *Proc Natl Acad Sci* 111(42):15202–15207.
- 743 75. McEachran AD, et al. (2015) Antibiotics, bacteria, and antibiotic resistance
744 genes: Aerial transport from cattle feed yards via particulate matter.
745 *Environ Health Perspect* 123(4):337–343.
- 746
- 747

748 *Figures:*
 749
 750 Figure 1.
 751




752

753 Figure 2




754
755

756 Figure 3
 757
 758



759

760 Figure 4
761



762

763 **Figure Legends:**

764

765 **Figure 1:** Fungal-to bacterial ratios (F:B) associated with sites receiving manure
 766 from cattle administered no antibiotics (Control), administered cephalosporin
 767 (bactericidal), or pirlimycin (bacteriostatic). Also shown is the ratio for sites
 768 receiving no manure (NMC). Bars represent the mean \pm 1 SEM. Letters indicate
 769 pairwise differences between treatments. 16S and ITS copies are indicated by
 770 open circles and filled squares, respectively.

771

772 **Figure 2:** Effect of manure treatments on soil prokaryotic community
 773 composition. **A)** Principal components analysis showing prokaryote community
 774 composition associated with the following treatments: Soil amended with no
 775 manure (NMC), soil amended with manure from cattle given no antibiotics
 776 (Control), and soil amended with manure from cattle given either a bactericidal
 777 antibiotic (Cephalosporin) or a bacteriostatic antibiotic (Pirlimycin). Centroids are
 778 indicated as symbols with central points and shown as the mean \pm 1 SE.
 779 Significant pairwise differences between centroids are denoted by different letters
 780 in the key. Additionally, lines connecting points indicate those treatments
 781 receiving antibiotics (solid line) versus those that did not receive antibiotics
 782 (dashed line). **B)** Relative abundance of OTUs that both contributed to
 783 dissimilarity between treatments (as determined via similarity percentages) and
 784 were statistically significant. Overall treatment statistical significance is indicated
 785 by *P*-values, and significant pairwise differences for within OTU comparisons are
 786 denoted by different letters.

787

788 **Figure 3:** Effect of manure and antibiotic treatments on the cycling of C through
 789 the above- and belowground pools across the following treatments: soil amended
 790 with no manure (NMC), soil amended with manure from cattle given no
 791 antibiotics (Control), and soil amended with manure from cattle given either a
 792 bactericidal antibiotic (Cephalosporin) or a bacteriostatic antibiotic (Pirlimycin). **a)**
 793 Ecosystem respiration dynamics across the 7-day sampling period are shown in
 794 the main panel. Points represent the mean \pm 1 SE (Treatment: $F_{3,61} = 5.3$, $P <$
 795 0.005 ; Time: $F_{2,61} = 26.6$, $P < 0.001$). The panel inset shows a boxplot of the
 796 cumulative ^{13}C respired across the entire pulse-chase (Treatment: $\chi^2 = 6.86$, $df =$
 797 3 , $P < 0.08$). While the cumulative ^{13}C respired is marginally significant, it
 798 represents a doubling of respired CO_2 , and is therefore ecologically meaningful.
 799 **b)** Total accumulation of ^{13}C in the mineral associated soil fraction by the end of
 800 the 7-day pulse chase event (Treatment: $F_{3,15} = 3.04$, $P = 0.06$). **c)** The ratio of
 801 ^{13}C fixed to ^{13}C respired, an indicator of whole ecosystem C-use efficiency
 802 (Treatment: $F_{3,15} = 3.65$, $P < 0.05$). Letters indicate pairwise differences between
 803 treatments.

804

805 **Figure 4:** Effect of manure and antibiotic treatments on the cycling of newly
 806 added N through the above- and belowground systems across the following
 807 treatments: soil amended with no manure (NMC), soil amended with manure
 808 from cattle given no antibiotics (Control), and soil amended with manure from

809 cattle given either a bactericidal antibiotic (Cephapirin) or a bacteriostatic
810 antibiotic (Pirlimycin), and across time. All panels show the proportion of ^{15}N
811 within each respective N pool. Error bars represent ± 1 SEM.

812 **Supplementary Materials:**

813

814 *Supplemental Methods:*

815

816 *Manure sourcing:*

817

818 Manure collection started by selecting two sets of cattle: 12 healthy, peak
 819 lactation dairy cows, and 6 cows at the end of their current lactation cycle (n=18).
 820 The latter group was treated with cephapirin (ToMORROW®; Boehringer
 821 Ingelheim Vetmedica, Inc., Duluth, GA, USA; intramammary dry cow therapy;
 822 single dose of 300-mg into each of four quarters). Half of the former group (n=6)
 823 was treated therapeutically with pirlimycin (Pirsue®; Zoetis, Parsippany, NJ, USA;
 824 intramammary dose typical for clinical mastitis; two doses of 50-mg each, 24-h
 825 apart). The remaining healthy lactating cows (n=6) were used for control manure
 826 and therefore were not treated with antibiotics. Experimental cows were selected
 827 for homogeneity of body weight and stage of lactation, and none had received
 828 previous antibiotic treatment in the current lactation.

829

830 All cattle were offered free choice water and *ad libitum* total mixed ration. Cows
 831 assigned to the same treatment were group housed in a single pen located in a
 832 free stall barn (*i.e.* total of three pens). On day 1 of the study, cows were treated
 833 with the assigned antibiotic. Manure (feces and urine mixed) accumulated over a
 834 24-h period was collected from the pen floor on day 2 and 3 post treatment.
 835 Manure from each pen was mixed separately to achieve homogeneous manure.
 836 All collected manure was then homogenized and stored at -20°C before being
 837 applied to the plots.

838

839 *Manure elemental properties:*

840

841 Some treatment differences in manure %C, %N, and C:N were observed when
 842 compared with linear models and ANOVAs (%C: $F_{2,33} = 12.525$, $P < 0.001$; %N:
 843 $F_{2,33} = 2.858$, $P < 0.072$; CN: $F_{2,33} = 4.256$, $P < 0.05$; Table S5). Con manure and
 844 Pir manure had a significantly greater %C than Ceph manure – although all
 845 manures were between 48-50.5% C. For manure %N a marginally significant
 846 treatment effect was found, although no significant pairwise differences between
 847 treatments were noted – all manures were between 3-4% N. Control manure had
 848 the greatest C:N, significantly greater than both of the other manures – all ratios
 849 were between 11.9 – 17.4.

850

851 *Quantification of antibiotics in manure:*

852

853 Manure samples were analyzed for cephapirin and pirlimycin using the methods
 854 described by Ray *et al.* (76, 77). These methods were modified to make them
 855 suitable for the quantification of cephapirin and pirlimycin in manure samples.
 856 Manure samples (sample size: 1 g) were extracted using 5 mL of extractant
 857 [methanol (70%) and phosphate buffer (50 mM at pH 8.5)]. Extraction was

858 followed by extract clean-up involving solid phase extraction (SPE) using OASIS
859 HLB Plus Short Cartridge (250 mg sorbent; Waters, Milford, MA). An aliquot of 1
860 mL clean extracts was dried to dryness at 35°C under N₂ gas using a Zipvap 20
861 evaporator (Glas-Col, Terre Haute, IN) and dissolved in 1 mL of methanol:water
862 (30:70, v/v) with 0.1% formic acid. Dissolved extracts were filtered through 0.2
863 µm PVDF syringe filter (Fisher, Pittsburgh, PA) into 1.5 mL amber glass HPLC
864 vials and analyzed for cephalosporin and pirlimycin using UPLC-MS/MS (Agilent
865 1290 UPLC coupled with Agilent 6490 Triple Quad tandem mass spectrometry).
866 A gradient elution program consisting of two mobile phases (mobile phase A:
867 0.1% formic acid in water; mobile phase B: 0.1% formic acid in methanol) were
868 used at a flow rate of 0.5 mL/min. The concentration of cephalosporin and pirlimycin
869 in manure samples was quantified using the calibration curve of seven matrix-
870 matched standards (0.5, 1, 2, 4, 5, 10, and 20 µg L⁻¹ matrix solution). Matrix-
871 match standards were prepared using the SPE cleaned-up extracts of blank
872 manure samples.

873 *Microbial community composition and statistical analysis:*

874 In order to determine the effect that antibiotic exposure has on microbial
875 community composition, bacterial and fungal DNA were analyzed. DNA was
876 extracted using DNeasy PowerSoil Kits (Qiagen, Hilden, Germany). The
877 community composition was determined by amplifying the V4 region of the
878 bacterial/archaeal 16S rRNA gene as well as the ITS1 region of the fungal ITS
879 spacer region using primer pairs 515FB/806RB, and ITS1f/ITS2 respectively.
880 Caporaso *et al.* (78) was followed for amplification of 16S and ITS regions.
881 Multiplexing and sequencing was carried out using an Illumina MiSeq, producing
882 250 base pair paired-end reads (78). A UPARSE pipeline was used for quality
883 filtering and for clustering into OTUs – operational taxonomic units (79).
884 Additionally, all chimeric sequences were identified and removed using UCHIME
885 (80). The Ribosomal Database Project Native Bayesian Classifier was used to
886 assign OTUs to 269 specific taxonomies with the OTU cutoff for clustering of
887 97% (81). This was carried out using the GreenGenes 13.8 reference database
888 for the bacteria and archaea (82) and the UNITE 6.97 database for fungi (83).
889 Rarefaction of OTU tables and alpha diversity estimations were carried out using
890 the QIIME pipeline (84).

891

892 For ITS, only forward reads were used due to the size variability of the ITS
893 region. We processed ITS read sequences using the DADA2 pipeline (85), which
894 is designed to resolve exact biological sequences from Illumina sequence data
895 and does not involve sequence clustering (86). Sequences were trimmed to
896 uniform lengths, dereplicated, and the unique sequence pairs were denoised
897 using the 'dada' function, accounting for errors through the model generated with
898 the 'learnErrors' command. We removed chimeras and then assigned taxonomy
899 using the UNITE dynamic general release (ver 01.12.2017; 83) for fungi. To
900 account for differences in sequencing depths, we rarefied fungal samples to
901 36195 sequences per sample.

902

903 We compared microbial community composition using Primer-E (Ver. 7.0.13).
904 Microbial community data were square-root transformed before calculating
905 community dissimilarity between each treatment using Bray-Curtis dissimilarity.
906 These distances were used to generate ordinations (principal coordinates
907 analysis, PCoA) for both bacteria and fungi. Next, we performed PERMANOVA
908 with the community distance matrices to compare community composition using
909 treatments as a fixed effect, and block as a random effect using Primer-E (9999
910 permutations, Ver. 7.0.13; 65). We tested for homogeneity of dispersions from
911 the centroids via betadisper tests (89). To determine the potential OTUs
912 responsible for treatment differences, we first determined the percentage
913 contribution of taxa to overall Bray-Curtis dissimilarity using the SIMPER
914 (similarity percentages) command in Primer. We then identified common OTUs
915 that contributed to the top 20% of dissimilarity between treatment pairs and
916 analyzed each via ANOVA. Analyses were conducted in Primer v6 except
917 ANOVA which was conducted in R.

918

919 *Additional soil and microbial parameters:*

920

921 Soil pH was measured using a SensION+ PH3 laboratory pH probe (Hach,
922 Loveland, CO, USA). In addition to the CFE procedure described above, soil
923 microbial biomass was determined via substrate-induced respiration (SIR). The
924 SIR method is modified from West and Sparling (90) according to Fierer *et al.*
925 (91) and is considered a measure of active microbial biomass, whereas CFE
926 measures the total standing stock of microbial biomass. Briefly, SIR biomass was
927 determined by pre-incubating 4-g of dry weight equivalent soil at 20°C for 24-h.
928 Next, an excess of autolyzed yeast substrate (792-mg of yeast in dissolved 8-ml
929 of DI water) was added to each sample. The sample was then homogenized and
930 shaken for 1-hour, before the sample was capped and the headspace flushed
931 with CO₂-free air. Samples were then incubated for 5-h at 20°C, before respired
932 CO₂ accumulated in the headspace of each sample is measured using a gas
933 syringe and a bench-top infrared gas analyzer (IRGA, LI-7000 CO₂ H₂O
934 Analyzer, Li-Cor, Lincoln, NE).

935

936 Using the same basic outline as the SIR protocol, C mineralization (CMin) (46)
937 and catabolic response profile (CRP) (92) were also measured. In order to
938 determine the rate of C mineralization taking place in the soil – a measurement of
939 microbially accessible C – 6-g of dry weight equivalent soil was incubated at
940 20°C for 60-d. During this time samples were flushed with CO₂-free air and
941 incubated for 24-h at 20°C. Headspace was then measured using a bench-top
942 IRGA. Over this time water holding capacity was monitored and maintained at
943 approximately 65% - which is advantageous to microbial function. The integrals
944 between these periodic measurements are then extrapolated to determine
945 cumulative C mineralized over the 60-d period.

946

947 Catabolic response profiles (CRP) were used to measure the range of substrate
948 utilization capabilities of a given microbial community. This assay provides a

949 profile of responses to substrates, and helps to describe the metabolic
 950 capabilities of a given microbial community. To accomplish this, soils were
 951 weighed, substrate added, sample and substrate homogenized, headspace
 952 flushed, incubated at 20°C, and headspace measured in a fashion similar to the
 953 SIR protocol. However, instead of an autolyzed yeast solution substrate –
 954 glucose, glycine, oxalic acid, cellulose, chitin and water were all used as single
 955 substrates in individual assays. Each substrate was pH adjusted to 6, and their
 956 respective incubation times varied according to recalcitrance (*i.e.* cellulose and
 957 chitin were incubated for 24-h, all others were incubated for 4-h after shaking).
 958 Finally, all measurements are standardized to calculate the amount of respired C
 959 per quantity of soil and unit of time.

960

961 *Works Cited*

962

- 963 46. Lal R (2001) Assessment methods for soil carbon (Lewis Publishers, Boca
 964 Raton).
- 965 76. Ray P, Knowlton KF, Shang C, Xia K (2014) Development and validation
 966 of a UPLC-MS/MS method to monitor cephalosporin excretion in dairy cows
 967 following intramammary infusion. *PLoS One* 9(11):e112343.
- 968 77. Ray P, Knowlton KF, Shang C, Xia K (2014) Method development and
 969 validation: solid phase extraction-ultra performance liquid chromatography-
 970 tandem mass spectrometry quantification of pirlimycin in bovine feces and urine.
 971 *J AOAC Int* 97(6):1730–1736.
- 972 78. Caporaso JG, et al. (2012) Ultra-high-throughput microbial community
 973 analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* 6(8):1621–1624.
- 974 79. Edgar RC (2013) UPARSE: highly accurate OTU sequences from
 975 microbial amplicon reads. *Nat Methods* 10:996.
- 976 80. Edgar RC, Haas BJ, Clemente JC, Quince C, Knight R (2011) UCHIME
 977 improves sensitivity and speed of chimera detection. *Bioinformatics* 27(16):2194–
 978 2200.
- 979 81. Wang Q, Garrity GM, Tiedje JM, Cole JR (2007) Naïve Bayesian classifier
 980 for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl*
 981 *Environ Microbiol* 73(16):5261–5267.
- 982 82. DeSantis TZ, et al. (2006) Greengenes, a chimera-checked 16S rRNA
 983 gene database and workbench compatible with ARB. *Appl Environ Microbiol*
 984 72(7):5069–5072.
- 985 83. Nilsson RH, et al. (2019) The UNITE database for molecular identification
 986 of fungi: Handling dark taxa and parallel taxonomic classifications. *Nucleic Acids*
 987 *Res* 47(D1):D259–D264.
- 988 84. Caporaso JG, et al. (2010) QIIME allows analysis of high-throughput
 989 community sequencing data. *Nat Methods* 7(5):335–336.
- 990 85. Callahan BJ, et al. (2016) DADA2: High-resolution sample inference from
 991 Illumina amplicon data. *Nat Methods* 13(7):581–583.
- 992 86. Leff JW, et al. (2018) Predicting the structure of soil communities from
 993 plant community taxonomy, phylogeny, and traits. *ISME J* 12(7):1794–1805.


- 994 87. Abarenkov K, et al. (2010) The UNITE database for molecular
995 identification of fungi - recent updates and future perspectives. *New Phytol*
996 186(2):281–285.
- 997 88. Anderson MJ, Gorley RN, Clarke KR (2008) PERMANOVA+ for PRIMER:
998 Guide to Software and Statistical Methods. Plymouth, UK
999 doi:10.13564/j.cnki.issn.1672-9382.2013.01.010.
- 1000 89. Clarke KR, Gorley RN (2006) PRIMER v6: User Manual/Tutorial. Prim
1001 Plymouth UK. doi:10.1111/j.1442-9993.1993.tb00438.x.
- 1002 90. West AW, Sparling GP (1986) Modifications to the substrate-induced
1003 respiration method to permit measurement of microbial biomass in soils of
1004 differing water contents. *J Microbiol Methods* 5(3–4):177–189.
- 1005 91. Fierer N, Schimel JP, Holden PA (2003) Variations in microbial community
1006 composition through two soil depth profiles. *Soil Biol Biochem* 35(1):167–176.
- 1007 92. Degens BP, Harris J a. (1997) Development of a physiological approach
1008 to measuring the catabolic diversity of soil microbial communities. *Soil Biol*
1009 *Biochem* 29(9–10):1309–1320.
- 1010
1011

1012 *Supplementary Figures:*
1013
1014 Figure S1.




1015

1016 Figure S2.
1017



1018
1019

1020 Figure S3.



1021
1022
1023

1024 Supplementary Tables:

1025

1026 Table S1.

Treatment	Cellulose	Chitin	Glucose	Glycine	Oxalic Acid	All Substrates	CMin
NMC	0.68±0.04	0.41±0.02	0.93±0.05	0.55±0.03	0.45±0.02	3.03±0.1	824.69±22.5
Con	0.84±0.03	0.48±0.02	1.18±0.06	0.62±0.03	0.55±0.02	3.66±0.1	1112.29±42.3
Ceph	0.89±0.05	0.54±0.02	1.34±0.07	0.68±0.03	0.54±0.02	4.00±0.1	1190.83±55.6
Pir	0.85±0.05	0.48±0.02	1.11±0.05	0.64±0.03	0.52±0.03	3.61±0.1	1144.89±49.4

1027

1028 Table S2

Treatment	Day	Aboveground – C	Aboveground – N	Aboveground – C:N	Belowground – C	Belowground – N	Belowground – C:N	Microbial – C	Microbial – N	Microbial – C:N	Respiration – C
NMC	1	235.3±33	8.2±0.9	28.3±1.2	327±46	11.9±1.9	27.6±0.9	1.5±0.2	0.3±0.03	7±0.8	2.8±0.3
	2	311.9±33	11.3±1.1	27.6±1.2	353.4±27	13.8±2.5	28±3.0	1.3±0.2	0.2±0.03	8.7±1.1	2.3±0.3
	7	362.7±62	12.6±2.6	29.6±1.0	532.2±63	13.8±2.7	42±5.7	1.5±0.1	0.2±0.03	7.8±0.5	3.6±0.3
Con	1	329.3±56	15.2±2.6	21.8±2.2	347.3±73	15.1±3.6	23.5±1.3	2.0±0.2	0.3±0.04	7.0±0.4	3.0±0.5
	2	377.5±42	16.7±1.7	22.7±1.6	417.1±29	16.5±1.5	25.9±2.1	1.3±0.2	0.2±0.03	8.9±1.3	2.1±0.1
	7	466.2±28	19±2.1	25.2±1.6	597.3±92	20.4±2.5	30.6±5.2	1.8±0.3	0.2±0.05	8.3±1.2	4.5±0.1
Ceph	1	270.3±42	10.9±1.7	25±1.0	318.9±55	13.3±2.3	24.3±1.3	1.9±0.4	0.3±0.06	10.1±2.5	2.6±0.6
	2	426.1±32	18.8±1.5	22.8±0.9	466±55	16.9±1.7	27.3±1.3	1.1±0.2	0.2±0.03	10.2±2.9	1.7±0.2
	7	448.8±62	20.7±1.7	21.4±1.9	497.8±45	20.4±2.1	24.7±1.1	1.5±0.1	0.2±0.02	6.7±0.6	4.9±0.3
Pir	1	308.8±47	13.8±1.6	22.1±1.2	367.2±52	15.5±2.6	24.2±1.3	1.5±0.2	0.2±0.02	11.4±1.4	3.2±0.5
	2	368.8±31	14±1.3	26.5±0.9	427.8±61	17.6±3.0	25.2±1.6	1.3±0.1	0.2±0.02	8.3±0.8	2.3±0.2
	7	495.0±44	19.9±1.4	25.2±2.5	657±109	20±3.5	34.4±4.1	1.6±0.2	0.2±0.03	7.7±0.4	5.4±0.5
Treat	<i>P</i>	0.047	<0.001	<0.001	0.553	0.065	0.022	0.071	0.080	$\chi^2=0.364$	0.044
Day	<i>P</i>	<0.001	<0.001	0.624	<0.001	0.018	0.001	<0.001	0.028	$\chi^2=0.036$	<0.001
T*D	<i>P</i>	0.869	--	0.137	--	0.940	--	--	--	--	0.068
Treatment Pairwise Significance		Con>NMC Ceph>NMC* Pir>NMC	All>NMC	NMC>Con NMC>Pir	--	Con>NMC Ceph>NMC Pir>NMC*	NMC>All	Con>NMC Con>Pir	Con>Pir Ceph>Pir	--	Pir>NMC Pir>Ceph*

1030
1031

Table S3:

Treatment	Day	POM – C	POM – N	POM –C:N	Mineral - C	Mineral - N	1032
							1033 Mineral – C:N
NMC	1	708.4±63.9	45.4±4.4	15.7±0.3	1607.8±156.4	166.1±16.9	9.7±0.1
	2	672.6±59.2	45.0±5.2	15.1±0.4	1586.1±150.3	163.7±15.4	9.7±0.1
	7	611.8±52.1	39.7±4.0	15.5±0.2	1419.6±76.2	150.7±7.8	9.4±0.1
Con	1	822.5±72.0	49.1±4.6	16.8±0.3	1440.2±113	148.9±12.2	9.7±0.1
	2	663.5±34.7	40.5±2.6	16.5±0.3	1422.4±71.6	144.5±7.1	9.8±0.1
	7	666.1±63.1	42.4±4.2	15.7±0.2	1453.8±131.8	153.6±11.3	9.4±0.2
Ceph	1	886.8±125.9	54.7±8.7	16.5±0.6	1669.8±109.1	169.7±11.2	9.9±0.1
	2	821.4±86.5	49.0±6.1	17.0±0.7	1489.5±142.8	149.8±13.7	9.9±0.1
	7	508.7±79.5	49.3±6.4	16.7±0.7	1485.6±124.2	148.2±14.3	10.1±0.5
Pir	1	731.6±56.0	42.2±3.6	17.4±0.4	1544.2±109.0	159.6±10.4	9.7±0.1
	2	774.6±55.4	44.0±2.9	17.6±0.2	1467.1±53.0	147.7±6.8	10.0±0.1
	7	756.7±62.0	46.0±4.4	16.6±0.4	1473.3±122.5	152.3±12.2	9.7±0.1
Treat	<i>P</i>	0.002	0.037	0.001	0.150	0.152	$\chi^2=0.182$
Day	<i>P</i>	0.115	0.311	0.168	0.058	0.037	--
T*D	<i>P</i>	0.575	0.605	0.369	0.459	0.273	--
Treatment Pairwise Significance (* Marginal)		Ceph>NMC Ceph>Pir	Ceph>NMC* Ceph>Pir	Con>NMC Pir>NMC Pir>Ceph*	--	--	--

1034 Table S4.

Treat	Day	AG- ¹³ C-C ⁻¹	AG- ¹⁵ N-N ⁻¹	BG- ¹³ C-C ⁻¹	BG- ¹⁵ N-N ⁻¹	POM- ¹³ C-C ⁻¹	POM- ¹⁵ N-N ⁻¹	Min- ¹³ C-C ⁻¹	Min- ¹⁵ N-N ⁻¹	Resp- ¹³ C-C ⁻¹
NMC	1	0.015±1.1E-3	0.267±0.03	2.1E-3±6.3E-4	0.167±0.03	9.9E-4±4.2E-4	0.019±5.0E-3	6.5E-4±2.2E-4	0.015±2.8E-3	0.077±0.1
	2	0.012±1.4E-3	0.387±0.05	2.8E-3±8.2E-4	0.218±0.04	5.6E-4±3.1E-4	0.019±3.4E-3	5.6E-4±2.5E-4	0.016±2.0E-3	1.14±0.1
	7	9.1E-3±1.2E-3	0.398±0.05	1.7E-3±5.6E-4	0.266±0.05	6.0E-4±3.1E-4	0.027±6.3E-3	6.7E-4±2.5E-4	0.016±4.4E-3	1.91±0.5
Con	1	0.017±2.1E-3	0.214±0.02	1.8E-3±4.0E-4	0.165±0.01	2.9E-4±2.1E-4	0.029±3.7E-3	2.6E-4±9.8E-5	0.024±3.0E-3	0.393±0.1
	2	0.012±6.2E-4	0.231±0.20	1.9E-3±6.4E-4	0.135±0.02	4.6E-4±3.5E-4	0.022±5.2E-3	4.4E-4±2.6E-4	0.017±2.4E-3	0.601±0.1
	7	7.3E-3±1.2E-3	0.250±0.04	8.4E-4±4.8E-4	0.134±0.04	3.6E-4±1.3E-4	0.018±4.4E-3	5.2E-4±1.2E-4	0.012±2.4E-3	1.54±0.5
Ceph	1	0.015±1.8E-3	0.183±0.02	3.4E-3±8.9E-4	0.135±0.02	6.3E-4±4.3E-4	0.023±5.4E-3	4.4E-4±2.0E-4	0.018±5.0E-3	0.316±0.1
	2	0.015±2.1E-3	0.238±0.02	3.6E-3±1.5E-3	0.154±0.02	1.1E-3±6.9E-4	0.017±2.8E-3	3.1E-4±1.7E-4	0.015±1.2E-3	0.607±0.2
	7	8.2E-3±1.3E-3	0.260±0.06	1.5E-3±5.5E-4	0.133±0.04	3.2E-4±2.3E-4	0.025±4.0E-3	3.0E-4±1.6E-4	0.013±1.9E-3	2.75±0.4
Pir	1	0.017±5.0E-4	0.226±0.02	1.1E-3±4.3E-4	0.139±0.03	2.2E-4±1.1E-4	0.028±5.0E-3	2.0E-4±9.7E-5	0.019±3.4E-3	0.993±0.2
	2	0.014±2.0E-3	0.277±0.03	3.3E-3±9.2E-4	0.177±0.03	9.0E-4±2.5E-4	0.033±4.9E-3	6.1E-4±9.5E-5	0.020±2.7E-3	1.44±0.2
	7	0.010±1.9E-3	0.334±0.03	1.2E-3±5.6E-4	0.277±0.05	2.1E-4±2.1E-4	0.039±6.9E-3	1.2E-4±5.9E-5	0.024±3.2E-3	2.44±0.5
Treat	<i>P</i> =	0.54	>0.001	0.150	0.009	0.480	0.005	0.041	0.068	0.027
Day	<i>P</i> =	>0.001	>0.005	0.010	0.088	0.261	0.397	0.621	0.403	0.001
Treatment Pairwise Significance (* Marginal)		--	<i>NMC>All</i> <i>Pir>Con*</i> <i>Pir>Ceph</i>	--	<i>NMC>Con</i> <i>NMC>Ceph</i> <i>Pir>Con</i> <i>Pir>Ceph</i>	--	<i>Pir>All</i>	<i>NMC>Con*</i> <i>NMC>Ceph</i> <i>NMC>Pir</i>	<i>Pir>NMC</i> <i>Pir>Ceph</i>	<i>NMC>Con</i> <i>Pir>Con</i> <i>Pir>Ceph</i>

1035

1036

1037 Table S5.

	%C	pw	%N	pw	C:N	pw	Cephapirin (ng g manure⁻¹)	Pirlimycin (ng g manure⁻¹)
Con	49.9	b	3.3	a	15.3	b	-	-
Ceph	49.0	a	3.4	a	14.3	a	Below detection (<0.36)	-
Pir	49.6	b	3.5	a	14.3	a	-	149 ± 3.38
SE	0.12		0.06		0.26			

1038

1039 **Supplementary Figure and Table Legends:**

1040

1041 **Figure S1:** Plexiglass box for ^{13}C pulse-chase experiment, with wooden frame
1042 and rubber liner for sealing the chamber to the ground. The wooden frame was
1043 trenched 10-cm into the ground to minimize any leakage of ^{13}C - CO_2 from the
1044 pulsing area, as well as any non-labeled CO_2 entering the pulsing area.

1045

1046 **Figure S2:** Effect of antibiotic exposure on microbial activity and active microbial
1047 biomass. **a)** Substrate induced respiration (SIR) by treatment, and **b)** Catabolic
1048 response profile (CRP) by substrate and treatment (N = 24 for each treatment).
1049 SIR showed a significant treatment effect ($F_{3,87} = 4.26$, $P < 0.01$), with treatments
1050 receiving manure from cephalosporin treated cattle having active microbial biomass
1051 significantly greater than the no-manure control (NMC; $P < 0.001$) and control
1052 manure (Con; $P < 0.05$), and marginally greater than manure from pirlimycin
1053 treated cattle (Pir; $P = 0.10$). A significant treatment effect was observed for
1054 glucose ($F_{3,92} = 8.27$, $P < 0.001$), cellulose ($F_{3,87} = 5.14$, $P < 0.005$), glycine ($F_{3,15}$
1055 $= 2.53$, $P = 0.097$), oxalic acid ($F_{3,15} = 4.02$, $P < 0.05$), and chitin ($F_{3,87} = 8.00$, $P <$
1056 0.001). Across CRP the Ceph treatment is higher than all other treatments,
1057 though not consistently significantly different when compared pairwise. This
1058 provides evidence that the exposure to manure from cephalosporin treated cattle can
1059 cause an increase in microbial activity – consistent with previous findings.
1060 Increased microbial activity has implications for microbial efficiency and soil C
1061 storage.

1062

1063 **Figure S3:** Nonmetric multidimensional scaling for fungal communities across
1064 antibiotic treatments. Distances are based on dissimilarity matrices of sequence-
1065 based Bray-Curtis distances. Fungal communities across treatments do not differ
1066 significantly from each other (PERMANOVA Fungal: pseudo- $F_{3,15} = 1.10$, $P =$
1067 0.18 , Stress = 0.16).

1068

1069 **Table S1:** Catabolic response profile (CRP) average respired C (mean; $\text{CO}_2\text{-C}$
1070 ($\mu\text{g g dry wt soil}^{-1} \text{h}^{-1}$)) by substrate type (cellulose, chitin, glucose, glycine, oxalic
1071 acid, and summed respiration across all substrates) and treatment (NMC: no
1072 manure control; Con: control manure; Ceph: manure from cephalosporin treated
1073 cattle; Pir: manure from pirlimycin treated cattle. Additionally, microbially
1074 mineralizable C (CMin; $\text{CO}_2\text{-C}$ ($\mu\text{g g dry wt soil}^{-1} \text{h}^{-1}$)). Error listed is standard
1075 error.

1076

1077 **Table S2:** C, N, and C:N of aboveground biomass (Aboveground; $\text{g}\cdot\text{m}^{-2}$),
1078 belowground biomass (Belowground; $\text{g}\cdot\text{m}^{-2}$), microbial biomass (Microbial; $\text{g}\cdot\text{m}^{-2}$),
1079 and ecosystem respiration ($\text{g CO}_2\text{-C m}^{-2} \text{h}^{-1}$). These were measured at three
1080 time points (Days 1, 2, and 7) across four treatments (NMC: no manure control;
1081 Con: control manure; Ceph: manure from cephalosporin treated cattle; Pir: manure
1082 from pirlimycin treated cattle). The values listed are means and associated
1083 standard errors. Linear mixed models (LMM) and type III analysis of variance
1084 (ANOVA) were used to analyze the data with the exception of the Mineral-C

1085 which was analyzed using generalized linear model (GLM) and type II ANOVA –
 1086 Wald χ^2 Test in order to address failure to pass assumptions necessary for LMM.
 1087 As a result, these values are reported as χ^2 as opposed to *P*-values. The additive
 1088 models all included treatment (Treat) and time (Day) as fixed effects and Block
 1089 as a random effect. Aboveground-N, Belowground-C, Belowground-C:N, and
 1090 Microbial-N data were log transformed to meet normality assumptions. Significant
 1091 and marginally significant *P*-values are designated by italic and bold font.

1092
 1093 **Table S3:** C, N, and C:N of particulate organic matter (POM; g-m⁻²) and mineral
 1094 associated (Mineral; g-m⁻²) soil pools. These were measured at three time points
 1095 (Days 1, 2, and 7) across four treatments (NMC: no manure control; Con: control
 1096 manure; Ceph: manure from cephalixin treated cattle; Pir: manure from pirlimycin
 1097 treated cattle). The values listed are means and associated standard errors.
 1098 Linear mixed models (LMM) and type III analysis of variance (ANOVA) were
 1099 used to analyze the data with the exception of the Mineral-C:N which was
 1100 analyzed using generalized linear model (GLM) and type II ANOVA – Wald χ^2
 1101 Test in order to address failure to pass assumptions necessary for LMM. As a
 1102 result, these values are reported as χ^2 as opposed to *P*-values. Models used
 1103 were either interactive or nested depending on the best model AIC. For factors
 1104 best analyzed with interactive models all *P*-values are reported – for nested
 1105 models only treatment *P*-value is reported. For interactive models treatment
 1106 (Treat) and time (Day) and interactive (T*D) are used as fixed effects and Block
 1107 as a random effect. For the nested model plot is nested within block, thus
 1108 accounting for the repeated sampling over time. POM-C:N data were log
 1109 transformed to meet normality assumptions. Significant and marginally significant
 1110 *P*-values are designated by italic and bold font.

1111
 1112 **Table S4:** Showing the fate of added isotopic C (¹³C) and N(¹⁵N) of the various
 1113 pools measured: aboveground biomass (AG) belowground biomass (BG)
 1114 particulate organic matter (POM), mineral associated (Min), and ecosystem
 1115 respiration (for C only). Isotopic content is reported as the amount of label
 1116 recovered as a proportion of the total element pool – as a proportion therefore
 1117 unitless. These were measured at three time points (days 1, 2, and 7) across four
 1118 treatments (NMC: no manure control; Con: control manure; Ceph: manure from
 1119 cephalixin treated cattle; Pir: manure from pirlimycin treated cattle). The values
 1120 listed are means and associated standard errors. Linear mixed models (LMM)
 1121 and type III analysis of variance (ANOVA) were used to analyze the data. The
 1122 models all included Treatment and Day as fixed effects and Block as a random
 1123 effect. AG-¹⁵N-N⁻¹ data were log transformed to pass normality assumptions.
 1124 Model quality analysis using AIC favored additive models over interactive ones.
 1125 Significant *P*-values are designated by bold and italic font.

1126
 1127 **Table S5:** Manure characteristics for control (Con), cephalixin (Ceph), and
 1128 pirlimycin (Pir) treated cattle. Pairwise comparisons are listed according to *P*-
 1129 values < 0.05. Model standard error is also noted (SE). Statistical tests were not
 1130 performed for antibiotic concentrations.