

Integration of motion and form cues for the perception of self-motion in the human brain

Article

Other

Figure 1

Kuai, S.-G., Shan, Z.-K.-D., Chen, J., Xu, Z.-X., Li, J.-M., Field, D. T. and Li, L. (2020) Integration of motion and form cues for the perception of self-motion in the human brain. *The Journal of Neuroscience*, 40 (5). pp. 1120-1132. ISSN 1529-2401 doi: <https://doi.org/10.1523/JNEUROSCI.3225-18.2019> Available at <http://centaur.reading.ac.uk/88279/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1523/JNEUROSCI.3225-18.2019>

Publisher: The Society for Neuroscience

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online




Figure S1. The classifier's decoding accuracy for the six stimuli as a function of the number of selected voxels (starting from 50 to the number that covers the minimum number of the activated voxels across all participants at a step of 10) for each ROI. The dotted lines represent the 95th percentile of the classifier's baseline decoding accuracies after 1000 shuffled tests. The solid lines represent the chance level. The error bars indicate SEs across 14 participants. *: $p < 0.05$ is from Tukey HSD tests that compared the decoding accuracy with the baseline decoding accuracy at each number of voxels.