

Enzymatic glycosylation involving fluorinated carbohydrates

Article

Accepted Version

Council, C. E., Kilpin, K. J., Gusthart, J. S. ORCID: <https://orcid.org/0000-0002-1529-9493>, Allman, S. A., Linclau, B. and Lee, S. S. (2020) Enzymatic glycosylation involving fluorinated carbohydrates. *Organic & Biomolecular Chemistry*, 18 (18). pp. 3423-3451. ISSN 1477-0520 doi: <https://doi.org/10.1039/D0OB00436G> Available at <https://centaur.reading.ac.uk/90489/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <http://dx.doi.org/10.1039/D0OB00436G>

To link to this article DOI: <http://dx.doi.org/10.1039/D0OB00436G>

Publisher: Royal Society of Chemistry

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Enzymatic glycosylation involving fluorinated carbohydrates

Claire E. Council^a, Kelly J. Kilpin^a, Jessica S. Gusthart^b, Sarah A. Allman^b, Bruno Linclau^a, and Seung Seo Lee^a

*

^a School of Chemistry, University of Southampton, Highfield, Southampton, SO17 1BJ, UK

^b Reading School of Pharmacy, University of Reading, Whiteknights, Reading, RG6 6UR, UK

Abstract

Fluorinated carbohydrates, where one (or more) fluorine atom(s) have been introduced into a carbohydrate structure, typically through deoxyfluorination chemistry, have a wide range of applications in the glycosciences. Fluorinated derivatives of galactose, glucose, *N*-acetylgalactosamine, *N*-acetylglucosamine, talose, fucose and sialic acid have been employed as either donor or acceptor substrates in glycosylation reactions. Fluorinated donors can be synthesised by synthetic methods or produced enzymatically from chemically fluorinated sugars. The latter process is mediated by enzymes such as kinases, phosphorylases and nucleotidyltransferases. Fluorinated donors produced by either method can subsequently be used in glycosylation reactions mediated by glycosyltransferases, or phosphorylases yielding fluorinated oligosaccharide or glycoconjugate products. Fluorinated acceptor substrates are typically synthesised chemically. Glycosyltransferases are most commonly used in conjunction with natural donors to further elaborate fluorinated acceptor substrates. Glycoside hydrolases are used with either fluorinated donors or acceptors. The activity of enzymes towards fluorinated sugars is often lower than towards the natural sugar substrates irrespective of donor or acceptor. This may be in part attributed to elimination of the contribution of the hydroxyl group to the binding of the substrate to enzymes. However, in many cases, enzymes still maintain a significant activity, and reactions may be optimised where necessary, enabling enzymes to be used more successfully in the production of fluorinated carbohydrates. This review describes the current state of the art regarding chemoenzymatic production of fluorinated carbohydrates, focusing specifically on examples of the enzymatic production of activated fluorinated donors and enzymatic glycosylation involving fluorinated sugars as either glycosyl donors or acceptors.

1 Introduction

Fluorinated carbohydrates have been employed for a variety of applications in the fields of biochemistry, structural biology, cell biology and immunology.¹⁻⁶ They have found uses as mechanistic probes for kinetic^{7,8} and structural studies⁹ to help understand protein-carbohydrate binding and catalytic mechanisms of enzymes. The *in-vitro* post-translational modification of proteins with fluorinated carbohydrates has been demonstrated to be an effective tool to examine the roles of glycans using ¹⁹F NMR.² Additionally, positron emission tomography (PET) allows non-invasive and non-destructive *in vivo* analysis of carbohydrates incorporating ¹⁸F in biological systems.⁵ Approaches such as these are advantageous as they allow the specific behaviour of the fluorinated species to be analysed without competing signals arising from other components of the biological milieu.^{10,11} Fluorination has also been shown to endow carbohydrate structures with increased resistance towards degradative enzymes.¹² This inherent stability makes fluorinated carbohydrates attractive components for glycoconjugate vaccines.¹³

Chemo-enzymatic approaches to the synthesis of natural and chemically modified carbohydrates are well established. The review of Li, McArthur and Chen¹⁴ provides an excellent summary of chemical and enzymatic transformations using natural and synthetic sugars. The reviews of Uhrig and Guo *et al.*¹⁵⁻¹⁶ provide overviews of synthetic strategies for fluorination of carbohydrates and methods of introducing stereoselective fluorination into monosaccharide structures. Enzymatic synthesis of fluorosugars has however not previously been the subject of a focussed extensive review. This work therefore focuses on examining the specific enzymes that accept fluorinated sugars as donors or acceptors and how such enzymes are employed to affect the transformation of larger and more complex carbohydrate structures.

The success of chemoenzymatic synthesis approaches with modified substrates relies upon the ability of the enzyme to both recognise and utilise such substrates despite changes in structure. Fluorine substitution has been shown to have only minimal effect on the conformation of sugars,¹⁷ and this has been shown to hold true even if sugars bear multiple fluorine substituents.¹⁸ However, as fluorine atoms function solely as (weak) hydrogen bond acceptors and not as a hydrogen bond donors this can influence the hydrogen bond properties of adjacent hydroxyl group.^{19,20}


Some enzymes display high substrate specificity. In some cases, even a single hydroxyl to fluorine substitution can prevent a sugar from acting as a substrate.²¹ Others are more tolerant to fluorinated sugars albeit with lower catalytic efficiency as discussed later in this article. Field and co-workers²² suggest that even at low efficiency, enzymatic biotransformations can still provide access to biomolecules that would be laborious to prepare via chemical synthesis as enzymes become more widely available.

Enzymes which catalyse glycosylation reactions typically require an activated donor such as a sugar-1-phosphate or sugar nucleotide mono- or diphosphate. The synthesis of fluorinated donor molecules can often also be achieved enzymatically, therefore enzymes which are capable of carrying out these conversions to achieve the full enzymatic conversion from fluorinated reducing sugar to glycan are also covered. It must also be noted these ‘intermediates’ are often interesting target molecules in their own right as many have been shown to be enzymatic inhibitors and have found applications as mechanistic probes.⁸

This review is organised by fluorinated sugar. For each named sugar, when and as appropriate, the enzymatic synthesis of activated sugar donors and their subsequent use in glycosylation reactions is first discussed, grouped by enzyme class, followed by glycosylation using the named sugar as acceptor, either as a monosaccharide or as part of a larger glycan, again organised by enzyme class. For the purposes of this review the nomenclature used for the fluorinated sugar derivatives does not distinguish between deoxy-fluorination and fluorination; 6FGal refers to 6-deoxy-6-fluorogalactose, whilst 6FFuc refers to fucose fluorinated at position 6. Where there is geminal difluorination, the original sugar name is maintained for clarity. Unless indicated, all sugars possess the D-configuration.

2 Glossary of glycoenzymes:

Carbohydrate synthesis and degradation *in vivo* is mediated by a suite of glycoenzymes. In this article, these glycoenzymes refer to any enzymes that synthesise or modify carbohydrate substrates and are limited to those that have been used with fluorinated carbohydrates. Thus, the enzyme groups presented in this glossary are classified differently from the CAZY database (<http://www.cazy.org>), and the classification follows the discussion in this article. The enzyme groups presented in this glossary are: kinases, pyrophosphorylases (or nucleotidyltransferases), glycosyltransferases, glycoside hydrolases and phosphorylases. A summary of the actions of key glycoenzymes in this article is presented in Scheme 1.



Scheme 1 Summary of key enzymes used in the formation of fluorinated donors and for enzymatic glycosylation.

Kinases

Kinases transfer a phosphate group from a high energy donor – typically adenosine triphosphate (ATP) – to the anomeric position of a reducing sugar with α -selectivity. Galactose kinase (GalK), N-acetylgalactosamine kinase (GalNAcK) and N-acetylhexosamine kinase (NahK) transfer phosphate to the 1-position of Gal, GalNAc and N-acetylhexoses respectively. Alternatively, hexokinase catalyses the transfer of phosphate to the 6-position of hexoses. This enzyme can be used with a phosphoglucomutase (PGM) which is capable of isomerising the phosphate group to the anomeric position, again with α -selectivity.

Pyrophosphorylases

Pyrophosphorylase (or nucleotidyltransferase) enzymes transfer nucleotide monophosphate (NMP) to the phosphosugar to give the nucleotide diphosphate (NDP) sugar. The mechanism involves the formation of a ternary complex where a nucleotide triphosphate (NTP) and the sugar-1P substrate successively bind into the active site. This allows nucleophilic attack by the anomeric phosphate moiety onto the first nucleotide linked phosphate group, which releases pyrophosphate and the sugar nucleotide.

Glycosyltransferases

Glycosyltransferases form a glycosylic bond between an activated sugar (usually nucleotidyl phosphate sugar) donor and a sugar acceptor such as monosaccharide, oligosaccharide or non-sugar molecule.

Glycosyltransferases are named for the donor sugar they transfer, so a galactosyltransferase (GalT) transfers activated galactose onto an acceptor, a sialyltransferase (SiaT) transfers activated sialic acid etc. Generally, GalTs utilise a uridine diphosphate (UDP) donor and SiaTs utilise cytosine monophosphate (CMP) -activated sialic acids as the sialic acid donor.

Glycoside hydrolases

As defined in CAZY database (<http://www.cazy.org>), glycoside hydrolases catalyse hydrolysis of glycosidic bonds. Although glycoside hydrolases are naturally degradative enzymes, under certain conditions some enzymes in this class can be directed to the synthetic reaction via reverse hydrolysis or transglycosidation.²³ Some glycoside hydrolases (such as the trans-sialidase from *Trypanosoma cruzi*) are known to have higher synthetic activity than hydrolytic activity, with the active site more suitably accommodating a non-water nucleophile such as a sugar molecule. Trans-sialidases transfer sialic acid from a donor oligosaccharide containing a terminal sialic acid, such as the heavily sialylated glycoprotein fetuin, to another Gal terminated acceptor with α -(2→3) configuration.²⁴ In synthetic applications, trans-sialidases have been used as an alternative to sialyltransferases to introduce sialic acid to carbohydrate structures.

Phosphorylases

Phosphorylases catalyse the reversible formation of a monosaccharide-1-phosphate from an oligosaccharide and an inorganic phosphate. The reverse reaction can be used to synthesize oligosaccharides.


3. Fluorinated Galactose

3.1 Fluorinated galactose-1-phosphate and UDP-galactose donors

Early evidence that deoxyfluorogalactose-1-phosphate sugar derivatives could be synthesised enzymatically from the corresponding deoxyfluorogalactopyranose came from work carried out by Kent and Wright.²⁵ In this study, they demonstrated the conversion of 6FGal to 6FGal-1P using a yeast-derived galactokinase (GalK) in the presence of ATP. Westwood *et al.*²⁶ later evidenced that 2FGal, 3FGal and 4FGal could be converted by the same enzyme to 2FGal-1P, 3FGal-1P and 4FGal-1P respectively. The results of their kinetic study suggests the yeast-derived GalK has lower affinity for the four fluorinated substrates when compared with the natural Gal substrate (in the order Gal > 2FGal > 3FGal ≈ 6FGal << 4FGal).


These early findings were later verified by other groups who exploited GalK to synthesise the anomeric phosphates 2FGal-1P, 3FGal-1P and 6FGal-1P on a larger scale.^{21-22, 27} In these studies, a one-pot, two-enzyme system was employed, with the anomeric phosphate converted directly to the UDP-sugar by the pyrophosphorylase Gal-1P uridylyltransferase (GalPUT).

Hayashi *et al.*²¹ reported that yeast GalK catalysed the synthesis of 2FGal-1P in preparative quantities from 2FGal (Scheme 2). Stereoselective conversion to the α -anomer was confirmed by NMR spectroscopy. The isolated 2FGal-1P was then converted to UDP-2FGal using GalPUT, coupled with a UDP-Glc regeneration cycle, which avoids the requirement for stoichiometric quantities of commercially available but costly UDP-Glc. The direct synthesis of α -UDP-2FGal from 2FGal was also possible without isolating the phosphate intermediate.



Scheme 2 Enzymatic synthesis of UDP-2FGal using a UDP-Glc regeneration cycle.²¹


Enzymatic synthesis of fluorinated nucleotide donors was also demonstrated by Barlow and Blanchard²⁸. In this work, the authors describe the synthesis of both UDP-2FGal and UDP-3FGal from 2FGal and 3FGal respectively in preparative scale at near quantitative yields (based upon UDP-Glc conversion). This study also employs an alternative strategy to overcome problems associated with the purification of the desired UDP-sugar from the UDP-Glc generated in the catalytic cycle (Scheme 3). By using phosphoglucomutase (PGM) in place of UDP-Glc pyrophosphorylase the reaction was driven toward UDP-2FGal. The UDP-Glc cofactor could be removed from the system via further enzymatic breakdown, using glucose-6-phosphate dehydrogenase. This system does however require near stoichiometric quantities of costly UDP-Glc.



Scheme 3 Enzymatic synthesis of UDP-2FGal and UDP-3FGal with enzymatic breakdown of UDP-Glc.²⁸

The utility and substrate scope of the GalK/GalPUT two-enzyme system was further expanded by Errey *et al.*²² In this work, the authors describe that UDP-6FGal can be synthesised from 6FGal using the GalK/GalPUT system (Scheme 4). Under identical reaction conditions, yields of UDP-2FGal were comparable to those achieved for the synthesis of UDP-Gal. In both cases near quantitative conversion was observed. In comparison, UDP-3FGal (64%) and UDP-6FGal (37%) were isolated in lower yields than the UDP-2FGal and the nonfluorinated product. Higher enzyme concentrations and longer reaction times were required for the synthesis of both UDP-3FGal and UDP-6FGal than for UDP-Gal and UDP-2FGal. It was unclear to the authors from their one-pot experiments which enzyme was rate limiting, but further experiments using chemically synthesised or commercially available sugar-1-phosphates of non-galactose sugars indicated that GalPUT has a relaxed substrate specificity and is able to turn over substrates not accepted by the two-enzyme system. This suggests GalK to be the rate limiting enzyme in the two-enzyme system. Indeed, these observations correlate with the


early kinetic findings presented by Bessell *et al.*,²⁹ who found GalK to have a greater affinity for 2FGal than 3FGal and 6FGal. In addition to using the UDP-Glc regeneration cycle to aid product purification, Errey and coworkers²² made use of GalK and GalPUT immobilised onto resin surfaces which offered the additional benefits of allowing facile enzyme recovery and simplified purification methods. This approach also meant the enzymes could be used in excess, allowing even reactions with low efficiency compared to natural substrates to be successfully achieved on a preparative scale.



Scheme 4 Enzymatic synthesis of UDP-2FGal UDP-3FGal and UDP-6FGal using immobilised enzymes and the UDP-Glc regeneration cycle.²²


The synthesis of UDP-4FGal from 4FGal was reported in the 2012 patent application of Nishimura *et al.* in two steps using *E. coli* derived GalK (*EcGalK*) and GalPUT (Scheme 5).³⁰ The authors noted that the conversion for the second step (i.e. the formation of UDP-4FGal from 4FGal-1P) was 30% lower if using GalPUT from yeast instead of *E. coli* was used, which highlights the value in considering enzymes from different sources.

Likewise, although initial kinetic studies using yeast GalK indicated a four-fold increase in K_m for 4FGal relative to Gal and a V_{max} of 10% relative to Gal²⁹, the authors reported a surprisingly short reaction time (1 h) using *E. coli* GalK.




Scheme 5 Enzymatic synthesis of UDP-4FGal.³⁰

6,6-Difluoro-galactose has been shown to be a substrate for GalK from *Lactococcus lactis*, a GalK demonstrated to have good substrate promiscuity against a large panel of sugars, with modifications at the 2, 3, 4 or 6 position (Scheme 6).³¹



Scheme 6 Enzymatic phosphorylation of 6,6-difluoro-galactose.³¹

A series of furanosyl sugar nucleotides have been synthesised using a bacterial α -D-glucopyranosyl-1-phosphate thymidyltransferase, Cps2L from *Streptococcus pneumoniae*.³² In these reactions, chemically synthesised 6-fluoro-galactofuranosyl-1-phosphate (anomeric mixture $\alpha/\beta = 1.7/1$) was successfully converted into deoxythymidine diphosphogalactofuranose (dTDP-Galf) in the presence of Cps2L, deoxythymidine 5'-triphosphate, Mg^{2+} and inorganic pyrophosphatase (Scheme 7). The conversion was moderate, only 19% in a 24 hour reaction. Later, the authors engineered Cps2L to expand its repertoire of nucleotide substrates including uridine and guanidine.³³ One of mutants, Cps2L Q24S showed an enhanced activity in the conversion of 6-fluoro-galactofuranosyl-1-phosphate ($\alpha/\beta = 1.7/1$) to dTDP-Galf with a conversion rate of c.a. 30% in 18 hours although its intended activity of uridylyltransfer was not detected with 6FGalf (Scheme 7).



Scheme 7 Enzymatic synthesis of dTDP-6FGalf.³²⁻³³


3.2 Enzymatic glycosylation using fluorinated galactose donors

3.2.1 Galactosyltransferases

An understanding of the likely catalytic mechanism for galactosyltransferase enzymes along with metabolic studies using 2FGal (which causes inhibition of glycosylation),³⁴ lead Hayashi *et al.*²¹ to propose that a strongly electron withdrawing fluorine atom at C-2 would result in enzyme inhibition. They report the K_i ($149 \pm 18 \mu\text{M}$) versus K_m ($127 \pm 9 \mu\text{M}$) for UDP-Gal indicating that the affinity of UDP-2FGal for the enzyme is comparable to the natural substrate and demonstrate UDP-2FGal to be a competitive inhibitor of β -(1 \rightarrow 4) galactosyltransferase (GalT), with respect to UDP-Gal, fluorine at the 2-position unaffected binding to the enzyme, but making the glycosidic bond inert to cleavage.


Whilst β -(1 \rightarrow 4) GalT is inhibited by UDP-2FGal, it is capable of processing both the UDP-4FGal and UDP-6FGal donors. To the best of our knowledge, there are no published examples UDP-3FGal being used as a donor in glycosyltransferase catalysed reactions.

4FGal and its applications, including the synthesis of UDP-4FGal detailed earlier, are reported in the previously mentioned 2012 patent application³⁰ in which the use of UDP-4FGal as a donor for human derived β -(1 \rightarrow 4) GalT is described. Incubation of the enzyme with a variety of β -glycosides of GlcNAc, followed by mass spectral characterisation indicated the formation of a peak corresponding to the desired disaccharides. However, the regio- and stereoselectivity was not unambiguously confirmed nor reaction conditions optimised (Scheme 8).



Scheme 8 β -(1 \rightarrow 4) GalT enzymatic synthesis of disaccharides containing 4FGal.³⁰

UDP-6FGal (obtained by chemical methods) was shown by Kodama *et al.* to be a substrate for β -(1 \rightarrow 4) GalT from bovine milk, albeit at rates significantly lower than of UDP-Gal (Scheme 9).³⁵ Nonetheless, increasing the enzyme concentration and reaction time allowed the enzyme to transfer UDP-6FGal to the natural GlcNAc acceptor to produce the fluorinated LacNAc analogue in 59% yield, with the regio- and stereoselectivity confirmed by proton NMR spectroscopy. In an independent study³⁶ using the same enzyme and acceptor, the authors were unable to detect transfer of UDP-6FGal to GlcNAc and demonstrated its ability as a competitive inhibitor of the enzyme. However, they noted that enzyme concentrations and reaction times were significantly reduced compared with the earlier study. These conflicting observations highlight the importance of screening multiple reaction conditions, and show that although reaction rates can be low compared to natural sugars, under optimised conditions use of enzymes can be a viable method of synthesis.




Scheme 9 β -(1 \rightarrow 4) GalT enzymatic synthesis of disaccharides containing 6FGal.³⁵

3.2.2 Phosphorylases

Galactosyl 1-phosphate donors can be utilised by a phosphorylase from *Bifidobacterium infantis* (*BiGalHexNAcP*) for the synthesis of β -(1 \rightarrow 3) linked disaccharides. Furthermore, the phosphorylase enzyme along with a suitable kinase such as *EcGalK* can be used in a one-pot, two-enzyme system to afford the disaccharide directly from the reducing sugar.³⁷ This system has been successfully employed with certain monofluorinated sugars.


Enzymatically synthesised 6FGal-1P was shown by Yan *et al.* to be substrate for *BiGalHexNAcP* and was able to be transferred to the 3-OH of a GalNAc acceptor with β -selectivity (Scheme 10).³⁸ Preparative scale yields were high (ca. 80%) and comparable to the natural donor Gal-1P. Furthermore, the enzyme was demonstrated to transfer 6FGal-1P to the unnatural fluorinated 6FGalNAc acceptor at yields comparable to GalNAc, yielding disaccharide fluorinated in both sugar units (see section 5). *BiGalHexNAcP* has optimum activity at pH 5.0 to 6.5, whereas *EcGalK* has optimum activity at pH >7.0. When using natural donors and acceptors, pH 6.5 was a

good compromise giving reasonable overall yields, however for the fluorinated donors and acceptors, to maximise yields, a two-step process was used, at pH 7.5 for the kinase reaction, and then once the reaction was complete as monitored by TLC, at pH 6.0 with addition of *BiGalHexNAcP*.



Scheme 10 One-pot two-step disaccharide synthesis using 6FGal.³⁸


Whilst the authors initially reported that 2FGal-1P is not a substrate for the enzyme they later demonstrated that it can be transferred to either GalNAc or GlcNAc acceptors.³⁹ As the *Bifidobacterium infantis* kinase *BiGalK* has relaxed substrate specificity and good activity at pH 6.5 this removed the need for a two-step reaction (Scheme 11). Under the same reaction conditions, isolated yields (26-28%) were much lower than for Gal-1P and 6FGal-1P however this activity and transformation is important, as 2-fluorinated UDP donors are inhibitors of glycosyltransferases as previously described, therefore the enzymatic routes to the synthesis of disaccharides containing 2-deoxy-2-fluoro sugar moiety at the reducing end are limited. Furthermore, it is worthwhile noting that in this two-enzyme reaction, the low conversion is due to the phosphorylase enzyme rather than the kinase, as 2FGal-1P can be synthesised by both *EcGalK* and *BiGalK* in yields greater than 90% under the conditions used in this system.



Scheme 11 One-pot disaccharide synthesis using 2FGal.³⁹

3.2.3 Glycoside hydrolases

As fluorinated carbohydrates have been reported to act as mechanism-based inhibitors of glycoside hydrolases particularly as a donor,^{6, 12} it is not surprising to be able to find only few examples. O'Donohue et al. used p-nitrophenyl β -D-galactofuranoside (pNP- β Gal f) as a substrate of arabinofuranosidase from *Thermobacillus xylanilyticus* (AbfD3) to conduct transglycosidation reaction.⁴⁰ To avoid extensive hydrolysis, the authors stopped the reaction in 30 minutes, and detected two transglycosidation product, β -(1 \rightarrow 2) and β -(1 \rightarrow 3) disaccharides, both of which were spectroscopically characterised (Scheme 12). Although the conversion rates or yields were not reported, the HPLC chromatogram showed that hydrolysis products were predominant. In another example, Ferrieres et al. conducted an enzymatic reaction with arabinofuranosidase (Araf51) from *Clostridium thermocellum* containing galactosyl furanose derivatives as substrates.⁴¹ The original intention was to use pNP- β Gal f as donor and octyl 6-fluoro- β -D-galactofuranoside (Oct 6FGal f) as acceptor, but the reaction yielded a mixture of various length oligosaccharides with both substrates as either donor or acceptor (Scheme 12). A few oligosaccharide products showed that Oct 6FGal f acted as donor albeit with the yield of only 1 – 6%, which indicates that fluorination is not accepted well by the enzyme.




Scheme 12 Use of arabinofuranosidase to synthesise fluorinated di-galactofuranosides.⁴⁰⁻⁴¹

3.3 Fluorinated galactose acceptors

Fluorinated galactose has also been used as an acceptor in enzymatic glycosylation, not as the monosaccharide but as a non-reducing galactose moiety as part of a disaccharide or larger oligosaccharide.

3.3.1 Galactosyl- and *N*-acetylgalactosaminyltransferases

The disaccharide Fuc- α -(1 \rightarrow 2)Gal is a motif common to both A and B blood group antigens. In biosynthetic pathways, α -(1 \rightarrow 3) galactosyltransferase (α -(1 \rightarrow 3)GalT) is responsible for transferring a Gal residue from UDP-Gal to this motif to give the B group antigen whereas α -(1 \rightarrow 3) *N*-acetylgalactosaminyltransferase (α -(1 \rightarrow 3) GalNAcT) transfers a GalNAc residue to give the A antigen.⁴² Following chemical synthesis of the fluorinated acceptor analogue Fuc- α -(1 \rightarrow 2) 6FGal- β -O(CH₂)₇CH₃, biosynthesis of the modified antigens was achieved using both natural donors (Scheme 13),⁴³ however the enzymes α -(1 \rightarrow 3)GalT and α -(1 \rightarrow 3)GalNAcT had a lower affinity for the fluorinated acceptor than the natural acceptor (α -(1 \rightarrow 3)GalT 30% activity, α -(1 \rightarrow 3)GalNAcT 43% activity relative to the native disaccharide). Fuc- α -(1 \rightarrow 2)4FGal was not a substrate (or inhibitor) for either enzyme, demonstrating the importance of the 4-position hydroxyl for these enzymatic transformations. Fuc- α -(1 \rightarrow 2)-3FGal was shown to be a weak competitive inhibitor of both enzymes.




Scheme 13 6FGal as acceptor in enzymatic synthesis using galactosyl and *N*-acetylgalactosamine transferases.⁴³

3.3.2 Sialyltransferases and trans-sialidases

Fluorinated sugars have been utilised as acceptor substrates for sialylation using both sialyltransferases (SiaT) and trans-sialidases. The enzymatic incorporation of fluorinated sialic acid derivatives as donors is discussed in section 9.


The disaccharide 2FGal β -(1 \rightarrow 3) GalNAc is a suitable acceptor substrate for the α -(2 \rightarrow 3) *Pasteurella multocida* SiaT 1 (*PmST1*). The trisaccharide Neu5Ac α -(2 \rightarrow 3) 2FGal β -(1 \rightarrow 3) GalNAc α -O-(CH₂)₃N₃ was synthesised in preparative scale in excellent yields from the corresponding disaccharide acceptor 2FGal β -(1 \rightarrow 3) GalNAc α -O-(CH₂)₃N₃ and Neu5Ac using this enzyme, coupled with the CMP sialic acid synthetase from *Neisseria meningitidis* (*NmCSS*) in a one-pot, two-enzyme system (Scheme 14). Yields were comparable to the natural acceptor Gal β -(1 \rightarrow 3) GalNAc under the same reaction conditions.³⁹



Scheme 14 2FGal as acceptor in enzymatic synthesis using SiaT *PmST1*.³⁹


Incubation of dodecyl 2FGal, dodecyl 4FGal and dodecyl 6FGal with mouse melanoma B16 cells showed toxicity with 6FGal and 4FGal, however 2FGal produced a glycosylated product, corresponding to a sialylated product by mass spectrometric analysis.⁴⁴ This observation was confirmed through use of a α -(2 \rightarrow 3) sialidase from *S. typhimurium* (cloned in *E. coli*) to be a fluorinated analogue of the ganglioside GM-4 (Neu5Ac α -(2 \rightarrow 3))

Gal β ceramide) (Scheme 15), indicating that cellular α -(2 \rightarrow 3)-SiaT are able to utilise 2FGal as an acceptor for the synthesis.



Scheme 15 2FGal as acceptor for cellular α -(2 \rightarrow 3) SiaT from *S. typhimurium*.⁴⁴


Disaccharides terminated with 2FGal have also been shown to be suitable acceptors for the *T. cruzi* trans-sialidase (*TcTs*) mediated transfer of Neu5Ac from the heavily sialylated mammalian protein fetuin. A crude cell lysate can be used for the transformation, and using *TcTs* with fetuin as the Neu5Ac donor removes the requirement for CMP-Neu5Ac and the enzymes associated with its synthesis. Using this system, Allman *et al.*⁴⁵ demonstrated the synthesis of the trisaccharides Neu5Ac α -(2 \rightarrow 3) 2FGal β -(1 \rightarrow 4) GlcNAc and Neu5Ac α -(2 \rightarrow 3) 2FGal β -(1 \rightarrow 3) GlcNAc, from the corresponding chemically synthesised disaccharides, with the regio- and stereo-selectivity of the enzyme confirmed via NMR spectroscopy (Scheme 16). For both β -(1 \rightarrow 3) and β -(1 \rightarrow 4) linked disaccharides, yields for the sialylation were less than obtained for the natural disaccharides, although in the case of the natural β -(1 \rightarrow 4) disaccharide, the use of an OBn anomeric protecting group instead of OTBDMS at the reducing end could also affect the affinity of the enzyme for the acceptor.



Scheme 16 2FGal as acceptor in enzymatic synthesis using *TcTs*.⁴⁵


4FGal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2) Man was shown to be a substrate for both human and recombinant rat liver α -(2 \rightarrow 6) SiaT with a good sialylation rate relative to the natural trisaccharide Gal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2)

Man, and also a poor substrate for recombinant rat liver α -(2 \rightarrow 3) SiaT.⁴⁶ 3FGal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2) Man showed low-level sialylation with both the human and rat liver α -(2 \rightarrow 6) SiaT in screening reactions (Scheme 17). In scale-up synthesis, the fluorinated tetrasaccharides Neu5Ac α -(2 \rightarrow 3) 4FGal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2) Man and Neu5Ac α -(2 \rightarrow 6) 4FGal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2) Man were able to be synthesised in low milligram quantities from 4FGal β -(1 \rightarrow 4) GlcNAc β -(1 \rightarrow 2) Man using rat liver recombinant α 2-3 SiaT or rat liver α 2-6 SiaT respectively. Reactions were performed over 7 days with CMP-Neu5Ac added in batches every 2 days, along with calf intestinal alkaline phosphatase to break down the released CMP, which otherwise inhibits the sialyl transferase.⁴⁷




Scheme 17 Trisaccharides containing 3FGal or 4FGal as acceptors in enzymatic synthesis using α -(2 \rightarrow 3) and α -(2 \rightarrow 6) SiaTs.⁴⁶

Rat liver α -(2 \rightarrow 3) - and α -(2 \rightarrow 6) - SiaTs also tolerate 4-deoxy-4-fluorogalactoside acceptors with further modifications at the reducing end. The Neu5Ac α -(2 \rightarrow 3) 4FGal and Neu5Ac α -(2 \rightarrow 6) 4FGal glycosides depicted in Scheme 18 were detected via HPLC and mass spectrometry following reaction with the appropriate enzyme and CMP-Neu5Ac, however the connectivity was not unambiguously determined. In all examples, kinetic data revealed the reaction to proceed slower for the fluorinated donor than for the unmodified.³⁰



Scheme 18 Disaccharide containing 4FGal as acceptor in enzymatic synthesis using α -(2 \rightarrow 3) and α -(2 \rightarrow 6) SiaTs.³⁰

Completing the fluorinated galactose sugars, disaccharides containing 6FGal were shown to be excellent acceptors for the α -(2 \rightarrow 3) SiaT *PmST1* (Scheme 19). Combined with *NmCSS* in the one-pot, two-enzyme system introduced above, the α -(2 \rightarrow 3) linked sialosides have been synthesised in milligram quantities, in yields comparable to the natural acceptor under the same reaction conditions. Stereo- and regioselectivity of the sialylation were confirmed by spectroscopic methods.³⁸




Scheme 19 3FGal and 4FGal containing trisaccharides as acceptors in enzymatic synthesis using α -(2 \rightarrow 3) and α -(2 \rightarrow 6) SiaTs.³⁸

4. Fluorinated Glucose

The synthetic strategy toward deoxyfluoro- α -D-glucopyranosyl phosphates and UDP-deoxyfluoro- α -D-glucopyranoside donors differs from that described above for their galactosyl analogues. Although GalK tolerates the introduction of fluorine, it has a high stringency toward galactose stereochemistry and Westwood demonstrated many years ago that 4FGlc was neither a substrate nor an inhibitor of yeast GalK.²⁶ Even a mutant GalK, which is known to possess relatively relaxed substrate specificity at the C-4 position was unable to efficiently phosphorylate 3FGlc, 4FGlc or 6FGlc.⁴⁸ Indeed, to the best of our knowledge, no kinases have been reported in the literature that are capable of efficiently processing deoxyfluorinated glucopyranosides. With the lack of suitable kinases, other enzymes have been evaluated in order to enzymatically access deoxyfluoro- α -D-glucopyranosyl phosphates. In particular, the hexokinase/phosphoglucomutase approach has shown promise. Early kinetic studies revealed 2FGlc and 2,2FFGlc are good substrates for yeast hexokinase with affinities similar to Glc.²⁹ A hexokinase/phosphoglucomutase approach, followed by conversion with UDP-glucose-1-pyrophosphorylase, has been used in the synthesis of UDP-2FGlc (Scheme 20).¹¹ The UDP-glucose-1-pyrophosphorylase mediated step was investigated further, and it was found that the activity of this enzyme decreases by a factor of 30 for the 2FGlc-1P compared with Glc-1P, and a six-day reaction time was


required to achieve 60% conversion of the 2FGlc. Following purification, UDP-2FGlc could be isolated in milligram quantities in 38% yield. The authors then attempted to extend this methodology to the synthesis of ^{18}F radio-labelled UDP-2FGlc. However, limited radiochemical yields led them to employ a chemo-enzymatic protocol whereby the 2FGlc-1P was first synthesised chemically. Enzymatic conversion to UDP-2FGlc was carried out in the void volume of an anion exchange cartridge, and using increased concentrations of UDP-glucose-1-pyrophosphorylase, the product was isolated with a radiochemical yield of 20% with a total synthesis time of 110 minutes from 2FGlc.



Scheme 20 Enzymatic synthesis of UDP-2FGlc using hexokinase, phosphoglucomutase and UDP-Glc pyrophosphorylase.¹¹


On the other hand, 3FGlc and 4FGlc are poor substrates for yeast hexokinase.²⁹ Indeed, the hexokinase/PGM approach to synthesising 3FGlc-1P and 4FGlc-1P from 3FGlc and 4FGlc respectively was evaluated by Caputi *et al.* who found only a low conversion to the anomeric phosphates, and abandoned this pathway in favour of other more synthetically viable routes. Namely, they successfully demonstrated the combination of UDP-Gal synthesis with uridine-5'-diphosphogalactose-4-epimerase (GalE) for the synthesis of fluorinated UDP-sugars with the glucose configuration. UDP-6FGal was enzymatically synthesised using the GalK/GalPUT system described earlier.²² Subsequent incubation with GalE resulted in inversion of the stereochemistry at C-4' to give UDP-6FGlc, with the equilibrium reached after 3 hrs in favour of the *glc*-configured product (UDP-6FGlc:UDP-

6FGal 3:1) (Scheme 21).⁴⁸ The extension to the GalE approach to other members of the deoxyfluoro glucosyl pyranosides is yet to be explored, however the synthesis of UDP-4FGlc is obviously not possible using this methodology.



Scheme 21 Enzymatic synthesis of UDP-6FGlc using uridine-5'-diphosphogalactose-4-epimerase (GalE).⁴⁸

The feasibility of UDP-6FGlc as a donor for glycosyltransferase mediated synthesis was demonstrated using the oat root transferase SAD10, whose physiological role is the generation of *O*-glucosyl *N*-methylantranilate ester. Indeed, UDP-6FGlc (synthesised from UDP-6FGal by GalE as described earlier)⁴⁸ was a substrate for the SAD10, which transferred the 6FGlc to *N*-methylantranilic acid (MNA-H). The reaction did not proceed to completion, but instead an equilibrium was reached within 20 hours, which showed approximately 31% conversion of UDP-6FGlc into 6FGlc-NMA. This ratio was similar for the natural donor UDP-Glc.⁴⁸ Certain glycosyltransferases from bacterial sources have been revealed to catalyse their traditional reaction in reverse, and this offers an alternative strategy toward the synthesis of NDP-sugars that does not require multiple enzymes (e.g. kinases, pyrophosphorylases, phosphatase) described above. However, careful selection of the activated aromatic glycoside donors is required to overcome thermodynamic limitations in order to drive the formation of NDP-sugar. Extensive screening of a range of aromatic glycosides and variants of oleandomycin glycosyltransferase (OleD) from *Streptomyces antibioticus* revealed a variant TDP16 and 2-chloro-4-nitrophenyl- β -D-glucopyranoside to be the best enzyme/donor combination for the reaction to occur in reverse. Furthermore, little to no donor hydrolysis was observed under the conditions. The authors went on to show TDP16 was able to convert 2-chloro-4-nitrophenyl-6-fluoro- β -D-glucopyranoside to either UDP-6FGlc or TDP-6FGlc when reacted in a 1:1 molar ratio with UDP or TDP, respectively (Scheme 22). For both NDPs, the conversion was greater than 90% after 21 h.⁴⁹




Scheme 22 Enzymatic synthesis of UDP- or TDP-6FGlc using the reverse reaction of a glycosyltransferase, TDP16, an engineered variant of OleD.⁴⁹

Further to this, it was shown that TDP16 could be used in a one-pot glycosylation procedure whereby the NDP-donor produced by the reverse reaction could be coupled *in situ* to an aglycone acceptor by the same or an additional glycosyltransferase working in the forward direction. Thus, from 2-chloro-4-nitrophenol-6-fluoro- β -D-glucopyranoside, the 6F-glucosides of both 4-methylumbelliferone (MBU) and vancomycin were detected using either a single enzyme (MBU) or double enzyme (vancomycin) coupled system in the presence of a molar equivalent of UDP. After a reaction time of 24 hours, 70% and 40% conversion from the donor were observed for the MBU and vancomycin glycosides respectively.⁴⁹


Cps2L (*Streptococcus pneumoniae*) is the enzyme that catalyses the synthesis of deoxythymidine diphosphate glucopyranose (dTDP-Glc), from glucose-1-phosphate. When evaluated against a panel of fluorinated glucose phosphates (2FGlc, 3FGlc, 4FGlc, 6FGlc, 4FGal, and 2,2F₂Glc) and 2FMan, it was shown to exhibit broad substrate specificity as it could convert all the substrates almost as effectively as for the natural substrate Glc-1P within 20-30 minutes (Scheme 23).⁵⁰ Kinetic analysis revealed 6FGlc behaved in a very similar manner to Glc, and this is consistent with previous studies that indicated that the 6'-hydroxyl group of Glc-1P made the smallest contribution to interactions with the enzyme⁵¹. Fluorination at the 3' position resulted in the most substantial decrease of k_{cat}/K_m compared to Glc-1P and other monofluorinated substrates, and to a somewhat lower extent, the enzyme did not tolerate well a change in stereochemistry at the 4 position. The 2,2 difluorinated version of Glc-1P maintained binding affinity, but with a decreased turnover number.⁵⁰ The same set of fluorinated sugars were evaluated with the enzyme guanosine diphosphate-mannopyranose pyrophosphorylase (GDP-ManPP) from *Salmonella enterica* (Scheme 23).⁵⁰ When given long enough reaction times, the enzyme was capable of turning over all the sugars assayed. 2FGlc-1P, and 2F2FGlc-1P were converted almost quantitatively within 2.5 h, which is consistent with the timeframe for the natural substrate Man-1P. 3FGlc-1P, 4FGlc-1P, 4FGal-1P and 6FGlc-1P required longer reaction times for similar rates of

conversion. The authors suggested the possible use of the enzymes and products in the synthesis of unnatural glycans, however, further results have not been reported yet.




Scheme 23 Enzymatic synthesis of a series of fluorinated TDP and GDP-sugars using Cps2L and GDP-mannopyranose pyrophosphorylase.⁵⁰

The use of fluorinated glucose as an acceptor of glycosyltransferases is rare. Notable examples are the synthesis of fluorinated trehalose analogues. Trehalose-6-phosphate (T6P) synthase (OtsA) from *E. coli* is a retaining glycosyltransferase that transfers glucose from UDP-glucose to the anomeric centre of glucose-6-phosphate (G6P) to synthesise a non-reducing disaccharide phosphate, trehalose-6-phosphate (Scheme 24). 2-Fluoro-, 3-fluoro- and 4-fluoro-glucose-6-phosphates and 2-fluoro-mannose-6-phosphate were used as the acceptor for kinetic studies, which revealed all the fluorinated acceptors acted as a substrate albeit with lower catalytic efficiencies.⁵² Products were not analysed spectroscopically except for 2-fluoro-trehalose-6-phosphate, which was fully characterised.⁵³ 2-F T6P then undergoes phosphate hydrolysis by T6P specific phosphatase, OtsB to yield 2-fluoro-trehalose (Scheme 24). Later, the Swarts group used a trehalose synthase (TreT) from *Thermoproteus tenax* to synthesise fluorinated trehalose analogues.⁵⁴ This enzyme is a retaining glycosyltransferase that uses UDP-glucose as a donor and glucose as acceptor, and thus able to synthesise trehalose in one step unlike the OtsA approach. The authors demonstrated that TreT could accept 2-fluoro-, 3-fluoro- and 6-fluoro-glucose as acceptor substrates while it did not react with 4-fluoro-glucose (Scheme 24). In contrast to OtsA, TreT showed comparable activities with fluorinated glucose acceptors to that of glucose, and particularly, the highest activity was measured with 2-fluoro-glucose. 2-Fluoro-, 3-fluoro- and 6-fluoro-trehalose analogues were fully characterised.



Scheme 24 Enzymatic synthesis of 2F-trehalose-6-phosphate using trehalose-6-phosphate synthase (OtsA)⁵³ and of fluorinated trehaloses using trehalose synthase (TreT)⁵⁴


In some cases, fluorinated glucose analogues have been used as an acceptor of transglycosidation by glycoside hydrolases (GHs). As mentioned earlier, synthetic activities of glycoside hydrolases are negatively offset by typically higher hydrolytic activity.²³ To address this, Withers et al. introduced an idea of glycosynthase, engineering a glycoside hydrolase to remove a nucleophile residue, and thus making the mutant non-hydrolytic while the intact original structure of the mutant allows the formation of the glycosidic bond.⁵⁵ In an early example of glycosynthase Abg E358A made from a glucosidase from *Agrobacterium* sp., α -glucosyl fluoride was used as a donor and 2,4-dinitrophenyl β -glucoside as an acceptor (Scheme 25).⁵⁵ The reaction yielded a mixture of β -(1 \rightarrow 4)-linked di- (38%), tri- (42%) and tetrasaccharides (4%). There are also examples that used wild type glycoside hydrolases in transglycosidation. α -Glucosyl fluoride and α -galactosyl fluoride were used to monitor transglycosidation kinetics of α -glycoside hydrolases including α -glucosidase from Baker's yeast *Saccharomyces cerevisiae* (Scheme 25) and α -galactosidase from either green coffee beans or *Bacillus stearothermophilus* with ¹⁹F NMR.⁵⁶ These reactions yielded mixtures of anomeric fluorides of mostly α -(1 \rightarrow 2) and α -(1 \rightarrow 6) disaccharides. Field et al. also applied ¹⁹F NMR to monitor the transglycosidation activity of three GH enzymes, glucanotranferases from *E. coli* (*EcMalQ*), *Manihot esculenta* (*MeDPE1*) and *Arabidopsis thaliana* (*AtDPE2*) using 2-fluoro-, 3-fluoro- and 6-fluoro-glucose as an acceptor and glycogen as a donor (Scheme 25).⁵⁷ These GHs are characterised by their high α -(1 \rightarrow 4) transglycosidation activity, thus given their name of glucanotransferase. All three fluorinated glucoses acted as an acceptor, yielding mixtures of maltooligosaccharides with different lengths in the reaction with the respective enzymes, and the conversion rates were 4 – 7% for *EcMalQ*, 12 – 17% for *MeDPE1* and 15 – 28% for *AtDPE2*.



Scheme 25 Use of fluorinated glucose analogues as an acceptor of transglucosidation reaction catalysed by glycoside hydrolases.⁵⁵⁻⁵⁷


5. Fluorinated *N*-acetyl-galactosamine

UDP-4FGalNAc is accessible from chemically synthesised 4FGalNAc-1P via a recombinant human enzyme, UDP-*N*-acetylglucosamine pyrophosphorylase (GlmU) (Scheme 26). Using a crude cell extract, an 18% yield of UDP-4GalNAc was isolated and structurally characterised after 2 h.⁵⁸



Scheme 26 Enzymatic synthesis of UDP-4FGalNAc and UDP-4FGlcNAc using human UDP-*N*-acetylglucosamine pyrophosphorylase (GlmU).⁵⁸

Fluorinated GalNAc residues were also recognised as acceptor substrates for the phosphorylase enzyme *BiGalHexNAcP* (Scheme 27). The disaccharide Gal β 1-3 6FGalNAc, with the fluorinated sugar moiety at the reducing end of the disaccharide can be obtained in milligram scale. The synthetic yield of 88% is in fact slightly higher than that with GalNAc as the acceptor (81%). However the anomeric substituent in the GalNAc acceptor is OH and the increased yield may be a result of more facile purification owing to the azidopropyl group in the fluorinated acceptor rather than increased activity of the enzyme toward the acceptor.³⁸




Scheme 27 Enzymatic synthesis of disaccharide Gal β -(1 \rightarrow 3) 6FGalNAc using a phosphorylase, BiGalHexNAcP.³⁸

6. Fluorinated *N*-acetyl-glucosamine


Like UDP-4FGalNAc, UDP-4FGlcNAc can be synthesised by chemoenzymatic methods. Chemically synthesised 4FGlcNAc-1P can be converted enzymatically to UDP-4FGlcNAc using a recombinant human enzyme, GlmU (Scheme 26).⁵⁸ Yields of UDP-4FGlcNAc are slightly better than the galacto-configured isomer (23% vs. 18%) over shorter times (1 h vs. 2 h), which is surprising given the natural substrate for the enzyme contains Gal rather than Glc stereochemistry.

Further to this, Amano *et al.*⁵⁹ were able to demonstrate the *in vivo* synthesis of UDP-4FGlcNAc in a human cancer cell line (PC-3) from acetylated 4FGlcNAc sugars (the acetylation was required for cellular uptake) via hijacking of the cellular hexosamine biosynthetic pathway. Further studies on cellular metabolic glycans revealed that the 4FGlcNAc moiety was not incorporated in glycans produced by the Golgi-*N*-glycosylation pathways and instead UDP-4FGlcNAc acted as an inhibitor of GlcNAc transferases present in the pathway. 5F-GlcNAc derivatives were initially synthesised to probe whether the introduction of fluorine at the 5-position would render such compounds enzyme inhibitors. Preliminary studies indicated that 5FGlcNAc-1P is a substrate for the enzyme GlmU and could be used for the synthesis of UDP-5FGlcNAc (Scheme 28).⁶⁰ UDP-5FGlcNAc is however an inhibitor of the enzyme UDP-GlcNAc 4 epimerase⁶⁰ and the glycosyltransferases, CLS⁶¹ and CgMshA.⁶²




Scheme 28 Enzymatic synthesis of UDP-5FGlcNAc using GlmU.⁶⁰

Conversely, the introduction of the fluorine at C-5 in acceptor substrates is recognised by the glycosyltransferases CLS and GalT (*bovine milk*), which transfer a galactosyl moiety to the 4-position, albeit at reduced efficiency when compared with the natural substrate GlcNAc for GalT (Scheme 29).⁶⁰⁻⁶¹ Kinetic parameters suggest the low efficiency is due to a lower affinity of the enzyme for the modified acceptor, suggesting a large binding energy from 6'-OH of the acceptor.




Scheme 29 Use of 5FGlcNAc as an acceptor of bovine milk β -1,4-galactosyltransferase.⁶⁰

A different type of fluorination, not directly part of the pyranose ring, but at the *N*-acetyl group, is possible in D-GlcNAc. In an early study, it was demonstrated that chemically synthesized UDP-*N*-trifluoroacetylglucosamine (UDP-GlcNTFA) was tolerated fairly well by two glucosaminyltransferases, ‘core-2’ GlcNAc transferase, involved in the biosynthesis of O-linked glycoproteins, and UDP-D-GlcNAc: α -D-Manp-(1 \rightarrow 6)- β -D-GlcNAc transferase V (GlcNAcT-V), a key biosynthetic enzyme controlling the branching pattern of cell surface complex Asn-linked oligosaccharides (Scheme 30).⁶³ Both enzymes were able to synthesize trisaccharide products with 62% and 56% isolated yield. Later, it was found that UDP-GlcNTFA was an excellent substrate for *Pasteurella multocida* strain P-1059 heparosan synthase 2 (*PmHS2*), a bifunctional enzyme that acts both as α -1,4-*N*-acetylglucosaminyltransferase and β -1,4-glucuronyltransferase, and *Escherichia coli* K5 α 1-4GlcNAcT.⁶⁴⁻⁶⁵



Scheme 30 Use of UDP-GlcNTFA as a donor of N-acetylglucosaminyltransferases, Core-2 GlcNAc transferase and GlcNAcT-V.⁶³

It was also demonstrated that the acceptor UDP-GlcNTFA could be synthesized enzymatically using *Bifidobacterium longum* ATCC55813 N-acetylhexosamine 1-kinase (NahK) and *Pasteurella multocida* strain P-1059 (ATCC15742) N-acetylglucosamine-1-phosphate uridylyltransferase (*PmGlmU*).^{64, 66} Along with these two enzymes, *PmHS2* was used in a one-pot multi-enzyme system for the synthesis of heparan sulfate oligosaccharides (Scheme 31). The trifluoroacetyl moiety was then hydrolysed in a mild basic condition, and the resulting free amine moiety sulfated either enzymatically or chemically, yielding natural heparan sulfate oligosaccharides.⁶⁷




Scheme 31 One-pot multi-enzyme system for the synthesis of heparin sulfate oligosaccharides containing NahK (*N*-acetylhexosamine 1-kinase), *PmGlmU* (*Pasteurella multocida* strain P-1059 (ATCC15742) *N*-acetylglucosamine-1-phosphate uridylyltransferase), inorganic pyrophosphatase, and *PmHS2* (heparosan synthase 2, bifunctional glycosyltransferase).⁶⁴

7. Fluorinated Talose

In addition to tolerating all the four isomers of FGal-1P, GalPUT is also able to process sugars which contain the talose stereochemistry (axial at C-2). Errey and co-workers demonstrated that the immobilised system described in section 3.1 is able to synthesise UDP-2FTal from 2FTal in 71% yield (Scheme 32).⁶⁸ This is greater


than for unmodified Tal (15%) and although yields are comparable to Gal and derivatives, longer reaction times (24 h vs. 8 h) are needed.



Scheme 32 Enzymatic synthesis of UDP-2F-talose using the GalK/GalPUT system.⁶⁸

8. Fluorinated Fucose


2-Deoxy-2-fluorinated analogues of sialyl Lewis a can be synthesised in milligram quantities from the donor GDP-2F-L-Fuc using the fucosyltransferase (FucT) III enzyme (Scheme 33).⁶⁹ Extended reaction times (up to one day) are needed to achieve a comparable yield (88%) to the natural donor GDP-L-Fuc (97%). However, the stereo- and regio-selectivity of the enzyme is not altered by the fluorinated donor. This result is remarkable since fluorination at the position adjacent to the anomeric centre normally creates a highly unfavourable environment for the glycosyl transfer reaction that involves an oxocarbenium ion-like transition state.⁷⁰ Usually, GDP-2F-L-Fuc has been observed to act as a competitive inhibitor of a number of human fucosyltransferases.⁸ ⁷¹ In line with this, Fuc-T VI does not accept GDP-2FFuc to synthesise sialyl Lewis x analogues⁶⁹ although the authors did not evaluate the fluorinated analogue as an inhibitor.



Scheme 33 Enzymatic synthesis of sialyl Lewis a containing 2-fluoro-L-fucose.⁶⁹


GDP-6F-L-Fuc has also been shown to be a competitive inhibitor toward the human fucosyltransferases III, V, VI and VII, however the 6F-L-Fuc moiety can be introduced into Lewis x analogues from 6F-L-Fuc making use of the bifunctional enzyme L-fucokinase/GDP-fucose pyrophosphorylase (FKP) and the bacterial α -(1 \rightarrow 3) fucosyltransferase (Scheme 34).⁷² Although FKP has a lower affinity for 6F-L-Fuc than L-Fuc, milligram quantities of GDP-6F-L-Fuc can be prepared in 94% yield from 6FFuc in 5-6 hours. When FKP is combined in

a one-pot, two-enzyme reaction with the fucosyltransferase the sialyl Lewis x derivative can be synthesised directly from 6F-L-Fuc, without the need to isolate GDP-6F-L-Fuc, in 90% yield. In both cases, the regio- and stereoselectivity of the enzymes was confirmed spectroscopically.



Scheme 34 Enzymatic synthesis of Lewis x derivative containing 6-fluoro-L-fucose.⁷²

When combined with the α-(1→2) fucosyltransferase FutC, FKP can also append a 6F-L-Fuc moiety to the non-reducing end of a Gb5 oligosaccharide to yield a derivative of Globo H (Scheme 35). Although FKP can be used to synthesise GDP-gem-6FF-Fuc from gem-6FF-Fuc, the former is not a good substrate for FutC, with only trace amounts of product formed.⁷³




Scheme 35 One-pot multi-enzyme system for the synthesis of fluorinated derivative of Globo-H.⁷³ FKP = L-fucokinase/GDP-fucose pyrophosphorylase, PPA – inorganic pyrophosphatase, Fut C = α -1,2-fucosyltransferase.

9. Fluorinated Sialic Acids

Some fluorinated sialic acid donors have been shown to inhibit SiaTs or TcTS, whilst others have been shown to act as substrates for sialic acid processing enzymes, many of which are discussed in the following section.

9.1 3-Fluorinated sialic acids

Watts and Withers demonstrated that CMP-3F^{ax}Neu5Ac could be synthesised in 91% yield from 3F^{ax}Neu5Ac and CTP using CMP-sialic acid synthase (CSS). To effect this transformation, the authors used a sialic acid aldolase (Scheme 36).⁷⁴ In this synthesis sialic acid aldolase catalyses the formation of Neu5Ac from ManNAc and pyruvate.⁷⁵ By using fluoropyruvate in place of pyruvate, 3F^{ax}Neu5Ac was successfully synthesised. When the two enzyme reactions were coupled together, CMP-3F^{ax}Neu5Ac was isolated in 84% yield over 2 steps from ManNAc. Interestingly, under these conditions only the isomer bearing the 3-fluorine in the axial position was obtained.⁷⁴




Scheme 36 The two step enzymatic synthesis of CMP-3F^{ax}Neu5Ac.⁷⁴

Chen *et al.*⁷⁶ later demonstrated that when the condensation was carried out with a recombinant aldolase from *E. coli* a mixture of both 3F axial and equatorial isomers were formed. These isomers could be which readily separated by flash chromatography and isolated in in 52 and 44% yield, respectively. Axial and equatorial isomers of 3-fluoro-*N*-glycolylneuraminic acid (3FNeu5Gc) and 3-fluoro-keto-deoxynonosonic acid (3FKDN)

were also formed from the aldolase promoted condensation of 3-fluoropyruvate and N-glycolylmannosamine (ManNGc) or Man respectively.


Despite early claims that neither 3F^{eq}Neu5Ac nor 3F^{ax}Neu5Ac are substrates for CSS from calf brain^{77,78}, Chen and colleagues went on to demonstrate the both 3F^{ax}Neu5Ac and 3F^{eq}Neu5Ac were substrates for CSS from *Neisseria meningitides* (NmCSS), and the CMP-sialic acids could be synthesised in excellent yields (Scheme 37)⁷⁶. Furthermore, Heise *et al.* reported that a C-5 carbamate increases the efficiency of the conversion of 3F sialic acids to the relevant CMP analogue.⁷⁹



Scheme 37 The enzymatic synthesis of CMP activated 3F sialic acid analogues.⁷⁶

The presence of the electronegative fluorine atom adjacent to the anomeric carbon has meant that CMP-3F^{ax}NeuAc has been utilised as an inhibitor of a commercial α -(2 \rightarrow 6) SiaT,⁷⁸ the SiaT Cst-II from *Campylobacter jejuni*⁸⁰ and TcTs.⁸¹ Nevertheless, Chen *et al.* demonstrated that CMP-3F^{ax}Neu5Ac and CMP-3F^{ax}Neu5Gc are substrates for the α -(2 \rightarrow 3) SiaT from *Pasteurella multocida* (PmST1). Fluorinated sialyl lactosides can be prepared in a one-pot, two-enzyme (NmCSS and PmST1) system from the corresponding 3F^{ax} sialic acid (Scheme 38). Under the same conditions, the sialyl lactosides containing the 3F^{eq} sialic acid could be obtained in higher yields. The α -(2 \rightarrow 6) SiaT from *Photobacterium damsela* (Pd2,6ST) does not appear to


tolerate the fluorinated donors as well as *PmST1* can, and corresponding yields of the α 2-6 fluorosialyl lactosides were much lower.⁷⁶



Scheme 38 The one-pot, two enzyme synthesis of 3F sialic acid-containing trisaccharides.⁷⁶

9.2 5-Fluorinated sialic acids


Wong *et al.*⁸² reported the synthesis of 5-fluorinated 5-epi KDN in 30% yield from 2-deoxy-2-fluoroglucose (2FGlc) and pyruvate using a sialic acid aldolase (Scheme 39). However, microbial CSS did not recognise 5F, 5-epi KDN as a substrate, possibly due to either the lack of 5-NHAc, or the differing stereochemistry (axial) at C-5 compared with Neu5Ac or KDN (equatorial).



Scheme 39 The enzymatic synthesis of 5, 5-epi FKDN from 2FGlc.⁸²

More recently, Chen *et al.*⁸³ reported the enzymatic synthesis of 5FKDN from 2-deoxy-2-fluoromannose (2FMan) using a sialic acid aldolase from *Pasteurella multocida* (*Pm* sialic acid aldolase). The authors utilised this transformation in a one-pot, three-enzyme system which also contained *NmCSS* and either *PmST1* or *Pd2,6ST* with a *para*-nitrophenyl galactoside (Gal β pNP) acceptor to give either the α 2-3 or α 2-6 fluorinated


disaccharide in yields of 74% and 79% respectively (Scheme 40). Despite these good yields, they were not as high as for the analogous reactions containing Neu5Gc derivatives.



Scheme 40 The one-pot, three-enzyme synthesis of 5FKDN-containing disaccharides.⁸³

9.3 7-Deoxy-7-fluoro sialic acids


The enzymatic synthesis of CMP-7FNeu5Ac from the corresponding 4FManNAc using sialic acid aldolase and CSS was initially explored by Dräger *et al.*⁸⁴ Using conditions which had resulted in full conversion of ManNAc to Neu5Ac, only 56% conversion to 7FNeu5Ac was obtained from 4FManNAc using a commercially available aldolase enzyme (Scheme 41). In order to shift the equilibrium of the aldolase catalysed reaction toward the sialic acid, coupling with the CSS from *Neisseria meningitidis* serogroup B (*NmBCSS*) was explored. This approach, however, did not significantly improve conversion to CMP-7FNeu5Ac as determined by LC-ESI-MS and both 4FManNAc and 7FNeu5Ac remained. Full conversion to CMP-7FNeu5Ac was only achieved when isolated 7FNeu5Ac was turned over by *NmBCSS* in the presence of pyrophosphatase. It is worthy to note that the authors investigated other sources of CSS (murine CSS or rainbow trout CMP-Kdn-synthetase), however best results were obtained with *NmBCSS*.



Scheme 41 The enzymatic synthesis of CMP-7FNeu5Ac.⁸⁴

Preparative scale synthesis of sialosides containing the 7FNeu5Ac from hexoses via a one-pot, three-enzyme system was later demonstrated by Khedri *et al.*⁸⁵ Using GalβpNP as an acceptor and employing *Pm* sialic acid aldolase, *NmCSS* and either *PmST1* or *Pd2,6ST*, milligram scale synthesis of the galactosyl α -(2→3) or α -


(2→6) sialosides could be isolated and fully characterised in 98 and 92% yields respectively (Scheme 42). Using the same enzymes, the α -(2→3) and α -(2→6) 7FKDN disaccharides could also be synthesised from 4FMan in 98 and 95% yields respectively. In both cases, isolation of the sialic acid or CMP-sialic acid was not required, and the stereo- and regioselectivity of the SiaTs was not altered with the fluorinated sugar.



Scheme 42 The one-pot, three enzyme system used to synthesise disaccharides containing 7F sialic acid moieties.⁸⁵

9.4 8-Deoxy-8-fluorosialic acids

CMP-8FNeu5Ac can be isolated in 82% yield from 8FNeu5Ac after 6 hours reaction with *NmCSS* and an inorganic pyrophosphatase (Scheme 43). Kinetic studies revealed that the α -(2→3) SiaT from *Campylobacter jejuni* (Cst-I) is able to transfer CMP-8FNeu5Ac to modified lactose acceptors (60 minutes), albeit with less efficiency than the natural donor CMP-Neu5Ac (2 minutes). Nonetheless, a sialyl lactose derivative could still be isolated in milligram quantities and 80% yield using this enzyme. Spectroscopic analysis confirmed the regio- and stereoselectivity of the enzyme was retained with the fluorinated donor. Under conditions optimal for α -(2→3)-sialyltransfer activity, the multifunctional enzyme *Pm0188h* (*Pasteurella multocida*) is also able to transfer CMP-8FNeu5Ac to a modified lactose acceptor, but unlike Cst-I the reaction does not proceed to completion, possibly due to hydrolysis of the sialylated product.⁸⁶ Indeed, the authors acknowledge *Pm0188h* has been shown to exhibit α 2,3-sialidase activity as well as α 2,3-trans-sialidase and α 2,6-sialyltransferase behaviour.⁸⁶⁻⁸⁷




Scheme 43 The enzymatic synthesis of an 8F sialyl lactose derivative.⁸⁶

9.5 9-Deoxy-9-fluorosialic acids

Early analytical studies suggested that chemically synthesised 9F-Neu5Ac was a substrate for CSS from calf brains, albeit kinetics showed the enzyme displayed a lower affinity toward the fluorinated substrate compared for the natural substrate Neu5Ac.^{77, 88}


9F-Neu5Ac has been synthesised enzymatically from 6FManNAc, and ¹⁴C-radio-labelling was introduced at the 2-position by Conrard *et al.* via the coupling of 6FManNAc and [2-¹⁴C]pyruvic acid using the enzyme acylneuraminate pyruvate-lyase from *Clostridium perfringens* with an overall yield of 60% (Scheme 44).

Although kinetic analysis revealed a higher *K_m* for 9F[2-¹⁴C]Neu5Ac than that of the natural substrate Neu5Ac, activation of the fluorinated substrate with CSS from rat liver gave CMP-9F [2-¹⁴C]Neu5Ac in 30% yield after purification. The authors went on to demonstrate CMP-9F [2-¹⁴C]Neu5Ac is a substrate for an α-(2→6) SiaT from rat liver, and can be transferred onto a galactosyl terminated asialo-α1-acid glycoprotein acceptor, albeit at a lower rate than CMP-Neu5Ac even at higher sugar nucleotide concentrations.⁸⁹




Scheme 44 The enzymatic synthesis of CMP-9F [2-¹⁴C]Neu5Ac, orange dot denotes the position of the isotopic label.⁸⁹

Liu *et al.*⁸² have also demonstrated the synthesis of 9-deoxy-9-fluorinated sialic acid derivatives of Neu5Ac and KDN from modified hexopyranose derivatives and pyruvate using sialic acid aldolase (Scheme 45). Kinetic studies revealed that CSS enzymes from *E. coli* (both a tagged and native enzyme) were able to turnover 9FNeu5Ac with parameters similar to those for the natural substrate Neu5Ac. Conversely, the 7F, 9F di-fluorinated 7-epi KDN derivative was not recognised by the enzyme, however the authors postulate this observation was due to the lack of a 5-NHAc group rather than the presence of the fluorine.




Scheme 45 The enzymatic synthesis of 9F derivatives of Neu5Ac and 7-epi KDN.⁸²

Using the same one-pot, three-enzyme system described for the synthesis of 7-fluorinated α -(2 \rightarrow 3) or α -(2 \rightarrow 6) sialosides, Khedri *et al.* extended the substrate scope to include 6FManNAc, 6FManNGc and 6FMan derivatives (Scheme 46). Yields of the corresponding sialosides were >95% in all cases, which is comparable or better than that observed with the natural substrates.⁹⁰



Scheme 46 The one-pot, three enzyme synthesis of sialosides containing 9F modified sialic acid derivatives.⁹⁰

Kajihara *et al.* presented a strategy to introduce ^{13}C labelling at the C-3 of modified Neu5Ac residues, taking advantage of the established synthetic routes used to obtain fluorinated Neu5Ac derivatives from Neu5Ac using selective protection and deprotection. This avoided the synthesis of the fluorinated ManNAc analogues, of which the authors argue the synthetic chemistry of which has been less explored. Thus, two aldolase reactions were performed, the first, coupled with pyruvate removal by lactate dehydrogenase, promoted the degradation of the fluorinated Neu5Ac derivative to the hexose (Scheme 47). After conversion to the hexose and switching off of the cofactor regeneration cycle, the second aldolase reaction with labelled pyruvate resulted in the labelled Neu5Ac analogue. Using this two-step, one-pot reaction, 9F[^{13}C]Neu5Ac was synthesised in milligram scales (62% yield) with 96% ^{13}C labelling incorporated.⁹¹ Subsequent chemical conversion of 9F [3- ^{13}C]Neu5Ac into CMP-9F [3- ^{13}C]Neu5Ac allowed for α -(2 \rightarrow 6) sialyl transfer (bacterial) onto a glycosylated protein.⁹²



Scheme 47 The enzymatic synthesis of 9F[3- ^{13}C]Neu5Ac from 9FNeu5Ac, orange dot denotes the position of the isotopic label.⁹¹

Conclusion

Glycosylation of fluorinated sugars using enzymatic methods has been achieved for a wide range of monofluorinated sugars and has been employed in the synthesis of many fluorinated oligosaccharides. Monofluorinated monosaccharides including galactose, glucose and sialic acid have been successfully utilised as donors and acceptors for glycosylation, for the production of a diverse range of fluorinated disaccharide and trisaccharide targets, and phosphorylated di- and tetrafluorinated galactose donors have been enzymatically synthesised, although not yet successfully used in enzymatic glycosylations. The incorporation of more heavily fluorinated monosaccharides into glycans through glycosylation remains challenging by enzymatic means, however enzymatic glycosylations to date have relied mainly on the use of natural enzymes, and the future development of engineered enzymes able to accept a wider range of unnatural substrates could provide new routes to glycans with more complicated fluorination patterns. Enzyme reaction rates using fluorinated sugars are often low in comparison to the natural substrates, but it has been demonstrated that where an unnatural sugar is able to be accepted by the enzyme, even sugars initially considered competitive inhibitors of the enzyme can be enzymatically transformed with careful optimisation of reaction conditions. One limitation of enzymatic synthesis is the low commercial availability of glyco-enzymes, with enzyme supply often reliant on transfers of enzymes, plasmids or expressed cells between researchers and in-house enzyme production, which restricts the uptake of these methods by chemists, who may choose a more familiar chemical glycosylation route over an enzymatic route that requires production of the specific enzyme for the transformation. The next generation of fluorinated glycans are likely to require a combination of chemical and enzymatic transformations to achieve more diverse fluorination patterns.

Conflicts of Interests

There are no conflicts to declare.

Acknowledgements

This work has been funded by the Industrial Biotechnology Catalyst (Innovate UK, BBSRC, EPSRC) to support the translation, development and commercialisation of innovative Industrial Biotechnology processes (BB/M028941/1 and BB/M028879/1).

References

1. Tyrikos-Ergas, T.; Fittolani, G.; Seeberger, P. H.; Delbianco, M., Structural Studies Using Unnatural Oligosaccharides: Toward Sugar Foldamers. *Biomacromolecules* **2019**, *21*, 18-29.
2. Ardá, A.; Jiménez-Barbero, J., The recognition of glycans by protein receptors. Insights from NMR spectroscopy. *Chemical Communications* **2018**, *54* (38), 4761-4769.
3. Bilska-Markowska, M.; Szwajca, A.; Marciniak, B., Design, properties and applications of fluorinated and fluoroalkylated N-containing monosaccharides and their analogues. *Journal of Fluorine Chemistry* **2019**, *227*, 109364.
4. Cavaliere, A.; Probst, K. C.; Westwell, A. D.; Slusarczyk, M., Fluorinated nucleosides as an important class of anticancer and antiviral agents. *Future Medicinal Chemistry* **2017**, *9* (15), 1809-1833.
5. van der Born, D.; Pees, A.; Poot, A. J.; Orru, R. V. A.; Windhorst, A. D.; Vugts, D. J., Fluorine-18 labelled building blocks for PET tracer synthesis. *Chemical Society Reviews* **2017**, *46* (15), 4709-4773.
6. Tysoe, C.; Withers, S. G., Fluorinated Mechanism-Based Inhibitors: Common Themes and Recent Developments. *Current Topics in Medicinal Chemistry* **2014**, *14* (7), 865-874.
7. Street, I. P.; Armstrong, C. R.; Withers, S. G., Hydrogen bonding and specificity. Fluorodeoxy sugars as probes of hydrogen bonding in the glycogen phosphorylase-glucose complex. *Biochemistry* **1986**, *25* (20), 6021-6027.
8. Burkart, M. D.; Vincent, S. P.; Düffels, A.; Murray, B. W.; Ley, S. V.; Wong, C.-H., Chemo-Enzymatic Synthesis of Fluorinated Sugar Nucleotide: Useful Mechanistic Probes for Glycosyltransferases. *Bioorganic & Medicinal Chemistry* **2000**, *8*, 1937-1946.
9. Matei, E.; André, S.; Glinschert, A.; Infantino, A. S.; Oscarson, S.; Gabius, H.-J.; Gronenborn, A. M., Fluorinated Carbohydrates as Lectin Ligands: Dissecting Glycan–Cyanovirin Interactions by Using ¹⁹F NMR Spectroscopy. *Chemistry – A European Journal* **2013**, *19* (17), 5364-5374.
10. Rajendran, J. G.; Wilson, D. C.; Conrad, E. U.; Peterson, L. M.; Bruckner, J. D.; Rasey, J. S.; Chin, L. K.; Hofstrand, P. D.; Grierson, J. R.; Eary, J. F.; Krohn, K. A., [¹⁸F]FMISO and [¹⁸F]FDG PET imaging in soft tissue sarcomas: correlation of hypoxia, metabolism and VEGF expression. *European Journal of Nuclear Medicine and Molecular Imaging* **2003**, *30* (5), 695-704.
11. Prante, O.; Hamacher, K.; Coenen, H. H., Chemoenzymatic n.c.a synthesis of the coenzyme UDP-2-deoxy-2-(¹⁸F)fluoro-α-D-glucopyranose as substrate of glycosyltransferases. *Journal of Labelled Compounds and Radiopharmaceuticals* **2007**, *50*, 55–63.
12. Williams, S. J.; Withers, S. G., Glycosyl fluorides in enzymatic reactions. *Carbohydrate Research* **2000**, *327* (1), 27-46.
13. Johannes, M.; Reindl, M.; Gerlitzki, B.; Schmitt, E.; Hoffmann-Röder, A., Synthesis and biological evaluation of a novel MUC1 glycopeptide conjugate vaccine candidate comprising a 4'-deoxy-4'-fluoro-Thomsen–Friedenreich epitope. *Beilstein Journal of Organic Chemistry* **2015**, *11*, 155-161.
14. Li, W.; McArthur, J. B.; Chen, X., Strategies for chemoenzymatic synthesis of carbohydrates. *Carbohydrate Research* **2019**, *472*, 86-97.
15. Uhrig, M. L.; Lantaño, B.; Postigo, A., Synthetic strategies for fluorination of carbohydrates. *Organic & Biomolecular Chemistry* **2019**, *17* (21), 5173-5189.

16. Cheng, Y.; Guo, A.-L.; Guo, D.-S., Recent Progress in Synthesis and Applications of Fluorinated Carbohydrates. *Current Organic Chemistry* **2010**, *14* (9), 977-999.
17. Withers, S. G.; Street, I. P.; Rettig, S. J., The preferred conformation of 2-fluoro-2-deoxy β -D-mannopyranosyl fluoride. An X-ray crystallographic and 2-dimensional proton nuclear magnetic resonance study. *Canadian Journal of Chemistry* **1986**, *64* (2), 232-236.
18. Linclau, B.; Golten, S.; Light, M.; Sebban, M.; Oulyadi, H., The conformation of tetrafluorinated methyl galactoside anomers: crystallographic and NMR studies. *Carbohydrate Research* **2011**, *346* (9), 1129-1139.
19. Graton, J.; Wang, Z.; Brossard, A.-M.; Gonçalves Monteiro, D.; Le Questel, J.-Y.; Linclau, B., An unexpected and significantly lower hydrogen-bond-donating capacity of fluorohydrins compared to nonfluorinated alcohols. *Angew Chem Int Ed Engl* **2012**, *51* (25), 6176-6180.
20. Graton, J.; Compain, G.; Besseau, F.; Bogdan, E.; Watts, J. M.; Mtashobya, L.; Wang, Z.; Weymouth-Wilson, A.; Galland, N.; Le Questel, J.-Y.; Linclau, B., Influence of Alcohol β -Fluorination on Hydrogen-Bond Acidity of Conformationally Flexible Substrates. *Chemistry – A European Journal* **2017**, *23* (12), 2811-2819.
21. Hayashi, T.; Murray, B. W.; Wang, R.; Wong, C.-H., A Chemoenzymatic Synthesis of UDP-(2-deoxy-2-fluoro)-galactose and Evaluation of its Interaction with Galactosyltransferase. *Bioorganic & Medicinal Chemistry* **1997**, *5*, 497-500.
22. Errey, J. C.; Mukhopadhyay, B.; Kartha, K. P. R.; Field, R. A., Flexible enzymatic and chemo-enzymatic approaches to a broad range of uridine-diphospho-sugars. *Chemical Communications* **2004**, (23), 2706-2707.
23. Danby, P. M.; Withers, S. G., Advances in Enzymatic Glycoside Synthesis. *ACS Chemical Biology* **2016**, *11* (7), 1784-1794.
24. Frasch, A. C. C., Functional Diversity in the Trans-sialidase and Mucin Families in *Trypanosoma cruzi*. *Parasitology Today* **2000**, *16* (7), 282-286.
25. Kent, P. W.; Wright, J. R., Fluorocarbohydrates. Part XXIV. Chemical and Enzymatic Reduction and Phosphorylation of 6-deoxy-6-fluoro- α -D-galactose. *Carbohydrate Research* **1972**, *22* (1), 193-200.
26. Thomas, P.; Bessell, E. M.; Westwood, J. H., The use of deoxyfluoro-D-galactopyranoses in a study of yeast galactokinase specificity. *Biochemical Journal* **1974**, *139* (3), 661-664.
27. Kean, E., CMP-sialic acid synthetase of the nucleus. *Biochimica et Biophysica Acta (BBA) - General Subjects* **2004**, *1673* (1-2), 56-65.
28. Barlow, J. N.; Blanchard, J. S., Enzymatic synthesis of UDP-(3-deoxy-3-fluoro)-D-galactose and UDP-(2-deoxy-2-fluoro)-D-galactose and substrate activity with UDP-galactopyranose mutase. *Carbohydrate Research* **2000**, *328* (4), 473-480.
29. Bessell, E. M.; Foster, A. B.; Westwood, J. H., The use of deoxyfluoro-D-glucopyranoses and related compounds in a study of yeast hexokinase specificity. *Biochemical Journal* **1972**, *128* (2), 199-204.
30. Nishimura, S.-I.; Nagahori, N.; Hamamoto, T.; Okuyama, K.; Noguchi, T. Sugar chain containing 4-position halogenated galactose and application thereof. 2012.
31. Yang, J.; Liu, L.; Thorson, J. S., Structure-Based Enhancement of the First Anomeric Glucokinase. *ChemBioChem* **2004**, *5* (7), 992-996.
32. Timmons, S. C.; Hui, J. P. M.; Pearson, J. L.; Peltier, P.; Daniellou, R.; Nugier-Chauvin, C.; Soo, E. C.; Syvitski, R. T.; Ferrières, V.; Jakeman, D. L., Enzyme-Catalyzed Synthesis of Furanosyl Nucleotides. *Organic Letters* **2008**, *10* (2), 161-163.

33. Jakeman, D. L.; Young, J. L.; Huestis, M. P.; Peltier, P.; Daniellou, R.; Nugier-Chauvin, C.; Ferrières, V., Engineering Ribonucleoside Triphosphate Specificity in a Thymidyltransferase. *Biochemistry* **2008**, *47* (33), 8719-8725.
34. Loch, N.; Geilen, C. C.; Spöndle, I.; Oberdorfer, F.; Keppler, D.; Tauber, R.; Reutter, W., 2-Deoxy-2-fluoro-D-galactose protein N-glycosylation. *FEBS Letters* **1991**, *294* (3), 217-220.
35. Kodama, H.; Kajihara, Y.; Endo, T.; Hashimoto, H., Synthesis of UDP-6-deoxy- and -6-fluoro-D-galactoses and their enzymatic glycosyl transfer to mono- and biantennary carbohydrate chains. *Tetrahedron Letters* **1993**, *34* (40), 6419-6422.
36. Schengrund, C.-L.; Kováč, P., UDP-6-deoxy-6-fluoro- α -D-galactose binds to two different galactosyltransferases, but neither can effectively catalyze transfer of the modified galactose to the appropriate acceptor. *Carbohydrate Research* **1999**, *319* (1-4), 24-28.
37. Yu, H.; Thon, V.; Lau, K.; Cai, L.; Chen, Y.; Mu, S.; Li, Y.; Wang, P. G.; Chen, X., Highly efficient chemoenzymatic synthesis of β 1-3-linked galactosides. *Chemical Communications* **2010**, *46* (40), 7507-7509.
38. Yan, J.; Chen, X.; Wang, F.; Cao, H., Chemoenzymatic synthesis of mono- and di-fluorinated Thomsen-Friedenreich (T) antigens and their sialylated derivatives. *Organic & Biomolecular Chemistry* **2013**, *11* (5), 842-848.
39. Li, L.; Liu, Y.; Li, T.; Wang, W.; Yu, Z.; Ma, C.; Qu, J.; Zhao, W.; Chen, X.; Wang, P. G., Efficient chemoenzymatic synthesis of novel galacto-N-biose derivatives and their sialylated forms. *Chemical Communications* **2015**, *51* (51), 10310-10313.
40. Euzen, R.; Lopez, G.; Nugier-Chauvin, C.; Ferrières, V.; Plusquellec, D.; Rémond, C.; O'Donohue, M., A Chemoenzymatic Approach for the Synthesis of Unnatural Disaccharides Containing D-Galacto- or D-Fucofuranosides. *European Journal of Organic Chemistry* **2005**, *2005* (22), 4860-4869.
41. Chlubnová, I.; Králová, B.; Dvořáková, H.; Spiwok, V.; Filipp, D.; Nugier-Chauvin, C.; Daniellou, R.; Ferrières, V., Biocatalyzed synthesis of difuranosides and their ability to trigger production of TNF- α . *Bioorganic & Medicinal Chemistry Letters* **2016**, *26* (6), 1550-1553.
42. Watkins, W. M., Chapter 5 Biosynthesis 5. Molecular Basis of Antigenic Specificity in the ABO, H and Lewis Blood-Group Systems. In *New Comprehensive Biochemistry*, Montreuil, J.; Vliegthart, J. F. G.; Schachter, H., Eds. Elsevier: 1995; Vol. 29, pp 313-390.
43. Lowary, T. L.; Hindsgaul, O., Recognition of synthetic deoxy and deoxyfluoro analogs of the acceptor α -l-Fucp-(1 \rightarrow 2)- β -D-Galp-OR by the blood-group A and B gene-specified glycosyltransferases. *Carbohydrate Research* **1993**, *249* (1), 163-195.
44. Kasuya, M. C.; Ito, A.; Hatanaka, K., Simple and convenient synthesis of a fluorinated GM4 analogue. *Journal of Fluorine Chemistry* **2007**, *128* (5), 562-565.
45. Allman, S. A.; Jensen, H. H.; Vijayakrishnan, B.; Garnett, J. A.; Leon, E.; Liu, Y.; Anthony, D. C.; Sibson, N. R.; Feizi, T.; Matthews, S.; Davis, B. G., Potent Fluoro-oligosaccharide Probes of Adhesion in Toxoplasmosis. *ChemBioChem* **2009**, *10* (15), 2522-2529.
46. Van Dorst, J. A. L. M.; Tikkanen, J. M.; Krezdorn, C. H.; Streiff, M. B.; Berger, E. C.; Van Kuik, J. A.; Kamerling, J. P.; Vliegthart, J. F. G., Exploring the substrate specificities of the α -2,6- and α -2,3-sialyltransferases using synthetic acceptor analogues. *European Journal of Biochemistry* **1996**, *242*, 674-681.
47. Unverzagt, C.; Kunz, H.; Paulson, J. C., High-efficiency synthesis of sialyloligosaccharides and sialoglycopeptides. *Journal of the American Chemical Society* **1990**, *112* (25), 9308-9309.

48. Caputi, L.; Rejzek, M.; Louveau, T.; O'Neill, E. C.; Hill, L.; Osbourn, A.; Field, R. A., A one-pot enzymatic approach to the O-fluoroglucoside of N-methylantranilate. *Bioorganic & Medicinal Chemistry* **2013**, *21*, 4762–4767.
49. Gantt, R. W.; Peltier-Pain, P.; Cournoyer, W. J.; Thorson, J. S., Using simple donors to drive the equilibria of glycosyltransferase-catalyzed reactions. *Nature Chemical Biology* **2011**, *7*, 685–691.
50. Zhu, J.-S.; McCormick, N. E.; Timmons, S. C.; Jakeman, D. L., Synthesis of α -Deoxymono and Difluorohexopyranosyl 1-Phosphates and Kinetic Evaluation with Thymidyl- and Guanidyltransferases. *The Journal of Organic Chemistry* **2016**, *81*, 8816–8825.
51. Jiang, J.; Biggins, J. B.; Thorson, J. S., A General Enzymatic Method for the Synthesis of Natural and “Unnatural” UDP- and TDP-Nucleotide Sugars. *Journal of the American Chemical Society* **2000**, *122* (28), 6803–6804.
52. Lee, S. S.; Hong, S. Y.; Errey, J. C.; Izumi, A.; Davies, G. J.; Davis, B. G., Mechanistic evidence for a front-side, S_Ni-type reaction in a retaining glycosyltransferase. *Nat Chem Biol* **2011**, *7* (9), 631–638.
53. Backus, K. M.; Boshoff, H. I.; Barry, C. S.; Boutureira, O.; Patel, M. K.; D'Hooge, F.; Lee, S. S.; Via, L. E.; Tahlan, K.; Barry, C. E.; Davis, B. G., Uptake of unnatural trehalose analogs as a reporter for Mycobacterium tuberculosis. *Nat Chem Biol* **2011**, *7* (4), 228–235.
54. Rundell, S. R.; Wagar, Z. L.; Meints, L. M.; Olson, C. D.; O'Neill, M. K.; Piligian, B. F.; Poston, A. W.; Hood, R. J.; Woodruff, P. J.; Swarts, B. M., Deoxyfluoro-d-trehalose (FDTre) analogues as potential PET probes for imaging mycobacterial infection. *Organic & Biomolecular Chemistry* **2016**, *14* (36), 8598–8609.
55. Mackenzie, L. F.; Wang, Q.; Warren, R. A. J.; Withers, S. G., Glycosynthases: Mutant Glycosidases for Oligosaccharide Synthesis. *Journal of the American Chemical Society* **1998**, *120* (22), 5583–5584.
56. André, C.; Spangenberg, P.; Gentil, E.; Rabiller, C., In situ ¹⁹F NMR spectroscopy study of enzymatic transglycosylation reactions using α -d-aldohexopyranosyl fluorides as donors and acceptors. *Tetrahedron: Asymmetry* **2001**, *12* (5), 779–783.
57. Tantanarat, K.; Rejzek, M.; O'Neill, E.; Ruzanski, C.; Hill, L.; Fairhurst, S. A.; Limpaseni, T.; Field, R. A., An expedient enzymatic route to isomeric 2-, 3- and 6-monodeoxy-monofluoro-maltose derivatives. *Carbohydrate Research* **2012**, *358*, 12–18.
58. Feng, F.; Okuyama, K.; Niikura, K.; Ohta, T.; Sadamoto, R.; Monde, K.; Noguchi, T.; Nishimura, S.-I., Chemo-enzymatic synthesis of fluorinated 2-N-acetamidoglycosyl nucleotides using UDP-GlcNAc pyrophosphorylase. *Organic & Biomolecular Chemistry* **2004**, *2* (11), 1617–1623.
59. Nishimura, S.-I.; Hato, M.; Hyugaji, S.; Feng, F.; Amano, M., Glycomics for Drug Discovery: Metabolic Perturbation in Androgen-Independent Prostate Cancer Cells Induced by Unnatural Hexosamine Mimics. *Angewandte Chemie International Edition* **2012**, *51*, 3386–3390.
60. Hartman, M. C. T.; Coward, J. K., Synthesis of 5-Fluoro N-Acetylglucosamine Glycosides and Pyrophosphates via Epoxide Fluoridolysis: Versatile Reagents for the Study of Glycoconjugate Biochemistry. *Journal of the American Chemical Society* **2002**, *124*, 10036–10053.
61. Hartman, M. C. T.; Jiang, S.; Rush, J. S.; Waechter, C. J.; Coward, J. K., Glycosyltransferase Mechanisms: Impact of a 5-Fluoro Substituent in Acceptor and Donor Substrates on Catalysis. *Biochemistry* **2007**, *46*, 11630–11638.

62. Frantom, P. A.; Coward, J. K.; Blanchard, J. S., UDP-(5F)-GlcNAc acts as a slow binding inhibitor of MShA, a retaining glycosyltransferase. *Journal of the American Chemical Society* **2010**, *132*, 6626–6627.
63. Sala, R. F.; MacKinnon, S. L.; Palcic, M. M.; Tanner, M. E., UDP-N-trifluoroacetylglucosamine as an alternative substrate in N-acetylglucosaminyltransferase reactions. *Carbohydrate Research* **1998**, *306* (1), 127-136.
64. Chen, Y.; Li, Y.; Yu, H.; Sugiarto, G.; Thon, V.; Hwang, J.; Ding, L.; Hie, L.; Chen, X., Tailored Design and Synthesis of Heparan Sulfate Oligosaccharide Analogues Using Sequential One-Pot Multienzyme Systems. *Angewandte Chemie International Edition* **2013**, *52* (45), 11852-11856.
65. Li, Y.; Yu, H.; Thon, V.; Chen, Y.; Muthana, M. M.; Qu, J.; Hie, L.; Chen, X., Donor substrate promiscuity of the N-acetylglucosaminyltransferase activities of *Pasteurella multocida* heparosan synthase 2 (PmHS2) and *Escherichia coli* K5 KfiA. *Applied Microbiology and Biotechnology* **2014**, *98* (3), 1127-1134.
66. Chen, Y.; Thon, V.; Li, Y.; Yu, H.; Ding, L.; Lau, K.; Qu, J.; Hie, L.; Chen, X., One-pot three-enzyme synthesis of UDP-GlcNAc derivatives. *Chemical Communications* **2011**, *47* (38), 10815-10817.
67. Zhang, X.; Lin, L.; Huang, H.; Linhardt, R. J., Chemoenzymatic Synthesis of Glycosaminoglycans. *Accounts of Chemical Research* **2019**.
68. Errey, J. C.; Mann, M. C.; Fairhurst, S. A.; Hill, L.; McNeil, M. R.; Naismith, J. H.; Percy, J. M.; Whitfield, C.; Field, R. A., Sugar nucleotide recognition by *Klebsiella pneumoniae* UDP-d-galactopyranose mutase: Fluorinated substrates, kinetics and equilibria. *Organic & Biomolecular Chemistry* **2009**, *7* (5), 1009-1016.
69. Baisch, G.; Öhrlein, R.; Katopodis, A.; Streiff, M.; Kolbinger, F., Synthetic potential of cloned fucosyl-transferase III and VI. *Bioorganic & Medicinal Chemistry Letters* **1997**, *7*, 2447–2450.
70. Lairson, L. L.; Henrissat, B.; Davies, G. J.; Withers, S. G., Glycosyltransferases: Structures, Functions and Mechanisms. *Annual Review of Biochemistry* **2008**, *77*, 521-555.
71. Murray, B. W.; Wittmann, V.; Burkart, M. D.; Hung, S.-C.; Wong, C.-H., Mechanism of Human α -1,3-Fucosyltransferase V: Glycosidic Cleavage Occurs Prior to Nucleophilic Attack. *Biochemistry* **1997**, *36* (4), 823-831.
72. Wang, W.; Hu, T.; Frantom, P. A.; Zheng, T.; Gerwe, B.; del Amo, D. S.; Garret, S.; Seidel, R. D.; Wu, P., Chemoenzymatic synthesis of GDP-l-fucose and the Lewis X glycan derivatives. *Proceedings of the National Academy of Sciences* **2009**, *106* (38), 16096-16101.
73. Lee, H.-Y.; Chen, C.-Y.; Tsai, T.-I.; Li, S.-T.; Lin, K.-H.; Cheng, Y.-Y.; Ren, C.-T.; Cheng, T.-J. R.; Wu, C.-Y.; Wong, C.-H., Immunogenicity Study of Globo H Analogues with Modification at the Reducing or Nonreducing End of the Tumor Antigen. *Journal of the American Chemical Society* **2014**, *136* (48), 16844-16853.
74. Watts, A. G.; Withers, S. G., The synthesis of some mechanistic probes for sialic acid processing enzymes and the labeling of a sialidase from *Trypanosoma rangeli*. *Canadian Journal of Chemistry* **2004**, *82* (11), 1581-1588.
75. Li, Y.; Yu, H.; Cao, H.; Lau, K.; Muthana, S.; Tiwari, V. K.; Son, B.; Chen, X., *Pasteurella multocida* sialic acid aldolase: a promising biocatalyst. *Applied Microbiology and Biotechnology* **2008**, *79* (6), 963-970.
76. Chokhawala, H. A.; Cao, H.; Yu, H.; Chen, X., Enzymatic Synthesis of Fluorinated Mechanistic Probes for Sialidases and Sialyltransferases. *Journal of the American Chemical Society* **2007**, *129* (35), 10630-10631.

77. Petrie III, C. R.; Sharma, M.; Slimmons, O. D.; Korytnyk, W., Synthesis of analogs of N-acetylneuraminic acid and their effect on CMP-sialate synthase. *Carbohydrate Research* **1989**, *186*, 326–334.
78. Burkart, M. D.; Vincent, S. P.; Wong, C.-H., An efficient synthesis of CMP-3-fluoroneuraminic acid. *Chemical Communications* **1999**, 1525-1526.
79. Heise, T.; Pijnenborg, J. F. A.; Büll, C.; Van Hilten, N.; Kers-Rebel, E. D.; Balneger, N.; Elferink, H.; Adema, G. J.; Boltje, T. J., Potent Metabolic Sialylation Inhibitors Based on C-5-Modified Fluorinated Sialic Acids. *Journal of Medicinal Chemistry* **2019**, *62* (2), 1014-1021.
80. Chiu, C. P. C.; Watts, A. G.; Lairson, L. L.; Gilbert, M.; Lim, D.; Wakarchuk, W. W.; Withers, S. G.; Strynadka, N. C. J., Structural analysis of the sialyltransferase CstII from *Campylobacter jejuni* in complex with a substrate analog. *Nat Struct Mol Biol* **2004**, *11* (2), 163-170.
81. Watts, A. G.; Damager, I.; Amaya, M. L.; Buschiazio, A.; Alzari, P. M.; Frasch, A. C.; Withers, S. G., *Trypanosoma cruzi* Trans-sialidase operates through a covalent sialyl-enzyme intermediate: Tyrosine is the Catalytic nucleophile. *Journal of the American Chemical Society* **2003**, *125*, 7532–7533.
82. Liu, J. L. C.; Shen, G. J.; Ichikawa, Y.; Rutan, J. F.; Zapata, G.; Vann, W. F.; Wong, C. H., Overproduction of CMP-sialic acid synthetase for organic synthesis. *Journal of the American Chemical Society* **1992**, *114* (10), 3901-3910.
83. Cao, H.; Li, Y.; Lau, K.; Muthana, S.; Yu, H.; Cheng, J.; Chokhawala, H. A.; Sugiarto, G.; Zhang, L.; Chen, X., Sialidase substrate specificity studies using chemoenzymatically synthesized sialosides containing C5-modified sialic acids. *Organic & Biomolecular Chemistry* **2009**, *7* (24), 5137-5145.
84. Hartlieb, S.; Günzel, A.; Gerardy-Schahn, R.; Münster-Kühnel, A. K.; Kirschning, A.; Dräger, G., Chemoenzymatic synthesis of CMP-N-acetyl-7-fluoro-deoxy-neuraminic acid. *Carbohydrate Research* **2008**, *343*, 2075–2082.
85. Khedri, Z.; Li, Y.; Muthana, S.; Muthana, M. M.; Hsiao, C.-W.; Yu, H.; Chen, X., Chemoenzymatic synthesis of sialosides containing C7-modified sialic acids and their application in sialidase substrate specificity studies. *Carbohydrate Research* **2014**, *389*, 100–111.
86. Morley, T. J.; Withers, S. G., Chemoenzymatic Synthesis and Enzymatic Analysis of 8-Modified Cytidine Monophosphate-Sialic Acid and Sialyl Lactose Derivatives. *Journal of the American Chemical Society* **2010**, *132* (27), 9430-9437.
87. Yu, H.; Chokhawala, H.; Karpel, R.; Yu, H.; Wu, B.; Zhang, J.; Zhang, Y.; Jia, Q.; Chen, X., A Multifunctional *Pasteurella multocida* Sialyltransferase: A Powerful Tool for the Synthesis of Sialoside Libraries. *Journal of the American Chemical Society* **2005**, *127* (50), 17618-17619.
88. Petrie III, C. R.; Korytnyk, W., A High-Performance Liquid Chromatography Method for the Assay of Cytidine Monophosphate-Sialic Acid Synthetase. *Analytical Biochemistry* **1983**, *131*, 153–159.
89. Conradt, H. S.; Bünsch, A.; Brossmer, R., Preparation of 9-fluoro-9-deoxy-N-[2-¹⁴C]acetylneuraminic acid. Activation and transfer onto asialo-a1-acid glycoprotein. *FEBS Letters* **1984**, *170* (2), 295–300.
90. Khedri, Z.; Muthana, M. M.; Li, Y.; Muthana, S.; Yu, H.; Cao, H.; Chen, X., Probe sialidase substrate specificity using chemoenzymatically synthesised sialosides containing C9-modified sialic acid. *Chemical Communications* **2012**, *48*, 3357–3359.

91. Miyazaki, T.; Sato, H.; Sakakibara, T.; Kajihara, Y., An Approach to the Precise Chemoenzymatic Synthesis of ^{13}C -Labeled Sialyloligosaccharide on an Intact Glycoprotein: A Novel One-Pot $[3\text{-}^{13}\text{C}]$ -Labeling Method for Sialic Acid Analogues by Control of the Reversible Aldolase Reaction, Enzymatic Synthesis of $[3\text{-}^{13}\text{C}]\text{-NeuAc-}\alpha\text{-(2}\rightarrow\text{3)-[U-}^{13}\text{C}]\text{-Gal-}\beta\text{-(1}\rightarrow\text{4)-GlcNAc-}\beta\text{-}$ Sequence onto Glycoprotein, and Its Conformational Analysis by Developed NMR Techniques. *Journal of the American Chemical Society* **2000**, *122* (24), 5678-5694.
92. Miyazaki, T.; Sakakibara, T.; Sato, H.; Kajihara, Y., Chemoenzymatic Synthesis of the 9-Deoxy- 9-fluoro- $[3\text{-}^{13}\text{C}]\text{-NeuAc-}\alpha\text{-(2}\rightarrow\text{6)-[U-}^{13}\text{C}]\text{-Gal-}\beta\text{-}$ Sequence on an Intact Glycoprotein. *Journal of the American Chemical Society* **1999**, *121* (6), 1411-1412.