

Behavioural validation of a parent-report measure of child food fussiness

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Rendall, S., Dodd, H. ORCID: <https://orcid.org/0000-0003-1446-5338> and Harvey, K. ORCID: <https://orcid.org/0000-0002-6819-0934> (2020) Behavioural validation of a parent-report measure of child food fussiness. *Appetite*, 154. 104796. ISSN 0195-6663 doi: 10.1016/j.appet.2020.104796 Available at <https://centaur.reading.ac.uk/91633/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.appet.2020.104796>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Running Title: Validation of the CEBQ FF

Behavioural validation of a parent-report measure of child food fussiness

Stella Rendall¹, Helen Dodd², Kate Harvey³

^{1,2,3} School of Psychology & Clinical Language Sciences, University of Reading, Reading,
RG6 6AL, UK

All correspondence should be addressed to: ³Email: k.n.harvey@reading.ac.uk (Kate Harvey)

Not for publication: Tel +44 (0)118 378 7524

School of Psychology & Clinical Language Science, University of Reading, Reading, RG6
6AL, UK.

Abstract

Food fussiness is the rejection of familiar and novel foods leading to consumption that is insufficient and/or inadequately varied. Its importance to children's nutrition and the development of food preferences means it has been the focus of extensive research. To measure food fussiness, research has predominantly relied on parent-report, though parents' reporting of their child's eating behaviour can be reliable, responses may also be subject to bias. Utilising data from video-recordings of sixty-seven mother-child dyads during a meal in the home environment, this study aimed to validate the most widely used parent-report questionnaire measuring food fussiness against independent observations of children's eating behaviour and, in so doing, determine its accuracy. Maternal reported food fussiness, assessed using the Food Fussiness subscale of the Children's Eating Behaviour Questionnaire (CEBQ; Wardle, Guthrie, Sanderson, & Rapoport, 2001) was compared to children's observed food rejection and acceptance behaviours. Bootstrapped Pearson's correlations revealed that maternal reports of food fussiness were significantly positively related to food rejection behaviours and significantly negatively related to food acceptance behaviours. Maternal reports of food fussiness were also found to be significantly negatively related to the proportion of familiar/appealing of familiar foods consumed by the child. There was no significant association between maternal reported food fussiness and the proportion of familiar/unappealing, unfamiliar/appealing and unfamiliar/unappealing foods consumed by the child or the meal duration. These findings support the CEBQ FF as a valid measure of food fussiness.

Keywords: Food fussiness, Child, Mother, Parent-report, Observation, Mealtime behaviours

1 INTRODUCTION

Food fussiness, characterised by the rejection of familiar and novel foods resulting in a diet that is insufficient and/or inadequately varied (Dovey, Staples, Gibson, & Halford, 2008) is a common childhood problem, with a prevalence of 50% in children's second year (Carruth, Ziegler, Gordon, & Barr, 2004). As children this age would be unreliable reporters of their eating behaviour, most research in this field has used parent-report to assess food fussiness (e.g., Carruth et al., 1998; Galloway, Fiorito, Lee, & Birch, 2005; Hafstad, Abebe, Torgersen, & von Soest, 2013; Haycraft, Farrow, Meyer, Powell, & Blissett, 2011). The cost effectiveness and ease with which parent-report questionnaires can be administered on a large scale makes them practical (Carnell & Wardle, 2007), however, parent-report can be subject to biases and inconsistencies (e.g., Boquin, Moskowitz, Donovan, & Lee, 2014; Goh & Jacob, 2012). Although evidence suggests that parents can be reliable informants of their children's eating behaviour (e.g., Cooper, Whelan, Woolgar, Morrell, & Murray, 2004), research validating parent-report against independent observations of children's eating behaviour is crucial to comprehensively evaluate its reliability.

The Food Fussiness (FF) subscale of the CEBQ (Wardle et al., 2001) is widely used to assess food fussiness in young children (Farrow & Coulthard, 2012; Hendy, Williams, Riegel, & Paul, 2010; Jansen et al., 2012; Tharner et al., 2015; van der Horst, 2012). It has good internal validity (e.g., Wardle et al., 2001) and responses on the FF subscale are related to other parent-report measures of food fussiness. For example, the CEBQ was found to be accurate at discriminating between fussy and non-fussy eaters who were categorised using a structured parent interview (Steinsbekk, Hamre Sveen, Fildes, Llewellyn, & Wichstrøm, 2017). Similarly, Rogers, Ramsey and Blissett (2018) found the CEBQ FF subscale to have good criterion validity with the Montreal Children's Hospital Feeding Scale (MCHFS;

Ramsay, Martel, Porporino & Zygmuntowicz, 2011), a brief 14 item parent-report measure of children's feeding problems.

A handful of studies have aimed to establish the reliability of the CEBQ FF by comparing it to observations of children's eating. In one, Fernandez et al. (2018) observed children's responses to two familiar and two unfamiliar vegetables in a laboratory setting. They found that maternal responses on the CEBQ FF scale were associated with observed food refusal behaviours characterised by children's consumption of fewer grams of food, fewer bites, more negative utterances about the food, less compliance with maternal encouragements to eat and longer observed latency to the first bite. In another, Werthmann et al., (2015) offered children variants of a well-known yoghurt whilst they were in day care, with texture, taste and colour manipulated. Food acceptance was measured via the amount consumed. In contrast to Fernandez et al's (2018) laboratory study, Werthmann and colleagues found that parental reports of food fussiness on the CEBQ FF scale were not related to observations of children's yoghurt acceptance. Similarly, Surette, Ward, Morin, Vatanparast, & Bélanger, (2017) found that observed food fussiness, established from children's plate waste after a meal in a day care setting, did not correspond to parental reported CEBQ FF scores. Thus, there is some inconsistency regarding how well the CEBQ FF scale aligns with observed fussy eating. It should be noted that there is considerable disparity regarding how food fussiness was determined in the observations across these studies which could account for some of this inconsistency. For example, Fernandez et al. (2018) determined food fussiness by fewer grams of food consumed as well as the child's hedonic rating of food while plate waste analysis was used to establish a proxy measure of food fussiness in Surette et al's (2017) study.

Inconsistent findings could also arise because of study limitations. While the laboratory setting used by Fernandez et al. (2018) has the advantage of ensuring control of extraneous

variables, the artificial environment may also have elicited behaviours from children that were not typical for them. Arguably, while the day-care centres used by Werthmann et al. (2015) and Surette et al. (2017) can be considered more naturalistic, the setting may still introduce bias. Day-care settings have been found to produce elevated stress levels in young children, as peer groups are a demanding context and have been shown to produce high emotional arousal (Vermeer & van IJzendoorn, 2006). It is therefore plausible that the day-care environment, like the laboratory setting, may also influence children's eating behaviour in unanticipated ways.

The majority of young children are most familiar with eating meals at home, and so it is in this naturalistic environment that researchers are most likely to be able to observe children's food fussiness. Recently, Fries, Martin, & van der Horst, (2017) validated parental report of food fussiness by comparing CEBQ FF scores with video-recorded observations of children's food refusal in a home environment. Fries et al. found no differences in overall food refusal between fussy and non-fussy groups as defined by the CEBQ FF, however they acknowledge a key weakness in the design of their study. Specifically, parents were not guided in which food they offered their child and it is plausible that parents of fussy eaters may have chosen to offer foods they judged their child more likely to accept, thus explaining why fussy eaters displayed few food refusal behaviours during the observed mealtime. This interpretation was supported by their finding from questionnaire items indicating that parents who tended to "give up" after their child had refused disliked foods and provide them with an alternative meal consisting of their favourite foods had children who made more refusals when presented with a novel food.

The current study aimed to establish the validity of the CEBQ FF subscale using observational data while aiming to address the weaknesses of existing studies. Specifically, the focus was on ensuring the study was as naturalistic as possible, by observing children

123 eating a meal at home in the presence of their parent. The food offered was manipulated to
 124 comprise familiar and unfamiliar foods as well as foods likely to be broadly appealing and
 125 unappealing to children. Foods differ in their level of appeal to young children according to
 126 sensory characteristics such as texture, colour and taste. For example, foods with slimy and
 127 mushy textures as well as green foods have been found to be unappealing to young children
 128 while brightly coloured foods have been found to be appealing (Russell & Worsley, 2013).
 129 The foods chosen for each child were based on information provided by his/her parents, and
 130 represented a plausible meal, comprising soup, bread, fruit/vegetables and a dessert. Children
 131 were given age-appropriate portion sizes and parents were asked to behave in the way they
 132 usually would when offering a meal.

133 The objective was to validate the food fussiness subscale of the CEBQ by observing
 134 children's rejection and acceptance of familiar and unfamiliar foods in a naturalistic setting.
 135 It was hypothesised that higher scores on the CEBQ FF would be associated with more
 136 observed food rejection behaviours and fewer food acceptance behaviours. It was also
 137 hypothesised that higher scores on the CEBQ FF will be associated with less consumption of
 138 all food types (familiar/appealing, familiar/unappealing, unfamiliar/appealing and
 139 unfamiliar/unappealing) and this association is expected to be strongest for
 140 unfamiliar/unappealing foods and weakest for familiar/appealing foods. Finally, it was further
 141 hypothesized that higher scores on the CEBQ FF will be associated with longer meal
 142 duration.

144 2 METHOD

145 2.1 Participants

146 Sixty-seven mother-child pairs took part in this study. It focused on children aged two to
 147 four years as this age range has been found to be associated with increased parent perception
 148 of child food fussiness (Carruth et al., 2004; Hafstad et al., 2013). Previous studies in this

field demonstrate that few fathers typically volunteer to participate in research of this kind (see Patrick & Nicklas, 2005; Holley, Haycraft & Farrow, 2017). To avoid the methodological limitation of having a mixed sex parental group, but insufficient fathers for sub-group analysis, it was decided that the eligibility criteria for the study would be mothers and their child aged from two to four years, therefore, only mothers were invited to participate. We acknowledge that this limits the conclusions we can draw from this study and discuss the implications of the decision below. The mean age of children who participated was 3 years (S.D = 1 year) and the sample consisted of 39 girls and 28 boys. Mothers' age ranged from 22 to 45 years (M = 36 years; S.D = 5 years); most were well-educated (65.7% had an undergraduate or postgraduate degree), the majority described themselves as white British (80.6%) (OPCS; 2003) and almost all were living with a spouse/partner (92.5%). Two exclusion criteria were employed. Firstly, because the foods selected for the mealtime observation could contain nuts and dairy, children were excluded if their mother reported diagnosed nut allergies or lactose intolerance. Secondly, children with developmental disorders may have unusual eating habits due to motor problems and/or sensory difficulties and so children were excluded if their mothers reported atypical development or failure to meet developmental milestones.

2.2 *Measures*

Mothers completed a background questionnaire which recorded the child's age and sex (male or female) as well as the mother's ethnicity, marital status, education and age. Maternal ethnicity was evaluated using the Office of Population Censuses and Surveys (OPCS; 2003) 17 group ethnic classification which combines ethnic and national group dimensions (e.g. White Irish, Black African, Asian Pakistan). Marital status was assessed using three categories (single, living with spouse/partner and not living with spouse/partner). Maternal

education was based on three stages of education in England; primary, secondary and higher. For higher education, the sub- categories were undergraduate and postgraduate degree qualification.

2.2.1 *CEBQ Food Fussiness Subscale CEBQ FF (Wardle et al., 2001)*

The CEBQ FF was used to assess mother's perception of their child's food fussiness. The subscale consists of six statements which evaluate whether the child eats a variety of foods, the child's interest in new foods and how difficult the child is to please with meals e.g. *my child decides he/she doesn't like a food, even without tasting it*. Three of the six statements which allude to food acceptance, e.g. *"my child is interested in tasting food he/she hasn't tasted before"* are reverse coded. Respondents rate on a 5-point Likert scale (1= never, 5= always) how applicable each statement is to their child. A global mean score is calculated which can range from one to five with higher scores reflecting greater child food fussiness. The CEBQ FF has been demonstrated as having high reliability with a Cronbach's alpha value of .91 (Wardle et al, 2001). For the current sample, Cronbach's alpha for food fussiness was 0.94.

2.2.2 *Food Checklist*

A food checklist was created to be completed by mothers with a view to providing a meal that represented a plausible meal (to include soup, fruit/vegetables, bread and dessert), which could be prepared in a standardised way and which was tailored for each child to include appealing and unappealing, familiar and unfamiliar foods. This was to ensure that children participating in the study were offered a meal that comprised liked and disliked, familiar and unfamiliar foods. This classification was done to delineate children's responses to each category. Foods to be included in the list were selected to be appealing or unappealing based on the characteristics of foods reported by parents of fussy eaters as being consistently avoided or preferred (Boquin, Smith-Simpson, Donovan, & Lee, 2014). Characteristics of

foods found to be unappealing to fussy eaters include foods with slippery and mushy textures, foods with sour and bitter tastes, food with strong aromas, mixed foods with complex ingredients, soups and most vegetables. Foods that appeal to fussy eaters were found to be sweet, crunchy, salty or have bland and simple flavours. These include desserts, milk, pastries and sweet fruits. The food items included in the checklist are shown in Table 1.

[Table 1 here]

2.3 Procedure

Ethical approval for this study was obtained from the local Research Ethics Committee (UREC 15/43/KH). Children were recruited from a university Child Development Group Database which contains the details of over 2000 families with children in this age group. The database comprised details of families from the Royal Berkshire Hospital in Reading who were invited to participate in future psychological research by joining the University of Reading Infant Panel. Potential participants are recruited via researchers making regular visits to the post-delivery ward, and parents who express an interest are added to the database (at this stage, they are consenting to being approached by researchers in the future). The database is representative of the local population in some respects, for example participants in the present study were predominantly White British (81%) which is also fairly representative of Reading's demographics.

Mothers were contacted either via email or telephone and given a brief overview of the study as well as the inclusion/exclusion criteria. Out of 375 mothers contacted, 23 confirmed that their child had been diagnosed as lactose intolerant or with nut allergies making them ineligible. Of the 352 eligible mothers, 195 did not respond and a further 68 responded to say that they were unavailable to participate (for example, they had moved out of the area). The remaining 89 mothers (25% of those eligible) agreed to participate and provided an email

address to receive a demographic and Food Fussiness questionnaire. Of these mothers, 22 did not participate because they could not be available for the observational study. Consistent with Research Ethics Committee directions, mothers were not required to explain non-participation in the observational study. Those who chose to so typically gave reasons such as their child being ill or other commitments meaning a convenient time for a home visit could not be arranged. The final sample comprised 67 mothers, 75% of those who agreed to participate and 19% of eligible mothers contacted. G*Power 3 (Faul, Erdfelder, Lang, & Buchner, 2007) was used to establish that the final sample of 67 participants was sufficient to meet Cohen's (1992) power recommendation and yield statistical β power of more than 0.80 (based on $\alpha = 0.05$) and to detect medium correlational effects ($r = 0.33$).

When mothers agreed to participate, they were emailed a checklist of nineteen foods and asked to indicate for each food whether their child was likely to find the food familiar and appealing, familiar and unappealing, unfamiliar and appealing or unfamiliar and unappealing. This classification was done to delineate children's responses to each category, as explained above. This was to ensure that children participating in the study were offered a meal that comprised liked and disliked, familiar and unfamiliar foods. To avoid the food checklist influencing their perception of their children's food fussiness, mothers completed the CEBQ before the food checklist. Upon completion of the questionnaire, researchers arranged a convenient date for a home visit. In advance of the home visit, mothers were informed of the food items that the researcher would be bringing for the child's lunch (based on their responses on the food checklist). For each child, the completed checklist was used to select one food for each of the following categories: familiar and appealing; familiar and unappealing; unfamiliar and appealing; or unfamiliar and unappealing). The researcher explained to mothers that their child needed to be observed eating the meal without the influence of family members eating at the same time and were asked to identify a mealtime

that would be most convenient; either lunch or evening meal. Mothers were asked not to feed their children for two hours prior to the meal with the aim of controlling for hunger.

2.4 *Mealtime Observation*

Children were observed in their homes during a typical meal. On arrival, following greetings, the researcher showed the mother the food items to be prepared for the child and assisted the mother in the meal preparation. Each child was provided with a meal comprising four food items two of which were familiar (appealing and unappealing) and two of which were unfamiliar (appealing and unappealing). An example of a meal might be 100g ready-made lentil dahl soup (unfamiliar and unappealing), one slice granary bread equivalent to 38g (familiar and unappealing), 16 seedless green grapes equivalent to 75g (familiar and appealing) and half a custard tart equivalent to 80g (unfamiliar and appealing) totalling about 420 kcal. To determine the proportion of food that the child had consumed, each portion of food was weighed by the researcher using a Salter digital kitchen weighing scale before it was placed on the child's plate and leftovers were weighed by the researcher after the child had finished eating. The proportion of food consumed was the amount of food eaten relative to the total amount of food presented. For example, if the food given to the child weighed 100g before and the leftovers weighed 80g, meaning the child consumed 20g, therefore the proportion of food consumed would be 20/100 which is 0.2. A video camera was used to capture the child's eating behaviour during the meal which was placed on a tripod and positioned in the dining area. To diminish social desirability effects, where the child might be inclined to behave differently because of the video camera, the camera was set up about 15-20 minutes prior to the meal and the researcher made conversation with the child with the intention of familiarising him/her to both the researcher and the video camera. During this time, the child was shown an age appropriate information sheet in the form of cartoon images

depicting the stages of the meal observation. The researcher explained to the child that they
 were first going to play a game that would be video recorded, thus explaining the presence of
 the camera. The game took place where the child would later eat his/her meal and involved a
 popular children's card game called "tummy ache". The researcher played this game with the
 child and the mother until the child felt at ease and was comfortable playing with the
 researcher alone at which point the mother took the opportunity to leave and prepare the
 child's meal. If the child was unwilling to play the game or too young to comprehend the
 game, he/she was invited to do a drawing of their favourite meal or indicate their favourite
 foods from the pack of cards. When the food had been prepared, the researcher left the room
 and the mother invited the child to eat. This was to ensure the meal was as typical as possible.
 Mothers were asked to behave as they usually would during a typical meal, for example,
 encouraging their child to eat if that is what they would typically do. Although, being seated
 and eating with their child may have been the norm for some mothers, they were asked not to
 eat at all, specifically asked not to eat from the presented food so that the amount of food
 eaten by the child could be accurately calculated. To ensure uniformity between meals,
 mothers were asked not to add to the meal, for example by offering butter, ketchup, cheese.
 Recording was stopped when mothers informed the researcher that the child had finished
 eating. Children were given stickers and thanked for participating while mothers were
 provided with a leaflet explaining the purpose of the study and thanked for their participation.

2.5 Coding Eating Behaviour

Video recordings of mealtimes were coded offline by the researcher using the Observer
 XT9 Software (<http://www.noldus.com/human-behaviourresearch/products/theobserver-xt-90>). Behavioural measures of food fussiness were obtained from previous literature (Fries et al., 2017; Klesges et al., 1983; Luchini, Lee, & Donovan, 2016; Timimi, Douglas, &

Tsiftsopoulou, 1997) which lists several mealtime behaviours that have been found to be associated with fussy eaters (see Table 2). As there was not an existing coding scheme that included all these behaviours together, one was adapted by integrating features from previously used coding schemes (e.g., Klesges et al., 19983; Luchini et al., 2016; Fries et al., 2017) and included a detailed description of the behaviours to be coded from the video recordings. The final inclusion of behaviours was informed by several pilot coding sessions.¹ Each behaviour was assigned a keyboard key and every time a particular behaviour was observed, it was scored by pressing the corresponding keyboard key. A second coder was trained by the first author until interrater reliability reached (calculated using the Observer XT9 software interrater reliability function) 90% agreement (Cohens $k = 0.896$, $p < 0.01$). The second coder subsequently coded 25% of the videos and reliability was high (percentage agreement between coders ranged from 79 - 92%).

[Table 2 here]

2.6 Data Analysis

The hypotheses and the data analytic plan were made prior to data collection and all data driven analyses are clearly identified and discussed accordingly. Correlation analyses were performed to test the hypotheses. Data were analysed using Statistical Package for Social Sciences (SPSS), version 23. Descriptive statistics were first computed. An examination of the normal probability plot and the histogram showed that the study variables were skewed and not normally distributed. Significant Shapiro-Wilk's tests for normality on all variables further indicated the violation of the assumption of normality making the data set unsuitable

¹ We acknowledge that child temperament in relation to child feeding is an important consideration and initially considered coding for emotional intensity such as crying and throwing tantrums as observed in a previous study (Fries et al, 2017). These behaviours, however, were not observed in any our pilot observations. Reviewing the videos, it can be confirmed it was rarely seen across our observations, and where it was observed, it was captured via existing codes.

for parametric analysis. The distribution of the variables was not improved using log, reciprocal or square root transformations, therefore a bootstrapping procedure to generate a 95% bias- corrected bootstrapped confidence intervals of the correlation coefficients (1000 samples, $N = 67$) was performed to test the study hypotheses. For child and maternal sociodemographic variables measured on a continuous scale (child age and maternal age), initial bootstrapped two-tailed Pearson's correlation analyses were conducted to check for significant associations between these variables with observed mealtime behaviours and food fussiness. For dichotomous child and maternal sociodemographic variables (child sex, maternal education, marital status and maternal ethnicity), bootstrapped independent samples t-tests were used to check if observed mealtime behaviours and food fussiness significantly differed by group. Significance levels were set at $p < .05$. Results indicated that the continuous sociodemographic variables were not significantly related to the study variables. For the dichotomous sociodemographic variables, results indicated that there was no significant difference between groups for observed mealtime behaviours and food fussiness. Therefore, sociodemographic variables were not included in further analyses (see Tables 1 and 2 in supplementary materials).

2.6.1 *Relationships between observed mealtime behaviours*

To explore relationships between observed mealtime behaviours, preliminary two -tailed bootstrapped Pearson's partial correlations controlling for mealtime duration were performed (see Table 3). An alpha of $p < 0.05$ was adopted for the analyses. Positive and negative child food comments were adjusted for total utterances by calculating a proportion score for positive and negative comments i.e. proportion of negative comments = negative comments/ (negative + positive comments). Results indicated that the majority of the mealtime observations associated with food rejection and avoidance namely food refusal, spitting out food, playing with food, licking food, touching food and child negative food comments were

all significantly positively correlated. The exception was smelling food followed by rejection, which was only significantly associated with food refusal, licking food and spitting food. However, like the majority of the behaviours associated with food rejection and avoidance, smelling food followed by rejection was significantly negatively related to mealtime behaviours associated with food acceptance. It was therefore decided to include smelling followed by food rejection as a food rejection mealtime behaviour. The results also indicated a significant positive relationship between the mealtime behaviours associated with food acceptance i.e. food consumption and child positive food comments.

2.6.2 Exploring relationships between CEBQ FF, observed mealtime behaviours and proportion of foods consumed

To test our main hypothesis, two-tailed bootstrapped Pearson's partial correlation analyses controlling for meal duration were used to investigate the relationship between mothers' responses on the CEBQ FF with observed food rejection and food acceptance mealtime behaviours, proportion of familiar/appealing, familiar/unappealing, unfamiliar/appealing and unfamiliar/unappealing foods consumed. Two tailed bootstrapped correlation analysis was also used to explore the relationships between maternal reported food fussiness and meal duration. Significance levels were set at $p < 0.05$.

3 RESULTS

Descriptive statistics for all measures and observed behaviours are displayed in Table 4. Mean scores on the CEBQ FF subscale for children in the current sample reflect those obtained from similar samples (e.g., de Barse et al., 2016; Holley, Farrow, & Haycraft, 2016). [Table 3 here]

[Table 4 here]

As indicated in Table 5, bootstrapped Pearson's partial correlation analyses revealed that maternal report of food fussiness was significantly positively correlated with the majority of mealtime behaviours associated with food rejection i.e. spitting food, playing with food, touching food, licking food, child negative food comments and food refusal. There was no correlation between maternal reported food fussiness and smelling food followed by rejection. Maternal reports of food fussiness were significantly negatively correlated to mealtime behaviours associated with food acceptance i.e. food consumption and child positive food comments. There was a significant negative correlation between maternal reports of food fussiness and the proportion of familiar/appealing foods consumed by the child. There was no significant correlation between maternal reported food fussiness and the proportion of familiar/unappealing foods, unfamiliar/appealing foods and unfamiliar/unappealing foods consumed. The correlation between maternal reported food fussiness and meal duration was also not significant which is included in Table 6 together with the correlations between meal duration and mealtime behaviours

[Table 5 here]

[Table 6 here]

4 DISCUSSION

The present study aimed to validate maternal reported child food fussiness using the Food Fussiness subscale of the CEBQ against independent observations of children's eating behaviour. Supporting the hypothesis, the results indicated that children whose mothers

reported greater levels of food fussiness exhibited more mealtime behaviours associated with food rejection and fewer mealtime behaviours associated with food acceptance. Maternal reported food fussiness was associated with more spitting food, touching food, licking food, food refusal, playing with food and more negative food comments by the child. Maternal reported food fussiness was also associated with less food consumption and fewer positive food comments by the child. This is consistent with previous findings where children categorised as fussy eaters have been reported to display more food rejection behaviours and less food acceptance behaviours during mealtimes in comparison to non-fussy eaters (e.g., Fries et al., 2017; Fernandez et al., 2018). Maternal reported food fussiness was not associated with smelling food followed by rejection contrary to the hypothesis.

In addition, as expected, children whose mothers reported greater levels of food fussiness consumed smaller proportions of familiar/appealing foods during the observed mealtime. However, the finding of a non-significant correlation between maternal reported food fussiness and the proportion of other food types consumed (i.e. familiar/unappealing, unfamiliar/appealing and unfamiliar/unappealing) does not support the hypothesis. These findings are contrary to the expectation of the strongest association between CEBQ FF scores and less consumption of unfamiliar/unappealing foods and weakest for familiar/appealing foods. Our findings show that the opposite- that maternal reported food fussiness is only associated with less consumption of familiar and appealing foods. These findings make sense given that children are considered fussy because they tend to dislike and refuse foods that children would usually eat. It is not unusual for children to refuse foods which are unfamiliar and unappealing to most children such as spinach and broccoli and they would not be labelled as fussy eaters as a result. Non-significant findings between maternal reported food fussiness and the proportion of familiar/unappealing and unfamiliar/unappealing foods consumed can also be attributed to floor effects, as the data indicate that children did not consume enough of

these food types for associations with food fussiness to be found (See figure 1 in supplementary material). This is plausible given that children regardless of whether they are fussy eaters are less likely to consume familiar and unfamiliar foods they consider to be unappealing.

Also contrary to expectations and to previous research where parents of fussy eaters have described their children as slow eaters who usually have prolonged feeding times (e.g., Reau, Senturia, Lebailly, & Christoffel, 1996; Timimi et al, 1997), the present study found that maternal reported food fussiness was not associated with mealtime duration. This finding is consistent with those of previous studies that have used observational approaches to investigate meal duration in fussy eaters (e.g., Fries et al., 2017; Jacobi, Agras, Bryson, & Hammer, 2003). It should be noted that studies that have found lengthened mealtimes to be a behavioural indicator of food fussiness have relied on parent-report. It is possible that the associations found in these studies may be explained by parents perceiving the mealtime as lasting longer because of their struggles to encourage food consumption. A possible explanation for the lack of association between food fussiness and meal duration in this study may be that fussy children rejected most of the food offered, curtailing the duration of the meal. In contrast, some less fussy children might have spent more time consuming the food, resulting in longer meal duration. The significant positive association between food consumption and mealtime duration in the present study as indicated in Table 6 lends support to this argument. In the present study, as mothers were asked to sit with their child during the meal, it is also possible that their expectations of whether their child was likely to consume the food might have affected the meal duration. For instance, it was observed that some mothers expected their children to eat some of the food and used verbal prompts and some pressure to encourage, resulting in longer meal durations. Other mothers did not expect their children to consume all/any of the food, did not encourage consumption, and did not resist

when the child refused the meal, thus ending the mealtime quickly. There is also the possibility that if mothers had provided and prepared the foods, they would have expected their child to like it and therefore used more strategies to encourage food consumption leading to longer meal durations. In the present context, however, mothers may have had no expectations for their child to consume the food given that it was provided and prepared by the researchers, therefore did not encourage food consumption when the child refused to eat resulting in shorter meal durations.

Mealtime food rejection behaviours found to be associated with food fussiness in previous studies (e.g., Boquin, Smith-Simpson, Donovan, & Lee, 2014; Fries et al., 2017; Klesges et al., 1983) were also observed in this study. Children were observed playing with food, verbally and physically refusing food, spitting food out, touching and licking food without consuming it and making negative comments about food. The non-significant association between smelling food followed by rejection and maternal reports of food fussiness in the present study is consistent with the findings of previous studies where smelling food was found to be unrelated to parent-reported food fussiness (e.g., Johnson, Davies, Boles, Gavin & Bellows, 2015; Momin et al., 2018). However, while smelling food has been reported to occur infrequently during mealtimes (e.g., Blissett, Bennett, Donohoe, Rogers, & Higgs, 2012), the present study found that smelling food occurred quite frequently during the mealtime observation. Children were observed to display this behaviour on occasions that led to both food rejection and food acceptance. Smelling followed by food rejection, however, was observed to occur more frequently than smelling followed by food acceptance and was found to be significantly negatively related to food acceptance behaviours i.e. food consumption and child positive food comments as indicated in Table 3. It is possible that smelling food may have been used as an exploratory strategy by children who were suspicious of some unfamiliar foods. Fussy eaters aged 2-5 years have been observed to

become suspicious and inspect food during mealtimes by touching and licking presented food (e.g., Boquin, Smith-Simpson, Donovan, & Lee, 2014; Luchini et al., 2016). In the present study, children's decision to accept or reject food following smelling may have been dependent on how appealing or unappealing they found the smell, with appealing smells resulting in food acceptance and unappealing smells in food rejection. While this proposed pattern could not be confirmed in the present study, future replications could determine whether smelling followed by food acceptance or food rejection is related to different foods, particularly foods children find appealing and unappealing. Given the findings of significant associations with food rejection and food acceptance behaviours, as well as its frequent occurrence during the observed meal, more research exploring smelling food as an important mealtime behaviour associated with food fussiness is warranted.

The main strength of this study is its use of a behavioural observation approach to explore children's eating behaviours in a naturalistic environment. This approach permitted objective measurement of the mealtime behaviours associated with food fussiness and offered insight into how maternal reported food fussiness relates to actual child mealtime behaviour. Observing children in their home environment, where they are likely to feel most at ease, minimises changes to behaviour that can arise in unfamiliar settings. Providing children with age-appropriate portion sizes representative of a plausible meal is another strength of this study and an improvement from methods where children's recommended portion sizes have been exceeded (e.g., Jacobi et al., 2003). Including familiar and unfamiliar foods from several food groups i.e. bread, vegetables, fruits, dessert, soup was an opportunity to observe how children approach a range of foods and provided the opportunity to observe food fussiness more broadly. This is an improvement from methods where familiar and unfamiliar foods have been limited to one food group (e.g., Fernandez et al., 2018).

499 Some limitations should be noted. First, the presence of the camera during the recorded
500 mealtime was likely to have affected children's behaviours. Although measures were taken to
501 ensure the child became accustomed to the presence of the camera before the mealtime
502 observation commenced, many children remained aware of its presence and this may have
503 altered their typical behaviours. Future replication where video-recording is unobtrusive
504 would address this limitation. Second, observation of children's eating behaviours was
505 limited to a single meal and it cannot be determined if the observed behaviours were typical
506 of the child. For example, some mothers commented on their child's unusual response to
507 some of the presented foods, for example "he/she usually likes avocados". Observing a
508 particular behaviour multiple times provides a more accurate representation (Young &
509 Drewett, 2000), therefore future research observing children on several occasions will help
510 improve reliability. Third, on reflection, offering all the food items at once is not
511 representative of a typical meal as children are not usually given their main meal together
512 with a dessert; indeed several mothers commented that they would not usually serve dessert
513 with the main meal. It is plausible that offering the dessert at the same time as the rest of the
514 food may have influenced children's decision to try the other food items. On subsequent
515 examination of the video recordings, it was observed that many children's attention was
516 initially drawn to the dessert as they found this most appealing. These children typically
517 consumed the dessert first and were then reluctant to try the other food items. It is unclear,
518 therefore, how children would have responded to these foods in the absence of the dessert.
519 Replication of this study where desserts are not included with other food items would help
520 provide a more accurate assessment of children's responses to familiar and unfamiliar food
521 items. Fourth, mothers were informed of the food items that the researcher brought for the
522 child's lunch prior to the mealtime observation (based on their responses on the food
523 checklist). Although it seems unlikely, it is possible that some mothers might have

subsequently exposed their children to some novel foods which may have influenced their children's responses to these foods during the observation. Future replications where mothers are not informed of the food their child will be eating during the observation would address this limitation. Fifth, the current study measured the frequencies of mealtime behaviours without accounting for their duration. For example, playing with food for 3 seconds was scored identically to playing with food for 15 seconds which is a limitation. However, as we were interested in the relationship between higher scores on the CEBQ FF and the number of occurrences of food rejection and acceptance behaviours during the recorded mealtime, measuring the presence or absence of a behaviour seemed more relevant than measuring its duration. Sixth, this study did not include a measure of neophobia which is a limitation given that children were asked to try unfamiliar foods. The inclusion of a food neophobia measure would have ascertained whether children with high food neophobia scores displayed more food rejection mealtime behaviours with unfamiliar foods. In addition, as food neophobia and food fussiness are considered as two separate constructs (Dovey et al., 2008), the inclusion of a food neophobia measure would have been useful to ascertain whether mothers conceptually differentiate between food fussiness and food neophobia. Such information would help determine if a mother's perception of food neophobia in her child also extends to the categorization of the child as a fussy eater on the CEBQ FF. Future replications would therefore benefit from an inclusion of a measure of food neophobia. Seventh, as is typical of research in this field (e.g., Powell, Farrow & Meyer, 2011; Farrow & Coulthard, 2012; Haycraft, Farrow & Blissett, 2013; Holley et al., 2017) the present findings cannot be generalised beyond the predominantly White British, well-educated mothers from two-parent households who agreed to participate in this study. The characteristics of our sample highlight the difficulty of recruiting participants with more diverse socio-demographic characteristics to research studies. Future studies should seek to replicate the findings with

other socio-demographic groups. For the reasons given above, fathers were not recruited to this study. This is typical of research in this field as participants in most studies using the CEBQ FF have been predominantly or exclusively mothers (e.g., Holley et al., 2017; Fernandez et al., 2018) either as a result of explicit inclusion criteria or because fathers are less likely to participate in research of this kind. While validating the subscale for mothers is of merit, it is important for research in this field to engage with fathers and their experiences of children's eating. This remains challenging given difficulty recruiting fathers into research as there have been reports of response rates of less than 10% from fathers when completing questionnaires directed at parents/caregivers (e.g., Patrick & Nicklas, 2005; Wardle, Carnell & Cooke, 2005). Finally, it should be noted that some mothers used some prompts to encourage food consumption in their children during the mealtime observation. It is possible that the use of prompts may have influenced child behaviour such that food refusal was in response to maternal control and not in response to the trait of food fussiness. Although this material falls beyond the scope of the present study which focuses on validating the CEBQ, further research investigating the relationship between maternal prompts and food fussiness is required.

5 CONCLUSIONS

Overall, the correspondence between independent observations of children's food rejection and acceptance behaviours with maternal reports of food fussiness suggests that mothers provide accurate and reliable information regarding their children's eating behaviour. These findings are plausible as mothers are often the main caregivers and tend to spend considerable time with their children in various settings, including mealtimes (Carnell & Wardle, 2007). The findings lend support to previous research that found maternal reports

of child eating to be a reliable reflection of independent observations (e.g., Carnell & Wardle, 2007; Fernandez et al., 2018) while improving on previous methods by observing children in a naturalistic setting and including a variety of foods. Importantly, these results validate the Food Fussiness subscale of the CEBQ as an accurate measure of child food fussiness that can be used by researchers and health practitioners with confidence.

All authors have reviewed and approved the complete manuscript and accept full responsibility for all aspects of the work described.

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declarations of interest: none

REFERENCES

- Blissett, J., Bennett, C., Donohoe, J., Rogers, S., & Higgs, S. (2012). Predicting Successful Introduction of Novel Fruit to Preschool Children. *Journal of the Academy of Nutrition and Dietetics*, 112(12), 1959–1967. <https://doi.org/10.1016/j.jand.2012.08.014>
- Boquin, M. M., Moskowitz, H. R., Donovan, S. M., & Lee, S. Y. (2014). Defining perceptions of picky eating obtained through focus groups and conjoint analysis. *Journal of Sensory Studies*. <https://doi.org/10.1111/joss.12088>
- Boquin, M., Smith-Simpson, S., Donovan, S. M., & Lee, S. Y. (2014). Mealtime Behaviors and Food Consumption of Perceived Picky and Nonpicky Eaters through Home Use Test. *Journal of Food Science*, 79(12), M2523-S2532. <https://doi.org/10.1111/1750-3841.12698>
- Carnell, S., & Wardle, J. (2007). Measuring behavioural susceptibility to obesity: Validation of the child eating behaviour questionnaire. *Appetite*, 48(1), 104–113. <https://doi.org/10.1016/j.appet.2006.07.075>
- Carruth, B. R., Skinner, J., Houck, K., Moran, J., Coletta, F., & Ott, D. (1998). The phenomenon of “picky eater”: a behavioral marker in eating patterns of toddlers. *Journal of the American College of Nutrition*. <https://doi.org/10.1080/07315724.1998.10718744>
- Carruth, B.R., Ziegler, P. J., Gordon, A., & Barr, S. I. (2004). Prevalence of picky eaters among infants and toddlers and their caregivers’ decisions about offering a new food. *Journal of the American Dietetic Association*, 104(SUPPL. 1), 57–64. <https://doi.org/10.1016/j.jada.2003.10.024>
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155.
- Cooper, P. J., Whelan, E., Woolgar, M., Morrell, J., & Murray, L. (2004). Association between childhood feeding problems and maternal eating disorder: Role of the family environment. *British Journal of Psychiatry*, 184(MAR.), 210–215. <https://doi.org/10.1192/bjp.184.3.210>

- de Barse, L. M., Cano, S. C., Jansen, P. W., Jaddoe, V. V. W., Verhulst, F. C., Franco, O. H.,
 ... Tharner, A. (2016). Are parents' anxiety and depression related to child fussy eating?
Archives of Disease in Childhood, 101(6). <https://doi.org/10.1136/archdischild-2015-309101>
- Dovey, T. M., Staples, P. A., Gibson, E. L., & Halford, J. C. G. (2008). Food neophobia and
 "picky/fussy" eating in children: A review. *Appetite*.
<https://doi.org/10.1016/j.appet.2007.09.009>
- Farrow, C. V., & Coulthard, H. (2012). Relationships between sensory sensitivity, anxiety
 and selective eating in children. *Appetite*, 58(3), 842–846.
<https://doi.org/10.1016/j.appet.2012.01.017>
- Faul, F., Erdfelder, E., Lang, a.-G., & Buchner, a. (2007). A Flexible statistical power
 analysis program for the social, behavioral, and biomedical sciences. *Behavioral Research*
Methods, 39(2), 175–191. <https://doi.org/10.3758/BF03193146>
- Fernandez, C., DeJesus, J. M., Miller, A. L., Appugliese, D. P., Rosenblum, K. L., Lumeng,
 J. C., & Pesch, M. H. (2018). Selective eating behaviors in children: An observational
 validation of parental report measures. *Appetite*, 127(May), 163–170.
<https://doi.org/10.1016/j.appet.2018.04.028>
- Fries, L. R., Martin, N., & van der Horst, K. (2017). Parent-child mealtime interactions
 associated with toddlers' refusals of novel and familiar foods. *Physiology and Behavior*, 176,
 93–100. <https://doi.org/10.1016/j.physbeh.2017.03.001>
- Galloway, A. T., Fiorito, L., Lee, Y., & Birch, L. L. (2005). Parental pressure, dietary
 patterns, and weight status among girls who are "picky eaters." *Journal of the American*
Dietetic Association, 105(4), 541–548. <https://doi.org/10.1016/j.jada.2005.01.029>
- Goh, D. Y., & Jacob, A. (2012). Perception of picky eating among children in Singapore and
 its impact on caregivers: a questionnaire survey. *Asia Pacific Family Medicine*, 11(1), 5.
<https://doi.org/10.1186/1447-056X-11-5>

- 645 Hafstad, G. S., Abebe, D. S., Torgersen, L., & von Soest, T. (2013). Picky eating in preschool
 646 children: The predictive role of the child's temperament and mother's negative affectivity.
 647 *Eating Behaviors*, 14(3), 274–277. <https://doi.org/10.1016/j.eatbeh.2013.04.001>
- 648 Haycraft, E., Farrow, C., Meyer, C., Powell, F., & Blissett, J. (2011). Relationships between
 649 temperament and eating behaviours in young children. *Appetite*, 56(3), 689–692.
 650 <https://doi.org/10.1016/j.appet.2011.02.005>
- 651 Haycraft, E.L., Farrow, C. & Blissett, J. (2013). Maternal symptoms of depression are related
 652 to observations of controlling feeding practices in mothers of young children. *Journal of*
 653 *Family Psychology*, 27 (1), 159-164.
- 654 Hendy, H. M., Williams, K. E., Riegel, K., & Paul, C. (2010). Parent mealtime actions that
 655 mediate associations between children's fussy-eating and their weight and diet. *Appetite*,
 656 54(1), 191–195. <https://doi.org/10.1016/j.appet.2009.10.006>
- 657 Holley, C. E., Farrow, C., & Haycraft, E. (2016). Investigating the role of parent and child
 658 characteristics in healthy eating intervention outcomes. *Appetite*, 105, 291–297.
 659 <https://doi.org/10.1016/j.appet.2016.05.038>
- 660 Holley, C.E., Haycraft, E., & Farrow, C. (2017). Predicting Children's Fussiness with
 661 vegetables: The role of feeding practices. *Maternal and Child Nutrition*, 14. [https://doi:](https://doi.org/10.1111/mcn.12442)
 662 10.1111/mcn.12442
- 663 Jacobi, C., Agras, W. S., Bryson, S., & Hammer, L. D. (2003). Behavioral validation,
 664 precursors, and concomitants of picky eating in childhood. *Journal of the American Academy*
 665 *of Child and Adolescent Psychiatry*, 42(1), 76–84. [https://doi.org/10.1097/00004703-](https://doi.org/10.1097/00004703-200306000-00025)
 666 200306000-00025
- 667 Jansen, P. W., Roza, S. J., Jaddoe, V. W., Mackenbach, J. D., Raat, H., Hofman, A., ...
 668 Tiemeier, H. (2012). Children's eating behavior, feeding practices of parents and weight
 669 problems in early childhood: results from the population-based Generation R Study. *The*

- 670 *International Journal of Behavioral Nutrition and Physical Activity*, 9(1), 130.
- 671 <https://doi.org/10.1186/1479-5868-9-130>
- 672 Johnson, S. L., Davies, P. L., Boles, R. E., Gavin, W. J., & Bellows, L. L. (2015). Young
- 673 Children's Food Neophobia Characteristics and Sensory Behaviors Are Related to Their
- 674 Food Intake. *Journal of Nutrition*, 145(11). <https://doi.org/10.3945/jn.115.217299>
- 675 Klesges, R. C., Coates, T. J., Brown, G., Sturgeon-Tillisch, J., Moldenhauer-Klesges, L. M.,
- 676 Holzer, B., ... Vollmer, J. (1983). Parental influences on children's eating behavior and
- 677 relative weight. *Journal of Applied Behavior Analysis*, 16(4), 371–378.
- 678 <https://doi.org/10.1901/jaba.1983.16-371>
- 679 Luchini, V., Lee, S. Y., & Donovan, S. (2016). Observed differences in child picky eating
- 680 behaviors between the child's home and centre-or home-based childcare. *FASEB Journal*, 30
- 681 (1). <https://doi.org/10.1016/j.appet.2017.04.021>
- 682 Momin, S. R., Hughes, S. O., Elias, C., Papaioannou, M. A., Phan, M., Vides, D., & Wood,
- 683 A. C. (2018). Observations of Toddlers' sensory-based exploratory behaviors with a novel
- 684 food. *Appetite*, 131(August), 108–116. <https://doi.org/10.1016/j.appet.2018.08.035>
- 685 Office of Population Censuses and Surveys (2003). *Census Ethnic Group and Country of*
- 686 *Birth (Census User Guide No. 9)*. London: HMSO
- 687 Patrick, H., & Nicklas, T. A. (2005). A review of family and social determinants of children's
- 688 eating patterns and diet quality. *Journal of the American College of Nutrition*, 24(2), 83–92.
- 689 <https://doi.org/24/2/83>
- 690 Powell, F. C., Farrow, C. V., & Meyer, C. (2011). Food avoidance in children. The influence
- 691 of maternal feeding practices and behaviours. *Appetite*, 57(3), 683–692.
- 692 <https://doi.org/10.1016/j.appet.2011.08.011>
- 693 Ramsay, M., Martel, C., Porporino, M., & Zygmuntowicz, C. (2011). The Montreal
- 694 children's hospital feeding scale: A brief bilingual screening tool for identifying feeding

- 695 problems. *Paediatrics and Child Health*, 16(3), 147–151.
- 696 <https://doi.org/10.1093/pch/16.3.147>
- 697 Reau, N. R., Senturia, Y. D., Lebailly, S. A., & Christoffel, K. K. (1996). Infant and toddler
- 698 feeding patterns and problems: Normative data and a new direction. *Journal of*
- 699 *Developmental and Behavioral Pediatrics*, Vol. 17, pp. 149–153.
- 700 <https://doi.org/10.1097/00004703-199606000-00002>
- 701 Rogers, S., Ramsey, M., & Blissett, J. (2018). The Montreal's children hospital feeding scale:
- 702 Relationship with parental report of child feeding behaviours and observed feeding
- 703 interactions. *Appetite*, 125, 201-209.
- 704 Russell, C.G. & Worsley, A. (2013). Why don't they like that? And can I do anything about
- 705 it? The nature and correlates of parents' attributions and self-efficacy beliefs about preschool
- 706 children's food preferences. *Appetite*, 66, 34-43.
- 707 Steinsbekk, S., Hamre Sveen, T., Fildes, A., Llewellyn, C., & Wichstrøm, L. (2017).
- 708 Screening for pickiness -- a validation study. *International Journal of Behavioral Nutrition &*
- 709 *Physical Activity*, 14, 1–4. <https://doi.org/10.1186/s12966-016-0458-7>
- 710 Surette, V., Ward, S., Morin, P., Vatanparast, H., & Bélanger, M. (2017). Food Reluctance of
- 711 Preschool Children Attending Daycare Centers Is Associated with a Lower Body Mass Index.
- 712 *Journal of the Academy of Nutrition and Dietetics*, 117(11), 1749–1756.
- 713 <https://doi.org/10.1016/j.jand.2017.07.007>
- 714 Tharner, A., Jansen, P. W., Kiefte-de Jong, J. C., Moll, H. a., Hofman, A., Jaddoe, V. W. V.,
- 715 ... Franco, O. H. (2015). Bidirectional Associations between Fussy Eating and Functional
- 716 Constipation in Preschool Children. *The Journal of Pediatrics*, 166(1), 91-96.e1.
- 717 <https://doi.org/10.1016/j.jpeds.2014.09.028>
- 718 Timimi, S., Douglas, J., & Tsiftopoulou, K. (1997). Selective eaters: a retrospective case
- 719 note study. *Child: Care, Health and Development*, 23(3), 265–278.

- 720 van der Horst, K. (2012). Overcoming picky eating. Eating enjoyment as a central aspect of
 721 children's eating behaviors. *Appetite*, 58(2), 567–574.
 722 <https://doi.org/10.1016/j.appet.2011.12.019>
- 723 van der Horst, K., Eldridge, A., Deming, D., & Reidy, K. (2014). Caregivers' perceptions
 724 about picky eating: associations with texture acceptance and food intake (379.3). *The FASEB*
 725 *Journal*, 28(1_supplement), 373–379.
- 726 Vermeer, H. J., & van IJzendoorn, M. H. (2006). Children's elevated cortisol levels at
 727 daycare: A review and meta-analysis. *Early Childhood Research Quarterly*, 21(3), 390–401.
 728 <https://doi.org/10.1016/j.ecresq.2006.07.004>
- 729 Wardle, J., Guthrie, C. A., Sanderson, S., & Rapoport, L. (2001). Development of the
 730 Children's Eating Behaviour Questionnaire. *Journal of Child Psychology and Psychiatry, and*
 731 *Allied Disciplines*, 42(7), 963–970. <https://doi.org/10.1111/1469-7610.00792>
- 732 Wardle, J., Carnell, S., & Cooke, L. (2005). Parental control over feeding and children's fruit
 733 and vegetable intake: How are they related? *Journal of the American Dietetic Association*,
 734 105(2), 227–232. <https://doi.org/10.1016/j.jada.2004.11.006>
- 735 Werthmann, J., Jansen, A., Havermans, R., Nederkoorn, C., Kremers, S., & Roefs, A. (2015).
 736 Bits and pieces. Food texture influences food acceptance in young children. *Appetite*, 84,
 737 181–187. <https://doi.org/10.1016/j.appet.2014.09.025>
- 738 Young, B., & Drewett, R. (2000). Eating behaviour and its variability in 1-year-old children.
 739 *Appetite*, 35(2), 171–177.
- 740

741 Table 1: List of food items included in food checklist

742

Soups	Wholegrain Breads
Sainsbury's Thai beetroot soup	Tesco Rye Bread
Sainsbury's Petits pois and ham soup	Hovis Country Granary Bread
Sainsbury's lentil dahl soup	Tesco Walnut Loaf
Desserts	Fruits and Vegetables
Tesco free crème caramel dessert	Grapes
Sainsbury's mango and coconut panna cotta	Pears
Tesco custard tarts	Gooseberry
Waitrose pistachio flavour macaroons	Carrots
Tesco profiteroles	Sweetcorn
Asda Kulfi-ice pistachio ice cream	Avocado

743

744

745

746 Table 2: List of behaviours coded from the mealtime observation

Observed Mealtime Behaviours	Description of Behaviour (References)
Food Refusal	The child refuses the presented food by pushing the food away, turning their head away when the food is presented by the parent, ignoring the presented food or by verbally refusing to try the food. ^{7, 10}
Spitting food	The child places the food in their mouth and spits it out or vomits. ^{1, 3, 5, 9, 10}
Playing with food	The child plays with food by messing, stirring, throwing and crumbling the food or treating the food as well as the utensils as a toy but does not consume the food. ^{2, 3, 8}
Licking food	The child licks the presented food but does not consume it. ^{8, 9}
Touching food	The child touches the presented food but does not consume it. ^{8, 9}
Smelling food followed by rejection	The child smells the presented food and refuses to consume it.
Child Positive food comments	Positive sounds and comments the child expresses towards the presented food, e.g. “I like this”, “this tastes nice”, and “yum!”
Child negative food comments	Negative sounds and comments the child expresses towards the presented food. This includes complaints and expressions of disgust, e.g. “this tastes disgusting”, “Yuk!” ⁹
Food consumption	The child consumes the presented food; putting food in the mouth and swallowing it ^{4, 6, 8.}

Note: Previous studies that have cited the above mealtime behaviours associated with food fussiness.

1. Klesges et al. (1983); 2. Sanders et al. (1993); 3. Timimi et al. (1997); 4. Jacobi et al. (2003); 5. Lewinsohn et al. (2005); 6. Galloway et al., (2005); 7. Dovey et al. (2008); 8. Boquin, Smith-Simpson, Donovan, & Lee, (2014); 9. Luchini et al. (2016); 10. Fries et al., (2017).

Table 3: Two tailed bootstrapped Pearson's partial correlations between observed mealtime behaviours

	Food Refusal	Spitting food	Licking Food	Touching food	Smelling food followed by rejection	Child negative food comments	Maternal positive comments	Maternal negative comments	Food Consumption	Child positive food comments
Spitting food	.59**									
Playing with food	.54**	.71**								
Licking food	.42**	.50**								
Touching food	.60**	.27*	.20*							
Smelling food followed by rejection	.30*	.42*	.25*	.14						
Child negative food comments	.63**	.56**	.28*	.67**	.13					
Maternal positive food comments	.37*	.33*	.09	.44**	-.005	.59**				
Maternal negative food comments	-.06	-.10	.03	-.18	.005	-.13	-.19			
Food Consumption	-.44**	-.50**	-.25*	-.14	-.31*	-.34**	-.12	-.09		
Child positive food comments	-.12	-.21	-.25*	.01	-.36**	.14	.30*	-.05	.33**	
Meal duration included as a covariate *p < 0.05, **p < 0.001										

Table 4: Descriptive statistics for food fussiness and observed mealtime behaviours.

Measure	Median (IQR)	Mean (SD)	Min/Max
CEBQ FF score	3.00 (1.30)	3.00 (1.00)	1.00/5.00
Food refusal	6.00 (9.00)	8.00 (5.60)	1.00/22.00
Spitting food	0.00 (2.00)	2.00 (3.50)	0.00/16.00
Playing with food	0.00 (2.00)	1.90 (3.20)	0.00/15.00
Licking food	2.00 (3.00)	2.00 (2.30)	0.00/9.00
Touching food	4.00 (4.00)	4.00 (3.40)	0.00/16.00
Smelling food followed by rejection	1.00 (2.00)	1.50 (1.30)	0.00/15.00
Food consumption	25.00 (14.00)	27.00 (13.00)	5.00/66.00
Child negative food comments	4.00 (8.00)	7.00 (5.50)	0.00/21.00
Child positive food comments	5.00 (6.00)	6.00 (4.20)	0.00/17.00
Maternal negative food comments	0.00 (0.00)	0.03 (0.17)	0.00/1.00
Maternal positive food comments	5.00 (8.00)	6.34 (5.78)	0.00/25.00
Proportion of familiar/appealing foods consumed	0.71(0.71)	0.65(0.35)	0.03/ 1.00
Proportion of familiar/unappealing foods consumed	0.07 (0.15)	0.14 (0.18)	0.00/0.88
Proportion of unfamiliar/appealing foods consumed	0.31 (0.48)	0.39 (0.33)	0.00/1.00
Proportion of unfamiliar/unappealing foods consumed	0.05 (0.48)	0.18 (0.28)	0.00/1.00
Meal duration	19.00 (6.00)	19.00 (4.90)	9.00/29.00

Note. IQR = interquartile range, SD = standard deviation

Table 5: Two-tailed Pearson's partial correlations and bootstrapped 95% confidence intervals for relationships between maternal reports of food fussiness and observed mealtime behaviours

Observed mealtime behaviour	r	p	CI^{95%}
Food refusal	.49	<.001	[.27, .67]
Spitting food	.44	<.001	[.22, .61]
Playing with food	.46	<.001	[.23, .62]
Licking food	.36	.003	[.09, .56]
Touching food	.47	<.001	[.28, .63]
Smelling food followed by rejection	.22	.057	[-.03, .41]
Child negative food comments	.46	<.001	[.24, .62]
Food consumption	-.24	.046	[-.45, -.01]
Child positive food comments	-.35	.004	[-.55, -.13]
Proportion familiar/appealing food consumed	-.39	.001	[-.59, -.17]

Proportion familiar/unappealing food consumed	-.10	.418	[-.34, .09]
Proportion unfamiliar/appealing food consumed	-.09	.479	[-.30, .15]
Proportion unfamiliar/unappealing food consumed	-.06	.642	[-.29, .19]

Meal duration included as a covariate. $CI^{95\%}$ = 95% confidence interval, lower, upper bound values.

Table 6: Two-tailed Pearson correlations and bootstrapped 95% confidence intervals for relationships between meal duration, maternal reports of food fussiness and observed mealtime behaviours.

	r	p	CI ^{95%}
Food Fussiness	-.03	.795	[-.28, .21]
Spitting food	-.02	.868	[-.32, .24]
Playing with food	-.04	.739	[-.30, .20]
Licking food	-.13	.309	[-.37, .15]
Touching food	-.11	.386	[-.37, .18]
Smelling food followed by rejection	-.14	.268	[-.36, .11]
Food consumption	.30	.013	[.07, .52]
Food refusal	-.07	.585	[-.31, .19]
Child negative food comments	-.91	.440	[-.36, .19]
Child positive food comments	-.13	.285	[-.11, .38]
Maternal negative food comments	-.11	.384	[-.01, .27]
Maternal positive food comments	-.03	.827	[-.24, .29]

CI^{95%} = 95% confidence interval, lower, upper bound values.

Supplementary MaterialTable 1: Bootstrapped Pearson's correlations between child age, maternal age with food fussiness and observed mealtime behaviours

	Child age	Maternal age
Food Fussiness	.15	-.23
Smelling food followed by rejection	-.206	-.04
Touching food	.20	.07
Licking food	.02	.12
Playing with food	.11	-.01
Food refusal	.16	-.07
Food consumption	.17	.13
Spitting food	-.02	-.08
Child negative food comments	.05	-.09
Child positive food comments	-.06	-.03
Maternal negative food comments	.06	.01
Maternal positive food comments	.02	-.13
Meal duration	.05	.13

Table 2: Bootstrapped Independent Samples t-tests comparing means of child sex, maternal ethnicity, maternal education, marital status with food fussiness and observed mealtime behaviours.

	Child sex					Maternal Education					Marital Status					Maternal Ethnicity				
	Females		Males			University Degree		No University Degree			Single mothers		Married/Living with partner			White British		Other Ethnicity		
	M	SE	M	SE	t	M	SE	M	SE	t	M	SE	M	SE	t	M	SE	M	SE	t
Food fussiness	3.3	.16	3.0	.14	-.15	3.1	.18	3.2	.93	-.32	3.9	.04	3.1	.11	1.94	3.1	.11	3.5	.27	-1.7
Food Consumption	25.5	1.95	28.2	2.66	.83	25.7	2.7	27.1	1.9	-.43	29.2	1.7	26.4	1.5	.24	26.5	1.8	26.9	3.63	-.095
Spitting food	2.7	.59	1.4	.60	-.12	2.4	.83	1.8	.34	-.61	5.2	3.07	1.7	.38	1.1	2.5	1.2	1.9	.44	-.49
Playing with food	2.0	.57	1.6	.51	-.42	1.3	.68	2.1	.48	-.103	4.2	.29	1.6	.35	.85	1.8	.44	2.2	.86	-.45
Licking food	2.7	.39	1.8	.35	-.16	1.8	.34	2.6	.37	-1.13	3.6	1.6	2.2	.28	1.31	2.5	.32	1.6	.43	1.27
Touching food	4.5	.62	3.5	.49	-.99	3.1	.55	4.4	.56	-1.52	5.0	.83	3.8	.45	.71	3.8	.48	4.6	.79	-.77
Smelling food followed by rejection	1.7	.22	1.14	.22	-.17	71.3	.26	1.5	.21	-.51	2.6	.68	1.4	.69	2.05	1.47	.18	1.46	.48	.003
Food refusal	8.2	1.02	8.0	.81	-.18	7.6	1.2	8.4	.83	-.55	11.4	2.9	7.8	.69	1.37	8.1	.78	8.0	1.43	.04
Child negative food comments	6.6	.91	6.5	.98	-.105	5.8	.95	7.0	.89	-.82	9.8	2.44	6.3	.68	1.37	6.0	.73	9.1	1.51	-1.87
Child positive food comments	5.1	.67	5.8	.81	.77	5.1	.96	5.8	.61	.61	6.8	2.9	5.5	.51	.68	5.5	.60	5.8	1.04	-.19
Maternal negative food comments	.02	.02	.03	.03	.23	<.001	<.001	.04	.03	-1.4	<.001	<.001	.03	.02	-.40	<.001	<.001	.03	.03	.69

Maternal positive food comments	6.1	.87	6.7	1.2	.39	7.2	.94	4.7	.93	-1.9	10.2	3.12	6.03	.71	1.6	6.35	.82	6.31	1.36	.02
Meal duration	19.3	.89	18.7	.68	-.53	19.0	.70	18.9	1.11	-.13	18.6	.75	19.0	.64	-.47	18.7	.67	20.5	1.18	-.13

788 There were no significant differences between groups across all analyses. M = mean, SE = standard error mean.

789

790

Figure 1: Scatterplots depicting association between food fussiness and proportion of foods consumed



