

Are remote sensing evapotranspiration models reliable across South American ecoregions?

Article

Accepted Version

Melo, D. C. D., Anache, J. A. A., Borges, V. P., Miralles, D. G., Martens, B., Fisher, J. B., Nobrega, R. L. B., Moreno, A., Cabral, O. M. R., Rodrigues, T. R., Bezerra, B., Silva, C. M. S., Meira Neto, A. A., Moura, M. S. B., Marques, T. V., Campos, S., Nogueira, J. S., Rosolem, R., Souza, R. M. S., Antonino, A. C. D., Holl, D., Galleguillos, M., Perez-Quesada, J. F., Verhoef, A. ORCID: <https://orcid.org/0000-0002-9498-6696>, Kutzbach, L., Lima, J. R. S., Souza, E. S., Gassman, M. I., Perez, C. F., Tonti, N., Posse, G., Rains, D., Oliveira, P. T. S. and Wendland, E. (2021) Are remote sensing evapotranspiration models reliable across South American ecoregions? *Water Resources Research*, 57 (11). e2020WR028752. ISSN 0043-1397 doi: 10.1029/2020WR028752 Available at <https://centaur.reading.ac.uk/101236/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1029/2020WR028752>

Publisher: American Geophysical Union

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Are remote sensing evapotranspiration models reliable across South American ecoregions?

D. C. D. Melo¹, J. A. A. Anache², V. P. Borges¹, D. G. Miralles³, B. Martens³, J. B. Fisher⁴, R. L. B. Nóbrega⁵, A. Moreno⁶, O. M. R. Cabral⁷, T. R. Rodrigues², B. Bezerra^{8,9}, C. M. S. Silva^{9,10}, A. A. Meira Neto¹⁰, M. S. B. Moura¹¹, T. V. Marques⁹, S. Campos⁹, J. S. Nogueira¹², R. Rosolem¹³, R. Souza¹⁴, A. C. D. Antonino¹⁵, D. Holl¹⁶, M. Galleguillos¹⁷, J. F. Perez-Quezada^{17,18}, A. Verhoef¹⁹, L. Kutzbach¹⁶, J. R. S. Lima²⁰, E. S. Souza²¹, M. I. Gassman^{22,23}, C. F. Perez^{22,23}, N. Tonti²², G. Posse²⁴, D. Rains³, and P. T. S. Oliveira², E. Wendland²⁵

¹Federal University of Paraíba, Areia, PB, Brazil

²Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

³Hydro-Climate Extremes Lab (H-CEL), Ghent University, Coupure Links 653, 9000 Ghent, Belgium

⁴Schmid College of Science and Technology, Chapman University, Orange, CA, USA

⁵Imperial College London, Department of Life Sciences, Silwood Park Campus, Buckhurst Road, Ascot, SL5 7PY, UK

⁶Numerical Terradynamic Simulation Group, University of Montana, Missoula, MT, USA

⁷Brazilian Agricultural Research Corporation, Embrapa Meio Ambiente, Jaguariúna, SP, Brazil

⁸Department of Atmospheric and Climate Sciences, Federal University of Rio Grande do Norte, Natal, RN, Brazil

⁹Climate Sciences Graduate Program, Federal University of Rio Grande do Norte, Natal, RN, Brazil

¹⁰Department of Hydrology and Atmospheric Sciences, The University of Arizona

¹¹Brazilian Agricultural Research Corporation – Embrapa Tropical Semi-arid, Petrolina, PE, Brazil

¹²Federal University of Mato Grosso, Cuiabá, MT, Brazil

¹³University of Bristol, BS7 8PD, UK

¹⁴Department of Biological and Agricultural Engineering, Texas A&M University, College Station, TX, USA

¹⁵Department of Nuclear Energy, Federal University of Pernambuco, Recife, PE, Brazil

¹⁶Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany

¹⁷Department of Environmental Science and Renewable Natural Resources, University of Chile, Santiago, Chile

¹⁸Institute of Ecology and Biodiversity, Santiago, Chile

¹⁹Department of Geography and Environmental Science, The University of Reading, Reading, UK

²⁰Federal University of the Agreste of Pernambuco, Garanhuns, PE, Brazil

²¹Federal Rural University of Pernambuco, Serra Talhada, PE, Brazil

²²Department of Atmospheric and Ocean Sciences, FCEN - UBA, Buenos Aires, Argentina

²³National Council for Scientific and Technical Research, (CONICET), Argentina

²⁴Instituto de Clima y Agua. Instituto Nacional de Tecnología Agropecuaria (INTA), Buenos Aires, Argentina

²⁵Department of Hydraulics and Sanitary Engineering, University of São Paulo, São Carlos, SP, Brazil

Key Points:

- Four remote sensing *ET* models were evaluated using data from 25 flux towers in South America
- GLEAM and PT-JPL provided a significantly greater number of daily outputs
- Comparisons with flux tower-based *ET* showed that GLEAM and PT-JPL produced higher correlations whereas *RMSE* was similar for all models
- No model outperformed the other for all biomes, climates or land uses

Corresponding author: Davi Diniz Melo, melo.dcd@gmail.com

Abstract

Many remote sensing-based evapotranspiration (RSBET) algorithms have been proposed in the past decades and evaluated using flux tower data, mainly over North America and Europe. Model evaluation across South America has been done locally or using only a single algorithm at a time. Here, we provide the first evaluation of multiple RSBET models, at a daily scale, across a wide variety of biomes, climate zones, and land uses in South America. We used meteorological data from 25 flux towers to force four RSBET models: Priestley–Taylor Jet Propulsion Laboratory (PT-JPL), Global Land Evaporation Amsterdam Model (GLEAM), Penman–Monteith Mu model (PM-MOD), and Penman–Monteith Nagler model (PM-VI). ET was predicted satisfactorily by all four models, with correlations consistently higher ($R^2 > 0.6$) for GLEAM and PT-JPL, and PM-MOD and PM-VI presenting overall better responses in terms of percent bias ($-10 < PBIAS < 10\%$). As for PM-VI, this outcome is expected, given that the model requires calibration with local data. Model skill seems to be unrelated to land-use but instead presented some dependency on biome and climate, with the models producing the best results for wet to moderately wet environments. Our findings show the suitability of individual models for a number of combinations of land cover types, biomes, and climates. At the same time, no model outperformed the other for all conditions, which emphasizes the need for adapting individual algorithms to take into account intrinsic characteristics of climates and ecosystems in South America.

1 Introduction

Land evaporation, or evapotranspiration (ET), is the phenomenon by which water is converted from a liquid into its vapor phase over land. It plays a significant role in the modulation of global climate feedbacks being a key driver of the Earth’s carbon, energy, and water cycles at local, regional, and global scales (Cao et al., 2010; Tong et al., 2017; Khosa et al., 2019; Valle Júnior et al., 2020; de Oliveira et al., 2021). *In situ* ET measurements can be obtained from micro-meteorological methods (e.g., eddy covariance, scintillometry, or Bowen ratio method) and those derived from the soil water balance (e.g., directly using lysimeters, or from changes in profile soil moisture content obtained gravimetrically, from neutron probes, or capacitance-based soil water monitoring equipment). Besides, plant physiological techniques such as sap flow methods, provide direct estimates of transpiration (Verhoef & Campbell, 2006; Allen et al., 2011; Fisher et al., 2011), but only the micro-meteorological methods provide ET data at the field to landscape (e.g., scintillometry) scale. Over the past three decades, eddy covariance (EC) systems have become the state-of-the-art and standard *in situ* method to quantify land surface energy and mass fluxes for different types of ecosystems (Restrepo-Coupe et al., 2013; Rodrigues et al., 2016; Campos et al., 2019; X. Wang et al., 2020). However, these techniques estimate fluxes for areas of relatively limited spatial dimensions ($\sim 1 \text{ km}^2$) depending on the heterogeneity of the landscape), and they are affected by specific local conditions, such as the occurrence of advection across sharp contrasts in vegetation and/or irrigation conditions, and those caused by topographic features, such as cold air drainage for sloping terrain (Allen et al., 2011; Rwasoka et al., 2011; Mutti et al., 2019; Rahimzadegan & Janani, 2019; Mauder et al., 2020).

During the 1990s and 2000s, remote sensing based ET (RSBET) algorithms, using information from visible, near-infrared, and thermal infrared bands, were developed, such as the Surface Energy Balance Algorithms for Land (SEBAL, Bastiaanssen et al., 1998), Simplified Surface Energy Balance Index (S-SEBI, Roerink et al. (2000)), Surface Balance Energy System (SEBS, Su (2002)), Simplified Surface Energy Balance (SSEB, Senay et al. (2007)), and Two-Source Energy Balance Model (TSEB, Norman et al. (1995); Kustas and Norman (1999)). These algorithms were developed for sub-regional applications, with a focus on irrigation or water resources management. Over South America, their predictive skills have been assessed quite extensively, mostly for irrigated crop-

land (Teixeira et al., 2009; Paiva et al., 2011; Poblete-Echeverría & Ortega-Farias, 2012; Bezerra et al., 2013, 2015; Olivera-Guerra et al., 2017; Lopes et al., 2019; Mutti et al., 2019). Studies show that these models perform well when compared to field observations of *ET* (Teixeira et al., 2009; Poblete-Echeverría & Ortega-Farias, 2012).

Since the late 2000s, algorithms such as PT-JPL (Fisher et al., 2008), PM-MOD (Mu et al., 2007, 2011), and GLEAM (Miralles et al., 2011; Martens et al., 2017) focused on the use of satellite-derived observations to create spatially coherent global *ET* estimates (Fisher et al., 2017). PT-JPL is at the core of the ECOSTRESS mission (Fisher et al., 2020), while PM-MOD is central to the global terrestrial MODIS *ET* product (MOD16). GLEAM is used for the annual State of the Climate report since 2015 (Blunden & Arndt, 2020).

Using flux tower data, previous studies conducted in South America evaluated GLEAM and MOD16 (Ruhoff et al., 2013; Moreira et al., 2019; Paca et al., 2019). However, these studies validated off-the-shelf *ET* datasets generated by these models, not the models themselves. Since such *ET* products are not produced using a common dataset of meteorological variables, a comparative evaluation cannot be made in terms of model structure. Rather, different model skills would be partially linked with the quality of the inputs. A multi-site tropical study, over several continents, validating the PT-JPL model at a regional scale on a monthly basis was presented by Fisher et al. (2009). However, to the best of our knowledge, studies assessing the daily predictive skills have only been conducted at the local scale (Teixeira et al., 2009, 2013; Miranda et al., 2017; B. S. Oliveira et al., 2018; V. d. A. Souza et al., 2019).

A major challenge to verify the results of these methods is the scarcity of ground-based observations, due to the uneven spatio-temporal distribution of the *ET* monitoring efforts. As a result, remote sensing *ET* methods are typically evaluated or parameterized using sites located only in North America, Europe (Ershadi et al., 2014; McCabe et al., 2016; Michel et al., 2016; Xu et al., 2019), Australia (Martens et al., 2016) and East Asia (Jang et al., 2013; Chang et al., 2018; Khan et al., 2018; Li et al., 2019). For example, Mu et al. (2011) proposed improvements to the PM-MOD *ET* global algorithm (Mu et al., 2007), based on comparisons with *ET* measurements from 46 AmeriFlux sites, 45 of them located in USA and Canada. Martens et al. (2017) evaluated the GLEAM algorithm with 91 worldwide FLUXNET sites; however, ~65 were located in the USA and in Europe. Therefore, these models might not satisfactorily represent *ET* in sparsely sampled regions with very different climate conditions such as South America, despite this continent representing ca. 12% of the total Earth’s terrestrial area.

South America spans two hemispheres, and four major climate zones, from near the equator to sub-Antarctic regions, which makes it a geographically unique continent (Goymer, 2017; Trajano, 2019). Biomes in this continent range from tropical to deciduous forests, and contain ecoregions with high sensitivity to variability in water (e.g., the Caatinga and Humid Pampas) and energy availability (e.g. the Amazon, Valdivian temperate and Magellanic subpolar forests) (Seddon et al., 2016). Also, five out of six of the terrestrial biomes not included in satellite-based *ET* algorithm evaluations at a global scale are found in South America (see Section 2.1). Thus, the evaluation of RS-BET methods for South America offers an opportunity to reduce the current research gap, in particular at large spatial scales.

FLUXNET provides a common framework for the verification of *ET* algorithms. Nevertheless, the available sites in the FLUXNET2015 database are not evenly distributed around the world (Pastorello et al., 2020). Validating global models in South America is challenging, mainly because the data from ~90% of its FLUXNET registered sites are not readily available to the scientific community: less than 50% of South American AmeriFlux sites are available for direct access. Additionally, flux towers in woody savannas

and evergreen broad-leaf forests account for nearly 65% of all Latin American FLUXNET sites while some of the biomes are not properly represented (Villarreal & Vargas, 2021).

The identification of scientific gaps and the proposed improvements are considered a priority for the future development of *ET* assessment methods from remote sensing (Fisher et al., 2017). Some of them include merging different *ET*-estimation methods, and the identification of their sources of uncertainty (Fisher et al., 2017; Y. Zhang et al., 2017; Paca et al., 2019). Indeed, despite the recent developments of remote sensing *ET* methods, there are still challenges concerning the refinement of those algorithms to remedy the lack of information on specific surface characteristics and fluxes of undersampled climate zones and vegetation types. In this context, one of the main sources of uncertainty in global satellite-based *ET* estimates are the fractional vegetation cover and net radiation (Ferguson et al., 2010; Vinukollu et al., 2011; Badgley et al., 2015)

We evaluated the predictive skills of four satellite-based *ET* models, designed for regional and continental scale applications, over South America. The main question we seek to answer is whether such models can be applied consistently to reliably capture *ET* in South America. Specific research questions include: (i) are the models capable of correctly estimating *ET* and its components? (ii) are the models predictive skills affected by climate, land cover type or biome?

2 Study area, data, and methods

2.1 South American biomes, flux tower-based *ET* and meteorological data

The study area encompasses five biomes (Table S1 in the Supporting Material – SM): Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Flooded Grasslands & Savannas (FGS); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS); Tropical & Subtropical Dry Broadleaf Forests (TSDBF) and Temperate Broadleaf & Mixed Forests (TBMF) (Olson et al., 2001).

We used daily meteorological data from 25 flux tower sites located across various South American biomes and land cover types to verify the predictive skill of the selected RSBET models (Figure 1a, Table S2 in SM). The time period considered for analysis was determined by the available time-series for each site (Figure S1 in SM). Further information about each biome is provided in SM. Ten sites are from FLUXNET (Pastorello et al., 2020), AmeriFlux networks (Novick et al., 2018) and Large-Scale Biosphere-Atmosphere Experiment in the Amazon (LBA) project (Saleska et al., 2013), while the remaining data were obtained from the respective principal investigators. Concerning towers sites not available in global networks, data handling included standard procedures to ensure quality data, including: detection of spikes caused by changes in the footprint or imprecise measurements; delay correction of H_2O/CO_2 in relation to the vertical wind component; correction of coordinates (2D rotation); correction of spectral loss; conversion of the buoyancy flux to sensible heat flux, known as SND-corrections (Schotanus et al., 1983); sonic virtual temperature correction; corrections for flux density fluctuations, known as WPL corrections (Webb. et al., 1980); incorporated frequency response correction. Additionally, we performed due corrections with respect to reduction of wind velocity or turbulence increase caused by the shadow of the tower and the sensor. Details about procedures carried out for data processing and filtering to implement these corrections can be found in Tonti et al. (2018); Holl et al. (2019); Campos et al. (2019); Cabral et al. (2020). We also emphasize that those data have been widely used in previously scientific publications (Rocha et al., 2009; Cabral et al., 2010, 2011; Restrepo-Coupe et al., 2013; Rodrigues et al., 2016; Arruda et al., 2016; Silva et al., 2017; Marques et al., 2020). The spatial patterns of mean annual precipitation (P), air temperature (T), and potential

evapotranspiration (PET) show that selected sites encompass a wide variety of climates (Figure 1b).

Figure 1. (a) Location of flux tower sites. Land cover types are indicated prior to tower names in the map: Croplands (CROP), Deciduous Needleleaf Forest (DNF), Evergreen Broadleaf Forest (EBF), Grasslands (GRA), Mixed Forest (MF), Permanent Wetland (PW), and Woody Savanna (WS); Biome types (Olson et al., 2001) are indicated by shades of green, yellow and blue on the map (see legend): Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Tropical & Subtropical Dry Broadleaf Forests (TSDBF); Temperate Broadleaf & Mixed Forests (TBMF); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS); Temperate Grasslands, Savannas & Shrublands (TGSS); Flooded Grasslands & Savannas (FGS); Montane Grasslands & Shrublands (MGS); Mediterranean Forests, Woodlands & Scrub (MFWS); Deserts & Xeric Shrublands (DXS); Climates across South America from selected representative sites are indicated by patterns on the map (see legend): Tropical savanna (Aw), Tropical monsoon (Am), Hot semi-arid (BSH), Cold semi-arid (BSk), Humid subtropical (Cfa), Temperate oceanic (Cfb), Dry-winter subtropical highland (Cwb), Polar Tundra (Td) (Peel et al., 2007). (b) Gridded annual average (AVG) and standard deviation (SD) for air temperature (T), rainfall (P), and potential evapotranspiration (PET) across South America and the monitored sites (Harris et al., 2020).

The closure of the energy budget is rarely observed with flux tower measurements (Wilson et al., 2002; Foken, 2008). Usually, the available energy flux ($Rn - G$) is greater than ($LE + H$), where Rn is the net radiation, G is the soil heat flux, LE is the latent heat flux and H is the sensible heat flux. The imbalances in the surface energy balance, reported here as an energy balance ratio, EBR (i.e. $(LE + H)/(Rn - G)$), range from 0.73 to 1.16 (mean ~ 0.90) (Table S2, SM). It is paramount that only high-quality data were used to run and assess the models. We computed daily EBR for each site and excluded days with $EBR \leq 0.75$ or ≥ 1.25 . Daily averages of meteorological variables were calculated from 30-min or hourly data only when at least 80% of the records per day were available. To obtain daytime and nighttime inputs for the MOD16 model (PM-MOD in this paper), we considered only days with a minimum of twenty 30-min daytime records and twenty during the night. As in Mu et al. (2011), the shortwave incoming radiation ($Rgs \downarrow$) was used to distinguish between daytime ($Rgs \downarrow > 10 \text{ W m}^{-2}$) and nighttime ($Rgs \downarrow < 10 \text{ W m}^{-2}$). Regarding the fluxes, we used quality-checked data that had not been gap-filled. Previous studies have shown that ET derived from the other energy balance fluxes, i.e. $LE = Rn - G - H$, can agree well with eddy covariance ET and lysimeter data (Amiro, 2009; Sánchez et al., 2019). Therefore, instead of using the EC-measured LE , to represent ET , we derived LE from the equation above. Such validation approach (i.e. comparing model ET with EB-derived LE , ET_{EB}) has been adopted in previous studies (Twine et al., 2000; Wilson et al., 2002; Stoy et al., 2013; Fisher et al., 2020). The results of using the eddy covariance ET (ET_{EC}) instead can be found in the SM (see Fig. S11-S13).

The quality control procedure described above was not adopted for the TF1, and TF2 towers (see Figure 1a). At those sites, horizontal advection plays an important role due to extreme weather variations throughout the year (Levy et al., 2020), such that the energy balance closure cannot be diagnosed by EBR, as described above. For instance, the SDF zone is known as an anticyclone pathway between the Pacific and Atlantic oceans, and TF1 and TF2 are located in the extreme southern parts of Patagonia, a region characterized by strong winds. Thus, for TF1 and TF2 sites, we used ET derived from measured LE .

2.2 Remote sensing-based vegetation indices

The required vegetation index (VI) to run PT-JPL, PM-MOD and PM-VI is the Enhanced Vegetation Index (*EVI*). Vegetation Optical Depth (*VOD*) is used in GLEAM. *EVI* was derived from the 16-day Level 3 Global product of the MODerate Resolution Imaging Spectroradiometer (MODIS), aboard the Terra and Aqua satellites (Huete et al., 2002). We used both MODIS VI products, i.e. MOD13Q1 (Terra) and MYD13Q1 (Aqua), at 250-m resolution, to derive daily composites of *EVI*. *VOD* was extracted from the product described in Moesinger et al. (2020). Fisher et al. (2008) used the Soil Adjusted Vegetation Index (*SAVI*) instead of *EVI* because the former does not require the blue reflectance (0.45–0.51 μm), however, the authors recognize that both indices are very similar. As we are interested in assessing the *ET* models rather than the products resulting from different forcing data, we used *EVI* in Fisher’s model (PT-JPL). Leaf area index (*LAI*) and other vegetation-related variables (e.g., fraction of Absorbed Photosynthetically Active Radiation, f_{PAR}) are handled differently in each model. For example, in PT-JPL, *LAI* is obtained from total fractional vegetation cover, whereas in PM-MOD the 1-km MODIS *LAI* (MOD15) product is adopted. The original procedures to obtain those variables were not changed here. The following treatment was applied to the MODIS-derived data. “Good quality” pixels were selected, based on the quality assurance (QA) flags. Next, an autoregressive model was applied to fill in the gaps (Akaike, 1969). The gap-filling procedure was applied to gaps smaller than 16 days, while gaps of longer periods were excluded from the analysis. Finally, we implemented a temporal filter to improve the f_{PAR} and *LAI* time series to reproduce precisely all pre-processing steps of the standard PM-MOD algorithm (Mu et al., 2011). Filtering of f_{PAR} and *LAI* allowed for the correction of underestimated values (abrupt and unrealistic decreases in the time series) that mostly originate from cloud contamination effects which were not correctly identified in the quality control fields.

2.3 Summary of remote sensing-based *ET* models

2.3.1 GLEAM

GLEAM is a semi-empirical/process-based model that estimates the total evaporative flux and its components. In this study, version 3 of the algorithm is used (Martens et al., 2017). The main aspects of the model are described briefly, while for details we refer to Miralles et al. (2011) and Martens et al. (2017). The model calculates potential evaporation for four sub-grid land cover fractions: (1) open water, (2) low vegetation, (3) tall vegetation, and (4) bare soil using the Priestley and Taylor (1972) equation. For tall and low vegetation cover fractions, potential transpiration is constrained using an empirical evaporative stress factor which is calculated as a function of soil moisture at root-zone depth and microwave *VOD* as described in Martens et al. (2017). *VOD* (Vegetation Optical Depth) accounts for the attenuation of microwaves through vegetation and can be used as a proxy for vegetation phenology. Thus, *VOD* is a microwave parameter closely linked to vegetation water content (Liu et al., 2013) and in GLEAM it is used to represent phenological changes in vegetation. The soil moisture in the root-zone is calculated with a multi-layer water-balance model forced by precipitation and satellite surface soil moisture retrievals. For bare soil, the evaporative stress factor is calculated as a function of surface soil moisture only, whereas for open water evaporation, no stress factor is applied. For the tall vegetation cover fraction, rainfall interception loss is estimated with Gash’s analytical model (Gash, 1979; Miralles et al., 2010). The *ET* is then calculated as the sum of low and tall vegetation transpiration, rainfall interception loss, bare soil evaporation, and open-water evaporation with each weighted by the respective fraction.

2.3.2 PT-JPL

The global ET model proposed by Fisher et al. (2008) is based on the Priestley and Taylor equation for potential ET (PET), which is partitioned into actual plant transpiration, soil evaporation, and interception evaporation, i.e. $E_{trans} + E_{soil} + E_{int}$. To reduce potential ET to actual ET , the PT-JPL model applies ecophysiological constraints based on land surface information such as vegetation properties and humidity/water vapor pressure deficit (VPD). Fisher et al. (2008) used $NDVI$ and $SAVI$ as a proxy for plant physiological status. We used EVI because it provides a better indication of green vegetation cover than $NDVI$, as acknowledged by Fisher et al. (2008). The model partitions available energy flux using four plant-related constraints: LAI , green canopy fraction, plant temperature, and plant moisture. Similar to PM-MOD (see next subsection), vegetation cover, canopy wetness, etc. determine how the available energy flux is partitioned among the ET terms. A unique aspect related to the plant temperature constraint is the determination of an optimal temperature, T_{opt} (Potter et al., 1993), which corresponds to an optimal stomatal conductance. The latter co-determines E_{trans} .

2.3.3 PM-MOD

The MOD16 ET model (PM-MOD) is based on the Penman-Monteith equation to produce a daily global ET product summing up daytime and nighttime ET (Mu et al., 2011). In this model, total ET is partitioned into E_{soil} , E_{int} , and E_{trans} . To compute E_{soil} , PM-MOD uses potential soil evaporation and a soil moisture constraint function based on VPD and air relative humidity (RH) (Fisher et al., 2008). The evaporation of the water intercepted by the canopy, E_{int} , is calculated using the relevant equations from a revised version of the Biome-BGC model (Thornton, 1998). The PM-MOD assumes that E_{int} occurs when the vegetation is covered with water, i.e. when the water cover fraction (f_{wet}) > 0 , which is constrained by RH (Mu et al., 2011). In the PM-MOD model f_{wet} is calculated as in the PT-JPL model: f_{wet} is set to 0 if $RH < 70\%$ and $f_{wet} = (RH/100)^4$ if $70 < RH < 100\%$ (Running et al., 2019). The PM-MOD model is designed to allow E_{trans} to occur during daytime and nighttime, by adding constraints to stomatal conductance for VPD and minimum air temperature, and ignoring constraints relating to high air temperature (Running et al., 2019). The partitioning of available energy flux into soil or interception evaporation is based on vegetation cover (Fc), which is assumed to be equal to f_{PAR} from the MODIS product MOD15A2 (Mu et al., 2011). Although this method is based on the PM equation, PM-MOD does neither require wind speed nor soil moisture data for the parameterization of aerodynamic and surface resistance. Further details about PM-MOD can be found in Mu et al. (2011) and Running et al. (2019). Note that some updates have been implemented in PM-MOD since Mu et al. (2011), which can be found in Running et al. (2019). These were also considered here in the implementation of PM-MOD.

2.3.4 PM-VI

This model relies upon the hypothesis that ET is mostly controlled by specific dominant processes, such as transpiration and photosynthesis, hence a good correlation between such processes and ET is necessary for good model performance (Nagler et al., 2007). There are several formulations to estimate ET from VIs (Nagler et al., 2005, 2009). In this study, we selected the algorithm proposed by Nagler et al. (2013), which estimates ET using the reference crop evapotranspiration, ET_0 , from the FAO-56 Penman-Monteith (PM) equation (Allen et al., 1998), and a crop coefficient, K_{cVI} , derived from a vegetation index. K_{cVI} can be calculated in different ways (Nagler et al., 2005, 2013). Following Nouri et al. (2016) and P. T. S. Oliveira et al. (2015), K_{cVI} was calculated as:

$$K_{cVI} = a(1 - e^{-b \times EVI}) - c \quad (1)$$

where a , b and c are fitted coefficients. We used a parameter optimization tool based on a genetic algorithm to optimise the coefficients to estimate ET values close to the measured ones (P. T. S. Oliveira et al., 2015). The fitting procedure minimizes the objective function (OF) given by the sum of squared differences between tower-based ET (ET_{obs}) and ET estimates from the models (ET_{sim}) at time i :

$$OF = \sum_{i=1}^n [ET_{obs}(i) - ET_{sim}]^2 \quad (2)$$

This model, herein referred to as PM-VI, has frequently been employed to estimate ET at local and regional scales (P. T. S. Oliveira et al., 2015; Nouri et al., 2016; Jarchow et al., 2017). Although obtaining ET_o requires a considerable amount of meteorological variables, the PM-VI implementation is easier and has a lower computational cost compared to other models. Unlike the three other models, PM-VI requires the calibration of the fitting coefficients, which can be a major issue for regions where ET and VI are poorly correlated or when correlations change over time (Chong et al., 1993). To calibrate the fitting coefficients, we randomly selected 20% of the available data at each site and used the remaining 80% to validate the model.

2.4 Quantifying model reliability

The model predictive skill was visually evaluated with scatter plots of measured versus modelled ET , as well as through the coefficient of determination (R^2), root mean square error ($RMSE$), percent bias ($PBIAS$), concordance correlation coefficient (ρ), slope (m), and intercept (b) of the linear regression. The data used in the analysis were filtered for rainy days ($P > 0.5$ mm). Our analysis proceeded from a general (no distinction among sites) to a site-by-site and group level analysis, i.e. per biome, climate, or land use. The number of flux towers assigned to each subgroup (i.e., the different biomes, climate, and land use classes) varied, and so did the record length per subgroup. To account for the different sizes, the following sampling procedure was performed, in which we computed the variability of each performance metrics for each group analysis (i.e., across its different subgroups): (i) A sample size N was defined as half of the record length of the shortest subgroup, among all models; (ii) for each model, samples of length N were taken from within each subgroup, and the performance metrics were computed; (iii) This procedure was repeated 1000 times, yielding a mean and standard deviation (SD) of the metrics at each subgroup, per model. The resulting SD are likely to be influenced by the choice of N , and other rationale for its choice could have been made. In this way the confidence bands reported here are to be seen as measures of relative variability, i.e., the variability between the models, and not as absolute uncertainty bounds for each of them. To establish a relationship between model predictive skill and water availability at individual tower sites, we obtained the aridity index ($AI = P/ET_o$) from the global dataset provided by Trabucco and Zomer (2019). For many tower sites, the available meteorological data (even from nearby meteorological stations) were not sufficient to provide a reliable AI ; hence the choice for a global dataset.

3 Results

3.1 ET partitioning

Partitioning of ET among the three components (E_{trans} , E_{int} and E_{soil}) exhibited more variation for the PT-JPL and PM-MOD models. On average, E_{trans} accounted for 60% (PT-JPL) and 56% (PM-MOD) of ET but, across sites, it presented a smaller range (30% to 85%) for PT-JPL than for PM-MOD (20 to 90%) (Figure 2, Table 1). GLEAM E_{trans} accounted for 82% of ET on average, varying between 60% and 95% across sites. Average interception across sites reached 9% (GLEAM), 13% (PT-JPL), and 24% (PM-

MOD) of total ET . E_{int} fractions range were similar for GLEAM and PT-JPL ($SD \approx 9\%$), whereas PM-MOD E_{int} varied more among sites ($SD = 18\%$). E_{int} was often correlated with LAI , especially for the GLEAM estimates ($R^2 = 0.57$, Figure S2 in SM). PT-JPL E_{soil} estimates exceeded the other models, particularly for sites with low LAI values (e.g., ESEC, CST, and USR).

Figure 2. Evaporation fractions estimated by the models at each site (stacked bars) and average partitioning of land evaporation per model (pie diagram). Black dots: LAI scaled between 0 and 1 based on the minimum and maximum values of LAI (from MODIS MDC15A2 product). Red \times : the concordance correlation coefficient between observed and simulated daily ET.

3.2 Overall model skills

Since each model requires a different input dataset (Table S3, SM), the data available to run and validate each model varied. GLEAM and PT-JPL provided a significantly greater number of daily outputs: 7301 (GLEAM), 7277 (PT-JPL), 5905 (PM-MOD), and 6638 (PM-VI). The complete data set was used to produce scatter plots of ET records and model simulations for each location (See Figures S4-S7 in SM). To allow a fair analysis, the results shown in the main text were obtained using data from days that were common across models, resulting in 4718 data points.

To illustrate the relative contribution of each site to the scatter plots in Figure 3, we display the regression lines (light grey lines) between model and tower-based ET for each tower site, and the mean metrics across individual sites. In general, ET was reasonably predicted by all models, as suggested by the relatively low spread of most points in the scatter plots, many regression lines close to the 1:1 line, mean determination coefficient, \bar{R}^2 , mean concordance correlation coefficient, $\bar{\rho}$, mostly above 0.65, and mean root mean square error (\bar{RMSE}) below 1 mm d^{-1} (Figure 3). Nevertheless there is some spread for a few sites, e.g., in the PT-JPL scatter plot that displays a few sites with large bias despite strong overall correlation and ρ .

The models slightly overestimate ET as suggested by higher density of points below the 1:1 line, except for GLEAM, which slightly underestimates. Correlations were similar between GLEAM and PT-JPL, with an average value of ~ 0.65 and the highest values at individual sites reaching close to 0.9, as indicated by the standard deviations (0.19 and 0.18, respectively). From Figures 3 and 4, it becomes evident that, despite the relatively lower spread of points for PM-VI, compared to the other models, this model (PM-VI) presented a less consistent performance across towers, as suggested by the contrasting slopes presented by the regression lines in that plot (e.g. reversed trend line at K77); hence the lower average determination coefficient (\bar{R}^2) and $\bar{\rho}$. Such contrasting aspect of the PM-VI model is also noted by the fact that a wide range of R^2 was found despite the similarity between mean simulated and observed ET (Figure 4).

Figure 3. Scatter plots of observed vs. simulated daily evapotranspiration at all flux tower sites, for each model. The light grey lines show the regression slope of individual sites. The coefficient of determination (R^2), root mean square error ($RMSE$) and concordance correlation coefficient (ρ) were averaged across towers and are displayed on the plots ($N = 4,718$).

Table 1. Comparison of evaporation fractions for several land uses between this study and field-based estimates. FE = field estimates. Land covers that present field data from the same modeling sites or same geographical region are indicated with “*”.

LULC	E_{trans} (%)				E_{soil} (%)				E_{int} (%)				References
	FE	GLEAM	PT-JPL	PM-MOD	FE	GLEAM	PT-JPL	PM-MOD	FE	GLEAM	PT-JPL	PM-MOD	
EBF*	80-84	74-79	47-63	31-88	NA	4-6	18-24	5-20	15-25	17-20	18-29	7-58	Leopoldo et al. (1995); Shuttleworth and Pereira (1988)
DNF*	50-81	84-94	64-84	78-90	NA	8-14	14-24	8-18	10	1-2	2-5	2-4	Gaj et al. (2016); Sun et al. (2019); de Queiroz et al. (2020)
CROP*	NA	93	63	70	20-4	6	31	21	10	1	6	9	Dennead et al. (1997); Cabral et al. (2012)
CROP*	85	88	55	69	NA	3	20	4	13	9	25	27	Cabral et al. (2010)
WS*	NA	86	76	78	NA	2	17	5	8	13	7	17	Cabral et al. (2015)
GRA	50-78	69-73	47-49	33-54	NA	25-30	32-46	30-52	NA	1-2	7-18	15-16	Ferretti et al. (2003); Sutanto et al. (2012); Z. Wang et al. (2014)
MF	36-74	82-88	63-75	58-71	19	11-16	23-30	28-29	NA	1	2-7	1-14	(Aron et al., 2020; Paul-Limoges et al., 2020)
PW	33-38	73-86	28-57	34-41	NA	0-20	32-57	16-54	NA	3-14	6-16	21-43	J. Zhang et al. (2018)

Figure 4. Comparison of mean observed and simulated ET . Circle colors vary according to individual model R^2 .

Figure 5. Model performance per biome, land use and climate. The error bars represent the standard deviation of the metrics within each class. Biome types: Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Tropical & Subtropical Dry Broadleaf Forests (TSDBF); Temperate Grasslands, Savannas & Shrublands (TGSS); Temperate Broadleaf & Mixed Forests (TBMF); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS); Flooded Grasslands & Savannas (FGS); Land use types: Cropland (CROP); Woodland Savanna (WS); Deciduous Needleleaf Forest (DNF); Evergreen Broadleaf Forest (EBF); Grasslands (GRA); Mixed Forest (MF); Permanent Wetland (PW); Deciduous Broadleaf Forest (DBF). Climate Zones: Tropical monsoon (Am); Tropical savanna (Aw); Hot semi-arid (BSh); Cold semi-arid (BSk); Temperate oceanic (Cfb); Dry-winter subtropical highland (Cwb); Polar Tundra (Td).

3.3 Model skills per biome, land use, and climate

Figure 5 presents ρ , $RMSE$, $PBIAS$, and R^2 for each model across six biomes, eight land use types, and seven climate classes in South America. Error bars are shown for all metrics, and they represent the standard deviation resulting from the resampling procedure outlined in 2.4. Note that the analysis about the FGS and TBMF biomes are based on one and three towers, respectively. For most biomes, $RMSE$ and R^2 did not significantly diverge. In general, TSGSS showed the best overall metrics for all models, while PM-VI in FGS (NPW site) presented the poorest ($\rho < 0.5$, $RMSE > 1.5 \text{ mm d}^{-1}$, and $R^2 < 0.25$). Model performance across towers within each biome did not vary much, as suggested by the relatively low range of the error bars for all metrics.

The central panels in Figure 5 provide evidence for the high variability of model predictive skills across different land uses (LU), which suggest that: (i) no model outperforms the others for all LU types, (ii) each model has intrinsic and in some cases exclusive characteristic that makes it more suitable for certain LU. Only for croplands (CROP) we found similar metrics among models ($\rho \approx 0.8$, $0.8 < RMSE < 1.2 \text{ mm d}^{-1}$, $-20\% < PBIAS < 10\%$, $0.6 < R^2 < 0.8$). Conversely, for most LU, the metrics variation is remarkable (e.g., DBF: $0.4 < \rho < 0.9$, $-50\% < PBIAS < 10\%$, $0.25 < R^2 < 0.80$). On average, each model has the best skills for two LU; e.g., ET prediction for GRA and DBF was best with PT-JPL ($\rho \approx 0.9$, $RMSE \approx 0.5 \text{ mm d}^{-1}$, $PBIAS \approx 0\%$, $R^2 > 0.75$) whereas PM-VI presented similar skills for estimation of ET for CROP and PW. Likewise, model skill is related to the climate type. The analysis of ρ and R^2 over semi-arid regions (BSk and BSh) indicates a relatively poor skill of all models (except PM-MOD for BSh climate). This is in contrast to the overall good performance over more humid environments (e.g., Aw and Cwb). The greatest divergence among model performances was found for the Polar Tundra (Td) climate zone, for which PM-VI presented the highest ρ and R^2 (both > 0.75), lowest $RMSE$ ($\sim 0.5 \text{ mm d}^{-1}$) and $PBIAS$ ($< 10\%$).

3.4 Individual sites

In this section, we explore the model performance at individual towers. Model skills for all individual sites are depicted in Figure 6. Sites with $N < 30$ (CAX and MCR) are not discussed here but are considered in the scatter plots shown in the SM (Figures S4-S7). To facilitate the comparison of our results with previous analyses using the same models, only three statistics are shown in Figure 6: $RMSE$, $PBIAS$, and R^2 . Other met-

rics are displayed in the scatter plots in Figures S4-S7 in the SM. In Figure 6, the metrics for the various towers are displayed in order of increasing aridity (varying from ~ 3 to 0, left to right), as suggested by the AI as described in Section 2.4). In general, there is a good agreement between the PM-based models in terms of $RMSE$ and $PBIAS$.

In terms of individual metrics, $RMSE$ values varied between ~ 0.5 and $\sim 1.5 \text{ mm d}^{-1}$ for all models, with $RMSE < 1 \text{ mm d}^{-1}$ for most sites. The boxplots show that $RMSE$ variation is similar among models, except for PT-JPL which presents the lowest $RMSE$ (e.g., K67). Figure 6 shows that $PBIAS$ for PM-VI varies around zero across sites, which is expected given the model requires calibration with local data. However, based on R^2 , it is apparent that this model's skill is quite limited for $AI > \sim 1.2$ and $AI < \sim 0.5$. In general, the PT-based models showed larger biases, with PT-JPL and GLEAM consistently overestimating and underestimating ET , respectively. In terms of R^2 , the PT-models ranked better than the PM-models for more than $\sim 50\%$ of the towers.

Figure 6. Comparison of statistics of the models in estimating evapotranspiration (ET) for the various flux towers used. (a) Sample size (N) used to compute the statistics; (b) $RMSE$ = Root Mean Square Error; (c) Percent Bias ($PBIAS$); (d) R^2 = coefficient of determination. A summary of each model's statistics is depicted in the boxplots: (e) $RMSE$; (f) $PBIAS$; (g) R^2 . Flux towers are arranged according to the aridity index (with aridity increasing from left to right). Sites with $N < 30$ (CAX and MCR) are not shown here

4 Discussion

4.1 General implications

We conducted the first multi-remote sensing ET model analysis in South America (SA) using a common set of forcing and validation data located on flux tower sites across a diverse range of land covers, climates, and biomes. Forcing data include both *in situ* (e.g., temperature and net radiation) and remote sensing data, mainly related to vegetation (e.g., LAI and EVI). To evaluate the models, energy balance-derived ET (ET_{EB}) was used as observation, instead of eddy covariance ET (ET_{EC}). Given the benefits and drawbacks of using either ET_{EB} or ET_{EC} , we compared both measures to verify whether such choices would lead to different results. As shown in the SM, for the great majority of tower sites, ET_{EC} and ET_{EB} are similar (Figure S10) and model statistics (R^2 , $RMSE$ and ρ) remained the same regardless of the ET approach or indicate a better performance when ET_{EB} was used (Figures S11-S13 in the SM). Many of the tower sites considered here are not yet available in flux network databases, including sites with land cover (deciduous needle-leaf forests, DNF), a biome (FGS), and two climate types (polar tundra, hot semi-arid) that have not been previously assessed in other regional studies on the performance of satellite-based ET models. Moreover, some classes included here were considered for validation of individual models only (e.g., semi-arid and tropical climate types, and TSDBF biome).

The fulfillment of such gaps (i.e. model evaluation across uncharted regions) is an important step because it allows a multitude of applications and studies relying on large scale ET mapping, such as: drought monitoring (Anderson et al., 2011, 2016), agricultural water management (Anderson et al., 2012), diagnosis of climate change (Mao et al., 2015). The current ability to map ET remotely at various spatial and temporal scales, could only be evaluated thanks to the vast number of eddy covariance towers available in continental and global flux networks. As shown in this study, a thorough assessment of RBSET models based solely on data from such networks would be challenging or insufficient for some regions or continents; hence the relevance of this study. Our analy-

sis provides essential information to identify model strengths and limitations across SA, allowing the users to identify which model is more suitable for them. Knowing under what circumstances, e.g., land use or climate, each model is more reliable is necessary to address remaining research and applied science gaps relative to *ET* at local, regional and global scales (Fisher et al., 2017). Despite the value of tower-based *ET* across SA, many of those questions persist due to our limited observational capabilities. According to Fisher et al. (2017), the way to begin answering those questions is producing high quality *ET* estimates, which includes acquiring accurate *ET* information at high temporal and spatial resolution with large spatial coverage for a sufficient long period.

4.2 Model performance, sources of errors and ET partitioning

Generally, model predictive skill over SA resembles what has been reported for other continents, including satisfactory values of coefficient of determination ($R^2 > 0.6$) of the models (except PM-VI) for most validation sites, and consistently better results for the GLEAM and PT-JPL models, with *RMSE* ranging from ~ 0.5 to 1.5 mm d^{-1} (McCabe et al., 2016). Also, in accordance with previous analysis, GLEAM and PT-JPL presented somewhat higher *RMSE* than PM-MOD but no clear evidence indicates decreasing performance with increasing aridity, as reported by McCabe et al. (2016); Michel et al. (2016). Nonetheless, the general analysis (Section 3.2) indicates that all models can be used reliably over most of the environmental conditions in SA covered in our study. The analysis across towers and groups (i.e., biome, land use type and climate, Section 3.3, Figure 5) identified considerable differences in terms of model skill.

Our results agree with previous studies from (Ershadi et al., 2014; McCabe et al., 2016; Michel et al., 2016; Miralles et al., 2016) who applied PM-MOD, GLEAM (except Ershadi et al. (2014)) and PT-JPL to sites located in Africa, Asia, Australia, Europe and Middle East and reported that PM-MOD showed, for most sites, lower correlations with measured *ET* compared to GLEAM and PT-JPL. Unlike previous analysis, our study agrees with Michel et al. (2016) in the sense that model skill seems to be unrelated to land cover. Michel et al. (2016) also reported a wide variation of R^2 (0.2–0.8) and *RMSE* (0.8–2 mm d^{-1}), for different sites under mixed forests. Conversely, contrasting results between our results and previous studies were found for woodland savanna. While we found $0.5 < R^2 < 0.8$ and $0.7 < RMSE < 1.5 \text{ mm d}^{-1}$, Michel et al. (2016) reported $R^2 < 0.2$ and $1 < RMSE < 3 \text{ mm d}^{-1}$.

Overall, our group-wise analysis based on climate agrees with previous studies. For example, the poor model skill found here for the cold semi-arid (Bsk) climate ($0.1 < R^2 < 0.5$) resembles that found by Michel et al. (2016) and McCabe et al. (2016) for several sites in the United States. While aridity could have played a role here, it could also be caused by the fact that semi-arid sites are covered with sparse canopies. Such canopies present challenges when it comes to the description of aerodynamic transfer for example and radiation partitioning (see e.g. Verhoef and Allen (2000)). Our findings also show a poor to moderate model skill for *ET* predictions for sites located in the Cfb climate zone, with PM-MOD having the worst performance. Conversely, PM-MOD presented the best predictive skill for the BSh climate, according to most metrics.

Besides the three RSBET models commonly assessed (GLEAM, PT-JPL, and PM-MOD), our analysis included the PM-VI model, which has been validated mostly for cropland or riparian ecosystems (Nagler et al., 2005, 2009, 2013; Jarchow et al., 2017). Here, we tested PM-VI for a much wider variety of biomes, climates and land uses, and found a poor predictive skill for several sites with $AI > 1.2$ (e.g. K67, K77, K83) or $AI < 0.5$ (e.g., CAA and SLU), even though the model accounts for a site-specific calibration. Considering the good results obtained for $\sim 50\%$ of the towers and the fact that, compared to the other models, PM-VI has a much simpler implementation, this model does have potential as long as sufficient data are available for calibration or, at least, valida-

tion. However, the need for local calibration is a hurdle for its implementation for most regions that are unsampled; therefore future studies are necessary to investigate which factors are most relevant in the determination of the model fitting coefficients, and to provide distributed reference values for its coefficients (e.g., based on land use dynamics).

We were able to identify a number of probable causes for poor model performance at individual sites, including (i) patch-scale heterogeneities; (ii) “mixed pixels”, i.e. mixed response of different vegetation types within a pixel; (iii) time-lag between ET_{obs} and EVI ; (iv) model sensitivity to individual inputs; (v) low correlation between ET and vegetation indices (see Section 3.0 in the SM for more details). Although we did not verify this in our study, we did not dismiss the possibility that known uncertainties in the estimation of site-specific vegetation characteristics (e.g., f_{PAR} and leaf stomatal conductance in the PM-MOD; Ershadi et al. (2014)) are further causes of lower model performance.

In our study, we used soil heat flux (G) which is generally measured below ground (usually at 5–20 cm deep) using soil heat flux plates. It could be argued that not correcting G for the heat storage between the plate and the soil surface could lead to sub-optimal estimates of ET when LE is calculated as the residual of the energy balance, especially for towers where the soil is bare or covered by sparse vegetation, where G can be relatively large. This, in turn, could lead to the conclusion that the models are performing worse than is actually the case. Although desirable, correcting G for heat storage is rarely possible due to data unavailability (few sites only measure soil moisture and temperature, which are required to estimate soil heat capacity, and heat storage using the calorimetric method). Moreover, at daily scales and for most sites, G is either negligible in SA (summer or winter, when the amount of heat stored during the day roughly equals that lost during the night) or represents a minor portion only (spring and autumn) of the energy balance. As detailed and discussed in Section S3.0 and Figure S7 in SM, it is highly unlikely that neglecting such corrections will have affected the results.

There are, however, some issues worth mentioning here. Cause (v), for instance, is a major issue for PM-VI, as expected because the model is highly dependent on VI dynamics (see Section 2.4) (Nagler et al., 2005). Regarding cause (iv), the superior performance of the PT models over PM-MOD at most sites is probably linked to uncertainties resulting from the estimation of aerodynamic resistance (Ershadi et al., 2014). In PM-MOD, the aerodynamic and surface resistances of each ET component (soil, interception, and transpiration) are parametrized based on biome-specific values of leaf-scale boundary layer conductance, for example (Mu et al., 2011). Compared to the previous version of PM-MOD (Mu et al., 2007), this new approach resulted in a perceptible improvement only for cropland and deciduous broadleaf forest flux tower sites, whereas for other land uses no meaningful change was reported (Ershadi et al., 2015). Conversely, PT models are highly dependent on Rn (causes iv and v); hence they often fail in dry environments (see metrics for $AI < \sim 0.6$ in Figure 6) where ET seasonality is dictated by P more than radiation, or in regions with low Rn (e.g., TF2). Poor model responses at K77 (cropland, Figure S9 in SM) were attributed to causes (i) and (ii), as remnants of forest and shrubs were identified within the tower footprint and within MODIS pixel. VI products with higher resolution than MODIS exist and have been used to estimate ET (Aragon et al., 2018; Fisher et al., 2020); thus offering a possible solution for causes (i) and (ii). Time lag between ET and EVI (cause iii) was identified at EUC, where EVI followed the decline of ET after ~ 1 – 2 months.

Regardless of all those potential causes for poor model response, it is also important to consider the role of the core formulation upon which those RBSET models are based, i.e. Penman-Monteith (PM) and Priestley and Taylor (PT) equations. A major problem of the PM equation refers to the linearization of the Clausius-Clapeyron relation, which has been addressed in a new version of that equation (McColl, 2020). The PT equa-

tion, in turn, implicitly assumes Rn and surface temperature (T_s) to be independent of evaporation. In reality, as shown by Yang and Roderick (2019), Rn not only decreases with increasing T_s (due to an increase of outgoing longwave radiation) but also a greater fraction of Rn becomes available for evaporation. Some of the deviations from the observations found in our analysis may happen due to such inconsistencies or simplifications. Here, we provide evidence to consider revisiting not just parameter values but the governing equations themselves and, ultimately, evaluate the benefits of such potential improvements in RBSET models.

Remote sensing based ET partitioning is expected to present some divergences from ground based measurements. This is the case especially for E_{soil} , because of the difficulty in obtaining remote sensing information on soil characteristics that drive E_{soil} , such as soil moisture and temperature (Talsma, Good, Jimenez, et al., 2018; Talsma, Good, Miralles, et al., 2018), in particular at high vegetation cover fractions. Globally, transpiration has been reported to account for 57–90% of global ET , based on *in situ* data and model outputs (Jasechko et al., 2013; Wei et al., 2017; Paschalis et al., 2018). Although these are global estimates, we expected E_{trans} to be the largest ET component also in SA due to its prevailing tropical climate and corresponding vegetation types. Our results show that this was indeed the case for GLEAM with an E_{trans}/ET ratio of $\sim 80\%$, and for PT-JPL and PM-MOD with values of 57 and 60%, respectively. Nonetheless, based on our findings, model predictive skill in estimating total ET is not necessarily associated with its ability to partition ET accurately.

Concomitantly, inconsistencies in ET partitioning do not necessarily translate into inaccurate model estimates of total ET : this depends on the modelling approach. On the one hand, if total ET results from the sum of ET components independently, then an under- or overestimation of ET components can reduce the overall model skill, or reasonable ET estimates can be achieved as the consequence of an occasional compensation of errors in E_{trans} , E_{soil} and E_{int} . On the other hand, if the ET partitioning is derived from the estimate of a proxy value for total ET , such as available energy flux (as in PM-MOD and PT-JPL), the ET partitioning is unlikely to influence the total ET estimates. Moreover, ET partitioning may be sensitive to certain model inputs. For example, contrasting ET fractions were estimated by PM-MOD for similar rain forest sites, i.e. K67 and K83 (Figure 2). The reason PM-MOD is returning that difference is because RH was estimated from actual (e_a) and saturation vapor pressure (e_s) data, as RH data is not available in K83 dataset. As a result, the difference between e_a -derived daytime and nighttime RH for K83 is greater than that for K67. In terms of daily averages, e_a -derived RH and measured RH are quite similar, which explains why the fractions for GLEAM and PT-JPL, at those two towers, are similar. Still, good estimates of ET components are important to differentiate the roles of vegetation and soil, i.e., how they contribute to vertical soil water fluxes and changes in profile soil water content. Reliable knowledge of the distribution between E_{soil} and E_{trans} is also important when this information is used in hydrological models to calculate other water balance components, such as runoff.

Ground-based ET partitioning data are generally not widely available. This also goes for most land cover types included in this study. We compared the models' outputs with field experiment studies that measured one or more ET components either at the same sites as those used here or within the same region (Table 1). ET partitioning values derived from GLEAM seem to be more consistent with ground-based information available for tropical rain forests, croplands and grasslands than for wetlands, and mixed and deciduous needle-leaf forests (Table 1). This also applies to PT-JPL with its ET partitioning agreeing reasonably well with observations made for both tropical rain- and dry forests. Note that PT-JPL (as well as PM-MOD) constrain E_{trans} based on f_{wet} . Hence, compared to GLEAM, transpiration will be lower under high RH in the model but ET can be high due to water availability in the soil and intercepted rainfall. Nonetheless,

the overall predictive skill of PT-JPL was satisfactory at such sites (Figure 6 and Figure S4 in SM). Regarding PM-MOD, the main inconsistency is the E_{inter} for tropical forests (Table 1). Despite the wide variability in E_{trans}/ET among models, their overall predictive skill was satisfactory, that is, not associated with their capability to correctly estimate each ET component individually (see SM for further discussion). No model was able to consistently capture the ET partitioning across all sites correctly, which is expected given the uncertainty of each ET component and the climate and land-cover variability in SA. However, the joint estimates of all models covered totally or partially all field-derived evidence on ET partitioning. This suggests that continental ET estimates for understudied regions, such as the SA, would benefit from merging ET outputs from models that are based on different methods (Paca et al., 2019).

Despite our efforts to gather as much tower data as possible, with the goal of having a common data set for all models, we faced several limitations including: differences in lengths of observational time series across towers (up to 3 years), as well as lack in overlap of these time series; uneven distribution of towers across groups (e.g., biomes); and, finally, South American geographical features that were not considered in this study (e.g., MGS biome or desert climate type, BWk). Thus, it was not possible to assess, for all towers, model responses during all seasons. Nonetheless, the fact that our dataset encompasses a wide variety of climates enabled us to evaluate, to a reasonable extent, model responses for contrasting seasons and fill in the gaps flagged by the literature, such as the absence, in a similar analysis, of towers in the tropical climate zone pointed out by McCabe et al. (2016).

5 Conclusion

Our results show that, in general, ET can be reasonably well predicted by all four models, despite an overall tendency of overestimation by PT-JPL and PM-MOD, and underestimation by GLEAM. Contrasting with results from other continents, we found no clear evidence linking model predictive skill with aridity. Our analysis emphasizes the need of improving model ET partitioning, although the link between flawed ET partitioning and poor model skill is not evident based on our results. Having reliable ET partitioning coefficients as part of the FLUXNET-type datasets would be valuable in this respect, but unfortunately such data are difficult to obtain, as they require labour-intensive and expensive methods (such as sapflow gauges and lysimeters), that also present problems with regards to upscaling from plot to field-scale.

Correlations are consistently higher for GLEAM and PT-JPL, with $R^2 > 0.5$ for most sites, whereas PM-MOD and PM-VI presented better performances in terms of PBIAS ($-10 < PBIAS < 10\%$ for most sites). As for PM-VI, the low PBIAS is expected, given the model requires calibration with local data.

The model skill for the various models seems to be unrelated to land cover type as we found a wide variability of metric values within the same class and across models. Conversely, a relatively lower performance was observed for most models in semi-arid regions, compared to an overall good performance for more humid environments. Except for the FGS biome, we found that skill across models was mostly similar within the same biome.

Despite the relatively high number of towers (compared to previous global analyses that used a similar amount of sites), gathering a balanced amount of data and uniform distribution of towers across different biomes and climate zones across the whole continent was challenging. Thus, there is a need to expand the flux tower network in South America as well as the formation of bilateral collaboration for future contributions. Previous studies (Michel et al., 2016; McCabe et al., 2016) have expressed the need to extend the evaluation of RSBE models to uncharted biomes and climate conditions. Our

analysis fills this gap by assessing the reliability of four RSBET models over South America. We provide benchmarking metrics that can serve the improvement of *ET* models for improved capturing of *ET* over this continent.

Acknowledgments

The data used in this study will be available through a data-sharing repository. Funding for AmeriFlux data resources was provided by the U.S. Department of Energy's Office of Science. Davi de C. D. Melo was supported by the São Paulo State Research Foundation (FAPESP) (grant 2016/23546-7), and by the Brazilian National Council for Scientific and Technological Development (CNPq) (project 409093/2018-1). Jamil A. A. Anache was supported by the Brazilian National Council for Scientific and Technological Development (CNPq) (project 150057/2018-0). Edson Wendland was supported by the São Paulo State Research Foundation (FAPESP) (grant 2015/03806-1). Paulo Tarso S. Oliveira was supported by the Brazilian National Council for Scientific and Technological Development (CNPq) (grants 441289/2017-7 and 306830/2017-5) and the CAPES Print program. Rafael Rosolem would like to acknowledge the Brazilian Experimental datasets for Multi-Scale interactions in the critical zone under Extreme Drought (BEMUSED) project [grant number NE/R004897/1] funded by the Natural Environment Research Council (NERC). Alvaro Moreno was financially supported by the NASA Earth Observing System MODIS project (grant NNX08AG87A) and the European Research Council (ERC) funding under the ERC Consolidator Grant 2014 SEDAL (Statistical Learning for Earth Observation Data Analysis, European Union) project under Grant Agreement 647423. Diego G. Miralles, Brecht Martens and Dominik Rains are supported by the European Research Council (ERC) DRY-2-DRY project (grant no. 715254) and the Belgian Science Policy Office (BELSPO) STEREO III ALBERI (grant no. SR/00/373) and ET-SENSE (grant. no SR/02/377) projects. Thiago R. Rodrigues was supported by the Brazilian National Council for Scientific and Technological Development (CNPq) with Bolsa de Produtividade em Pesquisa - PQ (Grant Number 308844/2018-1). Jorge Perez-Quezada and Mauricio Galleguillos were supported by the Chilean National Agency for Research and Development, grant FONDECYT 1211652. Rodolfo Nobrega and Anne Verhoef acknowledge support by the Newton/NERC/FAPESP Nordeste project (NE/N012526/1 ICL and NE/N012488/1 UoR). Gabriela Posse acknowledges support by AERN 3632 and PNNAT 1128023 INTA Projects. JBF was supported in part by NASA: ECOSTRESS and SUSMAP. Funding for site support:

- NPW tower: Brazilian National Institute for Science and Technology in Wetlands (INCT-INAU), Federal University of Mato Grosso (UFMT - PGFA and PGAT), University of Cuiabá (UNIC) and SESC-Pantanal;
- SDF tower: funded by the National Commission for Scientific and Technological Research (CONICYT, Chile) through grants FONDEQUIP AIC-37 and AFB170008 from the Associative Research Program;
- TF1 and TF2 towers: funded by the Deutsche Forschungsgemeinschaft (DFG) under Germany's Excellence Strategy - EXC 177 'CliSAP - Integrated Climate System Analysis and Prediction' - contributing to the Center for Earth System Research and Sustainability (CEN) of Universität Hamburg and by DFG project KU 1418/6-1;
- MCR and BAL towers: funded by the National Council for Scientific and Technological Research (CONICET, Argentina) grants PIP-11220100100044 and PIP-11220130100347CO, and by the National Agency for the Scientific and Technological Promotion (ANPCyT, Argentina) grant PICT 2010-0554;
- CAA, CST, and ESEC Towers: funded by National Observatory of Water and Carbon Dynamics in the Caatinga Biome (INCT-NOWCDCB), Federal University of Pernambuco (UFPE), FACEPE (Pernambuco State Research and Technology Foundation) through the Project Caatinga-FLUX APQ 0062-1.07/15.

References

- Akaike, H. (1969, December). Fitting autoregressive models for prediction. *Annals of the Institute of Statistical Mathematics*, 21(1), 243–247. Retrieved 2020-06-22, from <https://doi.org/10.1007/BF02532251> doi: 10.1007/BF02532251
- Allen, R. G., Pereira, L. S., Howell, T. A., & Jensen, M. E. (2011, April). Evapotranspiration information reporting: I. Factors governing measurement accuracy. *Agricultural Water Management*, 98(6), 899–920. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0378377411000023> doi: 10.1016/j.agwat.2010.12.015
- Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). *Crop evapotranspiration - Guidelines for computing crop water requirements - FAO Irrigation and drainage paper 56*. Rome: FAO - Food and Agriculture Organization of the United Nations. Retrieved 2021-04-15, from <http://www.fao.org/3/x0490e/x0490e00.htm>
- Amiro, B. (2009). Measuring boreal forest evapotranspiration using the energy balance residual. *Journal of Hydrology*, 366(1), 112–118. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0022169408006367> doi: <https://doi.org/10.1016/j.jhydrol.2008.12.021>
- Anderson, M. C., Allen, R. G., Morse, A., & Kustas, W. P. (2012). Use of landsat thermal imagery in monitoring evapotranspiration and managing water resources. *Remote Sensing of Environment*, 122, 50–65. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0034425712000326> (Landsat Legacy Special Issue) doi: <https://doi.org/10.1016/j.rse.2011.08.025>
- Anderson, M. C., Hain, C., Wardlow, B., Pimstein, A., Mecikalski, J. R., & Kustas, W. P. (2011). Evaluation of drought indices based on thermal remote sensing of evapotranspiration over the continental united states. *Journal of Climate*, 24(8), 2025 - 2044. Retrieved from <https://journals.ametsoc.org/view/journals/clim/24/8/2010jcli3812.1.xml> doi: 10.1175/2010JCLI3812.1
- Anderson, M. C., Zolin, C. A., Sentelhas, P. C., Hain, C. R., Semmens, K., Tugrul Yilmaz, M., ... Tetrault, R. (2016, March). The Evaporative Stress Index as an indicator of agricultural drought in Brazil: An assessment based on crop yield impacts. *Remote Sensing of Environment*, 174, 82–99. Retrieved 2019-10-17, from <http://www.sciencedirect.com/science/article/pii/S0034425715302212> doi: 10.1016/j.rse.2015.11.034
- Aragon, B., Houborg, R., Tu, K., Fisher, J. B., & McCabe, M. (2018, December). CubeSats Enable High Spatiotemporal Retrievals of Crop-Water Use for Precision Agriculture. *Remote Sensing*, 10(12), 1867. Retrieved 2020-09-01, from <https://www.mdpi.com/2072-4292/10/12/1867> (Number: 12 Publisher: Multidisciplinary Digital Publishing Institute) doi: 10.3390/rs10121867
- Aron, P. G., Poulsen, C. J., Fiorella, R. P., Matheny, A. M., & Veverica, T. J. (2020). An isotopic approach to partition evapotranspiration in a mixed deciduous forest. *Ecohydrology*, 13(6), e2229. Retrieved 2021-04-11, from <https://onlinelibrary.wiley.com/doi/abs/10.1002/eco.2229> (_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/eco.2229>) doi: <https://doi.org/10.1002/eco.2229>
- Arroyo, M. T. K., Pliscoff, P., Mihoc, M., & Arroyo-Kalin, M. (2005). The Magellanic moorland. In L. H. Fraser & P. A. Keddy (Eds.), *The World's Largest Wetlands: Ecology and Conservation* (pp. 424–445). Cambridge: Cambridge University Press. Retrieved 2020-08-15, from <https://www.cambridge.org/core/books/worlds-largest-wetlands/magellanic-moorland/B02EB2AE6B5BC3EB290B88FE794D3A77> doi: 10.1017/CBO9780511542091.013
- Arruda, P. H. Z., Vourlitis, G. L., Santanna, F. B., Jr., O. B. P., Lobo, F. A., & Nogueira, J. S. (2016). Large net co2 loss from a grass-dominated tropical savanna in south-central brazil in response to seasonal and interannual drought.

- Biogeosciences*, 121, 2110-2124. doi: 10.1002/2016JG003404
- Badgley, G., Fisher, J. B., Jiménez, C., Tu, K. P., & Vinukollu, R. (2015, August). On Uncertainty in Global Terrestrial Evapotranspiration Estimates from Choice of Input Forcing Datasets. *Journal of Hydrometeorology*, 16(4), 1449-1455. Retrieved 2020-09-01, from <https://journals.ametsoc.org/jhm/article/16/4/1449/6244/On-Uncertainty-in-Global-Terrestrial> (Publisher: American Meteorological Society) doi: 10.1175/JHM-D-14-0040.1
- Bai, J., Jia, L., Liu, S., Xu, Z., Hu, G., Zhu, M., & Song, L. (2015, May). Characterizing the Footprint of Eddy Covariance System and Large Aperture Scintillometer Measurements to Validate Satellite-Based Surface Fluxes. *IEEE Geoscience and Remote Sensing Letters*, 12(5), 943-947. (Conference Name: IEEE Geoscience and Remote Sensing Letters) doi: 10.1109/LGRS.2014.2368580
- Bastiaanssen, W. G. M., Menenti, M., Feddes, R. A., & Holtslag, A. A. M. (1998, December). A remote sensing surface energy balance algorithm for land (SEBAL). 1. Formulation. *Journal of Hydrology*, 212-213, 198-212. Retrieved 2020-06-15, from <http://www.sciencedirect.com/science/article/pii/S0022169498002534> doi: 10.1016/S0022-1694(98)00253-4
- Bezerra, B. G., Santos, C. A. C. d., Silva, B. B. d., Perez-Marin, A. M., Bezerra, M. V. C., Bezerra, J. R. C., & Rao, T. V. R. (2013, June). Estimation of soil moisture in the root-zone from remote sensing data. *Revista Brasileira de Ciência do Solo*, 37(3), 596-603. Retrieved 2021-04-15, from http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-06832013000300005&lng=en&nrm=iso&tlng=en (Publisher: Sociedade Brasileira de Ciência do Solo) doi: 10.1590/S0100-06832013000300005
- Bezerra, B. G., Silva, B. B. d., Santos, C. A. C. d., & Bezerra, J. R. C. (2015, July). Actual Evapotranspiration Estimation Using Remote Sensing: Comparison of SEBAL and SSEB Approaches. *Advances in Remote Sensing*, 4(3), 234-247. Retrieved 2021-04-15, from <http://www.scirp.org/Journal/Paperabs.aspx?paperid=59977> (Number: 3 Publisher: Scientific Research Publishing) doi: 10.4236/ars.2015.43019
- Blunden, J., & Arndt, D. S. (2020, August). State of the Climate in 2019. *Bulletin of the American Meteorological Society*, 101(8), S1-S429. Retrieved 2020-09-03, from <https://journals.ametsoc.org/bams/article/101/8/S1/353885/State-of-the-Climate-in-2019> (Publisher: American Meteorological Society) doi: 10.1175/2020BAMSSStateoftheClimate.1
- Borma, L. S., Rocha, H. R. d., Cabral, O. M., Randow, C. v., Collicchio, E., Kurzatkowski, D., ... Artaxo, P. (2009). Atmosphere and hydrological controls of the evapotranspiration over a floodplain forest in the Bananal Island region, Amazonia. *Journal of Geophysical Research: Biogeosciences*, 114(G1). Retrieved 2020-04-21, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000641> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2007JG000641>) doi: 10.1029/2007JG000641
- Cabral, O. M. R., da Rocha, H. R., Gash, J. H., Freitas, H. C., & Ligo, M. A. V. (2015, September). Water and energy fluxes from a woodland savanna (cerrado) in southeast Brazil. *Journal of Hydrology: Regional Studies*, 4, 22-40. Retrieved 2020-08-05, from <http://www.sciencedirect.com/science/article/pii/S2214581815000440> doi: 10.1016/j.ejrh.2015.04.010
- Cabral, O. M. R., Freitas, H. C., Cuadra, S. V., de Andrade, C. A., Ramos, N. P., Grutzmacher, P., ... Rossi, P. (2020, March). The sustainability of a sugarcane plantation in Brazil assessed by the eddy covariance fluxes of greenhouse gases. *Agricultural and Forest Meteorology*, 282-283, 107864. Retrieved 2021-10-01, from <https://www.sciencedirect.com/science/article/pii/S0168192319304800> doi: 10.1016/j.agrformet.2019.107864
- Cabral, O. M. R., Gash, J. H. C., Rocha, H. R., Marsden, C., Ligo, M. A. V., Fre-

- itas, H. C., ... Gomes, E. (2011, January). Fluxes of CO₂ above a plantation of Eucalyptus in southeast Brazil. *Agricultural and Forest Meteorology*, 151(1), 49–59. Retrieved 2020-07-24, from <http://www.sciencedirect.com/science/article/pii/S0168192310002480> doi: 10.1016/j.agrformet.2010.09.003
- Cabral, O. M. R., Rocha, H. R., Gash, J. H., Ligo, M. A. V., Tatsch, J. D., Freitas, H. C., & Brasílio, E. (2012). Water use in a sugarcane plantation. *GCB Bioenergy*, 4(5), 555–565. Retrieved 2020-07-28, from <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1757-1707.2011.01155.x> (eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1757-1707.2011.01155.x>) doi: 10.1111/j.1757-1707.2011.01155.x
- Cabral, O. M. R., Rocha, H. R., Gash, J. H. C., Ligo, M. A. V., Freitas, H. C., & Tatsch, J. D. (2010, July). The energy and water balance of a Eucalyptus plantation in southeast Brazil. *Journal of Hydrology*, 388(3), 208–216. Retrieved 2021-04-08, from <https://www.sciencedirect.com/science/article/pii/S0022169410002404> doi: 10.1016/j.jhydrol.2010.04.041
- Campos, S., Mendes, K. R., da Silva, L. L., Mutti, P. R., Medeiros, S. S., Amorim, L. B., ... Bezerra, B. G. (2019, June). Closure and partitioning of the energy balance in a preserved area of a Brazilian seasonally dry tropical forest. *Agricultural and Forest Meteorology*, 271, 398–412. Retrieved 2020-05-06, from <http://www.sciencedirect.com/science/article/pii/S0168192319301303> doi: 10.1016/j.agrformet.2019.03.018
- Cao, L., Bala, G., Caldeira, K., Nemani, R., & Ban-Weiss, G. (2010, May). Importance of carbon dioxide physiological forcing to future climate change. *Proceedings of the National Academy of Sciences*, 107(21), 9513–9518. Retrieved 2021-04-15, from <https://www.pnas.org/content/107/21/9513> (Publisher: National Academy of Sciences Section: Physical Sciences) doi: 10.1073/pnas.0913000107
- Chang, Y., Qin, D., Ding, Y., Zhao, Q., & Zhang, S. (2018, June). A modified MOD16 algorithm to estimate evapotranspiration over alpine meadow on the Tibetan Plateau, China. *Journal of Hydrology*, 561, 16–30. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0022169418302269> doi: 10.1016/j.jhydrol.2018.03.054
- Chen, B., Black, T. A., Coops, N. C., Hilker, T., (Tony) Trofymow, J. A., & Morgenstern, K. (2009, February). Assessing Tower Flux Footprint Climatology and Scaling Between Remotely Sensed and Eddy Covariance Measurements. *Boundary-Layer Meteorology*, 130(2), 137–167. Retrieved 2020-07-09, from <https://doi.org/10.1007/s10546-008-9339-1> doi: 10.1007/s10546-008-9339-1
- Chong, D. L. S., Mougin, E., & Gastellu-Etchegorry. (1993, May). Relating the Global Vegetation Index to net primary productivity and actual evapotranspiration over Africa. *International Journal of Remote Sensing*, 14(8), 1517–1546. Retrieved 2019-09-05, from <https://doi.org/10.1080/01431169308953984> doi: 10.1080/01431169308953984
- Costanza, R., d’Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., ... van den Belt, M. (1997, May). The value of the world’s ecosystem services and natural capital. *Nature*, 387(6630), 253–260. Retrieved 2021-04-22, from <https://www.nature.com/articles/387253a0> (Number: 6630 Publisher: Nature Publishing Group) doi: 10.1038/387253a0
- Curto, L., Covi, M., & Gassmann, M. I. (2019, December). Actual evapotranspiration and the pattern of soil water extraction of a soybean (*Glycine max*) crop. *Revista de la Facultad de Ciencias Agrarias UNCuyo*, 51(2), 125–141. Retrieved 2021-03-11, from <http://revistas.uncu.edu.ar/ojs/index.php/RFA/article/view/2615> (Number: 2)
- Dalmagro, H. J., Lathuillière, M. J., Hawthorne, I., Morais, D. D., Pinto Jr, O. B., Couto, E. G., & Johnson, M. S. (2018, June). Carbon biogeochemistry

- of a flooded Pantanal forest over three annual flood cycles. *Biogeochemistry*, 139(1), 1–18. Retrieved 2020-08-18, from <https://doi.org/10.1007/s10533-018-0450-1> doi: 10.1007/s10533-018-0450-1
- Denmead, O. T., Mayocchi, C. L., & Dunin, F. X. (1997). Does green cane harvesting conserve soil water? In *Proceedings of the Australian Society of Sugar Cane Technologists* (Vol. 19, pp. 139–146).
- de Oliveira, R. G., Valle Júnior, L. C. G., da Silva, J. B., Espíndola, D. A. L. F., Lopes, R. D., Nogueira, J. S., ... Rodrigues, T. R. (2021, May). Temporal trend changes in reference evapotranspiration contrasting different land uses in southern Amazon basin. *Agricultural Water Management*, 250, 106815. Retrieved 2021-04-01, from <https://www.sciencedirect.com/science/article/pii/S0378377421000809> doi: 10.1016/j.agwat.2021.106815
- de Queiroz, M. G., da Silva, T. G. F., Zolnier, S., de Souza, C. A. A., de Souza, L. S. B., do Nascimento Araújo, G., ... de Moura, M. S. B. (2020, April). Partitioning of rainfall in a seasonal dry tropical forest. *Ecohydrology & Hydrobiology*, 20(2), 230–242. Retrieved 2021-03-31, from <https://www.sciencedirect.com/science/article/pii/S1642359320300124> doi: 10.1016/j.ecohyd.2020.02.001
- Embry, J. L., & Nothnagel, E. A. (1994, September). Leaf Senescence of Post-production Poinsettias in Low-light Stress. *Journal of the American Society for Horticultural Science*, 119(5), 1006–1013. Retrieved 2021-04-22, from <https://journals.ashs.org/jashs/view/journals/jashs/119/5/article-p1006.xml> (Publisher: American Society for Horticultural Science Section: Journal of the American Society for Horticultural Science) doi: 10.21273/JASHS.119.5.1006
- Ershadi, A., McCabe, M. F., Evans, J. P., Chaney, N. W., & Wood, E. F. (2014, April). Multi-site evaluation of terrestrial evaporation models using FLUXNET data. *Agricultural and Forest Meteorology*, 187, 46–61. Retrieved 2020-01-15, from <http://www.sciencedirect.com/science/article/pii/S0168192313002980> doi: 10.1016/j.agrformet.2013.11.008
- Ershadi, A., McCabe, M. F., Evans, J. P., & Wood, E. F. (2015, June). Impact of model structure and parameterization on Penman–Monteith type evaporation models. *Journal of Hydrology*, 525, 521–535. Retrieved 2020-06-22, from <http://www.sciencedirect.com/science/article/pii/S0022169415002577> doi: 10.1016/j.jhydrol.2015.04.008
- Ferguson, C. R., Sheffield, J., Wood, E. F., & Gao, H. (2010, August). Quantifying uncertainty in a remote sensing-based estimate of evapotranspiration over continental USA. *International Journal of Remote Sensing*, 31(14), 3821–3865. Retrieved 2020-09-03, from <https://doi.org/10.1080/01431161.2010.483490> (Publisher: Taylor & Francis) eprint: <https://doi.org/10.1080/01431161.2010.483490> doi: 10.1080/01431161.2010.483490
- Ferretti, D. F., Pendall, E., Morgan, J. A., Nelson, J. A., LeCain, D., & Mosier, A. R. (2003, July). Partitioning evapotranspiration fluxes from a Colorado grassland using stable isotopes: Seasonal variations and ecosystem implications of elevated atmospheric CO₂. *Plant and Soil*, 254(2), 291–303. Retrieved 2021-04-19, from <https://doi.org/10.1023/A:1025511618571> doi: 10.1023/A:1025511618571
- Fisher, J. B., Lee, B., Purdy, A. J., Halverson, G. H., Dohlen, M. B., Cawse-Nicholson, K., ... Hook, S. (2020). ECOSTRESS: NASA's Next Generation Mission to Measure Evapotranspiration From the International Space Station. *Water Resources Research*, 56(4), e2019WR026058. Retrieved 2020-08-31, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019WR026058> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019WR026058>) doi: 10.1029/2019WR026058

- 10.1029/2019WR026058
- Fisher, J. B., Malhi, Y., Bonal, D., Rocha, H. R. D., Araújo, A. C. D., Gamo, M.,
 ... Randow, C. V. (2009). The land-atmosphere water flux in the tropics. *Global Change Biology*, 15(11), 2694–2714. Retrieved 2021-01-26, from <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2486.2008.01813.x> (eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2486.2008.01813.x>) doi: <https://doi.org/10.1111/j.1365-2486.2008.01813.x>
- Fisher, J. B., Melton, F., Middleton, E., Hain, C., Anderson, M., Allen, R.,
 ... Wood, E. F. (2017). The future of evapotranspiration: Global requirements for ecosystem functioning, carbon and climate feedbacks, agricultural management, and water resources. *Water Resources Research*, 53(4), 2618–2626. Retrieved 2020-08-31, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016WR020175> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2016WR020175>) doi: <https://doi.org/10.1002/2016WR020175>
- Fisher, J. B., Tu, K. P., & Baldocchi, D. D. (2008, March). Global estimates of the land-atmosphere water flux based on monthly AVHRR and ISLSCP-II data, validated at 16 FLUXNET sites. *Remote Sensing of Environment*, 112(3), 901–919. Retrieved 2018-08-09, from <http://www.sciencedirect.com/science/article/pii/S0034425707003938> doi: [10.1016/j.rse.2007.06.025](https://doi.org/10.1016/j.rse.2007.06.025)
- Fisher, J. B., Whittaker, R. J., & Malhi, Y. (2011). ET come home: potential evapotranspiration in geographical ecology. *Global Ecology and Biogeography*, 20(1), 1–18. Retrieved 2020-08-31, from <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1466-8238.2010.00578.x> (eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1466-8238.2010.00578.x>) doi: [10.1111/j.1466-8238.2010.00578.x](https://doi.org/10.1111/j.1466-8238.2010.00578.x)
- Foken, T. (2008). The Energy Balance Closure Problem: An Overview. *Ecological Applications*, 18(6), 1351–1367. Retrieved 2021-02-19, from <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/06-0922.1> (eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/06-0922.1>) doi: <https://doi.org/10.1890/06-0922.1>
- Gaj, M., Beyer, M., Koeniger, P., Wanke, H., Hamutoko, J., & Himmelsbach, T. (2016, February). In situ unsaturated zone water stable isotope (^2H and ^{18}O) measurements in semi-arid environments: a soil water balance. *Hydrology and Earth System Sciences*, 20(2), 715–731. Retrieved 2021-04-08, from <https://hess.copernicus.org/articles/20/715/2016/> (Publisher: Copernicus GmbH) doi: [10.5194/hess-20-715-2016](https://doi.org/10.5194/hess-20-715-2016)
- García, A. G., Di Bella, C. M., Houspanossian, J., Magliano, P. N., Jobbágy, E. G., Posse, G., ... Nasetto, M. D. (2017, December). Patterns and controls of carbon dioxide and water vapor fluxes in a dry forest of central Argentina. *Agricultural and Forest Meteorology*, 247, 520–532. Retrieved 2021-03-11, from <https://www.sciencedirect.com/science/article/pii/S0168192317302721> doi: [10.1016/j.agrformet.2017.08.015](https://doi.org/10.1016/j.agrformet.2017.08.015)
- García, M., Sandholt, I., Ceccato, P., Ridler, M., Mougin, E., Kergoat, L., ... Domingo, F. (2013, April). Actual evapotranspiration in drylands derived from in-situ and satellite data: Assessing biophysical constraints. *Remote Sensing of Environment*, 131, 103–118. Retrieved 2020-06-23, from <http://www.sciencedirect.com/science/article/pii/S0034425712004828> doi: [10.1016/j.rse.2012.12.016](https://doi.org/10.1016/j.rse.2012.12.016)
- Gash, J. H. C. (1979). An analytical model of rainfall interception by forests. *Quarterly Journal of the Royal Meteorological Society*, 105(443), 43–55. Retrieved 2021-04-15, from <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710544304> (eprint: <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/qj.49710544304>) doi: <https://doi.org/10.1002/qj.49710544304>

- Gash, J. H. C., Lloyd, C. R., & Lachaud, G. (1995, August). Estimating sparse forest rainfall interception with an analytical model. *Journal of Hydrology*, 170(1), 79–86. Retrieved 2020-07-28, from <http://www.sciencedirect.com/science/article/pii/002216949502697N> doi: 10.1016/0022-1694(95)02697-N
- Goymer, P. (2017, March). Spotlight on South America. *Nature Ecology & Evolution*, 1(4), 1–2. Retrieved 2020-07-08, from <https://www.nature.com/articles/s41559-017-0129> (Number: 4 Publisher: Nature Publishing Group) doi: 10.1038/s41559-017-0129
- Harris, I., Osborn, T. J., Jones, P., & Lister, D. (2020, April). Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Scientific Data*, 7(1), 109. Retrieved 2021-02-19, from <https://www.nature.com/articles/s41597-020-0453-3> (Number: 1 Publisher: Nature Publishing Group) doi: 10.1038/s41597-020-0453-3
- Hasler, N., & Avissar, R. (2007, June). What Controls Evapotranspiration in the Amazon Basin? *Journal of Hydrometeorology*, 8(3), 380–395. Retrieved 2020-03-23, from <https://journals.ametsoc.org/doi/full/10.1175/JHM587.1> (Publisher: American Meteorological Society) doi: 10.1175/JHM587.1
- Holl, D., Pancotto, V., Heger, A., Camargo, S., & Kutzbach, L. (2019). Cushion bogs are stronger carbon dioxide net sinks than moss-dominated bogs as revealed by eddy covariance measurements on tierra del fuego, argentina. *Biogeosciences*, 16, 3397–3423. doi: <https://doi.org/10.5194/bg-16-3397-2019>
- Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002, November). Overview of the radiometric and biophysical performance of the MODIS vegetation indices. *Remote Sensing of Environment*, 83(1), 195–213. Retrieved 2019-09-03, from <http://www.sciencedirect.com/science/article/pii/S0034425702000962> doi: 10.1016/S0034-4257(02)00096-2
- Hutyra, L. R., Munger, J. W., Saleska, S. R., Gottlieb, E., Daube, B. C., Dunn, A. L., ... Wofsy, S. C. (2007). Seasonal controls on the exchange of carbon and water in an Amazonian rain forest. *Journal of Geophysical Research: Biogeosciences*, 112(G3). Retrieved 2020-03-23, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JG000365> (_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JG000365>) doi: 10.1029/2006JG000365
- Jang, K., Kang, S., Lim, Y.-J., Jeong, S., Kim, J., Kimball, J. S., & Hong, S. Y. (2013). Monitoring daily evapotranspiration in North-east Asia using MODIS and a regional Land Data Assimilation System. *Journal of Geophysical Research: Atmospheres*, 118(23), 12,927–12,940. Retrieved 2021-04-15, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020639> (_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2013JD020639>) doi: <https://doi.org/10.1002/2013JD020639>
- Jarchow, C. J., Nagler, P. L., Glenn, E. P., Ramírez-Hernández, J., & Rodríguez-Burgueño, J. E. (2017, September). Evapotranspiration by remote sensing: An analysis of the Colorado River Delta before and after the Minute 319 pulse flow to Mexico. *Ecological Engineering*, 106, 725–732. Retrieved 2019-09-05, from <http://www.sciencedirect.com/science/article/pii/S0925857416305833> doi: 10.1016/j.ecoleng.2016.10.056
- Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y., & Fawcett, P. J. (2013, April). Terrestrial water fluxes dominated by transpiration. *Nature*, 496(7445), 347–350. Retrieved 2021-04-13, from <https://www.nature.com/articles/nature11983> (Number: 7445 Publisher: Nature Publishing Group) doi: 10.1038/nature11983
- Junk, W. J., Brown, M., Campbell, I. C., Finlayson, M., Gopal, B., Ramberg, L., & Warner, B. G. (2006, October). The comparative biodiversity of seven

- globally important wetlands: a synthesis. *Aquatic Sciences*, 68(3), 400–414. Retrieved 2021-04-22, from <https://doi.org/10.1007/s00027-006-0856-z> doi: 10.1007/s00027-006-0856-z
- Junk, W. J., da Cunha, C. N., Wantzen, K. M., Petermann, P., Strüßmann, C., Marques, M. I., & Adis, J. (2006, October). Biodiversity and its conservation in the Pantanal of Mato Grosso, Brazil. *Aquatic Sciences*, 68(3), 278–309. Retrieved 2021-04-22, from <https://doi.org/10.1007/s00027-006-0851-4> doi: 10.1007/s00027-006-0851-4
- Khan, M. S., Liaqat, U. W., Baik, J., & Choi, M. (2018, April). Stand-alone uncertainty characterization of GLEAM, GLDAS and MOD16 evapotranspiration products using an extended triple collocation approach. *Agricultural and Forest Meteorology*, 252, 256–268. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0168192318300224> doi: 10.1016/j.agrformet.2018.01.022
- Khosa, F. V., Feig, G. T., van der Merwe, M. R., Mateyisi, M. J., Mudau, A. E., & Savage, M. J. (2019, December). Evaluation of modeled actual evapotranspiration estimates from a land surface, empirical and satellite-based models using in situ observations from a South African semi-arid savanna ecosystem. *Agricultural and Forest Meteorology*, 279, 107706. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0168192319303223> doi: 10.1016/j.agrformet.2019.107706
- Kustas, W. P., & Norman, J. M. (1999, April). Evaluation of soil and vegetation heat flux predictions using a simple two-source model with radiometric temperatures for partial canopy cover. *Agricultural and Forest Meteorology*, 94(1), 13–29. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0168192399000052> doi: 10.1016/S0168-1923(99)00005-2
- Kutzbach, L. (2019a). Lars Kutzbach (2019), AmeriFlux AR-TF1 Rio Moat bog, Ver. 1-5, AmeriFlux AMP, (Dataset).. Retrieved 2021-04-15, from <https://ameriflux.lbl.gov/doi/AmeriFlux/AR-TF1> doi: <https://doi.org/10.17190/AMF/1543389>
- Kutzbach, L. (2019b). Lars Kutzbach (2019), AmeriFlux AR-TF2 Rio Pipo bog, Ver. 1-5, AmeriFlux AMP, (Dataset).. Retrieved 2021-04-15, from <https://ameriflux.lbl.gov/sites/siteinfo/AR-TF2> doi: <https://doi.org/10.17190/AMF/1543388>
- Leopoldo, P. R., Franken, W. K., & Villa Nova, N. A. (1995, May). Real evapotranspiration and transpiration through a tropical rain forest in central Amazonia as estimated by the water balance method. *Forest Ecology and Management*, 73(1), 185–195. Retrieved 2021-04-08, from <https://www.sciencedirect.com/science/article/pii/037811279403487H> doi: 10.1016/0378-1127(94)03487-H
- Levy, P., Drewer, J., Jammet, M., Leeson, S., Friborg, T., Skiba, U., & Oijen, M. v. (2020, January). Inference of spatial heterogeneity in surface fluxes from eddy covariance data: A case study from a subarctic mire ecosystem. *Agricultural and Forest Meteorology*, 280, 107783. Retrieved 2020-07-10, from <http://www.sciencedirect.com/science/article/pii/S0168192319303995> doi: 10.1016/j.agrformet.2019.107783
- Li, X., Long, D., Han, Z., Scanlon, B. R., Sun, Z., Han, P., & Hou, A. (2019, November). Evapotranspiration Estimation for Tibetan Plateau Headwaters Using Conjoint Terrestrial and Atmospheric Water Balances and Multisource Remote Sensing. *Water Resources Research*, 55(11), 8608–8630. Retrieved 2021-04-15, from <https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2019WR025196> (Publisher: John Wiley & Sons, Ltd) doi: 10.1029/2019WR025196
- Liu, Y. Y., Dijk, A. I. J. M. v., McCabe, M. F., Evans, J. P., & Jeu, R. A. M. d.

- (2013). Global vegetation biomass change (1988–2008) and attribution to environmental and human drivers. *Global Ecology and Biogeography*, 22(6), 692–705. Retrieved 2021-04-15, from <https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.12024> (eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/geb.12024>) doi: <https://doi.org/10.1111/geb.12024>
- Lopes, J. D., Rodrigues, L. N., Imbuzeiro, H. M. A., & Pruski, F. F. (2019, September). Performance of SSEBop model for estimating wheat actual evapotranspiration in the Brazilian Savannah region. *International Journal of Remote Sensing*, 40(18), 6930–6947. Retrieved 2021-04-15, from <https://doi.org/10.1080/01431161.2019.1597304> (Publisher: Taylor & Francis eprint: <https://doi.org/10.1080/01431161.2019.1597304>) doi: <https://doi.org/10.1080/01431161.2019.1597304>
- Machado, C. B., Lima, J. R. d. S., Antonino, A. C. D., Souza, E. S. d., Souza, R. M. S., Alves, E. M., ... Alves, E. M. (2016, September). Daily and seasonal patterns of CO₂ fluxes and evapotranspiration in maize-grass intercropping. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 20(9), 777–782. Retrieved 2021-03-11, from http://www.scielo.br/scielo.php?script=sci_abstract&pid=S1415-43662016000900777&lng=en&nrm=iso&tlng=en (Publisher: Departamento de Engenharia Agrícola - UFCG / Cnpq) doi: <https://doi.org/10.1590/1807-1929/agriambi.v20n9p777-782>
- Mao, J., Fu, W., Shi, X., Ricciuto, D. M., Fisher, J. B., Dickinson, R. E., ... Zhu, Z. (2015, sep). Disentangling climatic and anthropogenic controls on global terrestrial evapotranspiration trends. *Environmental Research Letters*, 10(9), 094008. Retrieved from <https://doi.org/10.1088/1748-9326/10/9/094008> doi: <https://doi.org/10.1088/1748-9326/10/9/094008>
- Marques, T. V., Mendes, K., Mutti, P., Medeiros, S., Silva, L., Perez-Marin, A. M., ... Bezerra, B. (2020, June). Environmental and biophysical controls of evapotranspiration from Seasonally Dry Tropical Forests (Caatinga) in the Brazilian Semiarid. *Agricultural and Forest Meteorology*, 287, 107957. Retrieved 2021-04-22, from <https://www.sciencedirect.com/science/article/pii/S0168192320300599> doi: <https://doi.org/10.1016/j.agrformet.2020.107957>
- Martens, B., Miralles, D., Lievens, H., Fernández-Prieto, D., & Verhoest, N. E. C. (2016, June). Improving terrestrial evaporation estimates over continental Australia through assimilation of SMOS soil moisture. *International Journal of Applied Earth Observation and Geoinformation*, 48, 146–162. Retrieved 2021-04-13, from <https://www.sciencedirect.com/science/article/pii/S0303243415300350> doi: <https://doi.org/10.1016/j.jag.2015.09.012>
- Martens, B., Miralles, D. G., Lievens, H., Schalie, R. v. d., Jeu, R. A. M. d., Fernández-Prieto, D., ... Verhoest, N. E. C. (2017, May). GLEAM v3: satellite-based land evaporation and root-zone soil moisture. *Geoscientific Model Development*, 10(5), 1903–1925. Retrieved 2020-07-01, from <https://gmd.copernicus.org/articles/10/1903/2017/> (Publisher: Copernicus GmbH) doi: <https://doi.org/10.5194/gmd-10-1903-2017>
- Mauder, M., Foken, T., & Cuxart, J. (2020, December). Surface-Energy-Balance Closure over Land: A Review. *Boundary-Layer Meteorology*, 177(2), 395–426. Retrieved 2021-02-19, from <https://doi.org/10.1007/s10546-020-00529-6> doi: <https://doi.org/10.1007/s10546-020-00529-6>
- McCabe, M. F., Ershadi, A., Jimenez, C., Miralles, D. G., Michel, D., & Wood, E. F. (2016, January). The GEWEX LandFlux project: evaluation of model evaporation using tower-based and globally gridded forcing data. *Geoscientific Model Development*, 9(1), 283–305. Retrieved 2019-08-29, from <https://www.geosci-model-dev.net/9/283/2016/> doi: <https://doi.org/10.5194/gmd-9-283-2016>
- McColl, K. A. (2020). Practical and Theoretical Benefits of an Alterna-

- tive to the Penman-Monteith Evapotranspiration Equation. *Water Resources Research*, 56(6), e2020WR027106. Retrieved 2021-09-30, from <https://onlinelibrary.wiley.com/doi/abs/10.1029/2020WR027106> (_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020WR027106>) doi: 10.1029/2020WR027106
- Michel, D., Jiménez, C., Miralles, D. G., Jung, M., Hirschi, M., Ershadi, A., ... Fernández-Prieto, D. (2016, February). The WACMOS-ET project – Part 1: Tower-scale evaluation of four remote-sensing-based evapotranspiration algorithms. *Hydrology and Earth System Sciences*, 20(2), 803–822. Retrieved 2019-08-29, from <https://www.hydrol-earth-syst-sci.net/20/803/2016/> doi: <https://doi.org/10.5194/hess-20-803-2016>
- Miralles, D. G., Gash, J. H., Holmes, T. R. H., Jeu, R. A. M. d., & Dolman, A. J. (2010). Global canopy interception from satellite observations. *Journal of Geophysical Research: Atmospheres*, 115(D16). Retrieved 2020-09-04, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2009JD013530> (_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JD013530>) doi: 10.1029/2009JD013530
- Miralles, D. G., Holmes, T. R. H., Jeu, R. A. M. D., Gash, J. H., Meesters, A. G. C. A., & Dolman, A. J. (2011, February). Global land-surface evaporation estimated from satellite-based observations. *Hydrology and Earth System Sciences*, 15(2), 453–469. Retrieved 2020-06-15, from <https://www.hydrol-earth-syst-sci.net/15/453/2011/hess-15-453-2011.html> (Publisher: Copernicus GmbH) doi: <https://doi.org/10.5194/hess-15-453-2011>
- Miralles, D. G., Jiménez, C., Jung, M., Michel, D., Ershadi, A., McCabe, M. F., ... Fernández-Prieto, D. (2016, February). The WACMOS-ET project – Part 2: Evaluation of global terrestrial evaporation data sets. *Hydrology and Earth System Sciences*, 20(2), 823–842. Retrieved 2019-08-29, from <https://www.hydrol-earth-syst-sci.net/20/823/2016/> doi: <https://doi.org/10.5194/hess-20-823-2016>
- Miranda, R. d. Q., Galv ncio, J. D., Moura, M. S. B. d., Jones, C. A., & Srinivasan, R. (2017). *Reliability of MODIS Evapotranspiration Products for Heterogeneous Dry Forest: A Study Case of Caatinga* [Research Article]. Retrieved 2020-05-06, from <https://www.hindawi.com/journals/amete/2017/9314801/> (ISSN: 1687-9309 Library Catalog: www.hindawi.com Pages: e9314801 Publisher: Hindawi Volume: 2017) doi: <https://doi.org/10.1155/2017/9314801>
- Moesinger, L., Dorigo, W., de Jeu, R., van der Schalie, R., Scanlon, T., Teubner, I., & Forkel, M. (2020, January). The global long-term microwave Vegetation Optical Depth Climate Archive (VODCA). *Earth System Science Data*, 12(1), 177–196. Retrieved 2021-04-13, from <https://essd.copernicus.org/articles/12/177/2020/> (Publisher: Copernicus GmbH) doi: 10.5194/essd-12-177-2020
- Moreira, A. A., Ruhoff, A. L., Roberti, D. R., Souza, V. d. A., da Rocha, H. R., & Paiva, R. C. D. d. (2019, August). Assessment of terrestrial water balance using remote sensing data in South America. *Journal of Hydrology*, 575, 131–147. Retrieved 2019-08-28, from <http://www.sciencedirect.com/science/article/pii/S0022169419304664> doi: 10.1016/j.jhydrol.2019.05.021
- Mu, Q., Heinsch, F. A., Zhao, M., & Running, S. W. (2007, December). Development of a global evapotranspiration algorithm based on MODIS and global meteorology data. *Remote Sensing of Environment*, 111(4), 519–536. Retrieved 2017-09-21, from <http://www.sciencedirect.com/science/article/pii/S0034425707001903> doi: 10.1016/j.rse.2007.04.015
- Mu, Q., Zhao, M., & Running, S. W. (2011, August). Improvements to a MODIS global terrestrial evapotranspiration algorithm. *Remote Sens-*

- ing of *Environment*, 115(8), 1781–1800. Retrieved 2020-06-21, from <http://www.sciencedirect.com/science/article/pii/S0034425711000691> doi: 10.1016/j.rse.2011.02.019
- Mutti, P. R., da Silva, L. L., Medeiros, S. d. S., Dubreuil, V., Mendes, K. R., Marques, T. V., ... Bezerra, B. G. (2019, March). Basin scale rainfall-evapotranspiration dynamics in a tropical semiarid environment during dry and wet years. *International Journal of Applied Earth Observation and Geoinformation*, 75, 29–43. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0303243418307244> doi: 10.1016/j.jag.2018.10.007
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Fonseca, G. A. B. d., & Kent, J. (2000, February). Biodiversity hotspots for conservation priorities. *Nature*, 403(6772), 853–858. Retrieved 2019-08-28, from <https://www.nature.com/articles/35002501> doi: 10.1038/35002501
- Nagler, P. L., Cleverly, J., Glenn, E., Lampkin, D., Huete, A., & Wan, Z. (2005, January). Predicting riparian evapotranspiration from MODIS vegetation indices and meteorological data. *Remote Sensing of Environment*, 94(1), 17–30. Retrieved 2020-06-19, from <http://www.sciencedirect.com/science/article/pii/S0034425704002615> doi: 10.1016/j.rse.2004.08.009
- Nagler, P. L., Glenn, E. P., Kim, H., Emmerich, W., Scott, R. L., Huxman, T. E., & Huete, A. R. (2007, August). Relationship between evapotranspiration and precipitation pulses in a semiarid rangeland estimated by moisture flux towers and MODIS vegetation indices. *Journal of Arid Environments*, 70(3), 443–462. Retrieved 2019-09-05, from <http://www.sciencedirect.com/science/article/pii/S0140196307000249> doi: 10.1016/j.jaridenv.2006.12.026
- Nagler, P. L., Glenn, E. P., Nguyen, U., Scott, R. L., & Doody, T. (2013, August). Estimating Riparian and Agricultural Actual Evapotranspiration by Reference Evapotranspiration and MODIS Enhanced Vegetation Index. *Remote Sensing*, 5(8), 3849–3871. Retrieved 2019-09-03, from <https://www.mdpi.com/2072-4292/5/8/3849> doi: 10.3390/rs5083849
- Nagler, P. L., Morino, K., Murray, R. S., Osterberg, J., & Glenn, E. P. (2009, December). An Empirical Algorithm for Estimating Agricultural and Riparian Evapotranspiration Using MODIS Enhanced Vegetation Index and Ground Measurements of ET. I. Description of Method. *Remote Sensing*, 1(4), 1273–1297. Retrieved 2019-09-03, from <https://www.mdpi.com/2072-4292/1/4/1273> doi: 10.3390/rs1041273
- Norman, J. M., Kustas, W. P., & Humes, K. S. (1995, December). Source approach for estimating soil and vegetation energy fluxes in observations of directional radiometric surface temperature. *Agricultural and Forest Meteorology*, 77(3), 263–293. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/016819239502265Y> doi: 10.1016/0168-1923(95)02265-Y
- Nouri, H., Glenn, E. P., Beecham, S., Chavoshi Boroujeni, S., Sutton, P., Alghamand, S., ... Nagler, P. (2016, June). Comparing Three Approaches of Evapotranspiration Estimation in Mixed Urban Vegetation: Field-Based, Remote Sensing-Based and Observational-Based Methods. *Remote Sensing*, 8(6), 492. Retrieved 2019-09-05, from <https://www.mdpi.com/2072-4292/8/6/492> doi: 10.3390/rs8060492
- Novick, K. A., Biederman, J. A., Desai, A. R., Litvak, M. E., Moore, D. J. P., Scott, R. L., & Torn, M. S. (2018, February). The AmeriFlux network: A coalition of the willing. *Agricultural and Forest Meteorology*, 249, 444–456. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0168192317303295> doi: 10.1016/j.agrformet.2017.10.009
- Oliveira, B. S., Moraes, E. C., Carrasco-Benavides, M., Bertani, G., & Mataveli, G. A. V. (2018, August). Improved Albedo Estimates Implemented in the

- METRIC Model for Modeling Energy Balance Fluxes and Evapotranspiration over Agricultural and Natural Areas in the Brazilian Cerrado. *Remote Sensing*, 10(8), 1181. Retrieved 2020-06-27, from <https://www.mdpi.com/2072-4292/10/8/1181> (Number: 8 Publisher: Multidisciplinary Digital Publishing Institute) doi: 10.3390/rs10081181
- Oliveira, P. T. S., Wendland, E., Nearing, M. A., Scott, R. L., Rosolem, R., & da Rocha, H. R. (2015, June). The water balance components of undisturbed tropical woodlands in the Brazilian cerrado. *Hydrology and Earth System Sciences*, 19(6), 2899–2910. Retrieved 2019-09-05, from <https://www.hydrol-earth-syst-sci.net/19/2899/2015/> doi: <https://doi.org/10.5194/hess-19-2899-2015>
- Olivera-Guerra, L., Mattar, C., Merlin, O., Durán-Alarcón, C., Santamaría-Artigas, A., & Fuster, R. (2017, June). An operational method for the disaggregation of land surface temperature to estimate actual evapotranspiration in the arid region of Chile. *ISPRS Journal of Photogrammetry and Remote Sensing*, 128, 170–181. Retrieved 2020-09-04, from <http://www.sciencedirect.com/science/article/pii/S0924271616303690> doi: 10.1016/j.isprsjprs.2017.03.014
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., ... Kassem, K. R. (2001, November). Terrestrial Ecoregions of the World: A New Map of Life on Earth A new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity. *BioScience*, 51(11), 933–938. Retrieved 2020-05-06, from <https://academic.oup.com/bioscience/article/51/11/933/227116> (Publisher: Oxford Academic) doi: 10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- Paca, V. H. d. M., Espinoza-Dávalos, G. E., Hessels, T. M., Moreira, D. M., Comair, G. F., & Bastiaanssen, W. G. M. (2019, February). The spatial variability of actual evapotranspiration across the Amazon River Basin based on remote sensing products validated with flux towers. *Ecological Processes*, 8(1), 6. Retrieved 2020-03-23, from <https://doi.org/10.1186/s13717-019-0158-8> doi: 10.1186/s13717-019-0158-8
- Paiva, C. M., França, G. B., Liu, W. T. H., & Filho, O. C. R. (2011, March). A comparison of experimental energy balance components data and SEBAL model results in Dourados, Brazil. *International Journal of Remote Sensing*, 32(6), 1731–1745. Retrieved 2021-04-15, from <https://doi.org/10.1080/01431161003623425> (Publisher: Taylor & Francis eprint: <https://doi.org/10.1080/01431161003623425>) doi: 10.1080/01431161003623425
- Paschalis, A., Fatichi, S., Pappas, C., & Or, D. (2018, October). Covariation of vegetation and climate constrains present and future T/ET variability. *Environmental Research Letters*, 13(10), 104012. Retrieved 2021-03-22, from <https://doi.org/10.1088/1748-9326/aae267> (Publisher: IOP Publishing) doi: 10.1088/1748-9326/aae267
- Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W., ... Papale, D. (2020, July). The FLUXNET2015 dataset and the ONE-Flux processing pipeline for eddy covariance data. *Scientific Data*, 7(1), 225. Retrieved 2021-04-21, from <https://www.nature.com/articles/s41597-020-0534-3> (Number: 1 Publisher: Nature Publishing Group) doi: 10.1038/s41597-020-0534-3
- Paul-Limoges, E., Wolf, S., Schneider, F. D., Longo, M., Moorcroft, P., Gharun, M., & Damm, A. (2020, January). Partitioning evapotranspiration with concurrent eddy covariance measurements in a mixed forest. *Agricultural and Forest Meteorology*, 280, 107786. Retrieved 2021-04-11, from <https://www.sciencedirect.com/science/article/pii/S0168192319304022> doi: 10.1016/j.agrformet.2019.107786

- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007, October). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences*, 11(5), 1633–1644. Retrieved 2020-05-06, from <https://www.hydrology-earth-syst-sci.net/11/1633/2007/> (Publisher: Copernicus GmbH) doi: <https://doi.org/10.5194/hess-11-1633-2007>
- Poblete-Echeverría, C., & Ortega-Farías, S. (2012, November). Calibration and validation of a remote sensing algorithm to estimate energy balance components and daily actual evapotranspiration over a drip-irrigated Merlot vineyard. *Irrigation Science*, 30(6), 537–553. Retrieved 2021-04-15, from <https://doi.org/10.1007/s00271-012-0381-x> doi: 10.1007/s00271-012-0381-x
- Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M., Mooney, H. A., & Klooster, S. A. (1993). Terrestrial ecosystem production: A process model based on global satellite and surface data. *Global Biogeochemical Cycles*, 7(4), 811–841. Retrieved 2019-12-20, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93GB02725> doi: 10.1029/93GB02725
- Priestley, C. H. B., & Taylor, R. J. (1972, February). On the Assessment of Surface Heat Flux and Evaporation Using Large-Scale Parameters. *Monthly Weather Review*, 100(2), 81–92. Retrieved 2021-04-15, from https://journals.ametsoc.org/view/journals/mwre/100/2/1520-0493_1972_100_0081_otaosh_2_3_co_2.xml (Publisher: American Meteorological Society Section: Monthly Weather Review) doi: 10.1175/1520-0493(1972)100<0081:OTAOSH>2.3.CO;2
- Rahimzadegan, M., & Janani, A. (2019, May). Estimating evapotranspiration of pistachio crop based on SEBAL algorithm using Landsat 8 satellite imagery. *Agricultural Water Management*, 217, 383–390. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S037837741831179X> doi: 10.1016/j.agwat.2019.03.018
- Restrepo-Coupe, N., da Rocha, H. R., Hutry, L. R., da Araujo, A. C., Borma, L. S., Christoffersen, B., ... Saleska, S. R. (2013, December). What drives the seasonality of photosynthesis across the Amazon basin? A cross-site analysis of eddy flux tower measurements from the Brasil flux network. *Agricultural and Forest Meteorology*, 182–183, 128–144. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0168192313001184> doi: 10.1016/j.agrformet.2013.04.031
- Rocha, H. R. d., Manzi, A. O., Cabral, O. M., Miller, S. D., Goulden, M. L., Saleska, S. R., ... Maia, J. F. (2009). Patterns of water and heat flux across a biome gradient from tropical forest to savanna in Brazil. *Journal of Geophysical Research: Biogeosciences*, 114(G1). Retrieved 2019-08-30, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000640> doi: 10.1029/2007JG000640
- Rodrigues, T. R., Vourlitis, G. L., Lobo, F. d. A., Oliveira, R. G. d., & Nogueira, J. d. S. (2014). Seasonal variation in energy balance and canopy conductance for a tropical savanna ecosystem of south central Mato Grosso, Brazil. *Journal of Geophysical Research: Biogeosciences*, 119(1), 1–13. Retrieved 2020-05-12, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JG002472> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2013JG002472>) doi: 10.1002/2013JG002472
- Rodrigues, T. R., Vourlitis, G. L., Lobo, F. d. A., Santanna, F. B., de Arruda, P. H. Z., & Nogueira, J. d. S. (2016, March). Modeling canopy conductance under contrasting seasonal conditions for a tropical savanna ecosystem of south central Mato Grosso, Brazil. *Agricultural and Forest Meteorology*, 218–219, 218–229. Retrieved 2020-08-31, from <http://www.sciencedirect.com/science/article/pii/S0168192315300216> doi:

- 10.1016/j.agrformet.2015.12.060
- Roerink, G. J., Su, Z., & Menenti, M. (2000, January). S-SEBI: A simple remote sensing algorithm to estimate the surface energy balance. *Physics and Chemistry of the Earth, Part B: Hydrology, Oceans and Atmosphere*, 25(2), 147–157. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S1464190999001288> doi: 10.1016/S1464-1909(99)00128-8
- Ruhoff, A. L., Paz, A. R., Aragao, L. E. O. C., Mu, Q., Malhi, Y., Collischonn, W., ... Running, S. W. (2013, November). Assessment of the MODIS global evapotranspiration algorithm using eddy covariance measurements and hydrological modelling in the Rio Grande basin. *Hydrological Sciences Journal*, 58(8), 1658–1676. Retrieved 2017-09-21, from <http://dx.doi.org/10.1080/02626667.2013.837578> doi: 10.1080/02626667.2013.837578
- Running, S. W., Mu, Q., Zhao, M., & Moreno, A. (2019, January). *User's Guide: MODIS Global Terrestrial Evapotranspiration (ET) Product, Version 2.0*. Retrieved from <https://modis-land.gsfc.nasa.gov/pdf/MOD16UsersGuideV2.022019.pdf>
- Rwasoka, D. T., Gumindoga, W., & Gwenzi, J. (2011, January). Estimation of actual evapotranspiration using the Surface Energy Balance System (SEBS) algorithm in the Upper Manyame catchment in Zimbabwe. *Physics and Chemistry of the Earth, Parts A/B/C*, 36(14), 736–746. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S1474706511001513> doi: 10.1016/j.pce.2011.07.035
- Saleska, S. R., Da Rocha, H. R., Huete, A. R., Nobre, A. D., Artaxo, P. E., & Shimabukuro, Y. E. (2013, July). LBA-ECO CD-32 Flux Tower Network Data Compilation, Brazilian Amazon: 1999-2006. *ORNL DAAC*. Retrieved 2020-05-06, from https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=1174 doi: <https://doi.org/10.3334/ORNLDAAC/1174>
- Schotanus, P., Nieuwstadt, F., & De Bruin, H. (1983, May). Temperature measurement with a sonic anemometer and its application to heat and moisture fluxes. *Boundary-Layer Meteorology*, 26(1), 81–93. Retrieved 2021-10-01, from <https://doi.org/10.1007/BF00164332> doi: 10.1007/BF00164332
- Seddon, A. W. R., Macias-Fauria, M., Long, P. R., Benz, D., & Willis, K. J. (2016, March). Sensitivity of global terrestrial ecosystems to climate variability. *Nature*, 531(7593), 229–232. Retrieved 2020-07-08, from <https://www.nature.com/articles/nature16986> (Number: 7593 Publisher: Nature Publishing Group) doi: 10.1038/nature16986
- Senay, G. B., Budde, M., Verdin, J. P., & Melesse, A. M. (2007, June). A Coupled Remote Sensing and Simplified Surface Energy Balance Approach to Estimate Actual Evapotranspiration from Irrigated Fields. *Sensors*, 7(6), 979–1000. Retrieved 2021-04-15, from <https://www.mdpi.com/1424-8220/7/6/979> (Number: 6 Publisher: Molecular Diversity Preservation International) doi: 10.3390/s7060979
- Shuttleworth, W. J., & Pereira, H. C. (1988, April). Evaporation from Amazonian rainforest. *Proceedings of the Royal Society of London. Series B. Biological Sciences*, 233(1272), 321–346. Retrieved 2021-04-08, from <https://royalsocietypublishing.org/doi/10.1098/rspb.1988.0024> (Publisher: Royal Society) doi: 10.1098/rspb.1988.0024
- Silva, P. F. d., Lima, J. R. d. S., Antonino, A. C. D., Souza, R., Souza, E. S. d., Silva, J. R. I., & Alves, E. M. (2017, December). Seasonal patterns of carbon dioxide, water and energy fluxes over the Caatinga and grassland in the semi-arid region of Brazil. *Journal of Arid Environments*, 147, 71–82. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0140196317301672> doi: 10.1016/j.jaridenv.2017.09.003
- Sánchez, J. M., López-Urrea, R., Valentín, F., Caselles, V., & Galve, J. M. (2019). Lysimeter assessment of the simplified two-source energy balance model

- and eddy covariance system to estimate vineyard evapotranspiration. *Agricultural and Forest Meteorology*, 274, 172–183. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0168192319301777> doi: <https://doi.org/10.1016/j.agrformet.2019.05.006>
- Souza, L. S. B. d., Moura, M. S. B. d., Sediya, G. C., Silva, T. G. F. d., Souza, L. S. B. d., Moura, M. S. B. d., ... Silva, T. G. F. d. (2015, August). Balanço de energia e controle biofísico da evapotranspiração na Caatinga em condições de seca intensa. *Pesquisa Agropecuária Brasileira*, 50(8), 627–636. Retrieved 2020-08-31, from http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-204X2015000800627&lng=en&nrm=iso&tlng=pt (Publisher: Embrapa Informação Tecnológica) doi: 10.1590/S0100-204X2015000800001
- Souza, V. d. A., Roberti, D. R., Ruhoff, A. L., Zimmer, T., Adamatti, D. S., Gonçalves, L. G. G. d., ... Moraes, O. L. L. d. (2019, September). Evaluation of MOD16 Algorithm over Irrigated Rice Paddy Using Flux Tower Measurements in Southern Brazil. *Water*, 11(9), 1911. Retrieved 2020-06-23, from <https://www.mdpi.com/2073-4441/11/9/1911> (Number: 9 Publisher: Multidisciplinary Digital Publishing Institute) doi: 10.3390/w11091911
- Stoy, P. C., El-Madany, T. S., Fisher, J. B., Gentine, P., Gerken, T., Good, S. P., ... Wolf, S. (2019, October). Reviews and syntheses: Turning the challenges of partitioning ecosystem evaporation and transpiration into opportunities. *Biogeosciences*, 16(19), 3747–3775. Retrieved 2021-03-23, from <https://bg.copernicus.org/articles/16/3747/2019/> (Publisher: Copernicus GmbH) doi: 10.5194/bg-16-3747-2019
- Stoy, P. C., Mauder, M., Foken, T., Marcolla, B., Boegh, E., Ibrom, A., ... Varlagin, A. (2013, April). A data-driven analysis of energy balance closure across FLUXNET research sites: The role of landscape scale heterogeneity. *Agricultural and Forest Meteorology*, 171–172, 137–152. Retrieved 2021-02-19, from <https://www.sciencedirect.com/science/article/pii/S0168192312003413> doi: 10.1016/j.agrformet.2012.11.004
- Su, Z. (2002, February). The Surface Energy Balance System (SEBS) for estimation of turbulent heat fluxes. *Hydrology and Earth System Sciences*, 6(1), 85–100. Retrieved 2021-04-15, from <https://hess.copernicus.org/articles/6/85/2002/> (Publisher: Copernicus GmbH) doi: 10.5194/hess-6-85-2002
- Sun, X., Wilcox, B. P., & Zou, C. B. (2019, September). Evapotranspiration partitioning in dryland ecosystems: A global meta-analysis of in situ studies. *Journal of Hydrology*, 576, 123–136. Retrieved 2021-04-08, from <https://www.sciencedirect.com/science/article/pii/S0022169419305736> doi: 10.1016/j.jhydrol.2019.06.022
- Sutanto, S. J., Wenninger, J., Coenders-Gerrits, A. M. J., & Uhlenbrook, S. (2012, August). Partitioning of evaporation into transpiration, soil evaporation and interception: a comparison between isotope measurements and a HYDRUS-1D model. *Hydrology and Earth System Sciences*, 16(8), 2605–2616. Retrieved 2021-04-11, from <https://hess.copernicus.org/articles/16/2605/2012/> (Publisher: Copernicus GmbH) doi: 10.5194/hess-16-2605-2012
- Talsma, C. J., Good, S. P., Jimenez, C., Martens, B., Fisher, J. B., Miralles, D. G., ... Purdy, A. J. (2018, October). Partitioning of evapotranspiration in remote sensing-based models. *Agricultural and Forest Meteorology*, 260–261, 131–143. Retrieved 2020-09-01, from <http://www.sciencedirect.com/science/article/pii/S016819231830162X> doi: 10.1016/j.agrformet.2018.05.010
- Talsma, C. J., Good, S. P., Miralles, D. G., Fisher, J. B., Martens, B., Jimenez, C., & Purdy, A. J. (2018, October). Sensitivity of Evapotranspiration Components in Remote Sensing-Based Models. *Remote Sensing*, 10(10), 1601. Retrieved 2020-06-29, from <https://www.mdpi.com/2072-4292/10/10/1601> (Number: 10 Publisher: Multidisciplinary Digital Publishing Institute) doi:

- 10.3390/rs10101601
- Teixeira, A. H. d. C., Bastiaanssen, W. G. M., Ahmad, M. D., & Bos, M. G. (2009, March). Reviewing SEBAL input parameters for assessing evapotranspiration and water productivity for the Low-Middle São Francisco River basin, Brazil: Part A: Calibration and validation. *Agricultural and Forest Meteorology*, 149(3), 462–476. Retrieved 2020-06-26, from <http://www.sciencedirect.com/science/article/pii/S0168192308002566> doi: 10.1016/j.agrformet.2008.09.016
- Teixeira, A. H. d. C., Scherer-Warren, M., Hernandez, F. B. T., Andrade, R. G., & Leivas, J. F. (2013, November). Large-Scale Water Productivity Assessments with MODIS Images in a Changing Semi-Arid Environment: A Brazilian Case Study. *Remote Sensing*, 5(11), 5783–5804. Retrieved 2021-04-15, from <https://www.mdpi.com/2072-4292/5/11/5783> (Number: 11 Publisher: Multidisciplinary Digital Publishing Institute) doi: 10.3390/rs5115783
- Tetens, O. (1930). *Über einige meteorologische Begriffe*. Friedrich Vieweg & Sohn Akt.- Gesellschaft. Retrieved from <https://books.google.com.br/books?id=ey5UtAEACAAJ>
- Thornton, P. E. (1998). *Regional ecosystem simulation: Combining surface- and satellite-based observations to study linkages between terrestrial energy and mass budgets* (phdthesis). The University of Montana, Missoula, MT,.
- Tong, X., Zhang, J., Meng, P., Li, J., & Zheng, N. (2017, February). Environmental controls of evapotranspiration in a mixed plantation in North China. *International Journal of Biometeorology*, 61(2), 227–238. Retrieved 2021-04-15, from <https://doi.org/10.1007/s00484-016-1205-0> doi: 10.1007/s00484-016-1205-0
- Tonti, N. E., Gassmann, M. I., & Pérez, C. F. (2018, December). First results of energy and mass exchange in a salt marsh on southeastern South America. *Agricultural and Forest Meteorology*, 263, 59–68. Retrieved 2021-03-11, from <https://www.sciencedirect.com/science/article/pii/S0168192318302612> doi: 10.1016/j.agrformet.2018.08.001
- Trabucco, A., & Zomer, R. (2019, January). *Global Aridity Index and Potential Evapotranspiration (ET0) Climate Database v2*. Retrieved 2020-05-06, from https://figshare.com/articles/Global_Aridity_Index_and_Potential_Evapotranspiration_ET0_Climate_Database_v2/7504448 doi: 10.6084/m9.figshare.7504448.v3
- Trajano, E. (2019, January). Chapter 20 - Biodiversity in South America. In W. B. White, D. C. Culver, & T. Pipan (Eds.), *Encyclopedia of Caves (Third Edition)* (pp. 177–186). Academic Press. Retrieved 2020-07-08, from <http://www.sciencedirect.com/science/article/pii/B9780128141243000194> doi: 10.1016/B978-0-12-814124-3.00019-4
- Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers, T. P., ... Wesely, M. L. (2000, June). Correcting eddy-covariance flux underestimates over a grassland. *Agricultural and Forest Meteorology*, 103(3), 279–300. Retrieved 2017-09-21, from <http://www.sciencedirect.com/science/article/pii/S0168192300001234> doi: 10.1016/S0168-1923(00)00123-4
- Valente, F., David, J. S., & Gash, J. H. C. (1997, March). Modelling interception loss for two sparse eucalypt and pine forests in central Portugal using reformulated Rutter and Gash analytical models. *Journal of Hydrology*, 190(1), 141–162. Retrieved 2020-07-28, from <http://www.sciencedirect.com/science/article/pii/S0022169496030661> doi: 10.1016/S0022-1694(96)03066-1
- Valle Júnior, L. C. G., Ventura, T. M., Gomes, R. S. R., de S. Nogueira, J., de A. Lobo, F., Vourlitis, G. L., & Rodrigues, T. R. (2020, April). Comparative assessment of modelled and empirical reference evapotranspiration methods for a brazilian savanna. *Agricultural Water Management*, 232, 106040. Retrieved 2020-08-31, from <http://www.sciencedirect.com/science/article/pii/>

- S0378377419314489 doi: 10.1016/j.agwat.2020.106040
- Verhoef, A., & Allen, S. J. (2000, March). A SVAT scheme describing energy and CO₂ fluxes for multi-component vegetation: calibration and test for a Sahelian savannah. *Ecological Modelling*, 127(2), 245–267. Retrieved 2021-04-10, from <https://www.sciencedirect.com/science/article/pii/S0304380099002136> doi: 10.1016/S0304-3800(99)00213-6
- Verhoef, A., & Campbell, C. L. (2006). Evaporation Measurement. In *Encyclopedia of Hydrological Sciences*. American Cancer Society. Retrieved 2021-02-19, from <https://onlinelibrary.wiley.com/doi/abs/10.1002/0470848944.hsa043> (Section: 40 _eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/0470848944.hsa043>) doi: 10.1002/0470848944.hsa043
- Villarreal, S., & Vargas, R. (2021). Representativeness of FLUXNET sites across Latin America. *Journal of Geophysical Research: Biogeosciences*, 126(3), e2020JG006090. Retrieved 2021-02-19, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JG006090> doi: <https://doi.org/10.1029/2020JG006090>
- Vinukollu, R. K., Wood, E. F., Ferguson, C. R., & Fisher, J. B. (2011, March). Global estimates of evapotranspiration for climate studies using multi-sensor remote sensing data: Evaluation of three process-based approaches. *Remote Sensing of Environment*, 115(3), 801–823. Retrieved 2017-09-21, from <http://www.sciencedirect.com/science/article/pii/S0034425710003251> doi: 10.1016/j.rse.2010.11.006
- von Randow, C., Manzi, A. O., Kruijt, B., de Oliveira, P. J., Zanchi, F. B., Silva, R. L., ... Kabat, P. (2004, June). Comparative measurements and seasonal variations in energy and carbon exchange over forest and pasture in South West Amazonia. *Theoretical and Applied Climatology*, 78(1), 5–26. Retrieved 2020-03-19, from <https://doi.org/10.1007/s00704-004-0041-z> doi: 10.1007/s00704-004-0041-z
- Vourlitis, G. L., Nogueira, J. d. S., Lobo, F. d. A., Sendall, K. M., Paulo, S. R. d., Dias, C. A. A., ... Andrade, N. L. R. d. (2008). Energy balance and canopy conductance of a tropical semi-deciduous forest of the southern Amazon Basin. *Water Resources Research*, 44(3). Retrieved 2020-05-12, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006WR005526> (_eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006WR005526>) doi: 10.1029/2006WR005526
- Wang, X., Huo, Z., Shukla, M. K., Wang, X., Guo, P., Xu, X., & Huang, G. (2020, February). Energy fluxes and evapotranspiration over irrigated maize field in an arid area with shallow groundwater. *Agricultural Water Management*, 228, 105922. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0378377419311023> doi: 10.1016/j.agwat.2019.105922
- Wang, Z., Schaaf, C. B., Strahler, A. H., Chopping, M. J., Román, M. O., Shuai, Y., ... Fitzjarrald, D. R. (2014, January). Evaluation of MODIS albedo product (MCD43A) over grassland, agriculture and forest surface types during dormant and snow-covered periods. *Remote Sensing of Environment*, 140, 60–77. Retrieved 2019-06-14, from <http://www.sciencedirect.com/science/article/pii/S0034425713002836> doi: 10.1016/j.rse.2013.08.025
- Wardlow, B. D., Egbert, S. L., & Kastens, J. H. (2007, June). Analysis of time-series MODIS 250 m vegetation index data for crop classification in the U.S. Central Great Plains. *Remote Sensing of Environment*, 108(3), 290–310. Retrieved 2020-06-22, from <http://www.sciencedirect.com/science/article/pii/S0034425706004949> doi: 10.1016/j.rse.2006.11.021
- Webb, E., Pearman, G., & Leuning, R. (1980). Correction of flux measurements for

- density effects due to heat and water vapour transfer. *Quarterly Journal of the Royal Meteorological Society*, 106, 85–100.
- Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S., & Lee, X. (2017). Revisiting the contribution of transpiration to global terrestrial evapotranspiration. *Geophysical Research Letters*, 44(6), 2792–2801. Retrieved 2021-03-22, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL072235> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2016GL072235>) doi: <https://doi.org/10.1002/2016GL072235>
- Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., ... Verma, S. (2002, December). Energy balance closure at FLUXNET sites. *Agricultural and Forest Meteorology*, 113(1), 223–243. Retrieved 2020-04-21, from <http://www.sciencedirect.com/science/article/pii/S0168192302001090> doi: 10.1016/S0168-1923(02)00109-0
- Xu, T., Guo, Z., Xia, Y., Ferreira, V. G., Liu, S., Wang, K., ... Zhao, C. (2019, November). Evaluation of twelve evapotranspiration products from machine learning, remote sensing and land surface models over conterminous United States. *Journal of Hydrology*, 578, 124105. Retrieved 2021-04-15, from <https://www.sciencedirect.com/science/article/pii/S0022169419308406> doi: 10.1016/j.jhydrol.2019.124105
- Yang, Y., & Roderick, M. L. (2019). Radiation, surface temperature and evaporation over wet surfaces. *Quarterly Journal of the Royal Meteorological Society*, 145(720), 1118–1129. Retrieved from <https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3481> doi: <https://doi.org/10.1002/qj.3481>
- Zhang, J., Zhang, S., Zhang, W., Liu, B., Gong, C., Jiang, M., ... Sheng, L. (2018, August). Partitioning daily evapotranspiration from a marsh wetland using stable isotopes in a semiarid region. *Hydrology Research*, 49(4), 1005–1015. Retrieved 2021-04-11, from <https://doi.org/10.2166/nh.2017.005> doi: 10.2166/nh.2017.005
- Zhang, Y., Chiew, F. H. S., Peña-Arancibia, J., Sun, F., Li, H., & Leuning, R. (2017). Global variation of transpiration and soil evaporation and the role of their major climate drivers. *Journal of Geophysical Research: Atmospheres*, 122(13), 6868–6881. Retrieved 2020-07-11, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027025> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JD027025>) doi: 10.1002/2017JD027025
- Zhou, S., Yu, B., Zhang, Y., Huang, Y., & Wang, G. (2016). Partitioning evapotranspiration based on the concept of underlying water use efficiency. *Water Resources Research*, 52(2), 1160–1175. Retrieved 2020-07-11, from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015WR017766> (eprint: <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015WR017766>) doi: 10.1002/2015WR017766