

Are remote sensing evapotranspiration models reliable across South American ecoregions?

Article

Accepted Version

Melo, D. C. D., Anache, J. A. A., Borges, V. P., Miralles, D. G., Martens, B., Fisher, J. B., Nobrega, R. L. B., Moreno, A., Cabral, O. M. R., Rodrigues, T. R., Bezerra, B., Silva, C. M. S., Meira Neto, A. A., Moura, M. S. B., Marques, T. V., Campos, S., Nogueira, J. S., Rosolem, R., Souza, R. M. S., Antonino, A. C. D., Holl, D., Galleguillos, M., Perez-Quesada, J. F., Verhoef, A., Kutzbach, L., Lima, J. R. S., Souza, E. S., Gassman, M. I., Perez, C. F., Tonti, N., Posse, G., Rains, D., Oliveira, P. T. S. and Wendland, E. (2021) Are remote sensing evapotranspiration models reliable across South American ecoregions? *Water Resources Research*, 57 (11). e2020WR028752. ISSN 0043-1397 doi: <https://doi.org/10.1029/2020WR028752> Available at <https://centaur.reading.ac.uk/101236/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1029/2020WR028752>

Publisher: American Geophysical Union

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Are remote sensing evapotranspiration models reliable across South American ecoregions?

D. C. D. Melo¹, J. A. A. Anache², V. P. Borges¹, D. G. Miralles³, B. Martens³, J. B. Fisher⁴, R. L. B. Nóbrega⁵, A. Moreno⁶, O. M. R. Cabral⁷, T. R. Rodrigues², B. Bezerra^{8,9}, C. M. S. Silva^{9,10}, A. A. Meira Neto¹⁰, M. S. B. Moura¹¹, T. V. Marques⁹, S. Campos⁹, J. S. Nogueira¹², R. Rosolem¹³, R. Souza¹⁴, A. C. D. Antonino¹⁵, D. Holl¹⁶, M. Galleguillos¹⁷, J. F. Perez-Quezada^{17,18}, A. Verhoef⁹, L. Kutzbach¹⁶, J. R. S. Lima²⁰, E. S. Souza²¹, M. I. Gassman^{22,23}, C. F. Perez^{22,23}, N. Tonti²², G. Posse²⁴, D. Rains³, and P. T. S. Oliveira², E. Wendland²⁵

¹Federal University of Paraíba, Areia, PB, Brazil

²Federal University of Mato Grosso do Sul, Campo Grande, MS, Brazil

³Hydro-Climate Extremes Lab (H-CEL), Ghent University, Coupure Links 653, 9000 Ghent, Belgium

⁴Schmid College of Science and Technology, Chapman University, Orange, CA, USA

⁵Imperial College London, Department of Life Sciences, Silwood Park Campus, Buckhurst Road, Ascot, SL5 7PY, UK

⁶Numerical Terradynamic Simulation Group, University of Montana, Missoula, MT, USA

⁷Brazilian Agricultural Research Corporation, Embrapa Meio Ambiente, Jaguariúna, SP, Brazil

⁸Department of Atmospheric and Climate Sciences, Federal University of Rio Grande do Norte, Natal, RN, Brazil

⁹Climate Sciences Graduate Program, Federal University of Rio Grande do Norte, Natal, RN, Brazil

¹⁰Department of Hydrology and Atmospheric Sciences, The University of Arizona

¹¹Brazilian Agricultural Research Corporation – Embrapa Tropical Semi-arid, Petrolina, PE, Brazil

¹²Federal University of Mato Grosso, Cuiabá, MT, Brazil

¹³University of Bristol, BS7 8PD, UK

¹⁴Department of Biological and Agricultural Engineering, Texas A&M University, College Station, TX, USA

¹⁵Department of Nuclear Energy, Federal University of Pernambuco, Recife, PE, Brazil

¹⁶Center for Earth System Research and Sustainability (CEN), Universität Hamburg, Hamburg, Germany

¹⁷Department of Environmental Science and Renewable Natural Resources, University of Chile, Santiago, Chile

¹⁸Institute of Ecology and Biodiversity, Santiago, Chile

¹⁹Department of Geography and Environmental Science, The University of Reading, Reading, UK

²⁰Federal University of the Agreste of Pernambuco, Garanhuns, PE, Brazil

²¹Federal Rural University of Pernambuco, Serra Talhada, PE, Brazil

²²Department of Atmospheric and Ocean Sciences, FCEN - UBA. Buenos Aires, Argentina

²³National Council for Scientific and Technical Research, (CONICET), Argentina

²⁴Instituto de Clima y Agua. Instituto Nacional de Tecnología Agropecuaria (INTA), Buenos Aires, Argentina

²⁵Department of Hydraulics and Sanitary Engineering, University of São Paulo, São Carlos, SP, Brazil

Key Points:

- Four remote sensing *ET* models were evaluated using data from 25 flux towers in South America
- GLEAM and PT-JPL provided a significantly greater number of daily outputs
- Comparisons with flux tower-based *ET* showed that GLEAM and PT-JPL produced higher correlations whereas *RMSE* was similar for all models
- No model outperformed the other for all biomes, climates or land uses

Corresponding author: Davi Diniz Melo, melo.dcd@gmail.com

Abstract

Many remote sensing-based evapotranspiration (RSBET) algorithms have been proposed in the past decades and evaluated using flux tower data, mainly over North America and Europe. Model evaluation across South America has been done locally or using only a single algorithm at a time. Here, we provide the first evaluation of multiple RSBET models, at a daily scale, across a wide variety of biomes, climate zones, and land uses in South America. We used meteorological data from 25 flux towers to force four RSBET models: Priestley–Taylor Jet Propulsion Laboratory (PT-JPL), Global Land Evaporation Amsterdam Model (GLEAM), Penman–Monteith Mu model (PM-MOD), and Penman–Monteith Nagler model (PM-VI). ET was predicted satisfactorily by all four models, with correlations consistently higher ($R^2 > 0.6$) for GLEAM and PT-JPL, and PM-MOD and PM-VI presenting overall better responses in terms of percent bias ($-10 < PBIAS < 10\%$). As for PM-VI, this outcome is expected, given that the model requires calibration with local data. Model skill seems to be unrelated to land-use but instead presented some dependency on biome and climate, with the models producing the best results for wet to moderately wet environments. Our findings show the suitability of individual models for a number of combinations of land cover types, biomes, and climates. At the same time, no model outperformed the other for all conditions, which emphasizes the need for adapting individual algorithms to take into account intrinsic characteristics of climates and ecosystems in South America.

1 Introduction

Land evaporation, or evapotranspiration (ET), is the phenomenon by which water is converted from a liquid into its vapor phase over land. It plays a significant role in the modulation of global climate feedbacks being a key driver of the Earth’s carbon, energy, and water cycles at local, regional, and global scales (Cao et al., 2010; Tong et al., 2017; Khosa et al., 2019; Valle Júnior et al., 2020; de Oliveira et al., 2021). *In situ* ET measurements can be obtained from micro-meteorological methods (e.g., eddy covariance, scintillometry, or Bowen ratio method) and those derived from the soil water balance (e.g., directly using lysimeters, or from changes in profile soil moisture content obtained gravimetrically, from neutron probes, or capacitance-based soil water monitoring equipment). Besides, plant physiological techniques such as sap flow methods, provide direct estimates of transpiration (Verhoef & Campbell, 2006; Allen et al., 2011; Fisher et al., 2011), but only the micro-meteorological methods provide ET data at the field to landscape (e.g., scintillometry) scale. Over the past three decades, eddy covariance (EC) systems have become the state-of-the-art and standard *in situ* method to quantify land surface energy and mass fluxes for different types of ecosystems (Restrepo-Coupe et al., 2013; Rodrigues et al., 2016; Campos et al., 2019; X. Wang et al., 2020). However, these techniques estimate fluxes for areas of relatively limited spatial dimensions ($\sim 1 \text{ km}^2$) depending on the heterogeneity of the landscape), and they are affected by specific local conditions, such as the occurrence of advection across sharp contrasts in vegetation and/or irrigation conditions, and those caused by topographic features, such as cold air drainage for sloping terrain (Allen et al., 2011; Rwasoka et al., 2011; Mutti et al., 2019; Rahimzadegan & Janani, 2019; Mauder et al., 2020).

During the 1990s and 2000s, remote sensing based ET (RSBET) algorithms, using information from visible, near-infrared, and thermal infrared bands, were developed, such as the Surface Energy Balance Algorithms for Land (SEBAL, (Bastiaanssen et al., 1998)), Simplified Surface Energy Balance Index (S-SEBI, Roerink et al. (2000)), Surface Balance Energy System (SEBS, Su (2002)), Simplified Surface Energy Balance (SSEB, Senay et al. (2007)), and Two-Source Energy Balance Model (TSEB, Norman et al. (1995); Kustas and Norman (1999)). These algorithms were developed for sub-regional applications, with a focus on irrigation or water resources management. Over South America, their predictive skills have been assessed quite extensively, mostly for irrigated crop-

land (Teixeira et al., 2009; Paiva et al., 2011; Poblete-Echeverría & Ortega-Farias, 2012; Bezerra et al., 2013, 2015; Olivera-Guerra et al., 2017; Lopes et al., 2019; Mutti et al., 2019). Studies show that these models perform well when compared to field observations of *ET* (Teixeira et al., 2009; Poblete-Echeverría & Ortega-Farias, 2012).

Since the late 2000s, algorithms such as PT-JPL (Fisher et al., 2008), PM-MOD (Mu et al., 2007, 2011), and GLEAM (Miralles et al., 2011; Martens et al., 2017) focused on the use of satellite-derived observations to create spatially coherent global *ET* estimates (Fisher et al., 2017). PT-JPL is at the core of the ECOSTRESS mission (Fisher et al., 2020), while PM-MOD is central to the global terrestrial MODIS *ET* product (MOD16). GLEAM is used for the annual State of the Climate report since 2015 (Blunden & Arndt, 2020).

Using flux tower data, previous studies conducted in South America evaluated GLEAM and MOD16 (Ruhoff et al., 2013; Moreira et al., 2019; Paca et al., 2019). However, these studies validated off-the-shelf *ET* datasets generated by these models, not the models themselves. Since such *ET* products are not produced using a common dataset of meteorological variables, a comparative evaluation cannot be made in terms of model structure. Rather, different model skills would be partially linked with the quality of the inputs. A multi-site tropical study, over several continents, validating the PT-JPL model at a regional scale on a monthly basis was presented by Fisher et al. (2009). However, to the best of our knowledge, studies assessing the daily predictive skills have only been conducted at the local scale (Teixeira et al., 2009, 2013; Miranda et al., 2017; B. S. Oliveira et al., 2018; V. d. A. Souza et al., 2019).

A major challenge to verify the results of these methods is the scarcity of ground-based observations, due to the uneven spatio-temporal distribution of the *ET* monitoring efforts. As a result, remote sensing *ET* methods are typically evaluated or parameterized using sites located only in North America, Europe (Ershadi et al., 2014; McCabe et al., 2016; Michel et al., 2016; Xu et al., 2019), Australia (Martens et al., 2016) and East Asia (Jang et al., 2013; Chang et al., 2018; Khan et al., 2018; Li et al., 2019). For example, Mu et al. (2011) proposed improvements to the PM-MOD *ET* global algorithm (Mu et al., 2007), based on comparisons with *ET* measurements from 46 AmeriFlux sites, 45 of them located in USA and Canada. Martens et al. (2017) evaluated the GLEAM algorithm with 91 worldwide FLUXNET sites; however, ~ 65 were located in the USA and in Europe. Therefore, these models might not satisfactorily represent *ET* in sparsely sampled regions with very different climate conditions such as South America, despite this continent representing ca. 12% of the total Earth's terrestrial area.

South America spans two hemispheres, and four major climate zones, from near the equator to sub-Antarctic regions, which makes it a geographically unique continent (Goymer, 2017; Trajano, 2019). Biomes in this continent range from tropical to deciduous forests, and contain ecoregions with high sensitivity to variability in water (e.g., the Caatinga and Humid Pampas) and energy availability (e.g. the Amazon, Valdivian temperate and Magellanic subpolar forests) (Seddon et al., 2016). Also, five out of six of the terrestrial biomes not included in satellite-based *ET* algorithm evaluations at a global scale are found in South America (see Section 2.1). Thus, the evaluation of RS-BET methods for South America offers an opportunity to reduce the current research gap, in particular at large spatial scales.

FLUXNET provides a common framework for the verification of *ET* algorithms. Nevertheless, the available sites in the FLUXNET2015 database are not evenly distributed around the world (Pastorello et al., 2020). Validating global models in South America is challenging, mainly because the data from $\sim 90\%$ of its FLUXNET registered sites are not readily available to the scientific community: less than 50% of South American AmeriFlux sites are available for direct access. Additionally, flux towers in woody savannas

151 and evergreen broad-leaf forests account for nearly 65% of all Latin American FLUXNET
152 sites while some of the biomes are not properly represented (Villarreal & Vargas, 2021).

153 The identification of scientific gaps and the proposed improvements are considered
154 a priority for the future development of *ET* assessment methods from remote sensing
155 (Fisher et al., 2017). Some of them include merging different *ET*-estimation methods,
156 and the identification of their sources of uncertainty (Fisher et al., 2017; Y. Zhang et al.,
157 2017; Paca et al., 2019). Indeed, despite the recent developments of remote sensing *ET*
158 methods, there are still challenges concerning the refinement of those algorithms to reme-
159 dy the lack of information on specific surface characteristics and fluxes of undersam-
160 pled climate zones and vegetation types. In this context, one of the main sources of un-
161 certainty in global satellite-based *ET* estimates are the fractional vegetation cover and
162 net radiation (Ferguson et al., 2010; Vinukollu et al., 2011; Badgley et al., 2015)

163 We evaluated the predictive skills of four satellite-based *ET* models, designed for
164 regional and continental scale applications, over South America. The main question we
165 seek to answer is whether such models can be applied consistently to reliably capture
166 *ET* in South America. Specific research questions include: (i) are the models capable
167 of correctly estimating *ET* and its components? (ii) are the models predictive skills af-
168 fected by climate, land cover type or biome?

169 2 Study area, data, and methods

170 2.1 South American biomes, flux tower-based *ET* and meteorological 171 data

172 The study area encompasses five biomes (Table S1 in the Supporting Material –
173 SM): Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Flooded Grasslands &
174 Savannas (FGS); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS);
175 Tropical & Subtropical Dry Broadleaf Forests (TSDBF) and Temperate Broadleaf & Mixed
176 Forests (TBMF) (Olson et al., 2001).

177 We used daily meteorological data from 25 flux tower sites located across various
178 South American biomes and land cover types to verify the predictive skill of the selected
179 RSBET models (Figure 1a, Table S2 in SM). The time period considered for analysis was
180 determined by the available time-series for each site (Figure S1 in SM). Further infor-
181 mation about each biome is provided in SM. Ten sites are from FLUXNET (Pastorello
182 et al., 2020), AmeriFlux networks (Novick et al., 2018) and Large-Scale Biosphere-Atmosphere
183 Experiment in the Amazon (LBA) project (Saleska et al., 2013), while the remaining data
184 were obtained from the respective principal investigators. Concerning towers sites not
185 available in global networks, data handling included standard procedures to ensure qual-
186 ity data, including: detection of spikes caused by changes in the footprint or imprecise
187 measurements; delay correction of H₂O/CO₂ in relation to the vertical wind component;
188 correction of coordinates (2D rotation); correction of spectral loss; conversion of the buoy-
189 ancy flux to sensible heat flux, known as SND-corrections (Schotanus et al., 1983); sonic
190 virtual temperature correction; corrections for flux density fluctuations, known as WPL
191 corrections (Webb. et al., 1980); incorporated frequency response correction. Addition-
192 ally, we performed due corrections with respect to reduction of wind velocity or turbu-
193 lence increase caused by the shadow of the tower and the sensor. Details about proce-
194 dures carried out for data processing and filtering to implement these corrections can
195 be found in Tonti et al. (2018); Holl et al. (2019); Campos et al. (2019); Cabral et al. (2020).
196 We also emphasize that those data have been widely used in previously scientific pub-
197 lications (Rocha et al., 2009; Cabral et al., 2010, 2011; Restrepo-Coupe et al., 2013; Ro-
198 driguez et al., 2016; Arruda et al., 2016; Silva et al., 2017; Marques et al., 2020). The
199 spatial patterns of mean annual precipitation (P), air temperature (T), and potential

200 evapotranspiration (PET) show that selected sites encompass a wide variety of climates
 201 (Figure 1b).

Figure 1. (a) Location of flux tower sites. Land cover types are indicated prior to tower names in the map: Croplands (CROP), Deciduous Needleleaf Forest (DNF), Evergreen Broadleaf Forest (EBF), Grasslands (GRA), Mixed Forest (MF), Permanent Wetland (PW), and Woody Savanna (WS); Biome types (Olson et al., 2001) are indicated by shades of green, yellow and blue on the map (see legend): Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Tropical & Subtropical Dry Broadleaf Forests (TSDBF); Temperate Broadleaf & Mixed Forests (TBMF); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS); Temperate Grasslands, Savannas & Shrublands (TGSS); Flooded Grasslands & Savannas (FGS); Montane Grasslands & Shrublands (MGS); Mediterranean Forests, Woodlands & Scrub (MFWS); Deserts & Xeric Shrublands (DXS); Climates across South America from selected representative sites are indicated by patterns on the map (see legend): Tropical savanna (Aw), Tropical monsoon (Am), Hot semi-arid (BSh), Cold semi-arid (BSk), Humid subtropical (Cfa), Temperate oceanic (Cfb), Dry-winter subtropical highland (Cwb), Polar Tundra (Td) (Peel et al., 2007). (b) Gridded annual average (AVG) and standard deviation (SD) for air temperature (T), rainfall (P), and potential evapotranspiration (PET) across South America and the monitored sites (Harris et al., 2020).

202 The closure of the energy budget is rarely observed with flux tower measurements
 203 (Wilson et al., 2002; Foken, 2008). Usually, the available energy flux ($Rn-G$) is greater
 204 than ($LE+H$), where Rn is the net radiation, G is the soil heat flux, LE is the latent
 205 heat flux and H is the sensible heat flux. The imbalances in the surface energy balance,
 206 reported here as an energy balance ratio, EBR (i.e. $(LE+H)/(Rn-G)$), range from
 207 0.73 to 1.16 (mean ~ 0.90) (Table S2, SM). It is paramount that only high-quality data
 208 were used to run and assess the models. We computed daily EBR for each site and ex-
 209 cluded days with $EBR \leq 0.75$ or ≥ 1.25 . Daily averages of meteorological variables were
 210 calculated from 30-min or hourly data only when at least 80% of the records per day were
 211 available. To obtain daytime and nighttime inputs for the MOD16 model (PM-MOD in
 212 this paper), we considered only days with a minimum of twenty 30-min daytime records
 213 and twenty during the night. As in Mu et al. (2011), the shortwave incoming radiation
 214 ($Rgs\downarrow$) was used to distinguish between daytime ($Rgs\downarrow > 10 \text{ W m}^{-2}$) and nighttime ($Rgs\downarrow <$
 215 10 W m^{-2}). Regarding the fluxes, we used quality-checked data that had not been gap-
 216 filled. Previous studies have shown that ET derived from the other energy balance fluxes,
 217 i.e. $LE = Rn - G - H$, can agree well with eddy covariance ET and lysimeter data
 218 (Amiro, 2009; Sánchez et al., 2019). Therefore, instead of using the EC-measured LE ,
 219 to represent ET , we derived LE from the equation above. Such validation approach (i.e.
 220 comparing model ET with EB-derived LE , ET_{EB}) has been adopted in previous stud-
 221 ies (Twine et al., 2000; Wilson et al., 2002; Stoy et al., 2013; Fisher et al., 2020). The
 222 results of using the eddy covariance ET (ET_{EC}) instead can be found in the SM (see
 223 Fig. S11-S13).

224 The quality control procedure described above was not adopted for the TF1, and
 225 TF2 towers (see Figure 1a). At those sites, horizontal advection plays an important role
 226 due to extreme weather variations throughout the year (Levy et al., 2020), such that the
 227 energy balance closure cannot be diagnosed by EBR, as described above. For instance,
 228 the SDF zone is known as an anticyclone pathway between the Pacific and Atlantic oceans,
 229 and TF1 and TF2 are located in the extreme southern parts of Patagonia, a region char-
 230 acterized by strong winds. Thus, for TF1 and TF2 sites, we used ET derived from mea-
 231 sured LE .

232

2.2 Remote sensing-based vegetation indices

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

The required vegetation index (VI) to run PT-JPL, PM-MOD and PM-VI is the Enhanced Vegetation Index (*EVI*). Vegetation Optical Depth (*VOD*) is used in GLEAM. *EVI* was derived from the 16-day Level 3 Global product of the MODerate Resolution Imaging Spectroradiometer (MODIS), aboard the Terra and Aqua satellites (Huete et al., 2002). We used both MODIS VI products, i.e. MOD13Q1 (Terra) and MYD13Q1 (Aqua), at 250-m resolution, to derive daily composites of *EVI*. *VOD* was extracted from the product described in Moesinger et al. (2020). Fisher et al. (2008) used the Soil Adjusted Vegetation Index (*SAVI*) instead of *EVI* because the former does not require the blue reflectance (0.45–0.51 μm), however, the authors recognize that both indices are very similar. As we are interested in assessing the *ET* models rather than the products resulting from different forcing data, we used *EVI* in Fisher’s model (PT-JPL). Leaf area index (*LAI*) and other vegetation-related variables (e.g., fraction of Absorbed Photosynthetically Active Radiation, f_{PAR}) are handled differently in each model. For example, in PT-JPL, *LAI* is obtained from total fractional vegetation cover, whereas in PM-MOD the 1-km MODIS *LAI* (MOD15) product is adopted. The original procedures to obtain those variables were not changed here. The following treatment was applied to the MODIS-derived data. “Good quality” pixels were selected, based on the quality assurance (QA) flags. Next, an autoregressive model was applied to fill in the gaps (Akaike, 1969). The gap-filling procedure was applied to gaps smaller than 16 days, while gaps of longer periods were excluded from the analysis. Finally, we implemented a temporal filter to improve the f_{PAR} and *LAI* time series to reproduce precisely all pre-processing steps of the standard PM-MOD algorithm (Mu et al., 2011). Filtering of f_{PAR} and *LAI* allowed for the correction of underestimated values (abrupt and unrealistic decreases in the time series) that mostly originate from cloud contamination effects which were not correctly identified in the quality control fields.

258

2.3 Summary of remote sensing-based *ET* models

259

2.3.1 GLEAM

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

GLEAM is a semi-empirical/process-based model that estimates the total evaporative flux and its components. In this study, version 3 of the algorithm is used (Martens et al., 2017). The main aspects of the model are described briefly, while for details we refer to Miralles et al. (2011) and Martens et al. (2017). The model calculates potential evaporation for four sub-grid land cover fractions: (1) open water, (2) low vegetation, (3) tall vegetation, and (4) bare soil using the Priestley and Taylor (1972) equation. For tall and low vegetation cover fractions, potential transpiration is constrained using an empirical evaporative stress factor which is calculated as a function of soil moisture at root-zone depth and microwave *VOD* as described in Martens et al. (2017). *VOD* (Vegetation Optical Depth) accounts for the attenuation of microwaves through vegetation and can be used as a proxy for vegetation phenology. Thus, *VOD* is a microwave parameter closely linked to vegetation water content (Liu et al., 2013) and in GLEAM it is used to represent phenological changes in vegetation. The soil moisture in the root-zone is calculated with a multi-layer water-balance model forced by precipitation and satellite surface soil moisture retrievals. For bare soil, the evaporative stress factor is calculated as a function of surface soil moisture only, whereas for open water evaporation, no stress factor is applied. For the tall vegetation cover fraction, rainfall interception loss is estimated with Gash’s analytical model (Gash, 1979; Miralles et al., 2010). The *ET* is then calculated as the sum of low and tall vegetation transpiration, rainfall interception loss, bare soil evaporation, and open-water evaporation with each weighted by the respective fraction.

281

2.3.2 PT-JPL

282

283

284

285

286

287

288

289

290

291

292

293

294

295

The global ET model proposed by Fisher et al. (2008) is based on the Priestley and Taylor equation for potential ET (PET), which is partitioned into actual plant transpiration, soil evaporation, and interception evaporation, i.e. $E_{trans} + E_{soil} + E_{int}$. To reduce potential ET to actual ET , the PT-JPL model applies ecophysiological constraints based on land surface information such as vegetation properties and humidity/water vapor pressure deficit (VPD). Fisher et al. (2008) used $NDVI$ and $SAVI$ as a proxy for plant physiological status. We used EVI because it provides a better indication of green vegetation cover than $NDVI$, as acknowledged by Fisher et al. (2008). The model partitions available energy flux using four plant-related constraints: LAI , green canopy fraction, plant temperature, and plant moisture. Similar to PM-MOD (see next subsection), vegetation cover, canopy wetness, etc. determine how the available energy flux is partitioned among the ET terms. A unique aspect related to the plant temperature constraint is the determination of an optimal temperature, T_{opt} (Potter et al., 1993), which corresponds to an optimal stomatal conductance. The latter co-determines E_{trans} .

296

2.3.3 PM-MOD

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

The MOD16 ET model (PM-MOD) is based on the Penman-Monteith equation to produce a daily global ET product summing up daytime and nighttime ET (Mu et al., 2011). In this model, total ET is partitioned into E_{soil} , E_{int} , and E_{trans} . To compute E_{soil} , PM-MOD uses potential soil evaporation and a soil moisture constraint function based on VPD and air relative humidity (RH) (Fisher et al., 2008). The evaporation of the water intercepted by the canopy, E_{int} , is calculated using the relevant equations from a revised version of the Biome-BGC model (Thornton, 1998). The PM-MOD assumes that E_{int} occurs when the vegetation is covered with water, i.e. when the water cover fraction (f_{wet}) > 0 , which is constrained by RH (Mu et al., 2011). In the PM-MOD model f_{wet} is calculated as in the PT-JPL model: f_{wet} is set to 0 if $RH < 70\%$ and $f_{wet} = (RH/100)^4$ if $70 < RH < 100\%$ (Running et al., 2019). The PM-MOD model is designed to allow E_{trans} to occur during daytime and nighttime, by adding constraints to stomatal conductance for VPD and minimum air temperature, and ignoring constraints relating to high air temperature (Running et al., 2019). The partitioning of available energy flux into soil or interception evaporation is based on vegetation cover (Fc), which is assumed to be equal to f_{PAR} from the MODIS product MOD15A2 (Mu et al., 2011). Although this method is based on the PM equation, PM-MOD does neither require wind speed nor soil moisture data for the parameterization of aerodynamic and surface resistance. Further details about PM-MOD can be found in Mu et al. (2011) and Running et al. (2019). Note that some updates have been implemented in PM-MOD since Mu et al. (2011), which can be found in Running et al. (2019). These were also considered here in the implementation of PM-MOD.

319

2.3.4 PM-VI

320

321

322

323

324

325

326

327

328

This model relies upon the hypothesis that ET is mostly controlled by specific dominant processes, such as transpiration and photosynthesis, hence a good correlation between such processes and ET is necessary for good model performance (Nagler et al., 2007). There are several formulations to estimate ET from VIs (Nagler et al., 2005, 2009). In this study, we selected the algorithm proposed by Nagler et al. (2013), which estimates ET using the reference crop evapotranspiration, ET_0 , from the FAO-56 Penman-Monteith (PM) equation (Allen et al., 1998), and a crop coefficient, K_{cVI} , derived from a vegetation index. K_{cVI} can be calculated in different ways (Nagler et al., 2005, 2013). Following Nouri et al. (2016) and P. T. S. Oliveira et al. (2015), K_{cVI} was calculated as:

$$K_{cVI} = a(1 - e^{-b \times EVI}) - c \quad (1)$$

329 where a , b and c are fitted coefficients. We used a parameter optimization tool based on
 330 a genetic algorithm to optimise the coefficients to estimate ET values close to the mea-
 331 sured ones (P. T. S. Oliveira et al., 2015). The fitting procedure minimizes the objec-
 332 tive function (OF) given by the sum of squared differences between tower-based ET (ET_{obs})
 333 and ET estimates from the models (ET_{sim}) at time i :

$$OF = \sum_{i=1}^n [ET_{obs}(i) - ET_{sim}]^2 \quad (2)$$

334 This model, herein referred to as PM-VI, has frequently been employed to estimate
 335 ET at local and regional scales (P. T. S. Oliveira et al., 2015; Nouri et al., 2016; Jarchow
 336 et al., 2017). Although obtaining ET_o requires a considerable amount of meteorologi-
 337 cal variables, the PM-VI implementation is easier and has a lower computational cost
 338 compared to other models. Unlike the three other models, PM-VI requires the calibra-
 339 tion of the fitting coefficients, which can be a major issue for regions where ET and VI
 340 are poorly correlated or when correlations change over time (Chong et al., 1993). To cal-
 341 ibrate the fitting coefficients, we randomly selected 20% of the available data at each site
 342 and used the remaining 80% to validate the model.

343 2.4 Quantifying model reliability

344 The model predictive skill was visually evaluated with scatter plots of measured
 345 versus modelled ET , as well as through the coefficient of determination (R^2), root mean
 346 square error ($RMSE$), percent bias ($PBIAS$), concordance correlation coefficient (ρ),
 347 slope (m), and intercept (b) of the linear regression. The data used in the analysis were
 348 filtered for rainy days ($P > 0.5$ mm). Our analysis proceeded from a general (no dis-
 349 tinction among sites) to a site-by-site and group level analysis, i.e. per biome, climate,
 350 or land use. The number of flux towers assigned to each subgroup (i.e., the different biomes,
 351 climate, and land use classes) varied, and so did the record length per subgroup. To ac-
 352 count for the different sizes, the following sampling procedure was performed, in which
 353 we computed the variability of each performance metrics for each group analysis (i.e.,
 354 across its different subgroups): (i) A sample size N was defined as half of the record length
 355 of the shortest subgroup, among all models; (ii) for each model, samples of length N were
 356 taken from within each subgroup, and the performance metrics were computed; (iii) This
 357 procedure was repeated 1000 times, yielding a mean and standard deviation (SD) of the
 358 metrics at each subgroup, per model. The resulting SD are likely to be influenced by the
 359 choice of N , and other rationale for its choice could have been made. In this way the con-
 360 fidence bands reported here are to be seen as measures of relative variability, i.e., the vari-
 361 ability between the models, and not as absolute uncertainty bounds for each of them.
 362 To establish a relationship between model predictive skill and water availability at in-
 363 dividual tower sites, we obtained the aridity index ($AI = P/ET_o$) from the global dataset
 364 provided by Trabucco and Zomer (2019). For many tower sites, the available meteoro-
 365 logical data (even from nearby meteorological stations) were not sufficient to provide a
 366 reliable AI ; hence the choice for a global dataset.

367 3 Results

368 3.1 ET partitioning

369 Partitioning of ET among the three components (E_{trans} , E_{int} and E_{soil}) exhib-
 370 ited more variation for the PT-JPL and PM-MOD models. On average, E_{trans} accounted
 371 for 60% (PT-JPL) and 56% (PM-MOD) of ET but, across sites, it presented a smaller
 372 range (30% to 85%) for PT-JPL than for PM-MOD (20 to 90%) (Figure 2, Table 1). GLEAM
 373 E_{trans} accounted for 82% of ET on average, varying between 60% and 95% across sites.
 374 Average interception across sites reached 9% (GLEAM), 13% (PT-JPL), and 24% (PM-

375 MOD) of total ET . E_{int} fractions range were similar for GLEAM and PT-JPL ($SD \approx$
 376 9%), whereas PM-MOD E_{int} varied more among sites ($SD = 18\%$). E_{int} was often cor-
 377 related with LAI , especially for the GLEAM estimates ($R^2 = 0.57$, Figure S2 in SM).
 378 PT-JPL E_{soil} estimates exceeded the other models, particularly for sites with low LAI
 379 values (e.g., ESEC, CST, and USR).

Figure 2. Evaporation fractions estimated by the models at each site (stacked bars) and av-
 erage partitioning of land evaporation per model (pie diagram). Black dots: LAI scaled between
 0 and 1 based on the minimum and maximum values of LAI (from MODIS MDC15A2 product).
 Red \times : the concordance correlation coefficient between observed and simulated daily ET.

380 3.2 Overall model skills

381 Since each model requires a different input dataset (Table S3, SM), the data avail-
 382 able to run and validate each model varied. GLEAM and PT-JPL provided a significantly
 383 greater number of daily outputs: 7301 (GLEAM), 7277 (PT-JPL), 5905 (PM-MOD), and
 384 6638 (PM-VI). The complete data set was used to produce scatter plots of ET records
 385 and model simulations for each location (See Figures S4-S7 in SM). To allow a fair anal-
 386 ysis, the results shown in the main text were obtained using data from days that were
 387 common across models, resulting in 4718 data points.

388 To illustrate the relative contribution of each site to the scatter plots in Figure 3,
 389 we display the regression lines (light grey lines) between model and tower-based ET for
 390 each tower site, and the mean metrics across individual sites. In general, ET was rea-
 391 sonably predicted by all models, as suggested by the relatively low spread of most points
 392 in the scatter plots, many regression lines close to the 1:1 line, mean determination co-
 393 efficient, \bar{R}^2 , mean concordance correlation coefficient, $\bar{\rho}$, mostly above 0.65, and mean
 394 root mean square error (\bar{RMSE}) below 1 mm d^{-1} (Figure 3). Nevertheless there is some
 395 spread for a few sites, e.g., in the PT-JPL scatter plot that displays a few sites with large
 396 bias despite strong overall correlation and ρ .

397 The models slightly overestimate ET as suggested by higher density of points be-
 398 low the 1:1 line, except for GLEAM, which slightly underestimates. Correlations were
 399 similar between GLEAM and PT-JPL, with an average value of ~ 0.65 and the highest
 400 values at individual sites reaching close to 0.9, as indicated by the standard deviations
 401 (0.19 and 0.18, respectively). From Figures 3 and 4, it becomes evident that, despite the
 402 relatively lower spread of points for PM-VI, compared to the other models, this model
 403 (PM-VI) presented a less consistent performance across towers, as suggested by the con-
 404 trasting slopes presented by the regression lines in that plot (e.g. reversed trend line at
 405 K77); hence the lower average determination coefficient (\bar{R}^2) and $\bar{\rho}$. Such contrasting
 406 aspect of the PM-VI model is also noted by the fact that a wide range of R^2 was found
 407 despite the similarity between mean simulated and observed ET (Figure 4).

Figure 3. Scatter plots of observed vs. simulated daily evapotranspiration at all flux tower
 sites, for each model. The light grey lines show the regression slope of individual sites. The co-
 efficient of determination (R^2), root mean square error ($RMSE$) and concordance correlation
 coefficient (ρ) were averaged across towers and are displayed on the plots ($N = 4,718$).

Table 1. Comparison of evaporation fractions for several land uses between this study and field-based estimates. FE = field estimates. Land covers that present field data from the same modeling sites or same geographical region are indicated with “*”.

LULC	E_{trans} (%)			E_{soil} (%)			E_{vint} (%)			References			
	FE	GLEAM PT- JPL	PM- MOD	FE	GLEAM PT- JPL	PM- MOD	FE	GLEAM PT- JPL	PM- MOD				
EBF*	80-84	74-79	47-63	31-88	NA	4-6	18-24	5-20	15-25	17-20	18-29	7-58	Leopoldo et al. (1995); Shuttleworth and Pereira (1988)
DNF*	50-81	84-94	64-84	78-90	NA	8-14	14-24	8-18	10	1-2	2-5	2-4	Gaj et al. (2016); Sun et al. (2019); de Queiroz et al. (2020)
CROP*	NA	93	63	70	20-4	6	31	21	10	1	6	9	Denmead et al. (1997); Cabral et al. (2012)
CROP*	85	88	55	69	NA	3	20	4	13	9	25	27	Cabral et al. (2010)
WS*	NA	86	76	78	NA	2	17	5	8	13	7	17	Cabral et al. (2015)
GRA	50-78	69-73	47-49	33-54	NA	25-30	32-46	30-52	NA	1-2	7-18	15-16	Ferretti et al. (2003); Sutantanto et al. (2012); Z. Wang et al. (2014)
MF	36-74	82-88	63-75	58-71	19	11-16	23-30	28-29	NA	1	2-7	1-14	(Aron et al., 2020; Paul-Limoges et al., 2020)
PW	33-38	73-86	28-57	34-41	NA	0-20	32-57	16-54	NA	3-14	6-16	21-43	J. Zhang et al. (2018)

Figure 4. Comparison of mean observed and simulated *ET*. Circle colors vary according to individual model R^2 .

Figure 5. Model performance per biome, land use and climate. The error bars represent the standard deviation of the metrics within each class. Biome types: Tropical & Subtropical Moist Broadleaf Forests (TSMBF); Tropical & Subtropical Dry Broadleaf Forests (TSDBF); Temperate Grasslands, Savannas & Shrublands (TGSS); Temperate Broadleaf & Mixed Forests (TBMF); Tropical & Subtropical Grasslands, Savannas & Shrublands (TSGSS); Flooded Grasslands & Savannas (FGS); Land use types: Cropland (CROP); Woodland Savanna (WS); Deciduous Needleleaf Forest (DNF); Evergreen Broadleaf Forest (EBF); Grasslands (GRA); Mixed Forest (MF); Permanent Wetland (PW); Deciduous Broadleaf Forest (DBF). Climate Zones: Tropical monsoon (Am); Tropical savanna (Aw); Hot semi-arid (BSh); Cold semi-arid (BSk); Temperate oceanic (Cfb); Dry-winter subtropical highland (Cwb); Polar Tundra (Td).

3.3 Model skills per biome, land use, and climate

Figure 5 presents ρ , $RMSE$, $PBIAS$, and R^2 for each model across six biomes, eight land use types, and seven climate classes in South America. Error bars are shown for all metrics, and they represent the standard deviation resulting from the resampling procedure outlined in 2.4. Note that the analysis about the FGS and TBMF biomes are based on one and three towers, respectively. For most biomes, $RMSE$ and R^2 did not significantly diverge. In general, TSGSS showed the best overall metrics for all models, while PM-VI in FGS (NPW site) presented the poorest ($\rho < 0.5$, $RMSE > 1.5 \text{ mm d}^{-1}$, and $R^2 < 0.25$). Model performance across towers within each biome did not vary much, as suggested by the relatively low range of the error bars for all metrics.

The central panels in Figure 5 provide evidence for the high variability of model predictive skills across different land uses (LU), which suggest that: (i) no model outperforms the others for all LU types, (ii) each model has intrinsic and in some cases exclusive characteristic that makes it more suitable for certain LU. Only for croplands (CROP) we found similar metrics among models ($\rho \approx 0.8$, $0.8 < RMSE < 1.2 \text{ mm d}^{-1}$, $-20\% < PBIAS < 10\%$, $0.6 < R^2 < 0.8$). Conversely, for most LU, the metrics variation is remarkable (e.g., DBF: $0.4 < \rho < 0.9$, $-50\% < PBIAS < 10\%$, $0.25 < R^2 < 0.80$). On average, each model has the best skills for two LU; e.g., *ET* prediction for GRA and DBF was best with PT-JPL ($\rho \approx 0.9$, $RMSE \approx 0.5 \text{ mm d}^{-1}$, $PBIAS \approx 0\%$, $R^2 > 0.75$) whereas PM-VI presented similar skills for estimation of *ET* for CROP and PW. Likewise, model skill is related to the climate type. The analysis of ρ and R^2 over semi-arid regions (BSk and BSh) indicates a relatively poor skill of all models (except PM-MOD for BSh climate). This is in contrast to the overall good performance over more humid environments (e.g., Aw and Cwb). The greatest divergence among model performances was found for the Polar Tundra (Td) climate zone, for which PM-VI presented the highest ρ and R^2 (both > 0.75), lowest $RMSE$ ($\sim 0.5 \text{ mm d}^{-1}$) and $PBIAS$ ($< 10\%$).

3.4 Individual sites

In this section, we explore the model performance at individual towers. Model skills for all individual sites are depicted in Figure 6. Sites with $N < 30$ (CAX and MCR) are not discussed here but are considered in the scatter plots shown in the SM (Figures S4-S7). To facilitate the comparison of our results with previous analyses using the same models, only three statistics are shown in Figure 6: $RMSE$, $PBIAS$, and R^2 . Other met-

441 metrics are displayed in the scatter plots in Figures S4-S7 in the SM. In Figure 6, the met-
 442 rics for the various towers are displayed in order of increasing aridity (varying from ~ 3
 443 to 0, left to right), as suggested by the AI as described in Section 2.4). In general, there
 444 is a good agreement between the PM-based models in terms of $RMSE$ and $PBIAS$.

445 In terms of individual metrics, $RMSE$ values varied between ~ 0.5 and ~ 1.5 mm d^{-1}
 446 for all models, with $RMSE < 1$ mm d^{-1} for most sites. The boxplots show that $RMSE$
 447 variation is similar among models, except for PT-JPL which presents the lowest $RMSE$
 448 (e.g., K67). Figure 6 shows that $PBIAS$ for PM-VI varies around zero across sites, which
 449 is expected given the model requires calibration with local data. However, based on R^2 ,
 450 it is apparent that this model's skill is quite limited for $AI > \sim 1.2$ and $AI < \sim 0.5$. In
 451 general, the PT-based models showed larger biases, with PT-JPL and GLEAM consis-
 452 tently overestimating and underestimating ET , respectively. In terms of R^2 , the PT-models
 453 ranked better than the PM-models for more than $\sim 50\%$ of the towers.

Figure 6. Comparison of statistics of the models in estimating evapotranspiration (ET) for
 the various flux towers used. (a) Sample size (N) used to compute the statistics; (b) $RMSE$
 = Root Mean Square Error; (c) Percent Bias ($PBIAS$); (d) R^2 = coefficient of determination.
 A summary of each model's statistics is depicted in the boxplots: (e) $RMSE$; (f) $PBIAS$; (g)
 R^2 . Flux towers are arranged according to the aridity index (with aridity increasing from left to
 right). Sites with $N < 30$ (CAX and MCR) are not shown here

454 4 Discussion

455 4.1 General implications

456 We conducted the first multi-remote sensing ET model analysis in South Amer-
 457 ica (SA) using a common set of forcing and validation data located on flux tower sites
 458 across a diverse range of land covers, climates, and biomes. Forcing data include both
 459 *in situ* (e.g., temperature and net radiation) and remote sensing data, mainly related
 460 to vegetation (e.g., LAI and EVI). To evaluate the models, energy balance-derived ET
 461 (ET_{EB}) was used as observation, instead of eddy covariance ET (ET_{EC}). Given the ben-
 462 efits and drawbacks of using either ET_{EB} or ET_{EC} , we compared both measures to ver-
 463 ify whether such choices would lead to different results. As shown in the SM, for the great
 464 majority of tower sites, ET_{EC} and ET_{EB} are similar (Figure S10) and model statistics
 465 (R^2 , $RMSE$ and ρ) remained the same regardless of the ET approach or indicate a bet-
 466 ter performance when ET_{EB} was used (Figures S11-S13 in the SM). Many of the tower
 467 sites considered here are not yet available in flux network databases, including sites with
 468 land cover (deciduous needle-leaf forests, DNF), a biome (FGS), and two climate types
 469 (polar tundra, hot semi-arid) that have not been previously assessed in other regional
 470 studies on the performance of satellite-based ET models. Moreover, some classes included
 471 here were considered for validation of individual models only (e.g., semi-arid and trop-
 472 ical climate types, and TSDBF biome).

473 The fulfillment of such gaps (i.e. model evaluation across uncharted regions) is an
 474 important step because it allows a multitude of applications and studies relying on large
 475 scale ET mapping, such as: drought monitoring (Anderson et al., 2011, 2016), agricul-
 476 tural water management (Anderson et al., 2012), diagnosis of climate change (Mao et
 477 al., 2015). The current ability to map ET remotely at various spatial and temporal scales,
 478 could only be evaluated thanks to the vast number of eddy covariance towers available
 479 in continental and global flux networks. As shown in this study, a thorough assessment
 480 of RBSET models based solely on data from such networks would be challenging or in-
 481 sufficient for some regions or continents; hence the relevance of this study. Our analy-

482 sis provides essential information to identify model strengths and limitations across SA,
 483 allowing the users to identify which model is more suitable for them. Knowing under what
 484 circumstances, e.g., land use or climate, each model is more reliable is necessary to ad-
 485 dress remaining research and applied science gaps relative to ET at local, regional and
 486 global scales (Fisher et al., 2017). Despite the value of tower-based ET across SA, many
 487 of those questions persist due to our limited observational capabilities. According to Fisher
 488 et al. (2017), the way to begin answering those questions is producing high quality ET
 489 estimates, which includes acquiring accurate ET information at high temporal and spa-
 490 tial resolution with large spatial coverage for a sufficient long period.

491 4.2 Model performance, sources of errors and ET partitioning

492 Generally, model predictive skill over SA resembles what has been reported for other
 493 continents, including satisfactory values of coefficient of determination ($R^2 > 0.6$) of the
 494 models (except PM-VI) for most validation sites, and consistently better results for the
 495 GLEAM and PT-JPL models, with $RMSE$ ranging from ~ 0.5 to 1.5 mm d^{-1} (McCabe
 496 et al., 2016). Also, in accordance with previous analysis, GLEAM and PT-JPL presented
 497 somewhat higher $RMSE$ than PM-MOD but no clear evidence indicates decreasing per-
 498 formance with increasing aridity, as reported by McCabe et al. (2016); Michel et al. (2016).
 499 Nonetheless, the general analysis (Section 3.2) indicates that all models can be used re-
 500 liably over most of the environmental conditions in SA covered in our study. The anal-
 501 ysis across towers and groups (i.e., biome, land use type and climate, Section 3.3, Fig-
 502 ure 5) identified considerable differences in terms of model skill.

503 Our results agree with previous studies from (Ershadi et al., 2014; McCabe et al.,
 504 2016; Michel et al., 2016; Miralles et al., 2016) who applied PM-MOD, GLEAM (except
 505 Ershadi et al. (2014)) and PT-JPL to sites located in Africa, Asia, Australia, Europe and
 506 Middle East and reported that PM-MOD showed, for most sites, lower correlations with
 507 measured ET compared to GLEAM and PT-JPL. Unlike previous analysis, our study
 508 agrees with Michel et al. (2016) in the sense that model skill seems to be unrelated to
 509 land cover. Michel et al. (2016) also reported a wide variation of R^2 (0.2–0.8) and $RMSE$
 510 ($0.8\text{--}2 \text{ mm d}^{-1}$), for different sites under mixed forests. Conversely, contrasting results
 511 between our results and previous studies were found for woodland savanna. While we
 512 found $0.5 < R^2 < 0.8$ and $0.7 < RMSE < 1.5 \text{ mm d}^{-1}$, Michel et al. (2016) reported
 513 $R^2 < 0.2$ and $1 < RMSE < 3 \text{ mm d}^{-1}$.

514 Overall, our group-wise analysis based on climate agrees with previous studies. For
 515 example, the poor model skill found here for the cold semi-arid (Bsk) climate ($0.1 < R^2 <$
 516 0.5) resembles that found by Michel et al. (2016) and McCabe et al. (2016) for several
 517 sites in the United States. While aridity could have played a role here, it could also be
 518 caused by the fact that semi-arid sites are covered with sparse canopies. Such canopies
 519 present challenges when it comes to the description of aerodynamic transfer for exam-
 520 ple and radiation partitioning (see e.g. Verhoef and Allen (2000)). Our findings also show
 521 a poor to moderate model skill for ET predictions for sites located in the Cfb climate
 522 zone, with PM-MOD having the worst performance. Conversely, PM-MOD presented
 523 the best predictive skill for the BSh climate, according to most metrics.

524 Besides the three RSBET models commonly assessed (GLEAM, PT-JPL, and PM-
 525 MOD), our analysis included the PM-VI model, which has been validated mostly for crop-
 526 land or riparian ecosystems (Nagler et al., 2005, 2009, 2013; Jarchow et al., 2017). Here,
 527 we tested PM-VI for a much wider variety of biomes, climates and land uses, and found
 528 a poor predictive skill for several sites with $AI > 1.2$ (e.g. K67, K77, K83) or $AI <$
 529 0.5 (e.g., CAA and SLU), even though the model accounts for a site-specific calibration.
 530 Considering the good results obtained for $\sim 50\%$ of the towers and the fact that, com-
 531 pared to the other models, PM-VI has a much simpler implementation, this model does
 532 have potential as long as sufficient data are available for calibration or, at least, valida-

tion. However, the need for local calibration is a hurdle for its implementation for most regions that are unsampled; therefore future studies are necessary to investigate which factors are most relevant in the determination of the model fitting coefficients, and to provide distributed reference values for its coefficients (e.g., based on land use dynamics).

We were able to identify a number of probable causes for poor model performance at individual sites, including (i) patch-scale heterogeneities; (ii) “mixed pixels”, i.e. mixed response of different vegetation types within a pixel; (iii) time-lag between ET_{obs} and EVI ; (iv) model sensitivity to individual inputs; (v) low correlation between ET and vegetation indices (see Section 3.0 in the SM for more details). Although we did not verify this in our study, we did not dismiss the possibility that known uncertainties in the estimation of site-specific vegetation characteristics (e.g., f_{PAR} and leaf stomatal conductance in the PM-MOD; Ershadi et al. (2014)) are further causes of lower model performance.

In our study, we used soil heat flux (G) which is generally measured below ground (usually at 5–20 cm deep) using soil heat flux plates. It could be argued that not correcting G for the heat storage between the plate and the soil surface could lead to sub-optimal estimates of ET when LE is calculated as the residual of the energy balance, especially for towers where the soil is bare or covered by sparse vegetation, where G can be relatively large. This, in turn, could lead to the conclusion that the models are performing worse than is actually the case. Although desirable, correcting G for heat storage is rarely possible due to data unavailability (few sites only measure soil moisture and temperature, which are required to estimate soil heat capacity, and heat storage using the calorimetric method). Moreover, at daily scales and for most sites, G is either negligible in SA (summer or winter, when the amount of heat stored during the day roughly equals that lost during the night) or represents a minor portion only (spring and autumn) of the energy balance. As detailed and discussed in Section S3.0 and Figure S7 in SM, it is highly unlikely that neglecting such corrections will have affected the results.

There are, however, some issues worth mentioning here. Cause (v), for instance, is a major issue for PM-VI, as expected because the model is highly dependent on VI dynamics (see Section 2.4) (Nagler et al., 2005). Regarding cause (iv), the superior performance of the PT models over PM-MOD at most sites is probably linked to uncertainties resulting from the estimation of aerodynamic resistance (Ershadi et al., 2014). In PM-MOD, the aerodynamic and surface resistances of each ET component (soil, interception, and transpiration) are parametrized based on biome-specific values of leaf-scale boundary layer conductance, for example (Mu et al., 2011). Compared to the previous version of PM-MOD (Mu et al., 2007), this new approach resulted in a perceptible improvement only for cropland and deciduous broadleaf forest flux tower sites, whereas for other land uses no meaningful change was reported (Ershadi et al., 2015). Conversely, PT models are highly dependent on Rn (causes iv and v); hence they often fail in dry environments (see metrics for $AI < \sim 0.6$ in Figure 6) where ET seasonality is dictated by P more than radiation, or in regions with low Rn (e.g., TF2). Poor model responses at K77 (cropland, Figure S9 in SM) were attributed to causes (i) and (ii), as remnants of forest and shrubs were identified within the tower footprint and within MODIS pixel. VI products with higher resolution than MODIS exist and have been used to estimate ET (Aragon et al., 2018; Fisher et al., 2020); thus offering a possible solution for causes (i) and (ii). Time lag between ET and EVI (cause iii) was identified at EUC, where EVI followed the decline of ET after ~ 1 – 2 months.

Regardless of all those potential causes for poor model response, it is also important to consider to role of the core formulation upon which those RBSET models are based, i.e. Penman-Monteith (PM) and Priestley and Taylor (PT) equations. A major problem of the PM equation refers to the linearization of the Clausius-Claperyron relation, which has been addressed in a new version of that equation (McColl, 2020). The PT equa-

tion, in turn, implicitly assumes Rn and surface temperature (T_s) to be independent of evaporation. In reality, as shown by Yang and Roderick (2019), Rn not only decreases with increasing T_s (due to an increase of outgoing longwave radiation) but also a greater fraction of Rn becomes available for evaporation. Some of the deviations from the observations found in our analysis may happen due to such inconsistencies or simplifications. Here, we provide evidence to consider revisiting not just parameter values but the governing equations themselves and, ultimately, evaluate the benefits of such potential improvements in RBSET models.

Remote sensing based ET partitioning is expected to present some divergences from ground based measurements. This is the case especially for E_{soil} , because of the difficulty in obtaining remote sensing information on soil characteristics that drive E_{soil} , such as soil moisture and temperature (Talsma, Good, Jimenez, et al., 2018; Talsma, Good, Miralles, et al., 2018), in particular at high vegetation cover fractions. Globally, transpiration has been reported to account for 57–90% of global ET , based on *in situ* data and model outputs (Jasechko et al., 2013; Wei et al., 2017; Paschalis et al., 2018). Although these are global estimates, we expected E_{trans} to be the largest ET component also in SA due to its prevailing tropical climate and corresponding vegetation types. Our results show that this was indeed the case for GLEAM with an E_{trans}/ET ratio of $\sim 80\%$, and for PT-JPL and PM-MOD with values of 57 and 60%, respectively. Nonetheless, based on our findings, model predictive skill in estimating total ET is not necessarily associated with its ability to partition ET accurately.

Concomitantly, inconsistencies in ET partitioning do not necessarily translate into inaccurate model estimates of total ET : this depends on the modelling approach. On the one hand, if total ET results from the sum of ET components independently, then an under- or overestimation of ET components can reduce the overall model skill, or reasonable ET estimates can be achieved as the consequence of an occasional compensation of errors in E_{trans} , E_{soil} and E_{int} . On the other hand, if the ET partitioning is derived from the estimate of a proxy value for total ET , such as available energy flux (as in PM-MOD and PT-JPL), the ET partitioning is unlikely to influence the total ET estimates. Moreover, ET partitioning may be sensitive to certain model inputs. For example, contrasting ET fractions were estimated by PM-MOD for similar rain forest sites, i.e. K67 and K83 (Figure 2). The reason PM-MOD is returning that difference is because RH was estimated from actual (e_a) and saturation vapor pressure (e_s) data, as RH data is not available in K83 dataset. As a result, the difference between e_a -derived daytime and nighttime RH for K83 is greater than that for K67. In terms of daily averages, e_a -derived RH and measured RH are quite similar, which explains why the fractions for GLEAM and PT-JPL, at those two towers, are similar. Still, good estimates of ET components are important to differentiate the roles of vegetation and soil, i.e., how they contribute to vertical soil water fluxes and changes in profile soil water content. Reliable knowledge of the distribution between E_{soil} and E_{trans} is also important when this information is used in hydrological models to calculate other water balance components, such as runoff.

Ground-based ET partitioning data are generally not widely available. This also goes for most land cover types included in this study. We compared the models' outputs with field experiment studies that measured one or more ET components either at the same sites as those used here or within the same region (Table 1). ET partitioning values derived from GLEAM seem to be more consistent with ground-based information available for tropical rain forests, croplands and grasslands than for wetlands, and mixed and deciduous needle-leaf forests (Table 1). This also applies to PT-JPL with its ET partitioning agreeing reasonably well with observations made for both tropical rain- and dry forests. Note that PT-JPL (as well as PM-MOD) constrain E_{trans} based on f_{wet} . Hence, compared to GLEAM, transpiration will be lower under high RH in the model but ET can be high due to water availability in the soil and intercepted rainfall. Nonetheless,

639 the overall predictive skill of PT-JPL was satisfactory at such sites (Figure 6 and Fig-
 640 ure S4 in SM). Regarding PM-MOD, the main inconsistency is the E_{inter} for tropical
 641 forests (Table 1). Despite the wide variability in E_{trans}/ET among models, their over-
 642 all predictive skill was satisfactory, that is, not associated with their capability to cor-
 643 rectly estimate each ET component individually (see SM for further discussion). No model
 644 was able to consistently capture the ET partitioning across all sites correctly, which is
 645 expected given the uncertainty of each ET component and the climate and land-cover
 646 variability in SA. However, the joint estimates of all models covered totally or partially
 647 all field-derived evidence on ET partitioning. This suggests that continental ET esti-
 648 mates for understudied regions, such as the SA, would benefit from merging ET outputs
 649 from models that are based on different methods (Paca et al., 2019).

650 Despite our efforts to gather as much tower data as possible, with the goal of hav-
 651 ing a common data set for all models, we faced several limitations including: differences
 652 in lengths of observational time series across towers (up to 3 years), as well as lack in
 653 overlap of these time series; uneven distribution of towers across groups (e.g., biomes);
 654 and, finally, South American geographical features that were not considered in this study
 655 (e.g., MGS biome or desert climate type, BWk). Thus, it was not possible to assess, for
 656 all towers, model responses during all seasons. Nonetheless, the fact that our dataset en-
 657 compasses a wide variety of climates enabled us to evaluate, to a reasonable extent, model
 658 responses for contrasting seasons and fill in the gaps flagged by the literature, such as
 659 the absence, in a similar analysis, of towers in the tropical climate zone pointed out by
 660 McCabe et al. (2016).

661 5 Conclusion

662 Our results show that, in general, ET can be reasonably well predicted by all four
 663 models, despite an overall tendency of overestimation by PT-JPL and PM-MOD, and
 664 underestimation by GLEAM. Contrasting with results from other continents, we found
 665 no clear evidence linking model predictive skill with aridity. Our analysis emphasizes the
 666 need of improving model ET partitioning, although the link between flawed ET parti-
 667 tioning and poor model skill is not evident based on our results. Having reliable ET par-
 668 titioning coefficients as part of the FLUXNET-type datasets would be valuable in this
 669 respect, but unfortunately such data are difficult to obtain, as they require labour-intensive
 670 and expensive methods (such as sapflow gauges and lysimeters), that also present prob-
 671 lems with regards to upscaling from plot to field-scale.

672 Correlations are consistently higher for GLEAM and PT-JPL, with $R^2 > 0.5$ for
 673 most sites, whereas PM-MOD and PM-VI presented better performances in terms of PBIAS
 674 ($-10 < PBIAS < 10\%$ for most sites). As for PM-VI, the low PBIAS is expected,
 675 given the model requires calibration with local data.

676 The model skill for the various models seems to be unrelated to land cover type
 677 as we found a wide variability of metric values within the same class and across mod-
 678 els. Conversely, a relatively lower performance was observed for most models in semi-
 679 arid regions, compared to an overall good performance for more humid environments.
 680 Except for the FGS biome, we found that skill across models was mostly similar within
 681 the same biome.

682 Despite the relatively high number of towers (compared to previous global anal-
 683 yses that used a similar amount of sites), gathering a balanced amount of data and uni-
 684 form distribution of towers across different biomes and climate zones across the whole
 685 continent was challenging. Thus, there is a need to expand the flux tower network in South
 686 America as well as the formation of bilateral collaboration for future contributions. Pre-
 687 vious studies (Michel et al., 2016; McCabe et al., 2016) have expressed the need to ex-
 688 tend the evaluation of RSBET models to uncharted biomes and climate conditions. Our

689 analysis fills this gap by assessing the reliability of four RSBET models over South Amer-
 690 ica. We provide benchmarking metrics that can serve the improvement of *ET* models
 691 for improved capturing of *ET* over this continent.

692 **Acknowledgments**

693 The data used in this study will be available through a data-sharing repository. Fund-
 694 ing for AmeriFlux data resources was provided by the U.S. Department of Energy's Of-
 695 fice of Science. Davi de C. D. Melo was supported by the São Paulo State Research Foun-
 696 dation (FAPESP) (grant 2016/23546-7), and by the Brazilian National Council for Sci-
 697 entific and Technological Development (CNPq) (project 409093/2018-1). Jamil A. A. Anache
 698 was supported by the Brazilian National Council for Scientific and Technological Devel-
 699 opment (CNPq) (project 150057/2018-0). Edson Wendland was supported by the São
 700 Paulo State Research Foundation (FAPESP) (grant 2015/03806-1). Paulo Tarso S. Oliveira
 701 was supported by the Brazilian National Council for Scientific and Technological Devel-
 702 opment (CNPq) (grants 441289/2017-7 and 306830/2017-5) and the CAPES Print pro-
 703 gram. Rafael Rosolem would like to acknowledge the Brazilian Experimental datasets
 704 for Multi-Scale interactions in the critical zone under Extreme Drought (BEMUSED)
 705 project [grant number NE/R004897/1] funded by the Natural Environment Research Coun-
 706 cil (NERC). Alvaro Moreno was financially supported by the NASA Earth Observing
 707 System MODIS project (grant NNX08AG87A) and the European Research Council (ERC)
 708 funding under the ERC Consolidator Grant 2014 SEDAL (Statistical Learning for Earth
 709 Observation Data Analysis, European Union) project under Grant Agreement 647423.
 710 Diego G. Miralles, Brecht Martens and Dominik Rains are supported by the European
 711 Research Council (ERC) DRY-2-DRY project (grant no. 715254) and the Belgian Sci-
 712 ence Policy Office (BELSPO) STEREO III ALBERI (grant no. SR/00/373) and ET-
 713 SENSE (grant. no SR/02/377) projects. Thiago R. Rodrigues was supported by the Brazil-
 714 ian National Council for Scientific and Technological Development (CNPq) with Bolsa
 715 de Produtividade em Pesquisa - PQ (Grant Number 308844/2018-1). Jorge Perez-Quezada
 716 and Mauricio Galleguillos were supported by the Chilean National Agency for Research
 717 and Development, grant FONDECYT 1211652. Rodolfo Nobrega and Anne Verhoef ac-
 718 knowledge support by the Newton/NERC/FAPESP Nordeste project (NE/N012526/1
 719 ICL and NE/N012488/1 UoR). Gabriela Posse acknowledges support by AERN 3632 and
 720 PNNAT 1128023 INTA Projects. JBF was supported in part by NASA: ECOSTRESS
 721 and SUSMAP. Funding for site support:

- 722 • NPW tower: Brazilian National Institute for Science and Technology in Wetlands
 723 (INCT-INAU), Federal University of Mato Grosso (UFMT - PGFA and PGAT),
 724 University of Cuiabá (UNIC) and SESC-Pantanal;
- 725 • SDF tower: funded by the National Commission for Scientific and Technological
 726 Research (CONICYT, Chile) through grants FONDEQUIP AIC-37 and AFB170008
 727 from the Associative Research Program;
- 728 • TF1 and TF2 towers: funded by the Deutsche Forschungsgemeinschaft (DFG) un-
 729 der Germany's Excellence Strategy - EXC 177 'CliSAP - Integrated Climate Sys-
 730 tem Analysis and Prediction' - contributing to the Center for Earth System Re-
 731 search and Sustainability (CEN) of Universität Hamburg and by DFG project KU
 732 1418/6-1;
- 733 • MCR and BAL towers: funded by the National Council for Scientific and Tech-
 734 nological Research (CONICET, Argentina) grants PIP-11220100100044 and PIP-
 735 11220130100347CO, and by the National Agency for the Scientific and Techno-
 736 logical Promotion (ANPCyT, Argentina) grant PICT 2010-0554;
- 737 • CAA, CST, and ESEC Towers: funded by National Observatory of Water and Car-
 738 bon Dynamics in the Caatinga Biome (INCT-NOWCDCB), Federal University
 739 of Pernambuco (UFPE), FACEPE (Pernambuco State Research and Technology
 740 Foundation) through the Project Caatinga-FLUX APQ 0062-1.07/15.

References

- 741 Akaike, H. (1969, December). Fitting autoregressive models for prediction. *Annals of*
 742 *the Institute of Statistical Mathematics*, 21(1), 243–247. Retrieved 2020-06-22,
 743 from <https://doi.org/10.1007/BF02532251> doi: 10.1007/BF02532251
 744 Allen, R. G., Pereira, L. S., Howell, T. A., & Jensen, M. E. (2011, April). Evap-
 745 otranspiration information reporting: I. Factors governing measurement
 746 accuracy. *Agricultural Water Management*, 98(6), 899–920. Retrieved
 747 2021-04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0378377411000023)
 748 [S0378377411000023](https://www.sciencedirect.com/science/article/pii/S0378377411000023) doi: 10.1016/j.agwat.2010.12.015
 749 Allen, R. G., Pereira, L. S., Raes, D., & Smith, M. (1998). *Crop evapotranspira-*
 750 *tion - Guidelines for computing crop water requirements - FAO Irrigation and*
 751 *drainage paper 56*. Rome: FAO - Food and Agriculture Organization of the
 752 United Nations. Retrieved 2021-04-15, from [http://www.fao.org/3/x0490e/](http://www.fao.org/3/x0490e/x0490e00.htm)
 753 [x0490e00.htm](http://www.fao.org/3/x0490e/x0490e00.htm)
 754 Amiro, B. (2009). Measuring boreal forest evapotranspiration using the energy bal-
 755 ance residual. *Journal of Hydrology*, 366(1), 112–118. Retrieved from [https://](https://www.sciencedirect.com/science/article/pii/S0022169408006367)
 756 [www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0022169408006367)
 757 [S0022169408006367](https://www.sciencedirect.com/science/article/pii/S0022169408006367) doi:
 758 <https://doi.org/10.1016/j.jhydrol.2008.12.021>
 759 Anderson, M. C., Allen, R. G., Morse, A., & Kustas, W. P. (2012). Use of
 760 landsat thermal imagery in monitoring evapotranspiration and manag-
 761 ing water resources. *Remote Sensing of Environment*, 122, 50–65. Re-
 762 trieved from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0034425712000326)
 763 [S0034425712000326](https://www.sciencedirect.com/science/article/pii/S0034425712000326) (Landsat Legacy Special Issue) doi: [https://doi.org/](https://doi.org/10.1016/j.rse.2011.08.025)
 764 [10.1016/j.rse.2011.08.025](https://doi.org/10.1016/j.rse.2011.08.025)
 765 Anderson, M. C., Hain, C., Wardlow, B., Pimstein, A., Mecikalski, J. R., & Kustas,
 766 W. P. (2011). Evaluation of drought indices based on thermal remote sensing
 767 of evapotranspiration over the continental united states. *Journal of Climate*,
 768 24(8), 2025 - 2044. Retrieved from [https://journals.ametsoc.org/view/](https://journals.ametsoc.org/view/journals/clim/24/8/2010jcli3812.1.xml)
 769 [journals/clim/24/8/2010jcli3812.1.xml](https://journals.ametsoc.org/view/journals/clim/24/8/2010jcli3812.1.xml) doi: 10.1175/2010JCLI3812.1
 770 Anderson, M. C., Zolin, C. A., Sentelhas, P. C., Hain, C. R., Semmens, K., Tu-
 771 grul Yilmaz, M., ... Tetrault, R. (2016, March). The Evaporative Stress
 772 Index as an indicator of agricultural drought in Brazil: An assessment based
 773 on crop yield impacts. *Remote Sensing of Environment*, 174, 82–99. Retrieved
 774 2019-10-17, from [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S0034425715302212)
 775 [S0034425715302212](http://www.sciencedirect.com/science/article/pii/S0034425715302212) doi: 10.1016/j.rse.2015.11.034
 776 Aragon, B., Houborg, R., Tu, K., Fisher, J. B., & McCabe, M. (2018, December).
 777 CubeSats Enable High Spatiotemporal Retrievals of Crop-Water Use for Pre-
 778 cision Agriculture. *Remote Sensing*, 10(12), 1867. Retrieved 2020-09-01, from
 779 <https://www.mdpi.com/2072-4292/10/12/1867> (Number: 12 Publisher:
 780 Multidisciplinary Digital Publishing Institute) doi: 10.3390/rs10121867
 781 Aron, P. G., Poulsen, C. J., Fiorella, R. P., Matheny, A. M., & Veverica, T. J.
 782 (2020). An isotopic approach to partition evapotranspiration in a mixed
 783 deciduous forest. *Ecohydrology*, 13(6), e2229. Retrieved 2021-04-11,
 784 from <https://onlinelibrary.wiley.com/doi/abs/10.1002/eco.2229>
 785 (_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1002/eco.2229>) doi:
 786 <https://doi.org/10.1002/eco.2229>
 787 Arroyo, M. T. K., Pliscoff, P., Mihoc, M., & Arroyo-Kalin, M. (2005). The Magel-
 788 lanic moorland. In L. H. Fraser & P. A. Keddy (Eds.), *The World's Largest*
 789 *Wetlands: Ecology and Conservation* (pp. 424–445). Cambridge: Cambridge
 790 University Press. Retrieved 2020-08-15, from [https://www.cambridge](https://www.cambridge.org/core/books/worlds-largest-wetlands/magellanic-moorland/B02EB2AE6B5BC3EB290B88FE794D3A77)
 791 [.org/core/books/worlds-largest-wetlands/magellanic-moorland/](https://www.cambridge.org/core/books/worlds-largest-wetlands/magellanic-moorland/B02EB2AE6B5BC3EB290B88FE794D3A77)
 792 [B02EB2AE6B5BC3EB290B88FE794D3A77](https://www.cambridge.org/core/books/worlds-largest-wetlands/magellanic-moorland/B02EB2AE6B5BC3EB290B88FE794D3A77) doi: 10.1017/CBO9780511542091.013
 793 Arruda, P. H. Z., Vourlitis, G. L., Santanna, F. B., Jr., O. B. P., Lobo, F. A., &
 794 Nogueira, J. S. (2016). Large net co2 loss from a grass-dominated tropical sa-
 795 vanna in south-central brazil in response to seasonal and interannual drought.

- 796 *Biogeosciences*, 121, 2110-2124. doi: 10.1002/2016JG003404
- 797 Badgley, G., Fisher, J. B., Jiménez, C., Tu, K. P., & Vinukollu, R. (2015, Au-
798 gust). On Uncertainty in Global Terrestrial Evapotranspiration Estimates
799 from Choice of Input Forcing Datasets. *Journal of Hydrometeorology*, 16(4),
800 1449–1455. Retrieved 2020-09-01, from [https://journals.ametsoc.org/
801 jhm/article/16/4/1449/6244/On-Uncertainty-in-Global-Terrestrial](https://journals.ametsoc.org/jhm/article/16/4/1449/6244/On-Uncertainty-in-Global-Terrestrial)
802 (Publisher: American Meteorological Society) doi: 10.1175/JHM-D-14-0040.1
- 803 Bai, J., Jia, L., Liu, S., Xu, Z., Hu, G., Zhu, M., & Song, L. (2015, May). Charac-
804 terizing the Footprint of Eddy Covariance System and Large Aperture Scintil-
805 lometer Measurements to Validate Satellite-Based Surface Fluxes. *IEEE Geo-
806 science and Remote Sensing Letters*, 12(5), 943–947. (Conference Name: IEEE
807 Geoscience and Remote Sensing Letters) doi: 10.1109/LGRS.2014.2368580
- 808 Bastiaanssen, W. G. M., Menenti, M., Feddes, R. A., & Holtslag, A. A. M. (1998,
809 December). A remote sensing surface energy balance algorithm for land (SE-
810 BAL). 1. Formulation. *Journal of Hydrology*, 212-213, 198–212. Retrieved
811 2020-06-15, from [http://www.sciencedirect.com/science/article/pii/
812 S0022169498002534](http://www.sciencedirect.com/science/article/pii/S0022169498002534) doi: 10.1016/S0022-1694(98)00253-4
- 813 Bezerra, B. G., Santos, C. A. C. d., Silva, B. B. d., Perez-Marin, A. M., Bez-
814 erra, M. V. C., Bezerra, J. R. C., & Rao, T. V. R. (2013, June). Estima-
815 tion of soil moisture in the root-zone from remote sensing data. *Revista
816 Brasileira de Ciência do Solo*, 37(3), 596–603. Retrieved 2021-04-15, from
817 [http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100
818 -06832013000300005&lng=en&nrm=iso&tlng=en](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-06832013000300005&lng=en&nrm=iso&tlng=en) (Publisher: Sociedade
819 Brasileira de Ciência do Solo) doi: 10.1590/S0100-06832013000300005
- 820 Bezerra, B. G., Silva, B. B. d., Santos, C. A. C. d., & Bezerra, J. R. C. (2015,
821 July). Actual Evapotranspiration Estimation Using Remote Sensing: Com-
822 parison of SEBAL and SSEB Approaches. *Advances in Remote Sensing*,
823 4(3), 234–247. Retrieved 2021-04-15, from [http://www.scirp.org/Journal/
824 Paperabs.aspx?paperid=59977](http://www.scirp.org/Journal/Paperabs.aspx?paperid=59977) (Number: 3 Publisher: Scientific Research
825 Publishing) doi: 10.4236/ars.2015.43019
- 826 Blunden, J., & Arndt, D. S. (2020, August). State of the Climate in 2019. *Bulletin
827 of the American Meteorological Society*, 101(8), S1–S429. Retrieved 2020-09-
828 03, from [https://journals.ametsoc.org/bams/article/101/8/S1/353885/
829 State-of-the-Climature-in-2019](https://journals.ametsoc.org/bams/article/101/8/S1/353885/State-of-the-Climature-in-2019) (Publisher: American Meteorological Soci-
830 ety) doi: 10.1175/2020BAMSSstateoftheClimate.1
- 831 Borma, L. S., Rocha, H. R. d., Cabral, O. M., Randow, C. v., Collicchio, E.,
832 Kurzatkowski, D., ... Artaxo, P. (2009). Atmosphere and hydrologi-
833 cal controls of the evapotranspiration over a floodplain forest in the Ba-
834 nanal Island region, Amazonia. *Journal of Geophysical Research: Bio-
835 geosciences*, 114(G1). Retrieved 2020-04-21, from [https://agupubs
836 .onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000641](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000641) (eprint:
837 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2007JG000641>) doi:
838 10.1029/2007JG000641
- 839 Cabral, O. M. R., da Rocha, H. R., Gash, J. H., Freitas, H. C., & Ligo, M. A. V.
840 (2015, September). Water and energy fluxes from a woodland savanna (cer-
841 rado) in southeast Brazil. *Journal of Hydrology: Regional Studies*, 4, 22–
842 40. Retrieved 2020-08-05, from [http://www.sciencedirect.com/science/
843 article/pii/S2214581815000440](http://www.sciencedirect.com/science/article/pii/S2214581815000440) doi: 10.1016/j.ejrh.2015.04.010
- 844 Cabral, O. M. R., Freitas, H. C., Cuadra, S. V., de Andrade, C. A., Ramos, N. P.,
845 Grutzmacher, P., ... Rossi, P. (2020, March). The sustainability of a sugar-
846 cane plantation in Brazil assessed by the eddy covariance fluxes of greenhouse
847 gases. *Agricultural and Forest Meteorology*, 282-283, 107864. Retrieved
848 2021-10-01, from [https://www.sciencedirect.com/science/article/pii/
849 S0168192319304800](https://www.sciencedirect.com/science/article/pii/S0168192319304800) doi: 10.1016/j.agrformet.2019.107864
- 850 Cabral, O. M. R., Gash, J. H. C., Rocha, H. R., Marsden, C., Ligo, M. A. V., Fre-

- 851 itas, H. C., ... Gomes, E. (2011, January). Fluxes of CO₂ above a plantation
852 of Eucalyptus in southeast Brazil. *Agricultural and Forest Meteorology*, 151(1),
853 49–59. Retrieved 2020-07-24, from [http://www.sciencedirect.com/science/
854 article/pii/S0168192310002480](http://www.sciencedirect.com/science/article/pii/S0168192310002480) doi: 10.1016/j.agrformet.2010.09.003
- 855 Cabral, O. M. R., Rocha, H. R., Gash, J. H., Ligo, M. A. V., Tatsch, J. D., Freitas,
856 H. C., & Brasilio, E. (2012). Water use in a sugarcane plantation. *GCB Bioen-
857 ergy*, 4(5), 555–565. Retrieved 2020-07-28, from [https://onlinelibrary
858 .wiley.com/doi/abs/10.1111/j.1757-1707.2011.01155.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1757-1707.2011.01155.x) (eprint:
859 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1757-1707.2011.01155.x>)
860 doi: 10.1111/j.1757-1707.2011.01155.x
- 861 Cabral, O. M. R., Rocha, H. R., Gash, J. H. C., Ligo, M. A. V., Freitas, H. C., &
862 Tatsch, J. D. (2010, July). The energy and water balance of a Eucalyp-
863 tus plantation in southeast Brazil. *Journal of Hydrology*, 388(3), 208–216.
864 Retrieved 2021-04-08, from [https://www.sciencedirect.com/science/
865 article/pii/S0022169410002404](https://www.sciencedirect.com/science/article/pii/S0022169410002404) doi: 10.1016/j.jhydrol.2010.04.041
- 866 Campos, S., Mendes, K. R., da Silva, L. L., Mutti, P. R., Medeiros, S. S., Amorim,
867 L. B., ... Bezerra, B. G. (2019, June). Closure and partitioning of the en-
868 ergy balance in a preserved area of a Brazilian seasonally dry tropical forest.
869 *Agricultural and Forest Meteorology*, 271, 398–412. Retrieved 2020-05-06, from
870 <http://www.sciencedirect.com/science/article/pii/S0168192319301303>
871 doi: 10.1016/j.agrformet.2019.03.018
- 872 Cao, L., Bala, G., Caldeira, K., Nemani, R., & Ban-Weiss, G. (2010, May). Im-
873 portance of carbon dioxide physiological forcing to future climate change. *Pro-
874 ceedings of the National Academy of Sciences*, 107(21), 9513–9518. Retrieved
875 2021-04-15, from <https://www.pnas.org/content/107/21/9513> (Publisher:
876 National Academy of Sciences Section: Physical Sciences) doi: 10.1073/pnas
877 .0913000107
- 878 Chang, Y., Qin, D., Ding, Y., Zhao, Q., & Zhang, S. (2018, June). A modified
879 MOD16 algorithm to estimate evapotranspiration over alpine meadow on
880 the Tibetan Plateau, China. *Journal of Hydrology*, 561, 16–30. Retrieved
881 2021-04-15, from [https://www.sciencedirect.com/science/article/pii/
882 S0022169418302269](https://www.sciencedirect.com/science/article/pii/S0022169418302269) doi: 10.1016/j.jhydrol.2018.03.054
- 883 Chen, B., Black, T. A., Coops, N. C., Hilker, T., (Tony) Trofymow, J. A., & Mor-
884 genstern, K. (2009, February). Assessing Tower Flux Footprint Clima-
885 tology and Scaling Between Remotely Sensed and Eddy Covariance Mea-
886 surements. *Boundary-Layer Meteorology*, 130(2), 137–167. Retrieved
887 2020-07-09, from <https://doi.org/10.1007/s10546-008-9339-1> doi:
888 10.1007/s10546-008-9339-1
- 889 Chong, D. L. S., Mougin, E., & Gastellu-Etchegorry. (1993, May). Relating the
890 Global Vegetation Index to net primary productivity and actual evapotranspi-
891 ration over Africa. *International Journal of Remote Sensing*, 14(8), 1517–1546.
892 Retrieved 2019-09-05, from <https://doi.org/10.1080/01431169308953984>
893 doi: 10.1080/01431169308953984
- 894 Costanza, R., d’Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., ...
895 van den Belt, M. (1997, May). The value of the world’s ecosystem services
896 and natural capital. *Nature*, 387(6630), 253–260. Retrieved 2021-04-22, from
897 <https://www.nature.com/articles/387253a0> (Number: 6630 Publisher:
898 Nature Publishing Group) doi: 10.1038/387253a0
- 899 Curto, L., Covi, M., & Gassmann, M. I. (2019, December). Actual evapotranspi-
900 ration and the pattern of soil water extraction of a soybean (*Glycine max*)
901 crop. *Revista de la Facultad de Ciencias Agrarias UNCuyo*, 51(2), 125–141.
902 Retrieved 2021-03-11, from [http://revistas.uncu.edu.ar/ojs/index.php/
903 RFCA/article/view/2615](http://revistas.uncu.edu.ar/ojs/index.php/RFCA/article/view/2615) (Number: 2)
- 904 Dalmagro, H. J., Lathuillière, M. J., Hawthorne, I., Morais, D. D., Pinto Jr, O. B.,
905 Couto, E. G., & Johnson, M. S. (2018, June). Carbon biogeochemistry

- 906 of a flooded Pantanal forest over three annual flood cycles. *Biogeochem-*
 907 *istry*, 139(1), 1–18. Retrieved 2020-08-18, from [https://doi.org/10.1007/](https://doi.org/10.1007/s10533-018-0450-1)
 908 [s10533-018-0450-1](https://doi.org/10.1007/s10533-018-0450-1) doi: 10.1007/s10533-018-0450-1
- 909 Denmead, O. T., Mayocchi, C. L., & Dunin, F. X. (1997). Does green cane harvest-
 910 ing conserve soil water? In *Proceedings of the Australian Society of Sugar Cane*
 911 *Technologists* (Vol. 19, pp. 139–146).
- 912 de Oliveira, R. G., Valle Júnior, L. C. G., da Silva, J. B., Espíndola, D. A. L. F.,
 913 Lopes, R. D., Nogueira, J. S., ... Rodrigues, T. R. (2021, May). Temporal
 914 trend changes in reference evapotranspiration contrasting different land uses
 915 in southern Amazon basin. *Agricultural Water Management*, 250, 106815.
 916 Retrieved 2021-04-01, from [https://www.sciencedirect.com/science/](https://www.sciencedirect.com/science/article/pii/S0378377421000809)
 917 [article/pii/S0378377421000809](https://www.sciencedirect.com/science/article/pii/S0378377421000809) doi: 10.1016/j.agwat.2021.106815
- 918 de Queiroz, M. G., da Silva, T. G. F., Zolnier, S., de Souza, C. A. A., de Souza,
 919 L. S. B., do Nascimento Araújo, G., ... de Moura, M. S. B. (2020, April).
 920 Partitioning of rainfall in a seasonal dry tropical forest. *Ecology*
 921 *& Hydrobiology*, 20(2), 230–242. Retrieved 2021-03-31, from [https://](https://www.sciencedirect.com/science/article/pii/S1642359320300124)
 922 www.sciencedirect.com/science/article/pii/S1642359320300124 doi:
 923 10.1016/j.ecohyd.2020.02.001
- 924 Embry, J. L., & Nothnagel, E. A. (1994, September). Leaf Senescence of Post-
 925 production Poinsettias in Low-light Stress. *Journal of the American So-*
 926 *ciety for Horticultural Science*, 119(5), 1006–1013. Retrieved 2021-04-22,
 927 from [https://journals.ashs.org/jashs/view/journals/jashs/119/5/](https://journals.ashs.org/jashs/view/journals/jashs/119/5/article-p1006.xml)
 928 [article-p1006.xml](https://journals.ashs.org/jashs/view/journals/jashs/119/5/article-p1006.xml) (Publisher: American Society for Horticultural Sci-
 929 ence Section: Journal of the American Society for Horticultural Science) doi:
 930 10.21273/JASHS.119.5.1006
- 931 Ershadi, A., McCabe, M. F., Evans, J. P., Chaney, N. W., & Wood, E. F. (2014,
 932 April). Multi-site evaluation of terrestrial evaporation models using FLUXNET
 933 data. *Agricultural and Forest Meteorology*, 187, 46–61. Retrieved 2020-
 934 01-15, from [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S0168192313002980)
 935 [S0168192313002980](http://www.sciencedirect.com/science/article/pii/S0168192313002980) doi: 10.1016/j.agrformet.2013.11.008
- 936 Ershadi, A., McCabe, M. F., Evans, J. P., & Wood, E. F. (2015, June). Impact of
 937 model structure and parameterization on Penman–Monteith type evaporation
 938 models. *Journal of Hydrology*, 525, 521–535. Retrieved 2020-06-22, from
 939 <http://www.sciencedirect.com/science/article/pii/S0022169415002577>
 940 doi: 10.1016/j.jhydrol.2015.04.008
- 941 Ferguson, C. R., Sheffield, J., Wood, E. F., & Gao, H. (2010, August). Quan-
 942 tifying uncertainty in a remote sensing-based estimate of evapotranspi-
 943 ration over continental USA. *International Journal of Remote Sens-*
 944 *ing*, 31(14), 3821–3865. Retrieved 2020-09-03, from [https://doi.org/](https://doi.org/10.1080/01431161.2010.483490)
 945 [10.1080/01431161.2010.483490](https://doi.org/10.1080/01431161.2010.483490) (Publisher: Taylor & Francis _eprint:
 946 <https://doi.org/10.1080/01431161.2010.483490>) doi: 10.1080/01431161.2010
 947 .483490
- 948 Ferretti, D. F., Pendall, E., Morgan, J. A., Nelson, J. A., LeCain, D., & Mosier,
 949 A. R. (2003, July). Partitioning evapotranspiration fluxes from a Colorado
 950 grassland using stable isotopes: Seasonal variations and ecosystem implica-
 951 tions of elevated atmospheric CO₂. *Plant and Soil*, 254(2), 291–303. Re-
 952 trieved 2021-04-19, from <https://doi.org/10.1023/A:1025511618571> doi:
 953 10.1023/A:1025511618571
- 954 Fisher, J. B., Lee, B., Purdy, A. J., Halverson, G. H., Dohlen, M. B.,
 955 Cawse-Nicholson, K., ... Hook, S. (2020). ECOSTRESS: NASA’s
 956 Next Generation Mission to Measure Evapotranspiration From the
 957 International Space Station. *Water Resources Research*, 56(4),
 958 e2019WR026058. Retrieved 2020-08-31, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019WR026058)
 959 [.onlinelibrary.wiley.com/doi/abs/10.1029/2019WR026058](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2019WR026058) (_eprint:
 960 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2019WR026058>) doi:

- 10.1029/2019WR026058
- 961
962 Fisher, J. B., Malhi, Y., Bonal, D., Rocha, H. R. D., Araújo, A. C. D., Gamo, M.,
963 ... Randow, C. V. (2009). The land-atmosphere water flux in the tropics.
964 *Global Change Biology*, 15(11), 2694–2714. Retrieved 2021-01-26, from
965 <https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1365-2486.2008.01813.x>
966 (.eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1365-2486.2008.01813.x>) doi: <https://doi.org/10.1111/j.1365-2486.2008.01813.x>
967
- 968 Fisher, J. B., Melton, F., Middleton, E., Hain, C., Anderson, M., Allen, R.,
969 ... Wood, E. F. (2017). The future of evapotranspiration: Global re-
970 quirements for ecosystem functioning, carbon and climate feedbacks,
971 agricultural management, and water resources. *Water Resources Re-*
972 *search*, 53(4), 2618–2626. Retrieved 2020-08-31, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016WR020175)
973 [.onlinelibrary.wiley.com/doi/abs/10.1002/2016WR020175](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016WR020175) (.eprint:
974 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2016WR020175>) doi:
975 10.1002/2016WR020175
- 976 Fisher, J. B., Tu, K. P., & Baldocchi, D. D. (2008, March). Global estimates of the
977 land-atmosphere water flux based on monthly AVHRR and ISLSCP-II data,
978 validated at 16 FLUXNET sites. *Remote Sensing of Environment*, 112(3),
979 901–919. Retrieved 2018-08-09, from [http://www.sciencedirect.com/](http://www.sciencedirect.com/science/article/pii/S0034425707003938)
980 [science/article/pii/S0034425707003938](http://www.sciencedirect.com/science/article/pii/S0034425707003938) doi: 10.1016/j.rse.2007.06.025
- 981 Fisher, J. B., Whittaker, R. J., & Malhi, Y. (2011). ET come home: potential
982 evapotranspiration in geographical ecology. *Global Ecology and Biogeog-*
983 *raphy*, 20(1), 1–18. Retrieved 2020-08-31, from [https://onlinelibrary](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1466-8238.2010.00578.x)
984 [.wiley.com/doi/abs/10.1111/j.1466-8238.2010.00578.x](https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1466-8238.2010.00578.x) (.eprint:
985 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/j.1466-8238.2010.00578.x>)
986 doi: 10.1111/j.1466-8238.2010.00578.x
- 987 Foken, T. (2008). The Energy Balance Closure Problem: An Overview. *Ecolog-*
988 *ical Applications*, 18(6), 1351–1367. Retrieved 2021-02-19, from [https://](https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/06-0922.1)
989 esajournals.onlinelibrary.wiley.com/doi/abs/10.1890/06-0922.1
990 (.eprint: [https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/06-](https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/06-0922.1)
991 [0922.1](https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1890/06-0922.1)) doi: <https://doi.org/10.1890/06-0922.1>
- 992 Gaj, M., Beyer, M., Koeniger, P., Wanke, H., Hamutoko, J., & Himmelsbach, T.
993 (2016, February). In situ unsaturated zone water stable isotope (^2H and
994 ^{18}O) measurements in semi-arid environments: a soil water balance. *Hy-*
995 *drology and Earth System Sciences*, 20(2), 715–731. Retrieved 2021-04-08,
996 from <https://hess.copernicus.org/articles/20/715/2016/> (Publisher:
997 Copernicus GmbH) doi: 10.5194/hess-20-715-2016
- 998 García, A. G., Di Bella, C. M., Houspanossian, J., Magliano, P. N., Jobbágy,
999 E. G., Posse, G., ... Nosetto, M. D. (2017, December). Patterns and
1000 controls of carbon dioxide and water vapor fluxes in a dry forest of central
1001 Argentina. *Agricultural and Forest Meteorology*, 247, 520–532. Retrieved
1002 2021-03-11, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192317302721)
1003 [S0168192317302721](https://www.sciencedirect.com/science/article/pii/S0168192317302721) doi: 10.1016/j.agrformet.2017.08.015
- 1004 García, M., Sandholt, I., Ceccato, P., Ridler, M., Mougin, E., Kergoat, L., ...
1005 Domingo, F. (2013, April). Actual evapotranspiration in drylands de-
1006 rived from in-situ and satellite data: Assessing biophysical constraints. *Re-*
1007 *remote Sensing of Environment*, 131, 103–118. Retrieved 2020-06-23, from
1008 <http://www.sciencedirect.com/science/article/pii/S0034425712004828>
1009 doi: 10.1016/j.rse.2012.12.016
- 1010 Gash, J. H. C. (1979). An analytical model of rainfall interception
1011 by forests. *Quarterly Journal of the Royal Meteorological Soci-*
1012 *ety*, 105(443), 43–55. Retrieved 2021-04-15, from [https://rmets](https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710544304)
1013 [.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710544304](https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.49710544304) (.eprint:
1014 <https://rmets.onlinelibrary.wiley.com/doi/pdf/10.1002/qj.49710544304>) doi:
1015 <https://doi.org/10.1002/qj.49710544304>

- 1016 Gash, J. H. C., Lloyd, C. R., & Lachaud, G. (1995, August). Estimat-
 1017 ing sparse forest rainfall interception with an analytical model. *Jour-*
 1018 *nal of Hydrology*, *170*(1), 79–86. Retrieved 2020-07-28, from [http://](http://www.sciencedirect.com/science/article/pii/002216949502697N)
 1019 www.sciencedirect.com/science/article/pii/002216949502697N doi:
 1020 10.1016/0022-1694(95)02697-N
- 1021 Goymer, P. (2017, March). Spotlight on South America. *Nature Ecology & Evo-*
 1022 *lution*, *1*(4), 1–2. Retrieved 2020-07-08, from [https://www.nature.com/](https://www.nature.com/articles/s41559-017-0129)
 1023 [articles/s41559-017-0129](https://www.nature.com/articles/s41559-017-0129) (Number: 4 Publisher: Nature Publishing
 1024 Group) doi: 10.1038/s41559-017-0129
- 1025 Harris, I., Osborn, T. J., Jones, P., & Lister, D. (2020, April). Version 4 of the
 1026 CRU TS monthly high-resolution gridded multivariate climate dataset. *Scien-*
 1027 *tific Data*, *7*(1), 109. Retrieved 2021-02-19, from [https://www.nature.com/](https://www.nature.com/articles/s41597-020-0453-3)
 1028 [articles/s41597-020-0453-3](https://www.nature.com/articles/s41597-020-0453-3) (Number: 1 Publisher: Nature Publishing
 1029 Group) doi: 10.1038/s41597-020-0453-3
- 1030 Hasler, N., & Avissar, R. (2007, June). What Controls Evapotranspiration in the
 1031 Amazon Basin? *Journal of Hydrometeorology*, *8*(3), 380–395. Retrieved 2020-
 1032 03-23, from <https://journals.ametsoc.org/doi/full/10.1175/JHM587.1>
 1033 (Publisher: American Meteorological Society) doi: 10.1175/JHM587.1
- 1034 Holl, D., Pancotto, V., Heger, A., Camargo, S., & Kutzbach, L. (2019). Cush-
 1035 ion bogs are stronger carbon dioxide net sinks than moss-dominated bogs
 1036 as revealed by eddy covariance measurements on tierra del fuego, argentina.
 1037 *Biogeosciences*, *16*, 3397-3423. doi: <https://doi.org/10.5194/bg-16-3397-2019>
- 1038 Huete, A., Didan, K., Miura, T., Rodriguez, E. P., Gao, X., & Ferreira, L. G. (2002,
 1039 November). Overview of the radiometric and biophysical performance of the
 1040 MODIS vegetation indices. *Remote Sensing of Environment*, *83*(1), 195–
 1041 213. Retrieved 2019-09-03, from [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S0034425702000962)
 1042 [article/pii/S0034425702000962](http://www.sciencedirect.com/science/article/pii/S0034425702000962) doi: 10.1016/S0034-4257(02)00096-2
- 1043 Hutyra, L. R., Munger, J. W., Saleska, S. R., Gottlieb, E., Daube, B. C., Dunn,
 1044 A. L., ... Wofsy, S. C. (2007). Seasonal controls on the exchange of carbon
 1045 and water in an Amazonian rain forest. *Journal of Geophysical Research:*
 1046 *Biogeosciences*, *112*(G3). Retrieved 2020-03-23, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JG000365)
 1047 [.onlinelibrary.wiley.com/doi/abs/10.1029/2006JG000365](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JG000365) (_eprint:
 1048 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006JG000365>) doi:
 1049 10.1029/2006JG000365
- 1050 Jang, K., Kang, S., Lim, Y.-J., Jeong, S., Kim, J., Kimball, J. S., & Hong,
 1051 S. Y. (2013). Monitoring daily evapotranspiration in North-
 1052 east Asia using MODIS and a regional Land Data Assimilation Sys-
 1053 tem. *Journal of Geophysical Research: Atmospheres*, *118*(23),
 1054 12,927–12,940. Retrieved 2021-04-15, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020639)
 1055 [.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020639](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JD020639) (_eprint:
 1056 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2013JD020639>) doi:
 1057 <https://doi.org/10.1002/2013JD020639>
- 1058 Jarchow, C. J., Nagler, P. L., Glenn, E. P., Ramírez-Hernández, J., & Rodríguez-
 1059 Burgueño, J. E. (2017, September). Evapotranspiration by remote sens-
 1060 ing: An analysis of the Colorado River Delta before and after the Minute
 1061 319 pulse flow to Mexico. *Ecological Engineering*, *106*, 725–732. Retrieved
 1062 2019-09-05, from [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S0925857416305833)
 1063 [S0925857416305833](http://www.sciencedirect.com/science/article/pii/S0925857416305833) doi: 10.1016/j.ecoleng.2016.10.056
- 1064 Jasechko, S., Sharp, Z. D., Gibson, J. J., Birks, S. J., Yi, Y., & Fawcett, P. J. (2013,
 1065 April). Terrestrial water fluxes dominated by transpiration. *Nature*, *496*(7445),
 1066 347–350. Retrieved 2021-04-13, from [https://www.nature.com/articles/](https://www.nature.com/articles/nature11983)
 1067 [nature11983](https://www.nature.com/articles/nature11983) (Number: 7445 Publisher: Nature Publishing Group) doi:
 1068 10.1038/nature11983
- 1069 Junk, W. J., Brown, M., Campbell, I. C., Finlayson, M., Gopal, B., Ramberg, L.,
 1070 & Warner, B. G. (2006, October). The comparative biodiversity of seven

- 1071 globally important wetlands: a synthesis. *Aquatic Sciences*, 68(3), 400–414.
 1072 Retrieved 2021-04-22, from <https://doi.org/10.1007/s00027-006-0856-z>
 1073 doi: 10.1007/s00027-006-0856-z
- 1074 Junk, W. J., da Cunha, C. N., Wantzen, K. M., Petermann, P., Strüßmann, C.,
 1075 Marques, M. I., & Adis, J. (2006, October). Biodiversity and its conservation
 1076 in the Pantanal of Mato Grosso, Brazil. *Aquatic Sciences*, 68(3), 278–309.
 1077 Retrieved 2021-04-22, from <https://doi.org/10.1007/s00027-006-0851-4>
 1078 doi: 10.1007/s00027-006-0851-4
- 1079 Khan, M. S., Liaqat, U. W., Baik, J., & Choi, M. (2018, April). Stand-alone uncer-
 1080 tainty characterization of GLEAM, GLDAS and MOD16 evapotranspiration
 1081 products using an extended triple collocation approach. *Agricultural and*
 1082 *Forest Meteorology*, 252, 256–268. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/S0168192318300224)
 1083 www.sciencedirect.com/science/article/pii/S0168192318300224 doi:
 1084 10.1016/j.agrformet.2018.01.022
- 1085 Khosa, F. V., Feig, G. T., van der Merwe, M. R., Mateyisi, M. J., Mudau, A. E.,
 1086 & Savage, M. J. (2019, December). Evaluation of modeled actual evap-
 1087 otranspiration estimates from a land surface, empirical and satellite-based
 1088 models using in situ observations from a South African semi-arid savanna
 1089 ecosystem. *Agricultural and Forest Meteorology*, 279, 107706. Retrieved
 1090 2021-04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192319303223)
 1091 [S0168192319303223](https://www.sciencedirect.com/science/article/pii/S0168192319303223) doi: 10.1016/j.agrformet.2019.107706
- 1092 Kustas, W. P., & Norman, J. M. (1999, April). Evaluation of soil and veg-
 1093 etation heat flux predictions using a simple two-source model with ra-
 1094 diometric temperatures for partial canopy cover. *Agricultural and For-*
 1095 *est Meteorology*, 94(1), 13–29. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/S0168192399000052)
 1096 www.sciencedirect.com/science/article/pii/S0168192399000052 doi:
 1097 10.1016/S0168-1923(99)00005-2
- 1098 Kutzbach, L. (2019a). Lars Kutzbach (2019), AmeriFlux AR-TF1 Rio Moat bog,
 1099 Ver. 1-5, AmeriFlux AMP, (Dataset).. Retrieved 2021-04-15, from [https://](https://ameriflux.lbl.gov/doi/AmeriFlux/AR-TF1)
 1100 ameriflux.lbl.gov/doi/AmeriFlux/AR-TF1 doi: [https://doi.org/10.17190/](https://doi.org/10.17190/AMF/1543389)
 1101 [AMF/1543389](https://doi.org/10.17190/AMF/1543389)
- 1102 Kutzbach, L. (2019b). Lars Kutzbach (2019), AmeriFlux AR-TF2 Rio Pipo bog,
 1103 Ver. 1-5, AmeriFlux AMP, (Dataset).. Retrieved 2021-04-15, from [https://](https://ameriflux.lbl.gov/sites/siteinfo/AR-TF2)
 1104 ameriflux.lbl.gov/sites/siteinfo/AR-TF2 doi: [https://doi.org/10.17190/](https://doi.org/10.17190/AMF/1543388)
 1105 [AMF/1543388](https://doi.org/10.17190/AMF/1543388)
- 1106 Leopoldo, P. R., Franken, W. K., & Villa Nova, N. A. (1995, May). Real evap-
 1107 otranspiration and transpiration through a tropical rain forest in central
 1108 Amazonia as estimated by the water balance method. *Forest Ecology*
 1109 *and Management*, 73(1), 185–195. Retrieved 2021-04-08, from [https://](https://www.sciencedirect.com/science/article/pii/037811279403487H)
 1110 www.sciencedirect.com/science/article/pii/037811279403487H doi:
 1111 10.1016/0378-1127(94)03487-H
- 1112 Levy, P., Drewer, J., Jammet, M., Leeson, S., Friborg, T., Skiba, U., & Oijen, M. v.
 1113 (2020, January). Inference of spatial heterogeneity in surface fluxes from
 1114 eddy covariance data: A case study from a subarctic mire ecosystem. *Agri-*
 1115 *cultural and Forest Meteorology*, 280, 107783. Retrieved 2020-07-10, from
 1116 <http://www.sciencedirect.com/science/article/pii/S0168192319303995>
 1117 doi: 10.1016/j.agrformet.2019.107783
- 1118 Li, X., Long, D., Han, Z., Scanlon, B. R., Sun, Z., Han, P., & Hou, A. (2019,
 1119 November). Evapotranspiration Estimation for Tibetan Plateau Headwa-
 1120 ters Using Conjoint Terrestrial and Atmospheric Water Balances and Mul-
 1121 tisource Remote Sensing. *Water Resources Research*, 55(11), 8608–8630.
 1122 Retrieved 2021-04-15, from [https://agupubs.onlinelibrary.wiley.com/](https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2019WR025196)
 1123 [doi/10.1029/2019WR025196](https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2019WR025196) (Publisher: John Wiley & Sons, Ltd) doi:
 1124 10.1029/2019WR025196
- 1125 Liu, Y. Y., Dijk, A. I. J. M. v., McCabe, M. F., Evans, J. P., & Jeu, R. A. M. d.

- 1126 (2013). Global vegetation biomass change (1988–2008) and attribu-
 1127 tion to environmental and human drivers. *Global Ecology and Bio-*
 1128 *geography*, 22(6), 692–705. Retrieved 2021-04-15, from [https://](https://onlinelibrary.wiley.com/doi/abs/10.1111/geb.12024)
 1129 onlinelibrary.wiley.com/doi/abs/10.1111/geb.12024 (_eprint:
 1130 <https://onlinelibrary.wiley.com/doi/pdf/10.1111/geb.12024>) doi: [https://](https://doi.org/10.1111/geb.12024)
 1131 doi.org/10.1111/geb.12024
- 1132 Lopes, J. D., Rodrigues, L. N., Imbuzeiro, H. M. A., & Pruski, F. F. (2019,
 1133 September). Performance of SSEBop model for estimating wheat actual
 1134 evapotranspiration in the Brazilian Savannah region. *International Jour-*
 1135 *nal of Remote Sensing*, 40(18), 6930–6947. Retrieved 2021-04-15, from
 1136 <https://doi.org/10.1080/01431161.2019.1597304> (Publisher: Tay-
 1137 lor & Francis _eprint: <https://doi.org/10.1080/01431161.2019.1597304>) doi:
 1138 [10.1080/01431161.2019.1597304](https://doi.org/10.1080/01431161.2019.1597304)
- 1139 Machado, C. B., Lima, J. R. d. S., Antonino, A. C. D., Souza, E. S. d., Souza,
 1140 R. M. S., Alves, E. M., ... Alves, E. M. (2016, September). Daily and sea-
 1141 sonal patterns of CO₂ fluxes and evapotranspiration in maize-grass intercrop-
 1142 ping. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 20(9), 777–782.
 1143 Retrieved 2021-03-11, from [http://www.scielo.br/scielo.php?script=](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S1415-43662016000900777&lng=en&nrm=iso&tlng=en)
 1144 [sci_abstract&pid=S1415-43662016000900777&lng=en&nrm=iso&tlng=en](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S1415-43662016000900777&lng=en&nrm=iso&tlng=en)
 1145 (Publisher: Departamento de Engenharia Agrícola - UFCG / Cnpq) doi:
 1146 [10.1590/1807-1929/agriambi.v20n9p777-782](https://doi.org/10.1590/1807-1929/agriambi.v20n9p777-782)
- 1147 Mao, J., Fu, W., Shi, X., Ricciuto, D. M., Fisher, J. B., Dickinson, R. E., ... Zhu,
 1148 Z. (2015, sep). Disentangling climatic and anthropogenic controls on global
 1149 terrestrial evapotranspiration trends. *Environmental Research Letters*, 10(9),
 1150 094008. Retrieved from <https://doi.org/10.1088/1748-9326/10/9/094008>
 1151 doi: [10.1088/1748-9326/10/9/094008](https://doi.org/10.1088/1748-9326/10/9/094008)
- 1152 Marques, T. V., Mendes, K., Mutti, P., Medeiros, S., Silva, L., Perez-Marin, A. M.,
 1153 ... Bezerra, B. (2020, June). Environmental and biophysical controls of evap-
 1154 otranspiration from Seasonally Dry Tropical Forests (Caatinga) in the Brazil-
 1155 ian Semiarid. *Agricultural and Forest Meteorology*, 287, 107957. Retrieved
 1156 2021-04-22, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192320300599)
 1157 [S0168192320300599](https://www.sciencedirect.com/science/article/pii/S0168192320300599) doi: [10.1016/j.agrformet.2020.107957](https://doi.org/10.1016/j.agrformet.2020.107957)
- 1158 Martens, B., Miralles, D., Lievens, H., Fernández-Prieto, D., & Verhoest, N. E. C.
 1159 (2016, June). Improving terrestrial evaporation estimates over continental
 1160 Australia through assimilation of SMOS soil moisture. *International Journal*
 1161 *of Applied Earth Observation and Geoinformation*, 48, 146–162. Retrieved
 1162 2021-04-13, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0303243415300350)
 1163 [S0303243415300350](https://www.sciencedirect.com/science/article/pii/S0303243415300350) doi: [10.1016/j.jag.2015.09.012](https://doi.org/10.1016/j.jag.2015.09.012)
- 1164 Martens, B., Miralles, D. G., Lievens, H., Schalie, R. v. d., Jeu, R. A. M. d.,
 1165 Fernández-Prieto, D., ... Verhoest, N. E. C. (2017, May). GLEAM v3:
 1166 satellite-based land evaporation and root-zone soil moisture. *Geosci-*
 1167 *entific Model Development*, 10(5), 1903–1925. Retrieved 2020-07-01, from
 1168 <https://gmd.copernicus.org/articles/10/1903/2017/> (Publisher: Coper-
 1169 nicus GmbH) doi: <https://doi.org/10.5194/gmd-10-1903-2017>
- 1170 Mauder, M., Foken, T., & Cuxart, J. (2020, December). Surface-Energy-Balance
 1171 Closure over Land: A Review. *Boundary-Layer Meteorology*, 177(2), 395–426.
 1172 Retrieved 2021-02-19, from <https://doi.org/10.1007/s10546-020-00529-6>
 1173 doi: [10.1007/s10546-020-00529-6](https://doi.org/10.1007/s10546-020-00529-6)
- 1174 McCabe, M. F., Ershadi, A., Jimenez, C., Miralles, D. G., Michel, D., & Wood,
 1175 E. F. (2016, January). The GEWEX LandFlux project: evaluation of model
 1176 evaporation using tower-based and globally gridded forcing data. *Geosci-*
 1177 *entific Model Development*, 9(1), 283–305. Retrieved 2019-08-29, from
 1178 <https://www.geosci-model-dev.net/9/283/2016/> doi: [https://doi.org/](https://doi.org/10.5194/gmd-9-283-2016)
 1179 [10.5194/gmd-9-283-2016](https://doi.org/10.5194/gmd-9-283-2016)
- 1180 McColl, K. A. (2020). Practical and Theoretical Benefits of an Alterna-

- 1181 tive to the Penman-Monteith Evapotranspiration Equation. *Water Re-*
 1182 *sources Research*, 56(6), e2020WR027106. Retrieved 2021-09-30, from
 1183 <https://onlinelibrary.wiley.com/doi/abs/10.1029/2020WR027106>
 1184 (_eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1029/2020WR027106>)
 1185 doi: 10.1029/2020WR027106
- 1186 Michel, D., Jiménez, C., Miralles, D. G., Jung, M., Hirschi, M., Ershadi, A., ...
 1187 Fernández-Prieto, D. (2016, February). The WACMOS-ET project –
 1188 Part 1: Tower-scale evaluation of four remote-sensing-based evapotranspiration
 1189 algorithms. *Hydrology and Earth System Sciences*, 20(2), 803–822. Retrieved
 1190 2019-08-29, from <https://www.hydrol-earth-syst-sci.net/20/803/2016/>
 1191 doi: <https://doi.org/10.5194/hess-20-803-2016>
- 1192 Miralles, D. G., Gash, J. H., Holmes, T. R. H., Jeu, R. A. M. d., & Dol-
 1193 man, A. J. (2010). Global canopy interception from satel-
 1194 lite observations. *Journal of Geophysical Research: Atmo-*
 1195 *spheres*, 115(D16). Retrieved 2020-09-04, from <https://agupubs>
 1196 [.onlinelibrary.wiley.com/doi/abs/10.1029/2009JD013530](https://onlinelibrary.wiley.com/doi/abs/10.1029/2009JD013530) (_eprint:
 1197 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2009JD013530>) doi:
 1198 10.1029/2009JD013530
- 1199 Miralles, D. G., Holmes, T. R. H., Jeu, R. A. M. D., Gash, J. H., Meesters,
 1200 A. G. C. A., & Dolman, A. J. (2011, February). Global land-surface evapora-
 1201 tion estimated from satellite-based observations. *Hydrology and Earth System*
 1202 *Sciences*, 15(2), 453–469. Retrieved 2020-06-15, from <https://www.hydrol>
 1203 [-earth-syst-sci.net/15/453/2011/hess-15-453-2011.html](https://www.hydrol-earth-syst-sci.net/15/453/2011/hess-15-453-2011.html) (Publisher:
 1204 Copernicus GmbH) doi: <https://doi.org/10.5194/hess-15-453-2011>
- 1205 Miralles, D. G., Jiménez, C., Jung, M., Michel, D., Ershadi, A., McCabe, M. F.,
 1206 ... Fernández-Prieto, D. (2016, February). The WACMOS-ET project
 1207 – Part 2: Evaluation of global terrestrial evaporation data sets. *Hy-*
 1208 *drology and Earth System Sciences*, 20(2), 823–842. Retrieved 2019-08-
 1209 29, from <https://www.hydrol-earth-syst-sci.net/20/823/2016/> doi:
 1210 <https://doi.org/10.5194/hess-20-823-2016>
- 1211 Miranda, R. d. Q., Galvncio, J. D., Moura, M. S. B. d., Jones, C. A., & Sрни-
 1212 vasan, R. (2017). *Reliability of MODIS Evapotranspiration Products for*
 1213 *Heterogeneous Dry Forest: A Study Case of Caatinga* [Research Article]. Re-
 1214 trieved 2020-05-06, from <https://www.hindawi.com/journals/amete/2017/>
 1215 9314801/ (ISSN: 1687-9309 Library Catalog: www.hindawi.com Pages:
 1216 e9314801 Publisher: Hindawi Volume: 2017) doi: <https://doi.org/10.1155/>
 1217 2017/9314801
- 1218 Moesinger, L., Dorigo, W., de Jeu, R., van der Schalie, R., Scanlon, T., Teub-
 1219 ner, I., & Forkel, M. (2020, January). The global long-term microwave
 1220 Vegetation Optical Depth Climate Archive (VODCA). *Earth System*
 1221 *Science Data*, 12(1), 177–196. Retrieved 2021-04-13, from <https://>
 1222 essd.copernicus.org/articles/12/177/2020/ (Publisher: Copernicus
 1223 GmbH) doi: 10.5194/essd-12-177-2020
- 1224 Moreira, A. A., Ruhoff, A. L., Roberti, D. R., Souza, V. d. A., da Rocha, H. R., &
 1225 Paiva, R. C. D. d. (2019, August). Assessment of terrestrial water balance
 1226 using remote sensing data in South America. *Journal of Hydrology*, 575, 131–
 1227 147. Retrieved 2019-08-28, from <http://www.sciencedirect.com/science/>
 1228 [article/pii/S0022169419304664](http://www.sciencedirect.com/science/article/pii/S0022169419304664) doi: 10.1016/j.jhydrol.2019.05.021
- 1229 Mu, Q., Heinsch, F. A., Zhao, M., & Running, S. W. (2007, December). Develop-
 1230 ment of a global evapotranspiration algorithm based on MODIS and global
 1231 meteorology data. *Remote Sensing of Environment*, 111(4), 519–536. Re-
 1232 trieved 2017-09-21, from <http://www.sciencedirect.com/science/>
 1233 [article/pii/S0034425707001903](http://www.sciencedirect.com/science/article/pii/S0034425707001903) doi: 10.1016/j.rse.2007.04.015
- 1234 Mu, Q., Zhao, M., & Running, S. W. (2011, August). Improvements to a
 1235 MODIS global terrestrial evapotranspiration algorithm. *Remote Sens-*

- 1236 *ing of Environment*, 115(8), 1781–1800. Retrieved 2020-06-21, from
 1237 <http://www.sciencedirect.com/science/article/pii/S0034425711000691>
 1238 doi: 10.1016/j.rse.2011.02.019
- 1239 Mutti, P. R., da Silva, L. L., Medeiros, S. d. S., Dubreuil, V., Mendes, K. R.,
 1240 Marques, T. V., ... Bezerra, B. G. (2019, March). Basin scale rainfall-
 1241 evapotranspiration dynamics in a tropical semiarid environment during
 1242 dry and wet years. *International Journal of Applied Earth Observation*
 1243 *and Geoinformation*, 75, 29–43. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/S0303243418307244)
 1244 www.sciencedirect.com/science/article/pii/S0303243418307244 doi:
 1245 10.1016/j.jag.2018.10.007
- 1246 Myers, N., Mittermeier, R. A., Mittermeier, C. G., Fonseca, G. A. B. d., & Kent, J.
 1247 (2000, February). Biodiversity hotspots for conservation priorities. *Nature*,
 1248 403(6772), 853–858. Retrieved 2019-08-28, from [https://www.nature.com/](https://www.nature.com/articles/35002501)
 1249 [articles/35002501](https://www.nature.com/articles/35002501) doi: 10.1038/35002501
- 1250 Nagler, P. L., Cleverly, J., Glenn, E., Lampkin, D., Huete, A., & Wan, Z. (2005,
 1251 January). Predicting riparian evapotranspiration from MODIS vegetation
 1252 indices and meteorological data. *Remote Sensing of Environment*, 94(1), 17–
 1253 30. Retrieved 2020-06-19, from [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S0034425704002615)
 1254 [article/pii/S0034425704002615](http://www.sciencedirect.com/science/article/pii/S0034425704002615) doi: 10.1016/j.rse.2004.08.009
- 1255 Nagler, P. L., Glenn, E. P., Kim, H., Emmerich, W., Scott, R. L., Huxman, T. E.,
 1256 & Huete, A. R. (2007, August). Relationship between evapotranspiration and
 1257 precipitation pulses in a semiarid rangeland estimated by moisture flux towers
 1258 and MODIS vegetation indices. *Journal of Arid Environments*, 70(3), 443–
 1259 462. Retrieved 2019-09-05, from [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S0140196307000249)
 1260 [article/pii/S0140196307000249](http://www.sciencedirect.com/science/article/pii/S0140196307000249) doi: 10.1016/j.jaridenv.2006.12.026
- 1261 Nagler, P. L., Glenn, E. P., Nguyen, U., Scott, R. L., & Doody, T. (2013, Au-
 1262 gust). Estimating Riparian and Agricultural Actual Evapotranspiration
 1263 by Reference Evapotranspiration and MODIS Enhanced Vegetation In-
 1264 dex. *Remote Sensing*, 5(8), 3849–3871. Retrieved 2019-09-03, from
 1265 <https://www.mdpi.com/2072-4292/5/8/3849> doi: 10.3390/rs5083849
- 1266 Nagler, P. L., Morino, K., Murray, R. S., Osterberg, J., & Glenn, E. P. (2009, De-
 1267 cember). An Empirical Algorithm for Estimating Agricultural and Riparian
 1268 Evapotranspiration Using MODIS Enhanced Vegetation Index and Ground
 1269 Measurements of ET. I. Description of Method. *Remote Sensing*, 1(4), 1273–
 1270 1297. Retrieved 2019-09-03, from [https://www.mdpi.com/2072-4292/1/4/](https://www.mdpi.com/2072-4292/1/4/1273)
 1271 [1273](https://www.mdpi.com/2072-4292/1/4/1273) doi: 10.3390/rs1041273
- 1272 Norman, J. M., Kustas, W. P., & Humes, K. S. (1995, December). Source ap-
 1273 proach for estimating soil and vegetation energy fluxes in observations
 1274 of directional radiometric surface temperature. *Agricultural and For-
 1275 est Meteorology*, 77(3), 263–293. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/016819239502265Y)
 1276 www.sciencedirect.com/science/article/pii/016819239502265Y doi:
 1277 10.1016/0168-1923(95)02265-Y
- 1278 Nouri, H., Glenn, E. P., Beecham, S., Chavoshi Boroujeni, S., Sutton, P., Alagh-
 1279 mand, S., ... Nagler, P. (2016, June). Comparing Three Approaches of
 1280 Evapotranspiration Estimation in Mixed Urban Vegetation: Field-Based, Re-
 1281 mote Sensing-Based and Observational-Based Methods. *Remote Sensing*, 8(6),
 1282 492. Retrieved 2019-09-05, from <https://www.mdpi.com/2072-4292/8/6/492>
 1283 [doi: 10.3390/rs8060492](https://www.mdpi.com/2072-4292/8/6/492)
- 1284 Novick, K. A., Biederman, J. A., Desai, A. R., Litvak, M. E., Moore, D. J. P., Scott,
 1285 R. L., & Torn, M. S. (2018, February). The AmeriFlux network: A coalition
 1286 of the willing. *Agricultural and Forest Meteorology*, 249, 444–456. Retrieved
 1287 2021-04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192317303295)
 1288 [S0168192317303295](https://www.sciencedirect.com/science/article/pii/S0168192317303295) doi: 10.1016/j.agrformet.2017.10.009
- 1289 Oliveira, B. S., Moraes, E. C., Carrasco-Benavides, M., Bertani, G., & Mataveli,
 1290 G. A. V. (2018, August). Improved Albedo Estimates Implemented in the

- 1291 METRIC Model for Modeling Energy Balance Fluxes and Evapotranspiration over Agricultural and Natural Areas in the Brazilian Cerrado. *Remote*
 1292 *Sensing*, 10(8), 1181. Retrieved 2020-06-27, from [https://www.mdpi.com/](https://www.mdpi.com/2072-4292/10/8/1181)
 1293 2072-4292/10/8/1181 (Number: 8 Publisher: Multidisciplinary Digital
 1294 Publishing Institute) doi: 10.3390/rs10081181
 1295
- 1296 Oliveira, P. T. S., Wendland, E., Nearing, M. A., Scott, R. L., Rosolem, R.,
 1297 & da Rocha, H. R. (2015, June). The water balance components of
 1298 undisturbed tropical woodlands in the Brazilian cerrado. *Hydrology*
 1299 *and Earth System Sciences*, 19(6), 2899–2910. Retrieved 2019-09-05,
 1300 from <https://www.hydro1-earth-syst-sci.net/19/2899/2015/> doi:
 1301 <https://doi.org/10.5194/hess-19-2899-2015>
- 1302 Olivera-Guerra, L., Mattar, C., Merlin, O., Durán-Alarcón, C., Santamaría-
 1303 Artigas, A., & Fuster, R. (2017, June). An operational method for
 1304 the disaggregation of land surface temperature to estimate actual evapo-
 1305 transpiration in the arid region of Chile. *ISPRS Journal of Photogram-*
 1306 *metry and Remote Sensing*, 128, 170–181. Retrieved 2020-09-04, from
 1307 <http://www.sciencedirect.com/science/article/pii/S0924271616303690>
 1308 doi: 10.1016/j.isprsjprs.2017.03.014
- 1309 Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell,
 1310 G. V. N., Underwood, E. C., ... Kassem, K. R. (2001, November). Terres-
 1311 trial Ecoregions of the World: A New Map of Life on Earth A new global map
 1312 of terrestrial ecoregions provides an innovative tool for conserving biodiver-
 1313 sity. *BioScience*, 51(11), 933–938. Retrieved 2020-05-06, from [https://](https://academic.oup.com/bioscience/article/51/11/933/227116)
 1314 academic.oup.com/bioscience/article/51/11/933/227116 (Publisher:
 1315 Oxford Academic) doi: 10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2
- 1316 Paca, V. H. d. M., Espinoza-Dávalos, G. E., Hessels, T. M., Moreira, D. M., Comair,
 1317 G. F., & Bastiaanssen, W. G. M. (2019, February). The spatial variability
 1318 of actual evapotranspiration across the Amazon River Basin based on remote
 1319 sensing products validated with flux towers. *Ecological Processes*, 8(1), 6.
 1320 Retrieved 2020-03-23, from <https://doi.org/10.1186/s13717-019-0158-8>
 1321 doi: 10.1186/s13717-019-0158-8
- 1322 Paiva, C. M., França, G. B., Liu, W. T. H., & Filho, O. C. R. (2011, March).
 1323 A comparison of experimental energy balance components data and SE-
 1324 BAL model results in Dourados, Brazil. *International Journal of Re-*
 1325 *remote Sensing*, 32(6), 1731–1745. Retrieved 2021-04-15, from [https://](https://doi.org/10.1080/01431161003623425)
 1326 doi.org/10.1080/01431161003623425 (Publisher: Taylor & Fran-
 1327 cis eprint: <https://doi.org/10.1080/01431161003623425>) doi: 10.1080/
 1328 01431161003623425
- 1329 Paschalis, A., Fatichi, S., Pappas, C., & Or, D. (2018, October). Covariation of veg-
 1330 etation and climate constrains present and future T/ET variability. *Environ-*
 1331 *mental Research Letters*, 13(10), 104012. Retrieved 2021-03-22, from [https://](https://doi.org/10.1088/1748-9326/aae267)
 1332 doi.org/10.1088/1748-9326/aae267 (Publisher: IOP Publishing) doi: 10
 1333 .1088/1748-9326/aae267
- 1334 Pastorello, G., Trotta, C., Canfora, E., Chu, H., Christianson, D., Cheah, Y.-W.,
 1335 ... Papale, D. (2020, July). The FLUXNET2015 dataset and the ONE-
 1336 Flux processing pipeline for eddy covariance data. *Scientific Data*, 7(1),
 1337 225. Retrieved 2021-04-21, from [https://www.nature.com/articles/](https://www.nature.com/articles/s41597-020-0534-3)
 1338 [s41597-020-0534-3](https://www.nature.com/articles/s41597-020-0534-3) (Number: 1 Publisher: Nature Publishing Group) doi:
 1339 10.1038/s41597-020-0534-3
- 1340 Paul-Limoges, E., Wolf, S., Schneider, F. D., Longo, M., Moorcroft, P., Gharun,
 1341 M., & Damm, A. (2020, January). Partitioning evapotranspiration with
 1342 concurrent eddy covariance measurements in a mixed forest. *Agricultural*
 1343 *and Forest Meteorology*, 280, 107786. Retrieved 2021-04-11, from [https://](https://www.sciencedirect.com/science/article/pii/S0168192319304022)
 1344 www.sciencedirect.com/science/article/pii/S0168192319304022 doi:
 1345 10.1016/j.agrformet.2019.107786

- 1346 Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007, October). Updated world
 1347 map of the Köppen-Geiger climate classification. *Hydrology and Earth System*
 1348 *Sciences*, *11*(5), 1633–1644. Retrieved 2020-05-06, from [https://www.hydrology-](https://www.hydrology-earth-syst-sci.net/11/1633/2007/)
 1349 [earth-syst-sci.net/11/1633/2007/](https://www.hydrology-earth-syst-sci.net/11/1633/2007/) (Publisher: Copernicus GmbH) doi:
 1350 <https://doi.org/10.5194/hess-11-1633-2007>
- 1351 Poblete-Echeverría, C., & Ortega-Farías, S. (2012, November). Calibration and val-
 1352 idation of a remote sensing algorithm to estimate energy balance components
 1353 and daily actual evapotranspiration over a drip-irrigated Merlot vineyard. *Irriga-*
 1354 *tion Science*, *30*(6), 537–553. Retrieved 2021-04-15, from [https://doi.org/](https://doi.org/10.1007/s00271-012-0381-x)
 1355 [10.1007/s00271-012-0381-x](https://doi.org/10.1007/s00271-012-0381-x) doi: 10.1007/s00271-012-0381-x
- 1356 Potter, C. S., Randerson, J. T., Field, C. B., Matson, P. A., Vitousek, P. M.,
 1357 Mooney, H. A., & Klooster, S. A. (1993). Terrestrial ecosystem pro-
 1358 duction: A process model based on global satellite and surface data.
 1359 *Global Biogeochemical Cycles*, *7*(4), 811–841. Retrieved 2019-12-20, from
 1360 <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/93GB02725>
 1361 doi: 10.1029/93GB02725
- 1362 Priestley, C. H. B., & Taylor, R. J. (1972, February). On the Assessment of Sur-
 1363 face Heat Flux and Evaporation Using Large-Scale Parameters. *Monthly*
 1364 *Weather Review*, *100*(2), 81–92. Retrieved 2021-04-15, from [https://](https://journals.ametsoc.org/view/journals/mwre/100/2/1520-0493_1972_100_0081_otaosh_2_3_co_2.xml)
 1365 [journals.ametsoc.org/view/journals/mwre/100/2/1520-0493_1972_100](https://journals.ametsoc.org/view/journals/mwre/100/2/1520-0493_1972_100_0081_otaosh_2_3_co_2.xml)
 1366 [_0081_otaosh_2_3_co_2.xml](https://journals.ametsoc.org/view/journals/mwre/100/2/1520-0493_1972_100_0081_otaosh_2_3_co_2.xml) (Publisher: American Meteorological Society
 1367 Section: Monthly Weather Review) doi: 10.1175/1520-0493(1972)100<0081:
 1368 OTAOSH>2.3.CO;2
- 1369 Rahimzadegan, M., & Janani, A. (2019, May). Estimating evapotranspiration
 1370 of pistachio crop based on SEBAL algorithm using Landsat 8 satellite im-
 1371 agery. *Agricultural Water Management*, *217*, 383–390. Retrieved 2021-
 1372 04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S037837741831179X)
 1373 [S037837741831179X](https://www.sciencedirect.com/science/article/pii/S037837741831179X) doi: 10.1016/j.agwat.2019.03.018
- 1374 Restrepo-Coupe, N., da Rocha, H. R., Hutryra, L. R., da Araujo, A. C., Borma,
 1375 L. S., Christoffersen, B., ... Saleska, S. R. (2013, December). What drives the
 1376 seasonality of photosynthesis across the Amazon basin? A cross-site analysis of
 1377 eddy flux tower measurements from the Brasil flux network. *Agricultural and*
 1378 *Forest Meteorology*, *182-183*, 128–144. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/S0168192313001184)
 1379 www.sciencedirect.com/science/article/pii/S0168192313001184 doi:
 1380 10.1016/j.agrformet.2013.04.031
- 1381 Rocha, H. R. d., Manzi, A. O., Cabral, O. M., Miller, S. D., Goulden, M. L.,
 1382 Saleska, S. R., ... Maia, J. F. (2009). Patterns of water and heat flux across a
 1383 biome gradient from tropical forest to savanna in Brazil. *Journal of Geophys-*
 1384 *ical Research: Biogeosciences*, *114*(G1). Retrieved 2019-08-30, from [https://](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000640)
 1385 agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007JG000640 doi:
 1386 10.1029/2007JG000640
- 1387 Rodrigues, T. R., Vourlitis, G. L., Lobo, F. d. A., Oliveira, R. G. d., &
 1388 Nogueira, J. d. S. (2014). Seasonal variation in energy balance and
 1389 canopy conductance for a tropical savanna ecosystem of south cen-
 1390 tral Mato Grosso, Brazil. *Journal of Geophysical Research: Biogeo-*
 1391 *sciences*, *119*(1), 1–13. Retrieved 2020-05-12, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JG002472)
 1392 [.onlinelibrary.wiley.com/doi/abs/10.1002/2013JG002472](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013JG002472) (eprint:
 1393 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2013JG002472>) doi:
 1394 10.1002/2013JG002472
- 1395 Rodrigues, T. R., Vourlitis, G. L., Lobo, F. d. A., Santanna, F. B., de Arruda,
 1396 P. H. Z., & Nogueira, J. d. S. (2016, March). Modeling canopy con-
 1397 ductance under contrasting seasonal conditions for a tropical savanna
 1398 ecosystem of south central Mato Grosso, Brazil. *Agricultural and For-*
 1399 *est Meteorology*, *218-219*, 218–229. Retrieved 2020-08-31, from [http://](http://www.sciencedirect.com/science/article/pii/S0168192315300216)
 1400 www.sciencedirect.com/science/article/pii/S0168192315300216 doi:

- 1401 10.1016/j.agrformet.2015.12.060
- 1402 Roerink, G. J., Su, Z., & Menenti, M. (2000, January). S-SEBI: A simple remote
1403 sensing algorithm to estimate the surface energy balance. *Physics and Chem-*
1404 *istry of the Earth, Part B: Hydrology, Oceans and Atmosphere*, 25(2), 147–
1405 157. Retrieved 2021-04-15, from [https://www.sciencedirect.com/science/](https://www.sciencedirect.com/science/article/pii/S1464190999001288)
1406 [article/pii/S1464190999001288](https://www.sciencedirect.com/science/article/pii/S1464190999001288) doi: 10.1016/S1464-1909(99)00128-8
- 1407 Ruhoff, A. L., Paz, A. R., Aragao, L. E. O. C., Mu, Q., Malhi, Y., Collischonn, W.,
1408 ... Running, S. W. (2013, November). Assessment of the MODIS global
1409 evapotranspiration algorithm using eddy covariance measurements and hy-
1410 drological modelling in the Rio Grande basin. *Hydrological Sciences Journal*,
1411 58(8), 1658–1676. Retrieved 2017-09-21, from [http://dx.doi.org/10.1080/](http://dx.doi.org/10.1080/02626667.2013.837578)
1412 [02626667.2013.837578](http://dx.doi.org/10.1080/02626667.2013.837578) doi: 10.1080/02626667.2013.837578
- 1413 Running, S. W., Mu, Q., Zhao, M., & Moreno, A. (2019, January). *User's*
1414 *Guide: MODIS Global Terrestrial Evapotranspiration (ET) Product, Ver-*
1415 *sion 2.0*. Retrieved from [https://modis-land.gsfc.nasa.gov/pdf/](https://modis-land.gsfc.nasa.gov/pdf/MOD16UsersGuideV2.022019.pdf)
1416 [MOD16UsersGuideV2.022019.pdf](https://modis-land.gsfc.nasa.gov/pdf/MOD16UsersGuideV2.022019.pdf)
- 1417 Rwasoka, D. T., Gumindoga, W., & Gwenzi, J. (2011, January). Estimation of
1418 actual evapotranspiration using the Surface Energy Balance System (SEBS)
1419 algorithm in the Upper Manyame catchment in Zimbabwe. *Physics and*
1420 *Chemistry of the Earth, Parts A/B/C*, 36(14), 736–746. Retrieved 2021-
1421 04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S1474706511001513)
1422 [S1474706511001513](https://www.sciencedirect.com/science/article/pii/S1474706511001513) doi: 10.1016/j.pce.2011.07.035
- 1423 Saleska, S. R., Da Rocha, H. R., Huete, A. R., Nobre, A. D., Artaxo, P. E., &
1424 Shimabukuro, Y. E. (2013, July). LBA-ECO CD-32 Flux Tower Network
1425 Data Compilation, Brazilian Amazon: 1999-2006. *ORNL DAAC*. Retrieved
1426 2020-05-06, from https://daac.ornl.gov/cgi-bin/dsvviewer.pl?ds_id=1174
1427 doi: <https://doi.org/10.3334/ORNLDAAC/1174>
- 1428 Schotanus, P., Nieuwstadt, F., & De Bruin, H. (1983, May). Temperature mea-
1429 surement with a sonic anemometer and its application to heat and moisture
1430 fluxes. *Boundary-Layer Meteorology*, 26(1), 81–93. Retrieved 2021-10-01, from
1431 <https://doi.org/10.1007/BF00164332> doi: 10.1007/BF00164332
- 1432 Seddon, A. W. R., Macias-Fauria, M., Long, P. R., Benz, D., & Willis, K. J.
1433 (2016, March). Sensitivity of global terrestrial ecosystems to climate
1434 variability. *Nature*, 531(7593), 229–232. Retrieved 2020-07-08, from
1435 <https://www.nature.com/articles/nature16986> (Number: 7593 Pub-
1436 lisher: Nature Publishing Group) doi: 10.1038/nature16986
- 1437 Senay, G. B., Budde, M., Verdin, J. P., & Melesse, A. M. (2007, June). A Coupled
1438 Remote Sensing and Simplified Surface Energy Balance Approach to Estimate
1439 Actual Evapotranspiration from Irrigated Fields. *Sensors*, 7(6), 979–1000.
1440 Retrieved 2021-04-15, from <https://www.mdpi.com/1424-8220/7/6/979>
1441 (Number: 6 Publisher: Molecular Diversity Preservation International) doi:
1442 10.3390/s7060979
- 1443 Shuttleworth, W. J., & Pereira, H. C. (1988, April). Evaporation from Ama-
1444 zonian rainforest. *Proceedings of the Royal Society of London. Series*
1445 *B. Biological Sciences*, 233(1272), 321–346. Retrieved 2021-04-08, from
1446 <https://royalsocietypublishing.org/doi/10.1098/rspb.1988.0024>
1447 (Publisher: Royal Society) doi: 10.1098/rspb.1988.0024
- 1448 Silva, P. F. d., Lima, J. R. d. S., Antonino, A. C. D., Souza, R., Souza, E. S. d.,
1449 Silva, J. R. I., & Alves, E. M. (2017, December). Seasonal patterns of carbon
1450 dioxide, water and energy fluxes over the Caatinga and grassland in the semi-
1451 arid region of Brazil. *Journal of Arid Environments*, 147, 71–82. Retrieved
1452 2021-04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0140196317301672)
1453 [S0140196317301672](https://www.sciencedirect.com/science/article/pii/S0140196317301672) doi: 10.1016/j.jaridenv.2017.09.003
- 1454 Sánchez, J. M., López-Urrea, R., Valentín, F., Caselles, V., & Galve, J. M. (2019).
1455 Lysimeter assessment of the simplified two-source energy balance model

- 1456 and eddy covariance system to estimate vineyard evapotranspiration. *Agri-*
 1457 *cultural and Forest Meteorology*, 274, 172-183. Retrieved from [https://](https://www.sciencedirect.com/science/article/pii/S0168192319301777)
 1458 www.sciencedirect.com/science/article/pii/S0168192319301777 doi:
 1459 <https://doi.org/10.1016/j.agrformet.2019.05.006>
- 1460 Souza, L. S. B. d., Moura, M. S. B. d., Sedyama, G. C., Silva, T. G. F. d.,
 1461 Souza, L. S. B. d., Moura, M. S. B. d., ... Silva, T. G. F. d. (2015, Au-
 1462 gust). Balanço de energia e controle biofísico da evapotranspiração na
 1463 Caatinga em condições de seca intensa. *Pesquisa Agropecuária Brasileira*,
 1464 50(8), 627-636. Retrieved 2020-08-31, from [http://www.scielo.br/](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-204X2015000800627&lng=en&nrm=iso&tlng=pt)
 1465 [scielo.php?script=sci_abstract&pid=S0100-204X2015000800627&lng=](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-204X2015000800627&lng=en&nrm=iso&tlng=pt)
 1466 [en&nrm=iso&tlng=pt](http://www.scielo.br/scielo.php?script=sci_abstract&pid=S0100-204X2015000800627&lng=en&nrm=iso&tlng=pt) (Publisher: Embrapa Informação Tecnológica) doi:
 1467 10.1590/S0100-204X2015000800001
- 1468 Souza, V. d. A., Roberti, D. R., Ruhoff, A. L., Zimmer, T., Adamatti, D. S.,
 1469 Gonçalves, L. G. G. d., ... Moraes, O. L. L. d. (2019, September). Eval-
 1470 uation of MOD16 Algorithm over Irrigated Rice Paddy Using Flux Tower
 1471 Measurements in Southern Brazil. *Water*, 11(9), 1911. Retrieved 2020-06-23,
 1472 from <https://www.mdpi.com/2073-4441/11/9/1911> (Number: 9 Publisher:
 1473 Multidisciplinary Digital Publishing Institute) doi: 10.3390/w11091911
- 1474 Stoy, P. C., El-Madany, T. S., Fisher, J. B., Gentine, P., Gerken, T., Good, S. P.,
 1475 ... Wolf, S. (2019, October). Reviews and syntheses: Turning the chal-
 1476 lenges of partitioning ecosystem evaporation and transpiration into oppor-
 1477 tunities. *Biogeosciences*, 16(19), 3747-3775. Retrieved 2021-03-23, from
 1478 <https://bg.copernicus.org/articles/16/3747/2019/> (Publisher: Coper-
 1479 nicus GmbH) doi: 10.5194/bg-16-3747-2019
- 1480 Stoy, P. C., Mauder, M., Foken, T., Marcolla, B., Boegh, E., Ibrom, A., ... Var-
 1481 lagin, A. (2013, April). A data-driven analysis of energy balance closure
 1482 across FLUXNET research sites: The role of landscape scale heterogeneity.
 1483 *Agricultural and Forest Meteorology*, 171-172, 137-152. Retrieved 2021-
 1484 02-19, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192312003413)
 1485 [S0168192312003413](https://www.sciencedirect.com/science/article/pii/S0168192312003413) doi: 10.1016/j.agrformet.2012.11.004
- 1486 Su, Z. (2002, February). The Surface Energy Balance System (SEBS) for estima-
 1487 tion of turbulent heat fluxes. *Hydrology and Earth System Sciences*, 6(1), 85-
 1488 100. Retrieved 2021-04-15, from [https://hess.copernicus.org/articles/6/](https://hess.copernicus.org/articles/6/85/2002/)
 1489 [85/2002/](https://hess.copernicus.org/articles/6/85/2002/) (Publisher: Copernicus GmbH) doi: 10.5194/hess-6-85-2002
- 1490 Sun, X., Wilcox, B. P., & Zou, C. B. (2019, September). Evapotranspiration par-
 1491 titioning in dryland ecosystems: A global meta-analysis of in situ studies.
 1492 *Journal of Hydrology*, 576, 123-136. Retrieved 2021-04-08, from [https://](https://www.sciencedirect.com/science/article/pii/S0022169419305736)
 1493 www.sciencedirect.com/science/article/pii/S0022169419305736 doi:
 1494 10.1016/j.jhydrol.2019.06.022
- 1495 Sutanto, S. J., Wenninger, J., Coenders-Gerrits, A. M. J., & Uhlenbrook, S. (2012,
 1496 August). Partitioning of evaporation into transpiration, soil evaporation and
 1497 interception: a comparison between isotope measurements and a HYDRUS-1D
 1498 model. *Hydrology and Earth System Sciences*, 16(8), 2605-2616. Retrieved
 1499 2021-04-11, from <https://hess.copernicus.org/articles/16/2605/2012/>
 1500 (Publisher: Copernicus GmbH) doi: 10.5194/hess-16-2605-2012
- 1501 Talsma, C. J., Good, S. P., Jimenez, C., Martens, B., Fisher, J. B., Miralles, D. G.,
 1502 ... Purdy, A. J. (2018, October). Partitioning of evapotranspiration in remote
 1503 sensing-based models. *Agricultural and Forest Meteorology*, 260-261, 131-
 1504 143. Retrieved 2020-09-01, from [http://www.sciencedirect.com/science/](http://www.sciencedirect.com/science/article/pii/S016819231830162X)
 1505 [article/pii/S016819231830162X](http://www.sciencedirect.com/science/article/pii/S016819231830162X) doi: 10.1016/j.agrformet.2018.05.010
- 1506 Talsma, C. J., Good, S. P., Miralles, D. G., Fisher, J. B., Martens, B., Jimenez,
 1507 C., & Purdy, A. J. (2018, October). Sensitivity of Evapotranspiration Com-
 1508 ponents in Remote Sensing-Based Models. *Remote Sensing*, 10(10), 1601.
 1509 Retrieved 2020-06-29, from <https://www.mdpi.com/2072-4292/10/10/1601>
 1510 (Number: 10 Publisher: Multidisciplinary Digital Publishing Institute) doi:

- 1511 10.3390/rs10101601
- 1512 Teixeira, A. H. d. C., Bastiaanssen, W. G. M., Ahmad, M. D., & Bos, M. G.
1513 (2009, March). Reviewing SEBAL input parameters for assessing evap-
1514 otranspiration and water productivity for the Low-Middle São Francisco
1515 River basin, Brazil: Part A: Calibration and validation. *Agricultural
1516 and Forest Meteorology*, *149*(3), 462–476. Retrieved 2020-06-26, from
1517 <http://www.sciencedirect.com/science/article/pii/S0168192308002566>
1518 doi: 10.1016/j.agrformet.2008.09.016
- 1519 Teixeira, A. H. d. C., Scherer-Warren, M., Hernandez, F. B. T., Andrade, R. G., &
1520 Leivas, J. F. (2013, November). Large-Scale Water Productivity Assessments
1521 with MODIS Images in a Changing Semi-Arid Environment: A Brazilian
1522 Case Study. *Remote Sensing*, *5*(11), 5783–5804. Retrieved 2021-04-15, from
1523 <https://www.mdpi.com/2072-4292/5/11/5783> (Number: 11 Publisher:
1524 Multidisciplinary Digital Publishing Institute) doi: 10.3390/rs5115783
- 1525 Tetens, O. (1930). *Über einige meteorologische Begriffe*. Friedrich Vieweg & Sohn
1526 Akt.- Gesellschaft. Retrieved from [https://books.google.com.br/books?id=](https://books.google.com.br/books?id=ey5UtAEACAAJ)
1527 [ey5UtAEACAAJ](https://books.google.com.br/books?id=ey5UtAEACAAJ)
- 1528 Thornton, P. E. (1998). *Regional ecosystem simulation: Combining surface- and
1529 satellite-based observations to study linkages between terrestrial energy and
1530 mass budgets* (phdthesis). The University of Montana, Missoula, MT,.
- 1531 Tong, X., Zhang, J., Meng, P., Li, J., & Zheng, N. (2017, February). Envi-
1532 ronmental controls of evapotranspiration in a mixed plantation in North
1533 China. *International Journal of Biometeorology*, *61*(2), 227–238. Retrieved
1534 2021-04-15, from <https://doi.org/10.1007/s00484-016-1205-0> doi:
1535 10.1007/s00484-016-1205-0
- 1536 Tonti, N. E., Gassmann, M. I., & Pérez, C. F. (2018, December). First results
1537 of energy and mass exchange in a salt marsh on southeastern South Amer-
1538 ica. *Agricultural and Forest Meteorology*, *263*, 59–68. Retrieved 2021-
1539 03-11, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0168192318302612)
1540 [S0168192318302612](https://www.sciencedirect.com/science/article/pii/S0168192318302612) doi: 10.1016/j.agrformet.2018.08.001
- 1541 Trabucco, A., & Zomer, R. (2019, January). *Global Aridity Index and Poten-
1542 tial Evapotranspiration (ET0) Climate Database v2*. Retrieved 2020-05-
1543 06, from [https://figshare.com/articles/Global_Aridity_Index_and
1544 _Potential_Evapotranspiration_ET0_Climate_Database_v2/7504448](https://figshare.com/articles/Global_Aridity_Index_and_Potential_Evapotranspiration_ET0_Climate_Database_v2/7504448) doi:
1545 10.6084/m9.figshare.7504448.v3
- 1546 Trajano, E. (2019, January). Chapter 20 - Biodiversity in South America. In
1547 W. B. White, D. C. Culver, & T. Pipan (Eds.), *Encyclopedia of Caves (Third
1548 Edition)* (pp. 177–186). Academic Press. Retrieved 2020-07-08, from [http://
1549 www.sciencedirect.com/science/article/pii/B9780128141243000194](http://www.sciencedirect.com/science/article/pii/B9780128141243000194)
1550 doi: 10.1016/B978-0-12-814124-3.00019-4
- 1551 Twine, T. E., Kustas, W. P., Norman, J. M., Cook, D. R., Houser, P. R., Meyers,
1552 T. P., ... Wesely, M. L. (2000, June). Correcting eddy-covariance flux under-
1553 estimates over a grassland. *Agricultural and Forest Meteorology*, *103*(3), 279–
1554 300. Retrieved 2017-09-21, from [http://www.sciencedirect.com/science/
1555 article/pii/S0168192300001234](http://www.sciencedirect.com/science/article/pii/S0168192300001234) doi: 10.1016/S0168-1923(00)00123-4
- 1556 Valente, F., David, J. S., & Gash, J. H. C. (1997, March). Modelling interception
1557 loss for two sparse eucalypt and pine forests in central Portugal using reformu-
1558 lated Rutter and Gash analytical models. *Journal of Hydrology*, *190*(1), 141–
1559 162. Retrieved 2020-07-28, from [http://www.sciencedirect.com/science/
1560 article/pii/S0022169496030661](http://www.sciencedirect.com/science/article/pii/S0022169496030661) doi: 10.1016/S0022-1694(96)03066-1
- 1561 Valle Júnior, L. C. G., Ventura, T. M., Gomes, R. S. R., de S. Nogueira, J., de
1562 A. Lobo, F., Vourlitis, G. L., & Rodrigues, T. R. (2020, April). Comparative
1563 assessment of modelled and empirical reference evapotranspiration methods for
1564 a brazilian savanna. *Agricultural Water Management*, *232*, 106040. Retrieved
1565 2020-08-31, from <http://www.sciencedirect.com/science/article/pii/>

- 1566 S0378377419314489 doi: 10.1016/j.agwat.2020.106040
- 1567 Verhoef, A., & Allen, S. J. (2000, March). A SVAT scheme describing energy
1568 and CO₂ fluxes for multi-component vegetation: calibration and test for a
1569 Sahelian savannah. *Ecological Modelling*, 127(2), 245–267. Retrieved 2021-
1570 04-10, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0304380099002136)
1571 S0304380099002136 doi: 10.1016/S0304-3800(99)00213-6
- 1572 Verhoef, A., & Campbell, C. L. (2006). Evaporation Measurement.
1573 In *Encyclopedia of Hydrological Sciences*. American Cancer Soci-
1574 ety. Retrieved 2021-02-19, from [https://onlinelibrary.wiley](https://onlinelibrary.wiley.com/doi/abs/10.1002/0470848944.hsa043)
1575 .com/doi/abs/10.1002/0470848944.hsa043 (Section: 40 _eprint:
1576 <https://onlinelibrary.wiley.com/doi/pdf/10.1002/0470848944.hsa043>) doi:
1577 10.1002/0470848944.hsa043
- 1578 Villarreal, S., & Vargas, R. (2021). Representativeness of FLUXNET sites
1579 across Latin America. *Journal of Geophysical Research: Biogeosciences*,
1580 126(3), e2020JG006090. Retrieved 2021-02-19, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020JG006090)
1581 .onlinelibrary.wiley.com/doi/abs/10.1029/2020JG006090 doi:
1582 <https://doi.org/10.1029/2020JG006090>
- 1583 Vinukollu, R. K., Wood, E. F., Ferguson, C. R., & Fisher, J. B. (2011, March).
1584 Global estimates of evapotranspiration for climate studies using multi-sensor
1585 remote sensing data: Evaluation of three process-based approaches. *Re-*
1586 *remote Sensing of Environment*, 115(3), 801–823. Retrieved 2017-09-21, from
1587 <http://www.sciencedirect.com/science/article/pii/S0034425710003251>
1588 doi: 10.1016/j.rse.2010.11.006
- 1589 von Randow, C., Manzi, A. O., Kruijt, B., de Oliveira, P. J., Zanchi, F. B., Silva,
1590 R. L., ... Kabat, P. (2004, June). Comparative measurements and seasonal
1591 variations in energy and carbon exchange over forest and pasture in South
1592 West Amazonia. *Theoretical and Applied Climatology*, 78(1), 5–26. Retrieved
1593 2020-03-19, from <https://doi.org/10.1007/s00704-004-0041-z> doi:
1594 10.1007/s00704-004-0041-z
- 1595 Vourlitis, G. L., Nogueira, J. d. S., Lobo, F. d. A., Sendall, K. M., Paulo,
1596 S. R. d., Dias, C. A. A., ... Andrade, N. L. R. d. (2008). En-
1597 ergy balance and canopy conductance of a tropical semi-deciduous
1598 forest of the southern Amazon Basin. *Water Resources Re-*
1599 *search*, 44(3). Retrieved 2020-05-12, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006WR005526)
1600 .onlinelibrary.wiley.com/doi/abs/10.1029/2006WR005526 (_eprint:
1601 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1029/2006WR005526>) doi:
1602 10.1029/2006WR005526
- 1603 Wang, X., Huo, Z., Shukla, M. K., Wang, X., Guo, P., Xu, X., & Huang, G.
1604 (2020, February). Energy fluxes and evapotranspiration over irrigated
1605 maize field in an arid area with shallow groundwater. *Agricultural Wa-*
1606 *ter Management*, 228, 105922. Retrieved 2021-04-15, from [https://](https://www.sciencedirect.com/science/article/pii/S0378377419311023)
1607 www.sciencedirect.com/science/article/pii/S0378377419311023 doi:
1608 10.1016/j.agwat.2019.105922
- 1609 Wang, Z., Schaaf, C. B., Strahler, A. H., Chopping, M. J., Román, M. O., Shuai, Y.,
1610 ... Fitzjarrald, D. R. (2014, January). Evaluation of MODIS albedo product
1611 (MCD43A) over grassland, agriculture and forest surface types during dormant
1612 and snow-covered periods. *Remote Sensing of Environment*, 140, 60–77. Re-
1613 trieved 2019-06-14, from [http://www.sciencedirect.com/science/article/](http://www.sciencedirect.com/science/article/pii/S0034425713002836)
1614 [pii/S0034425713002836](http://www.sciencedirect.com/science/article/pii/S0034425713002836) doi: 10.1016/j.rse.2013.08.025
- 1615 Wardlow, B. D., Egbert, S. L., & Kastens, J. H. (2007, June). Analysis of time-series
1616 MODIS 250 m vegetation index data for crop classification in the U.S. Central
1617 Great Plains. *Remote Sensing of Environment*, 108(3), 290–310. Retrieved
1618 2020-06-22, from [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/S0034425706004949)
1619 S0034425706004949 doi: 10.1016/j.rse.2006.11.021
- 1620 Webb., E., Pearman, G., & Leuning, R. (1980). Correction of flux measurements for

- 1621 density effects due to heat and water vapour transfer. *Quarterly Journal of the*
 1622 *Royal Meteorological Society*, 106, 85-100.
- 1623 Wei, Z., Yoshimura, K., Wang, L., Miralles, D. G., Jasechko, S., & Lee,
 1624 X. (2017). Revisiting the contribution of transpiration to global
 1625 terrestrial evapotranspiration. *Geophysical Research Letters*,
 1626 44(6), 2792–2801. Retrieved 2021-03-22, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL072235)
 1627 [.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL072235](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2016GL072235) (eprint:
 1628 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2016GL072235>) doi:
 1629 <https://doi.org/10.1002/2016GL072235>
- 1630 Wilson, K., Goldstein, A., Falge, E., Aubinet, M., Baldocchi, D., Berbigier, P., ...
 1631 Verma, S. (2002, December). Energy balance closure at FLUXNET sites. *Agricul-*
 1632 *tural and Forest Meteorology*, 113(1), 223–243. Retrieved 2020-04-21, from
 1633 <http://www.sciencedirect.com/science/article/pii/S0168192302001090>
 1634 doi: 10.1016/S0168-1923(02)00109-0
- 1635 Xu, T., Guo, Z., Xia, Y., Ferreira, V. G., Liu, S., Wang, K., ... Zhao, C. (2019,
 1636 November). Evaluation of twelve evapotranspiration products from ma-
 1637 chine learning, remote sensing and land surface models over contermi-
 1638 nous United States. *Journal of Hydrology*, 578, 124105. Retrieved 2021-
 1639 04-15, from [https://www.sciencedirect.com/science/article/pii/](https://www.sciencedirect.com/science/article/pii/S0022169419308406)
 1640 [S0022169419308406](https://www.sciencedirect.com/science/article/pii/S0022169419308406) doi: 10.1016/j.jhydrol.2019.124105
- 1641 Yang, Y., & Roderick, M. L. (2019). Radiation, surface temperature and evapora-
 1642 tion over wet surfaces. *Quarterly Journal of the Royal Meteorological Society*,
 1643 145(720), 1118-1129. Retrieved from [https://rmets.onlinelibrary.wiley](https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3481)
 1644 [.com/doi/abs/10.1002/qj.3481](https://rmets.onlinelibrary.wiley.com/doi/abs/10.1002/qj.3481) doi: <https://doi.org/10.1002/qj.3481>
- 1645 Zhang, J., Zhang, S., Zhang, W., Liu, B., Gong, C., Jiang, M., ... Sheng, L. (2018,
 1646 August). Partitioning daily evapotranspiration from a marsh wetland using
 1647 stable isotopes in a semiarid region. *Hydrology Research*, 49(4), 1005–1015.
 1648 Retrieved 2021-04-11, from <https://doi.org/10.2166/nh.2017.005> doi:
 1649 [10.2166/nh.2017.005](https://doi.org/10.2166/nh.2017.005)
- 1650 Zhang, Y., Chiew, F. H. S., Peña-Arancibia, J., Sun, F., Li, H., & Leuning, R.
 1651 (2017). Global variation of transpiration and soil evaporation and the role
 1652 of their major climate drivers. *Journal of Geophysical Research: Atmo-*
 1653 *spheres*, 122(13), 6868–6881. Retrieved 2020-07-11, from [https://agupubs](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027025)
 1654 [.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027025](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2017JD027025) (eprint:
 1655 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2017JD027025>) doi:
 1656 [10.1002/2017JD027025](https://doi.org/10.1002/2017JD027025)
- 1657 Zhou, S., Yu, B., Zhang, Y., Huang, Y., & Wang, G. (2016). Parti-
 1658 tioning evapotranspiration based on the concept of underlying wa-
 1659 ter use efficiency. *Water Resources Research*, 52(2), 1160–1175.
 1660 Retrieved 2020-07-11, from [https://agupubs.onlinelibrary](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015WR017766)
 1661 [.wiley.com/doi/abs/10.1002/2015WR017766](https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015WR017766) (eprint:
 1662 <https://agupubs.onlinelibrary.wiley.com/doi/pdf/10.1002/2015WR017766>)
 1663 doi: 10.1002/2015WR017766