

Whither second-sphere coordination?

Article

Published Version

Creative Commons: Attribution-Noncommercial 3.0

Open Access

Liu, W., Das, P. J., Colquhoun, H. M. and Stoddart, J. F. (2022) Whither second-sphere coordination? CCS Chemistry, 4 (3). pp. 755-784. ISSN 2096-5745 doi: 10.31635/ccschem.021.202101286 Available at <https://centaur.reading.ac.uk/101568/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <https://doi.org/10.31635/ccschem.021.202101286>

Identification Number/DOI: 10.31635/ccschem.021.202101286

<<https://doi.org/10.31635/ccschem.021.202101286>>

Publisher: Chinese Chemical Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Whither Second-Sphere Coordination?

Wenqi Liu¹, Partha J. Das¹, Howard M. Colquhoun^{2*} & J. Fraser Stoddart^{1,3,4,5*}


¹Department of Chemistry, Northwestern University, Evanston, IL 60208, ²Department of Chemistry, University of Reading, Reading RG6 6DX, ³School of Chemistry, University of New South Wales, Sydney, NSW 2052, ⁴Department of Chemistry, Stoddart Institute of Molecular Science, Zhejiang University, Hangzhou 310027, ⁵ZJU-Hangzhou Global Scientific and Technological Innovation Center, Hangzhou 311215

*Corresponding authors: h.m.colquhoun@rdg.ac.uk; stoddart@northwestern.edu

Cite this: *CCS Chem.* **2022**, 4, 755–784

DOI: 10.31635/ccschem.021.202101286

The properties of coordination complexes are dictated by both the metals and the ligands. The use of molecular receptors as second-sphere ligands enables significant modulation of the chemical and physical properties of coordination complexes. In this minireview, we highlight recent advances in functional systems based on molecular receptors as second-sphere coordination ligands, as applied in molecular recognition, synthesis of mechanically interlocked molecules, separation of metals, catalysis, and biomolecular chemistry. These functional systems demonstrate that second-sphere coordination is an emerging and very promising strategy for addressing societal challenges in health, energy, and the environment.



Keywords: catalysis, coordination complexes, mechanically interlocked molecules, metal recovery, molecular recognition, supramolecular chemistry

Introduction

Coordination chemistry,¹ the investigation of linking ligands directly to a central atom—most often a metal center—is at the heart of modern inorganic, organic, and materials chemistry. The specific combinations of ligands and metals dictate the structures and properties of the resulting coordination complexes, which are applied in numerous technologies, including metal-organic frameworks (MOFs),^{2–16} optical and magnetic materials,^{17,18} catalysis,^{19,20} and biomedical theranostics.^{21–24} In the resulting coordination complexes, the ligands linked directly to the metal center are referred to (Figure 1) as the first sphere of coordination for the metal.^{25–28} Meanwhile, another set of ligands can bind to the first-sphere ligands of the coordination complex through noncovalent bonding

interactions, leading to second-sphere adducts. Thus, second-sphere coordination of metal centers affords adducts that are essentially complexes of complexes.

The concept of second-sphere coordination (Figure 1c) was advanced in the first instance by Chemistry Nobel Prize Laureate Alfred Werner^{29,30} in 1913. He showed that this concept could explain a number of experimental observations that could not be understood solely on the basis of first-sphere coordination.²⁹ These observations included (1) the formation of adducts between amines and coordinatively saturated complexes such as $[M(acac)_3]^{n+}$, (2) the presence of solvents in the crystal structures of many coordination complexes, and (3) the solvent and counterion-dependent properties associated with the optical rotations of chiral coordination complexes.

DOI: 10.31635/ccschem.021.202101286

Corrected Citation: *CCS Chem.* **2022**, 4, 755–784

Previous Citation: *CCS Chem.* **2021**, 3, 3436–3465

Link to VoR: <https://doi.org/10.31635/ccschem.021.202101286>




Figure 1 | Structural formulas depicting the similarity of the three-point binding models for (a) a primary alkylammonium ion $[RNH_3]^+$ and (b) a transition-metal ammine. (c) Graphical representation of the concept of second-sphere coordination. A transition metal (M) is coordinated with first-sphere ligands L^1 , which interact with second-sphere ligands L^2 through noncovalent bonding interactions. Adapted with permission from ref 35. Copyright 1983 Royal Society of Chemistry.

Beginning in the early 1980s, we sought to gain insight into the nature of second-sphere coordination by investigating the interactions between transition-metal complexes and synthetic molecular receptors.³¹ Crown ethers are well-known to form complexes with primary alkylammonium ions, $[RNH_3]^+$, through hydrogen bonding and ion-dipole interactions.^{32–34} It occurred to us that the geometric and electronic (stereoelectronic) features of an $[RNH_3]^+$ ion closely resemble (Figures 1a and 1b) those of a transition-metal ammine complex,³⁵ $[ML_x(NH_3)]^{n+}$. The recognition of this similarity led to our initial investigations of crown ether receptors as second-sphere ligands, resulting in the isolation of a novel series of adducts between crown ethers and transition-metal complexes,^{36–39} whose superstructures were quickly

established by X-ray crystallography. Figure 2 presents a collage [(a)–(s)] of some of the solid-state superstructures that are reproduced from a review²⁵ that was published in *Angewandte Chemie* on “Second-Sphere Coordination—A Novel Rôle for Molecular Receptors” in 1986. The review was the first one to be published in the chemical literature with a copious use of color: at that time, it cost us the equivalent of £10,000. One of the most remarkable examples (Figures 2j and 2k) was the 1:1 adduct formed³⁸ between dibenzo[30]crown-10 (DB30C10) and $[Pt(bpy)(NH_3)_2]^{2+}$. The fascinating feature of this superstructure is that, in addition to the three pairs of $[N-H\cdots O]$ hydrogen bonds between the oxygen atoms in the crown ether and the hydrogen atoms of the NH_3 ligands, there are close, complementary $[\pi\cdots\pi]$ stacking (charge transfer) interactions between the two π -electron-rich aromatic (catechol) rings in the receptor molecule and the π -electron-poor 2,2'-bipyridine ligand in the platinum-based substrate. This second-sphere coordination adduct was shown by 1H NMR spectroscopy to exist as a stable complex in solution as well as in the solid state.³⁸ The high stability of this adduct is sustained by multiple weak interactions between the receptor and the ligands on the transition-metal complex. In contrast, conventional second-sphere ligands, such as water or alcohols, only form weak adducts with metal complexes through monovalent binding, and the well-defined superstructures observed crystallographically in the solid state are not sustained in solution. For example, in the crystal superstructure³⁸ of a lanthanum complex $[La(tptz)(NO_3)_3(H_2O)]\cdot 2C_2H_5OH$, the two ethanol molecules form hydrogen bonds with the coordinated H_2O molecule. Although the solid-state superstructure is well-defined, there is no suggestion that it exists in solution. Molecular receptors, when employed as second-sphere ligands, lead to the formation of stable, robust, and predictable adducts in contrast to those adducts formed by weakly binding ligands observed only in the solid state.

It may be noted that the superstructure of the bipyridine-based second-sphere adduct shown in Figure 2 as a space-filling representation (j) and as a ball-and-stick representation (k) provided (Box 1) inspiration for subsequent research into complex formation between dibenzo-crown ethers and the dicationic bipyridinium herbicides Diquat, derived from 2,2'-bipyridine, and Paraquat, derived from 4,4'-bipyridine. The success of the latter investigation in turn led to the discovery of the tetracationic cyclophane, cyclobis(paraquat-*p*-phenylene), known as Blue Box, which has since proved a hugely valuable component in developing the field of mechanically interlocked molecules (MIMs) and eventually artificial molecular machines (AMMs).

Following our initial work on using synthetic molecular receptors to bind coordination complexes, there was a surge in the study of second-sphere coordination. Numerous macrocyclic and polycyclic crown ethers,^{39–50} cyclodextrins (Figures 3a and 3b),^{51–65} calixarenes




Figure 2 | A sampling of solid-state superstructures of 1:1 adducts between transition-metal complexes and macrocyclic receptors taken directly from the 1986 review on second-sphere coordination published in *Angewandte Chemie*, where the superstructures are presented with space-filling or ball-and-stick models for (a) $[\{trans\text{-Pt(PMe}_3\text{)Cl}_2(\text{NH}_3)_2\}_2 \cdot 18\text{C6}]$; (b) $[\{cis\text{-Pt(NH}_3\text{)}_2\text{Cl}_2 \cdot dma\}_2 \cdot 18\text{C6}]$; (c and d) $[\text{Cu(NH}_3\text{)}_4(\text{H}_2\text{O}) \cdot 18\text{C6}]_n^{2n+}$; (e) $[\text{Rh(cod)(NH}_3\text{)}_2 \cdot \text{DB21C7}]^+$; (f) $[\text{Rh(nbd)(NH}_3\text{)}_2 \cdot \text{DB24C8}]^+$; (g) $[\text{diammine-1,1-cyclobutanecarboxylatoplatinum(II)-}\alpha\text{-CD}]$; (h) $[\text{Pt(en)}_2 \cdot 18\text{C6}]_n^{2n+}$; (i) diamminebis-(1,5-cyclooctadiene)(μ -1,4-10,13-tetraoxa-7,16-diazacyclooctadecane- N^7, N^{16})-dirhodium bis(hexafluorophosphate); (j and k) $[\text{Pt(bpy)(NH}_3\text{)}_2 \cdot \text{DB3OC10}]_n^{2n+}$; (l) $[\text{SnCl}_4(\text{H}_2\text{O})_2 \cdot 18\text{C6}][\text{H}_2\text{O}]_2$; (m) $[\{trans\text{-Ir(CO)(CH}_3\text{CN)(PPh}_3\text{)}_2\}_2 \cdot 18\text{C6}]_n^{2n+}$; (n) $[\text{Rh(cod)(NH}_3\text{)}_2 \cdot \alpha\text{-CD}]^+$; (o) $[\text{Mn(H}_2\text{O)}_6 \cdot 18\text{C6}]_n^{n+}$ (ClO_4^- salt); (p and q) $[\text{trans-Pt(PMe}_3\text{)Cl}_2(\text{NH}_3) \cdot \text{DB18C6}]$; (r) $[\text{Rh(cod)(NH}_3\text{)}_2 \cdot \text{DB24C8}]^+$; (s) $[\text{Rh(cod)(NH}_3\text{)}_2 \cdot \text{DB3OC10}]^+$. The X-ray superstructure (j and k) of $[\text{Pt(bpy)(NH}_3\text{)}_2 \cdot \text{DB3OC10}]_n^{2n+}$ inside the highlighted rectangle will be discussed in Box 1. Adapted with permission from ref 25. Copyright 1986 John Wiley and Sons.

(Figure 3e),^{66–71} resorcinarenes (Figure 3d),⁷² pillararenes (Figure 3f),⁷³ cucurbiturils (Figure 3c),⁷⁴ and cyclophanes^{75–79} all proved to be efficient second-sphere ligands and even, in some examples, acted simultaneously as both first- and second-sphere coordination ligands (Figure 2i).^{80–83} As a result, many second-sphere coordination adducts were discovered by exploiting the whole range of noncovalent bonding interactions, including hydrogen bonding, charge transfer, $[\pi \cdots \pi]$ stacking, hydrophobic effects, and van der Waals interactions. Summaries of these early investigations of second-sphere coordination chemistry may be found in several reviews,^{25–28,84–90} published by us as well as by other groups.


Over the past two decades, there has been a resurgence of interest in second-sphere coordination. Advances in supramolecular, organic, inorganic, biomolecular, and materials chemistry have created^{91,92} new demands for molecular receptors capable of binding a wide range of

substrates, constructing MIMs, separating different metals, and catalyzing organic reactions. In such contexts, second-sphere coordination has emerged as a valuable strategy for solving previously intractable problems. This minireview highlights many new adducts based on molecular receptors as second-sphere coordination ligands.

Molecular Recognition

Second-sphere coordination can be applied to the design and construction of new supramolecular complexes by incorporating noncovalent binding sites into ligands, followed by their coordination to transition metals. The metal ion can participate in molecular recognition by a receptor molecule either indirectly via second-sphere coordination or directly via simultaneous first- and second-sphere coordination, targeting specific substrates by tailored receptor-substrate and metal-ligand interactions.

Box 1. | A 40-year research trail illustrating how one observation in 1981 led to another line of research and so on—and how this dynamic is as sustained over four decades.






Figure 3 | Representative examples of solid-state superstructures for second-sphere coordination adducts reported by other researchers. (a) Tubular and space-filling representation of a 2:2:1 adduct of γ -cyclodextrin•12-crown-4• K^+ . (b) Tubular and space-filling representation of a 4:5 adduct of β -cyclodextrin•ferrocene. (c) Tubular and ball-and-stick representation of a 1:1 adduct of cucurbit[6]uril• $[Yb(OH_2)_8]^{3+}$ • $[ReO_4]^-$. (d) Tubular and space-filling representation of a 1:1 adduct of resorcinarene•ferrocene. (e) Tubular and space-filling representation of a 1:1 adduct of $[Cu(NC_5H_5)_2(H_2O)_4]$ •calix[4]arene sulfonate. (f) Tubular and space-filling representation of a 1:1 adduct of $(n\text{-octyl})MgBr(THF)_2$ •pillar[5]arene.

Loeb et al.^{50,80–83,93,94} have demonstrated the design and synthesis of several molecular receptors that interact with substrates through second-sphere interactions. One example involves^{80,94} a family of thiacyclophane receptors that recognize DNA nucleobases. The molecular recognition displayed by these receptors involves several different types of interaction, including (1) first-sphere coordination from a N atom of the nucleobase substrate to a Pd(II) metal center, (2) second-sphere hydrogen bonding between a NH_2 group of the substrate and polyether O atoms of the receptors, and (3) $[\pi\cdots\pi]$ stacking between the π -electron-rich aromatic spacing units of the receptors and the π -electron-poor aromatic rings of the substrates. Single-crystal X-ray analyses of two adducts involving a receptor binding either with adenine or guanine—the latter as a BF_3 adduct—reveals that all three interactions mentioned above exist in these adducts with near-ideal interaction parameters (Figures 4a and 4c). The molecular recognition of cytosine was realized by a receptor having coordination sites for Pd(II) and polyether functional groups. Single-crystal X-ray analysis (Figure 4b) reveals that adduct formation between this receptor and cytosine is sustained by direct first-sphere coordination from a N atom of cytosine to the Pd(II) metal center, aided and abetted by hydrogen-bonding interactions between the NH_2 group of cytosine and polyether O atoms of the macrocyclic receptor. These adducts between the DNA nucleobases and their receptors have been proved to exist both in the solid state and in solution. The same principles have been applied^{81–83} in designing synthetic receptors for molecular

recognition of other substrates, including barbiturates, amino acids, amines, and hydrazinium ions.

Huc et al.^{95–98} have presented (Figure 5a) the chemical community with several examples of molecular recognition through second-sphere coordination using molecular capsules formed by metal-coordination-directed folding of a helical oligomer. A key chain segment,




Figure 4 | Stick representations of the X-ray superstructures of thiacyclophane receptors complexed with (a) adenine, (b) cytosine, and (c) guanine (BF_3) through a combination of several interactions, including (1) first-sphere coordination from N atoms to Pd(II) metal center, (2) second-sphere hydrogen bonds between the NH_2 group of the substrate and polyether O atoms of the receptors, and (3) $[\pi\cdots\pi]$ stacking between the electron-rich aromatic spacing units of the receptors and the electron-poor aromatic rings of the substrates.




Figure 5 | (a) Schematic illustration of molecular recognition through second-sphere coordination using helical molecular capsules formed by metal-coordination-directed folding. (b) Stick-and-sphere representation of X-ray crystal superstructure of an adduct between a β -D-mannopyranose and a helical foldamer ligand associated with Cu^{2+} . (c) A zoomed-in view of the substrate associated with the capsule through second-sphere coordination, which involves hydrogen-bond (dashed lines) formation between the foldamer ligand and the hydroxyl groups of the β -D-mannopyranose. The Cu^{2+} center adopts a square pyramidal geometry, coordinating with three nitrogen atoms of the pyz-pyr-pyz segment and two O atoms of two H_2O molecules. Adapted with permission from ref 97. Copyright 2018 Royal Society of Chemistry.

pyridazine-pyridine-pyridazine (pyz-pyr-pyz), was introduced into the oligomer.⁹⁵ This segment exists in an *anti-anti* conformation that is favored by repulsion between the endocyclic N atoms, directing the oligomer into an extended conformation instead of a capsular shape. As a result, this oligomer displays no propensity for molecular recognition in the absence of metal ions. Coordination of transition-metal ions Cu^+ , Cu^{2+} , and Ag^+ , as well as the alkali metal ions Na^+ and K^+ , induces folding of the oligomer into a capsular shape by favoring a *syn-syn* conformation for the pyz-pyr-pyz segment.⁹⁶ In all cases, the metal ions were positioned on one side of the cavity wall, leaving its coordination sphere occupied partially by solvent molecules and available to bind a guest. The first-sphere coordination between guests and the metal centers can enhance the binding affinity of the helix, while the second-sphere interaction between the guest and the inner wall of the capsule leads to shape and size selectivity. Moreover, interaction of the oligomer with a hydrated Ca^{2+} or Ba^{2+} ion occurs *entirely* in the second sphere, via hydrogen bonding between ion-coordinated water and oligomeric N atoms, but still results⁹⁵ in a helical geometry for the oligomeric complex.

As an extension of these earlier reports, the same authors demonstrated molecular recognition of

carbohydrates⁹⁷ in organic solvents using foldamer-type molecular capsules, by taking advantage of first- and second-sphere coordination. Depending on the nature of the metal center, the molecular capsules show different binding affinities and diastereoselectivities toward a range of carbohydrates, including D/L-threitol, xylitol, D/L-mannopyranose, D/L-glucopyranose, D/L-galactopyranose, and D/L-fructopyranose. For example, the Cu^{2+} complex of the foldamer binds D-threitol 14 times more strongly than its K^+ counterpart. X-Ray crystal superstructure analysis of the adducts between a Cu^{2+} -coordinated capsule and D/L-mannopyranose reveals (Figures 5b and 5c) that the substrate is associated with the capsule through second-sphere coordination, which involves hydrogen bond formation between the foldamer ligand and the hydroxy groups of the mannopyranose. The Cu^{2+} center adopts a square pyramidal geometry, coordinating with three N atoms of the pyz-pyr-pyz segment and the O atoms of two H_2O molecules. The D-mannopyranose assumes the β -pyranose configuration in the solid state. In contrast, this molecule exists predominantly (97%) in the α -pyranose form in $\text{CHCl}_3/\text{Me}_2\text{SO}$ (4:1, v/v) solution. The authors also reported the single-crystal superstructure of an adduct between D/L-threitol and a smaller analog of the molecular capsule coordinated with a Cu^{2+} ion. Here, the Cu^{2+} center interacts with the substrate through second-sphere coordination involving solvents—a H_2O and a MeOH molecule—that are coordinated directly to the metal ion. As an extension of such molecular recognition, the authors also developed a new strategy⁹⁸ to prepare the foldamer capsule shells around a $[\text{2Fe-2S}]$ cluster. The foldamer shell influences the structural and spectroscopic properties of the metal cluster, including desymmetrization and confinement of part of its first coordination sphere within the foldamer cavity.

Leigh et al.⁹⁹⁻¹⁰² have reported molecular recognition of chloride anions through second-sphere coordination using molecular knots. These molecular knots were synthesized (Figure 6a) by transmetallation of a Zn(II) infused pentafoil (5_1) knot with tetrafluoroborate salts of Co(II), Cu(II), and Ni(II).⁹⁹ The resulting metallated knots exhibit second-sphere coordination of a single chloride anion within the central cavity of the knot by utilizing both multiple $[\text{CH}\cdots\text{Cl}^-]$ hydrogen bonds and electrostatic interactions. X-ray crystal superstructures of these transmetallated pentafoil knots reveal (Figures 6b-6d) that five of the 15 bipyridine groups form an inner cavity lined with 10 electron-poor H atoms that form an array of hydrogen bonds with the chloride anion located inside the cavity. This type of second-sphere coordination is the first example exhibited in molecular-knot geometry. The diameter of the central cavity varies depending on the metal cation: 3.3 Å for Fe(II), Zn(II), and Ni(II), and 3.5 Å for Co(II). The metal-to-metal distance in the distorted cobalt knot is, however, smaller than those in the case of iron, nickel, and




Figure 6 | (a) Graphical illustration and structural formulas of various pentafoil knotted complexes synthesized by transmetalation of Zn(II) infused pentafoil (5_1) knot with tetrafluoroborate salts of Co(II), Cu(II), and Ni(II). The resulting metallated knots exhibit second-sphere coordination of a single chloride ion within the central cavity of the knot, providing $[\text{CH} \cdots \text{Cl}^-]$ hydrogen bonding and electrostatic interactions. (b) Ball-and-stick representations of the X-ray crystal superstructures of chloride complex using the pentafoil knots sustained by the coordination with (b) Co^{2+} , (c) Ni^{2+} , and (d) Zn^{2+} . The binding affinities for Cl^- anions are determined by the second-sphere coordination and electrostatic interactions with different $\text{M}(\text{II})$ ions.

zinc pentafoil knots. In contrast, the metal-metal-metal angle is higher in cobalt compared with those in the other three metallic pentafoil knots. These conformational changes in the knotted ligands lead to different distances between the chloride anion and the metal ions in these knot complexes. As a result, the binding affinities for chloride anion through second-sphere coordination and electrostatic interaction vary with different $\text{M}(\text{II})$ ions over nearly three orders of magnitude, from $K_a = 8 \times 10^4 \text{ M}^{-1}$ (Cu^{2+} knot) to $K_a = 3.3 \times 10^7 \text{ M}^{-1}$ (Fe^{2+} knot) in MeCN.

Construction of Mechanically Interlocked Molecules

The synthesis of MIMs^{103–108} such as catenanes,^{109–122} rotaxanes,^{123–132} and suitanes^{133–139} has become a field with intense research activity on account of the potential of such molecules to act as molecular machines.^{140–153} Several effective synthetic strategies for synthesizing MIMs have been achieved, based on the template effect.^{154–156}

One of these strategies involves the coordination of organic ligands to transition metals to promote the formation of mechanically interlocked structures.^{157–168} Second-sphere coordination, as an extension of this strategy, can also be employed to advance the effective synthesis of MIMs. Wisner et al.^{169–173} demonstrated the use of second-sphere coordination to prepare several MIMs such as rotaxanes and catenanes. For example, this group described the synthesis of [2]pseudorotaxanes in a single step employing second-sphere coordination.¹⁶⁹ An isophthalamide-based tetralactam macrocycle, serving as the second-sphere ligand, establishes four sets of hydrogen bonds with the two chloride first-sphere ligands in a *trans*-palladium dichloride complex, leading to formation of a pseudorotaxane. It was found that the stability of the pseudorotaxane diminishes markedly as the size of the halide ligand increases from Cl < Br < I. In a subsequent investigation, these authors reported¹⁷⁰ the synthesis of a [2]rotaxane based on the same system. To produce the mechanically interlocked architecture shown in Figure 7a, 4-(3,5-di-*tert*-butylbenzyloxy)-pyridine was employed as the ligand, on the basis that its terminal groups are sterically demanding and function as stoppers to prevent the dethreading of the tetralactam macrocycle. The rotaxane was obtained in 89% isolated yield simply by mixing the tetralactam macrocycle, the ligands, and *trans*-bis(benzonitrile)palladium(II) dichloride in CHCl₃ at room temperature. The resulting rotaxane is soluble in CHCl₃ and remains stable over time as well as under column chromatographic conditions. The structure of the rotaxane was confirmed by X-ray crystallographic analysis, which revealed (Figure 7b) that the amide groups of the isophthalamide subunits in the macrocycle establish four sets of hydrogen bonds with the chloride ligands of the palladium(II) metal complex, providing the driving force for rotaxane formation. Following these investigations, these same authors explored¹⁷² the stabilities of a series of pseudorotaxanes by varying the metals and the *para*-substituted pyridine ligand. They found that the hydrogen bond-accepting ability of the chloride ligands can be tuned by varying the electron-donating/-withdrawing nature of the *para*-substituted pyridine coligands. Moreover, the stability of the pseudorotaxane in CHCl₃ solution decreases slightly when the Pd(II) ion is replaced by Pt(II). Replacing chloride by thiocyanate in the first coordination sphere led to the formation (Figure 7c) of a series of (pseudo)rotaxanes by second-sphere coordination (hydrogen bonding) between the thiocyanate ligands and the amide units of the macrocycle.¹⁷¹ The directional nature of the thiocyanate ligands affords a doubly degenerate binding geometry, leading to a rotaxane (Figure 7c) in which the ring shuttles back and forth 3300 times a second in solution at 15 °C.

In another example, Wisner et al.¹⁷³ demonstrated the construction of catenanes (Figure 8a) using

simultaneous first- and second-sphere coordination. An acyclic bidentate ligand that resembles a three-quarter unit of a previously employed tetralactam macrocycle was designed and synthesized. This ligand coordinates directly to a PdCl₂ unit in a *trans* arrangement, and the resulting product comprises (Figure 8a) a pair of catenated macrocycles, each incorporating the metal subunit *trans*-PdCl₂L₂ in its scaffold. The catenation is driven by mutual recognition of the PdCl₂L₂ subunit in one ring by its orthogonally positioned partner. The catenane was synthesized in 87% isolated yield simply by heating a solution of an equimolar amount of Pd(PhCN)₂Cl₂ and the *trans*-bidentate ligand. X-ray crystallographic analysis of the catenane reveals (Figure 8b) that a template effect is manifested in the interaction between both the PdCl₂ subunits and their opposing macrocyclic cavities. Eight pairs of [NH...Cl] hydrogen bonds were found between the first-sphere Cl[−] ligands and the amide units of the second-sphere ligand. A reversible transformation between catenane and macrocycle was also demonstrated by varying the solvent polarity. Transformation of the catenane into two individual macrocycles was observed when the sample was dissolved in the mixed solvent system (CD₃)₂SO/CDCl₃ as a result of competitive binding of the polar solvent with the NH groups of the macrocycle. The catenane re-forms when the macrocycle is redissolved in CDCl₃.

Metal Separation

Second-sphere coordination of transition-metal complexes could lead to changes in solubility of the resulting adducts, which can be utilized for the separation of metal ions. We demonstrated (Figure 9c) the separation of copper from cobalt using³⁵ this principle in 1983. We found that 18C6 formed (Figure 9a) a 1:1 polymeric adduct with [Cu(NH₃)₄(H₂O)][PF₆]₂, leading to precipitation from aqueous solution. X-ray structural investigation revealed that each macrocycle is involved in no fewer than 10 hydrogen bonds with the ammine ligands. A similar adduct obtained (Figure 9b) from the cobalt(III) complex [Co(NH₃)₆][PF₆]₃, requires the addition of a two-molar excess of 18C6 to its aqueous solution in order to form a crystalline orange precipitate. When one molar equivalent of 18C6 was added to a solution containing one molar equivalent of each of [Co(NH₃)₆][PF₆]₃ and [Cu(NH₃)₄(H₂O)][PF₆]₂, the 1:1 copper adduct was precipitated exclusively in >90% yield. A simple filtration separated the precipitated copper adduct from the cobalt-containing solution. The 18C6 ligand was recovered by treating the 1:1 adduct with HCl solution, yielding aqueous copper(II) and insoluble [NH₄][PF₆]₃•18C6. The ammonium hexafluorophosphate adduct with 18C6




Figure 7 | (a) Structural formula and (b) tubular and space-filling representations of X-ray solid-state superstructure of a rotaxane constructed by second-sphere coordination. The amide groups of the isophthalamide subunits in the macrocycle establish four sets of hydrogen bonds with the chloride ligands of the palladium(II) metal complex, providing the driving force for the [2]rotaxane formation. (c) X-ray solid-state structure of a [2]rotaxane formed by second-sphere coordination between the thiocyanate ligands and the amide units of the macrocycle. The directional nature of the thiocyanate ligands affords a degenerate binding geometry, leading to a degenerate [2]rotaxane shuttle where the ring moves back and forth along the dumb-bell shaped axle 3300 times a second in CDCl_3 solution at 15 °C. Adapted with permission from ref 170. Copyright 2006 Royal Society of Chemistry.

could be used again in separating $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})]^{2+}$ from $[\text{Co}(\text{NH}_3)_6]^{3+}$.

In a similar fashion, just as ammine ligands on transition metals bind to crown ethers, aqua ligands coordinate²⁵ to electropositive metals through oxygen $[\text{M} \leftarrow \text{OH}_2]^{n+}$, showing an enhanced dipole and increased acidity of the ligand. Crown ethers,^{25,174} macrocyclic polyamines,¹⁷⁵ and acyclic trisulfonate ligands^{176–178} have all been reported as receptors for aqua complexes of

Mn^{3+} , Co^{2+} , Th^{4+} , U^{4+} , Sn^{4+} , Er^{3+} , and so on. The resulting adducts show different solubilities and propensities for crystallization, properties that can be utilized for metal separation. Shimizu et al.¹⁷⁶ have shown (Figure 10) that 1,3,5-tris(sulfomethyl)benzene trianions can be employed as second-sphere ligands for aqua complexes. The methylsulfonate groups on this ligand adopt a *cis-cis-cis* orientation, that is, they are all on the same side of the benzene core. The convergently oriented




Figure 8 | (a) Structural formula and (b) tubular and space-filling representations of X-ray solid-state structure of a catenane constructed by second-sphere coordination. The catenation formation is driven by the mutual recognition of the PdCl_2L_2 subunit in one ring by its orthogonally disposed partner. Eight pairs of $[\text{NH}\cdots\text{Cl}]$ hydrogen bonds were found between the first-sphere Cl ligands and the amide units of the second-sphere ligand. Adapted with permission from ref 173. Copyright 2007 John Wiley and Sons.

sulfonate groups interact with aqua ligands through hydrogen bonds. The authors found^{177,178} that trivalent cations (Fe^{3+} , Cr^{3+} , and Al^{3+}) can be sandwiched (Figures 10a–10c) between two of the ligands, forming 1:2 adducts. Each ligand participants in the formation of six hydrogen bonds with the aqua ligands. This trisulfonate ligand also forms (Figures 10d–10f) 1:1 adducts when divalent cations (Cu^{2+} , Mg^{2+} , and Zn^{2+}) are used. The 1:2 adducts, however, showed a better propensity to form crystals. Competitive crystallization was performed when equal molar amounts of the ligand, $\text{Fe}(\text{NO}_3)_3$, and $\text{Zn}(\text{NO}_3)_2$ were combined in an aqueous solution. Diffusion of MeOH into this mixture afforded only yellow crystals of the 1:1 adduct between the ligand and Fe^{3+} aqua complex, leaving all the Zn^{2+} in solution. This group also has demonstrated^{179–181} the construction of porous materials by embracing second-sphere coordination to aqua ligands.

One of the most commercially successful processes for gold mining involves the cyanide protocol. A critical step in this protocol is to strip $\text{Au}(\text{CN})_2^-$ ions from the surface of carbon into an aqueous solution under harsh conditions,^{182–184} involving high temperatures (95–140 °C), high pressures (70–400 kbar), and concentrated solutions of cyanide and hydroxide. There is a demand to develop new technologies that allow the operation of this protocol to occur under much milder conditions. We have, at Northwestern University (NU), demonstrated¹⁸⁵ that second-sphere coordination between α -cyclodextrin (α -CD)

and $\text{KAu}(\text{CN})_2$ can be used to facilitate the separation. Thus, single-crystal X-ray analysis reveals that α -CD and $\text{KAu}(\text{CN})_2$ form a 1:1 adduct (Figures 11a and 11b) under slow evaporation conditions and a 2:1 adduct by slow diffusion of EtOH into an aqueous solution. In both cases, adduct formation in the solid state is sustained by multiple $[\text{C}-\text{H}\cdots\pi]$ and $[\text{C}-\text{H}\cdots\text{anion}]$ interactions between cyanide ligands in the first coordination sphere and α -CD in the second sphere. The K^+ counterions link the α -CD tori in the crystal by forming (Figure 11c) coordinative $[\text{K}^+\cdots\text{O}]$ bonds with glucose residues.

In aqueous solution, the existence of the 1:1 adduct between $\text{KAu}(\text{CN})_2$ and α -CD was demonstrated by ^1H NMR titration and isothermal titration calorimetry. The binding affinity was determined to be on the order of 10^4 M^{-1} . Adduct formation in an aqueous solution is driven by a favorable enthalpy change that overcomes a minor entropic penalty. This 1:1 adduct formation between $\text{KAu}(\text{CN})_2$ and α -CD in aqueous solution is highly selective. When β - or γ -CD are used, the binding affinities dropped to around 10^2 and 10^1 M^{-1} , respectively. Although $\text{KAg}(\text{CN})_2$ was also found to form a 1:1 adduct with α -CD in aqueous solution, the binding affinity is an order of magnitude lower than that of $\text{KAu}(\text{CN})_2$. This property of second-sphere coordination can be applied (Figure 11d) to the stripping of gold from the surface of activated carbon into aqueous solution at room temperature. Moreover, the stripping process is selective for $\text{Au}(\text{CN})_2^-$ in the presence of $\text{Ag}(\text{CN})_2^-$, making this new




Figure 9 | A cyclic process for the separation of Cu^{2+} from Co^{3+} based on second-sphere coordination. When one molar equivalent of 18C6 was added to a solution containing one molar equivalent each of $[\text{Co}(\text{NH}_3)_6]^{3+} [\text{PF}_6^-]_3$ and $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})]^{2+} [\text{PF}_6^-]_2$, the 1:1 copper adduct was precipitated exclusively in >90% yield. A simple filtration can separate the precipitated copper adducts from the cobalt-containing solution. The 18C6 ligand can be recovered by treating the 1:1 adduct with HCl solution, which yields aqueous copper(II) and insoluble $[\text{NH}_4]^+ [\text{PF}_6^-] \cdot 18\text{C6}$. The ammonium hexafluorophosphate adduct with 18C6 could be used again in separating $[\text{Cu}(\text{NH}_3)_4(\text{H}_2\text{O})]^{2+}$ from $[\text{Co}(\text{NH}_3)_6]^{3+}$. Adapted with permission from ref 25. Copyright 1986 John Wiley and Sons.

carbon-in-pulp process particularly attractive for mining gold ores with a high silver-content. In principle, molecular recognition processes could be integrated into commercial gold-mining protocols and lead to significantly reduced costs, energy consumption, and environmental impact.

We at NU have also developed^{186,187} an eco-friendly gold recovery protocol based on the serendipitous discovery of adduct formation between α -CD and KAuBr_4 . In an attempt to grow a CD-MOF^{188–191} using KAuX_4 ($\text{X} = \text{Cl}/\text{Br}$), we observed instantaneous formation of a brown precipitate (Figure 12a) on adding an aqueous solution of KAuBr_4 to a solution of α -CD in water. The X-ray superstructure of crystals from this precipitate revealed a 1:2 second-sphere adduct between KAuBr_4 and α -CD (superstructure at left in Figure 12b), forming a one-dimensional polymeric $\{[\text{K}(\text{OH}_2)_6][\text{AuBr}_4]\text{C}(\alpha\text{-CD})_2\}_n$ superstructure.¹⁸⁶ The first-sphere ligands H_2O and Br^- are encapsulated by the second-sphere ligand α -CD. The α -CDs are packed alternately in head-to-head and tail-to-tail fashion, leading to the formation of closely packed, one-dimensional channels. The K^+ is coordinated by six H_2O ligands, forming an octahedrally coordinated $[\text{K}(\text{OH}_2)_6]^+$ ion which is sandwiched between the secondary faces of two α -CDs. This type of encapsulation is unusual considering the fact that here a fully hydrated K^+ ion is buried inside a hydrophobic binding cavity. The hydrophobic AuBr_4^- ion with a square-planar coordination geometry is encapsulated between the primary faces of two α -CDs, where each primary face is occupied with a Br^- ligand. The hydrophobic effect, plus $[\text{C} \cdots \text{Br} \cdots \text{Au}]$ hydrogen bonds between AuBr_4^- and the 12 H-5 and H-6 atoms on the primary faces of α -CDs, provide the driving forces for second-sphere coordination of AuBr_4^- with α -CDs. It is also worth mentioning that, inside the one-dimensional channel formed by α -CD, the two first-sphere coordination complexes line up with alternate positive and negative charges, stabilizing the overall superstructures by electrostatic interactions.

The coprecipitation process between α -CD and KAuBr_4 is highly selective. No precipitate is obtained when α -CD is replaced with β - or γ -CD, which do not encapsulate the $[\text{K}(\text{OH}_2)_6]^+$ counterion complex. Although these larger cyclodextrins fail to induce formation of the low-solubility superstructure with KAuBr_4 , they do afford (Figure 12b) some more soluble complexes. Also, the use of KAuCl_4 does not induce coprecipitation with α , β , or γ -CD, though again, more soluble complexes can be isolated. The highly selective precipitation of KAuBr_4 with α -CD can even be achieved in the presence of other coordination complexes, including PtX_4^{2-} and PdX_4^{2-} ($\text{X} = \text{Cl}/\text{Br}$). The gold recovery efficiency is highly dependent on the counterions of AuBr_4^- , with the K^+ ion delivering by far the highest yield (~80%)




Figure 10 | Tubular and space-filling representation of 2:1 adducts between acyclic sulfonate ligand and (a) $[\text{Fe}(\text{OH}_2)_6]^{3+}$, (b) $[\text{Cr}(\text{OH}_2)_6]^{3+}$, and (c) $[\text{Al}(\text{OH}_2)_6]^{3+}$. Tubular and space-filling representation of 1:1 adducts between acyclic sulfonate ligand and (d) $[\text{Cu}(\text{OH}_2)_6]^{2+}$, (e) $[\text{Zn}(\text{OH}_2)_6]^{2+}$, and (f) $[\text{Mg}(\text{OH}_2)_6]^{2+}$. A competitive crystallization can be employed to separate Fe^{3+} from Zn^{2+} . Equimolar amounts of the acyclic sulfonate ligand, $\text{Fe}(\text{NO}_3)_3$, and $\text{Zn}(\text{NO}_3)_2$ were combined in an aqueous solution. Diffusion of methanol into this mixture afforded only yellow crystals of the 1:1 adduct between the ligand and Fe^{3+} aqua complex, leaving all the Zn^{2+} in solution.

of precipitated complex. Gold recovery decreases to 0%, 41%, and 61% for Na^+ , Rb^+ , and Cs^+ , respectively.¹⁸⁷ Hydrated counterions are not found in the crystal structures of these latter complexes, the alkali metal ions being instead coordinated by oxygen atoms of the cyclodextrin's glucose residues. Temperature also plays an important role in the recovery yield, with lower temperatures leading to decreased solubility of the adducts: the gold recovery yield reaches 94% when the coprecipitation is carried out at 0 °C.

A laboratory-scale gold-recovery process has been developed (Figure 13) based on this highly selective, simple, fast, and efficient coprecipitation method. Two scrap gold-bearing alloys containing 58 wt % of Au and 42 wt % of other metals (Zn, Cu, and Ag) were converted into HAuBr_4 by dissolving them in a mixture of the concentrated acids HBr and HNO_3 and then neutralizing the solutions with aqueous KOH. When α -CD was added to both solutions, coprecipitation occurred immediately in the presence of other metal ions, including Zn^{2+} and Cu^{2+} . The coprecipitated adduct—containing the recovered

gold—was separated by filtration and then treated with a reductant, specifically $\text{Na}_2\text{S}_2\text{O}_5$, to afford the recovered metal. The gold recovery from the two samples of gold scrap resulted in 89% yield with 97% purity and 92% yield with 95% purity.¹⁸⁶ For a larger, industrial-scale gold-recovery process, KAuBr_4 can be generated in situ using an etchant solution that is either composed of KBr, Br_2 , and H_2O (Figure 13a) or KBr, O_3 , and H_2O_2 (Figure 13b). This gold recovery process is much more environmentally benign in comparison with the universally accepted cyanide process. In view of the economic feasibility of the process, a startup company, Cycladex, was established in 2014. The Cycladex process has several advantages over the cyanide protocol, including faster leaching times, higher gold recovery efficiency, reduced environmental concerns, lower cost, and lower capital expenditure. The Tao group^{192,193} and our own NU-based team¹⁹⁴ discovered, simultaneously and independently, that cucurbit[n]urils ($\text{CB}[n]$, $n = 5-8$) are superior second-sphere ligands for AuX_4^- ($X = \text{Cl}/\text{Br}$) and lead¹⁹⁴ to instant coprecipitation (Figure 14a) of the adduct from aqueous solution. Instead




Figure 11 | (a) Space-filling and (b) tubular representations of the X-ray superstructure of an adduct between α -CD and the $\text{Au}(\text{CN})_2^-$ anion. (c) Stick and space-filling representation of the crystal packing between $\text{Au}(\text{CN})_2^-$ anions and α -CDs, showing the positions of K^+ cations and $\text{Au}(\text{CN})_2^-$ anions. (d) Graphical illustration of gold stripping from the surface of activated carbon into aqueous solution using α -CD at room-temperature. Adapted with permission from ref 185. Copyright 2021 American Chemical Society.

of encapsulating the AuX_4^- ions inside the binding cavity, cucurbiturils interact with AuX_4^- through their outer surfaces. The X-ray superstructure of an adduct between CB[6] and HAuCl_4 reveals that each CB[6] is associated (Figure 14b) with four AuCl_4^- ions through a combination of $[\text{Au}-\text{Cl}\cdots\text{C}=\text{O}]$ ion-dipole and $[\text{Au}-\text{Cl}\cdots\text{H}-\text{C}]$ hydrogen-bonding interactions between the Cl atoms in AuCl_4^- and the methine hydrogen atoms on the outer surface of CB[6]. The solid-state superstructure of the adduct between CB[6] and HAuBr_4 is quite different from that of the adduct between CB[6] and HAuCl_4 . The AuBr_4^- anions exhibit two types of binding with CB[6] molecules. One of the AuBr_4^- anion is connected with two CB[6] molecules located (Figure 14c) in adjacent layers, as a result of $[\text{Au}-\text{Br}\cdots\text{H}-\text{C}]$ interactions. The other AuBr_4^- anion interacts with five CB[6] molecules through $[\text{Au}-\text{Br}\cdots\text{H}-\text{C}]$ hydrogen-bonding and $[\text{Au}-\text{Br}\cdots\text{C}=\text{O}]$ interactions.

Gold recovery using CB[6] is highly efficient: the yield is between 93% and 96% depending on the halide ligands of AuX_4^- and its counter cations. Additionally, a laboratory-scale gold-recovery protocol has been developed using a

yellow gold-bearing alloy wire, which contains 58 wt % of Au and 42 wt % of Cu, Zn, and Ag. The gold portion of the alloy wire was converted to a solution of HAuCl_4 by dissolving it in a mixture of HCl and H_2O_2 . When an aqueous solution of CB[6] is added to this gold-containing solution, coprecipitation of the adduct between CB[6] and HAuCl_4 occurs immediately. Elemental analysis revealed that 99.8% of the gold present in the raw material was recovered, and its purity is 98.1%. The CB[6] can be recovered by recrystallization, providing yet another opportunity to develop more efficient processes for gold recovery.

Gong et al.¹⁹⁵ have reported a tetracationic imidazolium cyclophane, which acts (Figure 15a) as a receptor to coprecipitate square-planar coordination complexes of noble metals— AuCl_4^- , PtCl_4^{2-} , and PdCl_4^{2-} —from aqueous solution. Single-crystal X-ray superstructures (Figures 15b–15d) of the adducts formed between this receptor and AuCl_4^- , PtCl_4^{2-} , and PdCl_4^{2-} were obtained. Although the formation of these adducts involves $[\text{C}-\text{H}\cdots\text{Cl}-\text{M}]$ hydrogen-bonding, anion- π , and electrostatic interactions, the coordination complexes do not have strong directional preferences or shape selectivity for the structurally flexible cyclophane. The coprecipitation is thus driven primarily by the formation of hydrophobic ion pairs in aqueous solution.

Smith et al.¹⁹⁶ have developed a precise molecular recognition process (Figure 16) for the square planar ions $[\text{AuX}_4^-]$ ($\text{X} = \text{Cl}/\text{Br}$) using a durene-based tetralactam macrocycle. The cavity of this receptor is preorganized (Figures 16a and 16b) with four convergent hydrogen-bond donors to interact with the electronegative halide ligands. Meanwhile, the receptor possesses a pair of parallel arene units (durene residues) as π -electron donors to interact with the electropositive metal center. Several single-crystal X-ray superstructures were obtained (Figures 16c and 16d) of the adducts between this receptor and coordination complexes of precious metals, including AuX_4^- ($\text{X} = \text{Cl}/\text{Br}$), PtCl_4^{2-} , and $\text{Pd}_2\text{Cl}_6^{2-}$, which confirmed the presence of the hydrogen bonds and $[\text{metal}\cdots\pi]$ interactions. These second-sphere coordination adducts were also shown to exist in solution, with ^1H NMR titrations indicating that their stability in solution depends on the nature of both the halide ligands and the metals. The binding affinity of AuCl_4^- ($K_a = 7.8 \times 10^4 \text{ M}^{-1}$) is about 30 times stronger than that of AuBr_4^- ($K_a = 2.5 \times 10^3 \text{ M}^{-1}$). When the metal center is changed from Au(III) to Pt(II), the binding affinity ($\text{X} = \text{Cl}$) was decreased by a factor of 3900 to $K_a = 20 \text{ M}^{-1}$, highlighting the critical role of $[\text{metal}\cdots\pi]$ interactions. Addition of the receptor to a solution of HAuCl_4 or HAuBr_4 in an organic solvent led to instant coprecipitation of the adducts. In subsequent work, Jiang et al.¹⁹⁷ have developed a similar 2,3-dibutoxynaphthalene-based tetralactam macrocycle that can capture AuCl_4^- from organic solution. Molecular modeling suggests that




Figure 12 | (a) Selective formation of a coprecipitate $\alpha\text{-CD}\cdot\text{KAuBr}_4$. The brown precipitate formed instantly when adding an aqueous solution of KAuBr_4 into a solution of $\alpha\text{-CD}$ in H_2O . (b) Stick and space-filling representation of the X-ray superstructures of $\alpha\text{-CD}\cdot\text{KAuBr}_4$, $\beta\text{-CD}\cdot\text{KAuBr}_4$, $\gamma\text{-CD}\cdot\text{KAuBr}_4$, $\alpha\text{-CD}\cdot\text{KAuCl}_4$, $\beta\text{-CD}\cdot\text{KAuCl}_4$, and $\gamma\text{-CD}\cdot\text{KAuCl}_4$. The major reason for the adduct co-precipitation between $\alpha\text{-CD}$ and KAuBr_4 is that they form a one-dimensional polymeric $\{[\text{K}(\text{OH})_2]_6[\text{AuBr}_4]_2(\alpha\text{-CD})_2\}_n$ superstructure. Adapted with permission from ref 186. Copyright 2013 Springer Nature.




Figure 13 | Flow diagrams of gold recovery based on the formation of the insoluble 1:1 adduct between α -CD and $[K(OH)_2]_6^+[AuBr_4]^-$. $KAuBr_4$ can be generated in situ using an etchant solution that is either composed of (a) KBr, Br₂, and H₂O or a greener oxidant (b) KBr, O₃, and H₂O₂. Red arrows indicate the flow direction of the gold recovery. Na₂S₂O₅ = Sodium metabisulfite.

adduct formation is sustained by the same patterns of noncovalent bonding interactions.

In an effort to direct the molecular recognition process toward a practical gold-recovery technology, Smith et al.¹⁹⁸ have developed (Figures 16e and 16f) an acyclic system that can be made to work in a similar manner to their tetralactam macrocycle¹⁹⁶ for the coprecipitation of gold halides. The acyclic amides can be rendered soluble in either organic or aqueous solvents, depending on the constitution of the solubilizing R groups. These acyclic receptors and the gold(III) halide complexes coprecipitate instantly from solution. X-ray diffraction analysis of the recrystallized coprecipitates reveals (Figures 16g and 16h) a similar pattern of noncovalent-bonding interactions to the binding of the gold halides by the tetralactam macrocycle involving an array of hydrogen bonds in addition to $[Au\cdots\pi]$ interactions. These acyclic receptors with low molecular weights can be prepared cheaply on a large scale, rendering them attractive molecules for the development of next-generation gold-recovery technologies.

Catalysis

Natural enzymes perform catalytic reactions at lightning speeds and with high substrate selectivities using a potent catalytic center within a nanoconfined pocket^{91,199} constructed by chain-folding of a protein molecule. The superstructure of an enzyme-binding pocket not only dictates the substrate selectivity but also isolates the catalytic center from other reactive sites. This compartmentalization of different catalytic centers enhances²⁰⁰ the stability of a given catalytic center and extends its functional lifetime. Following pioneering research by

Breslow et al.^{201–208} in the 1970s using cyclodextrins and their derivatives, there have been continued efforts to develop enzyme analogues that allow catalytic reactions to be performed in confined nanoenvironments.^{91,199} Significant advances have been made in creating these artificial enzymes, as summarized^{209–219} in several excellent reviews. It turns out that second-sphere coordination has been exploited widely as an emerging strategy for designing artificial enzymes by encapsulating catalysts inside molecular containers, leading to high efficiencies and excellent regio- and stereoselectivities being achieved at the catalytic centers.

Sollogoub et al.²²⁰ have demonstrated a good example of regioselectivity that involves switching using covalent *N*-heterocyclic carbene-capped α - and β -cyclodextrins as second-sphere coordination ligands, encapsulating a Cu(I)Cl catalyst. The resulting catalysts, α - and β -CD-CuCl, were investigated (Figure 17a) for copper-catalyzed hydroboration of substituted and unsubstituted terminal and internal phenylalkynes. It was found that α -CD-CuCl results in excellent selectivity for the linear products (Figure 17b), yielding a branched-to-linear (B/L) ratio of 9:91. By contrast, the β -CD-CuCl catalyst leads mainly to branched products, with a B/L ratio of 90:10. The changed shape of the cavity promotes, not only a regioselectivity switch, but also a mechanistic switch. Density functional theory (DFT) calculations indicate that the cavity of α -CD-CuCl promotes an orthogonal approach of acetylene, leading to a linear product. The larger cavity of β -CD-CuCl favors instead a parallel approach, in which the alkyne projects directly into the cavity.

In a subsequent investigation, the same authors reported²²¹ another example (Figure 17c) of




Figure 14 | (a) Rapid formation of coprecipitates $\text{CB}[6] \cdot \text{MAuX}_4$ ($M = \text{H/K}$, $X = \text{Cl/Br}$) from a mixture of $\text{CB}[6]$ and MAuX_4 in aqueous solution. (b) Ball-and-stick representation of X-ray superstructure of the adduct between $\text{CB}[6]$ and AuCl_4^- . Each $\text{CB}[6]$ is associated with four AuCl_4^- anions through a combination of $[\text{Au}-\text{Cl} \cdots \text{C}=\text{O}]$ ion-dipole interactions and $[\text{Au}-\text{Cl} \cdots \text{H}-\text{C}]$ hydrogen-bonding interactions between the Cl atoms in AuCl_4^- and the methine hydrogen atoms on the outer surface of $\text{CB}[6]$. (c) Ball-and-stick representation of X-ray superstructure of the adduct between $\text{CB}[6]$ and AuBr_4^- . The AuBr_4^- anions exhibit two types of binding with $\text{CB}[6]$ molecules. One of the AuBr_4^- anion is associated with two $\text{CB}[6]$ molecules located in adjacent layers as a result of $[\text{Au}-\text{Br} \cdots \text{H}-\text{C}]$ interactions. The other AuBr_4^- anion interacts with five $\text{CB}[6]$ molecules through the $[\text{Au}-\text{Br} \cdots \text{H}-\text{C}]$ hydrogen-bonding and $[\text{Au}-\text{Br} \cdots \text{C}=\text{O}]$ interactions. Adapted with permission from ref 194. Copyright 2020 American Chemical Society.




Figure 15 | (a) Rapid formation of coprecipitates cyclophane• MX_4^{n-} ($M = Au/Pt/Pd$, $n = 1$ or 2) in aqueous solution. Ball-and-stick representations of X-ray superstructures between cyclophane and (b) $AuCl_4^-$, (c) $PtCl_4^{2-}$, and (d) $PdCl_4^{2-}$. These noble-metal coordination complexes do not have strong directional preference or shape selectivity for the structurally flexible cyclophane. The coprecipitation is driven primarily by the formation of hydrophobic ion pairs in an aqueous solution. Adapted with permission from ref 195. Copyright 2018 Elsevier.

regioselective catalysis using these two cyclodextrin-based catalysts for the hydrosilylation of α,β -unsaturated enones. The very reactive monomeric hydride (L)CuH is stabilized inside the cavity, leading to the high efficiencies exhibited by these two catalysts. Remarkably, exclusive 1,2-addition is observed using α -CD-CuCl, with good enantiomeric excesses. In contrast, the β -CD-CuCl catalyst, with its larger binding cavity, promotes the full reduction of the substrate. The difference in selectivity for these two catalysts can be attributed to the difference in the sizes of the binding cavities, which control the accessibility of the substrate to the catalytic Cu(I) center. Additionally, the interactions between CDs and their substrates favor 1,2-addition for α -CD-CuCl and 1,4-addition for β -CD-CuCl. This example demonstrates the ability of the two cyclodextrin-based ligands to stabilize the reactive catalytic center and select for different orientations of the substrate, resulting in highly effective catalysis with excellent regioselectivities.

The Raymond, Bergman, and Toste triumvirate^{211,216,222–224} at UC Berkeley have investigated numerous catalysts within confined nanospaces by exploiting a family of polyanionic tetrahedral cages that function as second-

sphere ligands for the encapsulation of catalysts composed of coordination complexes. For example, they have reported (Figure 18a) a rare example of site-selective hydrogenation of alkenes, promoted²²³ by a catalyst composed of a naphthalene-based gallium(III) cage ("Naph-Cage") encapsulating a cationic (DMPE)Rh(COD) precursor (DMPE = 1,2-bis(dimethylphosphino)ethane; COD = 1,5-cyclooctadiene). Using this cage-encapsulated catalyst, the authors observed high conversions of alkenes where the double bond is remote from the hydroxyl group, as in 5- and 4-hexen-1-ols. Conversely, little or no conversion of the double bonds more proximate to the hydroxy groups presented in 3- and 2-hexen-1-ols was observed under the same conditions. Moreover, the free Rh catalyst operating in bulk solution led to a quantitative conversion of all hexen-1-ol substrates, regardless of the position of the C=C double bond. The site-selectivity of this cage-encapsulated catalyst has been attributed to preferential binding of the more lipophilic alkyl end of the pendant alcohol substrate within the cavity. In a further demonstration of site-selective hydrogenation, the authors showed²²³ that the fatty alcohol of lineolenic acid (Figure 18b), which contains three points of unsaturation, undergoes only




Figure 16 | (a) Schematic illustration of binding AuX_4^- with cyclophane through two sets of directional interactions, that is, hydrogen bonding and Au-aromatic stacking. (b) The structural formula of a durene-based tetralactam macrocycle. Ball-and-stick representations of the X-ray solid-state superstructure of the adducts of macrocycle with (c) AuCl_4^- and (d) AuBr_4^- . (e) Schematic illustration of binding AuX_4^- with acyclic ligands through two sets of directional interactions (hydrogen bonding and Au-aromatic stacking). (f) The structural formula of a durene-based acyclic ligand. Ball-and-stick representations of the X-ray solid-state superstructure of the adducts of a durene-based acyclic ligand with (g) AuCl_4^- and (h) AuBr_4^- . Adapted with permission from ref 196. Copyright 2018 American Chemical Society.

monohydrogenation (74% yield) of the terminal double bond in the presence of a larger pyrene-based cage (“Pyr-Cage”). This example shows that supramolecular catalyst-directed regioselectivity, based on second-sphere coordination, can overcome any inherent selectivity deriving from the substrate or a nonencapsulated catalyst.

Second-sphere coordination of transition-metal catalysts has also been shown to enhance enantioselective transformations. Reek et al.²²⁵ have reported the encapsulation of a chiral Rh complex in an achiral supramolecular cage (Figure 19a) formed by coordination-driven self-assembly of a macrocyclic dipalladium complex and a tetracarboxylate zinc porphyrin. The resulting encapsulated catalyst is highly active in the case of the hydroformylation of styrene and derivatives (Figure 19b), which were converted into aldehydes with a much higher chiral induction—up to 71% e.e.—than with the free Rh catalyst. The stereoselectivity demonstrated by the encapsulated catalyst is based on controlling the second-sphere coordination by way of the structural

constraints imposed by the cage. The same group has also described several examples in which second-sphere coordination is able to control regioselectivity in hydroformylation reactions.^{226–230}

MOFs can be synthesized with a wide range of nodes and linkers,^{2–16} making them an attractive platform for systematic engineering of second-sphere interactions. Inspired by the effectiveness of biological catalytic conversion of carbon dioxide to the glucose precursor glyceraldehyde-3-phosphate, performed in compartmentalized organelles, Tsung et al.²³¹ have investigated (Figure 20) the influence of second-sphere coordination on catalytic reactivity in the hydrogenation of carbon dioxide to methanol, using MOF UiO-66 as the second-sphere ligand to encapsulate a transition-metal catalyst. This catalytic system is composed of three components: (1) a ruthenium PNP pincer complex (green) to carry out the hydrogenation of carbon dioxide to formic acid, (2) the $\text{Zr}_6(\text{OH})_4\text{O}_4$ metal nodes of the MOF to convert formic acid to formate ester, and (3) a second ruthenium




Figure 17 | (a) Covalent *N*-heterocyclic carbene-capped α - and β -CDs as second-sphere coordination ligands, which encapsulate a Cu(I)Cl catalyst. (b) Copper-catalyzed hydroboration with substituted and unsubstituted terminal and internal phenylalkynes using α - and β -CD-CuCl. The use of α -CD-CuCl results in excellent selectivity for the linear products, yielding a B/L ratio of 9:91. In contrast, the β -CD-CuCl catalyst leads to the branched products with a B/L ratio of 90:10. (c) Hydrosilylation of α,β -unsaturated enones catalyzed by α - and β -CD-CuCl. Exclusive 1,2-addition using α -CD-CuCl with good enantiomeric excesses was observed. The β -CD-CuCl catalyst was found to promote the full reduction of the substrate. Adapted with permission from ref 220. Copyright 2017 John Wiley and Sons.

PNN pincer complex (orange) to catalyze hydrogenation of the formate ester to methanol. The final product, methanol, was only observed in these cascade hydrogenation reactions when the two ruthenium complexes were separated from each other by encapsulating the PNP catalyst into the cavity of the MOF, illustrating the importance of component separation for efficient catalysis. The beneficial effects of the second-sphere coordination were investigated extensively for the first catalyst in this cascade process. The synthetic flexibility of the MOF UiO-66 allowed the authors to modify the linkers with different substituents, leading to a series of

functionalized MOF UiO-66. They found that when a primary ammonium functional group, [Ar-NH₃]⁺, is employed, the resulting UiO-66-NH₃⁺ significantly improves the catalytic efficiency for hydrogenation of carbon dioxide to formic acid. Mechanistic investigations revealed that the arylammonium group functions as a general Brønsted acid. Consequently, the resulting catalyst system provides access to high catalytic performance by manipulating second-sphere interactions. The authors obtained the highest reported turnover number (TON) (19,000) and turnover frequency (TOF) (9100 h⁻¹) for hydrogenation of carbon dioxide to




Figure 18 | (a) The structural formula of a Naph-Cage, which can encapsulate the Rh catalyst, (DMPE)Rh(COD), for the site-selective hydrogenation of olefins. High conversions of olefins in which the double bond is remote from the hydroxyl group (5- and 4-hexen-1-ol) are seen. Meanwhile, little or no conversion of more proximate double bonds (3- and 2-hexen-1-ol) was observed under the same catalytic conditions. (b) The structural formula of a Pyr-Cage which encapsulates a Rh catalyst, catalyzing monohydrogenation of the fatty alcohol of lineolenic acid that contains three points of unsaturation. Adapted with permission from ref 223. Copyright 2019 American Chemical Society.

methanol. Furthermore, the recyclable nature of this system allowed the authors to achieve a cumulative TON of 100,000 after 10 reaction cycles.

Biomolecular Chemistry

Biological macromolecules have long been known to bind to transition-metal complexes, most notably, perhaps, the metal-based anticancer drugs cisplatin,^{25,52} [cis-(Pt(NH₃)₂(Cl)₂], and carboplatin^{53,54} that complex with DNA and inhibit its replication and transcription. These drugs react with the guanine bases of DNA in the major groove, initially via aquation, with loss of halide or carboxylate ligands, followed by direct (first-sphere) coordination of the metal center by guanine nitrogen atoms that displace the coordinated water. Over the past decade, however, it has also been established that kinetically *inert* complexes of ruthenium(II) with heterocyclic ligands such as 2,2'-bipyridine (bipy), phenanthroline (phen), and dipyridophenazine (dppz) are able to bind to double-helical DNA by a *second-sphere* mechanism (Figure 21), involving intercalation of a heterocyclic ligand between successive pairs of nucleobases, without a great deal of base specificity.²³²

DOI: 10.31635/ccschem.021.202101286
Corrected Citation: CCS Chem. 2022, 4, 755–784
Previous Citation: CCS Chem. 2021, 3, 3436–3465
Link to VoR: <https://doi.org/10.31635/ccschem.021.202101286>




Figure 19 | (a) Ball-and-stick representation of a chiral Rh complex in an achiral supramolecular cage formed from coordination-driven self-assembly of macrocyclic dipalladium complex and tetracarboxylate zinc porphyrins. (b) Hydroformylation of styrene and derivatives, which were converted into formyl products with a very high chiral induction of up to 71% e.e. catalyzed by the encapsulated Rh catalyst, significantly outperforming the free Rh catalyst.




Figure 20 | A host-guest multicomponent catalyst system for the hydrogenation of carbon dioxide to MeOH engineered by second-sphere coordination. In this system, the green ruthenium complex 1@UiO-66-NH₃⁺ carries out the hydrogenation of CO₂ to HCO₂H, the blue Zr₆(OH)₄O₄ nodes of the MOF converts formic acid to formate ester, and the orange 2@UiO-66 carries out the hydrogenation of formate ester to MeOH. Adapted with permission from ref 231. Copyright 2021 American Chemical Society.




Figure 21 | Representative DNA-binding complexes $[Ru(LL)_3]^{2+}$, with illustrative modes of intercalation into DNA between nucleobase pairs. $L = Phen, bpy$.

Single-crystal X-ray studies of such systems, using double-stranded DNA oligomers (typically decamers or dodecamers), have revealed a wide range of binding geometries (Figure 22), depending to some extent on the base pairs involved but mainly on the detailed structure and chirality of the metal complex.^{233–235} Given the opposite charges of the DNA host (strongly anionic) and the metal complex guest (di-cationic), together with the relatively low polarity of the heterocyclic ligands and the coplanarity of the base pairs, it seems likely that electrostatic forces, hydrophobic effects, and $[\pi \cdots \pi]$ stacking interactions all play significant roles in the second-sphere binding of these complexes to DNA.

Ruthenium complexes of the type shown in Figure 21 are known sensitizers for photooxidation of DNA. This observation can be turned to advantage in photodynamic cancer therapy where only targeted tissue is destroyed. Recent studies of single crystals of a DNA oligomer bound to such a photosensitizing ruthenium complex, using both X-ray crystallography and ultra-fast (picosecond) infrared (IR) laser spectroscopy, have shown²³⁶ for the first time how light absorption by the metal complex and consequent electron transfer from the readily oxidized guanine base are related to DNA damage. Such damage represents the first step in killing a tumor cell. Guanine residues are also important in the specialized drug target known as the G-quadruplex, which has a large planar aromatic surface and which correspondingly has a strong affinity for large aromatic ligands. This type of unit is particularly easy to oxidize and so there is much scope for donor-acceptor matching in the design of new therapeutic agents.²³⁶




Figure 22 | Summary of the main binding modes of the two enantiomers of $[Ru(L)_2dppz]^{2+}$ with duplex DNA ($L = phenanthroline$ or tetraazaphenanthroline, TAP). Λ form is shown in blue and Δ form in green. The specific binding mode adopted is dependent on the DNA sequences involved, so that the symmetrical intercalation mode, for example, has so far been found only at the duplex sequence TA/TA. This sequence specificity arises from π - π stacking of phenanthroline with the adenosine (A) residues.

Summary and Outlook

Second-sphere coordination has emerged as a strategy to access many functional systems in molecular recognition, construction of MIMs, metal separations, catalysis, and biomolecular chemistry. The noncovalent bonding interactions between the second-sphere ligands and coordination complexes can fine-tune the chemical, physical, electrochemical, and photochemical properties of the resulting adducts, leading to numerous opportunities in creating new technologies.

As one of the future directions for this chemistry, second-sphere coordination can be employed to develop new protocols for the sustainable separation of metals. Currently, most of the effort has been focused on new strategies for gold separation by second-sphere coordination, but similar principles can undoubtedly be applied

to the separation of other metals. Second-sphere coordination can facilitate the crystallization, precipitation, leaching, and chromatographic separation in metal-purification processes, leading to selective, efficient, and green separation technologies for metals.

In the solid state, MOFs have provided a versatile platform for the operation of AMMs by harnessing the concept of second-sphere coordination. Some elegant examples of mechanically interlocked MOFs with machine-like dynamics have been demonstrated.^{237–241} The operation of these molecular machines in the solid state may well provide an important route for the realization of advanced technologies in gas storage, drug delivery, and molecular separations.

Looking to the longer term, the marriage between supramolecular and coordination chemistry can clearly generate new pathways for tackling societal challenges in health, energy, and the environment. We envision that this field will attract more and more attention from multidisciplinary researchers, who will unleash the potential of this concept to advance chemistry, medicine, materials, and environmental science.

Conflicts of Interest

The authors declare no conflict of interest.

Acknowledgments

The authors thank Dr. Margaret Schott for assistance with editing and proofing of the manuscript. The authors thank Northwestern University (NU) for their support of this work, which was also funded by the Center for Sustainable Separation of Metals (CSSM) and part of a National Science Foundation (NSF) Center for Chemical Innovation (CCI): grant number CHE1925708.

References

1. Lawrance, G. A. *Introduction to Coordination Chemistry*; Wiley: Chichester, **2013**.
2. Yaghi, O. M.; Li, G.; Li, H. Selective Binding and Removal of Guests in a Microporous Metal–Organic Framework. *Nature* **1995**, *378*, 703–706.
3. Rosi, N. L.; Eckert, J.; Eddaoudi, M.; Vodak, D. T.; Kim, J.; O’Keeffe, M.; Yaghi, O. M. Hydrogen Storage in Microporous Metal–Organic Frameworks. *Science* **2003**, *300*, 1127–1129.
4. Li, Q.; Zhang, W.; Miljanić, O. Š.; Sue, C.-H.; Zhao, Y.-L.; Liu, L.; Knobler, C. B.; Stoddart, J. F.; Yaghi, O. M. Docking in Metal–Organic Frameworks. *Science* **2009**, *325*, 855–859.
5. Kreno, L. E.; Leong, K.; Farha, O. K.; Allendorf, M.; Van Duyne, R. P.; Hupp, J. T. Metal–Organic Framework Materials as Chemical Sensors. *Chem. Rev.* **2012**, *112*, 1105–1125.
6. Furukawa, H.; Cordova, K. E.; O’Keeffe, M.; Yaghi, O. M. The Chemistry and Applications of Metal–Organic Frameworks. *Science* **2013**, *341*, 1230–1234.
7. Bobbitt, N. S.; Mendonca, M. L.; Howarth, A. J.; Islamoglu, T.; Hupp, J. T.; Farha, O. K.; Snurr, R. Q. Metal–Organic Frameworks for the Removal of Toxic Industrial Chemicals and Chemical Warfare Agents. *Chem. Soc. Rev.* **2017**, *46*, 3357–3385.
8. Kim, H.; Yang, S.; Rao, S. R.; Narayanan, S.; Kapustin, E. A.; Furukawa, H.; Umans, A. S.; Yaghi, O. M.; Wang, E. N. Water Harvesting from Air with Metal–Organic Frameworks Powered by Natural Sunlight. *Science* **2017**, *356*, 430–434.
9. Lazaro, I. A.; Forgan, R. S. Image-Guided Therapy Using Maghemite–MOF Nanovectors. *Chem* **2017**, *3*, 200–202.
10. Kung, C.-W.; Goswami, S.; Hod, I.; Wang, T. C.; Duan, J.; Farha, O. K.; Hupp, J. T. Charge Transport in Zirconium–Based Metal–Organic Frameworks. *Acc. Chem. Res.* **2020**, *53*, 1187–1195.
11. Goetjen, T. A.; Liu, J.; Wu, Y.; Sui, J.; Zhang, X.; Hupp, J. T.; Farha, O. K. Metal–Organic Framework (MOF) Materials as Polymerization Catalysts: A Review and Recent Advances. *Chem. Commun.* **2020**, *56*, 10409–10418.
12. Ji, Z.; Li, T.; Yaghi, O. M. Sequencing of Metals in Multivariate Metal–Organic Frameworks. *Science* **2020**, *369*, 674–680.
13. Chen, Z.; Li, P.; Anderson, R.; Wang, X.; Zhang, X.; Robinson, L.; Redfern, L. R.; Moribe, S.; Islamoglu, T.; Gómez-Gualdrón, D. A.; Yildirim, T.; Stoddart, J. F.; Farha, O. K. Balancing Volumetric and Gravimetric Uptake in Highly Porous Materials for Clean Energy. *Science* **2020**, *368*, 297–303.
14. Feng, L.; Wang, K. Y.; Willman, J.; Zhou, H. C. Hierarchy in Metal–Organic Frameworks. *ACS Cent. Sci.* **2020**, *6*, 359–367.
15. Islamoglu, T.; Chen, Z.; Wasson, M. C.; Buru, C. T.; Kirlikovali, K. O.; Afrin, U.; Mian, M. R.; Farha, O. K. Metal–Organic Frameworks against Toxic Chemicals. *Chem. Rev.* **2020**, *120*, 8130–8160.
16. Lin, R. B.; Xiang, S.; Zhou, W.; Chen, B. Microporous Metal–Organic Framework Materials for Gas Separation. *Chem* **2020**, *6*, 337–363.
17. Parker, D.; Suturina, E. A.; Kuprov, I.; Chilton, N. F. How the Ligand Field in Lanthanide Coordination Complexes Determines Magnetic Susceptibility Anisotropy, Paramagnetic NMR Shift, and Relaxation Behavior. *Acc. Chem. Res.* **2020**, *53*, 1520–1534.
18. Ramdass, A.; Sathish, V.; Babu, E.; Velayudham, M.; Thanasekaran, P.; Rajagopal, S. Recent Developments on Optical and Electrochemical Sensing of Copper(II) Ion Based on Transition Metal Complexes. *Coord. Chem. Rev.* **2017**, *343*, 278–307.
19. Jiao, K.-J.; Xing, Y.-K.; Yang, Q.-L.; Qiu, H.; Mei, T.-S. Site-Selective C–H Functionalization via Synergistic Use of Electrochemistry and Transition Metal Catalysis. *Acc. Chem. Res.* **2020**, *53*, 300–310.
20. Velazquez, H. D.; Verpoort, F. N-Heterocyclic Carbene Transition Metal Complexes for Catalysis in Aqueous Media. *Chem. Soc. Rev.* **2012**, *41*, 7032–7060.

21. Arrowsmith, R. L.; Pascu, S. I.; Smugowski, H. New Developments in the Biomedical Chemistry of Metal Complexes: From Small Molecules to Nanotheranostic Design. *Organomet. Chem.* **2012**, *38*, 1–35.
22. Ma, D.-L.; He, H.-Z.; Leung, K.-H.; Chan, D. S.-H.; Leung, C.-H. Bioactive Luminescent Transition-Metal Complexes for Biomedical Applications. *Angew. Chem. Int. Ed.* **2013**, *52*, 7666–7682.
23. Ahmedova, A. Biomedical Applications of Metallosupramolecular Assemblies—Structural Aspects of the Anticancer Activity. *Front. Chem.* **2018**, *6*, 620.
24. Ko, C.-N.; Li, G.; Leung, C.-H.; Ma, D.-L. Dual Function Luminescent Transition Metal Complexes for Cancer Theranostics: The Combination of Diagnosis and Therapy. *Coord. Chem. Rev.* **2019**, *381*, 79–103.
25. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J. Second-Sphere Coordination—A Novel Role for Molecular Receptors. *Angew. Chem. Int. Ed. Engl.* **1986**, *25*, 487–507.
26. Raymo, F. M.; Stoddart, J. F. Second-Sphere Coordination. *Chem. Ber.* **1996**, *129*, 981–990.
27. Stoddart, J. F.; Zarzycki, R. Cyclodextrins as Second-Sphere Ligands for Transition Metal Complexes. *Recl. Trav. Chim. Pays-Bas* **1988**, *107*, 515–528.
28. Liu, Z.; Schneebeli, S. T.; Stoddart, J. F. Second-Sphere Coordination Revisited. *Chimia* **2014**, *68*, 315–320.
29. Werner, A. Über Die Raumisomeren Kobaltverbindungen. *Justus Liebigs Ann. Chem.* **1912**, *386*, 1–272.
30. Werner, A. *Neuere Anschauungen Auf Dem Gebiete Der Anorganischen Chemie*; Vieweg: Braunschweig, **1913**.
31. Colquhoun, H. M.; Stoddart, J. F. Second-Sphere Coordination of Neutral and Cationic Transition Metal Complexes by Crown Ethers. *J. Chem. Soc. Chem. Commun.* **1981**, 612–613.
32. Pedersen, C. J. Cyclic Polyethers and Their Complexes with Metal Salts. *J. Am. Chem. Soc.* **1967**, *89*, 2495–2496.
33. Pedersen, C. J. Cyclic Polyethers and Their Complexes with Metal Salts. *J. Am. Chem. Soc.* **1967**, *89*, 7017–7036.
34. Pedersen, C. J. The Discovery of Crown Ethers (Nobel Lecture). *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 1021–1027.
35. Colquhoun, H. M.; Lewis, D. F.; Stoddart, J. F.; Williams, D. J. Crown Ethers as Second-Sphere Ligands. The Interactions of Transition-Metal Amines with 18-Crown-6 and Dibenzo-18-Crown-6. *J. Chem. Soc. Dalt. Trans.* **1983**, 607.
36. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J. Formation and X-Ray Crystal Structure of $[\text{Pt}(\text{H}_2\text{NCH}_2\text{CH}_2\text{NH}_2)_2 \cdot 18\text{-Crown-6}]^{2+}[\text{PF}_6]_2^{2-}$. A Hydrogen Bonded Stepped-Chain Copolymer. *J. Chem. Soc. Chem. Commun.* **1981**, 851–852.
37. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J. The Binding of Neutral Platinum Complexes by Crown Ethers. X-Ray Crystal Structures of $[\text{trans-PtCl}_2(\text{PMe}_3)\text{NH}_3 \cdot \text{Dibenzo-18-Crown-6}]$ and $[\{\text{trans-PtCl}_2(\text{PMe}_3)\text{NH}_3\}_2 \cdot 18\text{-Crown-6}]$. *J. Chem. Soc. Chem. Commun.* **1981**, 847–849.
38. Cotton, S. A.; Franckevicius, V.; Mahon, M. F.; Ooi, L. L.; Raithby, P. R.; Teat, S. J. Structures of 2,4,6-Tri- α -Pyridyl-1,3,5-Triazine Complexes of the Lanthanoid Nitrates: A Study in the Lanthanoid Contraction. *Polyhedron* **2006**, *25*, 1057–1068.
39. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J.; Wolstenholme, J. B.; Zarzycki, R. Second Sphere Coordination of Cationic Platinum Complexes by Crown Ethers—The X-Ray Crystal Structure of $[\text{Pt}(\text{Bpy})(\text{NH}_3)_2 \cdot \text{Dibenzo}[30]\text{Crown-10}]^{2+}[\text{PF}_6]_2 \cdot x\text{H}_2\text{O}$ ($x \approx 0.6$). *Angew. Chem. Int. Ed. Engl.* **1981**, *20*, 1051–1053.
40. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J. Isolation and X-Ray Crystal Structure of $[\text{Cu}(\text{NH}_3)_4\text{H}_2\text{O} \cdot 18\text{-Crown-6}]_n^{2+}[\text{PF}_6]_{2n}^{2-}$ a Linear Face-to-Face Hydrogen Bonded Chain Copolymer. *J. Chem. Soc. Chem. Commun.* **1981**, 849–850.
41. Colquhoun, H. M.; Stoddart, J. F.; Williams, D. J. Correlation of Ligating Acetonitrile by 18-Crown-6. X-Ray Crystal Structure of $\{\text{trans-Ir}(\text{CO})(\text{CH}_3\text{CN})(\text{PPh}_3)_2\}^{2+} \cdot 18\text{-Crown-6}\}[\text{PF}_6]^{2-} \cdot 2\text{CH}_2\text{Cl}_2$. *J. Am. Chem. Soc.* **1982**, *104*, 1426–1428.
42. Colquhoun, H. M.; Doughty, S. M.; Stoddart, J. F.; Williams, D. J. Second Sphere Coordination of Cationic Rhodium Complexes by Dibenzo[3n]Crown-n Ethers. *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 235–236.
43. Allwood, B. L.; Crosby, J.; Pears, D. A.; Stoddart, J. F.; Williams, D. J. Crown Ether Complexes of Sulfonium Salts—The X-Ray Crystal Structures of $[\text{PhCOCH}_2\text{SMe}_2 \cdot 18\text{-Crown-6}]_n[\text{PF}_6]_n$ and $[(\text{PhCOCHPhSMe}_2)_2 \cdot 18\text{-Crown-6}][\text{PF}_6]_2$. *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 977–979.
44. Allwood, B. L.; Colquhoun, H. M.; Crosby, J.; Pears, D. A.; Stoddart, J. F.; Williams, D. J. Crown Ether Complexes of Phosphonium Salts—The X-Ray Structure Analysis of $[(\text{Ph}_3\text{PMe})_2 \cdot 18\text{-Crown-6}][\text{PF}_6]_2$. *Angew. Chem. Int. Ed. Engl.* **1984**, *23*, 824–825.
45. Colquhoun, H. M.; Doughty, S. M.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J. Diamminebis(1,5-cyclooctadiene)(μ -1,4,10,13-tetraoxa-7,16-diazacyclooctadecane-N7, N16)dirhodiumbis-(hexafluorophosphate): An Example of Simultaneous First and Second Sphere Coordination. *Angew. Chem. Int. Ed. Engl.* **1985**, *24*, 135–136.
46. Alston, D. R.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J.; Zarzycki, R. Second Sphere Coordination of Tetraammineplatinum(II) by a Macropolycyclic Crown Ether Bisamide Receptor. *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 692–693.
47. Alston, D. R.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J.; Zarzycki, R. Macrobicyclic Polyethers as Second Sphere Ligands for Tetraammineplatinum(II). *Angew. Chem. Int. Ed. Engl.* **1987**, *26*, 693–696.
48. Ballardini, R.; Gandolfi, M. T.; Balzani, V.; Kohnke, F. H.; Stoddart, J. F. Second-Sphere Photochemistry and Photo-physics: Luminescence of the $[\text{Pt}(\text{Bpy})(\text{NH}_3)_2]^{2+}$ -Dibenzo [30]Crown-10 Adduct. *Angew. Chem. Int. Ed. Engl.* **1988**, *27*, 692–694.
49. Ballardini, R.; Gandolfi, M. T.; Prodi, L.; Ciano, M.; Balzani, V.; Kohnke, F. H.; Shahriari-Zavareh, H.; Spencer, N.; Stoddart, J. F. Supramolecular Photochemistry and Photophysics. Adducts of $\text{Pt}(\text{Bpy})(\text{NH}_3)_2^{2+}$ with Aromatic Crown Ethers. *J. Am. Chem. Soc.* **1989**, *111*, 7072–7078.
50. Cameron, B. R.; Corrent, S. S.; Loeb, S. J. Transition Metal Complexes as Both Host and Guest: Second-Sphere Coordination between a Pt-Azacrown Ether Host and a Pt-NH₃ Guest. *Angew. Chem. Int. Ed. Engl.* **1996**, *34*, 2689–2691.

51. Harada, A.; Takahashi, S. Preparation and Properties of Cyclodextrin-Ferrocene Inclusion Complexes. *J. Chem. Soc. Chem. Commun.* **1984**, 645–646.
52. Alston, D. R.; Stoddart, J. F.; Williams, D. J. The Isolation and X-Ray Structure of an Adduct Formation between 18-Crown-6 and Cisplatin. *J. Chem. Soc. Chem. Commun.* **1985**, 9, 532–533.
53. Alston, D. R.; Lilley, T. H.; Stoddart, J. F. The Binding of 1,1-Cyclo-Butanedicarboxylatodiammine Platinum(II) by α -Cyclodextrin in Aqueous Solution. *J. Chem. Soc. Chem. Commun.* **1985**, 1600–1602.
54. Alston, D. R.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J. The X-Ray Crystal Structure of a 1:1 Adduct between α -Cyclodextrin and Cyclobutane-1,1-Dicarboxylatodiammine-platinum(II). *J. Chem. Soc. Chem. Commun.* **1985**, 1602–1604.
55. Alston, D. R.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J. Cyclodextrins as Second Sphere Ligands for Transition Metal Complexes—The X-Ray Crystal Structure of [Rh(Cod)(NH₃)₂· α -Cyclodextrin][PF₆]₂·6H₂O. *Angew. Chem. Int. Ed. Engl.* **1985**, 24, 786–787.
56. Harada, A.; Takahashi, S. Preparation and Properties of Inclusion Compounds of Transition Metal Complexes of Cyclo-Octa-1,5-Diene and Norbornadiene with Cyclodextrins. *J. Chem. Soc. Chem. Commun.* **1986**, 1229.
57. Kamitori, S.; Hirotsu, K.; Higuchi, T. Crystal and Molecular Structures of Double Macrocyclic Inclusion Complexes Composed of Cyclodextrins, Crown Ethers, and Cations. *J. Am. Chem. Soc.* **1987**, 109, 2409–2414.
58. Alston, D. R.; Slawin, A. M. Z.; Stoddart, J. F.; Williams, D. J.; Zarzycki, R. Second Sphere Coordination Adducts of Phosphane-Transition Metal Complexes with β -Cyclodextrin and Its Methylated Derivative. *Angew. Chem. Int. Ed. Engl.* **1988**, 27, 1184–1185.
59. Alston, D. R.; Ashton, P. R.; Lilley, T. H.; Fraser Stoddart, J.; Zarzycki, R.; Slawin, A. M. Z.; Williams, D. J. Second-Sphere Co-Ordination of Carboplatin and Rhodium Complexes by Cyclodextrins (Cyclomalto-Oligosaccharides). *Carbohydr. Res.* **1989**, 192, 259–281.
60. Harada, A.; Yamamoto, S.; Takahashi, S. Preparation and Properties of Inclusion Compounds of Transition-Metal Complexes of Cycloocta-1,5-diene and Norbornadiene with Cyclodextrins. *Organometallics* **1989**, 8, 2560–2563.
61. Harada, A.; Shimada, M.; Takahashi, S. γ -Cyclodextrin as a Second Sphere Coordinating Ligand for Cobalt Cluster Complexes. *Chem. Lett.* **1989**, 18, 275–276.
62. Ando, I.; Ujimoto, K.; Kurihara, H. Effect of Second-Sphere Coordination 9. Partial Inclusion of Pentaammineruthenium(II) and Pentacyanoferrate(II) Complexes with Nitrogen Heterocycles into a Cyclodextrin Cavity. *Bull. Chem. Soc. Jpn.* **2001**, 74, 717–721.
63. Engeldinger, E.; Armspach, D.; Matt, D.; Jones, P. G.; Welter, R. A Cyclodextrin Diphosphane as a First and Second Coordination Sphere Cavitand: Evidence for Weak C-H...Cl-M Hydrogen Bonds within Metal-Capped Cavities. *Angew. Chem. Int. Ed.* **2002**, 41, 2593–2596.
64. Engeldinger, E.; Armspach, D.; Matt, D.; Jones, P. G. Cyclodextrin Phosphanes as First and Second Coordination Sphere Cavitands. *Chem. Eur. J.* **2003**, 9, 3091–3105.
65. Liu, Y.; Zhong, R.-Q.; Zhang, H.-Y.; Song, H.-B. A Unique Tetramer of 4:5 β -Cyclodextrin-Ferrocene in the Solid State. *Chem. Commun.* **2005**, 2211–2213.
66. Atwood, J. L.; Orr, G. W.; Hamada, F.; Vincent, R. L.; Bott, S. G.; Robinson, K. D. Second-Sphere Coordination of Transition-Metal Complexes by Calix[4]Arenes. *J. Am. Chem. Soc.* **1991**, 113, 2760–2761.
67. Atwood, J. L.; Orr, G. W.; Hamada, F.; Vincent, R. L.; Bott, S. G.; Robinson, K. D. Calixarenes as Second-Sphere Ligands for Transition Metal Ions. Synthesis and Crystal Structure of [(H₂O)₅Ni(NC₅H₅)₂](Na)[Calix[4]arene Sulfonate]·3.5 H₂O and [(H₂O)₄Cu(NC₅H₅)₂](H₃O)₃ [Calix[4]arene Sulfonate]·10 H₂O. *J. Incl. Phenom. Mol. Recognit. Chem.* **1992**, 14, 37–46.
68. Atwood, J. L.; Orr, G. W.; Robinson, K. D. First Structural Authentication of Third-Sphere Coordination: [P-Sulfonato-calix[4]arene]⁵⁻ as a Third-Sphere Ligand for Eu³⁺. *Supramol. Chem.* **1994**, 3, 89–91.
69. Beer, P. D.; Drew, M. G. B.; Ogden, M. I. First- and Second-Sphere Co-ordination of a Lanthanum Cation by a Calix[4]arene Tetraamide in the Partial-Cone Conformation. *J. Chem. Soc. Dalton Trans.* **1997**, 1489–1492.
70. Junk, P. C.; Atwood, J. L. Use of Metal Carbonyls in the Formation of [H₅O₂⁺·15-Crown-5][MOC₄(H₂O)]⁻, (M=Mo, W), and a Second Sphere Coordination Complex in [mer-CrCl₃(H₂O)₃·15-Crown-5]. *J. Organomet. Chem.* **1998**, 565, 179–186.
71. Belhamel, K.; Nguyen, T. K. D.; Benamor, M.; Ludwig, R. Design of Calixarene-Type Ligands for Second Sphere Complexation of Noble Metal Ions. *Eur. J. Inorg. Chem.* **2003**, 2003, 4110–4116.
72. MacGillivray, L. R.; Spinney, H. A.; Reid, J. L.; Ripmeester, J. A. Entrapment of Ferrocenes within Supramolecular, Deep-Cavity Resorcin[4]arenes. *Chem. Commun.* **2000**, 517–518.
73. Shen, L.; Zhao, Y.; Dai, D.; Yang, Y.-W.; Wu, B.; Yang, X.-J. Stabilization of Grignard Reagents by A Pillar[5]arene Host—Schlenk Equilibria and Grignard Reactions. *Chem. Commun.* **2020**, 56, 1381–1384.
74. Thuéry, P. Lanthanide Complexes with Cucurbit[*n*]uril (*n* = 5, 6, and 7) and Perrhenate Ligands: New Examples of Encapsulation of Perrhenate Anions. *Inorg. Chem.* **2009**, 48, 4497–4513.
75. Quan, M. L. C.; Cram, D. J. Constrictive Binding of Large Guests by a Hemiacarcerand Containing Four Portals. *J. Am. Chem. Soc.* **1991**, 113, 2754–2755.
76. Ashton, P. R.; Claessens, C. G.; Hayes, W.; Stoddart, J. F.; Menzer, S.; White, A. J. P.; Williams, D. J. Molecular Mosaics Formed by a Square Cyclophane and Its Inclusion Complex with Ferrocene. *Angew. Chem. Int. Ed. Engl.* **1995**, 34, 1862–1865.
77. Ashton, P. R.; Menzer, S.; Raymo, F. M.; Shimizu, G. K. H.; Stoddart, J. F.; Williams, D. J. The Template-Directed Synthesis of Cyclobis(paraquat-4,4'-biphenylene). *Chem. Commun.* **1996**, 487–490.
78. Raymo, F. M.; Stoddart, J. F. Self-Assembling Cyclobis(paraquat-4,4'-biphenylene). *Pure Appl. Chem.* **1996**, 68, 313–322.

79. Asakawa, M.; Ashton, P. R.; Menzer, S.; Raymo, F. M.; Stoddart, J. F.; White, A. J. P.; Williams, D. J. Cyclobis(paraquat-4,4'-biphenylene)—An Organic Molecular Square. *Chem. Eur. J.* **1996**, *2*, 877–893.
80. Kickham, J. E.; Loeb, S. J.; Murphy, S. L. Molecular Recognition of Nucleobases via Simultaneous First- and Second-Sphere Coordination. *J. Am. Chem. Soc.* **1993**, *115*, 7031–7032.
81. Kickham, J. E.; Loeb, S. J. Simultaneous First- and Second-Sphere Coordination. Organopalladium Metallo-receptors for Water, Ammonia, Amines, Hydrazine, and the Hydrazinium Ion. *Inorg. Chem.* **1995**, *34*, 5656–5665.
82. Kickham, J. E.; Loeb, S. J. Simultaneous First- and Second-Sphere Coordination. Organopalladium Crown Ether Complexes as Metallo-receptors for *o*-Aminopyridine Derivatives. *Inorg. Chem.* **1994**, *33*, 4351–4359.
83. Kickham, J. E.; Loeb, S. J. Receptors for the Hydrazinium Ion. Simultaneous First- and Second-Sphere Coordination inside Organopalladium Crown Ether Complexes. *J. Chem. Soc. Chem. Commun.* **1993**, 1848–1850.
84. Steed, J. W. First- and Second-Sphere Coordination Chemistry of Alkali Metal Crown Ether Complexes. *Coord. Chem. Rev.* **2001**, *215*, 171–221.
85. Beauchamp, D. A.; Loeb, S. J. Hydrogen-Bonded Networks through Second-Sphere Coordination. *Chem. Eur. J.* **2002**, *8*, 5084–5088.
86. Mercer, D. J.; Loeb, S. J. Metal-Based Anion Receptors: An Application of Second-Sphere Coordination. *Chem. Soc. Rev.* **2010**, *39*, 3612–3620.
87. Zhao, M.; Wang, H.-B.; Ji, L.-N.; Mao, Z.-W. Insights into Metalloenzyme Microenvironments: Biomimetic Metal Complexes with a Functional Second Coordination Sphere. *Chem. Soc. Rev.* **2013**, *42*, 8360–8375.
88. Guo, F.; Martí-Rujas, J. Second Sphere Coordination of Hybrid Metal-Organic Materials: Solid State Reactivity. *Dalton Trans.* **2016**, *45*, 13648–13662.
89. Therrien, B. The Role of the Second Coordination Sphere in the Biological Activity of Arene Ruthenium Metalla-Assemblies. *Front. Chem.* **2018**, *6*, 1–7.
90. Shaffer, C. C.; Smith, B. D. Macrocyclic and Acyclic Supramolecular Elements for Co-Precipitation of Square-Planar Gold(III) Tetrahalide Complexes. *Org. Chem. Front.* **2021**, *8*, 1294–1301.
91. Liu, W.; Stoddart, J. F. Emergent Behavior in Nanoconfined Molecular Containers. *Chem* **2021**, *7*, 919–947.
92. Liu, W.; Samanta, S. K.; Smith, B. D.; Isaacs, L. Synthetic Mimics of Biotin/(Strept)Avidin. *Chem. Soc. Rev.* **2017**, *46*, 2391–2403.
93. Cameron, B. R.; Loeb, S. J.; Yap, G. P. A. Calixarene Metallo-receptors. Synthesis and Molecular Recognition Properties of Upper-Rim Functionalized Calix[4]arenes Containing an Organopalladium Binding Site. *Inorg. Chem.* **1997**, *36*, 5498–5504.
94. Kickham, J. E.; Loeb, S. J.; Murphy, S. L. Molecular Receptors for Adenine and Guanine Employing Metal Coordination, Hydrogen-Bonding and π -Stacking Interactions. *Chem. Eur. J.* **1997**, *3*, 1203–1213.
95. Horeau, M.; Lautrette, G.; Wicher, B.; Blot, V.; Lebreton, J.; Pipelier, M.; Dubreuil, D.; Ferrand, Y.; Huc, I. Metal-Coordination-Assisted Folding and Guest Binding in Helical Aromatic Oligoamide Molecular Capsules. *Angew. Chem. Int. Ed.* **2017**, *129*, 6927–6931.
96. Mateus, P.; Wicher, B.; Ferrand, Y.; Huc, I. Alkali and Alkaline Earth Metal Ion Binding by a Foldamer Capsule: Selective Recognition of Magnesium Hydrate. *Chem. Commun.* **2017**, *53*, 9300–9303.
97. Mateus, P.; Wicher, B.; Ferrand, Y.; Huc, I. Carbohydrate Binding through First- and Second-Sphere Coordination within Aromatic Oligoamide Metallofoldamers. *Chem. Commun.* **2018**, *54*, 5078–5081.
98. Meunier, A.; Singleton, M. L.; Kauffmann, B.; Granier, T.; Lautrette, G.; Ferrand, Y.; Huc, I. Aromatic Foldamers as Scaffolds for Metal Second Coordination Sphere Design. *Chem. Sci.* **2020**, *11*, 12178–12186.
99. Zhang, L.; Stephens, A. J.; Lemonnier, J. F.; Pirvu, L.; Vitorica-Yrezabal, I. J.; Robinson, C. J.; Leigh, D. A. Coordination Chemistry of a Molecular Pentafoil Knot. *J. Am. Chem. Soc.* **2019**, *141*, 3952–3958.
100. Ayme, J.-F.; Beves, J. E.; Campbell, C. J.; Gil-Ramírez, G.; Leigh, D. A.; Stephens, A. J. Strong and Selective Anion Binding within the Central Cavity of Molecular Knots and Links. *J. Am. Chem. Soc.* **2015**, *137*, 9812–9815.
101. Marcos, V.; Stephens, A. J.; Jaramillo-García, J.; Nussbaumer, A. L.; Woltering, S. L.; Valero, A.; Lemonnier, J. F.; Vitorica-Yrezabal, I. J.; Leigh, D. A. Allosteric Initiation and Regulation of Catalysis with a Molecular Knot. *Science* **2016**, *352*, 1555–1559.
102. Fielden, S. D. P.; Leigh, D. A.; Woltering, S. L. Molecular Knots. *Angew. Chem. Int. Ed.* **2017**, *56*, 11166–11194.
103. Stoddart, J. F. Mechanically Interlocked Molecules (MIMs)—Molecular Shuttles, Switches, and Machines (Nobel Lecture). *Angew. Chem. Int. Ed.* **2017**, *56*, 11904–11125.
104. Bruns, C. J.; Stoddart, J. F. *The Nature of the Mechanical Bond*; Wiley: Hoboken, NJ, **2016**.
105. Stoddart, J. F. The Chemistry of the Mechanical Bond. *Chem. Soc. Rev.* **2009**, *38*, 1802–1820.
106. Stoddart, J. F.; Colquhoun, H. M. Big and Little Meccano. *Tetrahedron* **2008**, *64*, 8231–8263.
107. Philp, D.; Stoddart, J. F. Self-Assembly in Natural and Unnatural Systems. *Angew. Chem. Int. Ed. Engl.* **1996**, *35*, 1155–1196.
108. Amabilino, D. B.; Stoddart, J. F. Interlocked and Intertwined Structures and Superstructures. *Chem. Rev.* **1995**, *95*, 2725–2828.
109. Stoddart, J. F. Putting Mechanically Interlocked Molecules (MIMs) to Work in Tomorrow's World. *Angew. Chem. Int. Ed.* **2014**, *53*, 11102–11104.
110. Wasserman, E. The Preparation of Interlocking Rings: A Catenane. *J. Am. Chem. Soc.* **1960**, *82*, 4433–4434.
111. Schill, G.; Lüttringhaus, A. The Preparation of Catena Compounds by Directed Synthesis. *Angew. Chem. Int. Ed.* **1964**, *3*, 546–547.
112. Dietrich-Buchecker, C. O.; Sauvage, J.-P.; Kintzinger, J.-P. Une Nouvelle Famille de Molécules: Les Metallo-Catenanes. *Tetrahedron Lett.* **1983**, *24*, 5095–5098.

113. Johnston, A. G.; Leigh, D. A.; Pritchard, R. J.; Deegan, M. D. Facile Synthesis and Solid-State Structure of a Benzylic Amide [2]Catenane. *Angew. Chem. Int. Ed.* **1995**, *34*, 1209–1212.
114. Johnston, A. G.; Leigh, D. A.; Nezhat, L.; Smart, J. P.; Deegan, M. D. Structurally Diverse and Dynamically Versatile Benzylic Amide [2]Catenanes Assembled Directly from Commercially Available Precursors. *Angew. Chem. Int. Ed.* **1995**, *34*, 1212–1216.
115. Asakawa, M.; Ashton, P. R.; Balzani, V.; Credi, A.; Hamers, C.; Mattersteig, G.; Montalti, M.; Shipway, A. N.; Spencer, N.; Stoddart, J. F.; Tolley, M. S.; Venturi, M.; White, A. J. P.; Williams, D. J. A Chemically and Electrochemically Switchable [2]Catenane Incorporating a Tetrathiafulvalent Unit. *Angew. Chem. Int. Ed.* **1998**, *37*, 333–337.
116. Sauvage, J.-P. From Chemical Topology to Molecular Machines (Nobel Lecture). *Angew. Chem. Int. Ed.* **2017**, *56*, 11080–11093.
117. Fujita, M.; Ogura, K. Supramolecular Self-Assembly of Macrocycles, Catenanes, and Cages through Coordination of Pyridine-Based Ligands to Transition Metals. *Bull. Chem. Soc. Jpn.* **1996**, *69*, 1471–1482.
118. Nepogodiev, S. A.; Stoddart, J. F. Cyclodextrin-Based Catenanes and Rotaxanes. *Chem. Rev.* **1998**, *98*, 1959–1976.
119. Fujita, M. Self-Assembly of [2]Catenanes Containing Metals in Their Backbones. *Acc. Chem. Res.* **1999**, *32*, 53–61.
120. Faiz, J. A.; Heitz, V.; Sauvage, J.-P. Design and Synthesis of Porphyrin-Containing Catenanes and Rotaxanes. *Chem. Soc. Rev.* **2009**, *38*, 422–442.
121. Wu, Q.; Rauscher, P. M.; Lang, X.; Wojtecki, R. J.; de Pablo, J. J.; Hore, M. J. A.; Rowan, S. J. Poly[n]Catenanes: Synthesis of Molecular Interlocked Chains. *Science* **2017**, *358*, 1434–1439.
122. Erbas-Cakmak, S.; Fielden, S. D. P.; Karaca, U.; Leigh, D. A.; McTernan, C. T.; Tetlow, D. J.; Wilson, M. R. Rotary and Linear Molecular Motors Driven by Pulses of a Chemical Fuel. *Science* **2017**, *358*, 340–343.
123. Harrison, I. T.; Harrison, S. The Synthesis of a Stable Complex of a Macrocycle and a Threaded Chain. *J. Am. Chem. Soc.* **1967**, *89*, 5723–5724.
124. Schill, G.; Zollenkopf, H. Rotaxane-Verbindungen, I. *Liebigs Ann. Chem.* **1969**, *721*, 53–74.
125. Anelli, P. L.; Spencer, N.; Stoddart, J. F. A Molecular Shuttle. *J. Am. Chem. Soc.* **1991**, *113*, 5131–5133.
126. Gassensmith, J. J.; Baumes, J. M.; Smith, B. D. Discovery and Early Development of Squaraine Rotaxanes. *Chem. Commun.* **2009**, 6329–6338.
127. Bruns, C. J.; Stoddart, J. F. Rotaxane-Based Molecular Muscles. *Acc. Chem. Res.* **2014**, *47*, 2186–2199.
128. Durola, F.; Heitz, V.; Reviriego, F.; Roche, C.; Sauvage, J.-P.; Sour, A.; Trolez, Y. Cyclic [4]Rotaxanes Containing Two Parallel Porphyrinic Plates: Toward Switchable Molecular Receptors and Compressors. *Acc. Chem. Res.* **2014**, *47*, 633–645.
129. Langton, M. J.; Beer, P. D. Rotaxane and Catenane Host Structures for Sensing Charged Guest Species. *Acc. Chem. Res.* **2014**, *47*, 1935–1949.
130. Leigh, D. A.; Marcos, V.; Wilson, M. R. Rotaxane Catalysts. *ACS Catal.* **2014**, *4*, 4490–4497.
131. van Dongen, S. F. M.; Elemans, J. A. A. W.; Rowan, A. E.; Nolte, R. J. M. Processive Catalysis. *Angew. Chem. Int. Ed.* **2014**, *53*, 11420–11428.
132. Hashidzume, A.; Yamaguchi, H.; Harada, A. Cyclodextrin-Based Rotaxanes: From Rotaxanes to Polyrotaxanes and Further to Functional Materials. *Eur. J. Org. Chem.* **2019**, *2019*, 3344–3357.
133. Northrop, B. H.; Aricó, F.; Tangchiavang, N.; Badjić, J. D.; Stoddart, J. F. Template-Directed Synthesis of Mechanically Interlocked Molecular Bundles Using Dynamic Covalent Chemistry. *Org. Lett.* **2006**, *8*, 3899–3902.
134. Williams, A. R.; Northrop, B. H.; Chang, T.; Stoddart, J. F.; White, A. J. P.; Williams, D. J. Suitanes. *Angew. Chem. Int. Ed.* **2006**, *45*, 6665–6669.
135. Pun, A.; Hanifi, D. A.; Kiel, G.; O'Brien, E.; Liu, Y. Facile Route to an All-Organic, Triply Threaded, Interlocked Structure by Templated Dynamic Clipping. *Angew. Chem. Int. Ed.* **2012**, *51*, 13119–13122.
136. Zhu, K.; Baggi, G.; Vukotic, V. N.; Loeb, S. J. Reversible Mechanical Protection: Building a 3D “Suit” around a T-Shaped Benzimidazole Axle. *Chem. Sci.* **2017**, *8*, 3898–3904.
137. Liu, W.; Stern, C. L.; Stoddart, J. F. Suit[4]ane. *J. Am. Chem. Soc.* **2020**, *142*, 10273–10278.
138. Chen, X. Y.; Shen, D.; Cai, K.; Jiao, Y.; Wu, H.; Song, B.; Zhang, L.; Tan, Y.; Wang, Y.; Feng, Y.; Stern, C. L.; Stoddart, J. F. Suit[3]ane. *J. Am. Chem. Soc.* **2020**, *142*, 20152–20160.
139. Xu, H.; Lin, M.-D.; Yuan, J.; Zhou, B.; Mu, Y.; Huo, Y.; Zhu, K. Fluorescence Emission Enhancement of a T-Shaped Benzimidazole with a Mechanically-Interlocked ‘Suit.’ *Chem. Commun.* **2021**, *57*, 3239–3242.
140. Balzani, V.; Credi, A.; Raymo, F. M.; Stoddart, J. F. Artificial Molecular Machines. *Angew. Chem. Int. Ed.* **2000**, *39*, 3348–3391.
141. Collin, J.-P.; Heitz, V.; Bonnet, S.; Sauvage, J.-P. Transition Metal-Complexed Catenanes and Rotaxanes in Motion: Towards Molecular Machines. *Inorg. Chem. Commun.* **2005**, *8*, 1063–1074.
142. Browne, W. R.; Feringa, B. L. Making Molecular Machines Work. *Nat. Nanotechnol.* **2006**, *1*, 25–35.
143. Coskin, A.; Banaszak, M.; Astumian, R. D.; Stoddart, J. F.; Grzybowski, B. A. Great Expectations: Can Artificial Molecular Machines Deliver on Their Promise? *Chem. Soc. Rev.* **2012**, *41*, 19–30.
144. Qiu, Y.; Feng, Y.; Guo, Q.-H.; Astumian, R. D.; Stoddart, J. F. Pumps through the Ages. *Chem* **2020**, *6*, 1952–1977.
145. Feng, Y.; Ovalle, M.; Seale, J. S. W.; Lee, C. K.; Kim, D. J.; Stoddart, J. F. Molecular Pumps and Motors. *J. Am. Chem. Soc.* **2021**, *143*, 5569–5591.
146. Lewandowski, B.; De Bo, G.; Ward, J. W.; Papmeyer, M.; Kuschel, S.; Aldegunde, M. J.; Gramlich, P. M. E.; Heckmann, D.; Goldup, S. M.; Souza, D. M.; Fernandes, A. E.; Leigh, D. A. Sequence-Specific Peptide Synthesis by an Artificial Small-Molecule Machine. *Science* **2013**, *339*, 189–193.
147. Cheng, C.; Stoddart, J. F. Wholly Synthetic Molecular Machines. *ChemPhysChem* **2016**, *17*, 1780–1793.

148. Wilson, M. R.; Solà, J.; Carlone, A.; Goldup, S. M.; Leb-rasseur, N.; Leigh, D. A. An Autonomous Chemically Fuelled Small-Molecule Motor. *Nature* **2016**, *534*, 235–240.
149. Pezzato, C.; Cheng, C.; Stoddart, J. F.; Astumian, R. D. Mastering the Non-Equilibrium Assembly and Operation of Molecular Machines. *Chem. Soc. Rev.* **2017**, *46*, 5491–5507.
150. Kassem, S.; Lee, A. T. L.; Leigh, D. A.; Marcos, V.; Palmer, L. I.; Pisano, S. Stereodivergent Synthesis with a Program-mable Molecular Machine. *Nature* **2017**, *549*, 374–378.
151. Wang, Y.; Frascioni, M.; Stoddart, J. F. Introducing Stable Radicals into Molecular Machines. *ACS Cent. Sci.* **2017**, *3*, 927–935.
152. Qiu, Y.; Song, B.; Pezzato, C.; Shen, D.; Liu, W.; Zhang, L.; Feng, Y.; Guo, Q. H.; Cai, K.; Li, W.; Chen, H.; Nguyen, M. T.; Shi, Y.; Cheng, C.; Dean Astumian, R.; Li, X.; Stoddart, J. F. A Precise Polyrotaxane Synthesizer. *Science* **2020**, *368*, 1247–1253.
153. Corra, S.; Curcio, M.; Baroncini, M.; Silvi, S.; Credi, A. Photoactivated Artificial Molecular Machines that Can Per-form Tasks. *Adv. Mater.* **2020**, *32*, 1906064.
154. Vickers, M. S.; Beer, P. D. Anion Templated Assembly of Mechanically Interlocked Structures. *Chem. Soc. Rev.* **2007**, *36*, 211–225.
155. Griffiths, K. E.; Stoddart, J. F. Template-Directed Syn-thesis of Donor/Acceptor [2]Catenanes and [2]Rotaxanes. *Pure Appl. Chem.* **2008**, *80*, 485–506.
156. Mullen, K. M.; Beer, P. D. Sulfate Anion Templatation of Macrocycles, Capsules, Interpenetrated and Interlocked Structures. *Chem. Soc. Rev.* **2009**, *38*, 1701–1713.
157. Crowley, J. D.; Goldup, S. M.; Lee, A.-L.; Leigh, D. A.; McBurney, R. T. Active Metal Template Synthesis of Rotax-anes, Catenanes and Molecular Shuttles. *Chem. Soc. Rev.* **2009**, *38*, 1530–1541.
158. Aucagne, V.; Hänni, K. D.; Leigh, D. A.; Lusby, P. J.; Walker, D. B. Catalytic “Click” Rotaxanes: A Substoichio-metric Metal-Template Pathway to Mechanically Interlocked Architectures. *J. Am. Chem. Soc.* **2006**, *128*, 2186–2187.
159. Saito, S.; Takahashi, E.; Nakazono, K. Synthesis of [2] Rotaxanes by the Catalytic Reactions of a Macrocyclic Cop-per Complex. *Org. Lett.* **2006**, *8*, 5133–5136.
160. Crowley, J. D.; Hänni, K. D.; Lee, A.-L.; Leigh, D. A. [2] Rotaxanes through Palladium Active-Template Oxidative Heck Cross-Couplings. *J. Am. Chem. Soc.* **2007**, *129*, 12092–12093.
161. Berná, J.; Crowley, J. D.; Goldup, S. M.; Hänni, K. D.; Lee, A.-L.; Leigh, D. A. A Catalytic Palladium Active-Metal Tem-plate Pathway to [2]Rotaxanes. *Angew. Chem. Int. Ed.* **2007**, *46*, 5709–5713.
162. Aucagne, V.; Berná, J.; Crowley, J. D.; Goldup, S. M.; Hänni, K. D.; Leigh, D. A.; Lusby, P. J.; Ronaldson, V. E.; Slawin, A. M. Z.; Viterisi, A.; Walker, D. B. Catalytic “Active-Metal” Template Synthesis of [2]Rotaxanes, [3]Rotaxanes, and Mo-lecular Shuttles, and Some Observations on the Mechanism of the Cu(I)-Catalyzed Azide-Alkyne 1,3-Cycloaddition. *J. Am. Chem. Soc.* **2007**, *129*, 11950–11963.
163. Sato, Y.; Yamasaki, R.; Saito, S. Synthesis of [2]Cate-nanes by Oxidative Intramolecular Diyne Coupling Mediated by Macrocyclic Copper(I) Complexes. *Angew. Chem. Int. Ed.* **2009**, *48*, 504–507.
164. Goldup, S. M.; Leigh, D. A.; Long, T.; McGonigal, P. R.; Symes, M. D.; Wu, J. Active Metal Template Synthesis of [2] Catenanes. *J. Am. Chem. Soc.* **2009**, *131*, 15924–15929.
165. Barran, P. E.; Cole, H. L.; Goldup, S. M.; Leigh, D. A.; McGonigal, P. R.; Symes, M. D.; Wu, J.; Zengerle, M. Active-Metal Template Synthesis of a Molecular Trefoil Knot. *Angew. Chem. Int. Ed.* **2011**, *50*, 12280–12284.
166. Ibukuro, F.; Fujita, M.; Yamaguchi, K.; Sauvage, J.-P. Quantitative and Spontaneous Formation of a Doubly Inter-locking [2]Catenane Using Copper(I) and Palladium(II) as Templating and Assembling Centers. *J. Am. Chem. Soc.* **1999**, *121*, 11014–11015.
167. Dietrich-Buchecker, C.; Colasson, B.; Fujita, M.; Hori, A.; Geum, N.; Sakamoto, S.; Yamaguchi, K.; Sauvage, J.-P. Quan-titative Formation of [2]Catenanes Using Copper(I) and Palladium(II) as Templating and Assembling Centers: The Entwining Route and the Threading Approach. *J. Am. Chem. Soc.* **2003**, *125*, 5717–5725.
168. Liu, Y.; Zhao, Y.; Sun, X.; Furukawa, H.; Ma, Y.; Oleynikov, P.; Gandara, F.; Liu, Z.; Suenaga, K.; Zhu, H.; Zhu, C.; Alsham-mari, A. S.; Zhang, X.; Terasaki, O.; Yaghi, O. M. Weaving of Organic Threads into a Crystalline Covalent Organic Frame-work. *Science* **2016**, *351*, 365–369.
169. Blight, B. A.; Van Noortwyk, K. A.; Wisner, J. A.; Jen-nings, M. C. [2]Pseudorotaxanes Through Second-Sphere Coordination. *Angew. Chem. Int. Ed.* **2005**, *44*, 1499–1504.
170. Blight, B. A.; Wisner, J. A.; Jennings, M. C. Synthesis of a [2]Rotaxane through First- and Second-Sphere Coordina-tion. *Chem. Commun.* **2006**, 4593–4595.
171. Blight, B. A.; Wei, X.; Wiener, J. A.; Jennings, M. C. [2] Pseudorotaxane and [2]Rotaxane Molecular Shuttles: Self-Assembly through Second-Sphere Coordination of Thiocy-anate Ligands. *Inorg. Chem.* **2007**, *46*, 8445–8447.
172. Blight, B. A.; Wisner, J. A.; Jennings, M. C. Stability of [2] Pseudorotaxanes Templated through Second-Sphere Coordination. *Inorg. Chem.* **2009**, *48*, 1920–1927.
173. Blight, B. A.; Wisner, J. A.; Jennings, M. C. Reversible Formation of a [2]Catenane through First- and Second-Sphere Coordination. *Angew. Chem. Int. Ed.* **2007**, *46*, 2835–2838.
174. Amini, M. M.; Rheingold, A. L.; Taylor, R. W.; Zuckerman, J. J. A New Environment for Water. The First Authenticated Example of Water Molecules Engaged in Twin, Three-Center Hydrogen Bonds: The Crystal and Molecular Structure of $\{[(CH_3)_2SnCl_2 \cdot H_2O]_2 \cdot 18\text{-Crown-6}\}_n$. *J. Am. Chem. Soc.* **1984**, *106*, 7289–7291.
175. Zhu, X.; Ma, Z.; Bi, W.; Wang, Y.; Yuan, D.; Cao R. Forma-tion of a Sandwich-Type Supercomplex through Second-Sphere Coordination of Functionalized Macrocyclic Polya-mines. *CrystEngComm* **2008**, *10*, 19–22.
176. Dalrymple, S. A.; Parvez, M.; Shimizu, G. K. H. Supramo-lecular Encapsulation of Hexaaquo Metal Ions by Second Sphere Coordination. *Chem. Commun.* **2001**, 2672–2673.
177. Dalrymple, S. A.; Parvez, M.; Shimizu, G. K. H. Intra- and Intermolecular Second-Sphere Coordination Chemistry: Formation of Capsules, Half-Capsules, and Extended

Structures with Hexaaquo- and Hexaamminemetal Ions. *Inorg. Chem.* **2002**, *41*, 6986–6996.

178. Dalrymple, S. A.; Shimizu, G. K. H. Exploiting Complementary Second-Sphere Effects in Supramolecular Coordination Solids. *Supramol. Chem.* **2003**, *15*, 591–606.

179. Dalrymple, S. A.; Shimizu, G. K. H. An Open Channel Coordination Framework Sustained by Cooperative Primary and Secondary Sphere Interactions. *Chem. Commun.* **2002**, 2224–2225.

180. Reddy, D. S.; Duncan, S.; Shimizu, G. A Family of Supramolecular Inclusion Solids Based Upon Second-Sphere Interactions. *Angew. Chem.* **2003**, *115*, 1398–1402.

181. Taylor, J. M.; Dwyer, P. J.; Reid, J. W.; Gelfand, B. S.; Lim, D. Woon; Donoshita, M.; Veinberg, S. L.; Kitagawa, H.; Vukotic, V. N.; Shimizu, G. K. H. Holding Open Micropores with Water: Hydrogen-Bonded Networks Supported by Hexaaquachromium(III) Cations. *Chem* **2018**, *4*, 868–878.

182. Ubaldini, S.; Massidda, R.; Vegliò, F.; Beolchini, F. Gold Stripping by Hydro-Alcoholic Solutions from Activated Carbon: Experimental Results and Data Analysis by a Semi-Empirical Model. *Hydrometallurgy* **2006**, *81*, 40–44.

183. Soleimani, M.; Kaghazchi, T. Gold Recovery from Loaded Activated Carbon Using Different Solvents. *J. Chin. Inst. Chem. Eng.* **2008**, *39*, 9–11.

184. Bunney, K.; Jeffrey, M. I.; Pleyzier, R.; Breuer, P. L. Selective Elution of Gold, Silver and Mercury Cyanide from Activated Carbon. *Miner. Metall. Process.* **2010**, *27*, 205–211.

185. Liu, W.; Jones, L. O.; Wu, H.; Stern, C. L.; Sponenburg, R. A.; Schatz, G. C.; Stoddart, J. F. Supramolecular Gold Stripping from Activated Carbon Using α -Cyclodextrin. *J. Am. Chem. Soc.* **2021**, *143*, 1984–1992.

186. Liu, Z.; Frascioni, M.; Lei, J.; Brown, Z. J.; Zhu, Z.; Cao, D.; Iehl, J.; Liu, G.; Fahrenbach, A. C.; Botros, Y. Y.; Farha, O. K.; Hupp, J. T.; Mirkin, C. A.; Stoddart, J. F. Selective Isolation of Gold Facilitated by Second-Sphere Coordination with α -Cyclodextrin. *Nat. Commun.* **2013**, *4*, 1855.

187. Liu, Z.; Samanta, A.; Lei, J.; Sun, J.; Wang, Y.; Stoddart, J. F. Cation-Dependent Gold Recovery with α -Cyclodextrin Facilitated by Second-Sphere Coordination. *J. Am. Chem. Soc.* **2016**, *138*, 11643–11653.

188. Smaldone, R. A.; Forgan, R. S.; Furukawa, H.; Gassensmith, J. J.; Slawin, A. M. Z.; Yaghi, O. M.; Stoddart, J. F. Metal–Organic Frameworks from Edible Natural Products. *Angew. Chem. Int. Ed.* **2010**, *49*, 8630–8634.

189. Gassensmith, J. J.; Furukawa, H.; Smaldone, R. A.; Forgan, R. S.; Botros, Y. Y.; Yaghi, O. M.; Stoddart, J. F. Strong and Reversible Binding of Carbon Dioxide in a Green Metal–Organic Framework. *J. Am. Chem. Soc.* **2011**, *133*, 15312–15315.

190. Forgan, R. S.; Smaldone, R. A.; Gassensmith, J. J.; Furukawa, H.; Cordes, D. B.; Li, Q.; Wilmer, C. E.; Botros, Y. Y.; Snurr, R. Q.; Slawin, A. M. Z.; Stoddart, J. F. Nanoporous Carbohydrate Metal–Organic Frameworks. *J. Am. Chem. Soc.* **2012**, *134*, 406–417.

191. Roy, I.; Stoddart, J. F. Cyclodextrin Metal–Organic Frameworks and Their Applications. *Acc. Chem. Res.* **2021**, *54*, 1440–1453.

192. Chen, L. X.; Liu, M.; Zhang, Y. Q.; Zhu, Q. J.; Liu, J. X.; Zhu, B. X.; Tao, Z. Outer Surface Interactions to Drive

Cucurbit[8]uril-Based Supramolecular Frameworks: Possible Application in Gold Recovery. *Chem. Commun.* **2019**, 55, 14271–14274.

193. Lin, R.-L.; Dong, Y.-P.; Tang, M.; Liu, Z.; Tao, Z.; Liu, J.-X. Selective Recovery and Detection of Gold with Cucurbit[*n*]urils (*n* = 5–7). *Inorg. Chem.* **2020**, *59*, 3850–3855.

194. Wu, H.; Jones, L. O.; Wang, Y.; Shen, D.; Liu, Z.; Zhang, L.; Cai, K.; Jiao, Y.; Stern, C. L.; Schatz, G. C.; Stoddart, J. F. High-Efficiency Gold Recovery Using Cucurbit[6]uril. *ACS Appl. Mater. Interfaces* **2020**, *12*, 38768–38777.

195. Dong, C. C.; Xiang, J. F.; Xu, L. J.; Gong, H. Y. MCl_4^{n-} (AuCl_4^- , PtCl_4^{2-} , or PdCl_4^{2-}) Anion Extraction from Na_nMCl_4 in Water Using a Tetraimidazolium Macrocyclic Receptor. *Tetrahedron Lett.* **2018**, *59*, 264–267.

196. Liu, W.; Oliver, A. G.; Smith, B. D. Macrocyclic Receptor for Precious Gold, Platinum, or Palladium Coordination Complexes. *J. Am. Chem. Soc.* **2018**, *140*, 6810–6813.

197. Wang, L.-L.; Tu, Y.-K.; Yao, H.; Jiang, W. 2,3-Dibutoxy-naphthalene-Based Tetralactam Macrocycles for Recognizing Precious Metal Chloride Complexes. *Beilstein J. Org. Chem.* **2019**, *15*, 1460–1467.

198. Shaffer, C. C.; Liu, W.; Oliver, A. G.; Smith, B. D. Supramolecular Paradigm for Capture and Co-Precipitation of Gold(III) Coordination Complexes. *Chem. Eur. J.* **2021**, *27*, 751–757.

199. Grommet, A. B.; Feller, M.; Klajn, R. Chemical Reactivity Under Nanoconfinement. *Nat. Nanotech.* **2020**, *15*, 256–271.

200. Merlau, M. L.; Mejia, M. D. P.; Nguyen, S. T.; Hupp, J. T. Artificial Enzymes Formed through Directed Assembly of Molecular Square Encapsulated Epoxidation Catalysts. *Angew. Chem. Int. Ed.* **2001**, *40*, 4239–4242.

201. Breslow, R.; Overman, L. E. An “Artificial Enzyme” Combining a Metal Catalytic Group and a Hydrophobic Binding Cavity. *J. Am. Chem. Soc.* **1970**, *92*, 1075–1077.

202. Breslow, R.; Zhang, B. Cleavage of Phosphate Esters by a Cyclodextrin Dimer Catalyst That Binds the Substrates together with La^{3+} and Hydrogen Peroxide. *J. Am. Chem. Soc.* **1994**, *116*, 7893–7894.

203. Breslow, R. Biomimetic Chemistry and Artificial Enzymes: Catalysis by Design. *Acc. Chem. Res.* **1995**, *28*, 146–153.

204. Breslow, R.; Zhang, X.; Xu, R.; Maletic, M.; Merger, R. Selective Catalytic Oxidation of Substrates that Bind to Metalloporphyrin Enzyme Mimics Carrying Two or Four Cyclodextrin Groups and Related Metallosalens. *J. Am. Chem. Soc.* **1996**, *118*, 11678–11679.

205. Zhang, B.; Breslow, R. Ester Hydrolysis by a Catalytic Cyclodextrin Dimer Enzyme Mimic with a Metallobipyridyl Linking Group. *J. Am. Chem. Soc.* **1997**, *119*, 1676–1681.

206. Breslow, R.; Dong, S. D. Biomimetic Reactions Catalyzed by Cyclodextrins and Their Derivatives. *Chem. Rev.* **1998**, *98*, 1997–2011.

207. Baugh, S. D. P.; Yang, Z.; Leung, D. K.; Wilson, D. M.; Breslow, R. Cyclodextrin Dimers as Cleavable Carriers of Photodynamic Sensitizers. *J. Am. Chem. Soc.* **2001**, *123*, 12488–12494.

208. Breslow, R. Artificial Enzymes. *Science* **2006**, *218*, 532–537.

209. Koblenz, T. S.; Wassenaar, J.; Reek, J. N. H. Reactivity within a Confined Self-Assembled Nanospace. *Chem. Soc. Rev.* **2008**, *37*, 247–262.
210. Raynal, M.; Ballester, P.; Vidal-Ferran, A.; Van Leeuwen, P. W. N. M. Supramolecular Catalysis. Part 2: Artificial Enzyme Mimics. *Chem. Soc. Rev.* **2014**, *43*, 1734–1787.
211. Brown, C. J.; Toste, F. D.; Bergman, R. G.; Raymond, K. N. Supramolecular Catalysis in Metal-Ligand Cluster Hosts. *Chem. Rev.* **2015**, *115*, 3012–3035.
212. Roland, S.; Suarez, J. M.; Sollogoub, M. Confinement of Metal-*N*-Heterocyclic Carbene Complexes to Control Reactivity in Catalytic Reactions. *Chem. Eur. J.* **2018**, *24*, 12464–12473.
213. Jongkind, L. J.; Caumes, X.; Hartendorp, A. P. T.; Reek, J. N. H. Ligand Template Strategies for Catalyst Encapsulation. *Acc. Chem. Res.* **2018**, *51*, 2115–2128.
214. Tan, C.; Chu, D.; Tang, X.; Liu, Y.; Xuan, W.; Cui, Y. Supramolecular Coordination Cages for Asymmetric Catalysis. *Chem. Eur. J.* **2019**, *25*, 662–672.
215. Fang, Y.; Powell, J. A.; Li, E.; Wang, Q.; Perry, Z.; Kirchon, A.; Yang, X.; Xiao, Z.; Zhu, C.; Zhang, L.; Huang, F.; Zhou, H. C. Catalytic Reactions within the Cavity of Coordination Cages. *Chem. Soc. Rev.* **2019**, *48*, 4707–4730.
216. Morimoto, M.; Bierschenk, S. M.; Xia, K. T.; Bergman, R. G.; Raymond, K. N.; Toste, F. D. Advances in Supramolecular Host-Mediated Reactivity. *Nat. Catal.* **2020**, *3*, 969–984.
217. Wang, K.; Jordan, J. H.; Hu, X. Y.; Wang, L. Supramolecular Strategies for Controlling Reactivity within Confined Nanospaces. *Angew. Chem. Int. Ed.* **2020**, *59*, 13712–13721.
218. Mitschke, B.; Turberg, M.; List, B. Confinement as a Unifying Element in Selective Catalysis. *Chem* **2020**, *6*, 2515–2532.
219. Trouvé, J.; Gramage-Doria, R. Beyond Hydrogen Bonding: Recent Trends of Outer Sphere Interactions in Transition Metal Catalysis. *Chem. Soc. Rev.* **2021**, *50*, 3565–3584.
220. Zhang, P.; Meijide Suárez, J.; Driant, T.; Derat, E.; Zhang, Y.; Ménand, M.; Roland, S.; Sollogoub, M. Cyclodextrin Cavity-Induced Mechanistic Switch in Copper-Catalyzed Hydroboration. *Angew. Chem. Int. Ed.* **2017**, *129*, 10961–10965.
221. Xu, G.; Leloux, S.; Zhang, P.; Meijide Suárez, J.; Zhang, Y.; Derat, E.; Ménand, M.; Bistri-Aslanoff, O.; Roland, S.; Leysens, T.; Riant, O.; Sollogoub, M. Capturing the Monomeric (L) CuH in NHC-Capped Cyclodextrin: Cavity-Controlled Chemoselective Hydrosilylation of α,β -Unsaturated Ketones. *Angew. Chem. Int. Ed.* **2020**, *59*, 7591–7597.
222. Hong, C. M.; Bergman, R. G.; Raymond, K. N.; Toste, F. D. Self-Assembled Tetrahedral Hosts as Supramolecular Catalysts. *Acc. Chem. Res.* **2018**, *51*, 2447–2455.
223. Bender, T. A.; Bergman, R. G.; Raymond, K. N.; Toste, F. D. A Supramolecular Strategy for Selective Catalytic Hydrogenation Independent of Remote Chain Length. *J. Am. Chem. Soc.* **2019**, *141*, 11806–11810.
224. Bender, T. A.; Morimoto, M.; Bergman, R. G.; Raymond, K. N.; Toste, F. D. Supramolecular Host-Selective Activation of Iodoarenes by Encapsulated Organometallics. *J. Am. Chem. Soc.* **2019**, *141*, 1701–1706.
225. García-Simón, C.; Gramage-Doria, R.; Raoufmoghadam, S.; Parella, T.; Costas, M.; Ribas, X.; Reek, J. N. H. Enantioselective Hydroformylation by a Rh-Catalyst Entrapped in a Supramolecular Metallocage. *J. Am. Chem. Soc.* **2015**, *137*, 2680–2687.
226. Slagt, V. F.; Reek, J. N. H.; Kamer, P. C. J.; Van Leeuwen, P. W. N. M. Assembly of Encapsulated Transition Metal Catalysts. *Angew. Chem. Int. Ed.* **2001**, *40*, 4271–4274.
227. Bocokić, V.; Kalkan, A.; Lutz, M.; Spek, A. L.; Gryko, D. T.; Reek, J. N. H. Capsule-Controlled Selectivity of a Rhodium Hydroformylation Catalyst. *Nat. Commun.* **2013**, *4*, 1–9.
228. Nurttila, S. S.; Linnebank, P. R.; Krachko, T.; Reek, J. N. H. Supramolecular Approaches to Control Activity and Selectivity in Hydroformylation Catalysis. *ACS Catal.* **2018**, *8*, 3469–3488.
229. Slagt, V. F.; Kamer, P. C. J.; Van Leeuwen, P. W. N. M.; Reek, J. N. H. Encapsulation of Transition Metal Catalysts by Ligand-Template Directed Assembly. *J. Am. Chem. Soc.* **2004**, *126*, 1526–1536.
230. Kuil, M.; Soltner, T.; Van Leeuwen, P. W. N. M.; Reek, J. N. H. High-Precision Catalysts: Regioselective Hydroformylation of Internal Alkenes by Encapsulated Rhodium Complexes. *J. Am. Chem. Soc.* **2006**, *128*, 11344–11345.
231. Rayder, T. M.; Bensalah, A. T.; Li, B.; Byers, J. A.; Tsung, C. Engineering Second Sphere Interactions in a Host-Guest Multicomponent Catalyst System for the Hydrogenation of Carbon Dioxide to Methanol. *J. Am. Chem. Soc.* **2021**, *143*, 1630–1640.
232. Cardin, C. J.; Kelly, J. M.; Quinn, S. J. Photochemically Active DNA-Intercalating Ruthenium and Related Complexes—Insights by Combining Crystallography and Transient Spectroscopy. *Chem. Sci.* **2017**, *8*, 4705–4723.
233. Hall, J. P.; O'Sullivan, K.; Naseer, A.; Smith, J. A.; Kelly, J. M.; Cardin, C. J. Structure Determination of an Intercalating Ruthenium Dipyrrophenazine Complex which Kinks DNA by Semiintercalation of a Tetraazaphenanthrene Ligand. *Proc. Natl. Acad. Sci. U. S. A.* **2011**, *108*, 17610–17614.
234. Song, H.; Kaiser, J. K.; Barton, J. K. Crystal Structure of Δ -[Ru(bpy)₂dppz]²⁺ Bound to Mismatched DNA Reveals Side-by-side Metalloinsertion and Intercalation. *Nat. Chem.* **2012**, *4*, 615–620.
235. Niyazi, H.; Hall, J. P.; O'Sullivan, K.; Winter, G.; Sorensen, T.; Kelly, J. M.; Cardin, C. J. Crystal Structures of Λ -[Ru(phen)₂(dppz)]²⁺ with Oligonucleotides Containing TA/TA and AT/AT Steps Show Two Intercalation Modes. *Nat. Chem.* **2012**, *4*, 621–628.
236. Hall, J. P.; Poynton, F. E.; Keane, P. M.; Gurung, S. P.; Brazier, J. A.; Cardin, D. J.; Winter, G.; Gunnlaugsson, T.; Sazanovich, I. V.; Towrie, M.; Cardin, C. J.; Kelly, J. M.; Quinn, S. J. Monitoring One-Electron Photo-Oxidation of Guanine in DNA Crystals Using Ultrafast Infrared Spectroscopy. *Nat. Chem.* **2015**, *7*, 961–967.
237. Vukotic, V. N.; Harris, K. J.; Zhu, K.; Schurko, R. W.; Loeb, S. J. Metal-Organic Frameworks with Dynamic Interlocked Components. *Nat. Chem.* **2012**, *4*, 456–460.
238. Zhu, K.; Vukotic, V. N.; Okeefe, C. A.; Schurko, R. W.; Loeb, S. J. Metal-Organic Frameworks with Mechanically Interlocked Pillars: Controlling Ring Dynamics in the Solid-State via a Reversible Phase Change. *J. Am. Chem. Soc.* **2014**, *136*, 7403–7409.

239. Zhu, K.; O'Keefe, C. A.; Vukotic, V. N.; Schurko, R. W.; Loeb, S. J. A Molecular Shuttle that Operates inside a Metal–Organic Framework. *Nat. Chem.* **2015**, *7*, 514–519.
240. Wilson, B. H.; Loeb, S. J. Integrating the Mechanical Bond into Metal–Organic Frameworks. *Chem* **2020**, *6*, 1604–1612.
241. Wilson, B. H.; Vojvodin, C. S.; Gholami, G.; Abdulla, L. M.; O'Keefe, C. A.; Schurko, R. W.; Loeb, S. J. Precise Spatial Arrangement and Interaction between Two Different Mobile Components in a Metal–Organic Framework. *Chem* **2021**, *7*, 202–211.