

# *Interactions of umami with the four other basic tastes in equi-intense aqueous solutions*

Article

Accepted Version

Wang, S., Dermiki, M., Methven, L., Kennedy, O. B. and Cheng, Q. (2022) Interactions of umami with the four other basic tastes in equi-intense aqueous solutions. *Food Quality and Preference*, 98. 104503. ISSN 0950-3293 doi: <https://doi.org/10.1016/j.foodqual.2021.104503> Available at <https://centaur.reading.ac.uk/102035/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodqual.2021.104503>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

1 **Interactions of umami with the four other basic tastes in equi-intense aqueous**  
2 **solutions**

3 **Authors:** Sijia Wang<sup>a</sup>, Maria Dermiki<sup>b</sup>, Lisa Methven<sup>a</sup>, Orla B Kennedy<sup>a</sup>, Qiaofen  
4 Cheng<sup>a\*</sup>

5 <sup>a</sup>Department of Food and Nutritional Sciences, University of Reading, Whiteknights,  
6 Reading, RG6 6DZ, UK

7 <sup>b</sup>Department of **Health and Nutritional Sciences**, Institute of Technology Sligo, Ash  
8 Lane, Sligo, F91 YW50, Ireland

9 \*Corresponding author: Dr Qiaofen Cheng, Department of Food and Nutritional  
10 Sciences, University of Reading, Whiteknights, Reading, RG6 6DZ, UK. Tel.: +44  
11 118 378 8719. Email address: q.cheng@reading.ac.uk

12

13 **Highlight**

- 14 ● Umami taste does not enhance or suppress sweet, salty, sour or bitter in equi-  
15 intense solutions.
- 16 ● Sweet, salty, sour and bitter significantly suppress umami taste in equi-intense  
17 solutions.
- 18 ● Sodium chloride plus glutamate tastants maintained salty and savoury taste  
19 perception.

20

21 **Key words**

22 Umami, salty, taste, mixtures, suppression, equi-intense

23

24 **Abstract**

25 Previous research has shown that the addition of equi-intense concentrations of taste  
26 compounds leads to mixture suppression, with sweetness being the least suppressed  
27 taste while being the strongest suppressor of the other taste stimuli. However, perceived  
28 intensity of umami (savoury) within complex mixtures is less defined. Since  
29 maintaining savoury taste of foods at reduced salt levels is a growing need, this study  
30 aims to investigate the role of umami in complex taste systems. Initially the  
31 concentrations of single tastants were adjusted until a trained sensory panel rated them  
32 as equi-intense using general labelled magnitude scale (gLMS). In order to evaluate the  
33 impact of umami taste on other tastes, and vice versa, three sample sets were prepared  
34 as binary and quinary systems. The first two sets utilised monosodium glutamate  
35 (MSG) as the umami tastant; one set without balancing the sodium level in MSG  
36 (sodium unbalanced) and another set accounting for it by the addition of sodium at an  
37 equivalent molarity to all but the umami single tastant solution (sodium balanced). The  
38 third set used monopotassium L-glutamate monohydrate (MPG) as the source of umami  
39 to overcome the confounding influence of sodium. All samples were rated by trained  
40 sensory panellists. The results of the three studies conclude that umami taste does not  
41 enhance or suppress the perception of any other taste in binary aqueous taste systems  
42 ( $p > 0.05$ ); whereas sweet, salty, sour and bitter significantly suppress the perception of  
43 umami in both binary and quinary systems ( $p < 0.05$ ).

44

45

## 46 1. Introduction

47 Cross-modal interactions between two or more sensory modalities, have been  
48 investigated as a strategy for the reduction of salt and sugar (Ponzo *et al.*, 2021). For  
49 example, odour-taste interactions have been explored for the reduction of sugar  
50 (Velazquez *et al.*, 2020) and the reduction of salt (Thomas-Danguin, Guichard & Salles,  
51 2019; Emorine *et al.*, 2021). Mojet, Heidema and Christ-Hazelho (2004) described how  
52 taste-taste interactions influenced taste in various real foods, and found that tastants  
53 evoking salty, sweet, bitter or umami could alter the perception of one or more other  
54 taste qualities in the product which they had been added to. Such taste-taste interactions  
55 can be useful in salt reduction strategies. For example, where potassium chloride (KCl)  
56 is used to replace sodium chloride (NaCl) it can increase bitterness in the final product;  
57 however, Abu *et al.* (2018) found that adding sweetness (via trehalose or sucrose) to a  
58 KCl/NaCl mixture effectively reduced bitterness without changing saltiness. Therefore  
59 taste-taste interactions are of relevance to the food scientist, with applications in salt  
60 and sugar reduction continuing to be a growing interest.

61 Psychophysical functions are used to study and express relationships between a  
62 stimulus and a response, or perceived sensation, such as taste. For individual taste  
63 stimuli, as the physical concentration increases the perceived intensity elicited by that  
64 compound also increases, but the rate of increase is not always directly proportional. It  
65 is dependent on both the specific tastant and whether the concentration is at relatively  
66 low levels (just above threshold, accelerating relationship), moderate levels (linear  
67 relationship) or high levels (decelerating relationship) (Bartoshuk, 1975; McBride,  
68 1987).

69 Such **stimulus** response relationships are subsequently modified in tastant mixtures. In  
70 a previous review, Keast and Breslin (2002a) concluded that perception of binary taste  
71 mixtures is dependent on the position of the taste stimulus on the psychophysical curve.  
72 Whether the concentration is within the linear or decelerating (plateau) phase of the  
73 curve, helps predict whether a particular tastant would cause enhancement or

74 suppression within a tastant mixture. In an earlier paper, McBride (1993) noted that the  
75 binary mixing of two different tastants produces three senses: an overall total intensity  
76 and a sensation from each of the two components; he suggested that the total intensity  
77 would be determined only by the strength of the stronger components.

78 In the case of more complex ternary and quaternary taste combinations, Bartoshuk  
79 (1975) found that tastants suppressed each other. The extent of suppression was  
80 dependent upon the function of the individual tastant; tastes where perception increased  
81 sharply with increasing concentration tended to cause greater suppression. Similarly on  
82 studying a tertiary taste mixture's intensity of sucrose, fructose, and citric acid,  
83 McBride and Finlay (1990) found that the total perceived strength of the mixture was  
84 determined by the perceptual intensity of the individual stronger components, and the  
85 sweetness and sourness of the mixture tended to suppress each other. Taking a  
86 modelling approach to understand the psychophysics of taste interaction, Schifferstein  
87 and Frijters (1993) concluded that a summation model (addition of individual  
88 component intensities) was sufficiently able to predict total taste intensity of a mixture.

89 Since many foods are formulated with tastants at moderate and not extreme levels, it is  
90 likely that the influence of taste stimuli in the linear phase of the psychophysical curve  
91 might be the most relevant. The approach taken by Green *et al.* (2010) focused on taste  
92 mixtures combined at perceptually equi-intense moderate (not extreme) concentrations.  
93 They tested taste interactions in the four taste mixtures (salt, sweet, bitter and sour)  
94 using equi-intense concentrations of sodium chloride, sucrose, quinine sulfate and citric  
95 acid. Moreover, four tastes qualities in binary, ternary and quaternary mixtures were  
96 also investigated. They concluded that suppression between stimuli in binary mixtures  
97 could predict taste perception in more complex combinations. For example, the sweet  
98 taste of sucrose tended to be the least suppressed quality, whereas it was a potent  
99 suppressor to all other tastes.

100 Umami tastants are widely used as flavour enhancers in food products, and especially  
101 in developing salt-reduced foods. In practice such enhancement may result from

102 complex ingredients, such as yeast extracts, that comprise both amino acids (especially  
103 glutamate) and 5'- nucleotides. However, literature often focuses on the understanding  
104 of simpler systems. A review paper by Maluly et al (2017) recommended that  
105 monosodium glutamate (MSG) could be used to reduce NaCl in a broad range of foods.  
106 In specific applications, Yamaguchi and Takahashi (1984) demonstrated that MSG  
107 could be used to reduce NaCl in a Japanese soup (Sumash-Jiru). Where MSG is used  
108 in combination with 5'-nucleotides, such as inosine-5'-monophosphate (IMP) and  
109 guanosine-5'-monophosphate (GMP), a much stronger umami taste can be achieved.  
110 Yamaguchi and Kimizuka (1979) found that the perceived umami intensity was  
111 affected by the ratio of IMP to MSG, and more recently Yamaguchi summarized that  
112 maximum taste intensity could be achieved with a 70:30 ratio of IMP to MSG  
113 (Yamaguchi, 1998). In using a combination of umami tastants, Dos et al. (2014) found  
114 that MSG, disodium inosinate, disodium guanylate could enhance flavour and maintain  
115 saltiness at 50% reduced NaCl when added into fermented cooked sausages.

116 However, there is limited understanding about how MSG performs in mixture of  
117 tastants, and how it interacts with other tastants, especially at equi-intense levels.  
118 Indeed, some of the findings in the literature appear contradictory which is perhaps due  
119 to the differences in levels, compounds, and test strategies applied in the sensory test.  
120 The early study by Woskow (1969), investigated the effects of umami on other tastes,  
121 but not vice versa. The study used a series of 50:50 combination of disodium 5'-  
122 inosinate and disodium 5'-guanylate from low to moderate levels (0.1mM to 0.5mM),  
123 while MSG was not included. This umami combination was found to enhance  
124 sweetness and saltiness but suppress sourness and bitterness. Reporting on work from  
125 their laboratory in 1979, Yamaguchi (1998) noted that MSG slightly enhanced saltiness  
126 from NaCl, but only at high MSG concentrations, and found that NaCl had no  
127 substantial influence on the perception of umami, while all other tastes did suppress  
128 umami. Kemp and Beauchamp (1994) demonstrated that at threshold levels, MSG had  
129 no influence on sweet, salt, sour and bitter, while at supra-threshold concentrations it  
130 suppressed sweet and bitter tastes and enhanced salt perception.

131 The findings of Kemp and Beauchamp (1994) for bitterness suppression corroborates  
132 the work of Woskow (1969), which is perhaps unsurprising as the levels of bitter  
133 tastant, quinine sulfate, were relatively similar (0.007 and 0.025 mM respectively) in  
134 the two studies and the perceived intensity of MSG at the medium level was similar to  
135 the recorded umami intensity of the two ribonucleotides in the earlier study. However,  
136 for saltiness, Woskow (1969) concluded that ribonucleotides enhanced salty taste at  
137 moderate concentration ( $\geq 0.2\text{mM}$ ), whereas Kemp and Beauchamp (1994) reported  
138 the enhancement of umami taste on salty taste only happened at high concentration of  
139 MSG (0.032mM and 0.059mM), as also concluded by Yamaguchi (1998). In relation  
140 to sweet taste, the conflicting result is likely to be due to the difference in sucrose levels  
141 used between the two studies. Sweetness was enhanced when the sucrose levels was  
142 5% (w/v) or 0.16 M (Woskow, 1969), whereas it was suppressed when the level was  
143 three times lower at 0.05 M (Kemp & Beauchamp, 1994).

144 Bitterness suppression was later confirmed by Keast and Breslin (2002b), concluding  
145 that when using either MSG or adenosine monophosphate sodium salt (NaAMP), the  
146 bitter taste of any of five different bitter tastants was suppressed. However, according  
147 to the research by Fuke and Ueda (1996), NaAMP does not evoke umami taste alone,  
148 hence, inferring that taste suppression may not require the suppressing tastant to be  
149 perceived. Bitter and umami tastes are mediated via G-protein-coupled receptors, T1Rs  
150 and T2Rs which are found in type II taste receptor cells (Bachmanov & Beauchamp,  
151 2007). Kim *et al.* (2015) established that the suppression of bitter taste by umami could  
152 occur at a cellular level, by investigating umami-bitter taste interactions with a cell-  
153 based assay using hTAS2R16-expressing cells. They tested the effect of five umami  
154 peptides (Glu-Asp, Glu-Glu, Glu-Ser, Asp-Glu-Ser, and Glu-Gly-Ser) on the bitter  
155 tastant salicin and found that the glutamyl peptides inhibited the salicin-induced  
156 intracellular  $\text{Ca}^{2+}$  response. Specifically, the Glu-Glu peptide suppressed salicin-  
157 induced activation of hTAS2R16 to a greater extent compared with the probenecid, a  
158 specific antagonist of hTAS2R16.



159 Previous studies have considered taste-taste interactions within ternary and quaternary  
160 mixtures (Bartoshuk, 1975; Breslin & Beauchamp, 1997; Green *et al.*, 2010). Breslin  
161 and Beauchamp (1997) investigated the interaction between sweet, salt and bitter, and  
162 found that bitter (urea) and sweet (sucrose) suppressed each other when mixed together.  
163 However, when salt (sodium acetate) was added the bitterness substantially decreased  
164 and the sweetness increased. While these papers focused on complex tastant mixtures,  
165 umami tastants were not included, and there are few studies exploring the specific  
166 interaction between umami and saltiness along with other basic tastes i.e., sweet, bitter  
167 and sour. Therefore, in order to study the effect of umami on the perception of other  
168 taste stimuli and vice versa, aqueous model systems were established to assess the taste  
169 perception using equi-intense taste mixture combinations, where the intensity levels are  
170 realistic to levels typically present in foods. Progressing understanding from previous  
171 literature this study specifically hypothesised that at moderate levels of umami  
172 sensation, saltiness would not be enhanced but neither would saltiness suppress umami,  
173 anticipating therefore by the summation model that the overall taste perception of a  
174 savoury system would be increased.

175

## 176 **2. Materials and Methods**

### 177 2.1 Panelists

178 A total of 12 trained sensory panelists (11 females and 1 male, age 35 to 65) participated  
179 in all experiments. They were also screened for their detection, discrimination and  
180 description ability. All panelists were healthy and had no taste or olfactory defects or  
181 disorders. They were all employed as sensory panelists and provided consent through  
182 their employment to taste foods and for their data to be used.

### 183 2.2 Stimulus

184 The taste stimuli used (indicated in Table 1) were aqueous solutions of sucrose  
185 (granulated sugar, Co-op Food, Manchester, UK) for the taste quality sweet (S), sodium

186 chloride (table salt, Co-op Food, Manchester, UK) for salty (N), citric acid (Sigma-  
187 Aldrich, Gillingham, UK) for sour (C), quinine hemisulfate salt monohydrate (Sigma-  
188 Aldrich, Gillingham, UK) for bitter (Q), monosodium glutamate MSG and  
189 monopotassium L-glutamate monohydrate (MPG) (Ajinomoto, Paris, France) for the  
190 taste quality umami (U). Each tastant solution was prepared in mineral water (Harrogate  
191 Spa, UK) a day before the panel session and kept in the fridge (4 °C) overnight. All  
192 tastant solutions were taken out of the fridge prior to the test to equilibrate to ambient  
193 temperature, then 15 mL of the sample was poured into 20 mL transparent polystyrene  
194 cups labeled with three-digit random codes and were served to the panel at ambient  
195 temperature (22 ± 2 °C).

### 196 2.3 Training

197 Prior to the data collection, all panelists participated in training on the use of the general  
198 labelled magnitude scale (gLMS). Compared to labelled magnitude scale (LMS) **first**  
199 **developed by Green, Shafer, and Gilmore, (1993)**, the top of gLMS is defined as  
200 “strongest imaginable of any sensation”, which is more suitable for this experiment  
201 where intensity across modalities is compared (Bartoshuk *et al.*, 2004). The descriptors  
202 of the magnitude estimates were “barely detectable”, “weak”, “moderate”, “strong”,  
203 “very strong” and “strongest imaginable of any sensation” (anchored values on gLMS  
204 scale 0.14, 0.76, 1.12, 1.52, 1.70, 1.98; exponentiated values 1.38, 5.01, 15.9, 31.6, 50.1  
205 and 95 respectively) (Bartoshuk *et al.*, 2004).

206 During the training period, panelists were asked to rate the taste intensity of the five  
207 basic taste stimuli respectively. The concentration of each stimulus used in this  
208 experiment was finalized when each stimulus was perceived as equi-intense (within the  
209 range from ‘strong’ to ‘very strong’ sensation on gLMS) by the panel. The training for  
210 finalizing the choice of concentration for stimuli was completed in three days.

### 211 2.4 Tastants preparation

212 Each of the three experiments described below in detail, contained a total of 10 tastants,  
213 including five single tastant solutions and five tastant mixtures (four binary, one  
214 quinary). All 12 panelists took part in all three experiments. After the training session,  
215 the first set of solutions (Experiment 1) using MSG as the source of umami with sodium  
216 unbalanced (UB) was scored by the panel, which were followed by solutions using  
217 MSG as the source for umami with sodium balanced (B) (Experiment 2). Finally, the  
218 panel was required to taste the third set of solutions (Experiment 3) which were  
219 prepared using MPG as the source for umami. For the three experiments, scoring for  
220 the samples were completed within two days.

#### 221 2.4.1 Experiment 1: MSG as the source of umami with sodium unbalanced (UB)

222 Based on the training results to determine equi-intensity, the single stimulus was  
223 selected at concentrations with the mean panel scores being between strong and very  
224 strong on the gLMS. The concentration of each tastant was kept constant in each binary  
225 and quinary tastant mixture as seen in Table 1.

#### 226 2.4.2 Experiment 2: MSG as the source for umami with sodium balanced (B)

227 NaCl contains 39.34% (w/w) sodium whereas MSG contains 13.6% (w/w) sodium.  
228 Therefore, an experiment was designed to ensure that sodium levels were controlled so  
229 that a raised sodium level in samples were perceived as salty taste. To achieve this,  
230 0.015 M NaCl was added to all single tastants except MSG (Table 1). Based on the  
231 training results to determine equi-intensity, the single stimulus was selected at  
232 concentrations with the mean panel scores being between strong and very strong on the  
233 gLMS. The concentration of each tastant was kept constant in each binary and quinary  
234 tastant mixture as seen in Table 1.

#### 235 2.4.3 Experiment 3: MPG as the source for umami

236 In order to remove the possible influence of sodium in glutamate when evaluating  
237 saltiness, the source for the taste quality of umami was changed to MPG. The  
238 concentration of each tastant was also adjusted to achieve a slightly lower equi-intensity

**Table 1: Concentration of tastants used in binary and quinary mixture sets**

Sample*	Experiment 1: Concentration used in MSG (sodium unbalanced) set MSG (UB)	Experiment 2: Concentration used in MSG (sodium balanced) set MSG (B)	Experiment 3: Concentration used in MPG set
S	S 0.19 M	S 0.19 M + N 0.015M	S 0.10 M
N	N 0.08 M	N 0.08 M + N 0.015M	N 0.05 M
C	C 0.005 M	C 0.005 M + N 0.015M	C 0.004 M
Q	Q 0.025 mM	Q 0.025mM + N 0.015M	Q 0.02 mM
U	U 0.015 M	U 0.015M	U 0.01 M
U+S	S 0.19M, U 0.015M	S 0.19M, U 0.015M	S 0.10M, U 0.01M
U+N	N 0.08M, U 0.015M	N 0.08M, U 0.015M	N 0.05M, U 0.01M
U+C	C 0.005 M, U 0.015M	C 0.005 M, U 0.015M	C 0.004 M, U 0.01M
U+Q	Q 0.025mM, U 0.015M	Q 0.025mM, U 0.015M	Q 0.02mM, U 0.01M
U+S+N+C+Q	S 0.19M, N 0.08M, C 0.005 M, Q 0.025mM, U 0.015M	S 0.19M, N 0.08M, C 0.005 M, Q 0.025mM, U 0.015M	S 0.10M, N 0.05M, C 0.004 M, Q 0.02mM, U 0.01M

240 \*S = sucrose; N = sodium chloride; C = citric acid; Q = quinine hemisulfate salt monohydrate; U = monosodium glutamate (MSG) or potassium  
 241 L-glutamate monohydrate (MPG)

243 on the gLMS between the descriptors moderate and strong, which allows a linear  
244 relationship between stimuli and response on the psychophysical curve as the one  
245 achieved in experiments 1 and 2 (Table 1).

## 246 2.5 Sensory evaluation

247 The experiments were conducted within a standard sensory environment using  
248 individual sensory booths, artificial daylight and controlled room temperature ( $22 \pm$   
249  $2$  °C). All samples were blind-coded and presented monadically. During tasting  
250 sessions, panelists were instructed to sip and hold the stimulus in their mouths for five  
251 seconds before swallowing and rating six attributes for each sample as follows: overall  
252 taste intensity, sweet, salty, sour, bitter and umami intensity. Between samples, the  
253 panel was instructed to cleanse their palate with plain crackers and water (filtered tap  
254 water at room temperature) to return the mouth back to a neutral state; an automatic  
255 reminder appeared during the countdown of ninety seconds between each stimulus after  
256 evaluating consecutive taste samples. Within each experiment scoring sessions  
257 included 10 samples and 2 replicates scored across two days. Sample presentation order  
258 was balanced across panelists; they each received different sample orders between each  
259 other, between replicates and between experiments. Data were captured using the  
260 sensory software Compusense® (cloud version, Guelph, Ontario).

## 261 2.6 Data analysis

262 Data from each of the three experiments was analysed separately. Log data from each  
263 panelist from the gLMS were captured by Compusense®. Data were exponentiated.  
264 Two-way analysis of variance (ANOVA) was carried out using Senpaq (QI Statistics,  
265 Reading, UK) where panelists were treated as random effects and samples as fixed  
266 effects, main effects were tested against the assessor by sample interaction. Multiple  
267 pairwise comparisons were carried out using Tukey's HSD at a significance level of  
268 0.05.

269

### 270 3. Results

271 The mean scores of perceived taste intensity for all single tastes and taste mixtures are  
272 given in Figures 1 to 3 (further statistical details given in supplementary Table 1 to 3).  
273 The aim was to have all single tastants rated “strong to very strong” on the gLMS (1.52  
274 to 1.70 on the log scale, or 31.6 to 50.1 exponentiated values) in both the sodium  
275 unbalanced and balanced sets. Although panelists were extensively trained on each  
276 single tastant, saltiness and sourness were rated slightly lower than “strong”. However,  
277 the mean ratings (exponentiated data) only fell below this descriptor by a maximum of  
278 0.4 units, therefore it is suggested that this would not have greatly influenced the results.  
279 For samples using MPG as source of umami taste, all single tastants were rated as  
280 “moderate to strong” on the gLMS (1.21 to 1.52 on the log scale, or 15.85 to 31.62 as  
281 exponentiated values), while the concentration of tastants used was slightly lower in  
282 comparison to the MSG set samples.

#### 283 3.1 Intensity of umami

284 The ratings of perceived intensity of umami in the different experiments are presented  
285 in Figure 1. It is clear from this figure that the perception of umami was significantly  
286 suppressed by all other tastes in both the binary and quinary mixtures. In all experiment  
287 sets, all the taste mixtures containing MSG were significantly ( $p < 0.05$ ) lower in  
288 perceived umami intensity compared to MSG alone (U). The umami sensation was  
289 reduced from just above “strong” to “moderate” or “weak” in virtually all cases. The  
290 main exceptions were where the binary mixture was with sodium chloride (U+ N), this  
291 led to a lower reduction in umami, leading to “moderate” sensation rather than “weak”.  
292 The intensity of umami in the quinary taste systems (U+S+N+C+Q) was the lowest for  
293 all experiment sets.

#### 294 3.2 Intensity of other tastes

295 The ratings of perceived intensity of sweetness, saltiness, sourness and bitterness can  
296 be seen in Figure 2. The umami taste did not enhance or suppress the perceived intensity

297 of any other taste in the binary taste systems ( $p > 0.05$ ) (further statistical details given  
298 in supplementary Table 1 to 3). This is an unusual phenomenon as all other taste  
299 modalities will suppress each other when added together (Green *et al.*, 2010), and yet  
300 the addition of MSG as an umami tastant has neither suppressed, nor enhanced,  
301 perception of the other four tastes. Kemp and Beauchamp (1994) concluded that MSG  
302 at medium concentration (0.032M) suppressed sweet and bitter tastes and at higher  
303 concentrations (0.059M) enhanced salty taste. The MSG levels used by Kemp and  
304 Beauchamp (1994) are higher than the 0.015M used in the current study which may  
305 have partly led to the different findings. However, the main reason is likely to be the  
306 different concentration of the other tastants. The present study used 0.19 M sucrose and  
307 0.005 M citric acid for equi-intense perception of “strong to very strong”.

### 308 3.3 Overall taste intensity

309 The ratings of perceived intensity of overall taste in the different experiments are  
310 presented in Figure 3. Results indicated that the total taste intensity of binary mixtures  
311 was very similar to the total overall taste intensity of single tastants ( $p > 0.05$ ), except  
312 for quinine hemisulfate with umami mixture (U+Q) in the sodium balanced set and  
313 sodium chloride with umami mixture (U+N) in MPG set, where the binary mixture was  
314 significantly higher in overall taste intensity ( $P < 0.05$ ). The total taste intensity of the  
315 quinary solution had a higher mean rating than all binary mixtures. In particular, it had  
316 a significantly higher rating compared to the binary mixture with citric acid (U+C) in  
317 both MSG sessions, and the binary mixture with sodium chloride (U+N) in sodium  
318 balanced set and MPG set ( $p < 0.05$ ). The perception of all five tastes were all  
319 significantly and substantially lower in the quinary mixtures than as single tastants ( $p$   
320  $< 0.05$ ) in the sodium balanced set and MPG set. In the sodium unbalanced set, sour,  
321 bitter and umami tastes were similarly significantly lower in the quinary mixtures than  
322 as single tastants ( $p < 0.05$ ).

323 The binary mixture with quinine hemisulfate (U+Q) had a significantly higher overall  
324 taste intensity than the sample of quinine hemisulfate alone (Q) only in sodium balanced

325 set ( $p < 0.05$ ), but not in sodium unbalanced set and MPG set. This could possibly be  
326 due to the inclusion of 0.015mM NaCl in quinine solution in the sodium balanced set.  
327 Keast and Breslin (2002a) reported that NaCl has suppression effect on the bitterness  
328 perception at low, medium and high intensity level. Therefore, 0.015M salt addition  
329 would lead to a lower intensity of bitterness for quinine solution in sodium balanced set  
330 (Experiment 2), while it is not the case in sodium unbalanced set (Experiment 1) and  
331 MPG set (Experiment 3). As the total overall intensity is determined by the dominant  
332 taste (bitterness), as a result, a low overall taste intensity in quinine hemisulfate alone  
333 solution (Q) was expected compared with that in quinine hemisulfate with umami  
334 mixture (U+Q) in sodium balanced set. The binary mixture of MPG and NaCl (U+N)  
335 had a significantly higher overall taste intensity than the sample of NaCl (N) alone ( $p$   
336  $< 0.05$ ). This indicates that umami may enhance the total intensity of a salt solution  
337 without enhancing the specific taste modality (saltiness) in the MPG mixture. The  
338 binary mixtures of U+N in the MSG sample set had a similar trend, but the differences  
339 were not significant ( $p > 0.05$ ). These differences may be associated with the difference  
340 in concentrations used in the MSG and MPG sets (0.08M or 0.095M vs 0.05M). Finally,  
341 the total taste intensity of the quinary solution was the strongest, with all single tastants  
342 having a significantly and substantially lower overall taste intensity than the quinary  
343 mixtures except quinine hemisulfate ( $p < 0.05$ ).

#### 344 3.4 Taste interaction

345 The testing of the balanced sodium sample set allowed for an unbiased investigation of  
346 the influence of glutamate and the perception of all other tastes, and of the effect of  
347 sodium on glutamate, without the sodium within the MSG as a confounding factor. In  
348 conclusion, the results from both the sodium unbalanced and balanced trials were the  
349 same, increasing the confidence in the overall finding that umami from glutamate does  
350 not enhance or suppress other tastes when all tastes are presented at strong (but not  
351 excessive) intensity levels. The findings in this MPG set again confirmed that all other  
352 tastes suppressed umami ( $p < 0.05$ ), whereby all binary mixtures had significantly lower  
353 umami intensity than MPG alone ( $p < 0.05$ ), and the quinary mixture was significantly



354 and very substantially lower in umami taste ( $p < 0.05$ ). The results agree with the first  
355 two studies that the umami taste did not enhance or suppress the perceived intensity of  
356 any other taste in the binary taste systems ( $p > 0.05$ ), all other tastes could suppress the  
357 perception of umami taste in binary and quinary mixture ( $p < 0.05$ ).

#### 358 **4. Discussion**

359 Figure 4 summarizes the overall findings which were common to all three studies  
360 presented in this paper, illustrating the associations between umami and the other four  
361 basic tastes. As seen in this figure the addition of umami taste did not enhance or  
362 suppress any other taste, however, the addition of sweet, salty, sour and bitter do  
363 significantly suppress the umami taste.

364 Keast and Breslin (2002a) have shown that the concentration of taste stimuli, and the  
365 position on the concentration-intensity psychophysical curve could predict the  
366 interactions of tastes in taste mixtures. In the current study however, no matter whether  
367 it was in the “moderate” perceived intensity region or in “strong” perceived intensity  
368 region, the umami taste did not enhance or suppress the perceived intensity of any other  
369 taste in the binary taste systems; where sweet, salty, sour and bitter all significantly  
370 suppressed the perception of umami intensity in the binary and quinary taste systems.  
371 Previous research has tended to agree that umami enhances salt perception in aqueous  
372 solutions (Woskow, 1969; Kemp & Beauchamp, 1994) and in foods (Dermiki *et al.*,  
373 2013; Kremer *et al.*, 2013; Khetra *et al.*, 2019), and in recent years food manufacturers  
374 have been keen to use umami to enhance salty taste. However, the experimental results  
375 from this study conclude that umami taste did not affect the salty taste when presented  
376 at moderate or strong equi-intensities.

377 The disagreement between the current study and previous findings may be explained  
378 by the following factors: First, the levels of tastants used varies between studies.  
379 Compared to studies that previously used MSG, the 0.015M used in this study was  
380 lower than the levels found in the Kemp and Beauchamp study (1994) to enhance salty

381 taste (0.032 and 0.059M MSG), and the level of sodium chloride used in the previous  
382 study was much lower (0.025M compared to 0.08M in the present study).

383 **In addition, test procedure differences, i.e. a taste and spit procedure vs a taste and**  
384 **swallow procedure, are also** responsible for the conflict. Running and Hayes (2017)  
385 have previously concluded that taste ratings resulting from model solutions that had  
386 been spat out are lower than ratings for swallowed samples on a gLMS scale. Taken  
387 together these arguments might infer that umami may enhance salty perception where  
388 salty taste is lower. Kawasaki *et al.* (2016) **give** an insight into the time over which the  
389 different tastes are perceived, for example saltiness and sourness tend to be perceived  
390 as dominant before swallowing, whereas umami was dominant after swallowing. This  
391 finding highlights the effect of the test methodology on the perceived intensity of taste.  
392 The sip and spit method was used by Kemp and Beauchamp (1994), while Keast and  
393 Breslin (2002b) did not include swallowing. But solutions were swallowed in the  
394 present study. Therefore, it is difficult to compare the results of studies where the tests  
395 were not conducted in the same way. Kawasaki *et al.* (2016) also investigated the  
396 duration of impact of taste attributes of umami (MSG), salty (sodium chloride), sour  
397 (lactic acid) and their binary mixtures using temporal dominance of sensations  
398 methodology. They found that the presence of MSG increased the duration of NaCl  
399 saltiness but suppressed the sourness of lactic acid. On the other hand, the duration of  
400 umami taste of MSG was suppressed in the presence of NaCl but was not affected by  
401 lactic acid. This means that MSG could increase the duration of salty taste from NaCl  
402 rather than enhance the peak intensity. This might imply that where previous studies  
403 have reported an enhancement of salty taste, it could have been that the taste duration  
404 was extended rather than an increase in maximum intensity. However, our study was  
405 specifically set up to test maximum intensity following the sample remaining in the  
406 mouth for 5s, and so would not have captured an increase in duration that the Kawasaki  
407 study concluded.

408 A second explanation for such discrepancies might be that umami is a less recognised  
409 taste in Western countries and consumers may perhaps confuse it with salty perception,

410 despite it being one of the five basic tastes (Cecchini *et al.*, 2019). Although the  
411 panelists in this study were trained to recognise and score umami taste, they were UK  
412 assessors and as such they would not be habituated to umami taste throughout their  
413 lives, which might have affected their scoring. Certainly, in previous studies where  
414 functional magnetic resonance imaging (fMRI) was employed, it was confirmed that  
415 there was only a slight difference between the positions of the activation regions  
416 between umami and salty taste, which led to the conclusion that the basic perception  
417 system of umami taste was very similar to the basic perception system of salty taste  
418 (Nakamura *et al.*, 2011). Furthermore, Onuma, Maruyama, and Sakai (2018) had  
419 reported that the NaCl solutions with MSG increased responses in the frontal operculum  
420 but did not affect the hemodynamic salivary by functional near-infrared spectroscopy  
421 (fNIRS) data. This means that the umami induced saltiness enhancement effects occur  
422 in the central gustatory processing in the brain. Additionally, this might partly explain  
423 why umami, in the MPG model, was found to enhance the total taste intensity of the  
424 salt solution, without enhancing the specific taste modality (saltiness).

425 The type of panelist used in different studies should also be considered. Trained sensory  
426 panelists, such as the assessors in this study, “dissect” a product into its component  
427 attributes for rating, whereas consumers “synthesise” the information from the foods  
428 they are tasting (Ares & Varela, 2017). Compared with untrained consumers, trained  
429 panelists are more sensitive to taste discrimination, and they are significantly more  
430 aware of the flavour in the mixture and the intensity of suppression (McBride & Finlay,  
431 1989; Prescott, Ripandelli & Wakeling, 2001), although their hedonic perception of the  
432 product may not fully represent the wide and varied perceptions from untrained  
433 consumers (Ares & Varela, 2017). So, one might expect a consumer would synthesise  
434 congruent taste information in a way that a trained panelist might not, leading more  
435 readily to the conclusion that a salt reduced food that is higher in umami might have an  
436 overall similar salty perception as the two tastes are congruent. However, the previous  
437 studies which concluded that umami enhanced salty taste perception were all carried  
438 out with trained panelists (Woskow, 1969; Kemp & Beauchamp, 1994; Keast &

439 Breslin, 2002b), as employed in the current study; so, the differences in perception  
440 between trained panellists and consumers, does not lead to a satisfactory explanation of  
441 conflicting results.

442 When Green *et al.* (2010) studied binary, ternary and quaternary mixtures, they found  
443 that the overall perceived intensity of the mixtures was best predicted by perceptual  
444 additivity, the sum of the tastes perceived within the mixture (Green *et al.*, 2010). In  
445 fact, their study concluded the sum of the unmixed taste intensities to be much higher  
446 than the sum of the taste intensities in the mixture, or the overall taste intensity ratings,  
447 thus ruling out stimulus additivity (Keast & Breslin, 2002a). In the current study, it was  
448 consistent that the overall taste intensity was lower than both the sum of the unmixed  
449 taste intensities and the sum of the taste intensities in binary system and quinary mixture.  
450 However, it was relatively easy to distinguish each taste in the binary system but much  
451 more difficult to distinguish each taste in the quinary mixture system, which may lead  
452 to a great reduction in intensity compared to a single tastant.

453 One limitation of this work was that when the source of umami was changed from MSG  
454 to MPG, the concentration level did not remain in the same taste intensity level. It  
455 means the relationship between the five basic tastes is only valid at certain taste  
456 intensity level and for certain umami compound, i.e. from moderate to strong when  
457 MPG was used as the source of umami; from strong to very strong when MSG was used  
458 as the source of umami. Even if the results presented same trend (suppression), the  
459 impact of concentration range on perception was uncertain. However, it provides a  
460 prediction for the relationship of the five basic tastes when MSG is used as the source  
461 of umami at other concentration levels in the future.

462 In fact, taste interactions in a real food matrix are more complicated compared to  
463 aqueous solutions. This can explain why for example, MSG is added in variety of food  
464 products (e.g., soup, potato chips, sausage) to replace NaCl as well as to enhance  
465 flavour (Yamaguchi & Takahashi, 1984; Dos *et al.*, 2014; Maluly *et al.*, 2017).  
466 However, increasing saltiness perception using MSG in the aqueous model system of

467 the current study was not observed. The discrepancy could be explained due to the  
468 complexity of food matrices which affects the perception. In a real food there are cross-  
469 modal interactions between two or more sensory modalities such as taste-flavour or  
470 flavour-texture interactions. Additionally, ingredients used in food products are often  
471 added at much higher concentrations than in the aqueous model systems to achieve the  
472 required taste intensity, considering that the texture can reduce intensity. In general,  
473 meat products have a high sodium content, and the salt content is around 2% (Inguglia  
474 *et al.*, 2017), where only 0.29% or 0.55% salt was used in this study. Other research  
475 used higher MSG levels, 0.38% MSG was added to the sumashi-jiru (soup) to maintain  
476 the salty taste, and 0.3% MSG added to the sausage to compensate the saltiness loss  
477 caused by 50% salt reduction in low-sodium fish burgers (Quadros *et al.*, 2015). In  
478 contrast, only 0.19% or 0.25% MSG was used in this study. Therefore, the conclusions  
479 reached by investigating aqueous model solution may not be applicable to food systems  
480 directly, however they offer the basis for the design of further experiments in real foods.

481 The present study employed a trained sensory panel to investigate taste interactions,  
482 with limited variability in taste sensitivities. Prescott *et al.* (2001) concluded that  
483 perception of tastes and interaction between tastes in binary mixture are affected by the  
484 6-n-propylthiouracil (PROP) taster status, i.e. supertaster, medium taster and non-taster.  
485 However, the taste sensitivity is determined by many factors, such as genetic  
486 differences in taste receptors, including Tas2R38 gene that is predominantly  
487 responsible for PROP/PTC (phenylthiocarbamide) tasting (Hayes *et al.*, 2008), and  
488 single nucleotide polymorphisms (SNPs) for epithelial sodium channel (ENaC)  
489 (Chamoun *et al.*, 2021). For example, SNPs for the T1R receptors influence perception  
490 of sweet and umami taste. Therefore, to truly understand the influence of umami taste  
491 in taste mixtures for all consumers, a study considering taste sensitivities to basics tastes  
492 (each from more than one tastant) alongside genotyping would be needed in a large  
493 population cohort in the future.

494

495 **5. Conclusions**

496 The aim of this study was to investigate taste interactions in mixtures containing umami  
497 in the form of MSG and MPG. The result shows that the addition of umami taste did  
498 not enhance or suppress any other taste. Therefore, umami is dissimilar to other tastants  
499 which tend to suppress one another. However, the addition of sweet, salty, sour and  
500 bitter do significantly suppress the umami taste. The findings of this study are  
501 significant because they fill the gap that existed in the literature considering the effect  
502 of umami taste in taste mixture interactions and have an impact on our understanding  
503 of the underlying mechanisms of taste interactions that can be applied in food  
504 reformulation. Although umami was not found to enhance salty perception, as  
505 hypothesised, neither did it suppress it; hence when used together sodium chloride plus  
506 glutamate tastants maintained salty perception in addition to savoury taste perception,  
507 irrespective of the glutamate salt used. Overall, there is little evidence on the effect of  
508 umami on other taste stimuli, and the findings of the current study are difficult to  
509 compare directly with the limited information currently available in the literature. The  
510 reasons for this are the different sensory tests used (ranking vs gLMS), the different  
511 methodology (sip and spit vs swallowing), the different concentrations of tastants and  
512 the difference in perception of similar concentrations by the different groups studied.  
513 Further investigation is needed to determine whether these findings in aqueous  
514 solutions apply to real food systems where more complex and cross-modal interactions  
515 take place.

516

517 **Acknowledgement**

518 The sensory panelists are thanked for attending the sensory evaluation sessions  
519 Compusense are thanked for their provision of Compusense cloud software under their  
520 academic consortium agreement.

521

522 **References**

- 523 Ares, G. & Varela, P. (2017). Trained vs. consumer panels for analytical testing:  
524 Fueling a long lasting debate in the field. *Food Quality and Preference*, 61, 79-86.
- 525 Abu, N.B., Harries, D., Voet, H. and Niv, M.Y., (2018). The taste of KCl—What a  
526 difference a sugar makes. *Food chemistry*, 255, 165-173.
- 527 Bachmanov, A.A & Beauchamp, G.K. (2007). Taste Receptor Genes. *Annual Review*  
528 *of Nutrition*, 27, 389–414.
- 529 Bartoshuk, L. M. (1975). Taste mixtures: Is mixture suppression related to  
530 compression? *Physiology and Behavior*, 14(5), 643-649.
- 531 Bartoshuk, L.M., Duffy, V.B., Green, B.G., Hoffman, H.J., Ko, C.W., Lucchina, L.A.,  
532 Marks, L.E., Snyder, D.J. and Weiffenbach, J.M. (2004). Valid across-group  
533 comparisons with labeled scales: the gLMS versus magnitude matching. *Physiology &*  
534 *behavior*, 82(1), 109-114.
- 535 Breslin, P. A. S. & Beauchamp, G. K. (1997). Salt enhances flavour by suppressing  
536 bitterness. *Nature*, 387, 563-563.
- 537 Cecchini, M., Knaapila, A., Hoffmann, E., Boschi, F., Hummel, T. & Iannilli, E. (2019).  
538 A cross-cultural survey of umami familiarity in European countries. *Food Quality and*  
539 *Preference*, 74, 172-178.
- 540 Chamoun, E., Liu A.S. Duizer, L.M., Feng, Z., Darlington, G., Duncan, A.M., Haines,  
541 J., Ma, D.W.L. (2021). Single nucleotide polymorphisms in sweet, fat, umami, salt,  
542 bitter and sour taste receptor genes are associated with gustatory function and taste  
543 preferences in young adults. *Nutrition Research*, 85, 40-46.
- 544 Dermiki, M., Mounayar, R., Suwankanit, C., Scott, J., Kennedy, O., Mottram, D.,  
545 Gosney, M., Blumenthal, H. and Methven, L. (2013). Maximising umami taste in meat  
546 using natural ingredients: effects on chemistry, sensory perception and hedonic liking  
547 in young and old consumers. *Journal of the Science of Food and Agriculture*, 93(13),  
548 3312-3321.
- 549 Dos Santos, B., Campagnol, P., Morgano, M. & Pollonio, M. (2014). Monosodium  
550 glutamate, disodium inosinate, disodium guanylate, lysine and taurine improve the

551 sensory quality of fermented cooked sausages with 50% and 75% replacement of NaCl  
552 with KCl. *Meat Science*, 96(1), 509-513.

553 Emorine, M., Septier, C., Martin, C., Cordelle, S., Sémon, E., Thomas-Danguin, T. and  
554 Salles, C. (2021). Salt and aroma compound distributions influence flavour release and  
555 temporal perception while eating hot-served flans. *Molecules*, 26(5), 1300-1317.

556 Fuke, S. and Ueda, Y. (1996). Interactions between umami and other flavor  
557 characteristics. *Trends in Food Science & Technology*, 7(12), 407-411.

558 **Green, B.G., Shaffer, G.S. and Gilmore, M.M. (1993). Derivation and evaluation of a**  
559 **semantic scale of oral sensation magnitude with apparent ratio properties. *Chemical***  
560 ***senses*, 18(6), 683-702.**

561 Green, B. G., Lim, J., Osterhoff, F., Blacher, K., & Nachtigal, D. (2010). Taste mixture  
562 interactions: Suppression, additivity, and the predominance of sweetness. *Physiology*  
563 *& Behavior*, 101, 731-737.

564 Hayes, J.E., Bartoshuk, L.M., Kidd, J.R., Duffy, V.B. (2008). Supertasting and PROP  
565 bitterness depends on more than the TAS2R38 gene. *Chemical Senses*, 33 (3), 255-265.

566 Inguglia, E.S., Zhang, Z., Tiwari, B.K., Kerry, J.P. and Burgess, C.M., (2017). Salt  
567 reduction strategies in processed meat products—A review. *Trends in Food Science &*  
568 *Technology*, 59, 70-78.

569 Kawasaki, H., Sekizaki, Y., Hirota, M., Sekine-Hayakawa, Y. & Nonaka, M. (2016).  
570 Analysis of binary taste-taste interactions of MSG, lactic acid, and NaCl by temporal  
571 dominance of sensations. *Food Quality and Preference*, 52, 1-10.

572 Keast, R. S. J. & Breslin, P. A. S. (2002a). An overview of binary taste-taste  
573 interactions. *Food Quality and Preference*, 14, 111-124.

574 Keast, R. S. J. & Breslin, P. A. S. (2002b). Modifying the bitterness of selected oral  
575 pharmaceuticals with cation and anion series of salts. *Pharmaceutical Research*, 19,  
576 1019-1026.

577 Kemp, S. E. & Beauchamp, G. K. (1994). Flavor Modification by Sodium Chloride and  
578 Monosodium Glutamate. *Journal of Food Science*, 59, 682-686.



579 Khetra, Y., Kanawjia, S., Puri, R., Kumar, R. & Meena, G. (2019). Using taste-induced  
580 saltiness enhancement for reducing sodium in Cheddar cheese: Effect on physico-  
581 chemical and sensorial attributes. *International Dairy Journal*, 91, 165-171.

582 Kim, M., Son, H., Kim, Y., Misaka, T. & Rhyu, M. (2015). Umami–bitter interactions:  
583 The suppression of bitterness by umami peptides via human bitter taste receptor.  
584 *Biochemical and Biophysical Research Communications*, 456(2), 586-590.

585 Kremer, S., Shimojo, R., Holthuysen, N., Köster, E. & Mojet, J. (2013). Consumer  
586 acceptance of salt-reduced “soy sauce” foods over rapidly repeated exposure. *Food*  
587 *Quality and Preference*, 27(2), 179-190.

588 Maluly, H.D., Ariseto - Bragotto, A.P. and Reyes, F.G., (2017). Monosodium  
589 glutamate as a tool to reduce sodium in foodstuffs: Technological and safety aspects.  
590 *Food science & nutrition*, 5(6), 1039-1048.

591 McBride, R.L., (1987). Taste psychophysics and the Beidler equation. *Chemical*  
592 *Senses*, 12(2), 323-332.

593 McBride, R.L. and Finlay, D.C., (1989). Perception of taste mixtures by experienced  
594 and novice assessors 1. *Journal of Sensory Studies*, 3(4), 237-248.

595 McBride, R.L. and Finlay, D.C., (1990). Perceptual integration of tertiary taste  
596 mixtures. *Perception & psychophysics*, 48(4), 326-330.

597 McBride, R.L., (1993). Integration psychophysics: The use of functional measurement  
598 in the study of mixtures. *Chemical senses*, 18(2), 83-92.

599 Mojet, J., Heidema, J. and Christ-Hazelhof, E., (2004). Effect of concentration on taste–  
600 taste interactions in foods for elderly and young subjects. *Chemical senses*, 29(8), 671-  
601 681.

602 Nakamura, Y., Goto, T., Tokumori, K., Yoshiura, T., Kobayashi, K., Nakamura, Y.,  
603 Honda, H., Ninomiya, Y. & Yoshiura, K. (2011). Localization of brain activation by  
604 umami taste in humans. *Brain Research*, 1406, 18-29.

605 Onuma, T., Maruyama, H. and Sakai, N. (2018). Enhancement of Saltiness Perception  
606 by Monosodium Glutamate Taste and Soy Sauce Odor: A Near-Infrared Spectroscopy  
607 Study. *Chemical Senses*, 43(3), 151-167.

608 Prescott, J., Ripandelli, N. and Wakeling, I., (2001). Binary taste mixture interactions  
609 in prop non-tasters, medium-tasters and super-tasters. *Chemical senses*, 26(8), 993-  
610 1003.

611 Ponzio, V., Pellegrini, M., Costelli, P., Vázquez-Araújo, L., Gayoso, L., D'Eusebio, C.,  
612 Ghigo, E. and Bo, S., (2021). Strategies for reducing salt and sugar intakes in  
613 individuals at increased cardiometabolic risk. *Nutrients*, 13(1), 279-296.

614 Quadros, D. A., Rocha, I. F. O., Ferreira, S. M. R., & Bolini, H. M. A. (2015). Low-  
615 sodium fish burgers: Sensory profile and drivers of liking. *LWT - Food Science and*  
616 *Technology*, 63, 236–242.

617 Running, C. & Hayes, J. (2017). Sip and spit or sip and swallow: Choice of method  
618 differentially alters taste intensity estimates across stimuli. *Physiology & Behavior*,  
619 181, 95-99.

620 Schifferstein, H.N. and Frijters, J.E., (1993). Perceptual integration in heterogeneous  
621 taste percepts. *Journal of Experimental Psychology: Human Perception and*  
622 *Performance*, 19(3), 661-675.

623 Thomas-Danguin, T., Guichard, E. and Salles, C., (2019). Cross-modal interactions as  
624 a strategy to enhance salty taste and to maintain liking of low-salt food: A review. *Food*  
625 *& function*, 10(9), 5269-5281.

626 Velázquez, A.L., Vidal, L., Varela, P. and Ares, G., (2020). Cross-modal interactions  
627 as a strategy for sugar reduction in products targeted at children: Case study with vanilla  
628 milk desserts. *Food Research International*, 130, 108920.


629 Woskow, M. H. (1969). Selectivity in flavor modification by 5'-ribonucleotides. *Food*  
630 *Technology*, 23, 32-37.

631 Yamaguchi, S. & Kimizuka, A. (1979). Psychometric studies on the taste of  
632 monosodium glutamate. *Advances in biochemistry and physiology*, 35-54.

633 Yamaguchi, S., & Takahashi, C. (1984). Interactions of monosodium glutamate and  
634 sodium chloride on saltiness and palatability of clear soup. *Journal of Food Science*,  
635 49(1), 82– 85.

636 Yamaguchi, S. (1998). Basic properties of umami and its effect on food flavour. *Food*  
637 *Reviews International*, 14(2&3), 139-176.

638



639


640 Figure 1. Ratings of perceived intensity (exponentiated values) of umami in the  
641 sodium unbalanced and balanced sets and using MPG as source of umami taste set. S  
642 = sucrose; N = sodium chloride; C = citric acid; Q = quinine hemisulfate salt  
643 monohydrate; U = monosodium glutamate (MSG) or potassium L-glutamate  
644 monohydrate (MPG). **Within each sample set bars that do not share a common letter**  
645 **denote samples that differed significantly ( $p < 0.05$ ).** Lower case letters used for  
646 Experiment 1:MSG without salt balanced, upper case letters used for Experiment 2:  
647 MGS with salt balanced, and Greek letters used for Experiment 3: MPG.

648

649

650

651



652

653 Figure 2. Ratings of perceived intensity (exponentiated values) of sweetness (a),  
 654 saltiness (b), sourness (c), and bitterness (d) in the sodium unbalanced and balanced  
 655 sets and using MPG as source of umami taste set. S = sucrose; N = sodium chloride;  
 656 C = citric acid; Q = quinine hemisulfate salt monohydrate; U = monosodium  
 657 glutamate (MSG) or potassium L-glutamate monohydrate (MPG). **Within each sample**  
 658 **set bars that do not share a common letter denote samples that differed significantly ( $p$**   
 659  **$< 0.05$ ).** Lower case letters use for Experiment 1:MSG without salt balanced, upper  
 660 case letters use for Experiment 2: MGS with salt balanced, and Greek letters use for  
 661 Experiment 3: MPG.

662

663

664


665

666

667

668

669



670

671 Figure 3. Ratings of perceived intensity (exponentiated values) of overall taste in the  
 672 sodium unbalanced and balanced sets and using MPG as source of umami taste set. S  
 673 = sucrose; N = sodium chloride; C = citric acid; Q = quinine hemisulfate salt  
 674 monohydrate; U = monosodium glutamate (MSG) or potassium L-glutamate  
 675 monohydrate (MPG). **Within each sample set bars that do not share a common letter**  
 676 **denote samples that differed significantly ( $p < 0.05$ ).** Lower case letters use for  
 677 Experiment 1:MSG without salt balanced, upper case letters use for Experiment 2:  
 678 MGS with salt balanced, and Greek letters use for Experiment 3: MPG.

679

680

681

682

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708


709 **Figure 4. Binary interactions of taste qualities at equi-intense concentrations.**

710 **Direction of the arrow indicates the significant influence of primary taste on umami**

711 **( $p < 0.05$ ). It may be noted that there is no effect of umami on the primary tastes**

712 **including sweet, salty, sour and bitter.**

713



714 **Supplementary Table 1 for Experiment 1.** Ratings and significance testing (ANOVA)  
 715 results of perceived intensity (antilogged values) of overall taste, sweet, salty, sour,  
 716 bitter and umami where MSG was used as the umami tastant without sodium balance.

<b>Perceived intensity</b> (mean of antilogged gLMS intensity ratings)						
Sample	Total intensity	Sweet	Salty	Sour	Bitter	Umami
S	36.2 <sup>cd</sup>	34.7 <sup>a</sup>	2.5 <sup>c</sup>	2.2 <sup>c</sup>	1.9 <sup>c</sup>	1.2 <sup>d</sup>
<b>U+S</b>	45.1 <sup>abc</sup>	39.9 <sup>a</sup>	6.3 <sup>c</sup>	1.6 <sup>c</sup>	1.9 <sup>c</sup>	13.5 <sup>c</sup>
N	37.9 <sup>cd</sup>	1.1 <sup>b</sup>	31.4 <sup>ab</sup>	1.1 <sup>c</sup>	4.1 <sup>c</sup>	2.8 <sup>d</sup>
<b>U+N</b>	44.6 <sup>abc</sup>	4.5 <sup>b</sup>	32.8 <sup>a</sup>	1.3 <sup>c</sup>	2.5 <sup>c</sup>	23.5 <sup>b</sup>
C	38.7 <sup>cd</sup>	1.4 <sup>b</sup>	3.6 <sup>c</sup>	31.4 <sup>a</sup>	9.3 <sup>c</sup>	1.0 <sup>d</sup>
<b>U+C</b>	41.3 <sup>bcd</sup>	2.2 <sup>b</sup>	5.0 <sup>c</sup>	29.8 <sup>a</sup>	8.3 <sup>c</sup>	18.5 <sup>bc</sup>
Q	49.6 <sup>ab</sup>	1.0 <sup>b</sup>	1.1 <sup>c</sup>	1.9 <sup>c</sup>	45.6 <sup>a</sup>	1.0 <sup>d</sup>
<b>U+Q</b>	49.2 <sup>ab</sup>	1.1 <sup>b</sup>	2.7 <sup>c</sup>	1.4 <sup>c</sup>	43.6 <sup>a</sup>	16.6 <sup>bc</sup>
U	33.4 <sup>d</sup>	1.4 <sup>b</sup>	5.6 <sup>c</sup>	1.1 <sup>c</sup>	1.5 <sup>c</sup>	32.2 <sup>a</sup>
<b>U+S+N+C+Q</b>	53.2 <sup>a</sup>	39.1 <sup>a</sup>	24.7 <sup>b</sup>	11.7 <sup>b</sup>	14.2 <sup>b</sup>	5.3 <sup>d</sup>
<i>df</i> of Sample	9	9	9	9	9	9
<i>df</i> of Interaction	72	72	72	72	72	72
F-value of Sample Effect	4.08	80.81	24.8	29.93	25.45	19.22
Sample significance (p)	0.0003	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

717 <sup>abcde</sup> **Values within a column which don't share a common superscript are significantly**  
 718 **different in means ratings of the perceived magnitude** from Tukey's HSD test at the  
 719 95% confidence interval. S = sucrose; N = sodium chloride; C = citric acid; Q = quinine  
 720 hemisulfate salt monohydrate; U = monosodium glutamate (MSG). *df* = degrees of  
 721 freedom of interaction, noting that the main effect of sample (F-value of sample) was  
 722 determined by dividing the variance of sample by the variance of the interaction  
 723 (MS<sub>sample</sub>/MS<sub>interaction</sub>) hence both the *df* of sample and interaction are given.  
 724

725 **Supplementary Table 2 for Experiment 2.** Ratings and significance testing (ANOVA)  
 726 results of perceived intensity (antilogged values) of overall taste, sweet, salty, sour,  
 727 bitter and umami where MSG was used as the umami tastant with sodium balance.

<b>Perceived intensity</b> (mean of antilogged gLMS intensity ratings)						
Sample	Total intensity	Sweet	Salty	Sour	Bitter	Umami
S	43.5 <sup>bcd</sup>	41.9 <sup>a</sup>	4.9 <sup>c</sup>	1.2 <sup>c</sup>	1.1 <sup>c</sup>	1.0 <sup>e</sup>
<b>U+S</b>	49.9 <sup>ab</sup>	44.4 <sup>a</sup>	5.3 <sup>c</sup>	2.2 <sup>c</sup>	1.6 <sup>c</sup>	14.4 <sup>c</sup>
N	41.0 <sup>cde</sup>	2.1 <sup>c</sup>	35.4 <sup>a</sup>	2.8 <sup>c</sup>	3.0 <sup>c</sup>	6.2 <sup>e</sup>
<b>U+N</b>	47.5 <sup>bc</sup>	2.4 <sup>c</sup>	30.7 <sup>a</sup>	2.8 <sup>c</sup>	3.0 <sup>c</sup>	22.4 <sup>b</sup>
C	37.9 <sup>de</sup>	2.0 <sup>c</sup>	5.0 <sup>c</sup>	31.2 <sup>a</sup>	6.0 <sup>c</sup>	1.3 <sup>e</sup>
<b>U+C</b>	42.8 <sup>bcde</sup>	1.7 <sup>c</sup>	7.4 <sup>c</sup>	29.1 <sup>a</sup>	6.7 <sup>c</sup>	13.3 <sup>cd</sup>
Q	34.6 <sup>e</sup>	1.4 <sup>c</sup>	5.2 <sup>c</sup>	2.5 <sup>c</sup>	33.0 <sup>a</sup>	1.5 <sup>e</sup>
<b>U+Q</b>	50.4 <sup>ab</sup>	1.4 <sup>c</sup>	8.3 <sup>c</sup>	1.5 <sup>c</sup>	37.7 <sup>a</sup>	23.3 <sup>b</sup>
U	36.2 <sup>de</sup>	1.9 <sup>c</sup>	8.1 <sup>c</sup>	2.8 <sup>c</sup>	1.5 <sup>c</sup>	31.2 <sup>a</sup>
<b>U+S+N+C+Q</b>	57.5 <sup>a</sup>	32.3 <sup>b</sup>	20.1 <sup>b</sup>	10.5 <sup>b</sup>	23.1 <sup>b</sup>	7.4 <sup>de</sup>
<i>df</i> of Sample	9	9	9	9	9	9
<i>df</i> of Interaction	90	90	90	90	90	90
F-value of Sample Effect	2.64	113.66	23.5	28.39	21.03	21.16
Sample significance (p)	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

728 <sup>abcde</sup> **Values within a column which don't share a common superscript are significantly**  
 729 **different in means ratings of the perceived magnitude** from Tukey's HSD test at the  
 730 95% confidence interval. S = sucrose; N = sodium chloride; C = citric acid; Q = quinine  
 731 hemisulfate salt monohydrate; U = monosodium glutamate (MSG). *df* = degrees of  
 732 freedom of interaction, noting that the main effect of sample (F-value of sample) was  
 733 determined by dividing the variance of sample by the variance of the interaction  
 734 ( $MS_{\text{sample}}/MS_{\text{interaction}}$ ) hence both the *df* of sample and interaction are given.

735



736 **Supplementary Table 3 for Experiment 3.** Ratings and significance testing (ANOVA)  
 737 results of perceived intensity (antilogged values) of overall taste, sweet, salty, sour,  
 738 bitter and umami where MPG was used as the umami tastant.

Sample	<b>Perceived intensity</b> (mean of antilogged gLMS intensity ratings)					
	Total intensity	Sweet	Salty	Sour	Bitter	Umami
S	29.2 <sup>cd</sup>	28.6 <sup>a</sup>	2.2 <sup>c</sup>	1.2 <sup>c</sup>	1.3 <sup>c</sup>	1.1 <sup>d</sup>
<b>U+S</b>	35.3 <sup>abc</sup>	28.1 <sup>a</sup>	4.2 <sup>c</sup>	2.0 <sup>c</sup>	3.4 <sup>a</sup>	14.5 <sup>b</sup>
N	23.5 <sup>d</sup>	1.0 <sup>c</sup>	22.5 <sup>a</sup>	1.6 <sup>c</sup>	3.3 <sup>c</sup>	2.4 <sup>cd</sup>
<b>U+N</b>	34.2 <sup>bc</sup>	1.3 <sup>c</sup>	25.2 <sup>a</sup>	1.6 <sup>c</sup>	2.0 <sup>a</sup>	18.9 <sup>b</sup>
C	29.6 <sup>cd</sup>	1.4 <sup>c</sup>	1.5 <sup>c</sup>	26.3 <sup>a</sup>	5.8 <sup>bc</sup>	1.1 <sup>d</sup>
<b>U+C</b>	36.0 <sup>abc</sup>	1.3 <sup>c</sup>	3.7 <sup>c</sup>	28.8 <sup>a</sup>	6.1 <sup>bc</sup>	15.7 <sup>b</sup>
Q	32.8 <sup>bc</sup>	1.1 <sup>c</sup>	1.5 <sup>c</sup>	1.2 <sup>c</sup>	29.7 <sup>a</sup>	1.4 <sup>d</sup>
<b>U+Q</b>	38.5 <sup>ab</sup>	1.1 <sup>c</sup>	2.7 <sup>c</sup>	3.4 <sup>c</sup>	32.4 <sup>a</sup>	15.8 <sup>b</sup>
U	29.0 <sup>cd</sup>	1.3 <sup>c</sup>	3.1 <sup>c</sup>	1.3 <sup>c</sup>	3.8 <sup>bc</sup>	27.2 <sup>a</sup>
<b>U+S+N+C+Q</b>	42.2 <sup>a</sup>	21.6 <sup>b</sup>	16.7 <sup>b</sup>	13.2 <sup>b</sup>	9.7 <sup>b</sup>	7.2 <sup>c</sup>
<i>df</i> of Sample	9	9	9	9	9	9
<i>df</i> of Interaction	99	99	99	99	99	99
F-value of Sample Effect	3.98	65.36	34.69	37.19	26.64	21.49
Sample significance (p)	0.0002	<0.0001	<0.0001	<0.0001	<0.0001	<0.0001

739 <sup>abcde</sup> **Values within a column which don't share a common superscript are significantly**  
 740 **different in means ratings of the perceived magnitude** from Tukey's HSD test at the  
 741 95% confidence interval. S = sucrose; N = sodium chloride; C = citric acid; Q = quinine  
 742 hemisulfate salt monohydrate; U = potassium L-glutamate monohydrate (MPG). *df* =  
 743 degrees of freedom of interaction, noting that the main effect of sample (F-value of  
 744 sample) was determined by dividing the variance of sample by the variance of the  
 745 interaction (MS<sub>sample</sub>/MS<sub>interaction</sub>) hence both the *df* of sample and interaction are  
 746 given.

747

748