

Tracking the impacts of climate change on human health via indicators: lessons from the Lancet Countdown

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Di Napoli, C., McGushin, A., Romanello, M., Ayeb-Karlsson, S., Cai, W., Chambers, J., Dasgupta, S., Escobar, L. E., Kelman, I., Kjellstrom, T., Kniveton, D., Liu, Y., Liu, Z., Lowe, R., Martinez-Urtaza, J., McMichael, C., Moradi-Lakeh, M., Murray, K. A., Rabbaniha, M., Semenza, J. C., Shi, L., Tabatabaei, M., Trinanes, J. A., Vu, B. N., Brimicombe, C. and Robinson, E. J. (2022) Tracking the impacts of climate change on human health via indicators: lessons from the Lancet Countdown. BMC Public Health, 22 (1). ISSN 1471-2458 doi: 10.1186/s12889-022-13055-6 Available at https://centaur.reading.ac.uk/104552/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

Published version at: https://doi.org/10.1186/s12889-022-13055-6 To link to this article DOI: http://dx.doi.org/10.1186/s12889-022-13055-6

Publisher: Springer Nature

All outputs in CentAUR are protected by Intellectual Property Rights law,



including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

https://doi.org/10.1186/s12889-022-13055-6

(2022) 22:663

Di Napoli et al. BMC Public Health

Open Access

Tracking the impacts of climate change on human health via indicators: lessons from the *Lancet* Countdown

Claudia Di Napoli^{1,2*}, Alice McGushin³, Marina Romanello³, Sonja Ayeb-Karlsson^{4,5,6}, Wenjia Cai⁷, Jonathan Chambers⁸, Shouro Dasgupta^{9,10,11}, Luis E. Escobar¹², Ilan Kelman^{3,4,13}, Tord Kjellstrom¹⁴, Dominic Kniveton⁵, Yang Liu¹⁵, Zhao Liu⁷, Rachel Lowe^{16,17,18}, Jaime Martinez-Urtaza¹⁹, Celia McMichael²⁰, Maziar Moradi-Lakeh²¹, Kris A. Murray^{22,23}, Mahnaz Rabbaniha²⁴, Jan C. Semenza²⁵, Liuhua Shi¹⁵, Meisam Tabatabaei^{26,27}, Joaquin A. Trinanes²⁸, Bryan N. Vu¹⁵, Chloe Brimicombe² and Elizabeth J. Robinson⁹

Abstract

Background: In the past decades, climate change has been impacting human lives and health via extreme weather and climate events and alterations in labour capacity, food security, and the prevalence and geographical distribution of infectious diseases across the globe. Climate change and health indicators (CCHIs) are workable tools designed to capture the complex set of interdependent interactions through which climate change is affecting human health. Since 2015, a novel sub-set of CCHIs, focusing on climate change impacts, exposures, and vulnerability indicators (CCIEVIs) has been developed, refined, and integrated by Working Group 1 of the *"Lancet* Countdown: Tracking Progress on Health and Climate Change", an international collaboration across disciplines that include climate, geography, epidemiology, occupation health, and economics.

Discussion: This research in practice article is a reflective narrative documenting how we have developed CCIEVIs as a discrete set of quantifiable indicators that are updated annually to provide the most recent picture of climate change's impacts on human health. In our experience, the main challenge was to define globally relevant indicators that also have local relevance and as such can support decision making across multiple spatial scales. We found a hazard, exposure, and vulnerability framework to be effective in this regard. We here describe how we used such a framework to define CCIEVIs based on both data availability and the indicators' relevance to climate change and human health. We also report on how CCIEVIs have been improved and added to, detailing the underlying data and methods, and in doing so provide the defining quality criteria for *Lancet* Countdown CCIEVIs.

Conclusions: Our experience shows that CCIEVIs can effectively contribute to a world-wide monitoring system that aims to track, communicate, and harness evidence on climate-induced health impacts towards effective intervention strategies. An ongoing challenge is how to improve CCIEVIs so that the description of the linkages between climate change and human health can become more and more comprehensive.

Keywords: Climate change, Public health, Indicators, Climate data, Policy making

 $^{\rm 1}$ School of Agriculture, Policy and Development, University of Reading, Reading, UK

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/A.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/licenses/by/A.0/. The Creative Commons Public Domain Dedicated in a credit line to the data.



^{*}Correspondence: c.dinapoli@reading.ac.uk

Background

Climate change affects global health via multiple direct and indirect pathways [1, 2]. Every year, disasters involving weather- and climate-related hazards result in thousands of deaths worldwide and contribute to the global burden of disease [3, 4]. Direct health consequences may derive from changes in temperature and precipitation, and human exposure to heatwaves, wildfires, floods, and droughts [5-9]. These include an increase in cardiovascular mortality during events of extreme heat; a higher incidence of chronic kidney disease among outdoor workers in hot areas; and fatalities and multiple negative health consequences of fire smoke inhalation during wildfires [10–13]. Indirect impacts may be triggered by climate change-induced environmental and ecosystem alterations, such as crop failures, reduced marine food capture, geographic range expansion of disease vectors, and reduced labour capacity [14-17]. Other indirect effects may be mediated through social systems and responses to climate change. Examples include reduced labour capacity in vulnerable occupations due to increasing heat, migration of populations related to sea-level rise and food insecurity, as well as other drivers of human mobility [18-20].

As understanding and awareness grow regarding the health dimension of climate change, there is an increased demand for feasible and efficient tools that can inform climate change mitigation and adaptation strategies to deliver benefits to human health at all levels. Climatehealth linkages are complex, with multiple interactions, synergies, and feedback loops. Understanding their characteristics and relationships is challenging and poses a level of complexity that could render the development of such tools unfeasible. A reasonable alternative is to deploy a subset of measures describing the key aspects of the state, extent and change of the climate-health system [21].

Climate change and health indicators (CCHIs) are summary measures, often quantitative, that represent the heterogeneous set of factors and relationships linking climate change with health [22–26]. Their purpose is three-fold. First, to assess long-term trends and changes in climate-induced impacts on population health. Second, to effectively communicate relevant evidence to researchers, health professionals, policymakers, and the general public. Third, to support and monitor decision making for successful intervention strategies and plans that address the human health consequences of climate change.

In the past decade, CCHIs have represented the building blocks for a variety of initiatives tracking the climate-health system across multiple spatial and temporal scales [24, 25]. One high profile and current CCHI initiative is the *"Lancet* Countdown: Tracking Progress on Health and Climate Change". Since 2015 the *Lancet* Countdown has been providing a global monitoring system that tracks the complex, reflective ways in which climate change affects health [27–32]. It accomplishes this through a set of CCHIs spanning five domains of the climate-health system: climate change impacts, exposures, and vulnerability; adaptation planning and resilience for health; mitigation actions and health co-benefits; economics and finance; and public and political engagement.

One of the five Lancet Countdown thematic working groups, namely Working Group 1 (WG1), is tasked with identifying and tracking indicators that trace the links across "climate change impacts, exposures, and vulnerability". This research in practice article aims to provide guidance for the development and tracking of CCHIs through a detailed case study of how WG1 authors developed these indicators, referred to from here on as climate change impacts, exposures, and vulnerability indicators (CCIEVIs), that contribute to the *Lancet* Countdown's overall objectives. As such, this article is equally relevant for those individuals and policy makers wanting to develop their own climate change and health indicators specific to a country or region, and those wanting a more detailed understanding of specific indicators used in the Lancet Countdown, and particularly WG1. Hereafter, we describe (i) the aim, design, and characteristics of CCIEVIs, as employed in the Lancet Countdown, (ii) their main components, (iii) underlying data and methods, and (iv) plans for future improvements.

Climate change impacts, exposures, and vulnerability indicators Designing CCIEVIs

Lancet Countdown CCIEVIs have been developed to monitor the health impacts related to anthropogenic climate change, their trends to date, and the extent to which progress (or backsliding) is occurring over time. They are presented from a historical starting date up to the year for which the most recent complete data are available. The indicators provide evidence along different dimensions - some indicators are positioned closer to the climate change hazard and some closer to the health impact – as to whether, and to what extent, climate change is affecting health, for better or for worse. In the Lancet Countdown, CCIEVIs start tracking climate change-health linkages from the 1980s, and are grouped into five thematic clusters: heat; weather and climate extremes; infectious diseases; food security of terrestrial and marine assets; global mean sea-level rise.

To ensure that CCIEVIs are reproducible, relevant, and useful to scientific and policy-making communities, they must satisfy five quality criteria that mirror the World Meteorological Organisation guidelines for climate indicators [33]:

- 1. *Representativeness:* a CCIEVI should track an aspect of both climate change and health, particularly focusing on the relationship between the two; it should do this across a timescale and geographical coverage sufficient for long-term global trends to be observed.
- 2. *Relevance:* a CCIEVI should be clear and understandable to a broad range of audiences; global CCIEVIs may also have value at local (i.e. national, regional) level for policy and decision makers.
- 3. *Robustness:* a CCIEVI should use data and methods that are robust, reliable, and valid to track the relevant aspect of climate change and health; data from publicly available databases, and especially those developed by international organisations, governmental bodies, or academic institutions, are preferred; methods should be supported by a high standard of evidence from the scientific literature.
- 4. *Reproducibility:* a CCIEVI should be calculated using an internationally agreed and published scientific method as well as open-access and qualitycontrolled data; the methodology underlying an indicator should be clearly laid out, including details of the process of data collection and processing, which must be done in a systematic and unbiased way; statistical analysis of the data should be carried out to support the interpretation of the data.
- 5. *Timeliness:* a CCIEVI should be calculated regularly, with a short lag between the end of the period under consideration and the publication of the data; the calculation must be practicable with existing and future resources.

WG1 indicators have evolved over the five years that they have been reported in the annual Lancet Countdown publications so to fulfil above-listed criteria. For example, in 2020, there were twelve WG1 Lancet Countdown CCIEVIs, almost double the number of when they first appeared in 2016 [28, 32]. As well as new indicators being introduced each year, established indicators are revised to reflect the latest evidence in the scientific literature and the needs of new and emerging stakeholders [30, 31]. The revision of Lancet Countdown CCIEVIs occurs annually by experts across a broad range of relevant disciplines, including climate, geography, epidemiology, and occupation health. The publication of an appendix alongside each Lancet Countdown annual report ensures that the data and methods underlying every CCIEVI are explicitly cited and described along with possible caveats.

Main components of CCIEVIs

Lancet Countdown CCIEVIs focus on three main components: hazard, exposure, and vulnerability. These are defined according to the terminology provided by the United Nations Office for Disaster Risk Reduction and the Intergovernmental Panel on Climate Change [34, 35]. Specifically:

- Hazard is any physical event with the potential to cause disruption or damage in vulnerable and exposed elements;
- Exposure is who or what is present in the area where a hazard may occur;
- Vulnerability is the factors or constraints of an economic, social, physical, or geographic nature, which reduce the ability of exposed elements to prepare for and cope with the impact of the hazard.

In Lancet Countdown CCIEVIs, each of these components represents a dimension or layer of the multifacet climate-health system. For the hazards, the Lancet Countdown CCIEVIs target extremes in temperature and precipitation, namely heatwaves, drought, and floods, as well as phenomena mediated by these two environmental variables, i.e. wildfires and suitability of vector-borne diseases (malaria, dengue). Similarly, health hazards in the ocean system are identified in changes of global mean sea level as well as sea surface temperature and salinity. The second layer tracks the exposure of populations, cultivated areas, and health systems to hazards. This reflects the purpose of representing climate change-health pathways that are both direct (e.g. on people) and indirect (e.g. food mediated). Lancet Countdown CCIEVIs also track vulnerabilities to extreme heat, incorporating the proportion of the population over 65 years of age, the prevalence of predisposing chronic diseases (i.e. diabetes and cardiovascular, respiratory, and renal disease), the proportion of the population living in urban areas, and the number of outdoor workers. In the 2021 report, the vulnerability of children younger than 1 year to lifethreatening heatwaves was added [36]. Lancet Countdown CCIEVIs also consider vulnerability linked to the capacity of local health services to respond to public health risks and emergencies.

Data and methods for developing CCIEVIs

Hazards, exposures, and vulnerabilities change over time and across space as do their interactions as a result of changes in the climate. Consequently, climate change amplifies or diminishes existing health impacts, and induces or suppresses new ones with a high degree of spatial and temporal variability. Geographic information systems, and more specifically geospatial data, have long been identified as useful analytical tools for monitoring the health-climate system at all spatial levels, from local to global [37].

Many relevant sources for CCIEVIs provide the data as rasters, i.e. grids of cells. This allows information, such as variations in climate-relevant variables like air temperature, to be represented in a spatially continuous and consistent way across the Earth's surface. Usefully, rasters can be stacked to assess cumulative hazards in a given region or to produce spatially resolved time series of a given variable.

Environmental data are nowadays available as rasters across a wide range of spatial and temporal scales. This makes them an ideal tool for constructing relationships between hazards, exposed elements, and associated vulnerability anywhere in the world, low and middle income countries included [38]. For instance, one of the Lancet Countdown CCIEVIs tracks the climatic suitability for the transmission of *Plasmodium falciparum*, the parasite causing malaria. Following the work by Grover-Kopec and colleagues [39], the indicator tracks the number of months per year suitable for malaria transmission as the coincidence of precipitation accumulation greater than 80mm, average temperature between 18°C and 32°C, and relative humidity greater than 60%. The number of suitable months in a year is calculated for each grid cell by overlaying precipitation, temperature, and humidity raster layers across twelve months. Year-by-year changes in the number of suitable months generate a time series, which can be spatially aggregated (e.g. for highland vs lowland areas).

CCIEVIs can be explored via an online visualization platform. For example, maps represent a powerful way to visualise and communicate climate information, and harness the latter for action [40]. CCIEVIs can be displayed as a set of world-wide maps that can be navigated via an interactive, user-friendly interface to show year-specific data or to highlight countries or geographical areas of interest, as has been done by the *Lancet* Countdown [41].

Perspectives and improvements for CCIEVIs

Climate change is an evolving phenomenon and our understanding of it is also evolving. Only with the most up-to-date data, methodologies, and expertise, CCIEVIs can provide the quantitative underpinnings of a compelling narrative on the health impacts that climate change imposes across the globe. Based on our experience from the *Lancet* Countdown, we here report the main points CCIEVIs might consider to be meaningful and useful in their purpose.

1. *Partnership:* The cross-cutting nature of CCIEVIs demands a combination of skills, knowledge and data that span across institutions, disciplines, countries, and geographical regions. Creating and maintaining a long-term collaboration among a group of diverse experts is crucial to guaranteeing the robustness and reliability of developed CCIEVIs over time. Partner-

ship can also foster the development of CCIEVIbased initiatives in locations where these are currently missing but are perceived as useful [42–46]. Over the long term, partnerships can expand the areas where CCIEVIs drive decision-making as well as increase workforce preparedness via the inclusion of climate change curriculum into health professional education [47–50].

- 2. *Iterative process:* To satisfy the timeliness criteria of CCIEVIs, their design and utility must be revised periodically [21]. Every year, *Lancet* Countdown CCIEVIs indicators undergo a thorough quality check and improvement process before being considered for the annual report. In this process, independent experts assess the quality and suitability of each indicator and provide constructive feedback to aid their development and improvement. Additionally, new CCIEVIs can be developed and added to the original indicators suite to provide a more complete description of the climate-health system.
- 3. Going local: Because CCIEVIs are provided as geospatial data, they are down-scalable and able to identify priority areas for public health intervention across the globe. Location-specific CCIEVIs can assist local health departments in tracking variations in community exposure and vulnerability to climate change, uncovering health impacts at regional or sub-regional level (including their linkages to the surrounding urban/natural environment), designing interventions to enhance community resilience and evaluating the effectiveness of implemented interventions [51, 52]. With this motivation, the Lancet Countdown CCIEVIs structure has been replicated to produce the Australia and China reports with indicators being provided at national to regional and city scales [53, 54]. These local reports may serve as a guidance for disseminating progress on CCIEVIs from other countries in a uniform format. Every year, the Lancet Countdown report is also followed by a range of resources, such as briefs for policymakers and translations of the executive summaries, tailored to specific cities, countries and geographical areas [55-57].
- 4. *Data*: CCIEVIs typically have to rely on different sources for the provision of health, climate, and demographics data. They therefore may well differ in the spatial resolution, the time period covered, and the reference baseline used for the definition of extreme events. Notwithstanding this, consistency in climate data and demographics among *Lancet* Countdown CCIEVIs has been attempted and achieved wherever possible so that indicators are compatible and comparable with each other. It is

worth noting that protocols are in place for the effective archival, management, analysis, delivery, and use of climate data, whereas health data standardisation is an open topic [58, 59]. The collection and reporting of public health data, for instance, vary greatly across nations and healthcare organizations [60]. Data monitoring on disease incidence or health outcomes that are standardised at a global level is in general lacking. In most cases, health impacts are therefore tracked using epidemiological models rather than measured records. Furthermore, their geographical resolution is generally lower than climate data (i.e. country level rather than at raster cell level). Unlike climate data, health records are generally not publicly available. Creating open-access online databases for public health data would help foster knowledge exchange between the climate and the public health community, as well as promote its applicability at all scales.

CCIEVIs can be used as a starting point for broader explorations of the linkages between climate change and health that have relevance for policy making. Here we suggest six specific areas for future research, many of which are currently being pursued by WG1 authors.

- 1. Looking to the future: In the Lancet Countdown, CCIEVIs are used to track the current health impacts of climate change, and contextualise them with respect to the recent past. These indicators therefore hint at how the future will likely evolve without efforts to mitigate and adapt to climate change. However, the methodology described in this research in practice article is time-agnostic and can be applied to indicators predictive of the future. Designing, developing and implementing future projections of CCIEVIs presents additional research challenges, such as the choice of the climate prediction model and its verification, but has the potential to provide actionable information to support policymaking [61].
- 2. Identifying hotspots: Climate change often affects people and places in multiple ways. We define *climate-health hotspots* as locations where climate affects people negatively through multiple pathways. This definition builds upon work already established in the climatology field on mapping *hotspots*, i.e. geographical areas where the combined occurrence of multiple weather extremes (compound events) is observed [62, 63]. As adverse health effects occur in populations that are already at risk from climate extremes and lack adequate health infrastructure [64–66], the overlay method described in this work can be adapted to identify climate-health hotspots and trace them overtime by overlapping multiple lay-

ers of hazard, exposure and vulnerability. It can also be applied to different CCIEVIs to assess multiple climate change impacts in a single location. This could allow for a *one health* perspective if CCIEVIs at the human-animal-environment interface are implemented [67, 68]. For instance, concerning food security of terrestrial assets, a CCIEVI tracking changes in crop yield potential due to rising temperatures could be overlayed with CCIEVIs monitoring animal-source foods and/or the effects of environmental changes on diseases of livestock and crops. In either case, the identification and tracking of climate-health hotspots and incorporation into a monitoring system, could improve further public health action, particularly in the context of reducing the likelihood of systemic health crises [69].

- 3. *Inequality and inequity:* Global indicators generally do not provide a nuanced picture of the differentiated impacts of climate change on health by country and region, or across populations within countries. Disaggregating indicators by relevant socioeconomic and occupational characteristics (e.g. income, gender, race or ethnicity, disability, occupation, and age) is necessary to measure inequalities in exposure to climatic risk factors and health outcomes attributable to these exposures. For heatwaves, for instance, these may consider pregnancy and mental health [70, 71]. With this approach, climate change and climate-related risk factors are positioned as both environmental and social determinants of health.
- 4. Mental health: Mental health issues are closely intertwined with people's geographical, social, environmental, financial, and political context, and any climate-induced impact on physical health is also a risk factor for mental health [72]. As such, factors linking climate change to health may affect mental health and underlying mental health status can affect the capacity to adapt to (or to mitigate) climate change. The description of the climate-health system, therefore, requires the inclusion of CCIEVIs that address the mental health consequences of climate change. A growing body of literature is currently examining these linkages [73–79], inspiring the development of future CCIEVIS on the topic.
- 5. *Attribution*: In recent years, formal methods have been developed to identify changes in the occurrence of adverse health outcomes, and to determine the extent to which those changes may be attributed to climate change [80]. Multi-step attribution, for instance, consists of a) attributing an observed change in a variable of interest to a change in climate or other environmental variable and b) attributing the change in climate to external drivers such

as greenhouse gas emissions [81]. The 2020 *Lancet* Countdown report showcased the attribution of 76 extreme weather events to climate change, and the effects that a selection of these have had on the health of the population [32]. Future CCIEVIs could be based on attribution to quantitatively understand how climate change makes extreme events associated with health outcomes more likely.

6. *Embracing the unquantifiable*: There are important aspects of the relationship between climate change and health that cannot be quantified or are very difficult to quantify meaningfully. One example of this is human migration [18]. Local-level studies find worsening food security where people move away from places affected by sea level rise and saltwater intrusion, or adverse psychosocial impacts of disrupted lives, social networks and livelihoods for those on the move in a changing climate. It is extremely challenging, however, to attribute human mobility to climate change (and quantify its magnitude), or to develop a global CCIEVI that can quantify links between climate impacts, human mobility, and health.

Conclusions

In developing *climate change impacts, exposures, and vulnerability indicators* (CCIEVIs), several key challenges have to be addressed around the idea of representing highly complex systems by a discrete set of quantifiable indicators that are updated annually. One challenge is to find globally relevant indicators, with data available across a sufficiently large number of countries, that also have local relevance.

A hazard, exposure, and vulnerability framework has proven to be an effective structure within the *Lancet* Countdown initiative. Around such a framework, *Lancet* Countdown CCIEVIs have been crafted, based on both data availability and the indicators' relevance to climate change and human health. Over time these indicators have been improved and added to, as data become available and methodologies are refined. An ongoing challenge is how to incorporate those linkages between climate change and health that are clear and well documented but at the same time difficult to measure, quantify, and fit within this framework.

Together, *Lancet* Countdown CCIEVIs present a compelling visualisation of how climate change is increasingly exposing people to the negative health impacts of climate change, both direct and indirect, across land and water. Ultimately the impact of climate change on human health depends not only on the exposure and vulnerability of populations to climate hazards, but also on the extent to which individuals and countries are able to adapt and build resilience. By unveiling the challenges underpinning the evolution of the climate-health system in time and in space, CCIEVIs can be the pillars of a new public health that safeguards and advances global wellbeing in the decades to come.

Acknowledgments

The authors are grateful to the three anonymous reviewers and the editor for their constructive comments.

Authors' contributions

CDN, AM, MR, SAK, WC, JC, SD, LEE, IK, TK, DK, YL, ZL, RL, JMU, CM, MML, KAM, MR2, JCS, LS, MT, JAT, BNV, CB, and EJR authors planned and designed this manuscript. CDN prepared the first draft; CDN, AM, MR, SAK, WC, JC, DLEE, IK, TK, DK, YL, ZL, RL, JMU, CM, MML, KAM, MR2, JCS, LS, MT, JAT, BNV, CB, and EJR provided input into subsequent drafts. CDN, AM, MR, SAK, WC, JC, SD, LEE, IK, TK, DK, YL, ZL, RL, JMU, CM, MML, KAM, MR2, JCS, LS, MT, JAT, BNV, CB, and EJR authors read and approved the final manuscript.

Funding

This work is supported by an unrestricted grant from the Wellcome Trust (209734/Z/17/Z).

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

School of Agriculture, Policy and Development, University of Reading, Reading, UK.²Department of Geography and Environmental Science, University of Reading, Reading, UK. ³Institute for Global Health, University College London, London, UK. ⁴Institute for Risk and Disaster Reduction, University College London, London, UK. ⁵School of Global Studies, University of Sussex, Brighton Falmer, UK. ⁶United Nations University, Institute for Environment and Human Security, Bonn, Germany. ⁷Ministry of Education Key Laboratory for Earth System modeling, Department of Earth System Science, Tsinghua University, Beijing 100084, China.⁸Institute for Environmental Science, University of Geneva, Geneva, Switzerland. ⁹Grantham Research Institute on Climate Change and the Environment London School of Economics and Political Science (LSE), London, UK. ¹⁰Centro Euro-Mediterraneo sui Cambiamenti Climatici (CMCC), Venice, Italy.¹¹Università Ca' Foscari, Venice, Italy.¹²Department of Fish and Wildlife Conservation, Virginia Tech, Blacksburg, VA, USA. ¹³University of Agder, Kristiansand, Norway. ¹⁴Health and Environment International Trust, Nelson, New Zealand. ¹⁵Rollins School of Public Health, Emory University, Atlanta, USA.¹⁶Barcelona Supercomputing Center, Barcelona, Spain. ¹⁷Catalan Institution for Research and Advanced Studies (ICREA), Barcelona, Spain. ¹⁸Centre on Climate Change & Planetary Health and Centre for Mathematical Modelling of Infectious Diseases, London School of Hygiene & Tropical Medicine, London, UK. ¹⁹Department of Genetics and Microbiology, Universitat Autònoma de Barcelona (UAB), Barcelona, Spain.²⁰School of Geography, Earth and Atmospheric Sciences. The University of Melbourne, Melbourne, Australia.²¹ Preventive Medicine and Public Health Research Center, Psychosocial Health Research Institute, Iran University of Medical Sciences, Tehran, Iran. ²²MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK.²³MRC Unit The Gambia At London School of Hygiene and Tropical Medicine. Atlantic Boulevard, Fajara, The Gambia, ²⁴Iranian Fisheries Science Research Institute, Agricultural Research, Education, and Extension

Organisation, Tehran, Iran. ²⁵Heidelberg Institute of Global Health, University of Heidelberg, Heidelberg, Germany. ²⁶Higher Institution Centre of Excellence (HICoE), Institute of Tropical Aquaculture and Fisheries (AKUATROP), Universiti Malaysia Terengganu, 21030 Kuala Nerus, Terengganu, Malaysia. ²⁷Henan Province Forest Resources Sustainable Development and High-value Utilization Engineering Research Center, School of Forestry, Henan Agricultural University, Zhengzhou 450002, China. ²⁸Department of Electronics and Computer Science, Universidade de Santiago de Compostela, Santiago, Spain.

Received: 1 October 2021 Accepted: 22 March 2022 Published online: 06 April 2022

References

- Haines A, Kovats RS, Campbell-Lendrum D, Corvalan C. Climate change and human health: impacts, vulnerability and public health. Public Health. 2006;120(7):585–96.
- McMichael AJ, Haines J, Slooff R, Sari Kovats R, Organization WH. Climate change and human health: an assessment. World Health Organization; 1996.
- WHO. Climate change and health 2018 [Available from: https://www. who.int/news-room/fact-sheets/detail/climate-change-and-health.
- Confalonieri U, Menne B, Akhtar R, Ebi KL, Hauengue M, Kovats RS, et al. Human health. climate change 2007: impacts, adaptation and vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Cambridge University Press, Cambridge, UK; 2007.
- Xu R, Yu P, Abramson MJ, Johnston FH, Samet JM, Bell ML, et al. Wildfires, global climate change, and human health. N Engl J Med. 2020;383(22):2173–81.
- Alderman K, Turner LR, Tong S. Floods and human health: a systematic review. Environ Int. 2012;47:37–47.
- Stanke C, Kerac M, Prudhomme C, Medlock J, Murray V. Health effects of drought: a systematic review of the evidence. PLoS Curr. 2013;5.
- 8. Campbell S, Remenyi TA, White CJ, Johnston FH. Heatwave and health impact research: a global review. Health Place. 2018;53:210–8.
- Di Napoli C, Pappenberger F, Cloke HL. Assessing heat-related health risk in Europe via the universal thermal climate index (UTCI). Int J Biometeorol. 2018;62(7):1155–65.
- Gostimirovic M, Novakovic R, Rajkovic J, Djokic V, Terzic D, Putnik S, et al. The influence of climate change on human cardiovascular function. Arch Environ Occup Health. 2020;75(7):406–14.
- Nerbass FB, Pecoits-Filho R, Clark WF, Sontrop JM, McIntyre CW, Moist L. Occupational heat stress and kidney health: from farms to factories. Kidney Int Rep. 2017;2(6):998–1008.
- 12. Borchers Arriagada N, Bowman DM, Palmer AJ, Johnston FH. Climate change, wildfires, heatwaves and health impacts in Australia. Extreme weather events and human health: Springer; 2020. p. 99-116.
- Adetona O, Reinhardt TE, Domitrovich J, Broyles G, Adetona AM, Kleinman MT, et al. Review of the health effects of wildland fire smoke on wildland firefighters and the public. Inhal Toxicol. 2016;28(3):95–139.
- Ray DK, West PC, Clark M, Gerber JS, Prishchepov AV, Chatterjee S. Climate change has likely already affected global food production. PLoS One. 2019;14(5):e0217148.
- Brander KM. Global fish production and climate change. PNAS. 2007;104(50):19709–14.
- Kjellstrom T, Briggs D, Freyberg C, Lemke B, Otto M, Hyatt O. Heat, human performance, and occupational health: a key issue for the assessment of global climate change impacts. Annu Rev Public Health. 2016;37(1):97–112.
- Patz J, Githeko A, McCarty J, Hussein S, Confalonieri U, De Wet N. Climate change and infectious diseases. Climate change and human health: risks and responses 2003;2:103-132.
- McMichael C, Dasgupta S, Ayeb-Karlsson S, Kelman I. A review of estimating population exposure to sea-level rise and the relevance for migration. Environ Res Lett. 2020;15(12):123005.
- Schwerdtle PN, McMichael C, Mank I, Sauerborn R, Danquah I, Bowen KJ. Health and migration in the context of a changing climate: a systematic literature assessment. Environ Res Lett. 2020;15(10).

- Kjellstrom T, Holmer I, Lemke B. Workplace heat stress, health and productivity - an increasing challenge for low and middle-income countries during climate change. Glob Health Action. 2009;2.
- Kenney MA, Janetos AC, Gerst MD. A framework for national climate indicators. Clim Chang. 2018;163(4):1705–18.
- English PB, Sinclair AH, Ross Z, Anderson H, Boothe V, Davis C, et al. Environmental health indicators of climate change for the United States: findings from the state environmental health Indicator collaborative. Environ Health Perspect. 2009;117(11):1673–81.
- Navi M, Hansen A, Nitschke M, Hanson-Easey S, Pisaniello D. Developing health-related indicators of climate change: Australian stakeholder perspectives. Int J Environ Res Public Health. 2017;14(5).
- Murray KA, Escobar LE, Lowe R, Rocklov J, Semenza JC, Watts N. Tracking infectious diseases in a warming world. BMJ. 2020;371:m3086.
- Hambling T, Weinstein P, Slaney D. A review of frameworks for developing environmental health indicators for climate change and health. Int J Environ Res Public Health. 2011;8(7):2854–75.
- Liu AY, Trtanj JM, Lipp EK, Balbus JM. Toward an integrated system of climate change and human health indicators: a conceptual framework. Clim Change. 2021;166(49).
- Watts N, Adger WN, Agnolucci P, Blackstock J, Byass P, Cai W, et al. Health and climate change: policy responses to protect public health. Lancet. 2015;386(10006):1861–914.
- Watts N, Adger WN, Ayeb-Karlsson S, Bai Y, Byass P, Campbell-Lendrum D, et al. The lancet countdown: tracking progress on health and climate change. Lancet. 2017;389(10074):1151–64.
- Watts N, Amann M, Ayeb-Karlsson S, Belesova K, Bouley T, Boykoff M, et al. The lancet countdown on health and climate change: from 25 years of inaction to a global transformation for public health. Lancet. 2018;391(10120):581–630.
- Watts N, Amann M, Arnell N, Ayeb-Karlsson S, Belesova K, Berry H, et al. The 2018 report of the lancet countdown on health and climate change: shaping the health of nations for centuries to come. Lancet. 2018;392(10163):2479–514.
- Watts N, Amann M, Arnell N, Ayeb-Karlsson S, Belesova K, Boykoff M, et al. The 2019 report of the lancet countdown on health and climate change: ensuring that the health of a child born today is not defined by a changing climate. Lancet. 2019;394(10211):1836–78.
- Watts N, Amann M, Arnell N, Ayeb-Karlsson S, Beagley J, Belesova K, et al. The 2020 report of the lancet countdown on health and climate change: responding to converging crises. Lancet. 2021;397(10269):129–70.
- Williams M, Eggleston S. Using indicators to explain our changing climate to policymakers and the public. World Meteorological Organisation Bulletin. 2017;66(2).
- 34. UNDRR. Online glossary 2020 [Available from: https://www.undrr.org/ terminology.
- 35. Cardona OD, Van Aalst MK, Birkmann J, Fordham M, Mc Gregor G, Rosa P, et al. Determinants of risk: exposure and vulnerability. Managing the risks of extreme events and disasters to advance climate change adaptation: special report of the intergovernmental panel on climate change: Cambridge University Press; 2012. p. 65-108.
- Romanello M, McGushin A, Di Napoli C, Drummond P, Hughes N, Jamart L, et al. The 2021 report of the lancet countdown on health and climate change: code red for a healthy future. Lancet. 2021;398(10311):1619–62.
- Costello A, Abbas M, Allen A, Ball S, Bell S, Bellamy R, et al. Managing the health effects of climate change: Lancet and University College London Institute for Global Health Commission. Lancet. 2009;373(9676):1693–733.
- Sarkar S, Gangare V, Singh P, Dhiman RC. Shift in potential malaria transmission areas in India, using the Fuzzy-Based Climate Suitability Malaria Transmission (FCSMT) model under changing climatic conditions. Int J Environ Res Public Health. 2019;16(18):3474.
- Grover-Kopec EK, Blumenthal MB, Ceccato P, Dinku T, Omumbo JA, Connor SJ. Web-based climate information resources for malaria control in Africa. Malar J. 2006;5:38.
- Becsi B, Hohenwallner-Ries D, Grothmann T, Prutsch A, Huber T, Formayer H. Towards better informed adaptation strategies: co-designing climate change impact maps for Austrian regions. Clim Chang. 2019;158(3-4):393–411.

- The Lancet Countdown: Tracking progress on health and climate change. Data platform [Available from: https://www.lancetcountdown.org/dataplatform/.
- Healey Akearok G, Holzman S, Kunnuk J, Kuppaq N, Martos Z, Healey C, et al. Identifying and achieving consensus on health-related indicators of climate change in Nunavut. Arctic. 2019;72(3):289–99.
- Dovie DBK, Dzodzomenyo M, Ogunseitan OA. Sensitivity of health sector indicators' response to climate change in Ghana. Sci Total Environ. 2017;574:837–46.
- Houghton A, Austin J, Beerman A, Horton C. An approach to developing local climate change environmental public health indicators in a rural district. J Environ Public Health. 2017;2017:3407325.
- Doubleday A, Errett NA, Ebi KL, Hess JJ. Indicators to guide and monitor climate change adaptation in the US Pacific northwest. Am J Public Health. 2020;110(2):180–8.
- 46. Bozzi L, Dubrow R. Climate change and health in Connecticut: 2020 report. New Haven, Connecticut: Yale Center on Climate Change and Health.
- Bell EJ. Climate change: what competencies and which medical education and training approaches? BMC Med Educ. 2010;10:31.
- Maxwell J, Blashki G. Teaching about climate change in medical education: an opportunity. J Public Health Res. 2016;5(1):673.
- 49. Finkel ML. A call for action: integrating climate change into the medical school curriculum. Perspect Med Educ. 2019;8(5):265–6.
- Butterfield P, Leffers J, Vásquez MD. Nursing's pivotal role in global climate action. BMJ. 2021;373.
- Houghton A, English P. An approach to developing local climate change environmental public health indicators, vulnerability assessments, and projections of future impacts. J Environ Public Health. 2014;2014:132057.
- Singh P, Yadav Y, Saraswat S, Dhiman RC. Intricacies of using temperature of different niches for assessing impact on malaria transmission. Indian J Med Res. 2016;144(1):67.
- Zhang Y, Beggs PJ, McGushin A, Bambrick H, Trueck S, Hanigan IC, et al. The 2020 special report of the MJA-lancet countdown on health and climate change: lessons learnt from Australia's "Black Summer". Med J Aust. 2020;213(11):490–2 e10.
- Cai W, Zhang C, Suen HP, Ai S, Bai Y, Bao J, et al. The 2020 China report of the lancet countdown on health and climate change. Lancet Public Health. 2021;6(1):e64–81.
- 55. Robinson EJZ, Baldock J, Piears D, Di Napoli C, Dasgupta S. A public health and climate change "countdown" for Reading. Ethical Reading; 2020.
- The Lancet Countdown: Tracking progress on health and climate change. Resources [Available from: https://www.lancetcountdown.org/resources/.
- 57. Romanello M, van Daalen K, Anto JM, Dasandi N, Drummond P, Hamilton IG, et al. Tracking progress on health and climate change in Europe. The Lancet Public Health. 2021.
- 58. World Meteorological Organisation. Climate data management system specifications. Geneva, Switzerland; 2014.
- 59. World Health Organisation. WHO Forum on health data standardization and interoperability. Geneva, Switzerland; 2013.
- Schulz S, Stegwee R, Chronaki C. Standards in Healthcare Data. In: Kubben P, Dumontier M, Dekker A, editors. Fundamentals of Clinical Data Science. Cham (CH)2019. p. 19-36.
- 61. Nissan H, Conway D. From advocacy to action: projecting the health impacts of climate change. PLoS Med. 2018;15(7):e1002624.
- 62. Dilley M, Chen RS, Deichmann U, Lerner-Lam AL, Arnold M. Natural Disaster Hotspots: The World Bank; 2005.
- Zscheischler J, Martius O, Westra S, Bevacqua E, Raymond C, Horton RM, et al. A typology of compound weather and climate events. Nat Rev Earth Environ. 2020;1(7):333–47.
- 64. Patz JA, Kovats RS. Hotspots in climate change and human health. BMJ. 2002;325(7372):1094–8.
- 65. World Bank. Geographic hotspots for world bank action on climate change and health. Washington, DC: World Bank; 2017.
- Piontek F, Muller C, Pugh TA, Clark DB, Deryng D, Elliott J, et al. Multisectoral climate impact hotspots in a warming world. PNAS. 2014;111(9):3233–8.
- 67. Patz JA, Hahn MB. Climate change and human health: a one health approach. One Health: The Human-Animal-Environment Interfaces in Emerging Infectious Diseases 2012:141-171.

- Zinsstag J, Crump L, Schelling E, Hattendorf J, Maidane YO, Ali KO, et al. Climate change and one health. FEMS Microbiol Lett. 2018;365(11):fny085.
- Hui-Min L, Xue-Chun W, Xiao-Fan Z, Ye Q. Understanding systemic risk induced by climate change. Adv Clim Chang Res. 2021;12(3):384–94.
- Chersich MF, Pham MD, Areal A, Haghighi MM, Manyuchi A, Swift CP, et al. Associations between high temperatures in pregnancy and risk of preterm birth, low birth weight, and stillbirths: systematic review and meta-analysis. BMJ. 2020;371.
- Liu J, Varghese BM, Hansen A, Xiang J, Zhang Y, Dear K, et al. Is there an association between hot weather and poor mental health outcomes? A systematic review and meta-analysis. Environ Int 2021;153:106533.
- 72. Willox. Climate Change as the Work of Mourning. Ethics and the Environment. 2012;17(2).
- Stanley SK, Hogg TL, Leviston Z, Walker I. From anger to action: differential impacts of eco-anxiety, eco-depression, and eco-anger on climate action and wellbeing. J Clim Change Health. 2021;1.
- 74. Cunsolo A, Ellis NR. Ecological grief as a mental health response to climate change-related loss. Nat Clim Chang. 2018;8(4):275–81.
- Thompson R, Hornigold R, Page L, Waite T. Associations between high ambient temperatures and heat waves with mental health outcomes: a systematic review. Public Health. 2018;161:171–91.
- Ayeb-Karlsson S.'I do not like her going to the shelter': stories on gendered disaster (im)mobility and wellbeing loss in coastal Bangladesh. Int J Disaster Risk Reduct. 2020;50.
- Berry HL, Waite TD, Dear KBG, Capon AG, Murray V. The case for systems thinking about climate change and mental health. Nat Clim Chang. 2018;8(4):282–90.
- Kelman I, Ayeb-Karlsson S, Rose-Clarke K, Prost A, Ronneberg E, Wheeler N, et al. A review of mental health and wellbeing under climate change in small island developing states (SIDS). Environ Res Lett. 2021;16(3):033007.
- Hayward G, Ayeb-Karlsson S. 'Seeing with Empty Eyes': a systems approach to understand climate change and mental health in Bangladesh. Clim Chang. 2021;165(1-2).
- Ebi KL, Ogden NH, Semenza JC, Woodward A. Detecting and attributing health burdens to climate change. Environ Health Perspect. 2017;125(8):085004.
- Astrom C, Orru H, Rocklov J, Strandberg G, Ebi KL, Forsberg B. Heat-related respiratory hospital admissions in Europe in a changing climate: a health impact assessment. BMJ Open. 2013;3(1).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

