

Representation uncertainty in the earth sciences

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Bulgin, C. E. ORCID: <https://orcid.org/0000-0003-4368-7386>, Thomas, C. M., Waller, J. A. and Woolliams, E. R. (2022) Representation uncertainty in the earth sciences. *Earth and Space Science*, 9 (6). e2021EA002129. ISSN 2333-5084 doi: <https://doi.org/10.1029/2021EA002129> Available at <https://centaur.reading.ac.uk/105184/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1029/2021EA002129>

Publisher: American Geophysical Union

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Earth and Space Science



COMMENTARY

10.1029/2021EA002129

Key Points:

- Representation uncertainty can be defined as occurring at a comparison interface between two quantities
- A common definition of representation uncertainty can be applied across the Earth Science community
- Collaboration between scientists and metrologists is recommended to define an accessible and relevant vocabulary for uncertainty analysis

Correspondence to:

C. E. Bulgin,
c.e.bulgin@reading.ac.uk

Citation:

Bulgin, C. E., Thomas, C. M., Waller, J. A., & Woolliams, E. R. (2022). Representation uncertainty in the Earth sciences. *Earth and Space Science*, 9, e2021EA002129. <https://doi.org/10.1029/2021EA002129>

Received 12 NOV 2021

Accepted 16 MAY 2022

Author Contributions:

Conceptualization: C. E. Bulgin, C. M. Thomas, J. A. Waller, E. R. Woolliams

Funding acquisition: C. E. Bulgin

Methodology: C. E. Bulgin, C. M. Thomas, J. A. Waller, E. R. Woolliams

Writing – original draft: C. E. Bulgin, C. M. Thomas, J. A. Waller, E. R. Woolliams

Writing – review & editing: C. E. Bulgin, C. M. Thomas, J. A. Waller, E. R. Woolliams

© 2022. The Authors. Earth and Space Science published by Wiley Periodicals LLC on behalf of American Geophysical Union.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Representation Uncertainty in the Earth Sciences

C. E. Bulgin^{1,2} , C. M. Thomas³, J. A. Waller³ , and E. R. Woolliams⁴

¹University of Reading, Reading, UK, ²National Centre for Earth Observation, Leicester, UK, ³Met Office, Exeter, UK, ⁴National Physical Laboratory, Teddington, UK

Abstract The first Joint Workshop on Representation Uncertainty in the Earth Sciences was held in March 2021. This brought together the Earth observation, data assimilation and forecast verification and post-processing communities, alongside metrologists to discuss the definition and quantification of representation uncertainty within the Earth sciences. The aim of the workshop was to facilitate cross-disciplinary discussion, establishing where existing methodologies could be shared and to foster future collaboration. A key outcome of the workshop was a working definition of representation uncertainty applicable across all the Earth science communities, which is presented in this white paper. The cross-disciplinary discussions at the workshop highlighted the need for scientists to work with metrologists to establish a common vocabulary for uncertainties, accessible to Earth science applications. Further recommendations included regular workshops to discuss progress in defining and quantifying representation uncertainty and awareness of cross-disciplinary funding opportunities to further address representation uncertainty issues.

Plain Language Summary This paper describes a workshop which brought together experts from different Earth science disciplines to discuss and attempt to define the term “representation uncertainty”. We make observations of the Earth using satellites, ground based instruments (such as weather stations), air and sea-borne sensors. These observations are used in their own right, and also by computer models to generate weather forecasts. The observations themselves are imperfect and we quantify these imperfections using the term “uncertainty”. In this paper we discuss the uncertainty that occurs when we compare two different sets of observations, two different models, observations and models, or where there are differences in underlying assumptions. As well as the uncertainties inherent in models and observations, there is also an uncertainty due to the fact that the two things being compared are not representing a phenomenon in exactly the same way. For example, a satellite observation may represent an average value over a few hundred square meters, while an instrument on the surface measures only at a single point (typically one-square-meter or less), and the model represents an area of several square kilometres. Understanding those differences is essential to be able to properly combine different sets of observations, and observations with models.

1. Introduction

The term “representation uncertainty” is used in the Earth sciences to describe an uncertainty that occurs at the comparison interface of two different representations of the same physical quantity, although each sub-community has its own ways of characterizing and evaluating these quantities (Bulgin et al., 2016; Janjić et al., 2018). Uncertainty, as defined by the International Vocabulary of Metrology (VIM) (Joint Committee for Guides in Metrology JCGM, 2012) is a measure of the dispersion of the possible values that could be assigned to a measured quantity. Measurement error is the difference between the measured value and the true or reference value, and is unknown. Historically the terms “error” and “uncertainty” have mistakenly been used interchangeably and the need to establish a clear vocabulary to avoid confusion has been advocated in several recent publications (e.g., Merchant et al., 2017; Loew et al., 2017; Mittaz et al., 2019.) and throughout the workshop. Where not explicitly stated, VIM definitions have been used throughout this paper.

The first Joint Workshop on Representation Uncertainty in the Earth Sciences was held online from 23rd to 25th March 2021, sponsored by the National Centre for Earth Observation. This discussion-based workshop was open to scientists working in the fields of Earth observation (EO), data assimilation (DA), forecast verification and post-processing (FVPP), and metrology. The workshop was attended by 72 participants from a range of institutions including universities, the Met Office, the European Center for Medium Wave Forecasting, the UK national metrology institute and the National Oceanic and Atmospheric Administration. The workshop was designed to maximize discussion time, beginning with short presentations on where representation uncertainty might occur

within the Earth sciences and introducing the field of metrology, which underpins the science of uncertainty. Discussion groups were held first within each discipline, and then across the different disciplines, posing the questions “Where are we now (in the context of understanding and quantifying representation uncertainty)?” and “What could we do going forward?” Science talks and poster contributions were made, showcasing ongoing research toward quantifying representation uncertainty. The workshop concluded with a panel discussion summarizing the key outcomes identified within each discipline. The workshop promoted consistent use of terminology across all disciplines to facilitate discussion.

2. Workshop Outcomes

2.1. Defining Representation Uncertainty

One of the main challenges of the meeting was to find a definition of representation uncertainty that was widely applicable across the disciplines represented at the workshop. The “working definition” was regularly revisited throughout the meeting and during the concluding panel discussion, the following definition was proposed:

“Representation uncertainty is associated with a comparison of two quantities (that are themselves uncertain). This representation uncertainty occurs only at the comparison interface.”

Representation uncertainty is recognised as a wider term that includes many different sources of uncertainty, each of which is often described by a more specific term (for example, sampling uncertainty). This broad definition, examples of which are shown in Figure 1, should be applicable across all disciplines, as it allows the detail of how representation uncertainty arises to be different for each comparison, which we acknowledge is often complex.

2.2. Metrology and Uncertainty in Earth Science Applications

Metrology ensures international consistency and century-long stability of measurements for both science and trade by providing traceability to the International System of Units, the SI. More recently, metrologists have also started to apply the principles of metrological traceability to provide confidence in data derived from modeling. Metrological traceability is built on two core concepts: uncertainty analysis, defined by the Guide to the Expression of Uncertainty in Measurement (the GUM; JCGM 100:2008), and comparisons, formalized to validate uncertainty assessments. Metrologists have worked with the EO community for several decades but have had limited interactions with the DA and FVPP communities.

The Earth sciences often have data value chains, where one community's output becomes another community's input. Each stage of the chain, whether based on instruments or modeling, can be described by a “measurement model” that combines quantities from the previous stage's output, and new quantities introduced in this processing step. Additionally, there will be assumptions inherent in the form of the model which Mittaz et al. (2019) described as the “plus zero uncertainty”, written by including a + 0 into an equation form of the measurement model.

Thus there are two types of uncertainty associated with a value calculated by (or processed through) a “measurement model”. First, the propagated uncertainty comes from the input quantities, through the GUM methods, to an uncertainty associated with the calculated value. This describes the dispersion of probable values around the measured (or processed) value that can be reasonably attributed to the target quantity. Then there is the uncertainty associated with the extent to which the measurand defined by the model represents the quantity of interest – the + 0 term. This could include, but is not limited to, representation uncertainty.

2.3. Current Status Within Each Research Discipline

In this section we include some examples of where representation uncertainty occurs within the different disciplines and relate these back to the definition given in Section 2.1, using the shorthand notation (“a”) (“b”) (“c”) and (“d”) to refer to the four example types of representation uncertainty as defined in Figure 1. It should be noted that these types are not always mutually exclusive and a given comparison may include more than one of the examples we provide for illustration.




Figure 1. Schematic detailing examples of the comparison interface at which representation uncertainty occurs. At this interface, the quantity k is the result of a comparison of observed or modeled quantities, x and y . These quantities are transformed through the functions $f(x)$ and $g(y)$ to bring them to the comparison interface. The comparison is performed as some function of the difference between the transformed quantities, producing the result $k = h(f(x) - g(y))$. Both $f(x)$ and $g(y)$ have associated uncertainty and at the point of comparison, there is also the representation uncertainty, identified with the uncertainty on k at the comparison interface. The details of the representation uncertainty are specific to a given comparison, and may arise from (a) comparison of an observation to a geophysical model equivalent, (b) comparison of an observation to a coarser resolution dataset, (c) comparison of two realizations of the same physical quantity or (d) a definitional difference (which could be a comparison of a measured or modeled dataset to a conceptual, more complete dataset or the difference between what users would like and what data producers can provide). The schematics within each box provide just a single example of the possible comparisons.

2.3.1. Earth Observation (Remotely Sensed and In-Situ)

EO includes both in-situ and remote sensing measurements. There are “plus zero uncertainties” identified in EO data production that represent a conceptual comparison between the evaluated quantity and the desired quantity (“d”), or a definitional difference between what users want and what data producers can provide (“d”). Representation uncertainty may arise at the comparison interface between two datasets, for example, ground-based and satellite observations (“b” and/or “c”), where representation uncertainty arises both from making small-to-large area average comparisons, and because in-situ data are typically sparse; hence error statistics from comparisons with satellite data are not necessarily globally representative (despite a global distribution of observations). Error statistics may also mask data complexity (“d”, Povey and Grainger, 2019). Representation uncertainty is also inherent in the extent to which the measured variable matches the target geophysical quantity, either due to imperfections in the measurement equation or user requirements that do not match the “measurement model” (“d”, Stier, 2016).

Considering in-situ measurements (observations performed at the location of the phenomenon), representation uncertainty often occurs when a continuous data field in time or space is inferred from discrete measurements. This uncertainty may change temporally as sampling capacity for global measurements increases (Good, 2016). When filling data gaps to provide spatially complete data fields, representation uncertainty can be introduced

both by the choice of interpolation method (Dodd et al., 2015) and by the extent to which the available data represent areas that are not observed. This is commonly referred to as sampling uncertainty, but falls under the umbrella term of representation uncertainty. Work presented at the workshop highlighted the challenges for data producers in providing an uncertainty budget where the appropriate representation uncertainty has a dependence on how the user wants to use the data: point measurements, spatial or temporal averages, anomalies or trends.

In remote sensing data, representation uncertainty can occur as a result of instrument sampling (“d”), and gridding data (“d” or “b”). For some data streams, a sub-sample of the full resolution satellite data is passed to the ground receiving station introducing sampling uncertainties in the Level 1 (radiance) data (Belward et al., 1994). Representation uncertainty is also common at Level 3 (gridded products) when regularly gridding clear-sky only data; these products will include a sampling uncertainty when compared to all-sky observations (Bulgin et al., 2016). Further sources of representation uncertainty are associated with Level 4 products involving data composites from different instruments with different satellite overpass times (“c”) (Good et al., 2020).

2.3.2. Data Assimilation

Data assimilation is the process by which observations are combined with model data, weighted by their respective uncertainties and accounting for physical constraints, in order to provide an optimized estimate of the modeled state. Often the observations represent different variables from those modeled, so the model space must be mapped into observation space using an observation operator. There are two main comparison interfaces where representation uncertainty may arise: comparison of an observation with its model counterpart (“a”), and, if a processed form of an observation (e.g., a retrieval, super observation, or wind derived from cloud motion) is assimilated, the comparison of this processed observation to the conceptual perfectly processed observation (“d”). Note that the value of the conceptual perfectly processed observation is unlikely to be known since this would require the processing chain, for example, the calculation of a retrieval, to be perfectly known and to introduce no additional uncertainty (Janjić et al., 2018).

There are two main contributors to representation uncertainty at the observation-model interface (“a”). First is the error due to unresolved scales and processes, which arises when the observations represent different spatial and temporal scales than those of the assimilating model. Second is the observation operator error arising when the observation operator is approximated either to reduce computational complexity and cost or because of unknown parameters and processes. At the interface between the observation and its processed form (“d”), representation uncertainty will arise due to the errors introduced and propagated through the processing chain. Observations may also be subject to quality control procedures; inaccuracy or occasional failure of these can be an additional source of uncertainty.

Representation uncertainty is most commonly accounted for in DA through including the representation error covariance matrix, F , along with instrument error covariance matrix, E , in the observation error covariance matrix, $R = E + F$. For this reason, methods to estimate the full R matrix, where the measurement error covariance matrix is known, have been commonly used to isolate representation uncertainty (Desroziers et al. (2005); Hollingsworth and Lönnberg (1986)). Alternatively, individual sources of representation uncertainty can be estimated from the error statistics of a comparison between two values: for example, either two representations of a variable or using two observation operators with differing levels of approximation (Saunders et al., 2018; Schutgens et al., 2016; Waller et al., 2021). Although it is most common for the representation uncertainty to be included in R , work presented at the workshop showed that other methods exist that account for uncertainties via updates in small-scale background uncertainties and model uncertainties (e.g., Bell et al., 2020; Janjić and Cohn, 2006). These approaches highlight the difficulty in separating representation uncertainties from other types of uncertainty inherent in the DA process.

2.3.3. Forecast Verification and Post-Processing

Forecast post-processing attempts to correct, combine and exploit the information contained within existing forecasts to produce optimal products for dissemination to the public and other customers. The related field of forecast verification quantifies the success of a forecast by comparing its predictions to observations independent of those predictions. Both of these fields rely on the comparison of quantities at different spatial and temporal scales, and hence will have representation uncertainties associated with various steps in the processing chain. In addition, post-processing must often provide outputs at different scales depending on the user requirements (“d”). We

note that there are many similarities between the representation uncertainty that arises in DA and FVPP, though they may require different treatment since the two fields have different purposes.

As discussed in the workshop, the representation uncertainty related to the difference between observations based on different sampling (“a”) is an important topic in FVPP; scale mismatch uncertainties can occur due to sub-grid variability that is not modeled. A new probabilistic post-processing system incorporating verification, the Integrated Model post-PROcessing and VERification (Met Office, 2019), considers this representation uncertainty. Roberts et al. (pers. comm.) have demonstrated the importance of accounting for local topography when producing forecasts. Topography may not be well represented in a relatively coarse model, but accounting for this in the post-processing step can improve forecast skill. Adjusting the rate of change of temperature with height can also provide better agreement between locally observed and regionally averaged (modeled) values especially in areas with high relief. Ben Bouallegue et al., 2020 used a statistical parameterization to quantify the representation uncertainty related to the difference between locally observed and regionally averaged values. The results can be used in ensemble verification and to represent sub-grid variability that is not present in the model.

A further example of representation uncertainty in FVPP relates to differences in the spatial position between observations of local weather phenomena and the forecast equivalent. This is in distinction to scale mismatch uncertainties; a meteorological feature (e.g., a cloud) could be modeled to extremely high precision but be located in the incorrect position (“c”, as these are two different realizations of the same physical quantity). This is mitigated in FVPP by using neighborhood methods and forecast ensembles.

2.4. Existing Collaborations Between Disciplines

The Earth sciences are multidisciplinary, with one community's output often being another community's input. While different communities perform their own uncertainty analysis (to differing levels of formality), uncertainties may not be fully transferred and representation uncertainty generated at a comparison interface, may not be fully considered. Projects such as GAIA-CLIM have attempted to address these gaps by bringing the EO, DA and metrology communities together. Some of the residual-based methods commonly used in DA have begun to be used by the EO community. For example, Merchant et al. (2020) applied the diagnostics of Desroziers et al. (2005), along with additional bias correction, to estimate error covariance parameters for SST retrievals.

Another example is the combination of several EO products (remote sensing and/or in-situ) to quantify uncertainties in the Earth's energy and water cycles with inverse modeling methods (L'Ecuyer et al., 2015; Rodell et al., 2015; Thomas et al., 2020). The outputs of the inverse modeling procedure can potentially be used to evaluate the accuracy of Global Climate Model products, providing an opportunity to collaborate with the modeling community. For this evaluation to be effective, the uncertainties, including any representation uncertainties, must be accurately determined.

3. Future Opportunities for Collaboration and Community Requirements

One barrier to effective collaboration on uncertainty in the Earth sciences is communication, particularly where similar words take different meanings in different groups. Integrating metrologists into this multidisciplinary community can help to bridge this gap, clarifying vocabulary and the distinction between terms such as “error” and “uncertainty” (Mittaz et al., 2019). Regular communication on the definition, sources, quantification and mitigation of representation uncertainty will lead to more efficient transfer of information, better inventory of uncertainty (by considering other perspectives) and more traceability. Ideally the various communities will formulate a common language that can be used consistently in published papers. We can draw parallels with the adoption of the notation in Ide et al., 1997 by the DA community.

It was identified in the workshop that collaboration between the DA and FVPP communities could be beneficial, using statistical methods such as the Desroziers et al. (2005) diagnostics to account for temporal uncertainties in FVPP. These uncertainties are important given the serial correlation present in many atmospheric and oceanic phenomena. Furthermore, the acquisition of crowdsourced data (or in-situ measurement) to provide a “ground truth” for forecast verification was discussed. Such data are also of interest to the DA community but in order to properly estimate their uncertainty and use these data effectively, extensive quality control and validation would be required. Another example is the use of dynamical-model-dependent propagation of uncertainties for daily

satellite products. The ability to apply different methods to a single scenario would provide multiple estimates of representation uncertainty and could help provide confidence in those estimates.

The workshop has begun information dissemination between the participants, who represented several Earth science disciplines, and future workshops could benefit from inclusion of model developers/validators, instrument manufacturers, and space agencies. Model developers understand which scales and processes are poorly represented by models and may be able to aid quantification of representation uncertainties that are large during DA. Instrument manufacturers could also create common methods for documenting instrument calibration and data processing procedures to optimize data use including uncertainties, where commercial sensitivities allow.

4. Recommendations

We encourage all the communities represented at the workshop to continue to work toward a mutually agreed understanding of representation uncertainty, starting with the definition presented in this paper, and acknowledging in many cases more specialized terms will be used to characterize the individual component source(s) of uncertainty. Many of the discussions in the workshop highlighted the need to continue this conversation in a multidisciplinary format, perhaps facilitated by regular workshops, expanded to include other relevant communities. In continuing this conversation, the development of a consistent vocabulary on uncertainties that is relevant to the Earth science community is essential. Such a vocabulary would be supported by metrologists to aid communication and underpin future discussions and collaborations. A cross-community paper developing the ideas outlined in this white paper could be the first target for such future collaboration and could define both a formal vocabulary and make specific recommendations for collaborative research. As funding is often a barrier to inter-disciplinary research, the establishment of a community awareness of projects where representation uncertainty issues may be addressed would be beneficial and “sandpit events” (where scientists work together in a focused way to answer a specific challenge) may also be useful to foster in-depth consideration of inter-disciplinary knowledge transfer on representation uncertainty.

Conflict of Interest

The author declare no conflicts of interest relevant to this study.

Acknowledgments

We acknowledge the many excellent contributions from all workshop participants. In particular we would like to thank the panel, the speakers, session chairs and rapporteurs, and the poster authors. We would also like to thank Maurice Cox, John Eyre, Paul Green, Keith Haines, Christopher Merchant, Jonathan Mittaz, Simon Pinnock, Adam Povey, Nigel Roberts and Bruce Wright for their constructive feedback on the white paper draft. The organizing committee would like to thank the National Centre for Earth Observation (NCEO), which sponsored this event, providing technical and organizational support throughout from grant NE/R016518/1 of the Natural Environment Research Council. Emma Woolliams acknowledges funding in part from the Instrument Data quality Evaluation and Assessment Service - Quality Assurance for Earth Observation (IDEAS-QA4EO) contract funded by ESA-ESRIN (n. 4000128960/19/I-NS) and in part from UK Government's Department for Business, Energy and Industrial Strategy (BEIS) through the UK's National Measurement System programmes. © 2021 Crown copyright, Met Office.

Data Availability Statement

This article reports proceedings of a discussion-based online workshop. There are no associated datasets to be made available in conjunction with this commentary.

References

- Bell, Z., Dance, S. L., & Waller, J. A. (2020). Accounting for observation uncertainty and bias due to unresolved scales with the Schmidt-Kalman filter. *Tellus A: Dynamic Meteorology and Oceanography*, 72(1), 1–21. <https://doi.org/10.1080/16000870.2020.1831830>
- Belward, A. S., Kennedy, P. J., & Grégoire, J. M. (1994). The limitations and potential of AVHRR GAC data for continental scale fire studies. *International Journal of Remote Sensing*, 15(11), 2215–2234. <https://doi.org/10.1080/01431169408954239>
- Ben Bouallegue, Z., Haiden, T., Weber, N. J., Hamill, T. M., & Richardson, D. S. (2020). Accounting for representativeness in the verification of ensemble precipitation forecasts. *Monthly Weather Review*, 148(5), 2049–2062. <https://doi.org/10.1175/MWR-D-19-0323.1>
- Bulgín, C. E., Embury, O., & Merchant, C. J. (2016). Sampling uncertainty in gridded sea surface temperature products and advanced very high resolution radiometer (AVHRR) global area coverage (GAC) data. *Remote Sensing of Environment*, 117, 287–294. <https://doi.org/10.1016/j.rse.2016.02.021>
- Desroziers, G., Berre, L., Chapnik, B., & Poli, P. (2005). Diagnosis of observation, background and analysis-error statistics in observation space. *Quarterly Journal of the Royal Meteorological Society*, 131(613), 3385–3396. <https://doi.org/10.1256/qj.05.108>
- Dodd, E. M. A., Merchant, C. J., Rayner, N. A., & Morice, C. P. (2015). An investigation into the impact of using various techniques to estimate Arctic surface air temperature anomalies. *Journal of Climate*, 28(5), 1743–1763. <https://doi.org/10.1175/jcli-d-14-00250.1>
- Good, S. A. (2016). The impact of observational sampling on time series of a 0–700m ocean average temperature: A case study. *International Journal of Climatology*, 37(35), 2260–2268. <https://doi.org/10.1002/joc.4654>
- Good, S. A., Fiedler, E., Mao, C., Martin, M. J., Maycock, A., Ried, R., et al. (2020). The current configuration of the OSTIA system for operational production of foundation sea surface temperature and ice concentration analyses. *Remote Sensing*, 12(4), 720. <https://doi.org/10.3390/rs12040720>
- Hollingsworth, A., & Lönnberg, P. (1986). The statistical structure of short-range forecast errors as determined from radiosonde data. Part I: The wind field. *Tellus*, 38A(2), 111–136. <https://doi.org/10.1111/j.1600-0870.1986.tb00460.x>

- Ide, K., Courtier, P., Ghil, M., & Lorenc, A. C. (1997). Unified notation for data assimilation: Operational, sequential and variational (special issue: Data assimilation in meteorology and oceanography: Theory and practice). *Journal of the Meteorological Society of Japan. Ser. II*, 75(1B), 181–189. https://doi.org/10.2151/jmsj1965.75.1B_181
- Janjić, T., Bormann, N., Bocquet, M., Carton, J. A., Cohn, S. E., Dance, S. L., et al. (2018). On the representation error in data assimilation. *Quarterly Journal of the Royal Meteorological Society*, 144(713), 1257–1278. <https://doi.org/10.1002/qj.3130>
- Janjić, T., & Cohn, S. E. (2006). Treatment of observation error due to unresolved scales in atmospheric data assimilation. *Monthly Weather Review*, 134(10), 2900–2915. <https://doi.org/10.1175/mwr3229.1>
- JCGM 100. (2008). *Guide to the expression of uncertainty in measurement, GUM (sevres, Paris: BIPM) p 100*. Retrieved from www.bipm.org
- JCGM 200. (2012). *International vocabulary of metrology – basic and general concepts and associated terms (VIM)* (3rd ed.). Retrieved from https://www.bipm.org/documents/20126/2071204/JCGM_200_2012.pdf/f0e1ad45-d337-bbeb-53a6-15fe649d0ff1
- L'Ecuyer, T. S., Beaudoin, H. K., Rodell, M., Olson, W., Lin, B., Kato, S., et al. (2015). The observed state of the energy budget in the early twenty-first century. *Journal of Climate*, 28(21), 8319–8346. <https://doi.org/10.1175/JCLI-D-14-00556.1>
- Loew, A., Bell, W., Brocca, L., Bulgín, C. E., Burdanowitz, J., Calbet, X., et al. (2017). Validation practices for satellite-based Earth observation data across communities. *Reviews of Geophysics*, 55(3), 779–817. <https://doi.org/10.1002/2017RG000562>
- Merchant, C. J., Paul, F., Popp, T., Ablain, M., Bontemps, S., Defourny, P., et al. (2017). Uncertainty information in climate data records from Earth observation. *Earth System Science Data*, 9(2), 511–527. <https://doi.org/10.5194/essd-9-511-2017>
- Merchant, C. J., Saux-Picart, S., & Waller, J. (2020). Bias correction and covariance parameters for optimal estimation by exploiting matched in-situ references. *Remote Sensing of Environment*, 237, 111590. <https://doi.org/10.1016/j.rse.2019.111590>
- Met Office. (2019). *Improver: Probabilistic meteorological post-processing*. revision cc87e78d. Retrieved from <https://improver.readthedocs.io>
- Mittaz, J., Merchant, C. J., & Woolliams, E. R. (2019). Applying principles of metrology to historical Earth observations from satellites. *Metrologia*, 56(3), 032002. <https://doi.org/10.1088/1681-7575/ab1705>
- Povey, A. C., & Grainger, R. G. (2019). Toward more representative gridded satellite products. *IEEE Geoscience and Remote Sensing Letters*, 16(5), 672–676. <https://doi.org/10.1109/LGRS.2018.2881762>
- Rodell, M., Beaudoin, H. K., L'Ecuyer, T. S., Olson, W. S., Famiglietti, J. S., Houser, P. R., et al. (2015). The observed state of the water cycle in the early twenty-first century. *Journal of Climate*, 28(21), 8289–8318. <https://doi.org/10.1175/JCLI-D-14-00555.1>
- Saunders, R., Hocking, J., Turner, E., Rayer, P., Rundle, D., Brunel, P., et al. (2018). An update on the RTTOV fast radiative transfer model (currently at version 12). *Geoscientific Model Development*, 11(7), 2717–2737. <https://doi.org/10.5194/gmd-11-2717-2018>
- Schutgens, N. A. J., Gryspeerdt, E., Weigum, N., Tsyro, S., Goto, D., Schulz, M., & Stier, P. (2016). Will a perfect model agree with perfect observations? The impact of spatial sampling. *Atmospheric Chemistry and Physics*, 16(10), 6335–6353. <https://doi.org/10.5194/acp-16-6335-2016>
- Stier, P. (2016). Limitations of passive satellite remote sensing to constrain global cloud condensation nuclei. *Atmospheric Chemistry and Physics*, 16(10), 6595–6607. <https://doi.org/10.5194/acp-16-6595-2016>
- Thomas, C. M., Dong, B., & Haines, K. (2020). Inverse modeling of global and regional energy and water cycle fluxes using Earth observation data. *Journal of Climate*, 33(5), 1707–1723. <https://doi.org/10.1175/JCLI-D-19-0343.1>
- Waller, J. A., Dance, S. L., & Lean, H. W. (2021). Evaluating errors due to unresolved scales in convection permitting numerical weather prediction. *Quarterly Journal of the Royal Meteorological Society*, 147(738), 2657–2669. <https://doi.org/10.1002/qj.4043>