

# *Overcoming the disconnect between energy system and climate modelling*

Article

Accepted Version

Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Craig, Michael T., Wohland, Jan, Stoop, Lauren P., Kies, Alexander, Pickering, Bryn, Bloomfield, Hannah C. ORCID logo ORCID: <https://orcid.org/0000-0002-5616-1503>, Browell, Jethro, De Felice, Matteo, Dent, Chris J., Deroubaix, Adrien, Frischmuth, Felix, Gonzalez, Paula L. M. ORCID logo ORCID: <https://orcid.org/0000-0003-0154-0087>, Grochowicz, Aleksander, Gruber, Katarina, Härtel, Philipp, Kittel, Martin, Kotzur, Leander, Labuhn, Inga, Lundquist, Julie K., Pflugradt, Noah, van der Wiel, Karin, Zeyringer, Marianne and Brayshaw, David J. ORCID logo ORCID: <https://orcid.org/0000-0002-3927-4362> (2022) Overcoming the disconnect between energy system and climate modelling. *Joule*, 6 (7). pp. 1405-1417. ISSN 2542-4351 doi: <https://doi.org/10.1016/j.joule.2022.05.010> Available at <https://centaur.reading.ac.uk/105187/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.joule.2022.05.010>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

## **CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# Overcoming the disconnect between energy system and climate modelling

Michael T. Craig<sup>1†</sup>, Jan Wohland<sup>2,3\*†</sup>, Laurens P. Stoop<sup>4,5,6†</sup>, Alexander Kies<sup>7,27</sup>, Bryn Pickering<sup>8</sup>, Hannah C. Bloomfield<sup>9,10</sup>, Jethro Browell<sup>11</sup>, Matteo De Felice<sup>12</sup>, Chris J. Dent<sup>13,14</sup>, Adrien Deroubaix<sup>15</sup>, Felix Frischmuth<sup>16</sup>, Paula L.M. Gonzalez<sup>10</sup>, Aleksander Grochowicz<sup>17</sup>, Katharina Gruber<sup>18</sup>, Philipp Härtel<sup>16</sup>, Martin Kittel<sup>19</sup>, Leander Kotzur<sup>20</sup>, Inga Labuhn<sup>21</sup>, Julie K. Lundquist<sup>22,23,24</sup>, Noah Pflugradt<sup>25</sup>, Karin van der Wiel<sup>6</sup>, Marianne Zeyringer<sup>26</sup> and David J. Brayshaw<sup>10</sup>

<sup>1</sup>School for Environment and Sustainability, University of Michigan, USA.

<sup>2</sup>Institute for Environmental Decisions, ETH Zurich, Switzerland.

<sup>3\*</sup>Climate Service Center Germany (GERICS),  
Helmholtz-Zentrum Hereon, Germany.

<sup>4</sup>Information and Computing Science, and Copernicus Institute of Sustainable Development, Utrecht University, The Netherlands.

<sup>5</sup>TenneT TSO B.V., Arnhem, The Netherlands.

<sup>6</sup>Royal Netherlands Meteorological Institute, The Netherlands.

<sup>7</sup>Frankfurt Institute for Advanced Studies, Germany.

<sup>27</sup>Department of Electrical and Computer Engineering, Aarhus University, Denmark.

<sup>8</sup>ETH Zürich, Switzerland.

<sup>9</sup>School of Geographical Sciences, University of Bristol, UK.

<sup>10</sup>Department of Meteorology, University of Reading, UK.

<sup>11</sup>University of Glasgow, UK.

<sup>12</sup>European Commission, Joint Research Centre, The Netherlands.

<sup>13</sup>School of Mathematics, University of Edinburgh, UK.

<sup>14</sup>The Alan Turing Institute, UK.

<sup>15</sup>Max Planck Institute, Germany.

<sup>16</sup>Fraunhofer Institute for Energy Economics and Energy System Technology, Germany.

<sup>17</sup>Department of Mathematics, University of Oslo, Norway.

<sup>18</sup>University of Natural Resources and Life Sciences, Austria.

<sup>19</sup>German Institute for Economic Research, Germany.

<sup>20</sup>Institute of Energy and Climate Research, Techno-economic Systems Analysis, Germany.

<sup>21</sup>University of Bremen, Institute of Geography, Germany.

<sup>22</sup>Dept. of Atmospheric and Oceanic Science, University of Colorado Boulder, USA.

<sup>23</sup>National Renewable Energy Laboratory, USA.

<sup>24</sup>Renewable and Sustainable Energy Institute, USA.

<sup>25</sup>Forschungszentrum Jülich, Germany.

<sup>26</sup>Department of Technology Systems, University of Oslo, Norway.

\*Corresponding author(s). E-mail(s): [jan.wohland@hereon.de](mailto:jan.wohland@hereon.de);

†These authors contributed equally to this work.

### Abstract

Energy system models underpin decisions by energy system planners and operators. Energy system modelling faces a transformation: accounting for changing meteorological conditions imposed by climate change. To enable that transformation, a community of practice in energy-climate modelling has started to form that aims to better integrate energy system models with weather and climate models. Here, we evaluate the disconnects between the energy system and climate modelling communities, then lay out a research agenda to bridge those disconnects. In the near-term, we propose interdisciplinary activities for expediting uptake of future climate data in energy system modelling. In the long-term, we propose a transdisciplinary approach to enable development of (1) energy-system-tailored climate datasets for historical and future meteorological conditions and (2) energy system models that can effectively leverage those datasets. This agenda increases the odds of meeting ambitious climate mitigation goals by systematically capturing and mitigating climate risk in energy sector decision making.

## 1 Introduction

Transforming and decarbonising energy systems is necessary to meet ambitious climate change mitigation goals. This monumental task falls on energy system planners and operators around the world, who share the same primary goal:

to provide reliable, resilient, affordable, and clean energy to consumers. System transformation will increasingly couple electric power with other energy sectors [1]. Within individual sectors, system transformation will increasingly couple energy supply and demand with weather and climate. Integrated assessment models provide high-level guidance on the decarbonization needed, but poorly capture meteorological variability [2]. Higher resolution energy models are the key to capturing this variability within a strongly coupled system [3]. The resulting complexity of a strongly coupled system requires increasingly ambitious modelling efforts.

To guide their decisions, energy system planners and operators use energy system models of varying types. As wind and solar power have grown, energy system modelling has undergone a transformation to handle short- and/or long-term meteorological variability and uncertainty [4–6]. This transformation has increased our understanding of how current and future energy systems can be reliable and affordable while relying on variable and uncertain power generation from renewables. Further progress in this area holds one of the keys to successful climate change mitigation [1].

Despite the recent successes in representing the impact of present-day weather and climate, the energy system modelling community faces another transformation: accounting for the non-stationary meteorological conditions imposed by anthropogenic climate change [7–9] and long-term climate variability [10, 11]. An emerging body of literature is highlighting diverse threats that future weather might pose to reliable, resilient, affordable, and clean energy provision. Shifting meteorological and hydrological conditions can affect energy supply, e.g. from renewable [3, 8, 12–25] and thermal power generation [26–28], and energy demand [20, 24, 26, 29–32]. Many of these studies report climate change impacts on the order of  $\pm$  5-10% in long-term averages depending on the studied region, period, and part of the power system analysed [13, 15, 17, 18, 22, 24, 30–32]. Studies also report more pronounced impacts over shorter periods or smaller areas [13, 18, 22, 24, 32]. Although uncertainty surrounds climate change impacts on energy systems, ignoring climate change can be dangerous with respect to extremes and might lead to reduced system reliability [12, 33–35] and resiliency. The potential consequences of these threats are underscored by recent real-world events, like reliability failures during rolling blackouts in California and Texas in 2020 [36] and 2021 [37], respectively, and resiliency failures during wildfires in the Western United States, Australia, and elsewhere [38, 39]. While peer-reviewed climate change attribution studies [40, 41] have not been completed for these events, extreme weather driving these events (e.g., heat and drought) is projected to increase in severity and/or frequency under climate change in many parts of the world [42–45].

In recognition of these threats, a community of practice in energy-climate modelling has started to form that aims to better coordinate two types of models: (1) energy system models and (2) weather and climate models. The

community has formed around annual workshops [25, 46, 47]; monthly webinars [48]; overlapping research teams; and a virtual knowledge sharing platform enabling ongoing exchange about new research, code and data<sup>1</sup>. To expedite the development of this community, this perspective details the gaps between energy system and climate modellers, then suggests near- and long-term actions aimed at closing them. Our suggestions aim to bring the energy and climate modelling communities together to effectively and appropriately use climate information in guiding energy system design and operation.

## 2 The Disconnect between Energy System and Climate Modelling Communities

To highlight the gap between energy system and climate modelling communities, this section first describes the work of an illustrative member of each community. The description of the illustrative member is not meant to describe all members of each community, but is instead intended to provide a simplified representation of a typical member’s scope and priorities.

An illustrative energy system modeller uses one or more types of energy system models [4, 6] to better understand or advance the provision of reliable, affordable, and clean energy. Their energy system model is formulated to inform a particular decision, e.g. as an optimisation model to inform investment [49–51] or operational decisions [51] under various objectives or as a simulation model to investigate system behaviour under certain boundary conditions [52]. Given the scale and complexity of real-world energy systems, their model sacrifices model formulation and spatio-temporal resolution to maintain computational tractability [53–55]. Oftentimes, this means formulating a deterministic problem and simplifying meteorological data across space or time, e.g. by ignoring climate-related uncertainty, modelling a few days per year, assuming perfect foresight, or aggregating sites or time periods [56, 57]. As deterministic models, many energy system models return the optimal planning decision or operational strategy for a given set of meteorological and other inputs.

In selecting meteorological data, our representative energy system modeller looks for five key attributes in a dataset: (1) the relevant spatio-temporal resolution for the studied decision; (2) synchronous with other climate-sensitive inputs, e.g. renewable production synchronous with electricity demand; (3) convenient to process; (4) computationally manageable; and (5) a high-quality representation of meteorological phenomena relevant to energy system and technology performance (see also Figure 1). To satisfy these attributes, the status quo for meteorological data in energy system models is historical data, e.g. from reanalyses [58–63] or reanalysis-derived products (e.g. Renewable.ninja [64, 65], EMHIRES [66, 67], WIND Toolkit [68], NSRDB [69] and the PECD [70, 71]). Historical meteorological datasets provide hourly (or even

---

<sup>1</sup>The platform is hosted on Slack, see [https://join.slack.com/t/nextgenec/shared\\_invite/zt-13ylctgw5-OG7aWUAPQuC7dlRZkUsvvA](https://join.slack.com/t/nextgenec/shared_invite/zt-13ylctgw5-OG7aWUAPQuC7dlRZkUsvvA).

sub-hourly) data, which is the typical temporal resolution at which energy system models run when applied to planning and operational decisions [72]. Spatial resolutions in these datasets differ widely, but are generally on the order of 10 kms. With respect to synchronicity, many historical datasets include wind and solar resource data, which can be paired with historical electricity demand and hydropower time series. To process historical datasets, energy system modellers leverage widely-used, open-source code and/or tools (e.g., ECEM [73], Atlas [74], atlite [75], and SAM [76]) that convert meteorological data, e.g. wind speeds, into energy system model inputs, e.g. wind power output. To maintain computational manageability, they use sub-decadal — often even a single year — of data. Finally, to understand the quality of the dataset, our illustrative energy system modeller might review papers validating and bias-correcting relevant variables from datasets against empirical data [77–81]. They are, however, unlikely to extensively investigate the quality of the meteorological data for their specific purposes, nor to be fully aware of the in-depth meteorological literature assessing the representation of relevant meteorological processes and their vulnerability to climate change (e.g., [82] for a European perspective and [83], [84] and [77] for their connection to energy).

This selection process is much more daunting for meteorological datasets sourced from climate models seeking to represent the future (rather than historical) climate. To understand why, we first turn our attention to understanding the scope and priorities of an illustrative climate modeller.


Our illustrative climate modeller is primarily concerned with understanding and simulating long-term weather and climate change. They use general circulation models (GCMs), which employ a physics-based dynamical mathematical model of the circulation on Earth, to derive long-term climate and weather projections. They might further refine the accuracy and/or resolution of their weather projections by bias correcting and/or downscaling, which use statistical [85] or dynamical [86] models to capture the effect of geography or other factors on weather that GCMs are too coarse to account for. To them, an energy system modeller is often seen as a downstream user of climate data: the climate modeller’s primary focus is to generate high-quality meteorological data, and to provide access to the raw meteorological output for downstream users. As a result, our climate modeller has little knowledge about data needs specific to the energy sector and does not necessarily assess the climate model’s skill in the context of energy system applications. Instead, in generating and publishing data, our climate modeller has several objectives. First, while being interested in how their model performs relative to observations and other models (e.g., [87] and [88]), they likely care more about the model’s performance in purely meteorological-process terms than about energy-relevant surface climate variables (e.g., they may focus on wind-speed at 10 m — as at this level there are surface observations to validate models against — rather than estimating wind-power capacity factors at turbine hub-height). Second, though some surface climate variables (or “Essential Climate Variables”) are extensively evaluated [86] in climate models, that evaluation tends

to focus on static statistical properties (e.g., mean, variance, and probability distributions) of meteorological variables in isolation rather than on time series of and co-variability between energy-system-relevant variables (e.g., in a *kalte dunkelflaute* [89, 90]). Third, given the complex sources of uncertainty affecting climate models (e.g., scenario uncertainty, internal variability, and model error; [91]), our illustrative climate modeller typically views ensembles of many climate models as essential for understanding climate risk in present and future climate. Each simulation produced by them will therefore cover many decades including historical and future conditions. Ultimately, these datasets allow separation of fluctuations induced by climate change from natural climate fluctuations. Thus, overall our illustrative climate modeller might typically seek to draw conclusions from tens of model simulations (from one or more models), each of which will span 30 years or more for a historical and future climate (e.g., [92]).

The expertise, scope, and priorities of our illustrative climate modeller significantly differ from those of our illustrative energy system modeller, leading to a disconnect in information and data flows between the two. As a result, when considering using outputs from our climate modeller, our energy system modeller faces a steep challenge in satisfying their five meteorological data priorities (see Figure 1). This steep challenge could — and likely often does — deter our energy system modeller from engaging with state-of-the-art climate datasets, thus leading to an over-reliance on historical meteorology despite its increasingly poor representation of the future.

With respect to resolution, our illustrative energy system modeller wants hourly data fields at high spatial resolution. Such high-resolution data is rarely if ever available, including in the ongoing CMIP6 High Resolution Model Intercomparison Project [93], which currently provides typically no more than 3-hourly data [92]. Our illustrative energy system modeller also lacks the knowledge, expertise, and/or resources to consider, select, and run appropriate downscaling and bias adjustment methods. Finding synchronous datasets is generally easier than finding datasets with appropriate resolution, as many climate datasets include all types of meteorological variables of interest for energy systems. However, available meteorological variables in climate datasets are often deficient, as discussed below. Additionally, projecting electricity demand and hydropower generation with bottom-up models and climate model outputs is an area of active research. Diverse methods and assumptions exist to model electricity demand, e.g. as a function of surface air temperature, electrification, and societal trends, indicating substantial methodological uncertainty [94]. Providing accurate projections of hydropower generation potential given future meteorology requires complex hydrological modelling, but simpler approaches exist. Whether these simpler approaches are generalizable or remain valid in a changing climate is, however, often unclear. For instance, profiles of hydropower generation potentials have been inferred using runoff in upstream basins [95, 96], since generation potentials have been found to be highly correlated with reservoir inflow [97]. Finding convenient to process





**Fig. 1** Our illustrative energy system and climate modellers have different key priorities, leading to a disconnect (red arrows) between our illustrative energy system modeller needs and climate modeller outputs.

and computationally manageable datasets poses another significant challenge for the energy system modeller, to whom the concept of and processing tools for a meteorological ensemble are largely foreign. Standard climate change datasets of long-term ensembles radically differ from the single annual meteorological time series that our illustrative energy system modeller is trained to use. Finally, to understand whether a climate data set represents relevant meteorological phenomena well, an energy system modeller has two choices: (1) parse through literature, including documentation and journal articles, to understand the model’s skill with respect to each meteorological variable of interest and their co-variability, or (2) assess them themselves. The first choice is often not possible, as such information is typically unavailable and, where available, not quantified based on energy system model needs. But more importantly, our energy system modeller does not have the training or expertise to effectively carry forward either choice. Without familiarity with fundamental concepts in climate science, large parts of the existing literature are effectively useless.

Overall, these perspectives of an illustrative energy and climate modeller highlight several disconnects between energy and climate modelling that currently hinder effective usage of climate information in energy system modelling (see Figure 1). While some of these disconnects relate to data integration challenges (e.g., mismatches in spatio-temporal resolution), others relate to deeper challenges regarding the scope, objectives, expertise, and treatment of meteorological uncertainty embedded within each community. In the next section, we offer near- and long-term actions that can help bridge these disconnects.

### 3 Bridging the Divide through a New Approach to Data and Research

The problems identified above currently prevent the use of the full potential of climate expertise and information in energy system modelling. To address these

problems, we propose a set of near- and long-term interdisciplinary and transdisciplinary activities among the energy and climate modelling communities. In the near-term, our proposed interdisciplinary activities aim to expedite the use of future climate data in energy system modelling, generating much-needed insights for decision-makers. In the long term, our proposed transdisciplinary activities aim to enable two developments: (i) energy-system-tailored climate datasets for historical and future meteorological conditions, and (ii) energy system models that can effectively leverage those datasets. Proposed actions will require reframing and reconsidering methods and processes currently used to create and share data and knowledge between climate and energy modelling communities.

### 3.1 Align Climate Model Outputs with Energy System Model Needs

First, the climate modelling community should align the resolution of their outputs with energy system model needs. In future phases of coordinated climate simulations, climate modellers should output and save the following variables at hourly resolution to match the granularity of detailed energy system models: surface radiation (direct and diffuse); wind speeds and air density at multiple levels between 80 and 200m, including 100 m for intercomparison studies; surface air temperature; surface relative humidity; precipitation; runoff; and evaporation. With these variables, energy system modellers can estimate energy demand and potential electricity generation from wind, solar and hydropower with much greater fidelity (notwithstanding challenges associated with demand and hydropower projections discussed above). We acknowledge that for many current-generation global climate models, the benefits of producing hourly model output may be modest (compared to 3-hourly) but it is likely to increase in future models with higher resolution (as may already be the case for current regional climate models). More importantly, access to hourly surface weather variables from climate models would greatly simplify the pathway to uptake of climate model data by energy-system specialists by (i) improving comparability with historical reanalysis data like ERA5; (ii) easing integration with highly non-linear energy models (e.g., wind power has a cubic dependence on wind speed); and (iii) directly matching the time-granularity of energy system models used in operations and planning. Experience from previous projects, particularly the EU H2020 PRIMAVERA project, suggests that the additional storage costs of producing a small set of *single level* surface weather data are modest compared to, e.g., archival of 3-D cloud physics datasets CFMIP or ISCCP [98] or Lagrangian storm track diagnostics [99]. Moreover, a greater availability of high frequency surface variables would benefit climate impact analyses in energy system modelling and other communities. Adopting hourly surface weather data output in the ‘standard’ CMIP7 protocol would enable uptake in typical energy models and simplify comparison of system outcomes with historical versus future datasets, potentially enabling standardised energy risk assessments from CMIP7 onwards. As

energy systems evolve, the suggested list of variables should be adopted, for instance, to reflect future changes in turbine hub height.

### **3.2 Add Climate-Related Uncertainty Analysis to Default Energy System Modelling Toolbox**

Second, the energy system modelling community should add climate-related uncertainty quantification to their default toolbox. This requires accepting that there is not ‘one’ representative meteorological time series they can use. The use of a single year of meteorological data, whether historical or future, is invalidated by the pronounced inter-annual variability in surface climate [100–102]. Moreover, satellite-era reanalyses and other historical products are too short to capture multidecadal climate variability [11, 103]. Single time series of one or more years carry the risk of missing low probability extreme events. The problem is substantially exacerbated in climate change assessment because multiple potential climate realisations from single- or multi-model ensembles must be considered, along with the role of model error. Diverse sources of uncertainty accumulate in these potential future climates, and the lack of future observation makes validation infeasible by definition.

Several potential strategies exist to include climate-related uncertainty in energy system models. Sensitivity analysis [34, 104], advanced sampling procedures [53–55], and robust decision-making [105–107] can capture climate-related uncertainty without modifying energy system model formulations. Advanced sampling aims to minimise the number of energy system model runs by reducing the ensemble size while still capturing its variability. Conversely, sensitivity analysis and robust decision-making would run energy system models for individual ensemble members to identify trade-offs between objectives, e.g. between cost and reliability. Other strategies like stochastic and robust optimization [35, 108, 109] would embed climate uncertainty within the energy system model through major formulation changes. But these strategies face computational and parameterization challenges, constraining their applicability to real-world systems. With these enhanced uncertainty quantification strategies, the energy system community could begin to use ensembles of climate data [110], so the energy system community should prioritise addressing remaining conceptual and computational challenges. Distributed modelling teams that incorporate experts in uncertainty quantification and climate science could partially alleviate the challenges with implementation of these uncertainty quantification strategies [111].

### **3.3 Strengthen Transdisciplinary Collaborations between Energy System and Climate Modelling Communities**

Third, the energy system and climate modelling communities must engage more strongly in transdisciplinary collaboration (see Figure 2). Using the full potential of climate information to guide climate change mitigation and adaptation will require coordination between the energy and climate modelling


communities, taking into account uncertainties, limitations, and contexts. We acknowledge that climate services contribute to this coordination by seeking to provide useful climate information and context knowledge to impact modellers and policy makers through websites, interactive group activities and focused relationships (e.g., [112, 113]). Although these services aim to overcome the disconnect between users and providers of climate information, substantial issues persist. For instance, experts report that climate services mainly focus on delivering better data rather than advancing the needed research identified in this perspective [114]. In short, there is a need for the two communities to work together rather than merely seeking an efficient interface across which pre-packaged information is passed.

Sequential approaches that divide labour by expertise are problematic because they assume that the climate and energy steps can be separated. For example, when investigating the impacts of extreme events on an energy system, a climate modeller could try to identify the most and least extreme ensemble members which are subsequently used by the energy expert in their modelling. This approach, however, only works if the climate modeller's expertise is sufficient to predict what the energy model will consider extreme [110]. This is unlikely given the complexity of energy models with hundreds of millions of decision variables and interactions between technologies and sectors.

Instead of a sequential approach, we propose an iterative transdisciplinary framework (see Figure 2). In this framework, climate modellers would generate climate ensembles. Climate and energy modellers would then jointly evaluate those ensemble members to identify which ensembles, variables, locations, and meteorological phenomena yield extreme energy system outcomes (e.g., non-served or peak residual demand; [89]). This identification would occur by running each ensemble member through energy system models [110]. Identified variables and phenomena that yield extreme power system outcomes, as opposed to just extreme meteorological conditions, would be fed back to the climate modelling community to enable better skill assessment and targeted model development for energy system applications. Identified variables and phenomena would also guide downscaling activities to produce energy system inputs, which would be fed back into energy system model development.

Research funders can aid this transdisciplinary collaboration if the limitations of traditional paradigms are reflected in their priorities and objectives. Emphasis on enhanced interdisciplinary training and exchange, alongside career support for early- and mid-career researchers, would support those whose research already spans this interdisciplinary space. Moreover, funding agencies should push transdisciplinary research in energy and climate science, particularly where such research challenges the prevailing paradigms. To appropriately assess transdisciplinary proposals, funders should actively mitigate the risk that reviewers approach transdisciplinary work from the perspective of a single discipline by implementing policies that embed transdisciplinary diversity and awareness into the entire review process. Funding opportunities in this area should support fundamental scientific challenges

of translating climate risk into energy systems (e.g., maintaining computational tractability and understanding the roles of, as well as mitigating against, different sources of uncertainty) alongside larger-scale ‘applied’ or ‘solutions-oriented’ programmes



**Fig. 2** Current sequential approach and the proposed transdisciplinary framework for improving coordination between energy system and climate modelling communities.

## 4 Conclusion

The energy system modelling community faces a transformation: accounting for the non-stationary meteorological conditions associated with climate change. Succeeding in this transformation is crucial for future energy systems to provide reliable, affordable, and clean energy under an uncertain future climate. Yet, this transformation is held back by a disconnect between energy and climate modelling communities.

To bridge this disconnect and to overcome the challenges associated with modelling energy systems under climate change, this article explored the underlying drivers, then proposed three inter- and transdisciplinary actions: (1) the climate modelling community should align output variables and their spatio-temporal resolutions with energy system modelling needs; (2) the energy system modelling community should add climate-related uncertainty quantification to their default analytical toolbox; and (3) both communities should engage in a transdisciplinary approach that ensures the development and evaluation of climate information in line with energy sector needs. Ultimately, these actions point to the fact that effective energy system modelling in a

changing climate should not continue to treat climate and energy modelling sequentially. Rather, effective energy system modelling must understand climate and energy modelling as a transdisciplinary interactive endeavour. Given intensifying climate change and rapid decarbonization of energy systems, the time for this endeavour is now.

**Acknowledgments.** The hosting of the NextGenEC workshops – which led to the initiation and progression of this paper – was enabled with the kind support of the University of Reading and the National Centre for Atmospheric Science (e.g., through access to conferencing software), and the initial workshop concept was developed during the PRIMAVERA project (which received funding from the European Union’s Horizon 2020 Research and Innovation Programme under grant agreement no. 641727).

**Funding.** The authors would like to acknowledge the following sources of funding. During large parts of this work, Jan Wohland was funded through an ETH Postdoctoral Fellowship and acknowledged support from the ETH and Uniscientia foundations. Laurens P. Stoop received funding from the Netherlands Organisation for Scientific Research (NWO) under grant number 647.003.005 and is part of the IS-ENES3 project that has received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No. 824084. Chris Dent is part of the projects “Managing Uncertainty in Government Modelling” (sponsored by the Alan Turing Institute) and “Decision support under climate uncertainty for energy security and net zero ” (sponsored by the Alan Turing Institute and EPSRC). Alexander Grochowicz acknowledges UiO:Energi Thematic Research Group Spatial-Temporal Uncertainty in Energy Systems (SPATUS). Katharina Gruber is funded through the reFUEL project, a ERC-grant with No. ERC2017-STG 758149. Philip Härtel is part of the PROGRESS project funded by the Federal Ministry for Economic Affairs and Energy (BMWi.IIC5, funding reference 03EI1027). Julie K. Lundquist has funding provided by the US Department of Energy Office of Energy Efficiency and Renewable Energy Wind Energy Technologies Office. Copyright statement: This work was authored in part by the National Renewable Energy Laboratory, operated by Alliance for Sustainable Energy, LLC, for the U.S. Department of Energy (DOE) under Contract No. DE-AC36-08GO28308. The U.S. Government retains and the publisher, by accepting the article for publication, acknowledges that the U.S. Government retains a nonexclusive, paid-up, irrevocable, worldwide license to publish or reproduce the published form of this work, or allow others to do so, for U.S. Government purposes. David J. Brayshaw was a Co-Principal Investigator on the EU H2020 PRIMAVERA project (grant number 641727) and is the lead-convenor of the Next Generation Challenges in Energy Climate Modelling workshops from which this perspective was developed.

**Authors’ contributions.** Conceptualization: *David J. Brayshaw, Jan Wohland, Michael T. Craig, and Laurens P. Stoop*, Original draft: *Jan Wohland, Michael T. Craig, and Laurens P. Stoop*, Revision, review, and editing

core group: *David J. Brayshaw, Michael T. Craig, Alex Kies, Bryn Pickering, Laurens P. Stoop and Jan Wohland*, Revision and review: *All authors*. All authors (except L. Kotzur) were present at the Next Generation Energy Climate Modelling 2021 workshop that led to the conceptualization of this paper and was organized by David J. Brayshaw. The ordering of the first two authors was decided by a coin toss; these co-first authors can prioritise their names as first authors when adding this paper's reference to their résumés.

## References

- [1] Masson-Delmotte, V., Zhai, P., Pörtner, H.-O., Roberts, D., Skea, J., Shukla, P.R., Pirani, A., Moufouma-Okia, W., Péan, C., Pidcock, R., Connors, S., Matthews, J.B.R., Chen, Y., Zhou, X., Gomis, M.I., Lonnoy, E., Maycock, T., Tignor, M., Waterfield, T. (eds.): *Global Warming of 1.5 C*, (2018). <https://www.ipcc.ch/sr15/>
- [2] Collins, S., Deane, J.P., Poncelet, K., Panos, E., Pietzcker, R.C., Delarue, E., Ó Gallachóir, B.: Integrating short term variations of the power system into integrated energy system models: A methodological review. *Renewable and Sustainable Energy Reviews* (2017). <https://doi.org/10.1016/j.rser.2017.03.090>
- [3] Gernaat, D.E., de Boer, H.S., Daioglou, V., Yalew, S.G., Müller, C., van Vuuren, D.P.: Climate change impacts on renewable energy supply. *Nature Climate Change* (2021). <https://doi.org/10.1038/s41558-020-00949-9>
- [4] Pfenninger, S., Hawkes, A., Keirstead, J.: Energy systems modeling for twenty-first century energy challenges. *Renewable and Sustainable Energy Reviews* (2014). <https://doi.org/10.1016/j.rser.2014.02.003>
- [5] Lund, H., Arler, F., Østergaard, P.A., Hvelplund, F., Connolly, D., Mathiesen, B.V., Karnøe, P.: Simulation versus optimisation: theoretical positions in energy system modelling. *Energies* (2017). <https://doi.org/10.3390/en10070840>
- [6] Ringkjøb, H.-K., Haugan, P.M., Solbrekke, I.M.: A review of modelling tools for energy and electricity systems with large shares of variable renewables. *Renewable and Sustainable Energy Reviews* (2018). <https://doi.org/10.1016/j.rser.2018.08.002>
- [7] Craig, M.T., Cohen, S., Macknick, J., Draxl, C., Guerra, O.J., Sengupta, M., Haupt, S.E., Hodge, B.-M., Brancucci, C.: A review of the potential impacts of climate change on bulk power system planning and operations in the united states. *Renewable and Sustainable Energy Reviews* (2018). <https://doi.org/10.1016/j.rser.2018.09.022>

- [8] Cronin, J., Anandarajah, G., Dessens, O.: Climate change impacts on the energy system: a review of trends and gaps. *Climatic change* (2018). <https://doi.org/10.1007/s10584-018-2265-4>
- [9] Yalaw, S.G., van Vliet, M.T., Gernaat, D.E., Ludwig, F., Miara, A., Park, C., Byers, E., De Cian, E., Piontek, F., Iyer, G., *et al.*: Impacts of climate change on energy systems in global and regional scenarios. *Nature Energy* (2020). <https://doi.org/10.1038/s41560-020-0664-z>
- [10] Zeng, Z., Ziegler, A.D., Searchinger, T., Yang, L., Chen, A., Ju, K., Piao, S., Li, L.Z., Ciais, P., Chen, D., *et al.*: A reversal in global terrestrial stilling and its implications for wind energy production. *Nature Climate Change* (2019). <https://doi.org/10.1038/s41558-019-0622-6>
- [11] Wohland, J., Brayshaw, D., Pfenninger, S.: Mitigating a century of european renewable variability with transmission and informed siting. *Environmental Research Letters* (2021). <https://doi.org/10.1088/1748-9326/abff89>
- [12] Voisin, N., Kintner-Meyer, M., Skaggs, R., Nguyen, T., Wu, D., Dirks, J., Xie, Y., Hejazi, M.: Vulnerability of the us western electric grid to hydro-climatological conditions: How bad can it get? *Energy* (2016). <https://doi.org/10.1016/j.energy.2016.08.059>
- [13] Tobin, I., Jerez, S., Vautard, R., Thais, F., Van Meijgaard, E., Prein, A., Déqué, M., Kotlarski, S., Maule, C.F., Nikulin, G., *et al.*: Climate change impacts on the power generation potential of a european mid-century wind farms scenario. *Environmental Research Letters* (2016). <https://doi.org/10.1088/1748-9326/11/3/034013>
- [14] Reyers, M., Moemken, J., Pinto, J.G.: Future changes of wind energy potentials over europe in a large cmip5 multi-model ensemble. *International Journal of Climatology* (2016). <https://doi.org/10.1002/joc.4382>
- [15] Wohland, J., Reyers, M., Weber, J., Witthaut, D.: More homogeneous wind conditions under strong climate change decrease the potential for inter-state balancing of electricity in europe. *Earth System Dynamics* (2017). <https://doi.org/10.5194/esd-8-1047-2017>
- [16] Grams, C.M., Beerli, R., Pfenninger, S., Staffell, I., Wernli, H.: Balancing europe's wind-power output through spatial deployment informed by weather regimes. *Nature climate change* (2017). <https://doi.org/10.1038/nclimate3338>
- [17] Karauskas, K.B., Lundquist, J.K., Zhang, L.: Southward shift of the global wind energy resource under high carbon dioxide emissions. *Nature*



- Geoscience (2018). <https://doi.org/10.1038/s41561-017-0029-9>
- [18] Tobin, I., Greuell, W., Jerez, S., Ludwig, F., Vautard, R., Van Vliet, M., Breón, F.: Vulnerabilities and resilience of european power generation to 1.5 c, 2 c and 3 c warming. *Environmental Research Letters* (2018). <https://doi.org/10.1088/1748-9326/aab211>
- [19] Jerez, S., Tobin, I., Turco, M., Jiménez-Guerrero, P., Vautard, R., Montávez, J.: Future changes, or lack thereof, in the temporal variability of the combined wind-plus-solar power production in europe. *Renewable Energy* (2019). <https://doi.org/10.1016/j.renene.2019.02.060>
- [20] van Ruijven, B.J., De Cian, E., Wing, I.S.: Amplification of future energy demand growth due to climate change. *Nature communications* (2019). <https://doi.org/10.1038/s41467-019-10399-3>
- [21] Bartok, B., Tobin, I., Vautard, R., Vrac, M., Jin, X., Levvasseur, G., Denvil, S., Dubus, L., Parey, S., Michelangeli, P.-A., *et al.*: A climate projection dataset tailored for the european energy sector. *Climate services* (2019). <https://doi.org/10.1016/j.cliser.2019.100138>
- [22] Pryor, S.C., Barthelmie, R.J., Bukovsky, M.S., Leung, L.R., Sakaguchi, K.: Climate change impacts on wind power generation. *Nature Reviews Earth & Environment* (2020). <https://doi.org/10.1038/s43017-020-0101-7>
- [23] Hou, R., Li, S., Wu, M., Ren, G., Gao, W., Khayatnezhad, M., *et al.*: Assessing of impact climate parameters on the gap between hydropower supply and electricity demand by rcp scenarios and optimized ann by the improved pathfinder (ipf) algorithm. *Energy* (2021). <https://doi.org/10.1016/j.energy.2021.121621>
- [24] Bloomfield, H., Brayshaw, D., Troccoli, A., Goodess, C., De Felice, M., Dubus, L., Bett, P., Saint-Drenan, Y.-M.: Quantifying the sensitivity of european power systems to energy scenarios and climate change projections. *Renewable Energy* (2021). <https://doi.org/10.1016/j.renene.2020.09.125>
- [25] Bloomfield, H., Gonzalez, P., Lundquist, J.K., Stoop, L., Browell, J., Dargaville, R., De Felice, M., Gruber, K., Hilbers, A., Kies, A., *et al.*: The importance of weather and climate to energy systems: a workshop on next generation challenges in energy–climate modeling. *Bulletin of the American Meteorological Society* (2021). <https://doi.org/10.1175/BAMS-D-20-0256.1>
- [26] Van Vliet, M.T., Yearsley, J.R., Ludwig, F., Vögele, S., Lettenmaier,

- D.P., Kabat, P.: Vulnerability of us and european electricity supply to climate change. *Nature Climate Change* (2012). <https://doi.org/10.1038/nclimate1546>
- [27] Loew, A., Jaramillo, P., Zhai, H.: Marginal costs of water savings from cooling system retrofits: a case study for texas power plants. *Environmental Research Letters* (2016). <https://doi.org/10.1088/1748-9326/11/10/104004>
- [28] Miara, A., Macknick, J.E., Vörösmarty, C.J., Tidwell, V.C., Newmark, R., Fekete, B.: Climate and water resource change impacts and adaptation potential for us power supply. *Nature Climate Change* (2017). <https://doi.org/10.1038/nclimate3417>
- [29] Auffhammer, M., Baylis, P., Hausman, C.H.: Climate change is projected to have severe impacts on the frequency and intensity of peak electricity demand across the united states. *Proceedings of the National Academy of Sciences* (2017). <https://doi.org/10.1073/pnas.1613193114>
- [30] Fonseca, F.R., Jaramillo, P., Bergés, M., Severnini, E.: Seasonal effects of climate change on intra-day electricity demand patterns. *Climatic Change* (2019). <https://doi.org/10.1007/s10584-019-02413-w>
- [31] De Cian, E., Wing, I.S.: Global energy consumption in a warming climate. *Environmental and resource economics* (2019). <https://doi.org/10.1007/s10640-017-0198-4>
- [32] Deroubaix, A., Labuhn, I., Camredon, M., Gaubert, B., Monerie, P.-A., Popp, M., Ramarohetra, J., Ruprich-Robert, Y., Silvers, L.G., Siour, G.: Large uncertainties in trends of energy demand for heating and cooling under climate change. *Nature communications* (2021). <https://doi.org/10.1038/s41467-021-25504-8>
- [33] Harang, I., Heymann, F., Stoop, L.P.: Incorporating climate change effects into the european power system adequacy assessment using a post-processing method. *Sustainable Energy, Grids and Networks* (2020). <https://doi.org/10.1016/j.segan.2020.100403>
- [34] Ralston Fonseca, F., Craig, M., Jaramillo, P., Bergés, M., Severnini, E., Loew, A., Zhai, H., Cheng, Y., Nijssen, B., Voisin, N., *et al.*: Climate-induced tradeoffs in planning and operating costs of a regional electricity system. *Environmental Science & Technology* (2021). <https://doi.org/10.1021/acs.est.1c01334>
- [35] Bennett, J.A., Trevisan, C.N., DeCarolis, J.F., Ortiz-García, C., Pérez-Lugo, M., Etienne, B.T., Clarens, A.F.: Extending energy system modelling to include extreme weather risks and application to hurricane

- events in puerto rico. *Nature Energy* (2021). <https://doi.org/10.1038/s41560-020-00758-6>
- [36] California Independent System Operator, California Public Utilities Commission, California Energy Commission: Final: Root cause analysis; mid-august 2020 extreme heat wave. Technical report, California (January 2021). <https://www.caiso.com/Documents/Final-Root-Cause-Analysis-Mid-August-2020-Extreme-Heat-Wave.pdf>
- [37] Mays, J., Craig, M.T., Kiesling, L., Macey, J.C., Shaffer, B., Shu, H.: Private risk and social resilience in liberalized electricity markets. *Joule* (2022). <https://doi.org/10.1016/j.joule.2022.01.004>
- [38] Muhs, J.W., Parvania, M., Shahidehpour, M.: Wildfire risk mitigation: A paradigm shift in power systems planning and operation. *IEEE Open Access Journal of Power and Energy* (2020). <https://doi.org/10.1109/OAJPE.2020.3030023>
- [39] Chiu, B., Roy, R., Tran, T.: Wildfire resiliency: California case for change. *IEEE Power and Energy Magazine* (2022). <https://doi.org/10.1109/MPE.2021.3122730>
- [40] Stott, P.: How climate change affects extreme weather events. *Science* (2016). <https://doi.org/10.1126/science.aaf7271>
- [41] van Oldenborgh, G.J., van der Wiel, K., Kew, S., Philip, S., Otto, F., Vautard, R., King, A., Lott, F., Arrighi, J., Singh, R., *et al.*: Pathways and pitfalls in extreme event attribution. *Climatic Change* (2021). <https://doi.org/10.1007/s10584-021-03071-7>
- [42] Coumou, D., Rahmstorf, S.: A decade of weather extremes. *Nature climate change* (2012). <https://doi.org/10.1038/nclimate1452>
- [43] AghaKouchak, A., Cheng, L., Mazdidasni, O., Farahmand, A.: Global warming and changes in risk of concurrent climate extremes: Insights from the 2014 california drought. *Geophysical Research Letters* (2014). <https://doi.org/10.1002/2014GL062308>
- [44] Zhuang, Y., Fu, R., Santer, B.D., Dickinson, R.E., Hall, A.: Quantifying contributions of natural variability and anthropogenic forcings on increased fire weather risk over the western united states. *Proceedings of the National Academy of Sciences* (2021). <https://doi.org/10.1073/pnas.2111875118>
- [45] Seneviratne, S.I., Zhang, X., Adnan, M., Badi, W., Dereczynski, C., Luca, A.D., Ghosh, S., Iskandar, I., Kossin, J., Lewis, S., *et al.*: Weather and climate extreme events in a changing climate. IPCC (2021)

- [46] Brayshaw, D., Bloomfield, H., Browell, J., Dargaville, R., de Felice, M., Gonzalez, P., Gruber, K., Hilbers, A., Kies, A., Lundquist, J.K., Panteli, M., Price, J., Stoop, L.P., Thornton, H., Wohland, J., Zeyringer, M.: NextGenEC'20. In: Next Generation Challenges in Energy-Climate Modelling (2020). <https://research.reading.ac.uk/met-energy/next-generation-challenges-workshop/outputs/>
- [47] Brayshaw, D., Bloomfield, H., Browell, J., de Felice, M., Gonzalez, P., Gruber, K., Kies, A., Lundquist, J.K., Stoop, L.P., Thornton, H., Wohland, J., Zeyringer, M.: NextGenEC'21. In: Climate and Energy Systems: Foresight Days to Decades Ahead (2021). <https://research.reading.ac.uk/met-energy/next-generation-challenges-workshop/next-generation-energy-climate-modelling-2021/>
- [48] Brayshaw, D., Bloomfield, H., Browell, J., Craig, M., Dargaville, R., Deakin, M., de Felice, M., Gonzalez, P., Grams, C., Greenwood, D., Gruber, K., Hilbers, A., Kies, A., Lundquist, J.K., Panteli, M., Pickering, B., Stoop, L.P., Thornton, H., Victoria, M., van der Wiel, K., Wohland, J., Zeyringer, M.: NextGenEC Webinar series. In: NEXTGENEC COMMUNITY WEBINARS (2022). <https://research.reading.ac.uk/met-energy/next-generation-challenges-workshop/webinar/>
- [49] Pérez-Arriaga, I.J. (ed.): Regulation of the Power Sector. Springer, ??? (2013). <https://doi.org/10.1007/978-1-4471-5034-3>
- [50] Schill, W.-P., Zerrahn, A.: Long-run power storage requirements for high shares of renewables: Results and sensitivities. *Renewable and Sustainable Energy Reviews* (2018). <https://doi.org/10.1016/j.rser.2017.05.205>
- [51] Rubino, A., Sapio, A., La Scala, M. (eds.): *Handbook of Energy Economics and Policy*, (2021). <https://doi.org/10.1016/C2017-0-01718-4>
- [52] Murphy, S., Lavin, L., Apt, J.: Resource adequacy implications of temperature-dependent electric generator availability. *Applied Energy* (2020). <https://doi.org/10.1016/j.apenergy.2019.114424>
- [53] Nahmmacher, P., Schmid, E., Hirth, L., Knopf, B.: Carpe diem: A novel approach to select representative days for long-term power system modeling. *Energy* (2016). <https://doi.org/nahmmacher2016carpe>
- [54] Pfenninger, S.: Dealing with multiple decades of hourly wind and pv time series in energy models: A comparison of methods to reduce time resolution and the planning implications of inter-annual variability. *Applied energy* (2017). <https://doi.org/10.1016/j.apenergy.2017.03.051>
- [55] Hilbers, A.P., Brayshaw, D.J., Gandy, A.: Importance subsampling:

- improving power system planning under climate-based uncertainty. *Applied Energy* (2019). <https://doi.org/10.1016/j.apenergy.2019.04.110>
- [56] Hoffmann, M., Kotzur, L., Stolten, D., Robinius, M.: A review on time series aggregation methods for energy system models. *Energies* (2020). <https://doi.org/10.3390/en13030641>
- [57] Kotzur, L., Nolting, L., Hoffmann, M., Groß, T., Smolenko, A., Priesmann, J., Büsing, H., Beer, R., Kullmann, F., Singh, B., *et al.*: A modeler's guide to handle complexity in energy systems optimization. *Advances in Applied Energy* (2021). <https://doi.org/10.1016/j.adapen.2021.100063>
- [58] Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Bia-vati, G., Bidlot, J., Bonavita, M., De Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R.J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., de Ros-nay, P., Rozum, I., Vamborg, F., Villaume, S., Thépaut, J.-N.: The era5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society* (2020). <https://doi.org/10.1002/qj.3803>
- [59] Gelaro, R., McCarty, W., Suárez, M.J., Todling, R., Molod, A., Takacs, L., Randles, C.A., Darmenov, A., Bosilovich, M.G., Reichle, R., War-gan, K., Coy, L., Cullather, R., Draper, C., Akella, S., Buchard, V., Conaty, A., da Silva, A.M., Gu, W., Kim, G.-K., Koster, R., Lucchesi, R., Merkova, D., Nielsen, J.E., Partyka, G., Pawson, S., Putman, W., Rie-necker, M., Schubert, S.D., Sienkiewicz, M., Zhao, B.: The modern-era retrospective analysis for research and applications, version 2 (merra-2). *Journal of Climate* (2017). <https://doi.org/10.1175/JCLI-D-16-0758.1>
- [60] Ramon, J., Lledó, L., Torralba, V., Soret, A., Doblas-Reyes, F.J.: What global reanalysis best represents near-surface winds? *Quarterly Journal of the Royal Meteorological Society* (2019). <https://doi.org/10.1002/qj.3616>
- [61] Urraca, R., Huld, T., Gracia-Amillo, A., Martinez-de-Pison, F.J., Kaspar, F., Sanz-Garcia, A.: Evaluation of global horizontal irradiance estimates from era5 and cosmo-rea6 reanalyses using ground and satellite-based data. *Solar Energy* (2018). <https://doi.org/10.1016/j.solener.2018.02.059>
- [62] Kaspar, F., Niermann, D., Borsche, M., Fiedler, S., Keller, J., Potthast, R., Rösch, T., Spanghel, T., Tinz, B.: Regional atmospheric reanaly-sis activities at deutscher wetterdienst: review of evaluation results and

- application examples with a focus on renewable energy. *Advances in Science and Research* (2020). <https://doi.org/10.5194/asr-17-115-2020>
- [63] Bloomfield, H.C., Brayshaw, D.J., Gonzalez, P.L.M., Charlton-Perez, A.: Sub-seasonal forecasts of demand and wind power and solar power generation for 28 european countries. *Earth System Science Data* (2021). <https://doi.org/10.5194/essd-13-2259-2021>
- [64] Pfenninger, S., Staffell, I.: Long-term patterns of european pv output using 30 years of validated hourly reanalysis and satellite data. *Energy* (2016). <https://doi.org/10.1016/j.energy.2016.08.060>
- [65] Staffell, I., Pfenninger, S.: Using bias-corrected reanalysis to simulate current and future wind power output. *Energy* (2016). <https://doi.org/10.1016/j.energy.2016.08.068>
- [66] Iratxe, G.A., Zucker, A., Careri, F., Monforti, F., Huld, T., Badger, J.: Emhires dataset part i: Wind power generation. *European Meteorological derived HIGH resolution RES generation time series for present and future scenarios*. (2016). <https://doi.org/10.2790/831549>
- [67] Gonzalez Aparicio, I., Huld, T., Careri, F., Monforti, F., Zucker, A.: Emhires dataset part ii: Solar power generation. *European Meteorological derived HIGH resolution RES generation time series for present and future scenarios*. (2017). <https://doi.org/10.2760/044693>
- [68] Draxl, C., Clifton, A., Hodge, B.-M., McCaa, J.: The wind integration national dataset (wind) toolkit. *Applied Energy* (2015). <https://doi.org/10.1016/j.apenergy.2015.03.121>
- [69] Sengupta, M., Xie, Y., Lopez, A., Habte, A., Maclaurin, G., Shelby, J.: The national solar radiation data base (nsrdb). *Renewable and Sustainable Energy Reviews* (2018). <https://doi.org/10.1016/j.rser.2018.03.003>
- [70] Nuño, E., Maule, P., Hahmann, A., Cutululis, N., Sørensen, P., Karagali, I.: Simulation of transcontinental wind and solar pv generation time series. *Renewable Energy* (2018). <https://doi.org/10.1016/j.renene.2017.11.039>
- [71] De Felice, M.: ENTSO-E Pan-European Climatic Database (PECD 2021.3) in Parquet Format. <https://doi.org/10.5281/zenodo.5780185>
- [72] Deane, J., Drayton, G., Gallachóir, B.Ó.: The impact of sub-hourly modelling in power systems with significant levels of renewable generation. *Applied Energy* (2014). <https://doi.org/10.1016/j.apenergy.2013.07.027>

- [73] Troccoli, A., Goodess, C., Jones, P., Penny, L., Dorling, S., Harpham, C., Dubus, L., Parey, S., Claudel, S., Khong, D.-H., *et al.*: Creating a proof-of-concept climate service to assess future renewable energy mixes in europe: An overview of the c3s ecem project. *Advances in Science and Research* (2018). <https://doi.org/10.5194/asr-15-191-2018>
- [74] Ruohomäki, T., Airaksinen, E., Huuska, P., Kesäniemi, O., Martikka, M., Suomisto, J.: Smart city platform enabling digital twin. In: 2018 International Conference on Intelligent Systems (IS) (2018). <https://doi.org/10.1109/IS.2018.8710517>. IEEE
- [75] Hofmann, F., Hampp, J., Neumann, F., Brown, T., Hörsch, J.: Atlite: a lightweight python package for calculating renewable power potentials and time series. *Journal of Open Source Software* (2021). <https://doi.org/10.21105/joss.03294>
- [76] Blair, N., Diorio, N., Freeman, J., Gilman, P., Janzou, S., Neises, T., Wagner, M.: System advisor model (sam) general description (version 2017.9. 5). National Renewable Energy Laboratory Technical Report (2018)
- [77] Pickering, B., Grams, C.M., Pfenninger, S.: Sub-national variability of wind power generation in complex terrain and its correlation with large-scale meteorology. *Environmental Research Letters* (2020). <https://doi.org/10.1088/1748-9326/ab70bd>
- [78] Cannon, D.J., Brayshaw, D.J., Methven, J., Coker, P.J., Lenaghan, D.: Using reanalysis data to quantify extreme wind power generation statistics: A 33 year case study in great britain. *Renewable Energy* (2015). <https://doi.org/10.1016/j.renene.2014.10.024>
- [79] Habte, A., Sengupta, M., Lopez, A.: Evaluation of the national solar radiation database (nsrdb): 1998-2015. Technical report, National Renewable Energy Lab.(NREL), Golden, CO (United States) (2017). <https://doi.org/10.2172/1351859>
- [80] King, J., Clifton, A., Hodge, B.-M.: Validation of power output for the wind toolkit. Technical report, National Renewable Energy Lab.(NREL), Golden, CO (United States) (2014). <https://doi.org/10.2172/1159354>
- [81] Bloomfield, H.C., Brayshaw, D.J., Gonzalez, P.L., Charlton-Perez, A.: Pattern-based conditioning enhances sub-seasonal prediction skill of european national energy variables. *Meteorological Applications* (2021). <https://doi.org/10.1002/met.2018>
- [82] Woollings, T.: Dynamical influences on european climate: An uncertain future. *Philosophical Transactions of the Royal Society A: Mathematical,*

- Physical and Engineering Sciences (2010). <https://doi.org/10.1098/rsta.2010.0040>
- [83] Ravestein, P., Van der Schrier, G., Haarsma, R., Scheele, R., Van den Broek, M.: Vulnerability of european intermittent renewable energy supply to climate change and climate variability. *Renewable and Sustainable Energy Reviews* (2018). <https://doi.org/10.1016/j.rser.2018.08.057>
- [84] Gonzalez, P.L., Brayshaw, D.J., Zappa, G.: The contribution of north atlantic atmospheric circulation shifts to future wind speed projections for wind power over europe. *Climate Dynamics* (2019). <https://doi.org/10.1007/s00382-019-04776-3>
- [85] Abatzoglou, J.T., Brown, T.J.: A comparison of statistical downscaling methods suited for wildfire applications. *International journal of climatology* (2012). <https://doi.org/10.1002/joc.2312>
- [86] Losada Carreño, I., Craig, M.T., Rossol, M., Ashfaq, M., Batibeniz, F., Haupt, S.E., Draxl, C., Hodge, B.-M., Brancucci, C.: Potential impacts of climate change on wind and solar electricity generation in texas. *Climatic Change* (2020). <https://doi.org/10.1007/s10584-020-02891-3>
- [87] Eyring, V., Cox, P.M., Flato, G.M., Gleckler, P.J., Abramowitz, G., Caldwell, P., Collins, W.D., Gier, B.K., Hall, A.D., Hoffman, F.M., *et al.*: Taking climate model evaluation to the next level. *Nature Climate Change* (2019). <https://doi.org/10.1038/s41558-018-0355-y>
- [88] Hausfather, Z., Drake, H.F., Abbott, T., Schmidt, G.A.: Evaluating the performance of past climate model projections. *Geophysical Research Letters* (2020). <https://doi.org/10.1029/2019GL085378>
- [89] van der Wiel, K., Stoop, L.P., Van Zuijlen, B., Blackport, R., Van den Broek, M., Selten, F.: Meteorological conditions leading to extreme low variable renewable energy production and extreme high energy shortfall. *Renewable and Sustainable Energy Reviews* (2019). <https://doi.org/10.1016/j.rser.2019.04.065>
- [90] Li, B., Basu, S., Watson, S.J., Russchenberg, H.W.: Mesoscale modeling of a “dunkelflaute” event. *Wind Energy* (2021). <https://doi.org/10.1002/we.2554>
- [91] Hawkins, E., Sutton, R.: The potential to narrow uncertainty in regional climate predictions. *Bulletin of the American Meteorological Society* (2009). <https://doi.org/10.1175/2009BAMS2607.1>



- [92] Eyring, V., Bony, S., Meehl, G.A., Senior, C.A., Stevens, B., Stouffer, R.J., Taylor, K.E.: Overview of the coupled model intercomparison project phase 6 (cmip6) experimental design and organization. *Geoscientific Model Development* (2016). <https://doi.org/10.5194/gmd-9-1937-2016>
- [93] Haarsma, R.J., Roberts, M.J., Vidale, P.L., Senior, C.A., Bellucci, A., Bao, Q., Chang, P., Corti, S., Fučkar, N.S., Guemas, V., von Hardenberg, J., Hazeleger, W., Kodama, C., Koenigk, T., Leung, L.R., Lu, J., Luo, J.-J., Mao, J., Mizielinski, M.S., Mizuta, R., Nobre, P., Satoh, M., Scoccimarro, E., Semmler, T., Small, J., von Storch, J.-S.: High resolution model intercomparison project (highresmip v1.0) for cmip6. *Geoscientific Model Development* (2016). <https://doi.org/10.5194/gmd-9-4185-2016>
- [94] Chang, M., Thellufsen, J.Z., Zakeri, B., Pickering, B., Pfenninger, S., Lund, H., Østergaard, P.A.: Trends in tools and approaches for modelling the energy transition. *Applied Energy* (2021). <https://doi.org/10.1016/j.apenergy.2021.116731>
- [95] Tröndle, T., Lilliestam, J., Marelli, S., Pfenninger, S.: Trade-offs between geographic scale, cost, and infrastructure requirements for fully renewable electricity in europe. *Joule* (2020). <https://doi.org/10.1016/j.joule.2020.07.018>
- [96] Gøtske, E.K., Victoria, M.: Future operation of hydropower in europe under high renewable penetration and climate change. *Iscience* (2021). <https://doi.org/10.1016/j.isci.2021.102999>
- [97] Liu, H., Andresen, G.B., Brown, T., Greiner, M.: A high-resolution hydro power time-series model for energy systems analysis: Validated with chinese hydro reservoirs. *MethodsX* (2019). <https://doi.org/10.1016/j.mex.2019.05.024>
- [98] Webb, M.J., Andrews, T., Bodas-Salcedo, A., Bony, S., Bretherton, C.S., Chadwick, R., Chepfer, H., Douville, H., Good, P., Kay, J.E., *et al.*: The cloud feedback model intercomparison project (cfmip) contribution to cmip6. *Geoscientific Model Development* (2017). <https://doi.org/10.5194/gmd-10-359-2017>
- [99] Priestley, M.D., Ackerley, D., Catto, J.L., Hodges, K.I., McDonald, R.E., Lee, R.W.: An overview of the extratropical storm tracks in cmip6 historical simulations. *Journal of Climate* (2020). <https://doi.org/10.1175/JCLI-D-19-0928.1>
- [100] Bloomfield, H.C., Brayshaw, D.J., Shaffrey, L.C., Coker, P.J., Thornton, H.: Quantifying the increasing sensitivity of power systems to climate variability. *Environmental Research Letters* (2016). <https://doi.org/10.1088/1748-9324/11/1/014001>

[1088/1748-9326/11/12/124025](https://doi.org/10.1016/j.joule.2020.08.002)

- [101] Wohland, J., Reyers, M., Märker, C., Witthaut, D.: Natural wind variability triggered drop in german redispatch volume and costs from 2015 to 2016. *PloS one* (2018). <https://doi.org/10.1371/journal.pone.0190707>
- [102] Lombardi, F., Pickering, B., Colombo, E., Pfenninger, S.: Policy decision support for renewables deployment through spatially explicit practically optimal alternatives. *Joule* (2020). <https://doi.org/10.1016/j.joule.2020.08.002>
- [103] Wohland, J., Omrani, N.E., Keenlyside, N., Witthaut, D.: Significant multidecadal variability in german wind energy generation. *Wind Energy Science* (2019). <https://doi.org/10.5194/wes-4-515-2019>
- [104] Schyska, B.U., Kies, A., Schlott, M., von Bremen, L., Medjroubi, W.: The sensitivity of power system expansion models. *Joule* (2021). <https://doi.org/10.1016/j.joule.2021.07.017>
- [105] Weaver, C.P., Lempert, R.J., Brown, C., Hall, J.A., Revell, D., Sarewitz, D.: Improving the contribution of climate model information to decision making: the value and demands of robust decision frameworks. *Wiley Interdisciplinary Reviews: Climate Change* (2013). <https://doi.org/10.1002/wcc.202>
- [106] Shortridge, J.E., Guikema, S.D.: Scenario discovery with multiple criteria: An evaluation of the robust decision-making framework for climate change adaptation. *Risk Analysis* (2016). <https://doi.org/10.1111/risa.12582>
- [107] Lempert, R.J.: Robust decision making (rdm). In: *Decision Making Under Deep Uncertainty*, (2019). [https://doi.org/10.1007/978-3-030-05252-2\\_2](https://doi.org/10.1007/978-3-030-05252-2_2)
- [108] Kazemzadeh, N., Ryan, S.M., Hamzei, M.: Robust optimization vs. stochastic programming incorporating risk measures for unit commitment with uncertain variable renewable generation. *Energy Systems* (2019). <https://doi.org/10.1007/s12667-017-0265-5>
- [109] Perera, A., Nik, V.M., Chen, D., Scartezzini, J.-L., Hong, T.: Quantifying the impacts of climate change and extreme climate events on energy systems. *Nature Energy* (2020). <https://doi.org/10.1038/s41560-020-0558-0>
- [110] van der Wiel, K., Selten, F.M., Bintanja, R., Blackport, R., Screen, J.A.: Ensemble climate-impact modelling: extreme impacts from moderate meteorological conditions. *Environmental Research Letters* (2020).

<https://doi.org/10.1088/1748-9326/ab7668>

- [111] DeCarolis, J.F., Jaramillo, P., Johnson, J.X., McCollum, D.L., Trutnevyyte, E., Daniels, D.C., Akın-Olçum, G., Bergerson, J., Cho, S., Choi, J.-H., *et al.*: Leveraging open-source tools for collaborative macro-energy system modeling efforts. *Joule* (2020). <https://doi.org/10.1016/j.joule.2020.11.002>
- [112] Hewitt, C.D., Stone, R.C., Tait, A.B.: Improving the use of climate information in decision-making. *Nature Climate Change* (2017). <https://doi.org/10.1038/nclimate3378>
- [113] Fischer, L.J., Wernli, H., Bresch, D.N.: Widening the common space to reduce the gap between climate science and decision-making in industry. *Climate Services* (2021). <https://doi.org/10.1016/j.cliser.2021.100237>
- [114] Findlater, K., Webber, S., Kandlikar, M., Donner, S.: Climate services promise better decisions but mainly focus on better data. *Nature Climate Change* (2021). <https://doi.org/10.1038/s41558-021-01125-3>