

The insulin receptor family in the heart: new light on old insights

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open access

Clerk, A. ORCID: <https://orcid.org/0000-0002-5658-0708> and
Sugden, P. H. (2022) The insulin receptor family in the heart:
new light on old insights. Bioscience reports, 42 (7).
BSR20221212. ISSN 1573-4935 doi: 10.1042/BSR20221212
Available at <https://centaur.reading.ac.uk/106012/>

It is advisable to refer to the publisher's version if you intend to cite from the
work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1042/BSR20221212>

Publisher: Portland Press

All outputs in CentAUR are protected by Intellectual Property Rights law,
including copyright law. Copyright and IPR is retained by the creators or other
copyright holders. Terms and conditions for use of this material are defined in
the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Review Article

The insulin receptor family in the heart: new light on old insights

 **Angela Clerk and Peter H. Sugden**

School of Biological Sciences, University of Reading, Reading, U.K.

Correspondence: Angela Clerk (a.clerk@reading.ac.uk)



Insulin was discovered over 100 years ago. Whilst the first half century defined many of the physiological effects of insulin, the second emphasised the mechanisms by which it elicits these effects, implicating a vast array of G proteins and their regulators, lipid and protein kinases and counteracting phosphatases, and more. Potential growth-promoting and protective effects of insulin on the heart emerged from studies of carbohydrate metabolism in the 1960s, but the insulin receptors (and the related receptor for insulin-like growth factors 1 and 2) were not defined until the 1980s. A related third receptor, the insulin receptor-related receptor remained an orphan receptor for many years until it was identified as an alkali-sensor. The mechanisms by which these receptors and the plethora of downstream signalling molecules confer cardioprotection remain elusive. Here, we review important aspects of the effects of the three insulin receptor family members in the heart. Metabolic studies are set in the context of what is now known of insulin receptor family signalling and the role of protein kinase B (PKB or Akt), and the relationship between this and cardiomyocyte survival versus death is discussed. PKB/Akt phosphorylates numerous substrates with potential for cardioprotection in the contractile cardiomyocytes and cardiac non-myocytes. Our overall conclusion is that the effects of insulin on glucose metabolism that were initially identified remain highly pertinent in managing cardiomyocyte energetics and preservation of function. This alone provides a high level of cardioprotection in the face of pathophysiological stressors such as ischaemia and myocardial infarction.

The heart is a sophisticated pump essential for delivery of oxygenated blood and nutrients to the body. It relies on contractile cardiomyocytes, acting in synchrony, to pump blood around the body but, in adult mammals, there is negligible capacity for regeneration or replacement of these cells. Pathophysiological stresses imposed on cardiomyocytes can compromise their contractile ability or cause them to die. This reduces the ability of the heart to function properly and can result in heart failure, an increasing problem in society. Urgent remedies are needed to prevent cardiomyocyte dysfunction and death. Insulin has potent cardioprotective effects, particularly in the context of ischaemia with or without reperfusion, as occurs with myocardial infarction. Insulin acts via its cognate receptor, but there are two related receptors that activate similar signalling pathways. Here, we first provide an overview of the context of cardiac hypertrophy and heart failure, outlining the response of the heart to pathophysiological stresses. We then discuss the effects of insulin and the insulin family of receptors, describing key signalling elements and how these may affect cell survival. Finally, we discuss how insulin receptor family signalling may elicit cardioprotection.

Received: 29 May 2022
 Revised: 20 June 2022
 Accepted: 29 June 2022

Accepted Manuscript online:
 29 June 2022
 Version of Record published:
 18 July 2022

Cardiomyocyte growth and cardiac hypertrophy

Architecturally, the heart is fully formed at birth, but continues to grow through postnatal life, matching the increase in body size through to adulthood. In the embryo/foetus, cardiomyocytes proliferate to

generate sufficient cells for the heart to form, but they become terminally differentiated in the perinatal/postnatal period [1,2]. Subsequent maturational growth of the heart to the adult size requires an increase in size and myofibrillar content of individual cardiomyocytes (maturational growth or eutrophy). The adult heart experiences numerous pathophysiological stresses that may demand an increase in cardiomyocyte contractile function. This is accommodated by cardiomyocyte hypertrophy (i.e. an increase in cell size) with associated cardiac hypertrophy (enlargement of the heart) [3]. Physiological hypertrophy (e.g. as occurs in pregnancy or with endurance exercise) is clearly beneficial, necessary and largely reversible on cessation of the stress [4]. Pathological hypertrophy may develop in diseases such as hypertension which increase cardiac workload, or following myocardial infarction where loss of blood supply to part of the myocardium causes cardiomyocytes to die [5,6]. The resulting deposition of inelastic fibrotic scar tissue increases the workload of surviving cardiomyocytes and these hypertrophy. Such “compensated” cardiac hypertrophy initially facilitates survival, but may decompensate, leading to heart failure. Decompensated hypertrophy is associated with further cardiomyocyte death, with loss of capillaries, inflammation and fibrosis.

Cardiomyocyte growth, whether maturational, physiological or pathological is an anabolic process and, although associated with changes in gene expression, it requires an underpinning increase in protein accretion [7]. Insulin and insulin-like growth factors (IGFs) act through related receptor tyrosine kinases (RTKs), namely insulin receptors (INSRs) and Type 1 IGF1 receptors (IGF1Rs) [8]. As we describe below, both promote protein accumulation in the heart and are implicated in cytoprotection. The role of the third related receptor, the insulin receptor-related receptor (INSRR), identified in 1989 [9], is only now being revealed.

Regulation of cardiac metabolism by insulin and IGFs: impact on protein synthesis

One of the earliest effects of insulin to be established (and possibly the best characterised) was the stimulation of glucose uptake by insulin-sensitive tissues such as the heart [10]. In addition, insulin has a range of other effects on carbohydrate, lipid and protein metabolism: it increases the rate of glycolysis whilst promoting glycogen accumulation, it reduces the rate of lipolysis promoting fatty acid synthesis and decreasing fatty acid oxidation, it increases the rate of protein synthesis and it decreases the rate of protein degradation [11]. In the heart, there has been a focus on the effects of insulin to increase protein synthesis because it is a key feature of cardiac hypertrophy [7]. Thus, protein synthesis (measured by incorporation of the non-metabolised amino-acid phenylalanine) in ribosomes from rat hearts is stimulated by insulin and inhibited by diabetes [12], and cardiac protein synthesis is increased by perfusion of rat hearts with insulin (see, for example, [13,14]). Conversely, cardiac protein synthesis is inhibited by hypoinsulinaemia (diabetes, starvation) both *in vitro* and *in vivo* with loss of cardiac protein mass (see, for example, [14–17]). Analogous results are obtained in perfused hearts or isolated cardiomyocytes treated with IGF1 or IGF2 [18]. Insulin, IGF1 or IGF2 each increase the rate of protein synthesis to the maximum limit of up to ~80%, and the point of action is largely at the level of translational initiation [19]. In concert with its effects to increase the rate of protein synthesis, insulin inhibits global protein degradation [20,21], although the point of action remains ill-defined. Interestingly, degradation of the myofibrillar protein actin is increased by starvation in a manner which is unaffected by insulin [22]. This contrasts with phenylalanine release in the presence of cycloheximide (to inhibit protein synthesis) as a measure of global protein degradation, suggesting that insulin is selective in its effect on protein degradation. In conclusion, in the heart (as in other insulin-responsive tissues), insulin and IGFs act as ‘anabolic’ hormones.

Stimuli other than insulin increase the rate of protein synthesis in the heart and are overtly pro-hypertrophic causing an increase in cardiomyocyte size. These include α_1 -adrenergic receptor (α_1 -AR) agonists, but maximal stimulation of protein synthesis by, for example, the synthetic agonist phenylephrine, is less than that obtained by insulin [23]. Indeed, the only intervention capable of increasing cardiac protein synthesis to a similar degree as insulin or IGFs has so far proved to be exposure to pH_o above the ‘physiological’ value of 7.4, as shown with perfused rat hearts [24]. In contrast with insulin (which does not affect cardiac ‘contractility’), increased pH_o is positively inotropic, suggesting some mechanistic differences. This positive inotropic effect of increased pH_o could involve an effect on pH_i (pH_i is linearly coupled to pH_o with $\Delta pH_i/\Delta pH_o \sim 0.4\text{--}0.5$), since increased pH_i (by increasing the ionisation of the acidic amino acid Ca^{2+} -binding residues) increases Ca^{2+} -binding to troponin C, a key regulator of contraction. These studies were conducted in the 1980s prior to the identification of any of the downstream signalling events associated with insulin receptor signalling. The hypothesis at the time was that understanding the mechanism of action of the effect of increased pH_o on cardiac protein synthesis may assist in understanding the signalling pathways utilised by the insulin receptor, but it took over 20 years to revisit this question.

Insulin and IGF signalling in the heart

INSRs and IGF1Rs: receptors for insulin, IGF1 and IGF2

The general structure of insulin receptor (INSR) was established in 1980 [25,26], with subsequent studies of this and the related Type 1 IGF1 receptor (IGF1R) providing insight into how insulin and IGFs bind to these receptors [27,28]. Important progress has been made in recent years in elucidating the atomic detail of the insulin and IGF1 receptors, along with ligand interactions as reviewed in, for example, [29,30]. Since the molecular detail of the receptors or receptor–ligand interactions is not the focus of this review, we will only present an overview of receptor structure and, to avoid confusion, we will use Online Mendelian Inheritance in Man (OMIM[®]) terminology for the genes/proteins as far as possible. INSRs and IGF1Rs are heterotetrameric receptors, preformed prior to insertion in the plasma membrane (Figure 1A). Each half of the receptor (hemi-receptor) is derived from a single polypeptide. Proteolytic cleavage of an internal cleavage site generates α - and β -subunits held together by a disulphide bridge. Additional proteolysis removes the N-terminal signal peptide. The two extracellular α -subunits are also linked via disulphide bridges to form a stable heterotetrameric $\alpha_2\beta_2$ structure. The β -subunits each contain a single transmembrane domain and an intracellular protein tyrosine kinase. Insulin or IGFs interact with the α -subunits of INSRs or IGF1Rs, causing a conformational change that is transmitted through the transmembrane β -subunits to activate the innate protein tyrosine kinase. The β -chains transphosphorylate Tyr residues in the intracellular kinase domains to activate the kinase fully. Further phosphorylation of additional Tyr residues occurs, and other proteins are recruited to these sites, binding through SH2 or PTB domains (that recognise phospho-Tyr residues with C-terminal or N-terminal specificity determinants, respectively) [31]. Some are adapter proteins that become Tyr-phosphorylated by the receptor. This allows recruitment of further proteins to form a signalling complex or ‘signalosome’.

INSR and IGF1R have high sequence homology in the cytoplasmic tyrosine kinase domain, with greater divergence in the extracellular ligand-binding α -subunits [27,28]. Insulin and IGF1 can each bind to INSRs and IGF1Rs but, at physiological concentrations of circulating hormone, each receptor should only bind its cognate ligand. The mode of ligand binding to each receptor differs. A single molecule of insulin engages with both α -subunits of the INSR heterotetramer, and negative co-operativity between the hemi-receptors appears to reduce affinity for a second molecule [32]. In contrast, there is no co-operativity between IGF1R hemi-receptors, and IGF1R heterotetramers may bind one or two molecules of IGF1 [33,34]. This is important to consider since INSRs and IGF1Rs form hybrid receptors (Figure 1A), potentially a consequence of mass action during the process of receptor assembly [35–37]. These INSR/IGF1R hybrid receptors have similar affinity for IGF1 as IGF1R heterotetramers, and low affinity for insulin [38]. Since the hybrid receptors operate largely as IGF1Rs with respect to ligand binding, this increases the number of available receptors for IGF1 in the heart by almost 2-fold, most likely increasing sensitivity. A question remains as to whether signalling from INSR/IGF1R hybrid receptors is similar to IGF1R heterotetramers, given that downstream engagement of signalling intermediates to INSRs and IGF1Rs may differ, along with activation of protein kinases and effects on cell function [39,40]. Notably, approximately 90% of IGF1Rs in the heart form hybrid receptors with INSRs [41].

IGF2 is often considered to bind simply to the IGF2 receptor (IGF2R) [42]. This is the cation-independent mannose 6-phosphate receptor (CI-M6PR), a large glycoprotein that serves as a clearance receptor for IGF2 and other molecules, targeting them to the lysosome for degradation [43]. However, IGF2 binds with approximately equal affinity as IGF1 to IGF1Rs [44]. Apart from this, there are two isoforms of the INSR (INSR-A and INSR-B) generated by alternative splicing [45]. INSR-B contains an additional 12 residues derived from exon 11, towards the C-terminus of the α subunit and these receptors have high affinity for insulin whilst INSR-A receptors (lacking exon 11) have high affinity for IGF2 [46]. The ratio of INSR-A:INSR-B isoforms in the heart is approximately 1:2 in young, healthy rat hearts, but the proportion of INSR-A isoforms is higher in older animals and in diabetes [47]. Overall, there is potential for adult cardiac cells to respond to IGF2 via INSR-A receptors, IGF1Rs or hybrid INSR/IGF1Rs (Figure 1A). This potentially accounts for the similar effects of IGF1 and IGF2 on protein synthesis in the heart [18]. IGF2 is often considered as a developmental growth factor, but circulating levels of both IGF1 and IGF2 remain high in adults (IGF1: 100–300 ng/ml; IGF2: 400–1100 ng/ml [48,49]). However, the majority of both growth factors is associated with IGF binding proteins (IGFBPs) [50]. Free, unbound growth factors are unstable, having a short half-life (~10 min) and circulating at low levels of 1–2 ng/ml [49]. Release of IGFs from IGFBPs requires proteolysis of the binding proteins [51]. This presumably occurs within target tissues and the reaction could be increased in pathological states. Both IGF1 and IGF2 could also be produced locally in the heart under pathological conditions and, since IGF1 or IGF2 can bind to INSRs with low affinity, a question remains about whether local production of these growth factors may be sufficiently high to trigger INSRs in addition to IGF1Rs. Despite the abundance of both IGF2 and its active receptors, the role of IGF2 in the heart has been generally ignored relative to IGF1 or insulin.




Figure 1. Insulin receptor family signalling

(A) Insulin receptors (INSRs, black) and the related receptor family members, IGF1 receptors (IGF1Rs, blue) and insulin receptor-related receptors (INSRRs, yellow) are expressed at the cell surface as preformed heterotetramers, each comprising two hemi-receptors composed of one α - and one β -subunit. Each α - and β -subunit and the two α -subunits are cross-linked by disulphide bridges. The extracellular α subunits bind insulin or IGFs as indicated or are responsive to alkaline pH_o . The β -subunits have a tyrosine kinase domain (red). Alternative splicing generates two isoforms of INSR, INSR-A and INSR-B, without or with (respectively) exon 11 (pink). Hybrid receptors can form between the family members. IGF2 also binds to the cation-independent mannose 6-phosphate receptor (CI-M6PR) that directs IGF2 and other molecules to the lysosome for degradation. (B) Insulin or IGFs bind to INSRs/IGF1Rs causing receptor phosphorylation and full activation of the tyrosine kinase, with additional phosphorylation of tyrosine residues (pY) on the β -subunit. These recruit adapter proteins including insulin receptor substrates (IRS) that become phosphorylated. IRS proteins recruit phosphoinositide 3' kinase (PI3K) that stimulates production of phosphatidylinositol 3,4,5 tris phosphate in the membrane (red lipids), recruiting PDK1 and protein kinase B (PKB, also known as Akt). PDK1 phosphorylates PKB/Akt on Thr308. In some cells, INSRs/IGF1Rs also signal via Grb2 and the exchange factor Sos to the small G protein Ras. Activated Ras recruits Raf kinases that phosphorylate and activate mitogen-activated protein kinase kinases 1/2 (MKK1/2) that phosphorylate extracellular signal-regulated kinases 1 and 2 (ERK1/2). However, this pathway is not thought to be activated significantly in the heart and is shaded. (C) (i) PKB/Akt(T308) phosphorylates AS160 (Tbc1d4) to increase translocation of the glucose transporter GLUT4 to the membrane for glucose uptake. Phosphorylation and inhibition of glycogen synthase kinase 3 (GSK3) results in activation of glycogen synthase (GS) to increase glycogen synthesis. Other GSK3 substrates influence apoptosis, metabolism and cardiac hypertrophy. (ii) PKB/Akt(T308) activates mammalian target of rapamycin (mTOR) in mTOR complex 1 (mTORC1). Signalling via p70 ribosomal S6 kinase (p70S6K) increases ribosome biogenesis by increasing translation of RNAs with 5' terminal oligopyrimidine (TOP) sequences), phosphorylation of Rps6 and regulation of other eukaryotic initiation factors (eIFs). mTORC1-dependent phosphorylation of 4E-BP causes it to dissociate from eIF4E that is then available to bind to the 5' cap structure of mRNAs to increase translation and enhance protein synthesis. (iii) PKB/Akt(T308) is phosphorylated on Ser473 in mTORC2. Dually phosphorylated PKB/Akt(T308/S473) has greater activity and additional target specificity for forkhead transcription factors of the FoxO family. Phosphorylation of FoxO transcription factors inhibits their effects on gene expression, to influence apoptosis and metabolism.

As mentioned above, INSRs and IGF1Rs become Tyr-phosphorylated on the intracellular β -subunits following activation, recruiting other proteins to the complex. Two key proteins in the insulin signalling system are insulin receptor substrates (IRS), IRS1 and IRS2, large adapter proteins which bind to the receptors via PTB domains [52]. All potential phosphorylation sites and the protein interactome for each of the insulin receptor family β -subunits, IRS1 and IRS2 have been mapped [39]. The precise sites which are used and the proteins that interact presumably depend on protein expression profiles in specific tissues or cell types, in addition to considerations of subcellular localisation and compartmentalisation. INSRs and IGF1Rs signal through IRS1/IRS2 in the heart (as in other tissues) [53], extending the potential cardiac interactome of these receptors. Despite the high homology between the INSR and IGF1R β -subunits, insulin, IGF1 and IGF2 have different effects, both on the intracellular signalling pathways they activate and downstream consequences for the cell [39,40].

Protein kinase signalling from INSRs and IGF1Rs

A full summary the intracellular signalling pathways activated by INSRs and IGF1Rs is beyond the scope of this overview, and readers are referred to other reviews for full details [54–56]. Here, we aim to provide an outline of the signalling with reference to the knowledge of the events in the heart. Two key signalling pathways are potentially activated by the insulin receptor family (Figure 1B): phosphoinositide 3' kinase (PI3K) signalling through protein kinase B (PKB, also known as Akt); and the original mitogen-activated protein kinases (MAPKs), the extracellular signal-regulated kinases 1/2 (ERK1/2).

Signalling through PI3K and PKB/Akt

Activation of PKB/Akt

Of the proteins that bind to the insulin family receptors and IRS1/2, potentially the most prominent are the Type 1 PI3Ks [57,58]. INSRs and IGF1Rs have a direct docking site for PI3K at the C-terminus of the β -subunit (phospho-Y1351/1352), and also recruit PI3Ks via IRS1/2 [39]. PI3K phosphorylates phosphatidylinositol 4,5 bisphosphate (PIP₂) in the membrane to produce phosphatidylinositol 3,4,5 trisphosphate (PIP₃). This recruits the Ser/Thr protein kinases, phosphoinositide-dependent kinase 1 (PDK1) and PKB/Akt, allowing PDK1 to phosphorylate PKB/Akt on Thr308 to activate the kinase. PKB/Akt activates mammalian target of rapamycin (mTOR) within mTOR complex 1 (mTORC1). mTOR also operates in a separate complex, mTORC2, and promotes phosphorylation of PKB/Akt on Ser473. This increases PKB/Akt activity and extends the range of potential substrates [59]. As expected, insulin and IGF1 potently activate PKB/Akt in cardiomyocytes and the heart, with increased phosphorylation of both Thr308 and Ser473 [60–62], indicative of activation of mTORC1 and mTORC2 complexes. PKB/Akt serves as a hub for insulin family receptor signalling in the heart, as in other tissues, orchestrating a co-ordinated anabolic response to insulin associated with cell survival (Figure 1C) [55].

PKB/Akt substrates associated with metabolism and cell survival

Probably the best characterised direct substrates of PKB/Akt are glycogen synthase kinase 3 α and β (GSK3 α/β) that are inhibited by PKB/Akt-mediated phosphorylation of Ser21 and Ser9, respectively. GSK3 α/β phosphorylate various substrates including glycogen synthase, phosphorylation of which by GSK3 α/β is inhibitory [63], the net effect of PKB/Akt being to increase glycogen synthesis (by inhibition of inhibition). Phosphorylation of GSK3 α/β in response to insulin or IGF1 in the heart and cardiomyocytes is well-documented, although the emphasis is generally on its role in cardiac hypertrophy, rather than regulation of glycogen synthesis [64–66]. Overall, GSK3 α/β activity prevents cardiac and cardiomyocyte hypertrophy, acting through a variety of potential mechanisms. These include phosphorylation and inhibition of several transcription factors, prevention of opening of the mitochondrial permeability transition pore (mPTP) and apoptosis and/or phosphorylation and degradation of β -catenin.

PKB/Akt substrates account for other metabolic effects of insulin and IGF1: increased glucose uptake and enhanced glycolytic rate (Figure 2). The increase in glucose uptake induced by insulin is effected by increased translocation of storage vesicles containing GLUT4 glucose transporters to the plasma membrane. This process is regulated by Rab GTPases, inactivation of which is enhanced by Rab GTPase-activating proteins (RabGAPs). PKB/Akt phosphorylates two RabGAPs, Tbc1d1 and Tbc1d4 (more generally known as AS160) [67,68]. This enhances GLUT4 vesicle trafficking, probably by reducing binding of Tbc1d1/4 to Rabs. Insulin promotes phosphorylation of AS160/Tbc1d4 in the heart, and this is involved in translocation of GLUT4 and the fatty acid translocase CD36 to the cell surface [69]. The increase in the rate of glycolysis induced by insulin is mostly related to phosphorylation of the cardiac isoform of 6-phosphofructo-2-kinase/fructose 2,6-bisphosphatase (also known as phosphofructokinase 2; PFKB2) downstream of PDK1 and probably a result of direct phosphorylation by PKB/Akt [70–72]. PFKB2 is a bifunctional enzyme with an N-terminal kinase domain and C terminal phosphatase domain that regulates both production and degradation




Figure 2. Glucose metabolism through the glycolytic and pentose phosphate pathways: regulation by PKB/Akt

PKB/Akt phosphorylates RabGAPs (AS160/Tbc1d4, Tbc1d1) to prevent their inhibitory action (red type), increasing translocation of vesicles containing the glucose transporter GLUT4 to the membrane (blue arrow). This increases glucose uptake into the cell. Hexokinase II (HKII) converts glucose to glucose 6-phosphate (G6P). It is tethered to the mitochondrial membrane, interacting with the voltage-dependent anion channel (VDAC), allowing it to access ATP directly from the mitochondria, and facilitating direct transfer of ADP back into the mitochondria. HKII expression and activity is increased by PKB/Akt signalling (blue type) increasing production of G6P. Glycolysis (centre). G6P is converted to fructose 6-phosphate which is then converted to fructose 1,6-bisphosphate by phosphofructokinase (PFK) in the rate-limiting step of the pathway. Fructose 6-phosphate interconversion with fructose 2,6-bisphosphate is performed by the cardiac isoform of phosphofructokinase 2 (PFKB2), an enzyme with dual kinase (K) and phosphatase (P) activity. PKB/Akt phosphorylates and inhibits the phosphatase activity, resulting in accumulation of fructose 2,6-bisphosphate, an allosteric activator of PFK. This increases glycolytic flux to produce pyruvate for oxidative phosphorylation in the mitochondria. If oxidative phosphorylation is insufficient to manage additional pyruvate, pyruvate is converted to lactate by lactate dehydrogenase (LDH). This causes acidification and excess lactate is exported from the cell. Pentose phosphate pathway (left section). G6P is converted to 6-phospho-gluconolactone by G6P dehydrogenase (G6PDH), relying on production of G6P that is increased by PKB/Akt signalling. This reaction produces NADPH, required to reduce oxidised glutathione (GSSG) to GSH. GSH is a critical antioxidant in the cell, scavenging reactive oxygen species (ROS). 6-phospho-gluconolactone can be returned to the glycolytic pathway via further oxidative and non-oxidative reactions.

of fructose 2,6-bisphosphate (Figure 2). Phosphorylation of the C-terminal phosphatase domain by PKB/Akt selectively inhibits this activity, but the kinase remains active with continued production of fructose 2,6-bisphosphate, an allosteric regulator that promotes glycolysis and inhibits gluconeogenesis by acting on phosphofructokinase 1. PKB/Akt also phosphorylates hexokinase II (HKII) [73], the key enzyme required in insulin-sensitive tissues for production of glucose 6-phosphate in the first step of the glycolytic and pentose phosphate pathways. The latter contributes to the antioxidant capacity of the cell, in addition to provision of nucleosides. However, HKII also operates at the mitochondria to inhibit apoptosis [74], an area of considerable research in the heart. Indeed, studies of HKII in the heart have generally focused on cardioprotection and its role in preventing opening of the mitochondrial permeability transition pore (mPTP) with consequent inhibition of apoptosis [75], rather than its role in metabolism.

Other PKB/Akt substrates associated with cell survival

Apart from effects on metabolism, PKB/Akt phosphorylates a range of substrates with direct and/or indirect effects on cardiac cell survival (Figure 1C).

Endothelial nitric oxide synthase (eNOS) is phosphorylated by PKB/Akt on Ser1177 [76–78]. This residue is generally unphosphorylated in unstimulated cells and phosphorylation of endogenous enzyme increases production of NO [77]. This increased production of NO is associated with vasodilation of the blood vessels, reducing blood pressure and workload on the heart. It also reduces blood vessel reactivity, suppressing inflammation and reducing the potential for platelet activation [79]. Overall, this is expected to enhance cardiomyocyte survival. However, eNOS regulation is complex with an additional inhibitory phosphorylation site (Thr495) that may be constitutive in resting cells, and dephosphorylation of this site is required for activation. Moreover, the enzyme can become uncoupled and, instead of producing NO, eNOS can switch to production of potentially damaging $O_2^{\bullet -}$.

PKB/Akt phosphorylates two proteins in the pathway for interferon production. Here, activation of cGAS (cyclic GMP AMP synthase) by viral infection or DNA from damaged mitochondria leads to production of the second messenger cGAMP (cyclic guanosine monophosphate-adenosine monophosphate) [80,81]. cGAMP activates STING (stimulator of interferon genes) which signals via the protein kinase TBK1 (TANK binding kinase 1) to phosphorylate and activate IRF3 (interferon regulatory factor 3), a key transcription factor required for interferon production. cGAS is phosphorylated on Ser291 by PKB/Akt, reducing its activity and production of cGAMP [80]. PKB/Akt also phosphorylates and inhibits TBK1 on Ser172 [82]. The net effect of PKB/Akt is to reduce interferon production, potentially enhancing cell survival.

Other direct PKB/Akt substrates include Forkhead transcription factors of the FoxO family, phosphorylation of which by PKB/Akt results in their retention in the cytoplasm, reducing expression of pro-apoptotic genes [83,84]. Forkhead transcription factor phosphorylation is increased by insulin or IGF1 in cardiomyocytes and the heart, downstream of PKB/Akt [85] and, although this is associated with a reduction in cardiomyocyte death, FoxO transcription factors also affect cardiac metabolism potentially influencing glucose utilisation [86,87]. Phosphorylation of Forkhead transcription factors appears to be mediated by the dually-phosphorylated form of PKB/Akt(Thr308/Ser473) with a requirement for Ser473 phosphorylation to extend the PKB/Akt substrate profile [59].

PKB/Akt signalling via mTORC1

Other effects of PKB/Akt are mediated by mTORC1, particularly in relation to the regulation of protein synthesis and ribosome biogenesis (Figure 1C) [88–90]. mTORC1 acts in several ways to enhance mRNA translation. A key mediator is p70 ribosomal S6 kinase (p70S6K) which is phosphorylated and activated downstream of mTORC1 in cardiac systems, as in other tissues [91]. p70S6K not only phosphorylates the small ribosomal subunit RPS6 to increase translation but also enhances recruitment of mRNAs with a 5' terminal oligopyrimidine (TOP) sequence to polyribosomes for efficient translation. In cardiomyocytes, 5'-TOP mRNAs regulated by insulin are primarily those encoding the ribosomal subunits [92], so the net effect is to increase the translational machinery. Complementary effects of mTOR are to increase ribosomal RNA synthesis and processing [89]. p70S6K also influences the rate of initiation of translation of mRNAs by phosphorylating a number of eukaryotic initiation factors [90]. A second important group of substrates is the 4E-binding proteins (4E-BPs) which bind to eukaryotic initiation factor 4E (eIF4E). eIF4E interacts directly with the 7-methyl-GTP cap of mRNAs and eIF4E availability is a limiting factor. Phosphorylation of 4E-BPs in cardiomyocytes releases eIF4E, increasing availability of eIF4E for binding to the cap structure [60]. mTORC1 also increases the rate of protein synthesis by enhancing the rate of elongation [91]. Thus, mTORC1 signalling increases both the capacity and rate of protein synthesis in the heart. mTORC1 has a secondary effect to decrease autophagy, inhibiting ULK1 and preventing autophagic destruction of organelles [93]. This may account for the focused mechanism by which insulin inhibits protein degradation in the heart [20]. A final consideration for

mTORC1 is that it increases expression of HKII [73] which (as discussed above) may have a dual effect on metabolism, increasing production of glucose 6-phosphate for use in glycolysis or the pentose phosphate pathway (Figure 2).

Activation of the ERK1/2 cascade

Prior to the identification of PKB/Akt, microtubule-associated protein 2 [MAP2-kinase, later renamed mitogen-activated protein kinase (MAPK)] was identified by the Sturgill group as being associated with insulin signalling in 3T3-L1 adipocytes [94]. This particular 42 kDa protein became known as ERK2, and the related 44 kDa isoform as ERK1. INSRs and IGF1Rs activate the ERK1/2 cascade via guanine nucleotide exchange factors for the small G protein Ras (RasGEFs), including SOS1 and 2 that are recruited to the receptor complex by adapter proteins (e.g. Grb2 [95]) (Figure 1B). RasGEFs promote GTP-loading of Ras, which initiates signalling via RAF kinases. RAF kinases become activated and phosphorylate/activate MAPK kinases 1/2 (MKK1/2) which phosphorylate and activate ERK1/2 [96]. In contrast with their dominant effects on PKB/Akt, insulin and IGF1 have a minor effect on ERK1/2 activation in cardiomyocytes compared with other growth factors or hypertrophic agonists acting through G-protein-coupled receptors (GPCRs) [61]. It is not clear why insulin has a more pronounced effect on ERK1/2 activation in other systems, but it may relate to differentiation status, tissue-specificity or spatial organisation of intracellular signalling components as debated previously [97].

INSRRs: sensing alkaline pH

INSRR structure

INSRRs were identified soon after INSRs and IGF1Rs [9], but remained ‘orphan’ receptors for many years, having no known ligand. INSRRs have high homology with INSRs and IGF1Rs in the tyrosine kinase domain, but the extracellular α -subunit is divergent and does not bind insulin or IGFs (Figure 1A). The molecular phylogeny of the three insulin receptor family members indicates that whereas INSRs and IGF1Rs are expressed in all vertebrates, INSRRs are not found in teleost fish [98]. In addition, exon 11 is conserved between INSRRs and INSR-B receptors, but the alternatively spliced INSR-A receptor (that binds IGF2) appears to be exclusively expressed in mammals. The C-terminus of INSRRs is truncated compared with INSRs and IGF1Rs, producing a smaller β -subunit but, with this exception, the high homology in the intracellular region means that phospho-specific antibodies designed for conserved sites in the β -subunit recognise all three receptors and can be used to assess activation of INSRRs [62]. INSRR mRNA expression is highest in kidney, leading to a supposition that it was a kidney-specific receptor, but it is found at lower levels in other tissues including heart (~20% that of kidney), pancreas and brain [99–101]. The receptors themselves are expressed in specific cells and subdomains, and (for example) INSRRs are found at the basolateral surface of intercalated B cells in kidney [102]. With such tight localisation, overall tissue or cellular expression levels are not necessarily reflective of functional significance. Like IGF1Rs, INSRRs form hybrid receptors with INSRs and these hybrids are not transactivated by insulin [103,104].

Alkali-sensing role of INSRRs

Deyev et al. [105] established that INSRRs operate as alkali sensors: exposure to extracellular alkaline pH_o ($pH_o > 8.0$) causes a conformational change in the α subunit that activates the intracellular tyrosine kinase resulting in Tyr-phosphorylation of the β -subunit with activation of downstream signalling. INSRRs are important in maintaining acid-base balance. Thus, alkalosis in mice *in vivo* activates INSRRs in the kidney and, although INSRR null mice have no overt phenotypic abnormalities [106], they have an impaired renal response to HCO_3^- loading resulting in metabolic alkalosis [105]. The focus on INSRRs has been their function in kidney, but the fact that alkaline pH_o increases cardiac protein synthesis to the same degree as insulin [24] suggests that INSRRs are important in the heart. INSRRs are expressed in the heart, enriched in membranes associated with T-tubules (invaginations in the sarcolemma designed to maximise efficiency of ion-exchange during the cardiac cycle), and are activated in rat hearts perfused at alkaline pH_o [62]. As with IGF1Rs, INSRRs appear to operate in the heart as hybrid receptors with INSRs and/or IGF1Rs (these cannot be easily distinguished, having similar molecular weight), and alkaline pH_o causes Tyr-phosphorylation of both INSRRs and INSRs/IGF1Rs. On the basis of the relative expression levels of the three insulin receptor family members and, with 90% of IGF1Rs forming hybrid receptors with INSRs, it is predicted that virtually all of the INSRRs should be present in cardiomyocytes as INSR/INSRR hybrids. Overall, it is highly probable that INSRRs expressed in T-tubules in the heart respond to alkaline pH_o .

Activation of protein kinase signalling by INSRRs

As might be expected, given the homology between INSRRs and INSRs/IGF1Rs, INSRRs have similar effects on intracellular signalling and, in kidney, there is Tyr-phosphorylation of IRS1 with activation of PKB/Akt [105]. In the heart, perfusion at alkaline pH_o results in phosphorylation of PKB/Akt on Thr308 and Ser473 (indicative of activation of mTORC1 and mTORC2), with associated phosphorylation of GSK3 α/β , p70S6K and RPS6 [62]. The degree of activation is similar to that induced by insulin and presumably accounts for the effects of alkaline pH_o on cardiac protein synthesis [24]. The mechanism for triggering the PKB/Akt signalling pathway is not resolved, since INSRRs lack the direct binding sites for PI3Ks that are found at the C-terminus of INSRs/IGF1Rs [39]. It is also unclear if INSRRs recruit IRS1/2 directly. IRS1/2 bind to INSRs and IGF1Rs close to the transmembrane domain and, although the sequence for IRS1/2 binding is conserved in INSRRs, pull-down experiments failed to identify IRS1/2 binding to the peptide sequence of these receptors [39]. The most likely mechanism is via INSRs/IGF1Rs since INSRRs are almost certainly all expressed as hybrid receptors, and the conformational change induced in the INSRR hemi-receptor is potentially sufficient to trigger activation and transphosphorylation of its INSR/IGF1R partner. In contrast to insulin, perfusion of hearts at alkaline pH_o activates ERK1/2 to a significant degree [62]. This could be a direct INSRR-mediated effect, possibly mediated via the adapter protein Shc which has been detected in the INSRR interactome and can facilitate activation of Ras [39,107]. The site for interaction of Shc with INSRs/IGF1Rs is the same as for IRS1/2, so lack of binding of IRS1/2 may be permissive for Shc binding [39]. However, other stress-responsive MAPKs (JNKs and p38-MAPKs) are activated in hearts perfused at alkaline pH_o [62], so activation of all the MAPKs may be receptor-independent and simply reflect a degree of cellular stress.

Functional role of INSRRs in the heart

INSRRs have an obvious role in kidney as alkali sensors, but the potential role in heart is more obscure given that INSRRs are only activated when exposed to $\text{pH}_o > 8.0$ and blood pH as high as 7.6 is associated with mortality [108]. However, cardiac contraction is regulated by rapid ion fluxes throughout the cardiac cycle and it is essential that ion-balances, including intracellular and extracellular pH, are managed and normalised. INSRRs appear to be concentrated in transverse T tubule membranes [62]. The purpose of T tubules is to facilitate rapid contraction/relaxation, but the small volumes within these invaginations result in ion microdomains [109,110] that may generate a sufficiently high pH_o to activate INSRRs (Figure 3). Alternatively (or additionally), INSRRs (like INSRs or IGF1Rs) may be activated intracellularly. One mechanism may involve oxidative stress which can be generated intracellularly in microdomains by, for example, NADPH oxidases [111]. Oxidative stresses inhibit tyrosine phosphatases, with a net effect to increase Tyr-phosphorylation and activation of insulin family receptors, as occurs with the 'insulinomimetic' combination of H_2O_2 and/or vanadate treatment [112]. Furthermore, there could also be tyrosine phosphorylation of the β subunit by non-receptor tyrosine kinases (e.g. Src family kinases), potentially a consequence of cross-talk from other receptors resulting in transactivation of insulin family receptors [113].

Cross-talk from α_1 -adrenergic receptors to insulin family receptors in the heart

As anabolic hormones, insulin and IGF1 promote cardiac growth, playing a major role in coupling nutritional status to cell size rather than necessarily contributing to cardiac hypertrophy. The most potent hypertrophic agonists for the adult heart (e.g. endothelin-1, α_1 -AR agonists) act through Gq protein-coupled receptors (GqPCRs) [114]. RTKs and GPCRs are generally considered to activate discrete signalling pathways, with RTKs initiating the signal via tyrosine phosphorylation (as described above for the insulin receptor family) whilst GPCRs signal through heterotrimeric G proteins that initiate downstream events via second messenger systems: Gq signalling induces hydrolysis of PIP_2 to produce diacylglycerol and inositol 3,4,5 trisphosphate (Figure 3).

In many systems, there is cross-talk from GPCRs to RTKs, and downstream signalling emanates from RTKs [113]. In the heart, it is particularly well-recognised that epidermal growth factor receptors may be transactivated by different GPCRs [115]. There are also reports of GPCR transactivation of INSRs and IGF1Rs in proliferating cell lines with, for example, activation of INSRs by bradykinin or angiotensin II in INSR-overexpressing cells, possibly acting in a multireceptor complex and/or in association with β -arrestins [116,117], and transactivation of IGF1Rs by endothelin 1 in prostate cancer cells [118]. Recent data extend the concept to the heart, and activation of PKB/Akt by α_1 -AR agonists in perfused hearts or cardiomyocytes is inhibited by linsitinib [62], a highly selective inhibitor of the insulin receptor family tyrosine kinases [119]. Activation of ERK1/2 is also partially suppressed by linsitinib, suggesting that some of the signal is also mediated via one or more of the insulin family receptors. It appears that transactivation of insulin family receptors by α_1 -AR agonists is functionally important since linsitinib inhibits the increase in cell size




Figure 3. Cross-talk from α_1 -adrenergic receptors (α_1 -ARs) to INSRR receptors in the heart: possible mechanisms
(i) α_1 -ARs are G-protein-coupled receptors that couple through $G_{\alpha q}$ to activate phospholipase C (PLC) β and possibly $PLC\epsilon$. $PLC\beta/\epsilon$ hydrolyses phosphatidylinositol 4,5 bisphosphate (PIP_2) to produce diacylglycerol (DAG) which activates protein kinase C (PKC) family members and guanine nucleotide exchange factors for Ras (e.g. $RasGRP$). This results in activation of the $ERK1/2$ cascade that affects downstream signalling and gene expression. (ii) Activation of the small G protein $Rac1$ results in an increase in production of reactive oxygen species (ROS) via NADPH oxidases (NOX). ROS inhibit protein tyrosine phosphatases ($PTPs$), and this may lead to increased tyrosine phosphorylation of $INSRs/INSRRs$, with receptor activation and downstream signalling. (iii) Src family kinases ($SFKs$) may be activated, possibly by phosphatidic acid produced by phospholipase D (activated by PKC ; pathway not shown). $SFKs$ may phosphorylate $INSRs/INSRRs$ directly to cause receptor activation. (iv) α_1 -ARs increase intracellular pH_i via an unknown mechanism and this needs to be neutralised. $SFKs$ phosphorylate anion exchanger 3 ($AE3$) to increase exchange of Cl^- and HCO_3^- delivering HCO_3^- to the extracellular space. Intracellular HCO_3^- is probably generated by carbonic anhydrase II ($CAII$). Simultaneous production of H^+ reduces intracellular pH_i . Extracellular HCO_3^- may increase extracellular pH_o to activate $INSRRs$ via a conformational change of the extracellular α -subunit.

induced in cultured cardiomyocytes by α_1 -AR agonists [62]. Surprisingly, it enhances cardiac hypertrophy induced by α_1 -AR stimulation *in vivo* even in the short term, but this is associated with increased interstitial fibrosis indicative of a switch towards maladaptive hypertrophy predicted to predispose to heart failure. It could be argued that insulin family receptor signalling contributes independently to cardiac hypertrophy induced by α_1 -ARs. However, linsitinib alone has no baseline effect on cardiac function or dimensions, at least over the short term.

The mechanism for activation of insulin family receptors by α_1 -AR agonists remains to be clarified, but there are several possible mechanisms (Figure 3). As mentioned in the previous section in relation to INSRRs, one or more of the receptors may be activated as a consequence of intracellular signalling (rather than extracellular stimulation) that increase Tyr-phosphorylation and activation of the β -subunit. This may result from direct phosphorylation by other tyrosine kinases (e.g. Src family kinases [120]) or could be a consequence of oxidative stress whether resulting from increased flux through the mitochondrial electron transport chain as a consequence of increased contractility or activation of NADPH oxidases [121]. However, INSRRs may be functionally relevant as alkali sensors in the heart, being triggered by increased pH_o in the T-tubules, in which case it is necessary to consider the cellular physiology relating to α_1 -AR stimulation. α_1 -AR agonists increase cardiomyocyte contractility. This is, at least in part, a result of an increase in pH_i of the order of 0.1–0.2 pH unit, which increases myofilament sensitivity to Ca^{2+} , producing an increase in the force of contraction and rate of relaxation [122–125]. This increase in intracellular pH_i requires normalisation and one of the main systems involves anion exchanger 3 (AE3), a $\text{Cl}^-/\text{HCO}_3^-$ exchanger that extrudes HCO_3^- from the cell, with carbonic anhydrases providing a source of HCO_3^- [126,127]. AE3 is expressed in the T tubules in cardiomyocytes [128], and therefore has potential to increase pH_o in close proximity to INSRRs. Interestingly, cardiomyocytes from AE3-/- knockout mice, though smaller than those of wild-type littermates, do not hypertrophy in response to α_1 -AR stimulation with no increase in size, protein synthesis or hypertrophic gene markers [129]. Inhibitors of carbonic anhydrases also abolish α_1 -AR agonist-induced cardiomyocyte hypertrophy and the response is similarly ablated in cardiomyocytes from mice lacking an intracellular isoform of carbonic anhydrase, CAII [130,131]. There is, therefore, potential for INSRRs to become activated as a consequence of increased pH_o in T tubules.

Further research is clearly necessary to address a number of different questions that arise. With respect to α_1 -ARs, it is important to determine which of the insulin family receptors are transactivated in response to α_1 -AR stimulation, and if INSRRs are (as we predict) involved in the response *in vivo*. It is also important to establish if transactivation occurs intracellularly or via an extracellular system. It would be interesting to explore whether other GPCR agonists also transactivate insulin family receptors since, as mentioned above, endothelin-1 and bradykinin can activate INSRRs or IGF1Rs in other cells [116–118], and both have potent effects on cardiomyocyte signalling [132–134]. Apart from situations of receptor cross-talk, it is also worth considering how pathological stresses may influence insulin family receptor signalling. For example, pressure overload enhances insulin receptor signalling [135], and ischaemia/reperfusion (as occurs in myocardial infarction) is associated with activation of PKB/Akt [136]. In the latter case, acidosis that develops on ischaemia is rapidly normalised on reperfusion, in association with various changes in ion fluxes, and elevated oxidative stress [137]. This would be expected to inactivate tyrosine phosphatases, potentially resulting in activation of insulin receptor family members.

Insulin family receptor signalling and cardioprotection: when and how?

Insulin and IGF1 can be cardioprotective: insulin/IGF1 reduce infarct size in *ex vivo* models of myocardial infarction [138–143], and cardiomyocyte-specific expression of IGF1 in mice protects against myocardial infarction *in vivo* [144]. PKB/Akt signalling is particularly associated with cardioprotection [145,146]. Thus, insulin/IGF1 signalling via PKB/Akt promotes cell survival in the heart. However, hyperinsulinemia is not beneficial for the heart, being associated with diabetic cardiomyopathy and obesity-induced heart failure [147]. Moreover, not all studies are fully supportive of the concept that IGF1 is really cardioprotective [148]. Some of the discrepancies may reflect the increased reliance on genetically altered mice to assess the cardioprotective effects or otherwise of IGF1 or the different signalling intermediates and, even if a conditional and inducible system is used for gene manipulation, this results in changes of prolonged duration and abnormal stoichiometry that are not necessarily representative of the endogenous system. It is important to consider not just how insulin receptor family signalling is cardioprotective, but when it may be useful.

When?

Almost all studies of cardioprotection with insulin and IGF1 have used models of myocardial infarction with cardiomyocytes or the heart subjected to hypoxia/ischaemia followed by reoxygenation/reperfusion. In the ischaemic

phase, cardiomyocyte energy production via oxidative phosphorylation in the mitochondria is compromised reducing contractile ability, the cytosol becomes acidified and there is a low level of oxidative stress. Injury is exacerbated on reoxygenation/reperfusion, due in part to a substantial increase in reactive oxygen species [149]. Cytosolic pH recovers, but this is associated with further damage [150]. Cardiomyocytes in the ischaemic zone die via necrosis, but additional apoptotic cell death continues in the peri-infarct zone and even in the remote myocardium, contributing to cardiac dysfunction. Apoptotic cell death occurs largely through the mitochondrial death pathway with opening of the mPTP, triggering activation of the caspase 9 → caspase 3 death cascade [151]. This is an acute stress that requires an initial acute treatment. Points for intervention include maintenance of cardiac energetics, prevention of mitochondrial dysfunction and mitochondrial-induced cell death, reduction of inflammation and reduction of oxidative stress, all of which may be influenced by insulin receptor family signalling in the short term. Another scenario in which the insulin signalling pathway may be beneficial is myocarditis in which activation of PKB/Akt may reduce the degree of inflammation and prevent cardiomyocyte dysfunction or loss. Whether or not the benefits of insulin receptor family signalling may be extrapolated to chronic heart diseases is likely to depend on the cardioprotective mechanism of action of insulin receptor family signalling and the underlying mechanism of disease. Here, a consideration may be that prolonged activation of insulin receptor signalling may be associated with desensitisation of the pathway.

Mechanisms of cardioprotection?

PKB/Akt signalling may directly inhibit mitochondrial-induced cell death of cardiomyocytes and endothelial cells to maintain contraction and cardiac perfusion (Figure 4). As discussed above, direct phosphorylation of FoxO transcription factors by PKB/Akt (Figure 1C) is proposed to reduce synthesis of pro-apoptotic proteins [83–87]. This may provide longer term cardioprotection, but seems unlikely to account for acute protection afforded by insulin in models of myocardial infarction. A second system involves mitochondrial HKII that not only has a role in metabolism but also interacts with the mPTP and prevents channel opening [73]. Phosphorylation of HKII at the mitochondria by PKB/Akt may reduce the potential for apoptosis. In addition, mTORC1 increases HKII expression but, as with FoxO transcription factors, this seems unlikely to confer acute cardioprotection. mTORC1 can prevent autophagy and preserve organelles such as mitochondria in the cell [93]. Theoretically, this could be cardioprotective because potential energy production is preserved. However, autophagy is not necessarily detrimental and may even be beneficial because defective organelles (e.g. damaged mitochondria that may generate excessive reactive oxygen species) are cleared. Indeed, enhancement of autophagic processes is associated with increased longevity [152]. Possibly the strongest evidence for a direct effect of insulin receptor family signalling in preventing cell death relates to GSK3 α/β which, as discussed above, is phosphorylated and inhibited by PKB/Akt [63]. GSK3 α/β are associated with the mitochondria and GSK3 α/β activity increases the probability of mPTP opening and promotes cardiomyocyte apoptosis, so some of the protective effects of PKB/Akt may be via inhibition of GSK3-mediated apoptosis [153].

There is, therefore, evidence that insulin/IGFs may prevent cell death directly, but insulin and IGFs have profound effects on metabolism (Figure 2), and this may account for a large part of the cardioprotection they afford (Figure 4). Indeed, this was the basis for some of the earliest studies of the cardioprotective effects of insulin in the 1960s/1970s, demonstrating that glucose-insulin-potassium (GIK) treatment protects the myocardium from ischaemic damage in a dog model of myocardial infarction [154]. Under oxygenated conditions, cardiac energy production derives largely from fatty acid metabolism and oxidative phosphorylation, with heavy reliance on availability of oxygen. The hypothesis was that reducing oxygen consumption by the heart should limit ischaemic damage resulting from coronary artery occlusion and one way to achieve this is to enhance glycolysis using insulin with provision of additional glucose [155]. Increased glycolysis could potentially compensate for the reduction in mitochondrial ATP production resulting from even 50% oxygenation [156]. We now have some insight into the mechanism of action (PKB/Akt phosphorylation of RabGAPs promotes GLUT4 translocation to the membrane; phosphorylation of PFKB2 increases the rate of glycolysis (Figure 2)), but insulin was already known to increase glucose uptake and glycolytic flux, observations that continue to be upheld [157]. Whilst increasing glucose availability and glycolytic flux may not salvage cardiomyocytes in the full ischaemic zone, it may be sufficient to preserve cardiomyocytes in border zones for damage limitation.

HKII and GSK3 α/β are generally considered to exert their cardioprotective effects by preventing opening of the mPTP and suppressing apoptosis (Figure 4). However, they are key metabolism enzymes. HKII directs glucose into the glycolytic pathway (Figure 2), associating with the mPTP at the mitochondria where the enzyme has ready access to mitochondrial-generated ATP for production of glucose 6-phosphate [158]. It then allows the ADP that is produced to pass straight back into the mitochondrion for oxidative phosphorylation [159]. Glucose 6-phosphate is not just required for glycolysis but is also the first intermediate of the pentose phosphate pathway which produces NADPH required to maintain reduced glutathione levels, a vital component in the antioxidant repertoire of the cell [160]. This pathway is strictly regulated, with the next step of the pathway being highly dependent on the ratio of




Figure 4. Possible mechanisms of cardioprotection resulting from stimulation of insulin receptor family members

Activation of insulin receptor family members results in activation of protein kinase B (PKB, also known as Akt). (i) PKB/Akt signalling influences cell death by reducing mitochondrial-dependent apoptosis, increasing hexokinase II (HKII) and inhibiting glycogen synthase kinase 3 (GSK3) activities. This prevents opening of the mitochondrial permeability transition pore (mPTP). Inhibition of FoxO transcription factors reduces expression of pro-apoptotic proteins. Signalling via mTORC1 inhibits autophagy via Ulk1. Activation of eNOS may reduce cell death, acting to promote vasodilation and, thus, reduce workload. It also reduces vascular inflammation. Phosphorylation of cGAS and STING inhibits interferon production. Increased flux through the pentose phosphate pathway (PPP) increases antioxidant capacity in the cell by increasing production of NADPH and glutathione (GSH). (ii) PKB/Akt increases metabolism, providing the cell with options for reducing oxygen needs whilst sustaining energy production, and maintaining antioxidant capability via the NADPH/glutathione system. Glucose uptake is increased by translocation of GLUT4 to the membrane. HKII increases availability of glucose 6-phosphate (G6P) for both glycolysis and the pentose phosphate pathway (PPP). Phosphorylation of the cardiac isoform of phosphofructokinase 2 (PFKB2) inhibits the phosphatase activity, increasing production of fructose 2,6-bisphosphate, an allosteric activator of glycolysis. Glycogen storage is increased by GSK3 inhibition, providing a source of glucose during ischaemia. FoxO inhibition increases glucose metabolism.

NADPH:NADP⁺: as NADP⁺ rises (as a result of increased oxidative stress in the cell), flow through the pentose phosphate pathway is rapidly increased [161]. Lack of substrate compromises this response, resulting in enhanced oxidative stress and, potentially, causing activation of the mitochondrial death pathway via mPTP opening.

GSK3 α/β phosphorylates and inhibits glycogen synthase (Figure 1C), a metabolic effect that should not be ignored, and PKB/Akt phosphorylation/inhibition of GSK3 α/β has a net effect to increase glycogen levels in the cell [162]. The protective benefit of this is not immediately obvious, but glycogen is an important energy source, being mobilised in ischaemia as a result of increased phosphorylase kinase and glycogen phosphorylase activities [163]. This probably occurs as cytosolic Ca²⁺ concentrations increase and Ca²⁺ binding to the inhibitory δ subunit of phosphorylase kinase causes it to be released from the complex, increasing enzyme activity [164,165]. Phosphorylase kinase phosphorylates and activates glycogen phosphorylase which hydrolyses glycogen to produce glucose 1-phosphate. Glycogen stores provide a glycolytic reserve and potentially protect against cardiac damage in anoxia [166], so at least some of the protective effects associated with loss/inhibition of GSK3 activity, particularly in genetically altered mice, may be associated with increased glycogen stores, as in other models associated with increased glycogen accumulation (e.g. overexpression of AMPK γ [167]).

Insulin receptor family signalling clearly has acute effects on metabolism to provide cardioprotection. However, there are longer term effects on gene expression that also impact on metabolism. First, mTORC1-dependent up-regulation of HKII increases glucose utilisation for both glycolysis and to support antioxidant capacity of the cell via the pentose phosphate pathway [73]. In addition, whilst FoxO transcription factors are recognised as drivers of expression of pro-apoptotic genes, they also to have a significant impact on metabolism, and inhibition of FoxO may support a switch from fatty acid metabolism to glucose utilisation [168]. Such changes are likely to influence cardioprotection in the longer term, potentially during remodelling following myocardial infarction or, possibly, in other cardiac diseases.

Phosphorylation and activation of eNOS by PKB/Akt is often viewed as being cardioprotective (see, for example, [79]). eNOS is expressed at highest levels in endothelial cells, and increased production of NO in blood vessels to promote vasodilation is clearly potentially beneficial because of the reduced workload on the heart (Figure 4). However, in the context of high levels of oxidative stress, NO may combine with O₂^{•-} to produce the peroxynitrite anion (ONOO^{•-}), a highly reactive species associated with nitrosative stress that can cause cell death [169]. Furthermore, if eNOS becomes uncoupled, it may generate O₂^{•-} rather than NO, enhancing oxidative stress. Although eNOS is highly expressed in endothelial cells, it is also expressed in cardiomyocytes, and NO produced by either cell type in the heart may have autocrine/paracrine effects on cardiac function [170]. These effects may be beneficial. For example, most cardiac perfusion occurs during diastole and NO may reduce diastolic stiffening to improve cardiac filling and, therefore, perfusion [171]. However, the data are not always consistent and, as in blood vessels, the effects may depend on the context, particularly with respect to the degree of oxidative stress (which will be high following myocardial infarction) and disease state [170].

Therapeutic strategy for cardioprotection?

It is over 100 years since insulin was discovered and 40 years since the receptor structure was first investigated. The black box from 1979 now contains many signalling intermediates [172] and, though much of what was known about insulin effects in the 1970s is what we know now, we now have an idea of how it happens, even if much still remains to be learned. Irrespective of precise mechanism of action, there are many potential points of intervention in the black box which became the PKB/Akt signalling pathway that may be targeted for cardioprotection. Strategically, targeting an extracellular domain of a receptor is generally an easier option than developing small molecule inhibitors of sufficient specificity or selectivity for intracellular action. One problem for the insulin family receptors has been the pleiotropic effects of insulin and IGF1 throughout the body, so activating INSRs/IGF1Rs to protect the heart is potentially undesirable unless in the very short term. INSR expression is more restricted and the recent understanding of how their conformation is regulated by pH [173–176] will almost certainly provide insight into how they may best be activated. Developing synthetic agonists to target insulin receptor family members is not an easy option, and even a substitute for insulin has proved difficult to develop. However, as in cancer, one therapeutic option might be an activating monoclonal antibody to target INSRs at least one of which has previously been produced [50]. Peptide mimetics or aptamers may prove alternative viable options.

Summary and perspectives

Insulin and IGF1 are anabolic hormones and writing this overview enabled a rediscovery of their metabolic effects. As we describe above, many aspects of PKB/Akt signalling impinge on cardiac metabolism and, in that context, it is

not unreasonable that the major conclusion is that cardioprotection afforded by insulin and IGF1 is largely due to regulation of metabolism. Key effects are the increases in glucose uptake and glycolytic flux, and increasing the rate of glycolysis was suggested as a therapeutic option for improving cardiac function during cardiac anoxia as early as 1975 [156]. Interestingly, given that we now know that INSRRs are alkali sensors, another suggestion from Scheuer in the mid-1970s [156] was that ‘the protection of the ischemic or hypoxic heart by alkalosis may be a feasible approach’, because of the effects on glycolysis [177]. The greatest problem with increasing glycolytic flux in hypoxic conditions as a therapeutic option is probably the associated accumulation of lactic acid and intracellular acidification, but both are normalised on reperfusion/reoxygenation, and any interim damage may be offset by cardiomyocyte preservation. There are clearly other effectors of insulin signalling that may reduce cardiomyocyte or endothelial cell death, thus preserving cardiac function (Figure 4). Here, a consideration is whether preservation of cells that are clearly dysfunctional (or cell death programmes would not have been initiated) is genuinely beneficial in the absence of other systems to salvage and repair the cell. Clearly, indirect effects to reduce cardiac workload by, for example, vasodilation resulting from NO production or reducing inflammation will be beneficial to a heart in jeopardy. Perhaps the combinatorial effects of coupling bona fide protective systems affecting cardiac cell apoptosis with metabolic effects is the reason that insulin receptor family signalling is so potentially cardioprotective.

Competing Interests

The authors declare that there are no competing interests associated with the manuscript.

Funding

The authors declare that there are no sources of funding to be acknowledged.

Open Access

Open access for this article was enabled by the participation of University of Reading in an all-inclusive *Read & Publish* agreement with Portland Press and the Biochemical Society under a transformative agreement with JISC.

CRedit Author Contribution

Angela Clerk: Writing—original draft, Writing—review & editing. **Peter H. Sugden:** Writing—original draft, Writing—review & editing.

Acknowledgements

We thank the many researchers who have contributed to the vast literature on insulin and the insulin receptor family. It has been impossible to provide a full review of all the important studies relating to the physiological effects of insulin receptor family signalling, let alone the downstream signalling events. We have attempted to cite primary references for much of the work but have used many review articles, trying, where possible to cite those from the authors who conducted primary research in the field. We apologise that we could not reasonably cite more. We pay tribute to Alexander Petrenko (Sasha) with whom we planned future studies on INSRRs in the heart but who fell victim to COVID-19 [178]. We are immensely grateful to the authors and journals who have used Open Access for publication of primary research and reviews, in addition to the journals and publishers who made archived material available in electronic format for on-line access. We have drawn on a lifetime of literature, and this availability of information was a critical enabling factor in the development of this review.

Abbreviations

4E-BP, 4E-binding protein; AR, adrenergic receptor; CA, carbonic anhydrase; cGAMP, cyclic guanosine monophosphate-adenosine monophosphate; cGAS, cyclic GMP AMP synthase; eIF4E, eukaryotic initiation factor 4E; eNOS, endothelial nitric oxide synthase; ERK1/2, extracellular signal-regulated kinase 1/2; GAP, GTPase activating protein; GEF, guanine nucleotide exchange factor; GPCR, G-protein-coupled receptor; GSK, glycogen synthase kinase; HK, hexokinase; IGF, insulin-like growth factor; IGF1R, IGF1 receptor; IGFBP, IGF-binding protein; INSR, insulin receptor; INSRR, insulin receptor-related receptor; IRF3, interferon regulatory factor; IRS, insulin receptor substrate; M6PR, mannose 6 phosphate receptor; MAPK, mitogen-activated protein kinase; MKK, MAPK kinase; mPTP, mitochondrial permeability transition pore; mTOR, mammalian target of rapamycin; mTORC, mTOR complex; PDK1, phosphoinositide-dependent kinase 1; PI3K, phosphoinositide 3' kinase; PKB, protein kinase B; RTK, receptor tyrosine kinase; STING, stimulator of interferon genes; TBK1, TANK binding kinase 1.

References

- Ahuja, P., Sdek, P. and MacLellan, W.R. (2007) Cardiac myocyte cell cycle control in development, disease, and regeneration. *Physiol. Rev.* **87**, 521–544, <https://doi.org/10.1152/physrev.00032.2006>
- Broughton, K.M. and Sussman, M.A. (2019) Adult cardiomyocyte cell cycle detour: off-ramp to quiescent destinations. *Trends Endocrinol. Metab.* **30**, 557–567, <https://doi.org/10.1016/j.tem.2019.05.006>
- Dorn, II, G.W., Robbins, J. and Sugden, P.H. (2003) Phenotyping hypertrophy: eschew obfuscation. *Circ. Res.* **92**, 1171–1175, <https://doi.org/10.1161/01.RES.0000077012.11088.BC>
- Gibb, A.A. and Hill, B.G. (2018) Metabolic coordination of physiological and pathological cardiac remodeling. *Circ. Res.* **123**, 107–128, <https://doi.org/10.1161/CIRCRESAHA.118.312017>
- Messerli, F.H., Rimoldi, S.F. and Bangalore, S. (2017) The transition from hypertension to heart failure: contemporary update. *JACC Heart Fail.* **5**, 543–551, <https://doi.org/10.1016/j.jchf.2017.04.012>
- Heusch, G. (2022) Coronary blood flow in heart failure: cause, consequence and bystander. *Basic Res. Cardiol.* **117**, 1, <https://doi.org/10.1007/s00395-022-00909-8>
- Hannan, R.D., Jenkins, A., Jenkins, A.K. and Brandenburger, Y. (2003) Cardiac hypertrophy: a matter of translation. *Clin. Exp. Pharmacol. Physiol.* **30**, 517–527, <https://doi.org/10.1046/j.1440-1681.2003.03873.x>
- Seta, K.A., Kovacina, K.S. and Roth, R.A. (1993) The insulin receptor family. *Adv. Exp. Med. Biol.* **343**, 113–124, https://doi.org/10.1007/978-1-4615-2988-0_12
- Shier, P. and Watt, V.M. (1989) Primary structure of a putative receptor for a ligand of the insulin family. *J. Biol. Chem.* **264**, 14605–14608, [https://doi.org/10.1016/S0021-9258\(18\)63737-8](https://doi.org/10.1016/S0021-9258(18)63737-8)
- Morgan, H.E., Randle, P.J. and Regen, D.M. (1959) Regulation of glucose uptake by muscle. 3. The effects of insulin, anoxia, salicylate and 2:4-dinitrophenol on membrane transport and intracellular phosphorylation of glucose in the isolated rat heart. *Biochem. J.* **73**, 573–579, <https://doi.org/10.1042/bj0730573>
- Dimitriadis, G., Mitrou, P., Lambadiari, V., Maratou, E. and Raptis, S.A. (2011) Insulin effects in muscle and adipose tissue. *Diabetes Res. Clin. Pract.* **93**, S52–S59, [https://doi.org/10.1016/S0168-8227\(11\)70014-6](https://doi.org/10.1016/S0168-8227(11)70014-6)
- Rampersad, O.R. and Wool, I.G. (1965) Protein synthesis by ribosomes from heart muscle: effect of insulin and diabetes. *Science* **149**, 1102–1103, <https://doi.org/10.1126/science.149.3688.1102>
- Morgan, H.E., Jefferson, L.S., Wolpert, E.B. and Rannels, D.E. (1971) Regulation of protein synthesis in heart muscle. II. Effects of amino acid levels and insulin on ribosomal aggregation. *J. Biol. Chem.* **246**, 2163–2170, [https://doi.org/10.1016/S0021-9258\(19\)77203-2](https://doi.org/10.1016/S0021-9258(19)77203-2)
- Smith, D.M., Fuller, S.J. and Sugden, P.H. (1986) The effects of lactate, acetate, glucose, insulin, starvation and alloxan-diabetes on protein synthesis in perfused rat hearts. *Biochem. J.* **236**, 543–547, <https://doi.org/10.1042/bj2360543>
- Williams, I.H., Chua, B.H.L., Sahms, R.H., Siehl, D.L. and Morgan, H.E. (1980) Effects of diabetes on protein turnover in cardiac muscle. *Am. J. Physiol.* **239**, E178–E185, <https://doi.org/10.1152/ajpendo.1980.239.3.E178>
- Preedy, V.R., Smith, D.M., Kearney, N.F. and Sugden, P.H. (1984) Rates of protein turnover in vivo and in vitro in ventricular muscle of hearts from fed and starved rats. *Biochem. J.* **222**, 395–400, <https://doi.org/10.1042/bj2220395>
- Preedy, V.R., Paska, L., Sugden, P.H., Schofield, P.S. and Sugden, M.C. (1988) The effects of surgical stress and short-term fasting on protein synthesis in diverse tissues of the mature rat. *Biochem. J.* **250**, 179–188, <https://doi.org/10.1042/bj2500179>
- Fuller, S.J., Mynett, J.R. and Sugden, P.H. (1992) Stimulation of cardiac protein synthesis by insulin-like growth factors. *Biochem. J.* **282**, 85–90, <https://doi.org/10.1042/bj2820085>
- Flaim, K.E., Kochel, P.J., Kira, Y., Kobayashi, K., Fossel, E.T., Jefferson, L.S. et al. (1983) Insulin effects on protein synthesis are independent of glucose and energy metabolism. *Am. J. Physiol.* **245**, C133–C143, <https://doi.org/10.1152/ajpcell.1983.245.1.C133>
- Rannels, D.E., Kao, R. and Morgan, H.E. (1975) Effect of insulin on protein turnover in heart muscle. *J. Biol. Chem.* **250**, 1694–1701, [https://doi.org/10.1016/S0021-9258\(19\)41749-3](https://doi.org/10.1016/S0021-9258(19)41749-3)
- Sugden, P.H. and Smith, D.M. (1982) The effects of glucose, acetate, lactate and insulin on protein degradation in the perfused rat heart. *Biochem. J.* **206**, 467–472, <https://doi.org/10.1042/bj2060467>
- Smith, D.M. and Sugden, P.H. (1986) Contrasting response of protein degradation to starvation and insulin as measured by N¹⁵-methylhistidine or phenylalanine release from the perfused rat heart. *Biochem. J.* **237**, 391–395, <https://doi.org/10.1042/bj2370391>
- Fuller, S.J., Gaitanaki, C.J. and Sugden, P.H. (1990) Effects of catecholamines on protein synthesis in cardiac myocytes and perfused hearts isolated from adult rats. Stimulation of translation is mediated through the α_1 -adrenoceptor. *Biochem. J.* **266**, 727–736, <https://doi.org/10.1042/bj2660727>
- Fuller, S.J., Gaitanaki, C.J. and Sugden, P.H. (1989) Effects of increasing extracellular pH on protein synthesis and protein degradation in the perfused working rat heart. *Biochem. J.* **259**, 173–179, <https://doi.org/10.1042/bj2590173>
- Massagué, J. and Czech, M.P. (1980) Multiple redox forms of the insulin receptor in native liver membranes. *Diabetes* **29**, 945–947, <https://doi.org/10.2337/diab.29.11.945>
- Lang, U., Kahn, C.R. and Harrison, L.C. (1980) Subunit structure of the insulin receptor of the human lymphocyte. *Biochemistry* **19**, 64–70, <https://doi.org/10.1021/bi00542a010>
- De Meyts, P. and Whittaker, J. (2002) Structural biology of insulin and IGF1 receptors: implications for drug design. *Nat. Rev. Drug Discov.* **1**, 769–783, <https://doi.org/10.1038/nrd917>
- Lawrence, M.C., McKern, N.M. and Ward, C.W. (2007) Insulin receptor structure and its implications for the IGF-1 receptor. *Curr. Opin. Struct. Biol.* **17**, 699–705, <https://doi.org/10.1016/j.sbi.2007.07.007>

- 29 Lawrence, M.C. (2021) Understanding insulin and its receptor from their three-dimensional structures. *Mol. Metab.* **52**, 101255, <https://doi.org/10.1016/j.molmet.2021.101255>
- 30 Blyth, A.J., Kirk, N.S. and Forbes, B.E. (2020) Understanding IGF-II action through insights into receptor binding and activation. *Cells* **9**, 2276, <https://doi.org/10.3390/cells9102276>
- 31 Hunter, T. (2014) The genesis of tyrosine phosphorylation. *Cold Spring Harb. Perspect. Biol.* **6**, a020644, <https://doi.org/10.1101/cshperspect.a020644>
- 32 De Meyts, P. (2015) Insulin/receptor binding: the last piece of the puzzle? What recent progress on the structure of the insulin/receptor complex tells us (or not) about negative cooperativity and activation. *Bioessays* **37**, 389–397, <https://doi.org/10.1002/bies.201400190>
- 33 Tollefsen, S.E. and Thompson, K. (1988) The structural basis for insulin-like growth factor I receptor high affinity binding. *J. Biol. Chem.* **263**, 16267–16273, [https://doi.org/10.1016/S0021-9258\(18\)37588-4](https://doi.org/10.1016/S0021-9258(18)37588-4)
- 34 Feltz, S.M., Swanson, M.L., Wemmie, J.A. and Pessin, J.E. (1988) Functional properties of an isolated alpha beta heterodimeric human placenta insulin-like growth factor 1 receptor complex. *Biochemistry* **27**, 3234–3242, <https://doi.org/10.1021/bi00409a017>
- 35 Soos, M.A. and Siddle, K. (1989) Immunological relationships between receptors for insulin and insulin-like growth factor I. Evidence for structural heterogeneity of insulin-like growth factor I receptors involving hybrids with insulin receptors. *Biochem. J.* **263**, 553–563, <https://doi.org/10.1042/bj2630553>
- 36 Soos, M.A., Whittaker, J., Lammers, R., Ullrich, A. and Siddle, K. (1990) Receptors for insulin and insulin-like growth factor-I can form hybrid dimers. Characterisation of hybrid receptors in transfected cells. *Biochem. J.* **270**, 383–390, <https://doi.org/10.1042/bj2700383>
- 37 Moxham, C.P., Duronio, V. and Jacobs, S. (1989) Insulin-like growth factor I receptor beta-subunit heterogeneity. Evidence for hybrid tetramers composed of insulin-like growth factor I and insulin receptor heterodimers. *J. Biol. Chem.* **264**, 13238–13244, [https://doi.org/10.1016/S0021-9258\(18\)51620-3](https://doi.org/10.1016/S0021-9258(18)51620-3)
- 38 Soos, M.A., Field, C.E. and Siddle, K. (1993) Purified hybrid insulin/insulin-like growth factor-I receptors bind insulin-like growth factor-I, but not insulin, with high affinity. *Biochem. J.* **290**, 419–426, <https://doi.org/10.1042/bj2900419>
- 39 Hanke, S. and Mann, M. (2009) The phosphotyrosine interactome of the insulin receptor family and its substrates IRS-1 and IRS-2. *Mol. Cell. Proteomics* **8**, 519–534, <https://doi.org/10.1074/mcp.M800407-MCP200>
- 40 Nagao, H., Cai, W., Wewer Albrechtsen, N.J., Steger, M., Batista, T.M. et al. (2021) Distinct signaling by insulin and IGF-1 receptors and their extra- and intracellular domains. *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2019474118, <https://doi.org/10.1073/pnas.2019474118>
- 41 Bailyes, E.M., Nave, B.T., Soos, M.A., Orr, S.R., Hayward, A.C. and Siddle, K. (1997) Insulin receptor/IGF-I receptor hybrids are widely distributed in mammalian tissues: quantification of individual receptor species by selective immunoprecipitation and immunoblotting. *Biochem. J.* **327**, 209–215, <https://doi.org/10.1042/bj3270209>
- 42 Torrente, Y., Bella, P., Tripodi, L., Villa, C. and Farini, A. (2020) Role of insulin-like growth factor receptor 2 across muscle homeostasis: implications for treating muscular dystrophy. *Cells* **9**, 441, <https://doi.org/10.3390/cells9020441>
- 43 El-Shewy, H.M. and Luttrell, L.M. (2009) Insulin-like growth factor-2/mannose-6 phosphate receptors. *Vitam. Horm.* **80**, 667–697, [https://doi.org/10.1016/S0083-6729\(08\)00624-9](https://doi.org/10.1016/S0083-6729(08)00624-9)
- 44 Peyrat, J.P., Bonnetterre, J., Beuscart, R., Djiane, J. and Demaille, A. (1988) Insulin-like growth factor 1 receptors in human breast cancer and their relation to estradiol and progesterone receptors. *Cancer Res.* **48**, 6429–6433
- 45 Moller, D.E., Yokota, A., Caro, J.F. and Flier, J.S. (1989) Tissue-specific expression of two alternatively spliced insulin receptor mRNAs in man. *Mol. Endocrinol.* **3**, 1263–1269, <https://doi.org/10.1210/mend-3-8-1263>
- 46 Frasca, F., Pandini, G., Scalia, P., Sciacca, L., Mineo, R., Costantino, A. et al. (1999) Insulin receptor isoform A, a newly recognized, high-affinity insulin-like growth factor II receptor in fetal and cancer cells. *Mol. Cell. Biol.* **19**, 3278–3288, <https://doi.org/10.1128/MCB.19.5.3278>
- 47 Vidal, H., Auboeuf, D., Beylot, M. and Riou, J.P. (1995) Regulation of insulin receptor mRNA splicing in rat tissues. Effect of fasting, aging, and diabetes. *Diabetes* **44**, 1196–1201, <https://doi.org/10.2337/diab.44.10.1196>
- 48 Gatford, K.L., Heinemann, G.K., Thompson, S.D., Zhang, J.V., Buckberry, S., Owens, J.A. et al. (2014) Circulating IGF1 and IGF2 and SNP genotypes in men and pregnant and non-pregnant women. *Endocr. Connect.* **3**, 138–149, <https://doi.org/10.1530/EC-14-0068>
- 49 Frystyk, J., Skjaerbaek, C., Zapf, J. and Orskov, H. (1998) Increased levels of circulating free insulin-like growth factors in patients with non-islet cell tumour hypoglycaemia. *Diabetologia* **41**, 589–594, <https://doi.org/10.1007/s001250050951>
- 50 Humbel, R.E. (1990) Insulin-like growth factors I and II. *Eur. J. Biochem.* **190**, 445–462, <https://doi.org/10.1111/j.1432-1033.1990.tb15595.x>
- 51 van Doorn, J. (2020) Insulin-like growth factor-II and bioactive proteins containing a part of the E-domain of pro-insulin-like growth factor-II. *Biofactors* **46**, 563–578, <https://doi.org/10.1002/biof.1623>
- 52 Giovannone, B., Scaldaferrì, M.L., Federici, M., Porzio, O., Lauro, D., Fusco, A. et al. (2000) Insulin receptor substrate (IRS) transduction system: distinct and overlapping signaling potential. *Diabetes Metab. Res. Rev.* **16**, 434–441, [https://doi.org/10.1002/1520-7560\(2000\)9999:9999%3c::AID-DMRR159%3e3.0.CO;2-8](https://doi.org/10.1002/1520-7560(2000)9999:9999%3c::AID-DMRR159%3e3.0.CO;2-8)
- 53 Guo, C.A. and Guo, S. (2017) Insulin receptor substrate signaling controls cardiac energy metabolism and heart failure. *J. Endocrinol.* **233**, R131–R143, <https://doi.org/10.1530/JOE-16-0679>
- 54 White, M.F. and Kahn, C.R. (2021) Insulin action at a molecular level - 100 years of progress. *Mol. Metab.* **52**, 101304, <https://doi.org/10.1016/j.molmet.2021.101304>
- 55 Manning, B.D. and Toker, A. (2017) AKT/PKB Signaling: Navigating the Network. *Cell* **169**, 381–405, <https://doi.org/10.1016/j.cell.2017.04.001>
- 56 Liu, G.Y. and Sabatini, D.M. (2020) mTOR at the nexus of nutrition, growth, ageing and disease. *Nat. Rev. Mol. Cell Biol.* **21**, 183–203, <https://doi.org/10.1038/s41580-019-0199-y>
- 57 Vanhaesebroeck, B., Guillermet-Guibert, J., Graupera, M. and Bilanges, B. (2010) The emerging mechanisms of isoform-specific PI3K signalling. *Nat. Rev. Mol. Cell Biol.* **11**, 329–341, <https://doi.org/10.1038/nrm2882>

- 58 Oudit, G.Y., Sun, H., Kerfant, B.G., Crackower, M.A., Penninger, J.M. and Backx, P.H. (2004) The role of phosphoinositide-3 kinase and PTEN in cardiovascular physiology and disease. *J. Mol. Cell Cardiol.* **37**, 449–471, <https://doi.org/10.1016/j.yjmcc.2004.05.015>
- 59 Polak, P. and Hall, M.N. (2006) mTORC2 caught in a SInful Akt. *Dev. Cell.* **11**, 433–434, <https://doi.org/10.1016/j.devcel.2006.09.005>
- 60 Pham, F.H., Sugden, P.H. and Clerk, A. (2000) Regulation of protein kinase B and 4E-BP1 by oxidative stress in cardiac myocytes. *Circ. Res.* **86**, 1252–1258, <https://doi.org/10.1161/01.RES.86.12.1252>
- 61 Clerk, A., Aggeli, I.K.S., Stathopoulou, K. and Sugden, P.H. (2006) Peptide growth factors signal differentially through protein kinase C to extracellular signal-regulated kinases in neonatal cardiomyocytes. *Cell. Signal.* **18**, 225–235, <https://doi.org/10.1016/j.cellsig.2005.04.005>
- 62 Meijles, D., Fuller, S.J., Cull, J.J., Alharbi, H.O., Cooper, S.T.E., Sugden, P.H. et al. (2021) The insulin receptor family and protein kinase B (Akt) are activated in the heart by alkaline pH and α_1 -adrenergic receptors. *Biochem. J.* **478**, 2059–2079, <https://doi.org/10.1042/BCJ20210144>
- 63 Hermida, M.A., Dinesh Kumar, J. and Leslie, N.R. (2017) GSK3 and its interactions with the PI3K/AKT/mTOR signalling network. *Adv. Biol. Regul.* **65**, 5–15, <https://doi.org/10.1016/j.bior.2017.06.003>
- 64 Kerkelä, R., Woulfe, K. and Force, T. (2007) Glycogen synthase kinase-3 β - actively inhibiting hypertrophy. *Trends Cardiovasc. Med.* **17**, 91–96, <https://doi.org/10.1016/j.tcm.2007.01.004>
- 65 Lal, H., Ahmad, F., Woodgett, J. and Force, T. (2015) The GSK-3 family as therapeutic target for myocardial diseases. *Circ. Res.* **116**, 138–149, <https://doi.org/10.1161/CIRCRESAHA.116.303613>
- 66 Sugden, P.H., Fuller, S.J., Weiss, S.C. and Clerk, A. (2008) Glycogen synthase 3 as a point of integration in hypertrophic signalling in the heart: a critical analysis. *Br. J. Pharmacol.* **153**, S137–S153, <https://doi.org/10.1038/sj.bjp.0707659>
- 67 Roach, W.G., Chavez, J.A., Miinea, C.P. and Lienhard, G.E. (2007) Substrate specificity and effect on GLUT4 translocation of the Rab GTPase-activating protein Tbc1d1. *Biochem. J.* **403**, 353–358, <https://doi.org/10.1042/BJ20061798>
- 68 Chen, S., Murphy, J., Toth, R., Campbell, D.G., Morrice, N.A. and Mackintosh, C. (2008) Complementary regulation of TBC1D1 and AS160 by growth factors, insulin and AMPK activators. *Biochem. J.* **409**, 449–459, <https://doi.org/10.1042/BJ20071114>
- 69 Samovski, D., Su, X., Xu, Y., Abumrad, N.A. and Stahl, P.D. (2012) Insulin and AMPK regulate FA translocase/CD36 plasma membrane recruitment in cardiomyocytes via Rab GAP AS160 and Rab8a Rab GTPase. *J. Lipid Res.* **53**, 709–717, <https://doi.org/10.1194/jlr.M023424>
- 70 Deprez, J., Vertommen, D., Alessi, D.R. and Rider, M.H. (1997) Phosphorylation and activation of heart 6-phosphofructo-2-kinase by protein kinase B and other protein kinases of the insulin signaling cascades. *J. Biol. Chem.* **272**, 17269–17275, <https://doi.org/10.1074/jbc.272.28.17269>
- 71 Bertrand, L., Alessi, D.R., Deprez, J., Deak, M., Viaene, E., Rider, M.H. et al. (1999) Heart 6-phosphofructo-2-kinase activation by insulin results from Ser-466 and Ser-483 phosphorylation and requires 3-phosphoinositide-dependent kinase-1, but not protein kinase B. *J. Biol. Chem.* **274**, 30927–30933, <https://doi.org/10.1074/jbc.274.43.30927>
- 72 Mouton, V., Toussaint, L., Vertommen, D., Gueuning, M.A., Maisin, L., Havaux, X. et al. (2010) Heart 6-phosphofructo-2-kinase activation by insulin requires PKB (protein kinase B), but not SGK3 (serum- and glucocorticoid-induced protein kinase 3). *Biochem. J.* **431**, 267–275, <https://doi.org/10.1042/BJ20101089>
- 73 Roberts, D.J. and Miyamoto, S. (2015) Hexokinase II integrates energy metabolism and cellular protection: Aktting on mitochondria and TORCing to autophagy. *Cell Death Differ.* **22**, 248–257, <https://doi.org/10.1038/cdd.2014.173>
- 74 Gottlob, K., Majewski, N., Kennedy, S., Kandel, E., Robey, R.B. and Hay, N. (2001) Inhibition of early apoptotic events by Akt/PKB is dependent on the first committed step of glycolysis and mitochondrial hexokinase. *Genes Dev.* **15**, 1406–1418, <https://doi.org/10.1101/gad.889901>
- 75 Halestrap, A.P., Pereira, G.C. and Pasdois, P. (2015) The role of hexokinase in cardioprotection - mechanism and potential for translation. *Br. J. Pharmacol.* **172**, 2085–2100, <https://doi.org/10.1111/bph.12899>
- 76 Dimmeler, S., Fleming, I., Fisslthaler, B., Hermann, C., Busse, R. and Zeiher, A.M. (1999) Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* **399**, 601–605, <https://doi.org/10.1038/21224>
- 77 Fulton, D., Gratton, J.P., McCabe, T.J., Fontana, J., Fujio, Y., Walsh, K. et al. (1999) Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. *Nature* **399**, 597–601, <https://doi.org/10.1038/21218>
- 78 Michell, B.J., Griffiths, J.E., Mitchellhill, K.I., Rodriguez-Crespo, I., Tiganis, T., Bozinovski, S. et al. (1999) The Akt kinase signals directly to endothelial nitric oxide synthase. *Curr. Biol.* **9**, 845–848, [https://doi.org/10.1016/S0960-9822\(99\)80371-6](https://doi.org/10.1016/S0960-9822(99)80371-6)
- 79 Yu, Q., Gao, F. and Ma, X.L. (2011) Insulin says NO to cardiovascular disease. *Cardiovasc. Res.* **89**, 516–524, <https://doi.org/10.1093/cvr/cvq349>
- 80 Seo, G.J., Yang, A., Tan, B., Kim, S., Liang, Q., Choi, Y. et al. (2015) Akt kinase-mediated checkpoint of cGAS DNA sensing pathway. *Cell Rep.* **13**, 440–449, <https://doi.org/10.1016/j.celrep.2015.09.007>
- 81 Lei, C., Tan, Y., Ni, D., Peng, J. and Yi, G. (2022) cGAS-STING signaling in ischemic diseases. *Clin. Chim. Acta* **531**, 177–182, <https://doi.org/10.1016/j.cca.2022.04.003>
- 82 Wu, S., Zhang, Q., Zhang, F., Meng, F., Liu, S., Zhou, R. et al. (2019) HER2 recruits AKT1 to disrupt STING signalling and suppress antiviral defence and antitumour immunity. *Nat. Cell Biol.* **21**, 1027–1040, <https://doi.org/10.1038/s41556-019-0352-z>
- 83 Brunet, A., Bonni, A., Zigmond, M.J., Lin, M.Z., Juo, P., Hu, L.A. et al. (1999) Akt promotes cell survival by phosphorylating and inhibiting a Forkhead transcription factor. *Cell* **96**, 857–868, [https://doi.org/10.1016/S0092-8674\(00\)80595-4](https://doi.org/10.1016/S0092-8674(00)80595-4)
- 84 Rena, G., Guo, S., Cichy, S.C., Unterman, T.G. and Cohen, P. (1999) Phosphorylation of the transcription factor forkhead family member FKHR by protein kinase B. *J. Biol. Chem.* **274**, 17179–17183, <https://doi.org/10.1074/jbc.274.24.17179>
- 85 Skurk, C., Izumiya, Y., Maatz, H., Razeghi, P., Shiojima, I., Sandri, M. et al. (2005) The FOXO3a transcription factor regulates cardiac myocyte size downstream of AKT signaling. *J. Biol. Chem.* **280**, 20814–20823, <https://doi.org/10.1074/jbc.M500528200>
- 86 Menghini, R., Casagrande, V., Iuliani, G., Rizza, S., Mavilio, M., Cardellini, M. et al. (2020) Metabolic aspects of cardiovascular diseases: Is FoxO1 a player or a target? *Int. J. Biochem. Cell Biol.* **118**, 105659, <https://doi.org/10.1016/j.biocel.2019.105659>
- 87 Papanicolaou, K.N., Izumiya, Y. and Walsh, K. (2008) Forkhead transcription factors and cardiovascular biology. *Circ. Res.* **102**, 16–31, <https://doi.org/10.1161/CIRCRESAHA.107.164186>

- 88 Nandagopal, N. and Roux, P.P. (2015) Regulation of global and specific mRNA translation by the mTOR signaling pathway. *Translation (Austin)* **3**, e983402, <https://doi.org/10.4161/21690731.2014.983402>
- 89 Iadevaia, V., Liu, R. and Proud, C.G. (2014) mTORC1 signaling controls multiple steps in ribosome biogenesis. *Semin. Cell Dev. Biol.* **36**, 113–120, <https://doi.org/10.1016/j.semcdb.2014.08.004>
- 90 Proud, C.G. (2006) Regulation of protein synthesis by insulin. *Biochem. Soc. Trans.* **34**, 213–216, <https://doi.org/10.1042/BSR200340213>
- 91 Wang, L., Wang, X. and Proud, C.G. (2000) Activation of mRNA translation in rat cardiac myocytes by insulin involves multiple rapamycin-sensitive steps. *Am. J. Physiol. Heart Circ. Physiol.* **278**, H1056–H1068, <https://doi.org/10.1152/ajpheart.2000.278.4.H1056>
- 92 Markou, T., Marshall, A.K., Cullingford, T.E., Tham, E.L., Sugden, P.H. and Clerk, A. (2010) Regulation of the cardiomyocyte transcriptome vs translome by endothelin-1 and insulin: translational regulation of 5' terminal oligopyrimidine tract (TOP) mRNAs by insulin. *BMC Genomics* **11**, 343, <https://doi.org/10.1186/1471-2164-11-343>
- 93 Condon, K.J. and Sabatini, D.M. (2019) Nutrient regulation of mTORC1 at a glance. *J. Cell Sci.* **132**, jcs222570, <https://doi.org/10.1242/jcs.222570>
- 94 Ray, L.B. and Sturgill, T.W. (1987) Rapid stimulation of a serine/threonine protein kinase in 3T3-L1 adipocytes that phosphorylates microtubule-associated protein 2 in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **84**, 1502–1506, <https://doi.org/10.1073/pnas.84.6.1502>
- 95 Skolnik, E.Y., Batzer, A., Li, N., Lee, C.H., Lowenstein, E., Mohammadi, M. et al. (1993) The function of GRB2 in linking the insulin receptor to Ras signaling pathways. *Science* **260**, 1953–1955, <https://doi.org/10.1126/science.8316835>
- 96 Cargnello, M. and Roux, P.P. (2011) Activation and function of the MAPKs and their substrates, the MAPK-activated protein kinases. *Microbiol. Mol. Biol. Rev.* **75**, 50–83, <https://doi.org/10.1128/MMBR.00031-10>
- 97 Denton, R.M. and Tavaré, J.M. (1995) Does the mitogen-activated-protein kinase cascade have a role in insulin action? The cases for and against. *Eur. J. Biochem.* **227**, 597–611, <https://doi.org/10.1111/j.1432-1033.1995.tb20179.x>
- 98 Hernandez-Sanchez, C., Mansilla, A., de Pablo, F. and Zardoya, R. (2008) Evolution of the insulin receptor family and receptor isoform expression in vertebrates. *Mol. Biol. Evol.* **25**, 1043–1053, <https://doi.org/10.1093/molbev/msn036>
- 99 Shier, P. and Watt, V.M. (1992) Tissue-specific expression of the rat insulin receptor-related receptor gene. *Mol. Endocrinol.* **6**, 723–729
- 100 Kurachi, H., Jobo, K., Ohta, M., Kawasaki, T. and Itoh, N. (1992) A new member of the insulin receptor family, insulin receptor-related receptor, is expressed preferentially in the kidney. *Biochem. Biophys. Res. Commun.* **187**, 934–939, [https://doi.org/10.1016/0006-291X\(92\)91287-Z](https://doi.org/10.1016/0006-291X(92)91287-Z)
- 101 Mathi, S.K., Chan, J. and Watt, V.M. (1995) Insulin receptor-related receptor messenger ribonucleic acid: quantitative distribution and localization to subpopulations of epithelial cells in stomach and kidney. *Endocrinology* **136**, 4125–4132, <https://doi.org/10.1210/endo.136.9.7649121>
- 102 Ozaki, K., Takada, N., Tsujimoto, K., Tsuji, N., Kawamura, T., Muso, E. et al. (1997) Localization of insulin receptor-related receptor in the rat kidney. *Kidney Int.* **52**, 694–698, <https://doi.org/10.1038/ki.1997.384>
- 103 Kovacina, K.S. and Roth, R.A. (1995) Characterization of the endogenous insulin receptor-related receptor in neuroblastomas. *J. Biol. Chem.* **270**, 1881–1887, <https://doi.org/10.1074/jbc.270.4.1881>
- 104 Jui, H.Y., Accili, D. and Taylor, S.I. (1996) Characterization of a hybrid receptor formed by dimerization of the insulin receptor-related receptor (IRR) with the insulin receptor (IR): coexpression of cDNAs encoding human IRR and human IR in NIH-3T3 cells. *Biochemistry* **35**, 14326–14330, <https://doi.org/10.1021/bi9613032>
- 105 Deyev, I.E., Sohet, F., Vassilenko, K.P., Serova, O.V., Popova, N.V., Zozulya, S.A. et al. (2011) Insulin receptor-related receptor as an extracellular alkali sensor. *Cell Metab.* **13**, 679–689, <https://doi.org/10.1016/j.cmet.2011.03.022>
- 106 Kitamura, T., Kido, Y., Nef, S., Merenmies, J., Parada, L.F. and Accili, D. (2001) Preserved pancreatic beta-cell development and function in mice lacking the insulin receptor-related receptor. *Mol. Cell. Biol.* **21**, 5624–5630, <https://doi.org/10.1128/MCB.21.16.5624-5630.2001>
- 107 Saltiel, A.R. (2021) Insulin signaling in health and disease. *J. Clin. Invest.* **131**, e142241, <https://doi.org/10.1172/JCI142241>
- 108 Anderson, L.E. and Henrich, W.L. (1987) Alkalemia-associated morbidity and mortality in medical and surgical patients. *South. Med. J.* **80**, 729–733, <https://doi.org/10.1097/00007611-198706000-00016>
- 109 Hong, T. and Shaw, R.M. (2017) Cardiac T-tubule microanatomy and function. *Physiol. Rev.* **97**, 227–252, <https://doi.org/10.1152/physrev.00037.2015>
- 110 Ibrahim, M., Gorelik, J., Yacoub, M.H. and Terracciano, C.M. (2011) The structure and function of cardiac t-tubules in health and disease. *Proc. Biol. Sci.* **278**, 2714–2723, <https://doi.org/10.1098/rspb.2011.0624>
- 111 Zhang, M., Perino, A., Ghigo, A., Hirsch, E. and Shah, A.M. (2013) NADPH oxidases in heart failure: poachers or gamekeepers? *Antioxid. Redox Signal.* **18**, 1024–1041, <https://doi.org/10.1089/ars.2012.4550>
- 112 Jui, H.Y., Suzuki, Y., Accili, D. and Taylor, S.I. (1994) Expression of a cDNA encoding the human insulin receptor-related receptor. *J. Biol. Chem.* **269**, 22446–22452, [https://doi.org/10.1016/S0021-9258\(17\)31810-0](https://doi.org/10.1016/S0021-9258(17)31810-0)
- 113 Cattaneo, F., Guerra, G., Parisi, M., De Marinis, M., Tafuri, D., Cinelli, M. et al. (2014) Cell-surface receptors transactivation mediated by G protein-coupled receptors. *Int. J. Mol. Sci.* **15**, 19700–19728, <https://doi.org/10.3390/ijms151119700>
- 114 Clerk, A., Cullingford, T.E., Fuller, S.J., Giraldo, A., Markou, T., Pikkariainen, S. et al. (2007) Signaling pathways mediating cardiac myocyte gene expression in physiological and stress responses. *J. Cell. Physiol.* **212**, 311–322, <https://doi.org/10.1002/jcp.21094>
- 115 Palanisamy, S., Xue, C., Ishiyama, S., Prasad, S.V.N. and Gabrielson, K. (2021) GPCR-ErbB transactivation pathways and clinical implications. *Cell. Signal.* **86**, 110092, <https://doi.org/10.1016/j.cellsig.2021.110092>
- 116 Haxho, F., Haq, S. and Szewczuk, M.R. (2018) Biased G protein-coupled receptor agonism mediates Neu1 sialidase and matrix metalloproteinase-9 crosstalk to induce transactivation of insulin receptor signaling. *Cell. Signal.* **43**, 71–84, <https://doi.org/10.1016/j.cellsig.2017.12.006>
- 117 Kilpatrick, L.E. and Hill, S.J. (2021) Transactivation of G protein-coupled receptors (GPCRs) and receptor tyrosine kinases (RTKs): recent insights using luminescence and fluorescence technologies. *Curr. Opin. Endocr. Metab. Res.* **16**, 102–112, <https://doi.org/10.1016/j.coemr.2020.10.003>
- 118 Sumitomo, M., Milowsky, M.I., Shen, R., Navarro, D., Dai, J., Asano, T. et al. (2001) Neutral endopeptidase inhibits neuropeptide-mediated transactivation of the insulin-like growth factor receptor-Akt cell survival pathway. *Cancer Res.* **61**, 3294–3298

- 119 Mulvihill, M.J., Cooke, A., Rosenfeld-Franklin, M., Buck, E., Foreman, K., Landfair, D. et al. (2009) Discovery of OSI-906: a selective and orally efficacious dual inhibitor of the IGF-1 receptor and insulin receptor. *Fut. Med. Chem.* **1**, 1153–1171, <https://doi.org/10.4155/fmc.09.89>
- 120 Amatyia, N., Lin, D.Y. and Andreotti, A.H. (2019) Dynamic regulatory features of the protein tyrosine kinases. *Biochem. Soc. Trans.* **47**, 1101–1116, <https://doi.org/10.1042/BSR20180590>
- 121 Santos, C.X., Raza, S. and Shah, A.M. (2016) Redox signaling in the cardiomyocyte: From physiology to failure. *Int. Biochem. Cell Biol.* **74**, 145–151, <https://doi.org/10.1016/j.biocel.2016.03.002>
- 122 Terzic, A., Pucéat, M., Clement, O., Scamps, F. and Vassort, G. (1992) α_1 -adrenergic effects on intracellular pH and calcium and on myofilaments in single rat cardiac cells. *J. Physiol.* **447**, 275–292, <https://doi.org/10.1113/jphysiol.1992.sp019002>
- 123 Vila-Petroff, M., Perez, G.N., Alvarez, B., Cingolani, H.E. and Mattiazzi, A. (1996) Mechanism of negative lusitropic effect of α_1 -adrenoceptor stimulation in cat papillary muscles. *Am. J. Physiol.* **270**, H701–H709
- 124 Kim, D. and Smith, T.W. (1987) Altered Ca fluxes and contractile state during pH changes in cultured heart cells. *Am. J. Physiol.* **253**, C137–C146, <https://doi.org/10.1152/ajpcell.1987.253.1.C137>
- 125 Spitzer, K.W. and Bridge, J.H. (1992) Relationship between intracellular pH and tension development in resting ventricular muscle and myocytes. *Am. J. Physiol.* **262**, C316–C327, <https://doi.org/10.1152/ajpcell.1992.262.2.C316>
- 126 Casey, J.R., Grinstein, S. and Orlowski, J. (2010) Sensors and regulators of intracellular pH. *Nat. Rev. Mol. Cell Biol.* **11**, 50–61, <https://doi.org/10.1038/nrm2820>
- 127 Wang, H.S., Chen, Y., Vairamani, K. and Shull, G.E. (2014) Critical role of bicarbonate and bicarbonate transporters in cardiac function. *World J. Biol. Chem.* **5**, 334–345, <https://doi.org/10.4331/wjbc.v5.i3.334>
- 128 Pucéat, M., Korichneva, I., Cassoly, R. and Vassort, G. (1995) Identification of band 3-like proteins and Cl-/HCO₃- exchange in isolated cardiomyocytes. *J. Biol. Chem.* **270**, 1315–1322, <https://doi.org/10.1074/jbc.270.3.1315>
- 129 Sowah, D., Brown, B.F., Quon, A., Alvarez, B.V. and Casey, J.R. (2014) Resistance to cardiomyocyte hypertrophy in ae3-/- mice, deficient in the AE3 Cl. *BMC Cardiovasc. Disord.* **14**, 89, <https://doi.org/10.1186/1471-2261-14-89>
- 130 Alvarez, B.V., Johnson, D.E., Sowah, D., Soliman, D., Light, P.E., Xia, Y. et al. (2007) Carbonic anhydrase inhibition prevents and reverts cardiomyocyte hypertrophy. *J. Physiol.* **579**, 127–145, <https://doi.org/10.1113/jphysiol.2006.123638>
- 131 Brown, B.F., Quon, A., Dyck, J.R. and Casey, J.R. (2012) Carbonic anhydrase II promotes cardiomyocyte hypertrophy. *Can. J. Physiol. Pharmacol.* **90**, 1599–1610, <https://doi.org/10.1139/y2012-142>
- 132 Sugden, P.H. and Bogoyevitch, M.A. (1996) Endothelin-1-dependent signaling pathways in the myocardium. *Trends Cardiovasc. Med.* **6**, 87–94, [https://doi.org/10.1016/1050-1738\(96\)00013-8](https://doi.org/10.1016/1050-1738(96)00013-8)
- 133 Clerk, A., Gillespie-Brown, J., Fuller, S.J. and Sugden, P.H. (1996) Stimulation of phosphatidylinositol hydrolysis, protein kinase C translocation, and mitogen-activated protein kinase activity by bradykinin in ventricular myocytes. Dissociation from the hypertrophic response. *Biochem. J.* **317**, 109–118, <https://doi.org/10.1042/bj3170109>
- 134 Clerk, A. and Sugden, P.H. (1997) Regulation of phospholipases C and D in rat ventricular myocytes: stimulation by endothelin-1, bradykinin and phenylephrine. *J. Mol. Cell Cardiol.* **29**, 1593–1604, <https://doi.org/10.1006/jmcc.1997.0395>
- 135 Shimizu, I., Minamino, T., Toko, H., Okada, S., Ikeda, H., Yasuda, N. et al. (2010) Excessive cardiac insulin signaling exacerbates systolic dysfunction induced by pressure overload in rodents. *J. Clin. Invest.* **120**, 1506–1514, <https://doi.org/10.1172/JCI40096>
- 136 Hausenloy, D.J. and Yellon, D.M. (2004) New directions for protecting the heart against ischaemia- reperfusion injury: targeting the reperfusion Injury Salvage Kinase (RISK)-pathway. *Cardiovasc. Res.* **61**, 448–460, <https://doi.org/10.1016/j.cardiores.2003.09.024>
- 137 Turer, A.T. and Hill, J.A. (2010) Pathogenesis of myocardial ischemia-reperfusion injury and rationale for therapy. *Am. J. Cardiol.* **106**, 360–368, <https://doi.org/10.1016/j.amjcard.2010.03.032>
- 138 Jonassen, A.K., Sack, M.N., Mjos, O.D. and Yellon, D.M. (2001) Myocardial protection by insulin at reperfusion requires early administration and is mediated by Akt and p70s6 kinase cell-survival signaling. *Circ. Res.* **89**, 1191–1198, <https://doi.org/10.1161/hh2401.101385>
- 139 Fuglestad, B.N., Tiron, C., Jonassen, A.K., Mjos, O.D. and Ytrehus, K. (2009) Pretreatment with insulin before ischaemia reduces infarct size in Langendorff-perfused rat hearts. *Acta Physiol. (Oxf.)* **195**, 273–282, <https://doi.org/10.1111/j.1748-1716.2008.01901.x>
- 140 Jonassen, A.K., Brar, B.K., Mjos, O.D., Sack, M.N., Latchman, D.S. and Yellon, D.M. (2000) Insulin administered at reoxygenation exerts a cardioprotective effect in myocytes by a possible anti-apoptotic mechanism. *J. Mol. Cell Cardiol.* **32**, 757–764, <https://doi.org/10.1006/jmcc.2000.1118>
- 141 Baines, C.P., Wang, L., Cohen, M.V. and Downey, J.M. (1999) Myocardial protection by insulin is dependent on phosphatidylinositol 3-kinase but not protein kinase C or KATP channels in the isolated rabbit heart. *Basic Res. Cardiol.* **94**, 188–198, <https://doi.org/10.1007/s003950050142>
- 142 Duerr, R.L., Huang, S., Miraliakbar, H.R., Clark, R., Chien, K.R. and Ross, Jr, J. (1995) Insulin-like growth factor 1 enhances ventricular hypertrophy and function during the onset of experimental cardiac failure. *J. Clin. Invest.* **95**, 619–627, <https://doi.org/10.1172/JCI117706>
- 143 Duerr, R.L., McKirnan, M.D., Gim, R.D., Clark, R.G., Chien, K.R. and Ross, Jr, J. (1996) Cardiovascular effects of insulin-like growth factor-1 and growth hormone in chronic left ventricular failure in the rat. *Circulation* **93**, 2188–2196, <https://doi.org/10.1161/01.CIR.93.12.2188>
- 144 Santini, M.P., Tsao, L., Monassier, L., Theodoropoulos, C., Carter, J., Lara-Pezzi, E. et al. (2007) Enhancing repair of the mammalian heart. *Circ. Res.* **100**, 1732–1740, <https://doi.org/10.1161/CIRCRESAHA.107.148791>
- 145 Hausenloy, D.J. and Yellon, D.M. (2006) Survival kinases in ischemic preconditioning and postconditioning. *Cardiovasc. Res.* **70**, 240–253, <https://doi.org/10.1016/j.cardiores.2006.01.017>
- 146 Yao, H., Han, X. and Han, X. (2014) The cardioprotection of the insulin-mediated PI3K/Akt/mTOR signaling pathway. *Am. J. Cardiovasc. Drugs* **14**, 433–442, <https://doi.org/10.1007/s40256-014-0089-9>
- 147 Abel, E.D. (2021) Insulin signaling in the heart. *Am. J. Physiol. Endocrinol. Metab.* **321**, E130–E145, <https://doi.org/10.1152/ajpendo.00158.2021>

- 148 Fontana, L., Vinciguerra, M. and Longo, V.D. (2012) Growth factors, nutrient signaling, and cardiovascular aging. *Circ. Res.* **110**, 1139–1150, <https://doi.org/10.1161/CIRCRESAHA.111.246470>
- 149 Becker, L.B. (2004) New concepts in reactive oxygen species and cardiovascular reperfusion physiology. *Cardiovasc. Res.* **61**, 461–470, <https://doi.org/10.1016/j.cardiores.2003.10.025>
- 150 Lemasters, J.J., Bond, J.M., Chacon, E., Harper, I.S., Kaplan, S.H., Ohata, H. et al. (1996) The pH paradox in ischemia-reperfusion injury to cardiac myocytes. *EXS* **76**, 99–114, https://doi.org/10.1007/978-3-0348-8988-9_7
- 151 Clerk, A., Cole, S.M., Cullingford, T.E., Harrison, J.G., Jormakka, M. and Valks, D.M. (2003) Regulation of cardiac myocyte cell death. *Pharmacol. Ther.* **97**, 223–261, [https://doi.org/10.1016/S0163-7258\(02\)00339-X](https://doi.org/10.1016/S0163-7258(02)00339-X)
- 152 Green, C.L., Lamming, D.W. and Fontana, L. (2022) Molecular mechanisms of dietary restriction promoting health and longevity. *Nat. Rev. Mol. Cell Biol.* **23**, 56–73, <https://doi.org/10.1038/s41580-021-00411-4>
- 153 Miura, T. and Miki, T. (2009) GSK-3 β , a therapeutic target for cardiomyocyte protection. *Circ. J.* **73**, 1184–1192, <https://doi.org/10.1253/circj.CJ-09-0284>
- 154 Maroko, P.R., Libby, P., Sobel, B.E., Bloor, C.M., Sybers, H.D., Shell, W.E. et al. (1972) Effect of glucose-insulin-potassium infusion on myocardial infarction following experimental coronary artery occlusion. *Circulation* **45**, 1160–1175, <https://doi.org/10.1161/01.CIR.45.6.1160>
- 155 Braunwald, E. and Maroko, P.R. (1973) Investigations on protection of the ischemic myocardium. *Trans. Am. Clin. Climatol. Assoc.* **84**, 80–115
- 156 Scheuer, J. (1975) Some metabolic approaches to improving myocardial performance during cardiac hypoxia. *Recent Adv. Stud. Cardiac Struct. Metab.* **10**, 195–207
- 157 Van Rooyen, J., McCarthy, J. and Opie, L.H. (2002) Increased glycolysis during ischaemia mediates the protective effect of glucose and insulin in the isolated rat heart despite the presence of cardiodepressant exogenous substrates. *Cardiovasc. J. S. Afr.* **13**, 103–109
- 158 Wilson, J.E. (2003) Isozymes of mammalian hexokinase: structure, subcellular localization and metabolic function. *J. Exp. Biol.* **206**, 2049–2057, <https://doi.org/10.1242/jeb.00241>
- 159 Gerbitz, K.D., Gempel, K. and Brdiczka, D. (1996) Mitochondria and diabetes. Genetic, biochemical, and clinical implications of the cellular energy circuit. *Diabetes* **45**, 113–126, <https://doi.org/10.2337/diab.45.2.113>
- 160 Tran, D.H. and Wang, Z.V. (2019) Glucose metabolism in cardiac hypertrophy and heart failure. *J. Am. Heart Assoc.* **8**, e012673, <https://doi.org/10.1161/JAHA.119.012673>
- 161 Zimmer, H.G., Bunger, R., Koschine, H., Steinkopff, G. and Ibel, H. (1982) Possible role of oxidized glutathione for the regulation of the myocardial hexose monophosphate shunt. *Adv. Myocardiol.* **3**, 577–584, https://doi.org/10.1007/978-1-4899-5561-6_55
- 162 Wang, L., Li, J. and Di, L.J. (2022) Glycogen synthesis and beyond, a comprehensive review of GSK3 as a key regulator of metabolic pathways and a therapeutic target for treating metabolic diseases. *Med. Res. Rev.* **42**, 946–982, <https://doi.org/10.1002/med.21867>
- 163 Jennings, R.B. and Reimer, K.A. (1991) The cell biology of acute myocardial ischemia. *Annu. Rev. Med.* **42**, 225–246, <https://doi.org/10.1146/annurev.me.42.020191.001301>
- 164 Cohen, P., Burchell, A., Foulkes, J.G., Cohen, P.T., Vanaman, T.C. and Nairn, A.C. (1978) Identification of the Ca²⁺-dependent modulator protein as the fourth subunit of rabbit skeletal muscle phosphorylase kinase. *FEBS Lett.* **92**, 287–293, [https://doi.org/10.1016/0014-5793\(78\)80772-8](https://doi.org/10.1016/0014-5793(78)80772-8)
- 165 Cohen, P. (1980) The role of calcium ions, calmodulin and troponin in the regulation of phosphorylase kinase from rabbit skeletal muscle. *Eur. J. Biochem.* **111**, 563–574, <https://doi.org/10.1111/j.1432-1033.1980.tb04972.x>
- 166 Scheuer, J. and Stezoski, S.W. (1968) Effects of high-energy phosphate depletion and repletion on the dynamics and electrocardiogram of isolated rat hearts. *Circ. Res.* **23**, 519–530, <https://doi.org/10.1161/01.RES.23.4.519>
- 167 Ofir, M., Arad, M., Porat, E., Freimark, D., Chepurko, Y., Vidne, B.A. et al. (2008) Increased glycogen stores due to γ -AMPK overexpression protects against ischemia and reperfusion damage. *Biochem. Pharmacol.* **75**, 1482–1491, <https://doi.org/10.1016/j.bcp.2007.12.011>
- 168 Gopal, K., Saleme, B., Al Batran, R., Aburasayn, H., Eshreif, A., Ho, K.L. et al. (2017) FoxO1 regulates myocardial glucose oxidation rates via transcriptional control of pyruvate dehydrogenase kinase 4 expression. *Am. J. Physiol. Heart Circ. Physiol.* **313**, H479–H490, <https://doi.org/10.1152/ajpheart.00191.2017>
- 169 Wang, F., Yuan, Q., Chen, F., Pang, J., Pan, C., Xu, F. et al. (2021) Fundamental mechanisms of the cell death caused by nitrosative stress. *Front. Cell Dev. Biol.* **9**, 742483, <https://doi.org/10.3389/fcell.2021.742483>
- 170 Seddon, M., Shah, A.M. and Casadei, B. (2007) Cardiomyocytes as effectors of nitric oxide signalling. *Cardiovasc. Res.* **75**, 315–326, <https://doi.org/10.1016/j.cardiores.2007.04.031>
- 171 Paulus, W.J. and Bronzwaer, J.G. (2004) Nitric oxide's role in the heart: control of beating or breathing? *Am. J. Physiol. Heart Circ. Physiol.* **287**, H8–H13, <https://doi.org/10.1152/ajpheart.01147.2003>
- 172 Flier, J.S. (2019) C. Ronald Kahn: The Louisville Slugger of metabolic science. *J. Clin. Invest.* **129**, 5066–5070, <https://doi.org/10.1172/JCI133156>
- 173 Deyev, I.E., Mitrofanova, A.V., Zhevennev, E.S., Radionov, N., Berchatova, A.A., Popova, N.V. et al. (2013) Structural determinants of the insulin receptor-related receptor activation by alkali. *J. Biol. Chem.* **288**, 33884–33893, <https://doi.org/10.1074/jbc.M113.483172>
- 174 Deyev, I.E., Chachina, N.A., Shayahmetova, D.M., Serova, O.V. and Petrenko, A.G. (2015) Mapping of alkali-sensing sites of the insulin receptor-related receptor. The role of L2 and fibronectin domains. *Biochimie* **111**, 1–9, <https://doi.org/10.1016/j.biochi.2014.12.014>
- 175 Shtykova, E.V., Petoukhov, M.V., Mozhaev, A.A., Deyev, I.E., Dadinova, L.A., Loshkarev, N.A. et al. (2019) The dimeric ectodomain of the alkali-sensing insulin receptor-related receptor (ectolRR) has a droplike shape. *J. Biol. Chem.* **294**, 17790–17798, <https://doi.org/10.1074/jbc.RA119.010390>
- 176 Batishchev, O.V., Kuzmina, N.V., Mozhaev, A.A., Goryashchenko, A.S., Mileschina, E.D., Orsa, A.N. et al. (2021) Activity-dependent conformational transitions of the insulin receptor-related receptor. *J. Biol. Chem.* **296**, 100534, <https://doi.org/10.1016/j.jbc.2021.100534>
- 177 Scheuer, J. and Berry, M.N. (1967) Effects of alkalosis on glycolysis in the working heart. *Am. J. Physiol.* **213**, 1143–1148, <https://doi.org/10.1152/ajplegacy.1967.213.5.1143>

178 Ushkaryov, Y.A. and Langenhan, T. (2021) Alexander Georgievich Petrenko (1959–2021). *FEBS J.* **288**, 6680–6682,
<https://doi.org/10.1111/febs.16282>