

Evaluating winter precipitation over the western Himalayas in a high-resolution Indian regional reanalysis using multi-source climate datasets

Article

Accepted Version

Nischal, Attada, R. and Hunt, K. M. R. ORCID:
<https://orcid.org/0000-0003-1480-3755> (2022) Evaluating winter precipitation over the western Himalayas in a high-resolution Indian regional reanalysis using multi-source climate datasets. *Journal of Applied Meteorology and Climatology*, 61 (11). pp. 1613-1633. ISSN 1558-8432 doi: 10.1175/JAMC-D-21-0172.1 Available at <https://centaur.reading.ac.uk/106218/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/JAMC-D-21-0172.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Evaluating Winter Precipitation over the Western Himalayas in a High-Resolution Indian Regional Reanalysis using multi-source climate datasets

Nischal¹, Raju Attada^{1*}, Kieran M. R. Hunt²

¹Department of Earth and Environmental Sciences, Indian Institute of Science Education and Research Mohali, Punjab, India

²National Centre for Atmospheric Science, Department of Meteorology, University of Reading, UK

**Revision Submitted to
Journal of Applied Meteorology and Climatology**

***Corresponding author:**

Raju Attada
Department of Earth and Environmental Sciences,
Indian Institute of Science Education and Research Mohali, India.
Email: rajuattada@iisermohali.ac.in

ABSTRACT

1
2 Considerable uncertainties are associated with precipitation characteristics over the western
3 Himalayan region (WHR). These are due to typically small-scale but high intensity storms caused
4 by the complex topography which are und-er-resolved by a sparse gauge network. Additionally,
5 both satellite and gauge precipitation measurements remain subject to systematic errors, typically
6 resulting in underestimation over mountainous terrains. Reanalysis datasets provide prospective
7 alternative but are limited by their resolution, which has so far been too coarse to properly resolve
8 orographic precipitation. In this study, we evaluate and cross-compare Indian Monsoon Data
9 Assimilation and Analysis (IMDAA), the first high-resolution (12 km) regional reanalysis over
10 India, against various precipitation products during winter season over WHR. We demonstrate
11 IMDAA's efficiency in representing winter precipitation characteristics at seasonal, diurnal,
12 interannual scales, as well as heavy precipitation associated with western disturbances (WDs).
13 IMDAA shows closer agreement to other reanalyses than to gauge-based and satellite products in
14 error and bias analysis. Although depicting higher magnitudes, its fine resolution allows a much
15 closer insight into localized spatial patterns and diurnal cycle, a key advantage over other datasets.
16 Mean winter precipitation over WHR shows a significant decreasing trend in IMDAA, despite no
17 significant trend in the frequency of WDs tracked in either IMDAA or ERA5. The study also
18 exhibits the potential use of IMDAA for characterizing winter atmospheric dynamics, both for
19 climatological studies and during WD activity such as localized valley winds. Overall, these
20 findings highlight the potential utility for IMDAA in conducting monitoring and climate change
21 impact assessment studies over the fragile western Himalayan ecosystem.

22 **Keywords:** Precipitation, Western Himalayas, IMDAA reanalysis, Western Disturbances

23

24 1. Introduction

25 North India receives one-third of its annual precipitation (Dimri *et al.*, 2016; Hunt *et al.*, 2018)
26 during the winter season (December through March), in the form of snowfall and rainfall, primarily
27 associated with synoptic-scale extra-tropical cyclonic systems, known as western disturbances
28 (WDs) (Lang and Barros, 2004; Dimri and Mohanty, 2009; Yadav *et al.*, 2013; Dimri, 2013).
29 These extra-tropical cyclonic storms typically originate over the Mediterranean region and travel
30 eastward along the subtropical westerly jet (Dimri *et al.*, 2016, Hunt *et al.*, 2018). Moreover, winter
31 WDs are further known to contribute to approximately 40-50% of the total annual precipitation
32 over the western Himalayas (Madhura *et al.*, 2015; Cannon *et al.*, 2015; Krishnan *et al.*, 2019).
33 Winter precipitation is an important source of irrigation for Rabi crops and plays a critical role in
34 recharging the water resources in the northern plains as well as maintaining the snow cover of the
35 western Himalayan glaciers, which feed major north Indian rivers (Yadav *et al.*, 2013; Dimri *et*
36 *al.*, 2015). Accurate and precise measurement of precipitation is essential for studies involving
37 monitoring and assessing climate change impacts (Hussain *et al.*, 2017). Most products, including
38 gauge-based, satellite, and reanalysis datasets, often face difficulty estimating orographic
39 precipitation over the Himalayas (Sun *et al.*, 2018), since the complex and steep orography of the
40 Himalayas substantially modulates the spatio-temporal variability of regional precipitation on fine
41 scales (Andermann *et al.*, 2011). Thus, reliable precipitation measurements for such complex and
42 heterogeneous landscapes demand high-resolution datasets.

43 Conventionally, in-situ observational data – i.e. gauges – are considered to be one of the
44 most reliable and accurate measurements for precipitation fields at a point scale (e.g. Wang *et al.*,
45 2019). However, uncertainties resulting from measurement errors (e.g. Ye *et al.*, 2004), missing
46 data, insufficient spatial and temporal coverage, etc. (Dahri *et al.*, 2021) offer significant

47 challenges for station-based data analysis. Furthermore, underestimation of precipitation is a key
48 issue associated with rain gauges (e.g. Immerzeel *et al.*, 2015). This can be caused by instrument
49 wetting, pre-measurement evaporation (WMO, 2008), and wind-driven under-catchment, which
50 enhances up to 50 % during snowfall (Dahri *et al.*, 2018, Baudouin *et al.*, 2020). In addition, the
51 network of in-situ observations over the WHR is quite sparse due to varying topography and land
52 cover such as bare rocky surfaces where the possibility of natural hazards including rock falls,
53 avalanches or glacial lake outburst floods create difficulties for gauge installation (e.g. Juen 2006;
54 Carey 2010). Moreover, the spatial distribution of snow cover increases over the region during the
55 winter season and data is missing when stations are buried under snow (e.g. Joshi and Ganju, 2010;
56 Escher-Vetter *et al.*, 2012; Cullen and Conway, 2015; Choudhury *et al.* 2021). Moreover,
57 conventional rain gauges struggle to measure snowfall accurately (Strangeways 2004). Such
58 adverse conditions combined with large orographic variability make it difficult to set up and
59 maintain a dense precipitation-gauge network. As a result, these gauges are generally placed at the
60 foot of the mountains or in valleys, which are relatively drier than elevated regions (Singh and
61 Kumar, 1997; Winiger *et al.* 2005; Dimri and Ganju 2007; Immerzeel *et al.*, 2015; Dahri *et al.*,
62 2018), thus introducing additional uncertainties into gridded datasets, apart from the ones added
63 due to interpolation. Overall, the discontinuity and insufficiency of available observational data
64 reduces the representativeness of gridded observational precipitation products over the region and
65 hinders the production of accurate precipitation estimates and subsequent climate change impact
66 assessment studies over the area.

67 Various remotely sensed and reanalysis precipitation products have been used to
68 compensate for these disadvantages. Often, however, these datasets differ considerably in their
69 spatiotemporal resolution, making intercomparison challenging (Andermann *et al.*, 2011). Satellite

70 precipitation estimates are indirect and often associated with a large degree of variability (Sun *et*
71 *al.*, 2018). Although precipitation retrieval techniques in satellite products have evolved a lot in
72 recent decades (Maggioni *et al.*, 2016), the reliability and degree of precision for these datasets in
73 mountainous regions are still questionable (e.g. Meng *et al.*, 2014; Xia *et al.*, 2015, Xu *et al.*, 2017).
74 An underestimation of orographic precipitation is quite common in infrared (IR) retrievals, given
75 their inability to capture light precipitation events. Detection of cold season orographic
76 precipitation is also challenging for passive microwave retrievals (e.g. Derin and Yilmaz, 2014).
77 In addition, satellite-based microwave retrievals of precipitation rates are inaccurate above snow
78 cover (Derin *et al.*, 2016). Moreover, errors associated with sampling, geo-referencing, and applied
79 algorithms lead to various uncertainties and affect the accurate estimation of precipitation at higher
80 elevations (Hussain *et al.*, 2017). Besides, these products require rain-gauge calibration implying
81 a dependence on the quality and density of station data (Baudouin *et al.* 2020), and thus associated
82 discrepancy gets enhanced over orographic regimes like WHR, where station coverage is quite
83 sparse. In addition, these gridded datasets are typically available only at relatively coarse
84 resolutions, where the leeward and windward sides of mountain areas are generally embedded into
85 a single gridbox, exacerbating the unrepresentative nature of and uncertainties associated with
86 these measurements.

87 Reanalysis datasets provide prospective alternatives for estimating precipitation. These are
88 produced by assimilating observations from a wide range of sources into numerical weather
89 prediction models to generate atmospheric and surface fields. They provide significant advantages
90 in terms of data consistency, homogeneity and coherency, which makes them suitable for
91 atmospheric and climate research (Dee *et al.*, 2014). Globally, reanalysis datasets have been
92 extensively used in precipitation studies owing to their homogeneous nature (Trenberth and

93 Guillemot, 1998; Bengtsson *et al.*, 2004; Bao and Zhang, 2013; Murakami 2014). However, the
94 spatial resolution (more than tens of kilometres) of global reanalysis datasets is often inadequate
95 for effectively capturing localized and regional precipitation distribution, specifically over
96 complex topography, and thus relatively high-resolution regional reanalysis datasets are required
97 to adequately represent regional hydroclimate (Wang *et al.*, 2019; Ashrit *et al.*, 2020).

98 High-resolution data is particularly important in regions with complex topography due to
99 large spatial variability (Gampe *et al.*, 2017). Various studies have highlighted the enhancement
100 of temporal and intensity-related variability associated with WDs over the Himalayan regions due
101 to increased baroclinicity with adverse implications in terms of increased frequency and duration
102 of extreme precipitation events (e.g. Madhura *et al.*, 2015; Midhuna *et al.*, 2020). Such events are
103 often a result of supportive synoptic and mesoscale atmospheric conditions prevailing over the
104 region and the interaction of other processes, including intraseasonal oscillations, local convective
105 dynamics (Gouda *et al.*, 2018) and orographic forcing. The localized nature of these events hinders
106 their accurate assessment in coarse resolution datasets. Therefore, high resolution datasets are
107 crucial for precise understanding of extreme weather related hydrometeorological hazards such as
108 landslides, avalanches and floods over complex topography.

109 Recently, the first high resolution (12 km) regional atmospheric reanalysis dataset, Indian
110 Monsoon Data Assimilation and Analysis (IMDAA), focusing on the South Asian region, has been
111 released (Rani *et al.*, 2021). This state-of-the-art reanalysis is generated by the National Centre for
112 Medium Range Weather Forecasting (NCMRWF) in collaboration with the India Meteorological
113 Department (IMD) and UK Met Office under the National Monsoon Mission project, Government
114 of India. The reanalysis spans the modern meteorological satellite era (1979–present) and outputs
115 atmospheric data at 63 vertical levels. This is now one of the highest resolution regional reanalyses

116 available over India, providing the important advantage of improved representation of orographic
117 features. Ashrit *et al.* (2020) and Aggarwal *et al.* (2022) explored the efficiency of IMDAA in
118 representing precipitation characteristics as well as atmospheric thermodynamics and circulation
119 during the Indian summer monsoon, but wintertime precipitation characteristics are yet to be
120 evaluated using IMDAA.

121 Some validation studies of different multi-source datasets have been carried out previously
122 over the Himalayan region. Andermann *et al.* (2011) evaluated various precipitation datasets along
123 the Himalayan front and reported a significant variation in performance among the evaluated
124 datasets along the orography, however, the datasets show higher consistency, with respect to each
125 other, along with the lower relief realms. Palazzi *et al.* (2013) showed that various category gridded
126 precipitation datasets adequately captured the interannual variability of precipitation over the
127 Hindukush-Karakoram region. Dahri *et al.* (2018) reported underestimation in the rain gauge
128 measurements in the high-altitude Indus basin during the winter season and underpinned the
129 necessity of bias adjustment to reduce errors. A cross-validation for different categories of
130 precipitation datasets in the Indus River basin was performed by Baudouin *et al.* (2020), reporting
131 a large difference in average precipitation between the rain gauge and the reanalyses, most likely
132 resulting from opposite biases from both dataset types and not only from the reanalysis, as often
133 suggested. While precipitation in ERA-Interim (ERA-I) is found to be well correlated with
134 observational data across the Karakoram (Immerzeel *et al.*, 2015; Dahri *et al.*, 2016), Hussain *et*
135 *al.* (2017) reported a poor spatial correlation for TRMM-3B42 with in-situ observations, however,
136 an increase in the correlation was observed with decreasing temporal resolution from daily to
137 monthly scale. Furthermore, it was found that the estimation of summer precipitation compared to
138 winter precipitation in TRMM-3B42 was more accurate in the Hindu Kush-Karakoram Himalayan

139 region. The effect of elevation on the performance of gridded precipitation datasets has also been
140 observed (Andermann *et al.*, 2011; Hussain *et al.*, 2017). A statistical performance evaluation
141 study for different precipitation gridded datasets was conducted by Kanda *et al.* (2020), which
142 highlighted the need for bias correction in different datasets over the WHR.

143 The present study focuses on evaluating winter precipitation variability using high
144 resolution Indian reanalysis (IMDAA, 1979-2018) over diurnal, subseasonal, seasonal and
145 interannual timescales across the WHR. In this study, we will cross-compare and validate the
146 performance of IMDAA against a range of different precipitation datasets from various sources,
147 including gridded observational data, satellite-based and reanalysis products. The results will then
148 be contextualised by comparing the western Himalayan atmospheric dynamics in IMDAA and the
149 fifth generation European Centre for Medium-Range Weather Forecasts (ECMWF) Reanalysis
150 (ERA5) dataset (Hersbach *et al.*, 2018) during the winter monsoon, with an additional focus on
151 the trend and variability of WDs.

152 **2. Data and Methods**

153 *a. Data*

154 Our study focuses on the validation of IMDAA using different categories of gridded precipitation
155 datasets, discussed below, and summarized in Table 1, for the respective periods of their
156 availability from 1979-2018 during the winter season (December to March) over the WHR (29°N-
157 37.5°N and 72.5°E-80.5°E, see Figure 1a). The selection criteria for different categories of datasets
158 in this evaluation study are the availability of long-term (~20 years) precipitation estimates, and at
159 least daily temporal resolution, which is consistent with the availability period of IMDAA. The
160 types of datasets evaluated in the present study include reanalysis products, satellite and gridded

161 observational datasets, as well as combinations thereof. We also use various atmospheric fields
162 from IMDAA and ERA5 for the assessment of wintertime meteorological conditions.

163 **1) Gauge-based Datasets**

164 Three commonly used daily precipitation datasets generated through the interpolation of only rain-
165 gauge measurements have been utilised in our study. IMD precipitation data (0300 UTC-0300
166 UTC) provides daily gridded rainfall (Pai *et al.*, 2014) generated from a dense network of 6955
167 rain gauge stations with varying periods of availability. Daily rainfall estimates are interpolated
168 from gauges to a $0.25^\circ \times 0.25^\circ$ grid, following Shepard (1968). However, the station network suffers
169 from low spatial coverage of gauges over the western Himalayan belt, with almost non-existent
170 stations over the rugged terrains of the upper Himalayan and Karakoram ranges. We selected
171 another gauge-based daily precipitation product (0000 UTC-0000 UTC) provided by the Climate
172 Prediction Centre (CPC) of the National Oceanic and Atmospheric Administration (NOAA) which
173 is constructed from a global gauge-network of around 30,000 stations (Xie *et al.*, 2007; Chen *et*
174 *al.*, 2008). However, limited rain-gauge stations have been considered from the WHR. Covering
175 an extended period of over 50 years, the Asian Precipitation Highly-Resolved Observational Data
176 Integration towards Evaluation of Water Resources (APHRODITE) dataset provides long-term
177 daily precipitation data (0000 UTC-0000 UTC) generated from a dense network of in-situ rain
178 gauges (5000–12000 stations), interpolated at a resolution of 0.05° with an orographic correction
179 for precipitation, and further re-gridded to $0.25^\circ \times 0.25^\circ$ resolution using area-weighted mean
180 (Yatagai *et al.*, 2012). Our study combines APHRO_V1101 (1951–2007) and its extended version
181 APHRO-V1101EX_R1 (2007–2015) to obtain long term precipitation records over the region,
182 following previous literature (Ji *et al.*, 2020; Guan *et al.*, 2020; Lalande *et al.*, 2021; Liaqat *et al.*,
183 2021; Phung *et al.*, 2021).

184 **2) Reanalysis Datasets**

185 Precipitation estimates in different reanalyses can vary from each other based on the assimilation
186 scheme used, the underlying model (including parameterizations) and assimilated observations.
187 The Indian Monsoon Data Assimilation and Analysis (IMDAA) is a high resolution (12km)
188 regional atmospheric reanalysis over the South Asian region, developed by NCMRWF in
189 collaboration with UK Met Office and IMD. IMDAA obtains lateral boundary conditions from
190 ERAI and precipitation estimates are generated by the Unified atmospheric model and the four-
191 dimensional variational (4D-Var) data assimilation technique, which assimilates various
192 conventional and satellite observations from the ECMWF, NCMRWF and IMD archives,
193 including surface observations (land and ocean), aircraft data, upper air observations from
194 radiosondes and pilot balloons. However, no precipitation measurements are assimilated in
195 IMDAA. For further information, the reader is referred to Rani *et al.* (2021). Our study utilizes
196 IMDAA-generated precipitation at 0000 UTC. The dataset provides advantages in better
197 representation of orographic features owing to its high resolution.

198 We also used a state-of-the-art global reanalysis dataset, ERA5 (0000 UTC-0000 UTC),
199 developed by ECMWF with a new version of their NWP model (IFS Cycle 41r2; Hersbach *et al.*,
200 2020). Using 4D-Var, ERA5 assimilates observations from the ECMWF data archive, National
201 Centers for Environmental Prediction (NCEP), as well as other conventional datasets such as ISPD
202 and ICOADS, satellite observations, and precipitation measurements. ERAI, an older ECMWF
203 reanalysis product (Dee *et al.*, 2011) and predecessor of ERA5 has also been used in this study,
204 but only for tracking WDs since it is used to provide lateral boundary conditions in IMDAA. In
205 addition, NCEP's Climate Forecast System Reanalysis (CFSR, 0000 UTC-0000 UTC) is used,
206 which provides precipitation estimates using a coupled atmosphere–ocean model comprising the

207 Global Forecast System and the Geophysical Fluid Dynamics Laboratory Modular Ocean Model
208 (Saha *et al.*, 2010, 2014).

209 Another reanalysis used for validation is the Modern-Era Retrospective Analysis for
210 Research and Applications version-2 (MERRA-2, 0000 UTC-0000 UTC), published by the
211 National Aeronautics and Space Administration (NASA)'s Global Modeling and Assimilation
212 Office. This reanalysis is generated using the Goddard Earth Observing System Model-5, which
213 assimilates various land surface and satellite observations. The advanced data assimilation
214 techniques used in MERRA-2 provide an advantage over topographic regions with sparse gauges,
215 though an underestimation for winter seasonal precipitation has been reported (Hamal *et al.*, 2020).

216 **3) Satellite and Merged Datasets**

217 Our study also utilises five satellite datasets. The widely used Tropical Rainfall Measuring Mission
218 (TRMM)-Multi-Satellite Precipitation Analysis (0300 UTC-0300 UTC), is developed through a
219 collaboration between NASA and Japan's National Space Development Agency. It combines
220 precipitation data from various satellite instruments (TRMM Microwave Imager, Precipitation
221 Radar, Visible and IR Scanner, Special Sensor Microwave Imager), blended with geostationary IR
222 data, with further calibration using monthly gauge data (Huffman *et al.*, 2007).

223 We also used the Precipitation Estimation from Remotely Sensed Information using
224 Artificial Neural Networks-Climate Data Record (PERSIANN-CDR) developed by NOAA in
225 collaboration with the Centre for Hydrometeorology and Remote Sensing, University of
226 California, Irvine. The precipitation estimates (0000 UTC-0000 UTC) are generated by applying
227 the PERSIANN algorithm on GridSat-B1 IR satellite data followed by a training of the artificial
228 neural network using NCEP Stage IV hourly precipitation data and finally calibrating and
229 adjustment of biases is done using the Global Precipitation Climatology Project (GPCP) monthly

230 precipitation dataset (GPCPv2.2). Daily precipitation records are also produced by GPCP Version
231 1.3 (0000 UTC-0000 UTC), from the World Climate Research Programme, by merging estimates
232 from IR, microwave, and sounder data of precipitation-related satellites and gauge-based analyses.

233 A merged high-resolution satellite product, Integrated MultisatellitE Retrievals (V3) for
234 Global Precipitation Measurement (GPM), generates precipitation estimates (0000 UTC-0000
235 UTC) by the day-1 IMERG algorithm, through intercalibrating, merging, and interpolating
236 microwave and IR estimates of GPM satellite constellation with gauge-based observational data
237 (Huffman *et al.*, 2015). The data is specifically useful over regions with a lack of ground-based
238 precipitation-measuring instruments.

239 Lastly, we also use the Climate Hazards Group InfraRed Precipitation with Station data
240 (CHIRPS). This is a merged, daily, land-only precipitation product (0000 UTC-0000 UTC) by the
241 U.S. Geological Survey in association with Earth Resources Observation and Science Centre and
242 is prepared by merging $0.05^{\circ} \times 0.05^{\circ}$ and $0.25^{\circ} \times 0.25^{\circ}$ resolution satellite IR cold cloud duration
243 measurements with in-situ gauge observations.

244 *b. Methodology*

245 1) EVALUATION INDICES

246 Generally, precipitation datasets are utilised without any modifications or prior adjustments for
247 analysing localized precipitation patterns, and it has been now established that the spatial
248 resolution strongly affects dataset performance (Zandler *et al.*, 2019). Moreover, interpolation
249 techniques can generate substantial variations and uncertainties in the computed statistical metrics,
250 especially when precipitation is considered over a complex region. Therefore, we choose to include
251 this information in our evaluation, since the focus is on whether there are advantages offered by
252 the high resolution of IMDAA. We thus compare different datasets in their native resolutions

253 without any interpolation, similar to previous literature (e.g. Dinku *et al.*, 2008; Liu *et al.*, 2015;
254 Beck *et al.*, 2017; Bayissa *et al.*, 2017; Zandler *et al.*, 2019). Different statistical metrics, including
255 mean, standard deviation, coefficient of variation and pattern correlation coefficient (PCC), have
256 been computed to understand how the representation of regional precipitation varies between
257 datasets. All the metrics have been calculated for a common time period (2000–2015), using
258 regionally averaged seasonal mean precipitation at the native spatial resolution of each product,
259 except PCC, for which datasets have been re-gridded using bilinear interpolation to IMDAA’s
260 resolution. We also investigate the total number of heavy precipitation events per year exceeding
261 selected percentile (90th, 95th and 99th) thresholds in each dataset. The number of events has been
262 counted by considering precipitation estimates from all grids between the period 2000-2015,
263 therefore, the datasets have been re-gridded here to a common resolution of 0.25°, to avoid
264 additional counts from higher resolution datasets.

265 To quantify IMDAA’s skill, we further computed different skill scores of the model
266 agreement with the data, using area-averaged (though at original spatial resolution) daily time
267 series of precipitation in different datasets for a common time period 2000-2015. The definitions
268 for commonly known evaluation indices such as correlation coefficient, root mean square
269 difference (RMSD), mean absolute difference (MAD) and BIAS have been summarized in the
270 Appendix, except for less widely used metrics including index of agreement (IOA), adjusted R-
271 squared, and pattern correlation coefficient (PCC), which are discussed below.

272 (i) *Index of agreement*

273 Index of agreement (IOA) is widely used to measure how well model-produced estimates simulate
274 observed data (Willmott 1981) and thus has been used here to quantify the similarity between
275 IMDAA and other datasets (Gebregiorgis *et al.*, 2018). It is defined as:

276
$$IOA = 1 - \frac{\sum_{i=1}^n (x_i - y_i)^2}{\sum_{i=1}^n (|y_i - \bar{x}| + |x_i - \bar{x}|)^2} ,$$

277 Where, n is the total number of observations (daily time series of precipitation), y_i are the values
278 for target dataset (i.e., IMDAA) and, x_i and \bar{x} are the values and mean for the corresponding
279 evaluation dataset, respectively.

280 *(ii) Adjusted R-squared*

281 Adjusted R-squared is a statistical measure to indicate the amount of variance in the dependent
282 variable by the regression model in the population by replacing biased estimators with their
283 unbiased counterparts (Karch 2020). It considers the number of variables in the predictor
284 (IMDAA) and penalizes the supplementary explanatory variables by adjusting the degrees of
285 freedom while estimating the error variance. It provides the statistical quantification of variability
286 amounts associated with a dataset compared to a reference dataset with an adjustment for bias.
287 Additional details have been provided in the Appendix.

288 *(iii) Pattern Correlation Coefficient (PCC)*

289 PCC is the Pearson product-moment coefficient of linear correlation between two variables with
290 values at corresponding locations on two different maps (Anand *et al.*, 2018).

291 2) CLIMATOLOGY, VARIABILITY AND TRENDS

292 The spatial distribution of seasonal (DJFM) winter precipitation climatology and variability in all
293 datasets, considering their respective native spatial resolution and temporal availability between
294 1979–2018 over the study region has been investigated. Full available time periods for each dataset
295 between 1979-2018 have been considered for the evaluation of the geographical distribution of
296 precipitation estimates, since spatial patterns of precipitation tend to be less affected by the

297 averaging period. We further analysed spatial patterns of seasonal mean precipitation differences
298 in IMDAA with respect to different datasets, after rescaling all datasets to the grid used in IMDAA.
299 The spatial patterns on trends of seasonal mean precipitation are computed through seasonal mean
300 precipitation series at each grid point for different precipitation products for a common period of
301 35 years (1984–2018), at their native spatial resolutions, using the non-parametric Mann-Kendall
302 test (Roxy *et al.*, 2015). The datasets with at least an available time period of 35 years have been
303 selected for spatial trends considering that at least 30 years or more is a minimal period to provide
304 considerable climatological trend information. To investigate the interannual variability of the
305 seasonal mean precipitation, standardized precipitation anomalies over the WHR have been
306 utilized to categorize excess and deficit precipitation years, considering the standardized anomalies
307 exceeding ± 0.5 , respectively.

308 3) IDENTIFICATION OF WDs

309 The tracking of WDs over the study region was carried out using the WD tracking algorithm from
310 Hunt *et al.* (2018). The algorithm involves the computation of mean relative vorticity in the 450–
311 300 hPa layer followed by filtration of short-wavelength noise using the desired spectral truncation
312 (T63 and T42) and further identification of positive-definite vorticity regions to determine the
313 centroid locations for each candidate WD. The linkage of these centroids in time, with steering
314 wind-biased distance constraints, allows the identification of potential WD tracks. Lastly, the
315 database is refined by rejecting those tracks which do not pass-through north India (20°N – 36.5°N ,
316 60°E – 80°E), last less than 48 hours, or dissipate at a point westward of their genesis. WDs were
317 tracked in IMDAA (1980–2018) and ERA5 (1979–2018) at spectral truncations of T42 and T63
318 and in ERAI (1979–2015) at T63. The interannual frequency of identified winter WDs was
319 compared for IMDAA (T42 and T63), ERA5 (T42 and T63) and ERAI to analyse the interannual

320 variability of WDs at seasonal scale. ERAI is used as a lateral boundary condition in both IMDAA
321 (Rani *et al.*, 2021) thus, ERAI tracks have also been considered to analyse the frequency of WDs.
322 The T63 ERAI tracks are identical to those used in Hunt *et al.* (2018).

323 4) WINTER ATMOSPHERIC DYNAMICS

324 In order to evaluate the performance of IMDAA in capturing the mean circulation features during
325 the winter season, the climatology of various atmospheric variables such as upper-level winds (200
326 hPa), outgoing longwave radiation (OLR), geopotential heights (200 hPa, 500 hPa, 850 hPa) and
327 2-m air temperature have been compared with ERA5. Further, the fidelity of IMDAA in simulating
328 localized dynamics associated with WD activity over the WHR has been examined through a case
329 study for an intense WD which affected WHR during 16–19 February 2003 and caused widespread
330 precipitation over the region.

331 3. Results and Discussion

332 *a. Climatology and Variability*

333 The spatial distribution of multi-year seasonal mean precipitation in different gridded datasets over
334 WHR is presented in Figure 1 and regionally averaged seasonal precipitation estimates, including
335 mean and variability for a common temporal period (2000–2015) are provided in Table 2. Table 2
336 also details the pattern correlations and the number of heavy precipitation events (exceeding the
337 90th, 95th, and 99th percentiles) during the common period. Considerable heterogeneity for mean
338 winter precipitation amounts and variability is evident among different categories of datasets
339 (Table 2), with all reanalyses (IMDAA, ERA5, CFSR, MERRA-2) showing higher magnitudes
340 than the satellite products (GPCP, TRMM, PERSIANN, CHIRPS, IMERG) and gauge-based
341 datasets (IMD, CPC, APHRODITE). IMDAA, with a mean of 2.38 mm/day and standard deviation

342 of 0.74 mm/day (Table 2), shows a good agreement overall with the other reanalyses ERA5 and
343 CFSR, albeit with a slightly higher magnitude. Among the reanalyses, MERRA-2 has the lowest
344 mean and largest variability, which might be associated with the use of dry-biased CPC data for
345 surface precipitation flux (Reichle *et al.*, 2017; Baudouin *et al.*, 2020). The realistic representation
346 of the mean amounts of winter precipitation of IMDAA is evident through its observed similarity
347 to the ERA5 reanalysis, which has been revealed to perform best with observations in the
348 surrounding upper Indus Basin in previous studies (Baudouin *et al.*, 2020; Dahri *et al.*, 2021). It is
349 evident that all reanalyses, including IMDAA, are wetter and exhibit large spatial variability
350 compared to other types of datasets. Among the observational datasets, mean precipitation in
351 IMDAA is closest to IMD, followed by APHRODITE (Table 2), whereas it shows largest
352 differences compared to CPC. Insufficient observations and uncertainties arising from
353 interpolation techniques generally lead to dry biases in observational datasets (e.g. Dahri *et al.*,
354 2021). An overall general agreement within the satellite data category can be observed. However,
355 IMDAA shows an overestimation compared to satellite products, the exception being IMERG with
356 higher resolution. The precipitation amount and variability for IMERG is highest among satellite
357 datasets, much closer to IMDAA. As satellite observations generally exhibit higher accuracy for
358 convective precipitation over flat terrains (Ebert *et al.*, 2007; Baudouin *et al.*, 2020), their
359 usefulness is limited over the elevated western Himalayan terrain that typically receives more
360 stratiform precipitation (Fig. S2).

361 Wide discrepancies in the geographical seasonal mean precipitation patterns (Fig. 1) among
362 the different categories of datasets can be seen, though with an agreement across the regions of the
363 highest precipitation – the lower Himalayas and foothills. As indicated by the statistics (Table 2),
364 IMDAA stands much closer to all reanalyses compared to other data categories in depicting

365 regional spatial precipitation patterns (Fig. 1), having lower spatial precipitation differences with
366 IMDAA (Fig. 2). The IMD observational dataset shows a similar spatial distribution of
367 precipitation magnitude in the lower Himalaya and the western Karakoram range, but considerable
368 discrepancies are noticeable along the foothills and in eastern Ladakh (Fig. 1), where the IMD data
369 set has a marked positive difference compared to all other data sets (Fig. 2). Since IMD includes
370 almost no measurements over these locations (Kishore *et al.*, 2016), the precipitation estimates
371 here are derived from extrapolation of higher values downslope, resulting in higher precipitation
372 estimates. Relatively wetter patterns (Fig. 1) and a large positive difference are seen in IMDAA
373 (Fig. 2) compared to APHRODITE and CPC, both of which rely on the WMO Global
374 Telecommunication System, which covers few observations in WHR, generally collected from
375 stations present over dry valley locations in the study region, thus, additionally suffering from
376 under-catchment of solid precipitation at higher altitudes (Palazzi *et al.*, 2013; Dahri *et al.*, 2018).
377 This makes the validation of IMDAA precipitation over this region using gauge-based datasets
378 challenging. IMDAA exhibits a larger magnitude of precipitation than all the satellite datasets
379 (Table 3, Fig. 2), with the smallest differences over the foothills in IMERG, which has a similar
380 resolution to IMDAA. However, it can be noted from the previous studies (Dahri *et al.*, 2021;
381 Baudouin *et al.*, 2020) that precipitation from satellite products exhibit drier biases over the region.
382 Various uncertainties in terms of interpolation algorithms, calibration limitations due to poor
383 coverage of rain-gauge measurements, and inability to accurately retrieve orographic precipitation
384 have been denoted as causes for the same (Hussain *et al.*, 2017). However, differences in
385 precipitation magnitude are mainly found over lower Himalayas and foothills in IMDAA with all
386 datasets (Fig. 2). These regions receive maximum precipitation during winter, thus, such
387 differences highlight discrepancies and limitations associated with available data over the region.

388 Higher pattern correlations (Table 2, Fig. 1) between IMDAA and all other datasets are observed,
389 least being for GPM-IMERG, indicating an agreement on the spatial patterns of seasonal mean
390 precipitation for IMDAA with other datasets, even though significant differences in magnitudes
391 are present. Additionally, the spatial patterns of solid precipitation distributions in IMDAA are
392 similar to independent MODIS-satellite snow cover fractions (Fig. S6).

393 Summing up, the gridded products provide inconsistent precipitation amounts over the
394 region, but agree on the areas with the highest precipitation. Since different dataset categories are
395 generated with different input data and dissimilar developmental methods, presence of such similar
396 signals relates to the depiction of actual situation (Baudouin et al., 2020). Unlike satellite and
397 gauge-based products, reanalysis products are developed with different data assimilation
398 techniques and distinct atmospheric models (Ghodichore *et al.*, 2018) and are known to provide a
399 better depiction of frontal system precipitation in the winter season, specifically over high
400 elevations (Dahri et al., 2021; Beck et al., 2019), compared to other two product categories which
401 generally underestimate precipitation in such cases. IMDAA performs well in simulating regions
402 with precipitation maxima and allows a much closer look at the localized precipitation distribution
403 over the region. However, reanalyses generally tend to produce higher precipitation magnitudes
404 along with a depiction of larger variability, which in fact is highest in IMDAA, highlighting a key
405 limitation of the current IMDAA reanalysis.

406 Adjusted- R^2 values (Table 3) are computed for area-weighted daily precipitation values
407 during 2000-2015 and significance has been tested using Mann-Kendall test at a confidence level
408 of 95%. These also underpin the observed discrepancies among different datasets over the study,
409 with large variation for values observed over the panels of different datasets. Here, we carry out a
410 cross-validation by examining the association between individual datasets by using each dataset

411 as a reference. Highest agreements are observed within the reanalyses; however, MERRA-2 shows
412 lesser association supporting the negative differences observed earlier (Table 2). Observational
413 datasets IMD and CPC also show good agreement with each other, however, APHRODITE
414 diverges from the group and is also the dataset showing the least association with all other
415 categories too. Satellite products are in good agreement with each other and generally with datasets
416 from other categories too, except for MERRA-2 and APHRODITE; both of which also have lower
417 fitness magnitudes with all other datasets. The reanalyses, IMDAA, ERA5 and CFSR, depict a
418 significant relationship with all the datasets, specifically with each other and IMD. Considering
419 that reanalyses and observations generally rely on different data sources and developmental
420 methods, a significant relationship among these is a sign of quality of the datasets (Baudouin *et*
421 *al.*, 2020), thus, implying the reliability of their precipitation estimates.

422 The total number of heavy precipitation events per winter season exceeding the 90th, 95th,
423 and 99th percentiles (considering the wide range of discrepancies for precipitation amounts across
424 different datasets) considering all grid locations for each dataset during the period 2000-2015 have
425 also been evaluated (Table 2). For each dataset (re-gridded to common resolution of 0.25°), the
426 precipitation values during the common period are collapsed into a single vector to which the
427 percentile thresholds computed from area-averaged daily precipitation time series (2000-2015) are
428 applied and totals counts exceeding the thresholds are selected. It is to be noted that since large
429 uncertainties remain over the region regarding precipitation amounts, we use the percentile
430 thresholds rather than actual values. Overall, the obtained counts indicate close agreement of
431 IMDAA with all reanalyses. However, the number of events in MERRA-2 have lower counts
432 compared to other reanalyses at all three thresholds. An overall agreement can be observed within
433 the satellite category among TRMM, CHIRPS and PERSIANN-CDR, exceptions being IMERG

434 and GPCP (highest and lowest resolution data among satellite products, respectively), though
435 slight overestimation in IMDAA with respect to all satellite datasets (except IMERG) can be
436 observed. However, gauge-based datasets exhibit a sharp disagreement among each other and with
437 respect to other data categories too for all percentiles, with APHRODITE showing a sharp
438 overestimation as compared to all the evaluated datasets, including IMDAA.

439 The wintertime precipitation over the WHR comprises both snowfall and rainfall, with
440 snowfall constituting almost 80% of the total observed precipitation (e.g. Krishnan *et al.*, 2019).
441 Therefore, the differences between mean rainfall and snowfall in IMDAA and ERA5 have also
442 been analyzed. Compared to ERA5, IMDAA shows higher amount of snowfall (~3 mm/day) over
443 the Greater Himalayas and some regions of the Lower and Karakoram Himalayas (Fig. S1). The
444 variation in rainfall between IMDAA and ERA5 shows larger variations in IMDAA over some
445 regions of lower Himalayas. Along the foothills, the differences in rain and snow often have
446 different signs, implying that local temperature differences play a role.

447 The analysis of interannual variability patterns of seasonal mean winter precipitation in
448 different datasets has been used to identify pluvial and dry years (Fig. 3). IMDAA is able to
449 reproduce the pluvial and deficit precipitation years, corresponding well with other datasets. All
450 reanalyses as well as PERSIANN-CDR and IMD datasets depict an enhancement in the frequency
451 of dry periods in recent years. IMDAA, along with other datasets, agrees that 2001 was one of the
452 driest years. Decadal variability is evident in most of the analyzed datasets, which might be
453 attributable to forcing by large scale tropical climate drivers such as Arctic Oscillation, North
454 Atlantic Oscillation and Pacific Decadal Oscillation (e.g. Roy 2006). Overall, the interannual
455 variability of wet and dry years in IMDAA is in good agreement with the other datasets,
456 particularly IMD and ERA5.

457 *b. Trends*

458 Seasonal mean precipitation trends over the study region as a whole (Fig. 4i-k) reveals close
459 agreement among reanalyses, depicting similar values of mean and decreasing trend – although
460 IMDAA is the only one among them in which that trend is significant at the 95% confidence level
461 (Mann-Kendall test). IMDAA agrees with the gauge-based dataset IMD, with a significant
462 negative trend, but the trends for remaining gauge- and satellite-based datasets are not significant
463 and disagree with each other even on their sign. The spatial patterns of these trends computed over
464 a common period of 35 years (1984–2018) for eight datasets (Fig. 4a-h) corroborate the area-
465 weighted trends in the region. Most of the datasets agree on a negative trend over WHR, which is
466 strongest in IMDAA, IMD, and CFSR, but there is disagreement in terms of regional trends in
467 both the CHIRPS and CPC datasets, which are both predominated by positive trends. It is worth
468 noting that the observed negative trend for seasonal winter precipitation in most of the datasets
469 well agrees with the station based decreasing trends by Shekhar *et al.* (2010, 2017) who found
470 decreasing precipitation rates at different point stations over the region, confirming the reliability
471 of IMDAA data for trend analysis studies. The trend is statistically significant along the foothills
472 and lower Himalayan belt in IMDAA; mostly over western Jammu and Kashmir and western
473 Ladakh in CFSR; and in IMD, over eastern Himachal Pradesh and eastern and central Ladakh.
474 ERA5 agrees strongly with IMDAA on the spatial focus of negative trend, but without any
475 statistical significance. In contrast, positive precipitation trends observed in CPC and CHIRPS
476 show disagreement on where those trends are significant, and PERSIANN-CDR and MERRA-2
477 have regions of both significant positive and negative precipitation trends but disagree with each
478 other on the sign of those trends. A strong positive and statistically significant precipitation trend
479 is visible in CPC over Himachal Pradesh and Uttarakhand, whereas CFSR shows a strong yet

480 statistically insignificant trend is observed over some regions of eastern Jammu and Kashmir. The
481 decreasing spatial and temporal trends in the majority of the evaluated datasets support the earlier
482 observed increasing frequency of dry periods in recent years. Earlier, Dimri and Dash (2012)
483 observed a decreasing trend of winter precipitation over the WHR, and a decrease in the winter
484 snowfall has also been reported by Shekhar *et al.* (2010). Overall, the diversity in winter
485 precipitation trends among datasets is largely due to methodological constraints associated with
486 respective dataset development.

487 *c. Seasonal cycle of Precipitation*

488 A fine representation of seasonal variability in any dataset holds key significance as the seasonal
489 cycle of precipitation has important dynamical implications. Therefore, we also investigated the
490 seasonal cycle of winter precipitation over the study region by comparing area-averaged daily
491 climatology in all datasets for a common period of 2000-2015 for their individual spatial
492 resolutions (Figure 5). All reanalyses, including IMDAA, are in close agreement with each other
493 on the seasonal evolution of precipitation, with very little differences observed at daily
494 climatological scales. However, although gauge-based observations show differences in
495 magnitude, the representation of subseasonal variability is quite similar, with a notably stronger
496 agreement between IMD and IMDAA, compared to other gauge products. IMDAA overestimates
497 the magnitude of the seasonal cycle relative to all the satellite products, with least differences with
498 respect to IMERG, however, they do agree with each other on the form of the seasonal cycle.

499 Summarising so far, IMDAA is in agreement with all other reanalysis datasets, particularly
500 ERA5. All datasets agree that the heaviest precipitation falls in February and the weakest is in
501 December. Furthermore, the results reveal that all reanalyses and the gauge-based IMD dataset
502 produce similar patterns of regional daily precipitation variability.

503 *d. Skill Scores*

504 Various statistical skill scores (Table 4) have been computed for area-weighted daily time series
505 of precipitation for statistical evaluation of IMDAA in comparison to other datasets. IMDAA
506 shows a good correlation with all the datasets ($r > 0.6$), with the highest correlations found among
507 the reanalysis products ($r > 0.9$), followed by gridded observational datasets IMD, CPC and satellite
508 datasets ($r > 0.7$). IMDAA again depicts the lowest correlation with APHRODITE among all
509 datasets, in agreement with earlier computed statistics. Higher correlations among independent
510 datasets with different data sources and generation methods, are indicators of better precipitation
511 estimates in both the datasets (Baudouin *et al.*, 2020), thus good correlations obtained for IMDAA
512 highlight the reliability of its area-mean precipitation estimates. Overall, IMDAA has IOA values
513 closer to 1 with most datasets, with the highest values among reanalyses, supported by low RMSD,
514 MAD and bias with respect to ERA5 and CFSR, but MERRA-2 again diverges from the group.
515 IMDAA has an index of agreement greater than 89% with all reanalyses, around 71% with satellite
516 products, and greater than 70% with gridded gauge-based datasets. IMDAA shows relatively larger
517 positive differences (following Fig. 2) and higher error magnitudes with the gauge-based and
518 satellite datasets. A strong agreement between IMDAA and IMD is evident through all indices,
519 including highest correlation (0.90) and lowest RMSD, MAD and mean difference (bias). Indeed,
520 IMDAA shows positive differences in comparison to all the datasets, supporting the earlier
521 observed higher magnitudes of precipitation in IMDAA over other datasets (Fig. 2).

522 Fig. 6 shows the spatial maps of observed point-to-point correlation for daily precipitation
523 climatology between IMDAA and each of the other datasets for 2000–2015 at each grid point,
524 after regridding all datasets to a common spatial resolution of IMDAA. We see a high degree of
525 point-to-point correlation for all reanalyses over the majority of the region, with slightly lower

526 correlation with MERRA-2 over the Himalayan foothills. IMDAA shows a comparatively weak
527 pointwise correlation with many of the satellite datasets – TRMM, IMERG and GPCP; however,
528 the pointwise correlation is better between with PERSIANN-CDR and CHIRPS over the upper
529 Himalayan ranges. Among the observational products, IMDAA has a good correlation with IMD
530 and CPC over most of the study region, although some divergence from the group is evident in
531 APHRODITE, verifying the earlier obtained statistical metrics. It is promising that the region of
532 greatest agreement, especially among the reanalyses, is along the western Himalayas – implying
533 the existence of a common dynamical source of precipitation (e.g. WDs) in this region to which
534 the reanalyses respond relatively robustly; although it is also possible that this agreement is due to
535 common sources of biases between the reanalyses, such as the interaction between parameterised
536 convection and the orography.

537 *e. Diurnal cycle of Precipitation*

538 The diurnal cycle of different types of precipitation is one of the fundamental aspects of variability
539 but it is often neglected in validation studies. For those datasets with hourly data, we can also
540 compare their representation of the diurnal cycle of winter precipitation over the study region (Fig.
541 7). Realistic representation of the diurnal cycle of precipitation in reanalyses is often challenging
542 due to their reliance on convective parameterization, which typically causes precipitation to occur
543 too early in the day (Dirmeyer *et al.*, 2012). Among the four reanalyses considered here, IMDAA
544 performs best when compared with IMERG, capturing the early morning maximum along much
545 of the western Himalayas and relatively consistent late afternoon peak over the plains south of the
546 Himalaya, both of which are missed to at least some degree by the other reanalyses. IMDAA
547 struggles to simulate the correct diurnal cycle over Tibetan Plateau, with the peak occurring near
548 local midnight rather than the local noon seen in IMERG or mid-afternoon in the other reanalyses.

549 This is a region where much of the precipitation occurs in the form of stratiform from the
550 underlying model (Fig. S2), and thus generated by the microphysics scheme. Given, however, that
551 the peaks in both stratiform and convective precipitation (Fig. S3) appear mistimed here (at least
552 compared with ERA5), this may be a result of IMDAA assimilating far fewer observations outside
553 of India, rather than problems with the model physics. Analysis of precipitation contribution by
554 convective and stratiform fractions reveal a dominance by stratiform precipitation over the
555 Karakoram and Greater Himalayas in both IMDAA and ERA5, whereas the lower Himalayas
556 experience a mixed proportion of both types of precipitation (Fig. S2). The Himalayan foothills
557 and the plains observe more contribution through convective precipitation.

558 *f. Western Disturbances*

559 WDs are upper-level synoptic-scale cyclonic perturbations in the subtropical jet and are the
560 primary contributors of wintertime precipitation over the WHR. About 80% of observed winter
561 precipitation here occurs during days when a WD is active (Midhuna *et al.*, 2020), with the
562 remaining 20% typically being contributed by local scale convective systems. Given the general
563 agreement (Fig. 4) among datasets, including IMDAA, that winter precipitation is declining, and
564 that a large majority of winter precipitation occurs on WD days, we might ask whether the two are
565 linked by a decline in WD frequency. To test this, WDs were tracked in both ERA5 and IMDAA
566 datasets, using two values of spectral truncation (T42 and T63) to account for potential differences
567 in feature size. The trends of tracked WD frequency in these four datasets, along with an additional
568 set of T63 tracks computed using ERAI for Hunt *et al.* (2018), are shown in Figure 8(a). All
569 datasets show a weak and insignificantly increasing trend of winter WDs in the recent decades. An
570 agreement in the pattern of timeseries is observed for coarser grid resolution (T42), but with
571 differences in respective magnitude. The comparatively finer resolution truncation (T63) shows a

572 similarity in patterns for most years but IMDAA generally has more interannual variability
573 compared with ERA5. ERAI also has a similar interannual pattern, but with a reduced count
574 compared to the higher resolution reanalyses. The weaker, insignificant increase for WD frequency
575 indicates that seasonal occurrences of WDs has been almost constant during the last few decades,
576 or that the effect of interannual variability in WD frequency is much higher than the long-term
577 trends. Generally, the trend significance is measured against internal variability, the fact that the
578 trend fails a significance test indicates that the linear trend is weak compared to interannual
579 variability. Moreover, the standard deviation of the seasonal totals is clearly larger than the trend.
580 However, significant decreasing trend of seasonal mean precipitation has been observed (Fig. 4).
581 Similar results have been reported by Shekhar *et al.* (2010), concluding that seasonal (November-
582 April) occurrences of WDs (1984/85-2007/08) have less effect on snowfall patterns over the
583 western Himalayas, however, a decreasing trend for number of snowfall days was observed in their
584 study. On the contrary, Cannon *et al.* (2015) reported an increase in WD occurrences in the region,
585 while Madhura *et al.* (2015) highlighted an increase in the interannual variability of WD frequency
586 in recent decades. Our results highlight the challenges that remain in determining recent trends in
587 WD activity, a fact which is highlighted by disagreement between earlier studies.

588 Further, we carried out the analysis of WD-day precipitation composites (Fig. 8b-p) which
589 show that IMDAA performs well in capturing spatial precipitation details when compared with
590 ERA5, though the two reanalyses have higher magnitudes when compared with IMD. IMD shows
591 slight variations in terms of precipitation amount and location, which could be a result of the sparse
592 gauge density over the Himalayas. There is some variation across the season, with all three datasets
593 agreeing that February sees the heaviest WD-day precipitation and December the lightest, in
594 agreement with the results for the climatology discussed earlier.

595 WD-associated seasonal and monthly precipitation fractions have been plotted by dividing
596 the total precipitation observed during WD-days by the total seasonal precipitation over the study
597 region. The results reveal that 60–90% of wintertime precipitation over the Himalayan region can
598 be attributed to WD activity (Fig. 9), in agreement with previous studies (e.g. Hunt *et al.*, 2019).
599 IMDAA shows a strong agreement with IMD and ERA5, but the higher resolution of IMDAA
600 provides the advantage of a more detailed look into the localized precipitation fractions over the
601 domain, revealing elongated structures orientated northwest-southeast. These features are parallel
602 to the orography and have maximum values on the southern side of local ridges, consistent with
603 WDs bringing moisture flux from the southwest and orographic forcing from regional topography.
604 The attributable fractions for rain and snow (Fig. S4) in IMDAA and ERA5 reveal the dominance
605 of snowfall over the Greater and Karakoram Himalayas, whereas rainfall being the main observed
606 form of precipitation during wintertime over the lower Himalayas and foothills. IMDAA is able to
607 capture the localized variations in WD attributed precipitation percentages owing to its high
608 resolution, whereas ERA5 shows a comparatively homogeneous pattern.

609 *g. Evaluation of dynamical and thermodynamic conditions*

610 As we have seen, winter precipitation over the WHR is primarily associated with WDs. As they
611 are embedded in the large scale sub-tropical westerly jet (200 hPa), WD activity depends
612 significantly on its position and intensity (Krishnan *et al.*, 2019). A comparison of winds at 200
613 hPa for IMDAA with ERA5 shows that IMDAA is realistic in capturing the wintertime subtropical
614 westerly jet (SWJ) over the region (Fig. 10g-i). However, IMDAA produces a weaker SWJ over
615 WHR (Fig. 10i). An assessment of winter-mean climatological conditions of 2-m air temperature
616 (Fig. 10a-c), OLR (Fig. 10d-f) and geopotential heights (Fig. S5) in IMDAA compared to ERA5
617 has been carried out. IMDAA captures the spatial patterns of temperature and OLR over WHR

618 and shows potential in finely representing these dynamical features compared to ERA5. However,
619 IMDAA shows warmer temperatures (statistically significant differences) over the lower
620 Himalayas and foothills extending up to northeastern Himalayas and slightly colder temperatures
621 over some western regions of Greater Himalayas, central Tibetan plateau and WHR compared to
622 ERA5 (Fig. 10c). During winter, low OLR values are noticeable over WHR in both IMDAA (Fig.
623 10d) and ERA5 (Fig. 10e). Generally, lower magnitudes of mean OLR are observed over the region
624 during winter as compared to other seasons owing to the influence of convective activity and cloud
625 formation, although climatological mean OLR is also lower here due to the higher underlying
626 orography. IMDAA exhibits slightly higher OLR magnitudes than ERA5 along the south-eastern
627 Ladakh region and over the Himalayan foothills, extending up to the northeastern Himalayas. The
628 patterns of mean geopotential height (Fig. S5) at different pressure levels over the WHR during
629 the winter season seem to be well represented in both IMDAA and ERA5 with slightly higher
630 magnitudes in IMDAA over WHR. To sum up, IMDAA is capable of representing seasonal mean
631 dynamical and large-scale circulation patterns during winter.

632 *h. Case Study for western disturbance over WHR*

633 As one of the potential major advantages of the higher resolution of IMDAA is its ability to better
634 capture local orographically-driven dynamics, we also explore the representation of such dynamics
635 during the passage of WD over WHR. Here, this is accomplished through analysis of a case study
636 for an intense WD that occurred during 16-19 February 2003, which affected WHR and caused
637 widespread precipitation over the region. Overcast skies and enhanced cloud cover associated with
638 deeper convective activity are key features observed during the passage of WD over WHR (e.g.
639 Rao and Rao 1971), which is noticeable in satellite imagery from NASA EOSDIS (Fig. 11a-d).
640 IMDAA is realistically representing the daily evolution of the total cloud cover associated with

641 the intense WD over WHR (Fig. 11e-h), further supported by abrupt negative OLR anomalies (Fig.
642 11i) during the period indicating increased atmospheric convection. This convective activity plays
643 an essential role in developing secondary circulations during the propagation of WDs and results
644 in heavy localized precipitation over the WHR (e.g. Dimri *et al.*, 2016; Hunt *et al.*, 2019). Our
645 results demonstrate the ability of high resolution IMDAA in representation of realistic evolution
646 of the WD induced cloud cover patterns compared to remotely sensed observations.

647 Furthermore, the analysis of regional valley wind systems during the given period has been
648 carried out to validate the representation of local scale circulation patterns in IMDAA compared
649 to ERA5, which highlights the advantages offered by its high resolution. Figure 12 shows the
650 spatial and temporal patterns of local scale 10-m wind patterns at two different valley sites (the
651 Suru valley in Jammu and Kashmir and the Spiti Valley in Himachal Pradesh) before, during, and
652 after the passage of WD over WHR. IMDAA (Fig 12b) shows high fidelity in capturing the local
653 scale circulation features at the two valley sites, correctly capturing the maximum intensity of
654 winds during the in contrast to the more spatially homogeneous ERA5 wind speed.

655 The temporal evolution of these valley winds during the passage of WD is shown for both
656 valley sites in Fig. 12g-h, where an abrupt increase in magnitudes of valley wind speed is observed
657 on 16 February at both locations. This variability is captured in IMDAA, but not in ERA5 where
658 the wind speed is roughly constant during the passage of the WD. Additionally, the effect of the
659 WD passing over the study region is dynamically characterized by an increase of the local
660 minimum temperature (Fig. 12i-j) and a drop in the surface pressure (Fig. 12k-l), which
661 corroborates previous studies (e.g. Rao and Rao, 1971; Singh *et al.*, 2019). Overall, it is clear that
662 IMDAA shows high fidelity in representing local scale wind responses associated with WD
663 activity over WHR, a key advantage offered by its high resolution.

664 **4. Summary and Conclusions**

665 This study evaluated winter precipitation and its variability and trends in the recently introduced
666 high resolution reanalysis IMDAA using various gridded, satellite, and reanalysis datasets over
667 the western Himalayan region between 1979–2018, as per their respective period of availability.
668 Based on the findings, the following conclusions can be made:

- 669 1. IMDAA captures the spatial variability of winter precipitation over the western Himalayas
670 well, on both seasonal and interannual scales, and has climatological precipitation statistics
671 that are very similar to the other reanalyses considered. However, it shows higher
672 precipitation amounts compared to other datasets along the lower Himalayas and foothills.
- 673 2. IMDAA agrees with the other reanalysis datasets in showing a slight decline in winter
674 precipitation over the western Himalayas over recent decades, though it is the only
675 reanalysis dataset in which that trend is significant, in accordance with trends reported in
676 earlier studies in individual station data (Shekhar *et al.*, 2010, 2017). The IMD gauge-based
677 precipitation dataset also has a significant decline over the same period, but no other gauge-
678 based or satellite products have significant trends when averaged over the whole western
679 Himalayan region.
- 680 3. Tracking of WDs carried out in both ERA5 and IMDAA reanalyses showed a weak
681 insignificant increase in frequency over the study period (1979–2018). This agrees with
682 previous reanalysis-based studies; however, several observation-based studies have
683 reported a recent decline in WD frequency. Our results, thus, highlight the challenges that
684 remain in determining recent trends in WD activity over the region, making it an important
685 area of further work.

686 4. IMDAA shows potential in reproducing climatological winter circulation patterns and
687 surface conditions. However, the validation of high-resolution features such as valley wind
688 speeds and local dynamics during the passage of WDs over the region strongly emphasizes
689 advantages offered by IMDAA's high resolution compared to ERA5. The dataset showed
690 high potential in representing more localized ridge- and valley-scale features, offering a
691 better characterization of regional dynamics.

692 In summary, we acknowledge that low density of gauges and complex orography leads to high
693 discrepancies and uncertainties in the available data products over WHR. Overall, reanalyses
694 – including IMDAA – suffer from precipitation overestimation likely owing to errors in the
695 representation of parameterized convection. However, the findings in this study emphasize the
696 benefits provided by the high-resolution model and output of IMDAA, including
697 understanding the complex interplay between terrain and mountain meteorology over WHR,
698 even with its higher precipitation magnitudes. The reanalysis shows high fidelity in simulating
699 local-scale dynamics as well as large scale circulation features responsible for winter
700 precipitation over the region. Such findings strongly underpin the capabilities of IMDAA in
701 exploring the winter monsoon and its variability and in analysing the meteorological precursors
702 of precipitation extremes. Overall, IMDAA, despite its amplified magnitudes, is useful for
703 precipitation climatology, interannual variability and synoptic meteorology over WHR,
704 however, is still unable to capture diurnal peak precipitation realistically over the Tibetan
705 Plateau. Even with a resolution of 0.12° (~12 km), IMDAA still does not adequately resolve
706 regional orography, and this is likely to continue to result in discrepancies in precipitation
707 magnitude over the region.

708 Moreover, large uncertainties remain in understanding the spatio-temporal precipitation in
709 the WHR due to limited observations and thus, it becomes hard to obtain benchmark
710 precipitation trends, as well as verification of spatial precipitation amounts in IMDAA. Further
711 work is needed to constrain historical trends in winter precipitation in this region and link those
712 trends to changes in synoptic-scale activity, such as western disturbances. Finally, we note that
713 high-resolution simulations cannot replace ground-based in-situ observations and the lack of
714 gauge data over such terrains with high spatial variability adds challenges for accurate
715 precipitation measurements, which can be tackled through increased coverage of in-situ
716 stations over the region.

717 **Acknowledgments.**

718 This research work was supported by Science and Engineering Research Board, Department of
719 Science and Technology, Government of India under the “Start-up Research Grant (SRG) scheme”
720 (Grant reference: SRG/2020/001857). Authors gratefully acknowledge NCMRWF, Ministry of
721 Earth Sciences, Government of India, for IMDAA reanalysis. IMDAA reanalysis was produced
722 under the collaboration between UK Met Office, NCMRWF, and IMD with financial support from
723 the Ministry of Earth Sciences, under the National Monsoon Mission programme. Nischal
724 gratefully acknowledges the financial assistance from the Prime Minister’s Research Fellowship
725 (PMRF), Ministry of Education, Government of India. KMRH is supported through the Weather
726 and Climate Science for Service Partnership (WCSSP) India, a collaborative initiative between the
727 Met Office, supported by the U.K. Government’s Newton Fund, and the Indian Ministry of Earth
728 Sciences (MoES). The authors acknowledge and appreciate the editor and anonymous reviewers
729 for their constructive feedback, which helped in improving the quality of our work.

730 **Data Availability Statement**

731 All the data used in this study is publicly available and accessible. IMDAA data is available on the
732 RDS NCMRWF portal at <https://rds.ncmrwf.gov.in/datasets>. ECMWF fifth generation (ERA5)
733 data can be accessed through [https://www.ecmwf.int/en/forecasts/datasets/reanalysis-](https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5)
734 [datasets/era5](https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5). NCEP Climate Forecast System data is provided at <https://cfs.ncep.noaa.gov/cfsr>
735 and MERRA-2 is available at <https://gmao.gsfc.nasa.gov/reanalysis/MERRA-2>. The gauge-based
736 datasets can be accessed from
737 https://www.imdpune.gov.in/Clim_Pred_LRF_New/Grided_Data_Download.html (IMD),
738 <https://psl.noaa.gov/data/gridded/data.cpc.globalprecip.html> (CPC) and
739 <http://aphrodite.st.hirosaki-u.ac.jp/products.html> (APHRODITE). GPCP satellite data is available
740 at <https://www.ncei.noaa.gov/products/climate-data-records/precipitation-gpcp-daily> and
741 TRMM-3B42 at <https://disc.gsfc.nasa.gov/datasets/>. The remaining datasets used in the study are
742 publicly accessible at <https://www.chc.ucsb.edu/data> (CHIRPS),
743 <https://gpm.nasa.gov/data/directory> (GPM-IMERG) and <https://chrsdata.eng.uci.edu/>
744 (PERSIANN-CDR), respectively. WD days during DJFM over the study region are quantified
745 based on Indian Daily Weather Reports (IDWR) issued by the IMD.

746

747 **APPENDIX**

748 *Evaluation indices*

749 The quantitative assessment of IMDAA in comparison to different datasets was carried out using
750 a series of statistical indicators and skill scores for the daily time series of winter precipitation. The
751 indices used for statistical evaluation of the performance of IMDAA with other datasets are

752 discussed below where n is the total number of observations (daily time series of precipitation), x_i
 753 depicts the value for respective evaluation dataset, y_i depicts the value for IMDAA dataset, \bar{x} is
 754 the mean for evaluation dataset values and \bar{y} is the mean for IMDAA dataset values.

755 Pearson's correlation coefficient (r) is a quantitative measure of strength of linear
 756 agreement between two datasets which ranges from -1 to +1, with positive values indicating a
 757 positive correlation and vice versa.

$$758 \quad r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2}} \quad (1)$$

759
 760 The Root Mean Square Difference (RMSD) is a measure of overall difference associated with
 761 residuals in a predictor dataset in comparison to the validation dataset, whereas relative RMSD
 762 (rRMSD) is a normalized variant of RMSD with respect to the mean of the validation dataset.
 763 Lower values of RMSD and rRMSD indicate stronger association between datasets.

$$764 \quad RMSD = \sqrt{\frac{\sum_{i=1}^n (x_i - y_i)^2}{n}} \quad (2)$$

$$765 \quad RMSD_{rel} = \frac{RMSD}{\bar{x}} * 100 \quad (3)$$

766 Mean Absolute Difference (MAD) is a measure of the accuracy of a predictor dataset in terms of
 767 average magnitude of errors present in the predictions in comparison to the validation dataset.
 768 rMAD is a normalized variant of MAD.

$$769 \quad MAD = \frac{1}{n} \sum_{i=1}^n |x_i - y_i| \quad (4)$$

$$770 \quad rMAD = \frac{MAD}{\bar{x}} * 100 \quad (5)$$

771 BIAS is the quantitative measure to define the tendency of underestimation or overestimation in a
772 dataset with respect to a validation dataset where the negative values indicate underestimation and
773 vice versa.

$$774 \quad BIAS = \frac{1}{n} \sum_{i=1}^n (y_i - x_i) \quad (6)$$

$$775 \quad rBIAS = \frac{BIAS}{\bar{x}} * 100 \quad (7)$$

776 Adjusted R-squared is defined as:

$$777 \quad Adj. R^2 = 1 - (1 - R^2) \frac{n-1}{n-k-1}$$

778 where, k is the number of independent variables and n is the number of observations.

779 REFERENCES

780 Aggarwal, D., R. Attada., K.K. Shukla, R. Chakraborty, and R.K. Kunchala, 2022: Monsoon precipitation
781 characteristics and extreme precipitation events over Northwest India using Indian high-resolution
782 regional reanalysis. *Atmos. Res.*, 267, 105993, <https://doi.org/10.1016/j.atmosres.2021.105993>.

783 Anand, A. S. K. Mishra, S. Sahany, M. Bhowmick, J.S. Rawat, and S. K. Dash, 2018: Indian summer
784 monsoon simulations: Usefulness of increasing horizontal resolution, manual tuning, and semi-
785 automatic tuning in reducing present-day model biases. *Sci. rep.*, **8(1)**, 1-14.

786 Andermann, C., S. Bonnet, and R. Gloaguen, 2011: Evaluation of precipitation data sets along the
787 Himalayan front. *Geochem. Geophys. Geosyst.*, **12(7)**, Q07023.

788 Ashrit, R., and Coauthors, 2020: IMDAA Regional Reanalysis: Performance Evaluation during Indian
789 Summer Monsoon Season. *J. Geophys. Res. Atmos.*, **125(2)**, e2019JD030973,
790 <https://doi.org/10.1029/2019JD030973>.

791 Bao, X., and F. Zhang, 2013: Evaluation of NCEP-CFSR, NCEP-NCAR, ERA-Interim, and ERA-40
792 Reanalysis Datasets against Independent Sounding Observations over the Tibetan Plateau. *J.*
793 *Climate*, **26(1)**, 206-214.

794 Baudouin, J. P., M. Herzog, and C. A. Petrie, 2020: Cross-validating precipitation datasets in the Indus
795 River basin. *Hydrol. Earth Syst. Sci.*, **24(1)**, 427-450, <https://doi.org/10.5194/hess-24-427-2020>.

796 Bayissa, Y., T. Tadesse, G. Demisse and A. Shiferaw, 2017: Evaluation of Satellite-Based Rainfall
797 Estimates and Application to Monitor Meteorological Drought for the Upper Blue Nile Basin,
798 Ethiopia. *Remote Sens.*, 669.

799 Beck, H. E. and Co-authors, 2017: Global-scale evaluation of 22 precipitation datasets using gauge
800 observations and hydrological modeling. *Hydrol. Earth Syst. Sci.*, **21**, 6201–6217.

801 Bengtsson, L., S. Hagemann, and K. I. Hodges, 2004: Can Climate Trends be calculated from Re-Analysis
802 data?. *J. Geophys. Res. Atmos.*, **109(D11111)**.

803 Cannon, F., L. M. Carvalho, C. Jones, and B. Bookhagen, 2015: Multi-annual variations in winter westerly
804 disturbance activity affecting the Himalaya. *Climate Dyn.*, **44(1-2)**, 441-455.

805 Carey, M., 2010: In the Shadow of Melting Glaciers. Climate Change and Andean Society; Oxford
806 University Press: Oxford, UK, 288.

807 Chen, M., P. Xie, and Coauthors, 2008: CPC Unified Gauge-based Analysis of Global Daily Precipitation.
808 *Western Pacific Geophys. Meeting*, Cairns, Australia, 29 July-1 August, 2008.

809 Choudhury, A., A. C. Yadav, and S. Bonafoni, 2021: A response of snow cover to the climate in the
810 northwest himalaya (NWH) using satellite products. *Rem. Sens.*, **13(4)**, 655.

811 Cullen, N. J. and J. P. Conway, 2015: A 22-month record of surface meteorology and energy balance from
812 the ablation zone of Brewster Glacier, New Zealand. *J. Glaciol.*, **61**, 931–946.

- 813 Dahri, Z. H., and Co-authors, 2021: Spatio-temporal evaluation of gridded precipitation products for the
814 high-altitude Indus basin. *Int. J. Climatol.*, **41(8)**, 4283-4306.
- 815 Dahri, Z. H., E. Moors, F. Ludwig, S. Ahmad, Khan, A. Khan, I. Ali, and P. Kawat, 2018: Adjustment of
816 measurement errors to reconcile precipitation distribution in the high-altitude Indus basin. *Int. J.*
817 *Climatol.*, 38.
- 818 Dahri, Z. H., F. Ludwig, E. Moors, B. Ahmad, A. Khan, and P. Kabat, 2016: An appraisal of precipitation
819 distribution in the high-altitude catchments of the Indus basin. *Sci. Total Environ.*, **548**, 289-306.
- 820 Dee, D. P., M. Balmaseda, G. Balsamo, R. Engelen, A. J. Simmons, and J.-N. Thépaut, 2014: Toward a
821 consistent reanalysis of the climate system. *Bull. Amer. Meteor. Soc.*, **95(8)**, 1235–1248.
- 822 Dee, D. P., S. M. Uppala, A. J. Simmons, P. Berrisford, P. Poli, S. Kobayashi, and F. Vitart, 2011: The
823 ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q. J. R.*
824 *Meteorol. Soc.*, **137(656)**, 553-597.
- 825 Derin, Y., and K. K. Yilmaz, 2014: Evaluation of multiple satellite-based precipitation products over
826 complex topography. *J. Hydrometeor.*, **15(4)**, 1498-1516.
- 827 Derin, Y., E. Anagnostou, A. Berne, M. Borga, B. Boudevillain, W. Buytaert, and K. K. Yilmaz, 2016:
828 Multiregional satellite precipitation products evaluation over complex terrain. *J.*
829 *Hydrometeor.*, **17(6)**, 1817-1836.
- 830 Dimri, A. P., 2013: Intraseasonal oscillation associated with the Indian winter monsoon. *J. Geophys. Res.*
831 *Atmos.*, **118**, 1189– 1198.
- 832 Dimri, A. P., and A. Ganju, 2007: Wintertime seasonal scale simulation over Western Himalaya using
833 RegCM3. *Pure Appl. Geophys.*, **164(8)**, 1733-1746.
- 834 Dimri, A. P., and S. K. Dash, 2012: Wintertime climatic trends in the western Himalayas. *Climate*
835 *Change*, **111(3)**, 775-800.

836 Dimri, A. P., and U. C. Mohanty, 2009: Simulation of mesoscale features associated with intense Western
837 Disturbances over western Himalayas. *Meteorol. Appl.*, **16(3)**, 289-308.

838 Dimri, A. P., D. Niyogi, A. P. Barros, J. Ridley, U. C. Mohanty, T. Yasunari, and D. R. Sikka,
839 2015: Western Disturbances: A review. *Rev. Geophys.*, **53(2)**, 225– 246.

840 Dimri, A. P., T. Yasunari, B. S. Kotlia, U. C. Mohanty, and D. R. Sikka, 2016: Indian winter monsoon:
841 Present and past. *Earth Sci. Rev.*, **163(2)**, 297-322.

842 Dinku, T., S. J. Connor, P. Ceccato, and C.F. Ropelewski, 2008: Comparison of global gridded precipitation
843 products over a mountainous region of Africa. *Int. J. Climatol.*, **28**,1627–1638.

844 Dirmeyer, P. A., and Coauthors, 2012: Simulating the diurnal cycle of rainfall in global climate models:
845 resolution versus parameterization. *Climate Dyn.*, **39(1)**, 399-418.

846 Ebert, E. E., J. E. Janowiak, and C. Kidd, 2007: Comparison of near-real-time precipitation estimates from
847 satellite observations and numerical models. *Bull. Amer. Meteor. Soc.*, **88(1)**, 47-64.

848 Escher-Vetter, H., and Co-authors, 2012: Hydrological and Meteorological Records from the Vernagtferner
849 Basin—Vernagtbach Station, for the Years 1970 to 2001. PANGAEA: Bremen, Germany.

850 Forsythe N., S. Blenkinsop, and H. J. Fowler, 2015: Exploring objective climate classification for the
851 Himalayan arc and adjacent regions using gridded data sources. *Earth Syst. Dyn.*, **6(1)**, 311–326.

852 Funk, C., and Coauthors, 2015: The climate hazards infrared precipitation with stations-a new
853 environmental record for monitoring extremes. *Sci. Data*, **2**, 150066.

854 Gampe, D., and R. Ludwig, 2017: Evaluation of gridded precipitation data products for hydrological
855 applications in complex topography. *Hydrology*, **4(4)**, 53.

856 Gebregiorgis, A. S., and Co-authors, 2018: To what extent is the day 1 GPM IMERG satellite precipitation
857 estimate improved as compared to TRMM TMPA-RT?. *J. Geophys. Res.: Atmos.*, **123(3)**, 1694-
858 1707.

859 Gelaro, R., and Coauthors, 2017: The Modern-Era Retrospective Analysis for Research and Applications,
860 Version 2 (MERRA2). *J. Climate*, **30(14)**, 5419-5454.

861 Ghodichore, N., R. Vinnarasi, C. T. Dhanya, and S.B. Roy, 2018: Reliability of reanalyses products in
862 simulating precipitation and temperature characteristics over India. *J. Earth Syst. Sci.*, **127(8)**, 1-
863 21.

864 Gouda, K. C., S. Nahak, and P. Goswami, 2018: Evaluation of a GCM in seasonal forecasting of extreme
865 rainfall events over continental India. *Wea. Climate Extrem.*, **21**, 10-16.

866 Guan, X., and Co-authors, 2020: Evaluation of Precipitation Products by Using Multiple Hydrological
867 Models over the Upper Yellow River Basin, China. *Remote Sens.*, **12(24)**, 4023.

868 Hamal, K., and Coauthors, 2020: Evaluation of MERRA-2 Precipitation Products Using Gauge Observation
869 in Nepal. *Hydrology*, **7(3)**, 40, <https://doi.org/10.3390/hydrology7030040>.

870 Hersbach, H., and Coauthors, 2020: The ERA5 global reanalysis. *Quart. J. Roy. Meteor. Soc.*, **146(730)**,
871 1999-2049. <https://doi.org/10.1002/qj.3803>.

872 Huffman, G. J., and Coauthors, 2007: The TRMM Multisatellite Precipitation Analysis (TMPA): Quasi-
873 Global, Multiyear, Combined-Sensor Precipitation Estimates at Fine Scales. *J. Hydrometeorol.*, **8(1)**,
874 38-55.

875 Huffman, G. J., and Coauthors, 2015: NASA Global Precipitation Measurement (GPM) Integrated Multi-
876 satellite Retrievals for GPM (IMERG). Algorithm Theoretical Basis Document (ATBD)
877 Version, **4**, 26.

878 Huffman, G. J., R. F. Adler, M. M. Morrissey, D. T. Bolvin, S. Curtis, R. Joyce, B. McGavock, and J.
879 Susskind, 2001: Global precipitation at One-degree Daily Resolution from Multi-satellite
880 Observations. *J. Hydrometeor.*, **2(1)**, 36-50.

881 Hunt, K. M. R., A. G. Turner, and L. C. Shaffrey, 2018: The evolution, seasonality and impacts of western
882 disturbances. *Quart. J. Roy. Meteor. Soc.*, **144(710)**, 278-290.

883 Hunt, K. M. R., A. G. Turner, and L. C. Shaffrey, 2019: Representation of Western Disturbances in CMIP5
884 Models. *J. Climate*, **32(7)**, 1997-2011.

885 Hussain, S., X. Song, G. Ren, I. Hussain, H. Dongmei, and M. H. Zaman, 2017: Evaluation of gridded
886 precipitation data in the Hindu Kush–Karakoram–Himalaya mountainous area. *Hydrol. Sci.*
887 *J.*, **62(14)**, 2393-2405.

888 Immerzeel, W. W., N. Wanders, A. F. Lutz, J. M. Shea, and M. F. P. Bierkens, 2015: Reconciling high-
889 altitude precipitation in the upper Indus basin with glacier mass balances and runoff. *Hydrol. Earth*
890 *Syst. Sci.*, **19(11)**, 4673-4687.

891 Ji, X., and Co-authors, 2020: Evaluation of bias correction methods for APHRODITE data to improve
892 hydrologic simulation in a large Himalayan basin. *Atmos. Res.*, **242**, 104964.

893 Joshi, J. C., and A Ganju, 2010: Use of objective analysis to estimate winter temperature and precipitation
894 at different stations over western Himalaya. *J. Earth Syst. Sci.*, **119(5)**, 597-602.

895 Juen, I., 2006: Glacier Mass Balance and Runoff in the Cordillera Blanca, Peru. Ph.D. Thesis, University
896 of Innsbruck, Innsbruck, Austria.

897 Kanda, N., H. S. Negi, M. S. Rishi, and A. Kumar, 2020: Performance of various gridded temperature and
898 precipitation datasets over Northwest Himalayan Region. *Environ. Res. Commun.*, **2(8)**, 085002,
899 <https://doi.org/10.1088/2515-7620/ab9991>.

900 Karch, J., 2020: Improving on Adjusted R-Squared. *Collabra:Psychology*, **6(1)**, 45.

901 Kishore, P., S, Jyothi, G. Basha, S. Rao, M. Rajeevan, I. Velicogna, and T.C. Sutterley, 2016: Precipitation
902 climatology over India: validation with observations and reanalysis datasets and spatial trends.
903 *Climate Dyn.*, **46(1-2)**,541–556.

904 Krishnan, R., T. P. Sabin, R. K. Madhura, R. K. Vellore, M. Mujumdar, J. Sanjay, S. Nayak, and M.
905 Rajeevan, 2019: Non-monsoonal precipitation response over the Western Himalayas to climate
906 change. *Climate Dyn.*, **52(7)**, 4091-4109.

907 Lalande, M., M. Ménégoz, G. Krinner, K. Naegeli, and S. Wunderle, 2021: Climate change in the High
908 Mountain Asia in CMIP6. *Ear sys. dyn.*, **12(4)**, 1061-1098.

909 Lang, T. J., and A. P. Barros, 2004: Winter storms in Central Himalayas, *J. Meteor. Soc. Japan*, **82(3)**, 829–
910 844.

911 Liaqat, M. U., G. Grossi, S. ul Hasson, and R. Ranzi, 2022: Characterization of Interannual and Seasonal
912 Variability of Hydro-Climatic Trends in the Upper Indus Basin. *Theor. Appl. Climatol.*, **147**, 1163–
913 1184, <https://doi.org/10.1007/s00704-021-03850-3>.

914 Liu, M. and Co-authors, 2015: Evaluation of high-resolution satellite rainfall products using rain gauge data
915 over complex terrain in southwest China. *Theor. Appl. Climatol.* 119, 203–219.

916 Madhura, R. K., R. Krishnan, J. V. Revadekar, M. Mujumdar, and B. N. Goswami, 2015: Changes in
917 western disturbances over the Western Himalayas in a warming environment. *Climate Dyn.*, **44**,
918 1157-1168.

919 Maggioni, V., Meyers, P., & Robinson, M. (2016). A review of merged high resolution satellite
920 precipitation product accuracy during the Tropical Rainfall Measuring Mission (TRMM)-Era.
921 *Journal of Hydrometeorology*, 17(4), 1101–1117.

922 Meng, J., L. Li, Z. Hao, J. Wang, and Q. Shao, 2014: Suitability of TRMM satellite rainfall in driving a
923 distributed hydrological model in the source region of Yellow River. *J. Hydrol.*, **509**, 320-332.

924 Midhuna, T. M., P. Kumar, and A. P., Dimri, 2020: A new Western Disturbance Index for the Indian winter
925 monsoon. *J. Earth Syst. Sci.*, **129**, 59, <https://doi.org/10.1007/s12040-019-1324-1>.

926 Murakami, H., 2014: Tropical cyclones in reanalysis data sets. *Geophys. Res. Lett.*, **41(6)**, 2133-2141.

927 Nguyen, P., and Coauthors, 2019: The CHRS Data Portal, an easily accessible public repository for
928 PERSIANN global satellite precipitation data. *Sci. Data*, 6(1), 180296.

929 Nitu, R. and, Co-authors., 2018: WMO Solid Precipitation Intercomparison Experiment (SPICE) (2012–
930 2015). Instruments and Observing Methods Report No. 131, World Meteorological Organization,
931 Geneva. <https://www.wmo.int/pages/prog/www/IMOP/publications-IOM-series.html>.

932 Pai, D. S., L. Sridhar, M. Rajeevan, O. P. Sreejith, N. S. Satbhai, and B. Mukhopadhyay, 2014:
933 Development of a new high spatial resolution ($0.25^\circ \times 0.25^\circ$) long period (1901-2010) daily gridded
934 rainfall data set over India and its comparison with existing data sets over the
935 region. *Mausam*, **65(1)**, 1-18.

936 Palazzi, E., J. V. Hardenberg, and A. Provenzale, 2013: Precipitation in the Hindu-Kush Karakoram
937 Himalaya: Observations and future scenarios. *J. Geophys. Res.: Atmos.*, **118(1)**, 85-100.

938 Phung, D., T. Nguyen-Huy, N.N. Tran, D.N. Tran, S. Nghiem, N.H. Nguyen, and T. Bennett, 2021:
939 Hydropower dams, river drought and health effects: A detection and attribution study in the lower
940 Mekong Delta Region. *Clim. Risk Manag.*, **32**, 100280.

941 Rani S. I., T. Arulalan, J. P. George, E. N. Rajagopal, R. Renshaw, A. Maycock, D. M. Barker, and M.
942 Rajeevan, 2021: IMDAA: High-Resolution Satellite-Era Reanalysis for the Indian Monsoon
943 Region, *J. Climate*, **34(12)**, 5109-5133, <https://doi.org/10.1175/JCLI-D-20-0412.1>.

944 Rao, B. V., and Rao, T. S., 1971: A theoretical and synoptic study of western disturbances. *Pure Appl.*
945 *Geophys.*, **90(1)**, 193-208.

- 946 Reichle, R. H., Q. Liu, R. D. Koster, C. S. Draper, S. P. Mahanama, and G. S. Partyka, 2017: Land surface
947 precipitation in MERRA-2. *J. Clim.*, **30(5)**, 1643-1664.
- 948 Roxy, M., K. Ritika, P. Terray, R. Murtugudde, A. Karumuri and B. N. Goswami, 2015: Drying of Indian
949 subcontinent by rapid Indian Ocean warming and a weakening land-sea thermal gradient. *Nat.*
950 *Commun.*, 7423 (2015).
- 951 Roy, S. S., 2006: The impacts of ENSO, PDO, and local SSTs on winter precipitation in India. *Phy. Geog.*,
952 *27(5)*, 464-474.
- 953 Saha, S., and Coauthors, 2010: The NCEP Climate Forecast System Reanalysis. *Bull. Amer. Meteor.*
954 *Soc.*, **91(8)**, 1015-1058.
- 955 Saha, S., and Coauthors, 2014: The NCEP Climate Forecast System Version 2. *J. Climate*, **27(6)**, 2185-
956 2208.
- 957 Shekhar M. S., H. Chand, S. Kumar, K. Srinivasan, and A. Ganju, 2010: Climate change studies in the
958 western Himalaya; *Ann. Glaciol.*, **51(54)**, 105–112.
- 959 Shekhar, M. S., N. N. Rao, S. Paul, S.C. Bhan, G. P. Singh, and A. Singh, 2017: Winter precipitation
960 climatology over Western Himalaya: Altitude and Range wise study. *J. Ind. Geophys. Union*,
961 *21(2)*,148-152.
- 962 Singh, D., A. Kumar, M. S. Shekhar, and S. Garg, 2019: The impacts of the approaching western
963 disturbances (WDs) on the surface meteorological variables over the north-west Himalaya (NWH),
964 India. *J. Earth. Syst. Sci.*, 149.
- 965 Singh, P., and N. Kumar, 1997: Effect of orography on precipitation in the western Himalayan region. *J.*
966 *Hydrol.*, **199(1-2)**, 183-206.
- 967 Strangeways, I., 2004: Improving precipitation measurement. *Int. J. Climatol.*, **24(11)**, 1443–1460.

- 968 Sun, Q., C. Miao, Q. Duan, H. Ashouri, S. Sorooshian, and K.-L. Hsu, 2018: A Review of Global
969 Precipitation Data Sets: Data Sources, Estimation, and Intercomparisons. *Rev. Geophys.*, **56(1)**, 79-
970 107.
- 971 Trenberth, K. E., and C. J. Guillemot, 1998: Evaluation of the atmospheric moisture and hydrological cycle
972 in the NCEP/NCAR reanalyses. *Climate Dyn.*, **14(3)**, 213-231.
- 973 Wang, Y., J. Chen, and D. Yang, 2019: Bayesian Assimilation of Multiscale Precipitation Data and Sparse
974 Ground Gauge Observations in Mountainous Areas. *J. Hydrometeor.*, **20(8)**, 1473-1494.
- 975 Willmott, C. J., 1981: On the validation of models. *Phys. geo.*, **2(2)**, 184-194.
- 976 Winiger, M. Y., M. Gumpert, and H. Yamout, 2005: Karakorum–Hindukush–western Himalaya: assessing
977 high-altitude water resources. *Hydrol. Process.*, **19(12)**, 2329-2338.
- 978 Xia, T., Z. J. Wang, and H. Zheng, 2015: Topography and Data Mining Based Methods for Improving
979 Satellite Precipitation in Mountainous Areas of China. *Atmos.*, **6(8)**, 983-1005.
- 980 Xie, P., A. Yatagai, M. Chen, T. Hayasaka, Y. Fukushima, C. Liu, and S. Yang, 2007: A gauge-based
981 analysis of daily precipitation over East Asia, *J. Hydrometeor.*, **8**, 607-626.
- 982 Xu, R., F. Tian, L. Yang, H. Hu, H. Lu, and A. Hou, 2017: Ground validation of GPM IMERG and TRMM
983 3B42V7 rainfall products over southern Tibetan Plateau based on a high-density rain gauge
984 network. *J. Geophys. Res.: Atmos.*, **122(2)**, 910-924.
- 985 Yadav R. K., D. A. Ramu, and A. P. Dimri, 2013: On the relationship between ENSO patterns and winter
986 precipitation over North and Central India. *Glob. Planet. Change*, **107**, 50-58.
- 987 Yadav, R. K., R. Kumar, and M. Rajeevan, 2009: Increasing influence of ENSO and decreasing influence
988 of AO/NAO in the recent decades over northwest India winter precipitation. *J. Geophys. Res.:*
989 *Atmos.*, 114(D12).

- 990 Yatagai, A., K. Kamiguchi, O. Arakawa, A. Hamada, N. Yasutomi, and A. Kitoh, 2012: APHRODITE:
991 Constructing a Long-Term Daily Gridded Precipitation Dataset for Asia Based on a Dense Network
992 of Rain Gauges. *Bull. Amer. Meteor. Soc.*, **93(9)**, 1401-1415.
- 993 Ye, B., D. Yang, Y. Ding, T. Han, and T. Koike, 2004: A bias-corrected precipitation climatology for
994 china. *J. Hydrometeor.*, **5(6)**, 1147–1160.
- 995 Zandler, H., I. Haag, and C. Samimi, C., 2019: Evaluation needs and temporal performance differences of
996 gridded precipitation products in peripheral mountain regions. *Nature:Sci. rep.*, **9(1)**, 1-15.