

Gene-diet interactions on metabolic disease-related outcomes in Southeast Asian populations: a systematic review

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Ventura, E. F., Sekar, P., Dhanapal, A. C. T. A., Cheah, E. S. G. ORCID: https://orcid.org/0000-0001-7633-9455, Loganathan, A., Quen, P. L., Appukutty, M., Taslim, N. A., Hardinsyah, H., Noh, M. F. M., Lovegrove, J. A. ORCID: https://orcid.org/0000-0001-7633-9455, Givens, I. ORCID: https://orcid.org/0000-0002-6754-6935 and Vimaleswaran, K. S. ORCID: https://orcid.org/0000-0002-8485-8930 (2023) Gene-diet interactions on metabolic disease-related outcomes in Southeast Asian populations: a systematic review. Nutrients, 15 (13). 2948. ISSN 2072-6643 doi: 10.3390/nu15132948 Available at https://centaur.reading.ac.uk/112675/

It is advisable to refer to the publisher's version if you intend to cite from the work. See <u>Guidance on citing</u>.

To link to this article DOI: http://dx.doi.org/10.3390/nu15132948

Publisher: MDPI

All outputs in CentAUR are protected by Intellectual Property Rights law,



including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the End User Agreement.

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Remiern



Gene–Diet Interactions on Metabolic Disease-Related Outcomes in Southeast Asian Populations: A Systematic Review

Padmini Sekar¹, Eduard Flores Ventura¹, Anto Cordelia T. A. Dhanapal², Eddy Seong Guan Cheah², Annaletchumy Loganathan², Phoon Lee Quen², Mahenderan Appukutty^{3,4}, Nurpudji Astuti Taslim⁵, Hardinsyah Hardinsyah⁶, Mohd Fairulnizal Md Noh⁷, Julie A Lovegrove¹, Ian Givens⁸, and Karani Santhanakrishnan Vimaleswaran^{1,8,*}

- ¹ Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, Institute for Cardiovascular and Metabolic Research (ICMR), University of Reading, Reading RG6 6DZ, UK; p.sekar@pgr.reading.ac.uk (P.S.); eduard.flores.ventura@hotmail.com (E.F.V.); j.a.lovegrove@reading.ac.uk (J.A.L.)
- ² Centre for Biomedical and Nutrition Research, Universiti Tunku Abdul Rahman, Jalan Universiti, Bandar Barat, Kampar 31900, Malaysia; antoc@utar.edu.my (A.C.T.A.D.); cheahsg@utar.edu.my (E.S.G.C.); annal@utar.edu.my (A.L.); phoonlg@utar.edu.my (P.L.Q.)
- ³ Faculty of Sports Science and Recreation, Universiti Teknologi MARA, Shah Alam 40450, Malaysia; mahen@uitm.edu.my
- ⁴ Nutrition Society of Malaysia, Jalan PJS 1/48 off Jalan Klang Lama, Petaling Jaya 46150, Malaysia
- ⁵ Clinical Nutrition, Faculty of Medicine, Hasanuddin University, Makassar 90245, Indonesia; pudji_taslim@yahoo.com
- ⁶ Department of Community Nutrition, Faculty of Human Ecology, IPB University, Bogor 16680, Indonesia; hardinsyah2010@gmail.com
- ⁷ Institute for Medical Research, National Institutes of Health, Jalan Setia Murni U13/52, Seksyen U13 Setia Alam, Shah Alam 40170, Malaysia; fairulnizal@moh.gov.my
- Institute for Food, Nutrition and Health (IFNH), University of Reading, Reading RG6 6AH, UK; d.i.givens@reading.ac.uk
- * Correspondence: v.karani@reading.ac.uk

Abstract: Diabetes and obesity are chronic diseases that are a burden to low- and middle-income countries. We conducted this systematic review to understand gene-diet interactions affecting the Southeast Asian population's risk of obesity and diabetes. The literature search was performed on Google Scholar and MEDLINE (PubMed) search engines independently by four reviewers who evaluated the eligibility of articles based on inclusion criteria. Out of 19,031 articles, 20 articles examining gene-diet interactions on obesity and/or diabetes-related traits met the inclusion criteria. Three (Malaysia, Indonesia, and Singapore) out of eleven Association of Southeast Asian Nations (ASEAN) countries have conducted studies on gene-diet interactions on obesity and diabetes. From the 20 selected articles, the most common interactions were observed between macronutrients and genetic risk score (GRS) on metabolic disease-related traits in the Malay, Chinese, and Indian ethnicities. Overall, we identified 29 significant gene-diet interactions in the Southeast Asian population. The results of this systematic review demonstrate ethnic-specific gene-nutrient interactions on metabolicdisease-related traits in the Southeast Asian population. This is the first systematic review to explore gene-diet interactions on obesity and diabetes in the Southeast Asian population and further research using larger sample sizes is required for better understanding and framing nutrigenetic approaches for personalized nutrition.

Keywords: systematic review; nutrigenetics; Southeast Asia; genetics; gene–diet interaction; dietary intake; obesity; diabetes; metabolic disease



Citation: Sekar, P.; Ventura, E.F.; Dhanapal, A.C.T.A.; Cheah, E.S.G.; Loganathan, A.; Quen, P.L.; Appukutty, M.; Taslim, N.A.; Hardinsyah, H.; Md Noh, M.F.; et al. Gene–Diet Interactions on Metabolic Disease-Related Outcomes in Southeast Asian Populations: A Systematic Review. *Nutrients* **2023**, *15*, 2948. https://doi.org/10.3390/ nu15132948

Academic Editor: Aristides G. Eliopoulos

Received: 30 May 2023 Revised: 16 June 2023 Accepted: 21 June 2023 Published: 29 June 2023



Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// creativecommons.org/licenses/by/ 4.0/).

1. Introduction

Metabolic diseases such as obesity and diabetes are now considered epidemics rapidly spreading across developed and developing countries affecting both sexes, age, ethnicities, and socioeconomic groups [1]. This, in turn, has shown to compromise the quality of life that leads to potentially life-threatening conditions such as cancers, cardiovascular diseases, musculoskeletal disorders, and hypertension [2]. According to the 2021 reports from the World Health Organisation (WHO), worldwide obesity has tripled since 1975, with over 650 million obese adults and more than 340 million children and adolescents who are either overweight or obese [3]. By 2022, diabetes reports from WHO indicate that more than 422 million people are diabetic with rising prevalence in low-middle income group countries (LMIC) compared to developed countries [4]. Projected trends also show that diabetes and obesity are rapidly growing and will affect nearly two-thirds of the Southeast (SE) Asian population by the end of 2030, placing a burden on rural and low socioeconomic groups [5,6]. Recent reports from the Association of Southeast Asian Nations (ASEAN) show that the tripling rate of undernutrition has not improved and that obesity and diabetes are now a double burden for these countries [7,8].

Understanding gene–nutrient interactions provide insights regarding nutritional, genetic, and biochemical determinants to better understand complex interactions between environmental factors (including diet) and genes relevant to metabolic health and diseases [9,10]. Several studies have also reported the importance of physical activity and nutrient intake which potentially interact with genetic predispositions of a disease that promote the progression and pathogenesis of metabolic diseases [11]. Many studies have also reported the influence of certain gene–diet interactions on metabolic disease-related traits and emphasized the importance of a healthy lifestyle that may modify the outcome of the disease or its related parameters [10–15]. A better comprehension of the relationship between genes and diet is key to making correlations between nutrition and wellness, thereby allowing for specific nutritional suggestions that are tailored to individuals or genetic subgroups. This strategy presents an appropriate public health approach [9].

The increasing prevalence of diabetes and obesity in SE Asia can be understood by the nutrition transition phenomenon, environment multiplier theory, and the thrifty gene hypothesis [16,17]. These theories provide an understanding of the dietary shift from traditional high-carbohydrate, low-fat diets towards high-energy diets (high saturated fat, sugars, and salt), and the role of inherited genetic predispositions in over-nutrition-related diseases. Dietary factors can affect the outcome of a disease and there are ethnic-specific genetic variations that influence the mechanism of these nutrient interactions [2]. Furthermore, lifestyle/dietary factors could influence genetic predispositions of metabolic disorders, especially obesity, and diabetes [18–21], making nutrigenetics research a necessity in ethnically diverse populations such as SE Asia. Nutrition science along with a better understanding of nutrigenetics in different ethnic groups is important to improvise personal and societal health [2,20]. This would ultimately add to the efforts of implementing precision nutrition specific to the populations [22,23]. Hence, this systematic review examines gene-diet interactions on metabolic disease-related (diabetes and obesity) outcomes in the 11 SE Asian countries (Brunei, Burma (Myanmar), Cambodia, Timor-Leste, Indonesia, Laos, Malaysia, the Philippines, Singapore, Thailand, and Vietnam) that share sufficient socio-demographic and cultural similarities.

2. Materials and Methods

2.1. Study Identification and Source Strategy

To identify studies involving gene–diet interactions on metabolic disease-related outcomes, a literature search was undertaken until February 2023 using MEDLINE (via PubMed), and Google Scholar search engines (Supplementary Table S1). The reference lists of the included papers and independent search strings used by the researchers were examined until saturation. In PubMed, extensive search was performed using the search string: (polymorphism OR gene OR SNP OR single nucleotide polymorphism OR genetic variation OR genetic variant OR GRS OR genetic risk score OR PRS OR polygenic risk score) AND ("gene-diet interaction" OR "diet-gene interaction" OR SNP-diet interaction OR diet-SNP interaction OR "gene-nutrient interaction" OR "nutrient-gene interaction" OR "gene-lifestyle inter-action" OR "gene-environment interaction") AND (carbohydrate OR protein OR fat OR fiber OR sugar OR SFA OR saturated fat OR monounsaturated fat OR polyunsaturated fat OR MUFA OR PUFA OR diet OR B12 OR vitamin D OR amino acids OR polyphenols OR egg intake OR caffeine intake OR green tea OR alcohol intake OR meat intake OR energy intake OR food) AND (obesity OR weight OR BMI OR waist circumference OR waist hip ratio OR hip circumference OR adiposity OR metabolic diseases OR body fat OR body composition) AND (Southeast Asia OR Malay* OR Brunei* OR Burm* OR Cambodia* OR Timor* OR Indonesia* OR Laos OR Filipin* OR Philippine* OR Singapore* OR Thai* OR Vietnam*). In Google Scholar, an extensive search was performed using the search string: gene-diet interaction BMI Southeast Asia OR Malay* OR Brunei* OR Burm* OR Cambodia* OR Timor* OR Indonesia* OR Laos OR Filipin* OR Philippine* OR Singapore* OR Thai* OR Vietnam*. The literature search was restricted to studies involving human subjects only.

2.2. Data Extraction

The reviewers (E.V.F., P.S., A.C.T.A.D., and E.S.G.C.) ensured data consistency across the articles extracted for this study, and a narrative synthesis was conducted to compile the data sourced. Duplicate articles were eliminated using EndNote. Titles and abstracts were subjected to blind screening to assess the pre-established inclusion criteria, followed by full-text screening and discussion. The study protocol was submitted to PROSPERO (Identification number: CRD42022366475).

2.3. Study Selection: Inclusion and Exclusion Criteria

Related studies published in PubMed and Google Scholar in the English language were included. Only gene–nutrient interaction (nutrigenetic) studies examining the association between dietary factors and genes on diabetes and/or obesity-related outcomes were included. Eligible articles on clinical studies, multicentre studies, comparative studies, observational studies, and randomized controlled studies were included. Studies on patients, neonates, children, and pregnant women were included. The study included populations from ASEAN countries namely Brunei, Cambodia, Indonesia, Laos, Malay-sia, Myanmar, the Philippines, Singapore, Thailand, Timor-Leste, and Vietnam. Studies were excluded if they were (1) animal studies; (2) did not include gene–diet interactions; (3) the outcome was not diabetes- and/or obesity-related traits; (4) not examined in the SE Asian population; or (5) nutrigenomic studies (gene expression in response to dietary factors).

2.4. Data Items and Effect Measures

Obesity, diabetes, and parameters associated with anthropometric measurements and attributes were considered primary outputs of data extraction (Tables 1 and 2). The result of interactions between the exposure (genetic and dietary factors) and outcome (obesity- or diabetes-related traits) was estimated using *p*-values extracted from the included literature. Based on the statistical output, the interactions were considered significant if the $P_{interaction}$ values were below 0.05. In this study, a narrative synthesis was conducted to elaborate on the dietary factors, genetic variation, and disease traits.

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
					Mala	ysia				
FADS1	rs174547	C-S	200	Chinese	>18	Linoleic acid (7.9 \pm 3.6 g/day)	WC	0.177	No significant interaction; vegetarians with TT	[24]
mbor	1517 1517	CO	(69/131)	Indians	210	α-Linolenic Aaid (0.4–0.8 g/day)		0.258	of MetS, larger WC, and low HDL-c	[= 1]
Maternal VDR	rs2228570			Malay					Significant interaction	
Maternal GC	rs7041			Chinese		Maternal 250HD			Inverse association of	
Maternal GC	rs4588	C-S	217	Indians	28.9 ± 4.2	(< or >30 nmol/L),	Infant birth weight	0.018	maternal Vit D deficiency	[25]
Cord VDR	rs2228570	_	(107/110)	Bajau		infant 25OHD	0		anthropometry; neonatal G	
Cord GC	rs7041			Suluk Miyod Ethnia		(< 01 >30 111101/ L)			allele carriers associated	
Cord GC	rs4588			witzed Eurific					with higher birth weight	
	rs9930501						BMI	0.125	_	
FTO	rs9930506						Body weight	0.058	_	
	rs9932754		103	Malay		Hincref (high-protein	WC	0.224	_	
	rs1042713	RCT	(16/87)	Chinese Indians	>18	calorie-restricted) diet	WHR	0.369	No significant interactions	[26]
ADRB2							Fat mass	0.234	_	
	rs1042714						BFP	0.468	_	
							Muscle mass	0.068		
AGTR1	rs5186			Malay		Vegetables, fruits, soy diet (VFSD) in Malay		0.994	No significant interactions; Malay and Chinese	
nonn	100100	C-S	507 (154/353)	Chinese Indian	30-65	Rice, egg and fish diet (REFD) in Chinese	BMI	0.66	showed high risk for lipids with gene-diet interactions	[27]
AGTR2	rs1403543					VFSD in Chinese females	-	0.053	[–] but not Indians	
VEGFR2	Rs1870377	P-C	179	Chinese	30–65	Meat, rice, and noodles diet	BMI	0.408	No significant interactions	[28]

Table 1. Summary table of gene-diet interactions on obesity in populations from Southeast Asians by country.

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	P _{interaction}	Interpretation	Ref.
					Indor	lesia				
						Total energy intake (kcal)	Changes in body weight (kg)	0.263	_	
						$[2570 \pm 1066 \text{ (baseline)}]$ to 2120 ± 1042	Changes in % body fat	0.303	-	
						(after 2 years)]	Changes in WHR	0.464	-	
							Changes in body weight (kg)	0.896	-	
						Fat intake % [23.9 \pm 11.2]	Changes in % body fat	0.965	-	
							Changes in WHR	0.996	-	
	rs659366 (Carbohydrate intake %	Changes in body weight (kg)	0.433	-	
UCP2	AA + GA Genotype	P-C	203	Indonesian	20–56	$[63.9 \pm 11.2 \text{ (baseline) to} \\ 64.2 \pm 10.4 \\ (after 2 years)]$	Changes in % body fat	0.839	No significant interactions	[29]
						(arter = j care)]	Changes in WHR	0.665	-	
						Protein intake %	Changes in body weight (kg)	0.076	-	
						$[12.3 \pm 3.4$ (baseline) to 12.4 ± 3.7 (after 2 years)]	Changes in % body fat	0.360	-	
							Changes in WHR	0.355	-	
							Changes in body weight (kg)	0.251	-	
						Physical activity (MET-min/week)	Changes in % body fat	0.979	-	
							Changes in WHR	0.684	-	

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
						Total energy intake (kcal)	Changes in body weight (kg)	0.016		
						$[2570 \pm 1066 \text{ (baseline)})$ to 2120 ± 1042 (after 2 years)]	Changes in % body fat	0.034	-	
						(arter 2 years)]	Changes in WHR	0.070	-	
							Changes in body weight (kg)	0.682	-	
						Fat intake % [23.9 \pm 11.2]	Changes in % body fat	0.974	- Significant interaction	
							Changes in WHR	0.753	between UCP2 gene	
	rs659366					Carbohydrate intake %	Changes in body weight (kg)	0.580	 variation and total energy intake on body weight change and BFP 	
UCP2	(–866G/A) GG Genotype	P-C	120	Indonesian	20–56	$[63.9 \pm 11.2 \text{ (baseline) to} 64.2 \pm 10.4 \text{ (after 2 years)]}$	Changes in % body fat	0.771	Significant interaction	
					(after 2 years)]	(arter 2 years)]	Changes in WHR	0.826	 between UCP2 gene variation and physical 	
						Protein intake %	Changes in body weight (kg)	0.830	activity on WHR	
						$[12.3 \pm 3.4$ (baseline) to 12.4 ± 3.7 (after 2 years)]	Changes in % body fat	0.913	_	
							Changes of WHR	0.103		
						Physical activity	Changes in body weight (kg)	0.666	-	
				(MET-min/week)	Changes in % body fat	0.653				
							Changes in WHR	0.040	-	

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
GRS	Vitamin D GRS [DHCR7, CYP2R1, CYP24A1, GC, CASR]	C-S	110	Minangkabau women	25-60	Carbohydrate [(235.2 g ± 73.5)]	BFP	0.049	A significant interaction between Vitamin D GRS and carbohydrate intake on \log_{BFP} ; carriers of more than 2 risk alleles and consumed high carbohydrate amounts had significantly high \log_{BFP}	[12]
	Metabolic GRS [FTO, TCF7L2, MC4R, KCNQ1, CDKN2A/B]	-				Carbohydrate [(235.2 g ± 73.5)] and protein [77.87 g ± 220.5]	BMI WC BFP	0.997	No significant interactions	
UCP2	rs659366 (–866G/A)	C-C	261 (145/116)	Indonesian	15–21	Fat intake % [(>24.14 ± 5.95]	Obesity	0.006	A significant interaction between <i>UCP2</i> rs659366 (–866G/A) and fat intake on obesity	[30]
							BMI	0.214		
	rs659366					Coffee intake ml	Body fat	0.015	_	
	(-866 AA + GA)					(34.5 ± 89.6)	WC	0.302	Significant interaction	
UCP2		C-S	455 (223/232)	Indonesian	19–56		Hip circumference	0.253	between UCP2 rs659366	[31]
	(500//						BMI	0.231	(-866 AA + GA) and coffee intake on body fat	
	rs659366 (-866 GG)					Coffee intake ml	Body fat	0.313		
	()					(37.1 ± 106.4)	WC	0.510	_	
							Hip circumference	0.421		

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
							BMI	0.933		
						Fat (59.00 g \pm 33.10)	WC	0.444	_	
							BFP	0.275	_	
							BMI	0.685	_	
						Carbohydrate $(233 \circ + 71)$	WC	0.875	- A significant interaction	
	9-SNP-B12-GRS					$(200 g \pm 71)$	BFP	0.064	between <i>B12 GRS</i> and	
							BMI	0.993	 protein energy (%) on BFP 	
						Protein energy $(76.90 \text{ g} + 36.50)$	WC	0.395	_	
						(10170 g ± 00100)	BFP	0.034	_	
							BMI	0.155	_	
						Fiber energy $(8.80 \text{ g} + 4.50)$	WC	0.547	_	
GRS		C-S	117	Minangkabau women	25-60	$(0.00 \text{ g} \pm 1.00)$	BFP	0.697	_	[32]
		_		Wonten			BMI	0.422		
						Fat (59.00 g \pm 33.10)	WC	0.812	_	
							BFP	0.775	_	
							BMI	0.230	_	
						Carbohydrate $((233 \circ + 71))$	WC	0.072	A significant interaction	
	9-SNP- Metabolic-GRS					((100 g ± / 1)	BFP	0.844	between metabolic <i>GRS</i>	
	Weldbolle Cho						BMI	0.110	on WC	
						Protein energy $(76.90 \text{ g} + 36.50)$	WC	0.032	_	
						(10170 g ± 00100)	BFP	0.568	_	
							BMI	0.273	_	
						Fiber energy $(8.80 \text{ g} + 4.50)$	WC	0.648		
						(0.00 8 ± 1.00)	BFP	0.423	_	

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
	6SNP-Vitamin						Infant birth weight	0.611		
	D-GRS (≤3)	_					Infant birth length	0.065		
	6SNP-Vitamin	_			29.6 ± 5.56		Infant birth weight	0.872	_	
	D-GRS (≥4)	_		Minangkabau		Maternal carbohydrate 29.6 ± 5.56 intake during third trimester	Infant birth length	0.073		
	4-SNP- GRS[DHCR7, GC, CYP24A1 and CYP2R1]						Infant birth weight	0.841	Significant interactions between VDR GRS and carbohydrate intake on new-born birth length	
GRS	(<3)	- P-C	183				Infant birth length	0.256	 Pregnant women with a high genetic risk of vitamin D deficiency with high carbohydrate intake gave birth to babies with lower birth lengths 	[33]
GIG	4-SNP- GRS[DHCR7, GC, CYP24A1 and CYP2P11		100	women	22.0 ± 0.00		Infant birth weight	0.795		[00]
	(≥3)	_					Infant birth length	0.079		
	2-SNP-VDR- GRS						Infant birth weight	0.810		
	(<2)	_					Infant birth length	0.032		
	2-SNP-VDR-						Infant birth weight	0.775		
	(≥2)						Infant birth length	0.099		
						Carbohydrate	BMI	0.961	_	
						(53.97 ± 9.44)	WC	0.224	_	
	15-SNP-					Protein	BMI	0.282	Significant interaction	
GRS	GRS cardiometabolic C-S disease related traits-GRS	110	Minangkabau	25-60	$(13.51 \pm 1.18\% \text{ TEI})$	WC	0.002	between <i>GRS</i> and protein	[21]	
ene		disease related C-S 110 women 2 traits-GRS	women	20 00		BMI	0.721	intake on obesity- related outcome	[21]	
				Fat (28	Fat (28.95 ± 7.99)	WC	0.577			
					BMI	0.876				
						Fiber g (8.78 \pm 4.29)	WC	0.614		

Gene	Genetic Variation	Study Design	n (Men/Women)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
CCDC171	rs4740619	C-S P-C	7817	Chinese	24-95	Cholesterol (lowest consumption group 184.35 \pm 116.00 to highest consumption group 260.61 \pm 150.56)	BMI	0.043	Significant interaction was observed; <i>CCDC171</i> rs4740619 interaction with cholesterol showed increased BMI level in subjects	[34]
FADS	rs174570	P-C	5264	Chinese	30–55	Total fish (0.16 servings per day ± 0.07) Food sourced EPA + DHA (0.33 g/d ± 0.20)	BMI	0.035	Significant interaction was observed.; long-term BMI changes in people with high fish/n-3 PUFA intake carrying signature allele show increased weight gain and risk of obesity	[35]
APOA2	rs5082 (-265T > C)	C-S	3605 (1714/1891)	Chinese Malay Indian	18–69	SFA intake (22 g)	BMI	0.758	No significant interactions	[36]
PPAR-Ƴ	rs1801282 (Pro12Ala)	C-S	4038	Chinese Malay	18–69	PUFA/SFA	BMI	0.873	No significant interactions	[37]
	rs3856806(C1431T)		(1009/2109)	Indian				0.472	_	

Notes: Hipcref (high-protein calorie-restricted) Diet: energy deficit of 300–500 kcal/day, 30% energy from protein, 30% energy from fat, 40% energy from carbohydrate, vitamin $E \ge 15 \text{ mg/day}$, and fiber $\ge 25 \text{ g/day}$. Control diet: dietary advice on weight loss based on the Malaysian Dietary Guidelines 2010 (<1500 kcal/day with a macronutrient composition of approximately 10–15% energy from protein, 20–30% energy from fat, and 55–70% energy from carbohydrate) c PRS (polygenic risk score): *FTO* rs9930501, rs9930506, rs9932754 *ADRB2* rs1042713, rs1042714. PRAL: potential renal acid load = 0.49 protein (g/day) + 0.037 phosphorus (mg/day)—0.021 potassium (mg/day)—0.026 magnesium (mg/day)—0.013 calcium (mg/day). C-S, cross-sectional; P-C, prospective-cohort study; RCT, randomized control trial. Statistically significant gene-diet interactions are highlighted in bold.

2.5. Risk of Bias and Certainty Assessment

The appraisal tool for cross-sectional studies (AXIS) was used to assess the methodological quality and risk of bias (RoB) of the cross-sectional studies (Supplementary Section S1, Tables S2 and S3) [42]. The RoB in non-randomized studies of interventions (ROBINS-1) assessment tool was used for cohort, case–control, and non-randomized studies (Supplementary Section S1, Table S4) [43]. A revised Cochrane RoB tool for randomized trials (Rob2) was used for randomized control trials (Supplementary Section S1, Table S5) [41,44]. This review falls within the framework and guidelines from the synthesis without meta-analysis (SWiM) in systematic reviews [24,26].

3. Results and Discussion

3.1. Nutrigenetics Studies in Southeast Asia

Using PubMed and Google Scholar search engines, we found 19,031 articles matching the search strings. After the full-text screening, we included a total of 20 nutrigenetic studies related to obesity- and diabetes-related parameters carried out in SE Asia. Out of this, 16 studies examined obesity-related outcomes, 13 examined diabetes-related outcomes, and 9 studies observed both obesity, and diabetes-related outcomes. Figure 1 shows the selection of the 20 studies included in this systematic review. From the included studies, only three ASEAN countries (Malaysia, Indonesia, and Singapore) conducted studies to understand gene–diet interactions on metabolic disease-related traits.





3.2. Gene–Diet Interactions on Obesity-Related Outcomes in the Southeast Asian Population

Gene–diet interactions on obesity-related traits were observed in five Malaysian, seven Indonesian, and four Singaporean studies (Figure 2, Table 1).

3.2.1. Malaysia

The interaction between gene variants and dietary factors on obesity-related outcomes were examined in five Malaysian studies: four cross-sectional ([24,25,27,28]) and one randomized control trial (RCT) [26].

One cross-sectional study (n = 200) in Chinese and Indian ethnic groups living in Malaysia reported no significant interaction between *FADS1* SNP rs174547 and linoleic acid (LA) or α -linolenic acid (ALA) on WC but showed that vegetarians with TT genotype of *FADS1* gene had higher odds of metabolic diseases and larger WC [24]. The *FADS1* gene is involved in the lipid metabolic pathway to catalyze the biosynthesis of unsaturated fatty acids and is known to play a significant role in the maintenance of triglycerol and HDL-c levels [45].

Another cross-sectional study in 217 individuals showed significant interaction (P_{interaction} = 0.018) between maternal vitamin D deficiency and cord *VDR* SNP rs2228570 on neonatal birth weight and an inverse association of maternal vitamin D deficiency with neonatal birth weight indicating the importance of this gene–diet interaction on fetal anthropometry [25]. Vitamin D is a secosteroid and a prohormone that plays a pivotal role in embryogenesis, calcium homeostasis, and fetal bone development and its deficiency is associated with adverse fetal and maternal outcomes [46]. This explains the importance of understanding maternal vitamin D status along with a genetic factor, here cord VDR SNP rs2228570, to examine how gene–nutrient interaction can influence neonatal birth anthropometric outcomes.

The third cross-sectional study (n = 507) in the Malay, Chinese, and Indian ethnic groups analyzed the interactions between *AGTR1* and *AGTR2* gene variants and different dietary patterns on body mass index (BMI). This study revealed no significant interactions between *AGTR1* SNP rs5186 and the dietary patterns (vegetables, fruits, soy diet (VFSD) in Malays and rice, egg, and fish diet (REFD) in Chinese) on BMI. The same study also failed to show interactions between *AGTR2* SNP rs1403543 and VFSD on BMI in Chinese women. Interestingly, this study found that the Malay and Chinese ethnic groups were at a higher risk for elevated lipids compared with the Indian ethnic group [27]. A better understanding of these genes in the context of obesity in different ethnicities is important as the reninangiotensin system is an important regulator of adipose tissue metabolism, whole-body energy, and glucose homeostasis [47]. Previous studies have shown that overexpression of the adipose renin-angiotensin system could be associated with obesity [48].

The fourth cross-sectional study (n = 179) examined the interaction between *VEGFR2* SNP rs1870377 and meat, rice, and noodles diet on BMI in the Chinese population and showed no significant interaction [28]. Previous *in vivo* studies to understand the role of *VEGFR1* and *VEGFR2* in angiogenesis in diet-induced obesity have shown that *VEGFR2* antiangiogenic blockade may limit adipose tissue expansion in obesity [49]. Studies in larger populations of different ethnicities are needed to better understand this mechanism in the context of ethnic-specific gene–nutrient interaction as this may be a potential target for obesity prevention and treatment strategies.

The final study is an RCT (n = 128) in the Malaysian population that observed the interaction between *FTO* (rs9930501, rs9930506, rs9932754) and *ADRB2* (rs1042713, rs1042714) gene variants and Hipcref (high-protein, calorie-restricted, high-vitamin E, and high-fiber) diet pattern on BMI, body weight, WC, WHR, fat mass, body fat percentage (BFP) and muscle mass in the Chinese, Malay, and Indian ethnicities. However, this study failed to show any significant interaction between these genes and dietary patterns on the abovementioned obesity-related parameters [26]. The *FTO* and *ADRB2* genes are widely studied specifically in relation to obesity [26,50]. An interesting finding from previous literature showed that *FTO* carriers of heterozygous risk alleles could still have a protective effect against obesity when subjected to increased physical activity and by following an appropriate weight loss regimen. Homozygous carriers of the *ADRB2* allele (G > A genotype) have been linked to lower levels of lipid mobilization, which could provide insights into creating dietary plans and obesity prevention strategies for various ethnicities [26]. Further understanding of this concept in larger groups in the SE Asian population may yield promising outcomes in the field of nutrigenetics to target obesity.

3.2.2. Indonesia

Seven studies analyzed gene–diet interactions on obesity-related parameters in the Indonesian population: four cross-sectional [12,21,31,32], two prospective cohort studies [29,33] and one case–control study [30].



Figure 2. Gene–diet interactions on obesity-related traits. The figure shows the significant gene–diet interactions (p < 0.05) on obesity traits and the dietary factors that have influenced the risk of obesity in individuals carrying specific genetic variations in the Malaysian [25], Indonesian [12,21,29–31,33] and Singaporean [34,35] populations. GRS, genetic risk score; BMI, body mass index; EPA + DHA, eicosapentaenoic acid + docosahexaenoic; TEI, total energy intake; plus-minus symbol (\pm) indicates Standard deviation.

Genes	Genetic Variations	Study Design	n (Male/Female)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
						Malaysia				
	rs9930501						Fasting glucose	0.381	Significant interaction	
FTO	rs9930506	-	103	Malay	alay	Hipcref ^a	Easting insulin	0.101	observed;participants showed a greater	
	rs9932754	RCT	(16/87)	Chinese	>18	(high-protein calorie-restricted)	Fasting insum	0.121	reduction in hsCRP	[26]
ADRB2	rs1042713	_		mentans		control diet ^b	HOMA-IR	0.122	diet compared to	
MDRD2	rs1042714	_				_	hs-CRP	0.048	normal diet	
AGTR1	rs5186			Malay		Vegetables, fruits, and soy diet (VFSD) in Malays		0.537		
		C-S	507 (154/353)	Chinese Indian	30–65	Rice, egg, and fish diet (REFD) in Chinese	HbA1C	0.844	No significant interactions	[27]
AGTR2	rs1403543	-				VFSD in Chinese females		0.989	_	
VEGFR2	rs1870377	C-S	179	Chinese	30–65	Meat, rice, and	Blood glucose	p > 0.05	No significant	[28]
						noodles diet	HbA1c	p > 0.05	interactions	
FADS1	rs174547	C-S	200	Chinese	>18	Linoleic acid (7.9 \pm 3.6 g/day)	Log _{FBG}	0.807	No significant	[24]
			(69/131)	Indians		α-Linolenic acid (0.4–0.8 g/day)	0	0.293	interactions	
IGF1	rs35767	_ C-S	211	Chinese	66.7 ± 6	DAL (using PRAL)	FBG	NS	No significant interactions; study shows association between DAL and high	[38]
IGF1	rs7136446	_							FBG, indicating a	
IL6	rs1800796								for diabetes	
ADRB2	rs1042713	C-S	126	Malaysian Chinese	18–74	Saturated fat intake (<7.3% of total energy/day) PUFA intake	FBG	0.011	Significant gene diet interactions; G allele carriers of <i>ADRB2</i> rs1042713 are associated	[39]
				Indians		(≥0.8/day)	HOMA-IR	0.026	with increased odds of	
						$(\geq 6\% \text{ of TE/ day})$	Fasting insulin	0.036	insulin resistance	

Table 2. Summary table of gene-diet interactions on diabetes in Southeast Asians by country.

Genes	Genetic Variations	Study Design	n (Male/Female)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
						Indonesia				
							log _{HbA1c}	0.175		
						Fat (59.00 \pm 33.10)	log _{FBG}	0.374	_	
							log fasting serum insulin	0.757		
							log _{HbA1c}	0.091		
						Carbohydrate (233 ± 71)	log _{FBG}	0.260	_	
	9-SNP-B12-GRS						log fasting serum insulin	0.341	_	
						D	log _{HbA1c}	0.150		
						(76.90 ± 36.50)	log _{FBG}	0.368	_	
						· · · · · ·	log _{fasting serum insulin}	0.073	—	
						T"1 · · 1	log _{HbA1c}	0.042	_ between B12 GRS and	
				Minanakahau		Fiber intake (8.80 ± 4.50)	log _{FBG}	0.380	fiber intake on HbA1c	
GRS		C-S	117	women	25-60	· · · ·	log fasting serum insulin	0.215	≥ 9 risk alleles who	[32]
						T (log _{HbA1c}	0.298	consumed low fiber diet	
						Fat (59.00 ± 33.10)	log _{FBG}	0.634	HbA1c levels	
						· · · · ·	log fasting serum insulin	0.108		
							log _{HbA1c}	0.166	_	
						Carbohydrate (233 ± 71)	log _{FBG}	0.771	_	
	9-SNP- Metabolic-GRS					. ,	log fasting serum insulin	0.104	_	
							log _{HbA1c}	0.155	_	
						(76.90 ± 36.50)	log _{FBG}	0.929	_	
							log fasting serum insulin	0.890	_	
						Fiber intake (a/d) —	log _{HbA1c}	0.851	_	
						$(4.90 \pm 1.00 \text{ g/day})$	log _{FBG} log _{fasting} serum insulin	0.215 0.947		

Genes	Genetic Variations	Study Design	n (Male/Female)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.											
							Glucose	0.360													
						Carbohydrate (g/d) $=$ (233.7 \pm 75.1 g) $=$	HbA1c	0.780													
						() =	Fasting insulin	0.630													
							Glucose	0.560													
	Metabolic GRS					Protein (g/d) (77.2 ± 41.7 g)	HbA1c	0.680													
GRS	[FTO, TCF7L2, MC4R, KCNO1,	C-S	110	Minangkabau women	25–60		Fasting insulin	0.220	No significant interactions	[12]											
	CDKN2A/B]						T (1)	Glucose	0.700	-											
							Fat (g/d) (61.2 ± 36.1 g)	HbA1c	0.780												
						(01.2 ± 00.1 g) _	Fasting insulin	0.440													
							Glucose	0.830													
						Fiber (g/d) (8.6 ± 4.3 g)	HbA1c	0.530	-												
						(0.0 - 0.0 8) -	Fasting insulin	0.440													
									log _{Glucose}	0.882											
						Carbohydrates % (53.97 + 9.44)	log _{Insulin}	0.336	-												
						(***** = ****) =	log _{HbA1c}	0.766													
							log _{Glucose}	0.751													
	15-SNP-					Protein % $-$ (16.93 \pm 3.32)	log _{Insulin}	0.341													
GRS	cardiometabolic disease related	C-S	110	Minangkabau women	25-60	() =	log _{HbA1c}	0.638	No significant interactions	[21]											
	traits-GRS			women			log _{Glucose}	0.732	interactions												
						Fat % - (28.95 + 7.99)	log _{Insulin}	0.480													
					(1000 ± 100)	log _{HbA1c}	0.935														
							-											log _{Glucose}	0.833		
								Fiber g $-$ (8 78 + 4 29)	log _{Insulin}	0.216											
						(00 ± 1)) =	log _{HbA1c}	0.162													

Genes	Genetic Variations	Study Design	n (Male/Female)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
						Singapore				
GRS	DM 37-SNP	P-C	38,434	Chinese	30–79	Alcohol (men: 10–25 g/d; women: 5–15 g/d)	Diabetes risk	NS	No significant interaction; but a healthy lifestyle and any genetic	[40]
	GRS		,			VFSD pattern			risk category was associated with a	
						Meat—dim sum pattern			significantly lower risk of diabetes	
							Fasting glucose	0.425		
						Total fat % [–] (25.2–29.8 TEI)	Fasting insulin	0.01		
						(20.2 2).0 (20)	HOMA-IR	0.007		
	rs894160						Fasting glucose	0.004	_	
	(11482G > A)					SFA % - (9.4–11.8)	Fasting insulin	0.004	_	
						(*** -***) =	HOMA-IR	0.003		
							Fasting glucose	0.145	Significant interaction	
						Carbohydrates % (56.3–61.7)	Fasting insulin	0.007		
PLIN		C-S	4107	Chinese Malay	18-69	(**** ****)	HOMA-IR	0.004	between <i>PLIN</i> variants and dietary factors on	[41]
				Indian	10 07		Fasting glucose	0.448	diabetes	
						Total fat % (25.2–29.8 TEI)	Fasting insulin	0.014	related outcomes	
							HOMA-IR	0.012		
	rs1052700						Fasting glucose	0.009		
	(14995A > T)					SFA % (9.4–11.8)	Fasting insulin	0.014		
					(///)	HOMA-IR	0.005			
						Fasting glucose	0.293			
						Carbohydrates % — (56.3–61.7)	Fasting insulin	0.008		
						(56.3–61.7)	HOMA-IR	0.012		

Tabl	le 2.	Cont.

Genes	Genetic Variations	Study Design	n (Male/Female)	Ethnicity	Age (Years)	Dietary Factors	Outcomes	Pinteraction	Interpretation	Ref.
PPAR-7	rs1801282(Pro12Ala)) C-S	4038 1869/2169)	Chinese	ne 7 18–69 1	PUFA/SFA	Insulin	0.089	No significant	[37]
	rs3856806(C1431T)			Malay Indian				0.175	interactions	
APOA2	rs5082 (-265T > C)	C-S	3605 (1714/1891)	Chinese Malay Indian	18–69	SFA intake (22 g)	HOMA-IR	0.026	Significant interaction between APOA2 rs5082(–265T > C) and SFA intake on HOMA-IR	[36]

Notes. ^a Hipcref (high-protein calorie-restricted) Diet: energy deficit of 300–500 kcal/day, 30% energy from protein, 30% energy from fat, 40% energy from carbohydrate, vitamin $E \ge 15 \text{ mg/day}$, and fiber $\ge 25 \text{ g/day}$. ^b control diet: dietary advice on weight loss based on the Malaysian Dietary Guidelines 2010 (<1500 kcal/ day with a macronutrient composition of approximately 10–15% energy from protein, 20–30% energy from fat, and 55–70% energy from carbohydrate) ^c PRS: *FTO* rs9930501, rs9930506, rs9932754 *ADRB2* rs1042713, rs1042714. PRAL: potential renal acid load = 0.49 protein (g/day) + 0.037 phosphorus (mg/day)—0.021 potassium (mg/day)—0.026 magnesium (mg/day)—0.013 calcium (mg/day). C-S, cross-sectional; P-C, prospective-cohort study; RCT, randomized control trial. Statistically significant gene-diet interactions are highlighted in bold.

A cross-sectional study in 110 Minangkabau women of Indonesia showed significant interaction (Pinteraction = 0.049) between vitamin D GRS (DHCR7, CYP2R1, CYP24A1, GC, CASR) and carbohydrate intake on BFP. The results indicated that participants carrying more than two risk alleles and who consumed high carbohydrate intake had significantly higher BFP than participants with less than two risk alleles. However, there was no significant interaction between metabolic-GRS (FTO, TCF7L2, MC4R, KCNQ1, CDKN2A/B) and carbohydrate or protein intake on obesity-related parameters [12]. Another cross-sectional study in 117 Minangkabau women showed significant interaction ($P_{interaction} = 0.034$) between vitamin D-associated 9-SNP-B12-GRS and protein intake on BFP. The same study also indicated a significant interaction (Pinteraction = 0.032) between vitamin D-associated 9-SNP-metabolic-GRS and protein energy on WC, indicating that women consuming a low fiber diet (4.90 \pm 1.00 g/day) and harboring \geq 9 risk alleles for vitamin B12 deficiency had notably higher HbA1C levels than the others ($P_{interaction} = 0.025$) [32]. There are many mechanisms proposed to understand the role of Vitamin D levels and obesity including increased fat stores and increased vitamin D storage in adipose tissue. This also considers the lifestyle differences between obese and lean individuals along with combinatorial effects of dietary patterns to have a significant effect on obesity [51]. Further studies are needed to confirm the exact mechanism behind gene-diet interaction on these obesity-related parameters.

One study on 455 Indonesian adults from Yogyakarta examined that coffee consumption and carriers of *UCP2* SNP rs659366 AA + GA genotype had a negative correlation with BMI (P_{interaction} = 0.01) and body fat (kg) (P_{interaction} = 0.021) levels. The same study also showed that carriers of the *GG* genotype had no correlation with coffee consumption and obesity, indicating that gene variations and coffee intake influences obesity-related parameters [31]. Previous studies have elucidated the potential anti-obesity properties of tea and coffee [52]. Scientific evidence shows possible mechanisms of this activity via cell cycle regulation in adipocytes during adipogenesis, the effect on transcription factors involved in weight loss, and lipogenesis-related proteins [53]. However, further ethnic-specific research is needed to better understand these mechanisms as this proves to be a promising strategy to combat obesity due to the large global consumption of coffee.

Another study in 110 Minangkabau women observed that carriers of more than six risk alleles of a 15-SNP-GRS for cardiometabolic disease and consuming low protein intake had significantly (P_{interaction} = 0.002) lower WC compared to carriers of less than six risk alleles. In addition, the study also showed a significant influence of GRS on WC and triglyceride levels through a low-protein diet specifically in Minangkabau women [21]. One prospective cohort study in the Indonesian adult population examined the interaction between two UCP2 gene variations ((rs659366 (-866G/A) AA + GA Genotype) and rs659366 (-866G/A) GG Genotype) and dietary factors on obesity-related outcomes (body weight, BFP, waisthip ratio (WHR)). There were no significant interactions observed between UCP2 gene variations and any of the dietary factors on the obesity-related parameters; however, significant interaction depicting a positive correlation between UCP2 SNP rs659366 (-866G/A) GG genotype and total energy intake on body weight change (Pinteraction = 0.016) and BFP $(P_{interaction} = 0.034)$ was observed. The same study also showed significant interaction $(P_{interaction} = 0.040)$ between UCP2 SNP rs659366 (-866G/A) GG genotype and physical activity on WHR indicating that participants with increased physical activity and the UCP2 gene variant had lower WHR [29]. Further studies on larger populations are required to validate these results and elucidate the mechanisms.

Another study showed a significant interaction ($P_{interaction} = 0.032$) between 2-*SNP*-*GRS* and carbohydrate intake on infant birth length. Pregnant women with >2 risk alleles of *VDR-GRS* and low vitamin D status and who consumed a high carbohydrate diet (405.88 ± 57.16 g/day) during the third trimester gave birth to babies with a lower birth length. This could suggest the benefits of low carbohydrate intake in Indonesian women with >2 risk alleles of *VDR*-GRS and low vitamin D status, but would need validation from further studies [33]. A case–control study of 261 Indonesian adolescents showed significant interaction ($P_{interaction} = 0.006$) between fat intake and the *UCP2* SNP rs659366 on obesity risk. This study also indicated that carriers of *UCP2* SNP rs659366 who consumed a high-fat diet had a lower chance of becoming obese compared to non-carriers with normal fat intake [30]. *UCP2* gene variants are very commonly studied in association with obesity. The best--understood mechanisms of *UCP2*-mediated regulation of obesity include: (a) direct activation of melanocortin-4 receptor that increases energy expenditure and decreases food intake and (b) negative regulation of glucose-dependent insulin secretion in the beta cells of the pancreas and positive regulation of glucagon from the alpha cells [54]. It is also understood that *UCP2* expression has a positive correlation with weight loss [55].

3.2.3. Singapore

Four Singaporean studies examined gene–diet interactions on obesity-related outcomes [34–37]. A study on 7817 Singaporeans examining the interaction between *CCDC* SNP rs4740619 variant and cholesterol intake on BMI in the Chinese ethnicity showed a statistically significant interaction ($P_{interaction} = 0.043$) between SNP rs4740619 and cholesterol intake on change in BMI. This study also indicated that this new locus identified does not commonly interact with dietary factors but proves an association in the SE Asian population [34]. An initially proposed mechanism according to the HaploReg analysis (tool to investigate the non-coding genome annotations from published GWAS (genome-wide association study) or novel variants) describes a possible alteration in the binding affinity of peroxisome proliferator-activated receptors regulating multiple metabolic pathways in obesity, but further studies are required to precisely determine the actual mechanism of *CCDC* SNP rs4740619 variant in obesity [56].

A 10-year prospective cohort study on 5264 individuals of Chinese ethnicity examined the interactions between *FADS* SNP rs174570 and total fish, food-sourced eicosapentaenoic acid (EPA) + docosahexaenoic acid (DHA) on BMI. The study showed significant interaction ($P_{interaction} = 0.035$) and a long-term increase in BMI in individuals carrying the signature 'T' allele with high fish/n-3 polyunsaturated fatty acids (PUFA) intake [35]. It is also well understood that *FADS1* and *FADS2* are involved in the rate-limiting steps of the fatty acid metabolic pathway and are consistently associated with plasma and tissue levels of arachidonic acid and EPA [57].

APOA2 is associated with high-density lipoproteins, reverse cholesterol transport impairment, antioxidant properties, and fat distribution phenotypes that are associated with metabolic disease progression [58]. Here, a multi-ethnic cross-sectional study on 3605 Singaporeans of the Chinese, Malay, and Indian ethnicities examining saturated fatty acid (SFA) intake and *APOA2* SNP rs5082 on BMI showed no significant interaction [36]. A similar study in 4038 individuals also showed no significant interaction between PUFA/saturated fatty acids (SFA) and *PPAR-* γ SNPs rs1801282 and rs3856806 on BMI [37]. Further analysis of different *APOA2* gene variants may provide insights into gene–diet interactions specific to the SE Asian population.

3.3. Gene–Diet Interactions on Diabetes-Related Outcomes in the Southeast Asian Population

Gene–diet interactions on diabetes-related traits were observed in six Malaysian, three Indonesian, and four Singaporean studies (Figure 3, Table 2).



Figure 3. Gene–diet interactions on diabetes-related traits. The figure shows the significant gene–diet interactions (p < 0.05) on diabetes traits and the dietary factors that influenced the risk of diabetes in individuals carrying specific genetic variations in the Malysian [39], Indonesian [32] and Singaporean [36,41] populations. GRS, genetic risk score; BMI, body mass index; SFA, saturated fatty acids; PUFA, polyunsaturated fatty acids; HOMA-IR, homeostatic model assessment—insulin resistance; TE, total energy.

3.3.1. Malaysia

The gene–diet interactions on diabetes-related outcomes in the Malaysian population were investigated by six studies: one RCT [26] and five cross-sectional studies [24,27,28,38,39].

One RCT in 128 Malaysian participants examined the interaction between *FTO* (rs9930501, rs9930506, rs9932754) and *ADRB2* (rs1042713, rs1042714) gene variants and Hipcref diet on diabetes-related outcomes in the Malay, Chinese, and Indian ethnic groups. This study showed a significant interaction (P_{interaction} = 0.048) indicating that participants had a reduction in hs-CRP level in the Hipcref-*PRS* interventional diet compared to normal diets [26]. This is a notable association because the *FTO* gene variants are not only linked with obesity but also have a strong association with diabetes [59]. When the *FTO* gene is overexpressed in INS-1 pancreatic beta cells, it upregulates transcription factor 7-like 2 (*TCF7L2*) which is a key determinant of diabetes [60]. *ADRB2* is also another gene understood to play a pivotal role in glucose homeostasis. In vivo studies have shown that pancreas-specific deletion of the *ADRB2* gene in the pancreas impacts not only glucose secretion and tolerance, but also increases *VEGF-A* production. This has a direct effect on impaired insulin production, exocytosis, and accelerates the development of diabetes related complications like retinopathy and macular edema [61].

A cross-sectional study of 507 participants showed no significant interaction between *AGTR 1* SNP rs5186, *AGTR2* SNP rs1403543, and VFSD in Malay, REFD in Chinese, and VFSD in Chinese females on HbA1c levels [27]. A previous study reported several *AGTR1* gene variants expressed in several tissues such as blood vessels, kidneys, and lungs and once expressed, lead to water–sodium retention, elevated blood pressure, and microvascular disorders in diabetes [62]. Another study (n = 179) in the Chinese ethnic group examined

VEGFR2 SNP rs1870377 and meat, rice, and noodles diet on blood glucose and HbA1c also showed no significant interaction [28]. A better understanding of this relationship in the context of gene–diet interaction is important because inhibition/downregulation of the *VEGFR2* signaling axis is associated with endothelial dysfunction in diabetes [63].

One study conducted on 200 participants of the Chinese and Indian ethnic groups to understand the interaction of *FADS1* SNP rs174547 and LA, ALA intake on fasting blood glucose levels showed no statistically significant interactions [24]. A cross-sectional study (n = 211) performed on the Chinese ethnic group to understand the interaction of *IGF1* rs35767, *IGF1* rs7136446, *IL6* rs1800796, and DAL intake (using PRAL levels) on fasting blood glucose also showed no significant interaction [38]. *IGF1*, with structural homology to insulin, is responsible for increased peripheral glucose intake and reduction in hepatic glucose production for better insulin sensitivity. When *IGF1* levels are lower, it is often associated with higher anthropometric variables correlating with insulin resistance [64]. *IL6* is a pro-inflammatory cytokine with a known mechanism to develop insulin resistance and is involved in the pathogenesis of diabetes. This is often the result of its irregular expression (usually genetic) and long-term exposure leading to inflammation that induces insulin resistance and increases the overall risk of diabetes [65].

The fifth cross-sectional study (n = 126) on Malaysian, Chinese, and Indian ethnic groups to understand the interaction between *ADRB2* SNP rs1042713 and saturated fat, PUFA intake on diabetes-related outcomes showed significant gene–diet interactions on fasting blood glucose (P_{interaction} = 0.011), HOMA-IR (P_{interaction} = 0.026) and fasting insulin (P_{interaction} = 0.036). This study also showed that G allele carriers of *ADRB2* SNP rs1042713 were associated with increased odds of developing insulin resistance [39]. Understanding this gene–diet interaction in diabetes is important because *ADRB2* has shown close associations with diabetes by directly influencing anthropometric measures, fasting insulin level, and insulin resistance [61].

3.3.2. Indonesia

Three cross-sectional studies were performed on Minangkabau women of Indonesia to understand the significance of gene–diet interaction in diabetes-related outcomes.

One cross-sectional study (n = 117) showed significant interaction ($P_{interaction} = 0.042$) between vitamin B12 GRS and fiber intake on HbA1C levels. Interestingly, the study also showed that individuals with \geq 9 risk alleles who consumed a low-fiber diet had higher HbA1c levels indicating a significant interaction between *B12* GRS and dietary factor [32]. Vitamin B12 deficiency in diabetic patients with metformin is quite common compared to the relationship between vitamin B12 deficiency in individuals who are not administered metformin medication. It was interesting to note that a previous study in the Chinese population (n = 16,699) demonstrated that individuals who did not take metformin as a part of their treatment regime had significantly higher B12 deficiency [66]. This poses a need for further studies on larger groups of SE Asians to understand the interaction between B12 GRS and dietary factors on diabetes-related outcomes.

The second study (n = 110) showed no significant interaction between *Metabolic GRS* [*FTO*, *TCF7L2*, *MC4R*, *KCNQ1*, *CDKN2A*/*B*] and dietary factors on diabetes-related outcomes (glucose, HbA1C, fasting insulin) [12]. The third cross-sectional study (n = 110) on Minangkabau women who were carriers of more than six risk alleles of a 15-SNP-cardiometabolic disease GRS also showed no significant interaction with protein intake on glucose levels, HbA1c levels, and fasting insulin [21]. These studies in the Indonesian population show the importance of understanding gene–diet interaction on diabetes-related outcomes, but further analysis on larger populations is required to validate these results.

3.3.3. Singapore

A total of four studies were identified: three cross-sectional studies [36,37,41], and one prospective cohort study [40]. A total of fifteen gene–diet interactions on diabetes- related traits have been identified in these studies, mostly by Corella et. al [41]. This extensive

cross-sectional study on 4017 participants of the Chinese, Malay, and Indian ethnic groups showed several significant interactions between gene variants (*PLIN* rs894160, rs1052700) and dietary factors (total fat, carbohydrates, SFA) [41].

Another study (n = 3605) identified a significant interaction ($P_{interaction} = 0.026$) between apolipoprotein A2 (*APOA2*) SNP rs5082 and high SFA intake on homeostasis model assessment-estimated insulin resistance (HOMA-IR) in individuals who are carriers of the *CC* genotype of the variant in the Chinese, Asian, and Indian ethnicities [36]. These results are in line with another French–Caucasian case–control cohort (n = 12,387) study where the authors found an association between *APOA2* and diabetes, specifically the SNP rs5082 variant [67]. In vivo studies have also understood the mechanism of *APOA2* in diabetes where overexpression of the *APOA2* gene significantly resulted in elevated fasting blood glucose and a two-fold increase in plasma insulin levels that are key features of insulin resistance [68].

Peroxisome proliferator-activated receptor gamma (*PPAR-* γ) has been a prime subject of diabetes research because its ligands have been shown to be potential insulin sensitizers for the treatment of diabetes [69]. Contrastingly, Tai et. al (n = 4038), revealed that there were no interactions between the PUFA/SFA intake ratio and the *PPAR-* γ SNPs rs1801282 and rs3856806 on fasting insulin [37]. In addition, a prospective cohort study (n = 38,434) with a mean follow-up of 10.72 years examined DM-37-SNP GRS and dietary patterns (alcohol, vegetable-fruit-soy pattern, meat dim-sum pattern) on diabetes and found no significant interactions [40].

The above studies provide a complete picture of the nutrigenetic status of obesity and diabetes in the SE Asian population. Although the studies are in their infancy and are required to be understood by larger populations and all countries of the ASEAN, this research provides comparable results with similar gene–diet interactions with other parts of the world. A meta-analysis in the French population (n = 3069) confirmed an interaction (P_{interaction} = 0.0005) between low LA intake and *FADS1* rs174547 on low WC and BMI. The same study also indicated that minor allele carriers of *FADS1* SNP rs174547 benefitted from a lower dietary intake of LA [70]. A previous systematic analysis of a GLACIER study in the Swedish population (n = 5160) indicated that high PUFA intake modified the association between *FADS1,2,3* gene cluster variants (rs74771917, rs3168072, rs12577276, rs7115739, rs174602, and rs174570) and triglycerides [71]. Though not all *PLIN* variants have been associated with diabetes, some studies in American (n = 431) [72] and Chinese (n = 993) [73] women have shown a significant association between certain *PLIN* variants and the risk of diabetes. This emphasizes the need to validate the results of the above-mentioned gene–diet interaction on larger groups of the SE Asian population.

4. Precision Nutrition Approach for the Southeast Asian Population

The understanding of genetic diversity between individuals and among different ethnic groups should be established before designing dietary and nutritional requirements because different individuals respond differently to lifestyle interventions. Human genome sequencing plays a pivotal role in understanding genetic variations among different ethnic groups and has paved the way for the concept of personalized nutrition to frame effective lifestyle intervention strategies [74]. Developments in omics technology provide a better understanding of the whole genome of individuals as well as different ethnic groups along with the transcriptome, proteome, metabolome, and metagenome [75]. Integration of Artificial Intelligence along with gene nutrient analysis, especially in populations such as SE Asia will be useful to develop public health strategies and personalized nutrition plans for cardiometabolic diseases such as obesity and diabetes.

A high-throughput genetic screening has been developed to understand the role of SNPs in cardiometabolic diseases. However, molecular and pathophysiological mechanisms to understand gene–nutrient interactions and its influence on cardiometabolic diseases remain unexplored. In LMIC such as in SE Asia, nutrigenetics is still in its infancy and requires an evidence-based approach before framing precision nutrition strategies

for the population. Further, such studies on larger populations and ethnic groups as well as on different levels of nutrition transition, are crucial for the development of accurate and population-specific precision nutrition strategies effective to combat chronic, yet preventable diseases such as obesity and diabetes [2]. While the Western countries have shifted their approach towards nutrigenetics, developing countries like SE Asia still favor traditional methods for evaluating, categorizing, and managing obesity and diabetes. Costly gene testing, lack of knowledge, and experts in this field are the primary impediments of nutrigenetics implementation, particularly in LMIC. Even though this field is expanding globally, there are not many researchers in this discipline in SE Asia. Moving forward, the nutrigenetics approach should be considered for government health programs, particularly those aimed at noncommunicable diseases (NCDs). Currently, lifestyle diseases are a major burden to all countries, and long-term investments in accelerating nutrigenetics research and generating scientific evidence may provide a solution to obesity and its comorbidities through precision nutrition.

5. Limitations

Our study sought to analyze the gene-diet interactions on metabolic disease-related parameters in the SE Asian population. This review included twenty studies conducted in Malaysia, Indonesia, and Singapore (Figure 4) with four Indonesian studies focusing on Minangkabau women—a minority ethnic group. To the best of our knowledge, there are no articles examining gene-diet interactions on metabolic-disease-related outcomes in eight out of 11 ASEAN countries. Hence, the results of this review cannot be applied to the entire SE Asian population due to a lack of consistency and replication in dietary exposures and individual SNPs. Given these limitations, there was no possibility of a metaanalysis. Most studies performed had a cross-sectional study design and the sample size was insufficient to apply the results to the entirety of the SE Asian population. Some studies explored the relationship between SNPs and dietary factors, but there could have been an influence of other SNPs on the same outcome that remains unexplored. On performing ROBINS-1 risk assessment, one prospective cohort study was found to be at a higher risk of bias where participants were not screened for gestational diabetes which could have been a confounder in the study [33]. Future research needs to consider the general limitations highlighted in the present study and emphasize ethnic differences when looking at multi-ethnic populations.



Figure 4. Common genes involved in gene–diet interactions associated with obesity and diabetes in the Southeast Asian population (Malaysia, Indonesia, and Singapore). *AGTR1*, angiotensin II receptor type 1; *AGTR2*, angiotensin II receptor type 2; ADRB2, adrenoceptor beta 2; *FADS1*, fatty acid desaturase 1; *FTO*, fat mass and obesity-associated gene; *VEGFR2*, vascular endothelial growth factor receptor 2; *APOA2*, apolipoprotein A2; *PPARg*, peroxisome proliferator-activated receptor-gamma.

6. Conclusions

This is the first systematic analysis of the effects of gene-diet interactions on obesity and diabetes in the Southeast Asian population. This review highlights several population-, sex-, and ethnicity-specific gene-diet interactions that are significant in Malaysian, Indonesian, and Singaporean populations and provide a complete picture of nutrigenetic research conducted in SE Asia. The commonly reported interactions were between macronutrients and GRS such as B12-GRS, vitamin D GRS, and a metabolic-GRS, and there were multiple interactions between UCP2 SNP rs659366 and dietary factors on obesity traits in the Indonesian population, making the UCP2 gene a candidate for further studies to understand the mechanisms of interaction. A deeper understanding of the UCP2 gene–diet interaction and studies on larger groups of the SE Asian population may provide insights into personalized nutrition strategy development. Additionally, some Malaysian studies examined gene-diet interactions with specific dietary patterns in population subgroups including FTO, ADRB2, and Hipcref diet, AGTR1, AGTR2 genes, and VFSD, REFD diets, and VEGFR2 with a meat, rice, noodles diet to better understand its influence on obesity and diabetes [26–28]. Similar such studies are crucial in larger populations and ethnic groups for the development of accurate, population-specific precision nutrition strategies to effectively combat chronic, yet preventable diseases such as obesity and diabetes.

Supplementary Materials: The following supporting information can be downloaded at: https://www.mdpi.com/article/10.3390/nu15132948/s1, Table S1. Number of Hits and Search Strings per Database. Table S2. Summary Outcome of Assessment with the Appraisal Tool for Cross-Sectional Studies (AXIS). Table S3. Assessment with the Comments Appraisal Tool for Cross-Sectional Studies. Table S4. Assessment using the Risk of Bias in Non-Randomized Studies—of Interventions (ROBINS-I) [29,33–35,40]. Table S5. Assessment using RoB 2: A revised Cochrane risk-of-bias tool for randomized trials. Section S1—Risk of bias assessment.

Author Contributions: Conceptualization, K.S.V.; methodology, P.S., E.F.V., A.C.T.A.D. and E.S.G.C.; validation, K.S.V., E.F.V., A.L. and P.S.; writing—original draft preparation, P.S., E.F.V. and K.S.V.; writing—review and editing, P.S., E.F.V., K.S.V., J.A.L., A.C.T.A.D., E.S.G.C., A.L., P.L.Q., M.A., N.A.T., H.H., M.F.M.N. and I.G.; visualization, P.S., E.F.V. and K.S.V.; supervision, K.S.V.; project administration, K.S.V.; funding acquisition, K.S.V. and A.C.T.A.D. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by the British Council-Malaysian Industry Government Group for High Technology (MIGHT) BC-REL Going Global Partnership Grant (Grant ID: 871106314).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in this study are included in the Supplementary material. Further inquiries can be directed to the corresponding author.

Acknowledgments: We thank the funding body for their consistent support for the provisions to carry on with this review article.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript; or in the decision to publish the results.

References

- Di Cesare, M.; Sorić, M.; Bovet, P.; Miranda, J.J.; Bhutta, Z.; Stevens, G.A.; Laxmaiah, A.; Kengne, A.P.; Bentham, J. The epidemiological burden of obesity in childhood: A worldwide epidemic requiring urgent action. *BMC Med.* 2019, 17, 212. [CrossRef] [PubMed]
- Vimaleswaran, K.S. GeNuIne (gene-nutrient interactions) Collaboration: Towards implementing multi-ethnic population-based nutrigenetic studies of vitamin B(12) and D deficiencies and metabolic diseases. *Proc. Nutr. Soc.* 2021, 80, 1–11. [CrossRef] [PubMed]
- World Health Organisation. Obesity. Available online: https://www.who.int/news-room/fact-sheets/detail/obesity-andoverweight (accessed on 3 November 2022).

- World Health Organisation. Diabetes. Available online: https://www.who.int/news-room/fact-sheets/detail/diabetes (accessed on 3 November 2022).
- World Health Organisation. Obesity and Overweight in South East Asia. Available online: https://www.who.int/southeastasia/ health-topics/obesity (accessed on 3 November 2022).
- Biswas, T.; Townsend, N.; Magalhaes, R.J.S.; Islam, M.S.; Hasan, M.M.; Mamun, A. Current Progress and Future Directions in the Double Burden of Malnutrition among Women in South and Southeast Asian Countries. *Curr. Dev. Nutr.* 2019, *3*, nzz026. [CrossRef] [PubMed]
- Association of Southeast Asian Nations: ASEAN Food and Nutrition Security Report 2021, Volume 1. Available online: https://asean.org/book/asean-food-and-nutrition-security-report-2021-volume-1-the-asean-secretariat-jakarta/ (accessed on 4 November 2022).
- ASEAN Food and Nutrition Security Report 2021 Volume 2—Food and Nutrition Security Country Profiles. Available online: https: //asean.org/book/asean-food-and-nutrition-security-report-2021-volume-2-food-and-nutrition-security-country-profiles/ (accessed on 8 November 2022).
- Phillips, C.M. Nutrigenetics and metabolic disease: Current status and implications for personalised nutrition. *Nutrients* 2013, 5, 32–57. [CrossRef] [PubMed]
- Vimaleswaran, K.S.; Franks, P.W.; Barroso, I.; Brage, S.; Ekelund, U.; Wareham, N.J.; Loos, R.J. Habitual energy expenditure modifies the association between NOS3 gene polymorphisms and blood pressure. *Am. J. Hypertens.* 2008, *21*, 297–302. [CrossRef]
- Vimaleswaran, K.S.; Li, S.; Zhao, J.H.; Luan, J.; Bingham, S.A.; Khaw, K.T.; Ekelund, U.; Wareham, N.J.; Loos, R.J. Physical activity attenuates the body mass index-increasing influence of genetic variation in the FTO gene. *Am. J. Clin. Nutr.* 2009, 90, 425–428. [CrossRef]
- Alathari, B.E.; Aji, A.S.; Ariyasra, U.; Sari, S.R.; Tasrif, N.; Yani, F.F.; Sudji, I.R.; Lovegrove, J.A.; Lipoeto, N.I.; Vimaleswaran, K.S. Interaction between Vitamin D-Related Genetic Risk Score and Carbohydrate Intake on Body Fat Composition: A Study in Southeast Asian Minangkabau Women. *Nutrients* 2021, 13, 326. [CrossRef]
- 13. Bodhini, D.; Gaal, S.; Shatwan, I.; Ramya, K.; Ellahi, B.; Surendran, S.; Sudha, V.; Anjana, M.R.; Mohan, V.; Lovegrove, J.A.; et al. Interaction between TCF7L2 polymorphism and dietary fat intake on high density lipoprotein cholesterol. *PLoS ONE* **2017**, *12*, e0188382. [CrossRef]
- Vimaleswaran, K.S.; Bodhini, D.; Lakshmipriya, N.; Ramya, K.; Anjana, R.M.; Sudha, V.; Lovegrove, J.A.; Kinra, S.; Mohan, V.; Radha, V. Interaction between FTO gene variants and lifestyle factors on metabolic traits in an Asian Indian population. *Nutr. Metab.* 2016, *13*, 39. [CrossRef]
- 15. Vimaleswaran, K.S.; Cavadino, A.; Hyppönen, E. APOA5 genotype influences the association between 25-hydroxyvitamin D and high density lipoprotein cholesterol. *Atherosclerosis* **2013**, *228*, 188–192. [CrossRef]
- 16. Bishwajit, G. Nutrition transition in South Asia: The emergence of non-communicable chronic diseases. *F1000Res* **2015**, *4*, 8. [CrossRef] [PubMed]
- 17. Williams, R.; Periasamy, M. Genetic and Environmental Factors Contributing to Visceral Adiposity in Asian Populations. *Endocrinol Metab* 2020, *35*, 681–695. [CrossRef] [PubMed]
- 18. Wuni, R.; Adela Nathania, E.; Ayyappa, A.K.; Lakshmipriya, N.; Ramya, K.; Gayathri, R.; Geetha, G.; Anjana, R.M.; Kuhnle, G.G.C.; Radha, V.; et al. Impact of Lipid Genetic Risk Score and Saturated Fatty Acid Intake on Central Obesity in an Asian Indian Population. *Nutrients* **2022**, *14*, 2713. [CrossRef] [PubMed]
- 19. Wuni, R.; Kuhnle, G.G.C.; Wynn-Jones, A.A.; Vimaleswaran, K.S. A Nutrigenetic Update on CETP Gene-Diet Interactions on Lipid-Related Outcomes. *Curr. Atheroscler. Rep.* **2022**, *24*, 119–132. [CrossRef] [PubMed]
- 20. Vimaleswaran, K.S. A nutrigenetics approach to study the impact of genetic and lifestyle factors on cardiometabolic traits in various ethnic groups: Findings from the GeNuIne Collaboration. *Proc. Nutr. Soc.* **2020**, *79*, 194–204. [CrossRef]
- Alsulami, S.; Aji, A.S.; Ariyasra, U.; Sari, S.R.; Tasrif, N.; Yani, F.F.; Lovegrove, J.A.; Sudji, I.R.; Lipoeto, N.I.; Vimaleswaran, K.S. Interaction between the genetic risk score and dietary protein intake on cardiometabolic traits in Southeast Asian. *Genes Nutr.* 2020, 15, 19. [CrossRef]
- 22. Dhanapal, A.; Wuni, R.; Ventura, E.F.; Chiet, T.K.; Cheah, E.S.G.; Loganathan, A.; Quen, P.L.; Appukutty, M.; Noh, M.F.M.; Givens, I.; et al. Implementation of Nutrigenetics and Nutrigenomics Research and Training Activities for Developing Precision Nutrition Strategies in Malaysia. *Nutrients* **2022**, *14*, 5108. [CrossRef]
- 23. Thiruvenkataswamy, C.S.; Appukutty, M.; Vimaleswaran, K.S. Role of precision nutrition in improving military performance. *Pers. Med.* **2022**, *19*, 167–170. [CrossRef]
- 24. Ching, Y.K.; Chin, Y.S.; Appukutty, M.; Ramanchadran, V.; Yu, C.Y.; Ang, G.Y.; Gan, W.Y.; Chan, Y.M.; Teh, L.K.; Salleh, M.Z. Interaction of Dietary Linoleic Acid and α-Linolenic Acids with rs174547 in FADS1 Gene on Metabolic Syndrome Components among Vegetarians. *Nutrients* 2019, *11*, 1686. [CrossRef]
- 25. Lee, S.S.; Ling, K.H.; Tusimin, M.; Subramaniam, R.; Rahim, K.F.; Loh, S.P. Interplay between Maternal and Neonatal Vitamin D Deficiency and Vitamin-D-Related Gene Polymorphism with Neonatal Birth Anthropometry. *Nutrients* **2022**, *14*, 564. [CrossRef]

- 26. Tan, P.Y.; Mitra, S.R. The Combined Effect of Polygenic Risk from FTO and ADRB2 Gene Variants, Odds of Obesity, and Post-Hipcref Diet Differences. *Lifestyle Genom.* **2020**, *13*, 84–98. [CrossRef] [PubMed]
- Yap, R.W.K.; Shidoji, Y.; Yap, W.S.; Masaki, M. Association and Interaction Effect of AGTR1 and AGTR2 Gene Polymorphisms with Dietary Pattern on Metabolic Risk Factors of Cardiovascular Disease in Malaysian Adults. *Nutrients* 2017, 9, 853. [CrossRef] [PubMed]
- 28. Yap, R.W.; Shidoji, Y.; Hon, W.M.; Masaki, M. Interaction between VEGF receptor-2 gene polymorphisms and dietary patterns on blood glucose and lipid levels in Chinese Malaysian adults. *J. Nutr. Nutr.* **2011**, *4*, 309–321. [CrossRef]
- Muhammad, H.F.L.; Sulistyoningrum, D.C.; Huriyati, E.; Lee, Y.Y.; Muda, W. The interaction between energy intake, physical activity and UCP2 -866G/A gene variation on weight gain and changes in adiposity: An Indonesian Nutrigenetic Cohort (INDOGENIC). Br. J. Nutr. 2021, 125, 611–617. [CrossRef] [PubMed]
- 30. Huriyati, E.; Luglio, H.F.; Ratrikaningtyas, P.D.; Tsani, A.F.; Sadewa, A.H.; Juffrie, M. Dyslipidemia, insulin resistance and dietary fat intake in obese and normal weight adolescents: The role of uncoupling protein 2 -866G/A gene polymorphism. *TInt. J. Mol. Epidemiol. Genet.* **2016**, *7*, 67–73.
- Muhammad, H.F.L.; Sulistyoningrum, D.C.; Huriyati, E.; Lee, Y.Y.; Manan Wan Muda, W.A. The Interaction between Coffee: Caffeine Consumption, UCP2 Gene Variation, and Adiposity in Adults-A Cross-Sectional Study. *J. Nutr. Metab.* 2019, 2019, 9606054. [CrossRef]
- 32. Surendran, S.; Aji, A.S.; Ariyasra, U.; Sari, S.R.; Malik, S.G.; Tasrif, N.; Yani, F.F.; Lovegrove, J.A.; Sudji, I.R.; Lipoeto, N.I.; et al. A nutrigenetic approach for investigating the relationship between vitamin B12 status and metabolic traits in Indonesian women. *J. Diabetes Metab. Disord.* **2019**, *18*, 389–399. [CrossRef]
- 33. Aji, A.S.; Lipoeto, N.I.; Yusrawati, Y.; Malik, S.G.; Kusmayanti, N.A.; Susanto, I.; Nurunniyah, S.; Alfiana, R.D.; Wahyuningsih, W.; Majidah, N.M.; et al. Impact of maternal dietary carbohydrate intake and vitamin D-related genetic risk score on birth length: The Vitamin D Pregnant Mother (VDPM) cohort study. *BMC Pregnancy Childbirth* 2022, 22, 690. [CrossRef]
- 34. Chang, X.; Dorajoo, R.; Sun, Y.; Han, Y.; Wang, L.; Khor, C.C.; Sim, X.; Tai, E.S.; Liu, J.; Yuan, J.M.; et al. Gene-diet interaction effects on BMI levels in the Singapore Chinese population. *Nutr. J.* **2018**, *17*, 31. [CrossRef]
- 35. Huang, T.; Wang, T.; Heianza, Y.; Wiggs, J.; Sun, D.; Choi, H.K.; Chai, J.F.; Sim, X.; Khor, C.C.; Friedlander, Y.; et al. Fish and marine fatty acids intakes, the FADS genotypes and long-term weight gain: A prospective cohort study. *BMJ Open* **2019**, *9*, e022877. [CrossRef]
- Corella, D.; Tai, E.S.; Sorlí, J.V.; Chew, S.K.; Coltell, O.; Sotos-Prieto, M.; García-Rios, A.; Estruch, R.; Ordovas, J.M. Association between the APOA2 promoter polymorphism and body weight in Mediterranean and Asian populations: Replication of a gene-saturated fat interaction. *Int. J. Obes.* 2011, 35, 666–675. [CrossRef] [PubMed]
- Tai, E.S.; Corella, D.; Deurenberg-Yap, M.; Adiconis, X.; Chew, S.K.; Tan, C.E.; Ordovas, J.M. Differential effects of the C1431T and Pro12Ala PPARgamma gene variants on plasma lipids and diabetes risk in an Asian population. *J. Lipid Res.* 2004, 45, 674–685. [CrossRef] [PubMed]
- Lim, S.Y.; Chan, Y.M.; Ramachandran, V.; Shariff, Z.M.; Chin, Y.S.; Arumugam, M. Dietary Acid Load and Its Interaction with IGF1 (rs35767 and rs7136446) and IL6 (rs1800796) Polymorphisms on Metabolic Traits among Postmenopausal Women. *Nutrients* 2021, 13, 2161. [CrossRef] [PubMed]
- 39. Mitra, S.R.; Tan, P.Y.; Amini, F. Association of ADRB2 rs1042713 with Obesity and Obesity-Related Phenotypes and Its Interaction with Dietary Fat in Modulating Glycaemic Indices in Malaysian Adults. *J. Nutr. Metab.* **2019**, 2019, 8718795. [PubMed]
- Li, H.; Khor, C.C.; Fan, J.; Lv, J.; Yu, C.; Guo, Y.; Bian, Z.; Yang, L.; Millwood, I.Y.; Walters, R.G.; et al. Genetic risk, adherence to a healthy lifestyle, and type 2 diabetes risk among 550,000 Chinese adults: Results from 2 independent Asian cohorts. *Am. J. Clin. Nutr.* 2020, 111, 698–707. [PubMed]
- Corella, D.; Qi, L.; Tai, E.S.; Deurenberg-Yap, M.; Tan, C.E.; Chew, S.K.; Ordovas, J.M. Perilipin gene variation determines higher susceptibility to insulin resistance in Asian women when consuming a high-saturated fat, low-carbohydrate diet. *Diabetes Care* 2006, 29, 1313–1319. [CrossRef] [PubMed]
- 42. Downes, M.J.; Brennan, M.L.; Williams, H.C.; Dean, R.S. Development of a critical appraisal tool to assess the quality of cross-sectional studies (AXIS). *BMJ Open* **2016**, *6*, e011458. [CrossRef] [PubMed]
- Sterne, J.A.; Hernán, M.A.; Reeves, B.C.; Savović, J.; Berkman, N.D.; Viswanathan, M.; Henry, D.; Altman, D.G.; Ansari, M.T.; Boutron, I.; et al. ROBINS-I: A tool for assessing risk of bias in non-randomised studies of interventions. *BMJ* 2016, 355, i4919. [CrossRef]
- 44. RoB 2: A Revised Cochrane Risk-Of-Bias Tool for Randomized Trials | Cochrane Bias. Available online: https://methods.cochrane.org/bias/resources/rob-2-revised-cochrane-risk-bias-tool-randomized-trials (accessed on 24 October 2022).
- 45. Liu, S.J.; Zhi, H.; Chen, P.Z.; Chen, W.; Lu, F.; Ma, G.S.; Dai, J.C.; Shen, C.; Liu, N.F.; Hu, Z.B.; et al. Fatty acid desaturase 1 polymorphisms are associated with coronary heart disease in a Chinese population. *Chin. Med. J.* **2012**, *125*, 801–806.
- 46. Agarwal, S.; Kovilam, O.; Agrawal, D.K. Vitamin D and its impact on maternal-fetal outcomes in pregnancy: A critical review. *Crit. Rev. Food Sci. Nutr.* **2018**, *58*, 755–769. [CrossRef]
- Cohen, J.B.; Stephens-Shields, A.J.; Denburg, M.R.; Anderson, A.H.; Townsend, R.R.; Reese, P.P. Obesity, Renin-Angiotensin System Blockade and Risk of Adverse Renal Outcomes: A Population-Based Cohort Study. *Am. J. Nephrol.* 2016, 43, 431–440. [CrossRef] [PubMed]

- 48. Pahlavani, M.; Kalupahana, N.S.; Ramalingam, L.; Moustaid-Moussa, N. Regulation and Functions of the Renin-Angiotensin System in White and Brown Adipose Tissue. *Compr. Physiol.* **2017**, *7*, 1137–1150. [PubMed]
- Tam, J.; Duda, D.G.; Perentes, J.Y.; Quadri, R.S.; Fukumura, D.; Jain, R.K. Blockade of VEGFR2 and not VEGFR1 can limit dietinduced fat tissue expansion: Role of local versus bone marrow-derived endothelial cells. *PLoS ONE* 2009, *4*, e4974. [CrossRef] [PubMed]
- 50. Fawcett, K.A.; Barroso, I. The genetics of obesity: FTO leads the way. Trend Genet. 2010, 26, 266–274. [CrossRef]
- Abbas, M.A. Physiological functions of Vitamin D in adipose tissue. *J. Steroid Biochem. Mol. Biol.* 2017, 165, 369–381. [CrossRef]
 Sirotkin, A.V.; Kolesárová, A. The anti-obesity and health-promoting effects of tea and coffee. *Physiol. Res.* 2021, 70, 161–168. [CrossRef]
- 53. Pan, M.H.; Tung, Y.C.; Yang, G.; Li, S.; Ho, C.T. Molecular mechanisms of the anti-obesity effect of bioactive compounds in tea and coffee. *Food Funct.* **2016**, *7*, 4481–4491.
- Li, J.; Jiang, R.; Cong, X.; Zhao, Y. UCP2 gene polymorphisms in obesity and diabetes, and the role of UCP2 in cancer. *FEBS Lett.* 2019, 593, 2525–2534. [CrossRef]
- 55. Allister, E.M.; Robson-Doucette, C.A.; Prentice, K.J.; Hardy, A.B.; Sultan, S.; Gaisano, H.Y.; Kong, D.; Gilon, P.; Herrera, P.L.; Lowell, B.B.; et al. UCP2 regulates the glucagon response to fasting and starvation. *Diabetes* **2013**, *62*, 1623–1633. [CrossRef]
- 56. Evans, R.M.; Barish, G.D.; Wang, Y.X. PPARs and the complex journey to obesity. Nat. Med. 2004, 10, 355–361. [CrossRef]
- 57. Murff, H.J.; Edwards, T.L. Endogenous Production of Long-Chain Polyunsaturated Fatty Acids and Metabolic Disease Risk. *Curr. Cardiovasc. Risk Rep.* **2014**, *8*, 1–9. [CrossRef]
- 58. Zaki, M.E.; Amr, K.S.; Abdel-Hamid, M. APOA2 Polymorphism in Relation to Obesity and Lipid Metabolism. *Cholesterol* 2013, 2013, 289481. [CrossRef]
- 59. Wang, Q.; Wang, J.; Lin, H.; Huo, X.; Zhu, Q.; Zhang, M. Relationship between fat mass and obesity-associated gene expression and type 2 diabetes mellitus severity. *Exp. Ther. Med.* **2018**, *15*, 2917–2921. [CrossRef]
- Grant, S.F.; Thorleifsson, G.; Reynisdottir, I.; Benediktsson, R.; Manolescu, A.; Sainz, J.; Helgason, A.; Stefansson, H.; Emilsson, V.; Helgadottir, A.; et al. Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat. Genet.* 2006, 38, 320–323. [CrossRef] [PubMed]
- 61. Ceasrine, A.M.; Lin, E.E.; Lumelsky, D.N.; Iyer, R.; Kuruvilla, R. Adrb2 controls glucose homeostasis by developmental regulation of pancreatic islet vasculature. *Elife* **2018**, *7*, e39689. [CrossRef] [PubMed]
- 62. Zhuang, Y.; Niu, F.; Liu, D.; Sun, J.; Zhang, X.; Zhang, J.; Guo, S. Association between AGTR1 A1166C polymorphism and the susceptibility to diabetic nephropathy: Evidence from a meta-analysis. *Medicine* **2018**, *97*, e07689. [CrossRef]
- 63. Chou, E.; Suzuma, I.; Way, K.J.; Opland, D.; Clermont, A.C.; Naruse, K.; Suzuma, K.; Bowling, N.L.; Vlahos, C.J.; Aiello, L.P.; et al. Decreased cardiac expression of vascular endothelial growth factor and its receptors in insulin-resistant and diabetic States: A possible explanation for impaired collateral formation in cardiac tissue. *Circulation* **2002**, *105*, 373–379. [CrossRef] [PubMed]
- 64. Møller, N.; Jørgensen, J.O. Effects of growth hormone on glucose, lipid, and protein metabolism in human subjects. *Endocr. Rev.* **2009**, *30*, 152–177. [CrossRef]
- 65. Rehman, K.; Akash, M.S.H.; Liaqat, A.; Kamal, S.; Qadir, M.I.; Rasul, A. Role of Interleukin-6 in Development of Insulin Resistance and Type 2 Diabetes Mellitus. *Crit. Rev. Eukaryot. Gene Expr.* **2017**, *27*, 229–236. [CrossRef]
- 66. Liu, L.; Huang, X.; Wang, B.; Song, Y.; Lin, T.; Zhou, Z.; Wang, Z.; Wei, Y.; Guo, H.; Chen, P.; et al. Vitamin B(12) and risk of diabetes: New insight from cross-sectional and longitudinal analyses of the China Stroke Primary Prevention Trial (CSPPT). *BMJ Open Diabetes Res. Care* **2020**, *8*, e001423. [CrossRef]
- 67. Duesing, K.; Charpentier, G.; Marre, M.; Tichet, J.; Hercberg, S.; Balkau, B.; Froguel, P.; Gibson, F. Evaluating the association of common APOA2 variants with type 2 diabetes. *BMC Med. Genet.* **2009**, *10*, 13.
- Castellani, L.W.; Goto, A.M.; Lusis, A.J. Studies with apolipoprotein A-II transgenic mice indicate a role for HDLs in adiposity and insulin resistance. *Diabetes* 2001, 50, 643–651. [CrossRef] [PubMed]
- 69. Rangwala, S.M.; Lazar, M.A. Peroxisome proliferator-activated receptor gamma in diabetes and metabolism. *Trends Pharmacol. Sci.* **2004**, *25*, 331–336. [CrossRef]
- 70. Dumont, J.; Goumidi, L.; Grenier-Boley, B.; Cottel, D.; Marécaux, N.; Montaye, M.; Wagner, A.; Arveiler, D.; Simon, C.; Ferrières, J.; et al. Dietary linoleic acid interacts with FADS1 genetic variability to modulate HDL-cholesterol and obesity-related traits. *Clin. Nutr.* **2018**, *37*, 1683–1689. [CrossRef]
- 71. Chen, Y.; Estampador, A.C.; Keller, M.; Poveda, A.; Dalla-Riva, J.; Johansson, I.; Renström, F.; Kurbasic, A.; Franks, P.W.; Varga, T.V. The combined effects of FADS gene variation and dietary fats in obesity-related traits in a population from the far north of Sweden: The GLACIER Study. *Int. J. Obes.* 2019, 43, 808–820. [CrossRef]
- Qi, L.; Zhang, C.; Greenberg, A.; Hu, F.B. Common variations in perilipin gene, central obesity, and risk of type 2 diabetes in US women. *Obesity* 2008, 16, 1061–1065. [CrossRef] [PubMed]
- 73. Yu, D.; Li, C.; Xie, J.; Xu, G.; Li, Y.; Liu, J.; Chen, B.; Pan, J.; Shen, M.; Yang, L.; et al. Association between three genetic variants of the Perilipin Gene (PLIN) and glucose metabolism: Results from a replication study among Chinese adults and a meta-analysis. *Endocr. Res.* **2013**, *38*, 263–279. [CrossRef]

- 74. Ferguson, L.R.; De Caterina, R.; Görman, U.; Allayee, H.; Kohlmeier, M.; Prasad, C.; Choi, M.S.; Curi, R.; de Luis, D.A.; Gil, Á.; et al. Guide and Position of the International Society of Nutrigenetics/Nutrigenomics on Personalised Nutrition: Part 1Fields of Precision Nutrition. J. Nutr. Nutr. 2016, 9, 12–27. [CrossRef]
- 75. Maria Catherine, B. Otero LFB: Role of Nutrigenomics in Modern-Day Healthcare and Drug Discovery; Elsevier: Amsterdam, The Netherlands, 2023.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.