

Methane prediction equations including genera of rumen bacteria as predictor variables improve prediction accuracy

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Zhang, B., Lin, S., Moraes, L., Firkins, J., Hristov, A. N., Kebreab, E., Janssen, P. H., Bannink, A., Bayat, A. R., Crompton, L. A., Dijkstra, J., Eugène, M. A., Kreuzer, M., McGee, M., Reynolds, C. K. ORCID: <https://orcid.org/0000-0002-4152-1190>, Schwarm, A., Yáñez-Ruiz, D. R. and Yu, Z. (2023) Methane prediction equations including genera of rumen bacteria as predictor variables improve prediction accuracy. *Scientific Reports*, 13. 21305. ISSN 2045-2322 doi: [10.1038/s41598-023-48449-y](https://doi.org/10.1038/s41598-023-48449-y) Available at <https://centaur.reading.ac.uk/114218/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1038/s41598-023-48449-y>

Publisher: Nature Publishing Group

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in

the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



OPEN

Methane prediction equations including genera of rumen bacteria as predictor variables improve prediction accuracy

Boyang Zhang¹, Shili Lin²✉, Luis Moraes^{1,15}, Jeffrey Firkins¹, Alexander N. Hristov³, Ermias Kebreab⁴, Peter H. Janssen⁵, André Bannink⁶, Alireza R. Bayat⁷, Les A. Crompton⁸, Jan Dijkstra⁹, Maguy A. Eugène¹⁰, Michael Kreuzer¹¹, Mark McGee¹², Christopher K. Reynolds⁸, Angela Schwarm¹³, David R. Yáñez-Ruiz¹⁴ & Zhongtang Yu¹✉

Methane (CH₄) emissions from ruminants are of a significant environmental concern, necessitating accurate prediction for emission inventories. Existing models rely solely on dietary and host animal-related data, ignoring the predicting power of rumen microbiota, the source of CH₄. To address this limitation, we developed novel CH₄ prediction models incorporating rumen microbes as predictors, alongside animal- and feed-related predictors using four statistical/machine learning (ML) methods. These include random forest combined with boosting (RF-B), least absolute shrinkage and selection operator (LASSO), generalized linear mixed model with LASSO (glmmLasso), and smoothly clipped absolute deviation (SCAD) implemented on linear mixed models. With a sheep dataset (218 observations) of both animal data and rumen microbiota data (relative sequence abundance of 330 genera of rumen bacteria, archaea, protozoa, and fungi), we developed linear mixed models to predict CH₄ production (g CH₄/animal-d, ANIM-B models) and CH₄ yield (g CH₄/kg of dry matter intake, DMI-B models). We also developed models solely based on animal-related data. Prediction performance was evaluated 200 times with random data splits, while fitting performance was assessed without data splitting. The inclusion of microbial predictors improved the models, as indicated by decreased root mean square prediction error (RMSPE) and mean absolute error (MAE), and increased Lin's concordance correlation coefficient (CCC). Both glmmLasso and SCAD reduced the Akaike information criterion (AIC) and Bayesian information criterion (BIC) for both the ANIM-B and the DMI-B models, while the other two ML methods had mixed outcomes. By balancing prediction performance and fitting performance, we obtained one ANIM-B model (containing 10 genera of bacteria and 3 animal data) fitted using glmmLasso and one DMI-B model (5 genera of bacteria and 1 animal datum) fitted using SCAD. This study highlights the importance of incorporating rumen microbiota data in CH₄ prediction models to enhance accuracy and robustness. Additionally, ML methods facilitate the selection of microbial predictors from high-dimensional metataxonomic data of the rumen microbiota without overfitting. Moreover, the identified microbial predictors can serve as biomarkers of CH₄ emissions from sheep, providing valuable insights for future research and mitigation strategies.

¹Department of Animal Sciences, The Ohio State University, Columbus, OH 43210, USA. ²Department of Statistics, The Ohio State University, 2029 Fyffe Road, Columbus, OH 43210, USA. ³Department of Animal Science, The Pennsylvania State University, University Park, PA, USA. ⁴Department of Animal Science, University of California, Davis, CA, USA. ⁵AgResearch Limited, Grasslands Research Centre, Palmerston North 4442, New Zealand. ⁶Wageningen Livestock Research, Wageningen University & Research, Wageningen, The Netherlands. ⁷Milk Production, Production Systems, Natural Resources Institute Finland (Luke), 31600 Jokioinen, Finland. ⁸School of Agriculture, Policy, and Development, University of Reading, Reading, UK. ⁹Animal Nutrition Group, Wageningen University & Research, Wageningen, The Netherlands. ¹⁰INRAE UMR Herbivores, VetAgro Sup, Université Clermont Auvergne, Saint-Genès-Champanelle, France. ¹¹Institute of Agricultural Sciences, ETH Zurich, Zurich, Switzerland. ¹²Teagasc, AGRIC, Grange, Dunsany, CO., Meath, Ireland. ¹³Department of Animal and Aquacultural Sciences, Norwegian University of Life Sciences, Ås, Norway. ¹⁴Estación Experimental del Zaidin (CSIC), Granada, Spain. ¹⁵Present address: Consultoria, Piracicaba, SP, Brazil. ✉email: shili@stat.osu.edu; yu.226@osu.edu

Abbreviations

CH ₄	Methane
ANIM-B	Animal-based
A:P ratio	Acetate:propionate ratio
ITS1	Internal transcribed spacer 1
ASV	Amplicon sequence variant
VIF	Variance inflation factor
Uncl	Uncultured
BW	Body weight
DMI	Dry matter intake
DMI-B	Dry matter-based
ML	Machine learning
RF	Random forest
RF-B	Random forest combined with boosting
LASSO	Least absolute shrinkage and selection operator
glimmLasso	Generalized linear mixed with LASSO
SCAD	Smoothly clipped absolute deviation
AIC	Akaike information criterion
BIC	Bayesian information criterion
SE	Standard errors
SIS	Sure independence screening
MSPE	Mean square prediction error
RMSPE	Root mean square prediction error
MAE	Mean absolute errors
CCC	Concordance correlation coefficient

As a potent greenhouse gas that contributes significantly to climate change, methane (CH₄) emissions from ruminants pose a direct threat to the environment and sustainable agricultural production¹. Methane emissions from ruminants also lead to a waste of part of the ingested feed energy². In sheep, CH₄ emissions result in approximately a 7% loss of gross energy intake³. In recent years, there has been growing interest in prediction models for CH₄ emissions for several reasons. First, direct measurement of CH₄ emissions from large herds or flocks of ruminants is technically difficult and time-consuming, and thus it can be done practically only with small numbers of animals in research settings⁴. Second, it is not practical or feasible to establish regional, national, or global inventories of CH₄ emissions by direct measurement of CH₄ emissions from individual animals. Third, statistical models can assist in describing and understanding the relationship among CH₄ emissions, animal performance, and other factors (nutritional, microbial, and physiological).

A diverse set of prediction models has been developed to predict enteric CH₄ emissions from sheep⁵ and cattle^{6,7}. These prediction equations include predictor variables describing the dietary composition, feed intake, rumen fermentation profiles, and other animal traits such as body weight (BW), but not individual taxa (genera or species) of the rumen microbiota. These prediction models vary considerably in robustness and prediction accuracy^{8,9}. The rumen microbiota is solely responsible for CH₄ emissions from ruminants. Specifically, rumen bacteria, fungi, and protozoa digest and ferment the ingested feed and produce the substrates (methylamines, hydrogen, CO₂, and formate) that are then used by the rumen methanogens to produce CH₄¹⁰. Thus, the rumen microbiota is directly responsible for the CH₄ emitted by ruminants although it can be affected by several factors, including diet and host-related factors¹¹. Indeed, several studies have reported strong associations or correlations of some bacteria, methanogens, or protozoa with or qualitatively indicative of CH₄ production^{12,13}. In this context, we hypothesized that the inclusion of individual taxa of the ruminal microbiota as predictor variables, in addition to animal and feed-related predictor variables, in CH₄ prediction models would improve our ability to predict CH₄ emissions. Methane prediction equations or models containing rumen microbial taxa as predictor variables may also help understand their roles and importance in enteric CH₄ emissions. The objective of this study was to develop such prediction models using multiple statistical/machine learning (ML) methods and a dataset that contained both animal-related data and metataxonomic data of the rumen microbiota of a cohort of sheep that were not subjected to any dietary interventions.

Methods

Dataset

This study did not involve the use of any animals, and thus no approval of the Institutional Animal Care and Use Committee was required. The dataset (Supplementary Table 1) used in this study was generated in a previous study that compared the rumen microbiota and CH₄ emissions among a group of New Zealand sheep that were not subjected to any dietary interventions¹³. The animal experiment and analyses were detailed in another report¹⁴. Briefly, 1225 sheep were fed a pelleted alfalfa diet (without any anti-methane inhibitors) twice daily at two times their maintenance requirement for energy (based on BW). CH₄ emissions from 340 individual animals were measured with open-circuit respiration chambers twice in two independent measuring rounds (10–15 days apart) with each time lasting for two consecutive days¹⁴. CH₄ production (g CH₄/animal·d) and CH₄ yield (g CH₄/kg of dry matter intake, DMI) were calculated. Feed intake and BW of the 340 individual animals were also recorded. Rumen samples were collected from 118 of the 340 sheep, with 60 representing the highest and 58 representing the lowest CH₄ yield sheep, immediately after each CH₄ emission measurement (corresponding to 17–18 h after the last feeding) using the stomach tubing method. The two groups of sheep were

balanced by BW (24–60 kg), sex, and breeds (indicated below). The rumen fermentation characteristics, including acetate:propionate (A:P) ratio, were analyzed using gas–liquid chromatography. We provided the density plots of the CH₄ data (g/d and g/kg of DMI, Supplementary Fig. 1). Both density curves appeared to represent the mixture of the two groups, which is consistent with the data feature (high and low emission sheep) and with the patterns found in the previous study¹³.

Two rumen samples were collected (10 to 15 days apart) from each sheep and the composition and structure of the rumen microbiota were analyzed using metataxonomics sequencing the 16S rRNA genes of both bacteria and archaea, 18S rRNA gene of protozoa, and internal transcribed spacer 1 (ITS1) of fungi¹³. One sheep was excluded due to missing data, and 16 sheep were represented by only one sample that yielded complete metataxonomic data. This resulted in a cumulative 218 observations, with 101 sheep each contributing two observations and the remaining 16 sheep contributing one observation.

In this study, we retrieved the metataxonomic sequencing data of the sheep rumen samples from the EMBL database per the accession numbers (ERP003779 for Bacteria, ERP003773 for Archaea, ERP003772 for ciliate protozoa, and ERP003764 for anaerobic fungi). The sequences were then analyzed using DADA2¹⁵. Briefly, the amplicon sequences were clustered into amplicon sequence variants (ASVs), which were then taxonomically assigned to individual taxa using specialty databases: the Silva database (version 138.1) for bacteria, archaea, and protozoa¹⁶; and the UNITE database for fungi¹⁷. The relative sequence abundance of each genus was calculated as the sum of the relative sequence abundance of all the ASVs assigned to the respective genus. The sequences of the ASVs that were classified to an unclassified genus (Uncl_genus) are listed in Supplementary Table 2. Only the genera that each had a relative sequence abundance reaching > 1% in all the samples were used as microbial variables. For an Uncl_genus, its higher and known taxon name was used to show its taxonomic lineage. In total, 330 microbial genera were obtained and used, including 308 genera of bacteria, 9 genera of archaea, 5 genera of protozoa, and 8 genera of fungi. The relative sequence abundance data of the microbial genera were log-transformed before the modeling process. In addition to the rumen microbiota data and CH₄ emission data (CH₄ production, g CH₄/animal-d; CH₄ yield, g CH₄/kg of DMI), the dataset also contained the following five animal-related data of the individual sheep: DMI (kg/d), BW (kg), A:P ratio, sex, and breed (Coopworth, Corriedale, Finnish Landrace, Perendale, Romney, and Texel).

Linear mixed model

In our dataset, most animals (101 out of 118) had two records of CH₄ emissions and other data (except for sex and breed). These repeated measurements from the same sheep were likely to be more akin to each other than to the measurements from a different sheep. Thus, the two records from the same sheep were expected to be correlated. Linear mixed-effects models were used to model the dependence of the data and account for correlations between the two repeated measurements. The general structure of the model was described as follows:

$$y_{ij} = \mathbf{X}_i\boldsymbol{\beta} + \alpha_i + \varepsilon_{ij} \quad (1)$$

where y_{ij} is the observed CH₄ emissions from the i th animal and j th observation ($i = 1, \dots, 118$ and $j = 1, 2$), set as either g CH₄/animal-d (referred to as ANIM-B models) or g CH₄/kg of DMI (referred to as DMI-B models); $\mathbf{X}_i = (1, \text{DMI}, \text{BW}, \text{A:P ratio}, \text{sex}, \text{breed}, \text{microbial variables})$ is the design vector for the fixed effects containing the values of the predictor variables for the i th animal and the “1” entry for the intercept (DMI was not included as a variable in developing the DMI-B models because it is a dependent variable in those models); $\boldsymbol{\beta}$ is the coefficients vector of the fixed effects describing the relationships between CH₄ emissions and the predictor variables; α_i is the random animal effects to account for intra-animal correlation between the two measurements; and ε_{ij} is the random error. The α_i was assumed to be independent and identically distributed with mean zero and variance σ_α^2 , and the ε_{ij} errors were assumed to be independent and identically distributed with mean zero and variance σ^2 . Further, it was assumed that random effects and errors were mutually independent.

Machine learning methods

The rumen microbiota is diverse, containing many hundreds to thousands of distinct species of microbes at varying relative sequence abundance. The inclusion of such a large number of microbial variables in prediction models can lead to overfitting. Additionally, relative abundance data of microbiota are usually on different scales, zero-inflated, sparse, and over-dispersed. Thus, proper selection of microbial predictor variables to be included in prediction models is crucial in model development. In the present study, we used four ML methods to select microbial predictor variables: random forest combined with boosting (RF-B) using the randomForest and gbm packages¹⁸, least absolute shrinkage and selection operator (LASSO) using the glmnet package¹⁹, generalized linear mixed model with LASSO (glmLasso) using the glmLasso package²⁰, and smoothly clipped absolute deviation (SCAD) using the splmm package²¹. We opted for both LASSO and glmLasso because the former does not consider intra-animal correlation, while the latter takes these correlations into account by utilizing the linear mixed-effect model described in Eq. (1). Each of the ML methods was evaluated for both prediction performance and fitting performance. Specifically, to evaluate prediction performance (based on 200 evaluations with random data splitting into testing data and training data (see section ‘Evaluation of prediction performance’ below), each ML method was used to make variable selection (selection for both animal and microbial predictor variables). Using the lmer function of the lme4 package²², we also fitted conventional models utilizing solely the animal-related data (i.e., BW, A:P ratio, and DMI (DMI was used only in developing the ANIM-B models)) as predictors. The two categorical variables (sex and breed) were not included in the prediction models because their addition did not lead to improvement in prediction performance (data not shown). The sets of variables selected by glmLasso, LASSO, SCAD, RF-B, and conventional modeling were then separately fitted with a linear mixed model, and the resultant models were used to predict CH₄ emissions. Further, to evaluate fitting

performances, linear mixed models were fitted with all available data by selecting the variables using the ML and the conventional method. The Akaike information criterion (AIC) and Bayesian information criterion (BIC) of those models were computed using the ANOVA function in R v4.0.2²³. The estimated standard deviation of the animal random effects ($\hat{\sigma}_a$), coefficients, and associated standard errors (SE) of the predictor variables were computed using the lmer function²². The observations and residuals versus the model predictions were plotted to evaluate the linear mixed model assumptions for all the ML models.

Evaluation of prediction performance

Prediction performance was validated in five steps. Step 1 involved screening the independent variables using the sure independence screening (SIS) test implemented in the SIS package in R²⁴. This step was omitted for the RF-B method because this method can handle high-dimensional data without the need for data prescreening when building the decision trees. To check for potential multicollinearity, the variance inflation factor (VIF) was examined, and variables with a VIF < 5 were retained as recommended by Niu et al.⁷.

Step 2 began by randomly splitting the dataset into testing data (20% of the data) and training data (the remaining data). The training data were standardized by mean centering (subtracting each variable from the corresponding mean) and scaling (dividing each variable by its corresponding standard deviation). The testing data were standardized based on the centering and scaling from the training data. To prevent data leaking of the repeated records from the same animal, the two repeated records were always allocated to either the testing data or the training data.

In step 3, glmmLasso, LASSO, SCAD, and RF-B were individually used to select predictor variables from the training data. For glmmLasso and SCAD, the optimal tuning parameter λ was determined using BIC. Two increments of λ , 0.1 and 0.5, were also evaluated, but the results were virtually the same (data not shown) except for a substantial increase in computing time. For LASSO, tenfold cross-validation was used to determine the optimal tuning parameter λ . For RF-B, the top 10 important variables identified using the randomForest package were selected when the number of decision trees was set as 500, and the number of variables randomly sampled as candidates at each split (using mtry in the randomForest package) was set as 40. The top 10 important variables determined by boosting using the gbm package were also selected with the default settings applied. The union of the variables selected by random forest and boosting was kept.

Step 4 involved separately refitting a linear mixed model with the sets of selected variables using each of the ML methods and the training data. Predictions were then made using the values of the variables from the testing data. All the predictions and observations were recorded.

Step 5 repeated steps 2 to 4 for 200 times (sufficiently large and not too computationally intensive) by re-splitting the data into testing and training data in each iteration. Mean squared prediction error (MSPE), root MSPE (RMSPE), mean absolute errors (MAE), and Lin's concordance correlation coefficients (CCC) were calculated using the equations specified in the section 'Model evaluation equations' below. We utilized RMSPE, MAE, and CCC to provide a comprehensive evaluation of the model performance. In particular, MAE is a robust assessment that is less sensitive to extreme values compared to RMSPE, whereas CCC can provide information on the precision and accuracy of the prediction.

Model evaluation equations

The values of MSPE, RMSPE, and MAE of all the models were calculated using the equations shown below²⁵. The unit of MSPE was $(g/d)^2$ for the ANIM-B models and $(g/kg)^2$ for the DMI-B models. Lin's CCC was calculated by the epi.ccc function in the epiR package in R²⁶.

$$MSPE = \frac{\sum_{i=1}^N (O_i - P_i)^2}{N}$$

$$MAE = \frac{\sum_{i=1}^N |O_i - P_i|}{N}$$

$$RMSPE = \sqrt{MSPE}$$

$$CCC = \frac{2s_{op}}{s_o^2 + s_p^2 + (\bar{o} - \bar{p})^2}$$

where O_i is the CH_4 emissions of i th observation; P_i is CH_4 prediction from the model; N is the total number of observations; S_o is the standard deviation of the observations; S_p is the standard deviation of the predictions; \bar{o} is the mean of the observations; \bar{p} is the mean of the predictions; S_{op} is the covariance of the observations and the predictions.

The R codes we used in the present study were available at GitHub (https://github.com/yu2269/CH4_prediction_model).

Results

Model performance

Tables 1 and 2 summarize the prediction performances of the ANIM-B and the DMI-B models developed using both the ML and the conventional methods, based on random data splitting (200 times). The ML methods

	Conventional	glmLasso	LASSO	SCAD	RF-B
RMSPE	2.96	2.85	2.85	3.00	2.91
Reduction of RMSPE (%)	–	3.80	3.62	-1.25	1.52
MAE	2.29	2.18	2.17	2.29	2.21
Reduction of MAE (%)	–	4.60	5.08	-0.39	3.50
CCC	0.64	0.70	0.71	0.66	0.68
Increase of CCC (%)	–	9.49	9.80	2.64	5.29

Table 1. Prediction performance of the animal-based (ANIM-B) models (CH₄ production; g/d) developed using conventional method and four machine learning methods. *The conventional method only used animal-related data; the relative abundance of all the microbial data was log-transformed; glmLasso, generalized linear mixed model combined with LASSO; LASSO, least absolute shrinkage and selection operator; SCAD, smoothly clipped absolute deviation implemented on linear mixed models; RF-B, random forest combined with boosting. The data were randomly split into a training set and a testing set (80:20) 200 times and were standardized by mean centering and scaling (detailed in “Methods”).

	Conventional	glmLasso	LASSO	SCAD	RF-B
RMSPE	2.12	1.99	1.89	1.92	1.99
Reduction of RMSPE (%)	–	6.22	11.12	9.66	6.50
MAE	1.65	1.51	1.43	1.48	1.52
Reduction of MAE (%)	–	8.21	13.31	9.97	7.36
CCC	0.28	0.46	0.57	0.51	0.48
Increase of CCC (%)	–	68.73	106.18	84.36	73.09

Table 2. Prediction performance of the dry matter intake-based (DMI-B) models (CH₄ yield; g/kg DMI) developed using conventional method and four machine learning methods. See the notes of Table 1.

notably improved the performance of ANIM-B models, reducing RMSPE (by – 1.25% for SCAD and 3.8% for glmLasso) and MAE (by – 0.39% for SCAD and 5.08% for LASSO), while increasing CCC (by 2.64% for SCAD and 9.8% for LASSO), as compared to the conventional method. Similarly, the ML methods enhanced the performance of DMI-B models, reducing RMSPE (by 6.22% for glmLasso and 11.12% for LASSO) and MAE (by 7.36% for RF-B and 13.31% for LASSO) and increasing CCC (by 68.73% for glmLasso and 106.18% for LASSO). Tables 3 and 4 present the fitting performance of the ANIM-B and the DMI-B models developed using both the ML methods and the conventional method, based on the entire dataset. Compared to the conventional method, the ML methods decreased BIC (by – 33 for LASSO and 19 for glmLasso) and AIC (by – 152 for LASSO and 53 for glmLasso) for the ANIM-B models. Similarly, for the DMI-B models, the ML methods decreased BIC (by – 201 for RF-B and 5 for glmLasso and SCAD) and AIC (by – 157 for RF-B and 18 for glmLasso), as compared to the conventional method.

The final models were selected by balancing the prediction performance and the fitting performance. Among the ANIM-B models, the glmLasso model had the lowest BIC and AIC (Table 3). It also had the lowest RMSPE in prediction performance (Table 1). Thus, we selected the ANIM-B glmLasso model as the final model. For the prediction performance of the DMI-B models, the LASSO model had a lower RMSPE than the SCAD model (1.89 vs. 1.92), which was lower than that of the glmLasso model (Table 2), but the LASSO model had a higher BIC and AIC than the SCAD model and the glmLasso model (Table 4). The DMI-B SCAD model also had

	Conventional	glmLasso	LASSO	SCAD	RF-B
BIC	1070	1051	1103	1065	1095
Reduction of BIC	–	19	– 33	5	– 25
AIC	1050	997	1202	1028	1031
Reduction of AIC	–	53	– 152	22	19
Number of variables	3	13	27	8	17

Table 3. Fitting performance of the animal-based (ANIM-B) models (CH₄ production; g/d) fitted with the variables selected from all the available data by the conventional method and machine learning methods. See the notes of Table 1.

	Conventional	gmmLasso	LASSO	SCAD	RF-B
BIC	894	889	969	889	1095
Reduction of BIC	–	5	– 75	5	– 201
AIC	874	856	868	859	1031
Reduction of AIC	–	18	6	15	– 157
Number of variables	2	9	25	6	17

Table 4. Fitting performance of the dry matter intake-based (DMI-B) models (CH₄ yield; g/kg DMI) fitted with the variables selected from all the available data by the conventional method and machine learning methods. See the notes of Table 1.

the fewest variables (6 in total, Table 4) among the ML models. Thus, the DMI-B SCAD model was selected as the final DMI-B model.

We assessed the performance of the models by plotting the observed CH₄ emissions against the CH₄ emissions predicted using the final ANIM-B and the DMI-B models, along with the corresponding studentized residuals (Supplementary Figs. 2 and 3, respectively). Both the ANIM-B gmmLasso and the DMI-B SCAD models showed more tightly clustered and less scattered patterns in the observation vs. prediction plots, as compared to the ANIM-B and the DMI-B conventional models (Supplementary Figs. 2A,B and 3A,B, respectively). On the studentized residuals vs. predictions plots, the two ANIM-B and the DMI-B ML models also had tighter and more symmetric patterns around the regression lines that had smaller slopes compared to the ANIM-B and the DMI-B conventional models (Supplementary Figs. 2a,b and 3a,b, respectively).

Animal-based (g CH₄/animal-d) prediction models

The final ANIM-B gmmLasso model contained three animal-related predictor variables (DMI, A:P ratio, and BW) and 10 bacterial genera (or equivalents) of rumen bacteria out of the 308 genera (Table 5). None of the protozoal, fungal, or archaeal genera was selected as predictor variables. The three animal-related variables all had a positive coefficient. Of the 10 microbial predictor variables, four had a positive coefficient: 0.59 for the Uncl_genus of *Oscillospiraceae*, 0.49 for the Uncl_genus of the order *Clostridia*, 0.46 for the Uncl_Family of the order *Gastranaerophilales*, and 0.24 for the genus *Moryella*. The remaining six genera had a negative coefficient: Uncl_Family of the order *RF39* (–0.27), the genus *Prevotella_7* (–0.30), the Uncl_Genus of *Marinifilaceae* (–0.34), the genus *Syntrophococcus* (–0.37), the genus *Oribacterium* (–0.40), and the Uncl_Family of the order *Oscillospirales* (–0.44). The ANIM-B conventional model had the same three of the five animal-related data as predictor variables (DMI, A:P ratio, and BW). Compared to the ANIM-B conventional model, the ANIM-B gmmLasso model had a reduced $\hat{\sigma}_\alpha$ (1.59 vs. 2.04), RMSPE (by 3.8%), and MAE (by 4.6%), and increased CCC

	Predictor variables	gmmLasso ¹	Conventional ²
Animal	Intercept	21.9 (0.19)	21.9 (0.23)
	Body weight (kg)	1.17 (0.24)	1.48 (0.18)
	Dry matter intake (kg/d)	1.35 (0.24)	1.18(0.25)
	Acetate to propionate (A:P) ratio	1.17 (0.18)	1.29 (0.26)
Bacteria**	Uncl_Genus of <i>Oscillospiraceae</i>	0.59 (0.17)	
	Uncl_Genus of the order <i>Clostridia</i>	0.49 (0.16)	
	Uncl_Family of the order <i>Gastranaerophilales</i>	0.46 (0.17)	
	Genus <i>Moryella</i>	0.24 (0.15)	
	Uncl_Family of the order <i>RF39</i>	– 0.27 (0.16)	
	Genus <i>Prevotella_7</i>	– 0.30 (0.18)	
	Uncl_Genus of <i>Marinifilaceae</i>	– 0.34 (0.16)	
	Genus <i>Syntrophococcus</i>	– 0.37 (0.17)	
	Genus <i>Oribacterium</i>	– 0.40 (0.17)	
	Uncl_Family of <i>Oscillospirales</i>	– 0.44 (0.18)	
Number of predictors		13	3
$\hat{\sigma}_\alpha$		1.59	2.04

Table 5. Coefficients and the associated standard errors (SE) of the predictor variables of the final ANIM-B (CH₄ production; g/d) model. The bolded genera were shared between the ANIM-B and the DMI-B models. Uncl unclassified. For all the models, the predictor variables were centered and scaled to have a mean of 0 and variance of 1. **Log-transformed relative sequence abundance. ¹Developed using gmmLasso and all the available data after log transformation of the microbial data. ²Developed using conventional model and only animal-related data.

(by 9.49%). The ANIM-B glmLasso model also had a lower AIC and BIC than the ANIM-B conventional model (by 53 and 19, respectively).

Dry matter intake-based (g CH₄/kg of DMI) prediction model

The final DMI-B SCAD model had six predictor variables: one animal-related predictor variable (A:P ratio) and five bacterial genera as microbial predictor variables (Table 6). No protozoal, fungal, or archaeal genera were selected as predictor variables. The A:P ratio predictor variable had a positive coefficient. Except for *Uncl_Family* of the order *Gastranaerophilales* (with a coefficient of 0.50), the other four microbial predictor variables were all known genera, all of which had a negative coefficient: -0.24 for *Pseudoramibacter*, -0.33 for *Megasphaera*, -0.39 for *Selenomonas*, and -0.40 for *Oribacterium*. The DMI-B conventional model had two animal-related predictor variables (A:P ratio and BW), and both had a positive coefficient. Compared to the DMI-B conventional model, the DMI-B SCAD model had a reduced $\hat{\sigma}_\alpha$ (0.98 vs. 1.47), RMSPE (by 9.66%), and MAE (by 9.97%), and increased CCC (by 84.36%). Compared to the DMI-B conventional model, the DMI-B SCAD model reduced both AIC and BIC (by 15 and 5, respectively).

Discussion

The rumen microbiota is responsible for about 90% of the CH₄ emitted from sheep²⁷. With respect to CH₄ production, rumen microbes can be categorized as producers of methanogenesis substrates (primarily bacteria and protozoa), CH₄ producers (i.e., methanogens), and those that influence CH₄ production by interacting with the above two categories either positively (e.g., through mutualism and commensalism) or negatively (i.e., through amensalism, competition, and predation) interactions. Thus, some of the rumen microbes could have a quantitative relationship with the overall CH₄ emissions. Using four statistical/ML methods and one dataset (218 observations) that contains both animal-related data and metataxonomic data of the rumen microbiota (330 genera in total), we developed ML models for improved prediction of CH₄ production (g CH₄/animal-d) and CH₄ yield (g CH₄/kg of DMI) from sheep, which include rumen microbes as predictor variables.

New ML models including microbial predictor variables

As evaluated for prediction performance, it is evident that all the ML models, except for the ANIM-B SCAD model, improved prediction of CH₄ emissions in terms of error, bias, and accuracy when compared with the conventional models. Additionally, the improvement in prediction performance demonstrated in the evaluation with testing data randomly selected from the entire dataset suggests the potential usefulness of these models over a broad range of CH₄ emissions. The improvement in CH₄ prediction by the final ANIM-B glmLasso model and the final DMI-B SCAD model is also evident in the plots of the predictions vs. the observations along with the studentized residuals. In addition to the same animal-related predictor variables included in the conventional models (except for the exclusion of BW in the DMI-B SCAD model), the final ML included genera of bacteria as predictor variables. The improvement in prediction performance strongly suggests that the inclusion of these microbial predictor variables might have fine-tuned the models. Moreover, the improved predictions of the final ML models align well with the assumptions of linear mixed models. These findings suggest that microbial predictor variables can differentiate sheep based on CH₄ production levels, offering promising prospects for capturing variations in CH₄ emissions and helping achieve accurate CH₄ prediction.

The bacterial genera included in the ML models

Despite the dataset containing 330 genera of microbes, only a small number of genera were selected as microbial predictor variables, and their inclusion did not lead to model overfitting. Notably, all the microbial predictor

	Predictor variables	SCAD ¹	Conventional ²
Animal	Intercept	15.13 (0.13)	15.12 (0.17)
	Body weight (kg)	–	0.89 (0.13)
	Acetate to propionate (A:P) ratio	0.93 (0.12)	0.25 (0.16)
Bacteria**	Uncl_Family of the order <i>Gastranaerophilales</i>	0.50 (0.12)	
	Genus <i>Pseudoramibacter</i>	-0.24 (0.12)	
	Genus <i>Megasphaera</i>	-0.33 (0.12)	
	Genus <i>Selenomonas</i>	-0.39 (0.13)	
	Genus <i>Oribacterium</i>	-0.40 (0.12)	
Number of predictors		6	2
$\hat{\sigma}_\alpha$		0.98	1.47

Table 6. Coefficients and the associated standard errors (SE) of the predictor variables of the final DMI-B (CH₄ yield; g/kg DMI) model. The bolded genera were shared between the ANIM-B and the DMI-B models. *Uncl* unclassified. For all models, the predictor variables were centered and scaled to have a mean of 0 and variance of 1. **Log-transformed relative sequence abundance. ¹Developed using SCAD and all the available data after log-transformation of the microbial data. ²Developed using conventional model and only animal-related data except DMI.

variables in both the ANIM-B and the DMI-B prediction models are genera of bacteria, with the majority belonging to *Firmicutes*, an abundant phylum of rumen microbiota. Interestingly, no archaea, protozoa, or fungi were selected as predictor variables in any of the ML models. The absence of methanogen genera in the models might seem counterintuitive, given their role as CH₄ producers. However, previous studies have shown that the abundance of rumen methanogens, even that of the most abundant genus, *Methanobrevibacter*²⁸, has very a weak to no correlation with CH₄ production^{12,29}. Furthermore, changes in diets that significantly affected CH₄ production were not found to lead to substantial alterations in the relative sequence abundance of *Methanobrevibacter*³⁰. Thus, the lack of methanogens in the final prediction models is not surprising. At low abundance, methanogens, fungi, and protozoa might have little predicting power for CH₄ emissions.

Both the ANIM-B and DMI-B ML models included the genus *Oribacterium* and one unclassified genus of the order *Gastranaerophilales* (the Uncl_Family of the order *Gastranaerophilales*) with a negative and a positive coefficient, respectively. Although the function of this unclassified genus is not known, its large coefficients in both prediction models suggest a high predicting power. Interestingly, *Gastranaerophilales* has been shown to be negatively associated with milk yield in dairy cattle³¹ and negatively associated with growth in lambs³². Regarding the genus *Oribacterium*, it had a negative coefficient in both the final ANIM-B and the DMI-B models. This aligns with *Oribacterium* being a potential H₂ sink in the rumen of sheep³³ and its negative association with CH₄ emissions documented in dairy cows³⁴. Future research is warranted to explore and verify if *Oribacterium* has a quantitative relationship with rumen CH₄ production and can be used as a predictor variable or biomarker of CH₄ emissions.

In addition to the two genera shared with the DMI-B SCAD model, the ANIM-B glmLasso model included another eight genera, five of which were unclassified, and their functions remain unknown. Among these genera, *Moryella*, *Prevotella_7*, and *Syntrophococcus* are known genera of bacteria, but their association or correlation with CH₄ emissions in ruminants has not been reported in the literature. *Prevotella_7* is a newly established genus³⁵, consisting of *P. multiformis*, *P. multisaccharivorax*, and uncultured bacteria from the oral cavities and the rumen (<https://www.arb-silva.de/>). Studies have shown that both *P. multiformis* and *P. multisaccharivorax* were enriched by concentrate-based diets correlated with decreased CH₄ emissions^{36,37}. In buffalos, a higher relative sequence abundance of *Prevotella* (closely related to *Prevotella_7*) was associated with low CH₄ emissions³⁸. *Prevotella* is considered to be hydrogen-consuming³⁹, making the inclusion of *Prevotella_7* as a predictor variable with a negative coefficient not surprising. As for *Moryella* and *Syntrophococcus*, the literature has no information about their relationship with CH₄ emissions, except for one in vitro study that showed a positive correlation between *Syntrophococcus* and maximum CH₄ production, but a negative correlation with the appearance of peak CH₄ production⁴⁰. Future quantitative studies are needed to verify the qualitative relationship between these two genera and CH₄ emissions.

The DMI-B SCAD model did not include BW as a predictor variable, but the inclusion of bacterial predictor variables improved the prediction accuracy. This suggests that these bacterial predictor variables might have reduced the variation in CH₄ prediction introduced by BW. In addition to the two unknown genera discussed earlier, the final DMI-B SCAD model included three known genera (*Pseudoramibacter*, *Megasphaera*, and *Selenomonas*), which were not included in the final ANIM-B glmLasso model. While the literature has no information about any potential association between *Pseudoramibacter* and CH₄ emissions, *Pseudoramibacter* was reported to be positively correlated with feed efficiency in sheep⁴¹. Similarly, it remains unknown if *Megasphaera* or *Selenomonas* has any association or correlation with CH₄ emissions in the absence of anti-methane inhibitors. However, one in vitro study has shown that calcium salts of long-chain fatty acids⁴², tucuma oil⁴³, and ginkgo extract⁴⁴ decreased CH₄ emissions while increasing propionate and the abundance of *Megasphaera* and *Selenomonas*. Encapsulated nitrate has also been found to reduce CH₄ emissions while increasing *Selenomonas* in steers⁴⁵. Additionally, *Megasphaera* was more abundant in low than in high CH₄-yield sheep⁴⁶. Thus, the negative coefficients of both *Megasphaera* and *Selenomonas* in the DMI-B SCAD model align with their negative association with CH₄ emissions and their ability to consume hydrogen and produce propionate^{33,39,47,48}. It should be noted that many of these correlations were observed in ruminants that were subjected to dietary interventions using anti-methane inhibitors, which could directly or indirectly affect these genera, skewing their relationships with CH₄ production. Future research is warranted to determine their relationship with CH₄ emissions in the absence of anti-methane inhibitors and their predicting power of CH₄ emissions.

Potential limitations of the sheep dataset and the selected models

The dataset used in developing the prediction models was from one study using a relatively small number of sheep that were fed the same diet¹³. A larger database from multiple studies using different diets would improve prediction models and enhance their applicability across a broad range of feeding systems. However, currently, no such a database exists. A few studies in the literature have reported rumen microbiota data together with CH₄ emission data, and we attempted to combine these data into a larger dataset. However, most of the studies used anti-methane inhibitors, likely skewing the microbiota data. Additionally, different studies sequenced different regions of the 16S rRNA or 18S rRNA gene, making it impossible to combine the sequence data for microbiota analysis. To facilitate the combination of sequence data from multiple studies, a standardized metataxonomic protocol should be adopted.

Metataxonomic analysis of the rumen microbiota incurs additional cost, which may hinder the adoption and application in the livestock industry, but it is relatively small and justifiable for several reasons. First, amplicon sequences are routinely generated and analyzed in numerous studies to understand the microbiological underpinning of treatment effects or group differences, making sequencing cost a part of the study cost. Second, the cost of amplicon sequencing has been steadily decreasing, and with the advent of new sequencing technologies (e.g., the Illumina NovaSeq 6000), it is expected to further decrease. Third, accurate and precise CH₄ emission

prediction is becoming increasingly important for precision livestock farming, which is an essential approach to increase feed efficiency while reducing CH₄ emissions for a sustainable industry of ruminants. Improved prediction of CH₄ emissions can assist researchers and farmers in improving the overall feed efficiency and sustainability of livestock production.

Conclusion

Using machine learning methods, we successfully developed animal-based and dry matter intake-based CH₄ prediction models that include both animal data and microbial genera as predictor variables. The inclusion of microbial predictor variables improved CH₄ emission prediction in terms of error, bias, and accuracy compared to the conventional models that solely contain animal-related data, particularly for predicting CH₄ yield (g CH₄/kg of DMI). Notably, the coefficients (positive or negative) of some of the predictor variables from known bacterial genera (e.g., *Oribacterium*, *Prevotella_7*, *Megasphaera*, and *Selenomonas*) align well with their known associations or correlations (positive or negative) with CH₄ emissions and/or with their known functions in the rumen ecosystem, such as hydrogen consumption. However, the role and association of other predictor variables of known genera (e.g., *Moryella*, *Pseudoramibacter*, and *Syntrophococcus*), as well as the unclassified genera, with CH₄ emissions remain unknown.

While further improvements in CH₄ prediction models would benefit from extensive databases encompassing rumen bacteria, archaea, fungi, and protozoa, along with animal-related data from multiple studies, this study demonstrated the potential and feasibility to include rumen microbes as predictor variables to improve CH₄ predictions from sheep, likely from cattle also. A standardized approach to generate and analyze rumen microbial data is imperative to facilitate the compilation of multiple datasets from numerous studies. Such efforts will contribute to a better understanding of the rumen microbiota's impact on CH₄ emissions and improved prediction models for sustainable agricultural practices.

Data availability

The sequence and animal data used in this study were published by Kittelmann, et al.¹³ and can be downloaded from the EMBL database (accession numbers: ERP003779 for Bacteria, ERP003773 for Archaea, ERP003772 for ciliate protozoa, and ERP003764 for anaerobic fungi). The R codes used in this study are available at https://github.com/yu2269/CH4_prediction_model.

Received: 20 December 2022; Accepted: 27 November 2023

Published online: 02 December 2023

References

1. Masson-Delmotte, V. et al. Climate Change 2021: The Physical Science Basis. *Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change* (2021).
2. Johnson, K. A. & Johnson, D. E. Methane emissions from cattle. *J. Anim. Sci.* **73**, 2483–2492. <https://doi.org/10.2527/1995.7382483x> (1995).
3. Pelchen, A. & Peters, K. J. Methane emissions from sheep. *Small Rumin. Res.* **27**, 137–150 (1998).
4. Hammond, K. J. et al. Review of current in vivo measurement techniques for quantifying enteric methane emission from ruminants. *Anim. Feed Sci. Technol.* **219**, 13–30. <https://doi.org/10.1016/j.anifeeds.2016.05.018> (2016).
5. Zhao, Y. G., O'Connell, N. E. & Yan, T. Prediction of enteric methane emissions from sheep offered fresh perennial ryegrass (*Lolium perenne*) using data measured in indirect open-circuit respiration chambers. *J. Anim. Sci.* **94**, 2425–2435. <https://doi.org/10.2527/jas.2016-0334> (2016).
6. van Lingen, H. J. et al. Prediction of enteric methane production, yield and intensity of beef cattle using an intercontinental database. *Agric. Ecosyst. Environ.* **283**, 106575. <https://doi.org/10.1016/j.agee.2019.106575> (2019).
7. Niu, M. et al. Prediction of enteric methane production, yield, and intensity in dairy cattle using an intercontinental database. *Glob. Chang. Biol.* **24**, 3368–3389. <https://doi.org/10.1111/gcb.14094> (2018).
8. Appuhamy, J. A., France, J. & Kebreab, E. Models for predicting enteric methane emissions from dairy cows in North America, Europe, and Australia and New Zealand. *Glob. Chang. Biol.* **22**, 3039–3056. <https://doi.org/10.1111/gcb.13339> (2016).
9. Hristov, A. N. et al. Symposium review: Uncertainties in enteric methane inventories, measurement techniques, and prediction models. *J. Dairy Sci.* **101**, 6655–6674. <https://doi.org/10.3168/jds.2017-13536> (2018).
10. Patra, A. K. & Yu, Z. Combinations of nitrate, saponin, and sulfate additively reduce methane production by rumen cultures in vitro while not adversely affecting feed digestion, fermentation or microbial communities. *Bioresour. Technol.* **155**, 129–135. <https://doi.org/10.1016/j.biortech.2013.12.099> (2014).
11. Newbold, C. J. & Ramos-Morales, E. Review: Ruminal microbiome and microbial metabolome: Effects of diet and ruminant host. *Animal* **14**, s78–s86. <https://doi.org/10.1017/S1751731119003252> (2020).
12. Tapio, I., Snelling, T. J., Strozzi, F. & Wallace, R. J. The ruminal microbiome associated with methane emissions from ruminant livestock. *J. Anim. Sci. Biotechnol.* **8**, 7. <https://doi.org/10.1186/s40104-017-0141-0> (2017).
13. Kittelmann, S. et al. Two different bacterial community types are linked with the low-methane emission trait in sheep. *PLoS ONE* **9**, e103171. <https://doi.org/10.1371/journal.pone.0103171> (2014).
14. Pinares-Patino, C. S. et al. Heritability estimates of methane emissions from sheep. *Animal* **7**(Suppl 2), 316–321. <https://doi.org/10.1017/S1751731113000864> (2013).
15. Callahan, B. J. et al. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat. Methods* **13**, 581–583. <https://doi.org/10.1038/nmeth.3869> (2016).
16. Quast, C. et al. The SILVA ribosomal RNA gene database project: Improved data processing and web-based tools. *Nucleic Acids Res.* **41**, D590–596. <https://doi.org/10.1093/nar/gks1219> (2013).
17. Nilsson, R. H. et al. The UNITE database for molecular identification of fungi: Handling dark taxa and parallel taxonomic classifications. *Nucleic Acids Res.* **47**, D259–D264. <https://doi.org/10.1093/nar/gky1022> (2019).
18. Greg, R. *gbm: Generalized Boosted Regression Models v. 2.1.8.1* (2010).
19. Friedman, J., Hastie, T. & Tibshirani, R. Regularization paths for generalized linear models via coordinate descent. *J. Stat. Softw.* **33**, 1–22. <https://doi.org/10.18637/jss.v033.i01> (2010).
20. Groll, A. & Tutz, G. Variable selection for generalized linear mixed models by L1-penalized estimation. *Stat. Comput.* **24**, 137–154. <https://doi.org/10.1007/s11222-012-9359-z> (2012).

21. Yang, L. *Model-Based Clustering of Longitudinal Data in High Dimensions* Thesis (Ph.D.) thesis (University of Rochester, 2021).
22. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48. <https://doi.org/10.18637/jss.v067.i01> (2015).
23. Team, R. C. R. *A Language and Environment for Statistical Computing (v. 4.0.2) [Computer Software]* (R Foundation for Statistical Computing, 2020).
24. Saldana, D. F. & Feng, Y. SIS: An R package for sure independence screening in ultrahigh-dimensional statistical models. *J. Stat. Softw.* <https://doi.org/10.18637/jss.v083.i02> (2018).
25. James, G., Witten, D., Hastie, T., Tibshirani, R. & Taylor, J. *An Introduction to Statistical Learning with Applications in R* 2nd edn, 445 (Springer, 2021).
26. Stevenson, M. *et al.* *EpiR: An R Package for the Analysis of Epidemiological Data v. 2.0.62* (2013).
27. Murray, R. M., Bryant, A. M. & Leng, R. A. Rates of production of methane in the rumen and large intestine of sheep. *Br. J. Nutr.* **36**, 1–14. <https://doi.org/10.1079/bjn19760053> (1976).
28. Zhou, M., Hernandez-Sanabria, E. & Guan, L. L. Characterization of variation in rumen methanogenic communities under different dietary and host feed efficiency conditions, as determined by PCR-denaturing gradient gel electrophoresis analysis. *Appl. Environ. Microbiol.* **76**, 3776–3786. <https://doi.org/10.1128/AEM.00010-10> (2010).
29. Cunha, C. S. *et al.* Compositional and structural dynamics of the ruminal microbiota in dairy heifers and its relationship to methane production. *J. Sci. Food Agric.* **99**, 210–218. <https://doi.org/10.1002/jsfa.9162> (2019).
30. van Gastelen, S. *et al.* Linseed oil and DGAT1 K232A polymorphism: Effects on methane emission, energy and nitrogen metabolism, lactation performance, ruminal fermentation, and rumen microbial composition of Holstein-Friesian cows. *J. Dairy Sci.* **100**, 8939–8957. <https://doi.org/10.3168/jds.2016-12367> (2017).
31. Bach, A. *et al.* Changes in the rumen and colon microbiota and effects of live yeast dietary supplementation during the transition from the dry period to lactation of dairy cows. *J. Dairy Sci.* **102**, 6180–6198. <https://doi.org/10.3168/jds.2018-16105> (2019).
32. Perea, K. *et al.* Feed efficiency phenotypes in lambs involve changes in ruminal, colonic, and small-intestine-located microbiota. *J. Anim. Sci.* **95**, 2585–2592. <https://doi.org/10.2527/jas.2016.1222> (2017).
33. Greening, C. *et al.* Diverse hydrogen production and consumption pathways influence methane production in ruminants. *ISME J.* **13**, 2617–2632. <https://doi.org/10.1038/s41396-019-0464-2> (2019).
34. Lopez-Garcia, A. *et al.* Fungal and ciliate protozoa are the main rumen microbes associated with methane emissions in dairy cattle. *Gigascience* **11**, 88. <https://doi.org/10.1093/gigascience/giab088> (2022).
35. Henderson, G. *et al.* Improved taxonomic assignment of rumen bacterial 16S rRNA sequences using a revised SILVA taxonomic framework. *PeerJ* **7**, e6496. <https://doi.org/10.7717/peerj.6496> (2019).
36. Ellison, M. J. *et al.* Diet and feed efficiency status affect rumen microbial profiles of sheep. *Small Rumin. Res.* **156**, 12–19. <https://doi.org/10.1016/j.smallrumres.2017.08.009> (2017).
37. Hristov, A. N. *et al.* Special topics—mitigation of methane and nitrous oxide emissions from animal operations: I. A review of enteric methane mitigation options. *J. Anim. Sci.* **91**, 5045–5069. <https://doi.org/10.2527/jas.2013-6583> (2013).
38. Aguilar-Marin, S. B., Betancur-Murillo, C. L., Isaza, G. A., Mesa, H. & Jovel, J. Lower methane emissions were associated with higher abundance of ruminal Prevotella in a cohort of Colombian buffalos. *BMC Microbiol.* **20**, 364. <https://doi.org/10.1186/s12866-020-02037-6> (2020).
39. Denman, S. E., Martinez Fernandez, G., Shinkai, T., Mitsumori, M. & McSweeney, C. S. Metagenomic analysis of the rumen microbial community following inhibition of methane formation by a halogenated methane analog. *Front. Microbiol.* **6**, 1087. <https://doi.org/10.3389/fmicb.2015.01087> (2015).
40. Li, R. *et al.* Effect of different forage-to-concentrate ratios on ruminal bacterial structure and real-time methane production in sheep. *PLoS ONE* **14**, e0214777. <https://doi.org/10.1371/journal.pone.0214777> (2019).
41. McLoughlin, S. *et al.* Rumen microbiome composition is altered in sheep divergent in feed efficiency. *Front. Microbiol.* **11**, 1981. <https://doi.org/10.3389/fmicb.2020.01981> (2020).
42. Sato, Y. *et al.* Calcium salts of long-chain fatty acids from linseed oil decrease methane production by altering the rumen microbiome in vitro. *PLoS ONE* **15**, e0242158. <https://doi.org/10.1371/journal.pone.0242158> (2020).
43. Ramos, A. F. O. *et al.* Tucuma oil shifted ruminal fermentation, reducing methane production and altering the microbiome but decreased substrate digestibility within a RUSITEC fed a mixed hay–concentrate diet. *Front. Microbiol.* **9**, 1647. <https://doi.org/10.3389/fmicb.2018.01647> (2018).
44. Oh, S., Koike, S. & Kobayashi, Y. Effect of ginkgo extract supplementation on in vitro rumen fermentation and bacterial profiles under different dietary conditions. *Anim. Sci. J.* **88**, 1737–1743. <https://doi.org/10.1111/asj.12877> (2017).
45. Granja-Salcedo, Y. T. *et al.* Long-term encapsulated nitrate supplementation modulates rumen microbial diversity and rumen fermentation to reduce methane emission in grazing steers. *Front. Microbiol.* **10**, 614. <https://doi.org/10.3389/fmicb.2019.00614> (2019).
46. Kamke, J. *et al.* Rumen metagenome and metatranscriptome analyses of low methane yield sheep reveals a Sharpea-enriched microbiome characterised by lactic acid formation and utilisation. *Microbiome* **4**, 56. <https://doi.org/10.1186/s40168-016-0201-2> (2016).
47. Chen, L. *et al.* *Megasphaera elsdenii* lactate degradation pattern shifts in rumen acidosis models. *Front. Microbiol.* **10**, 162. <https://doi.org/10.3389/fmicb.2019.00162> (2019).
48. Marchandin, H., Juvonen, R. & Haikara, A. *Megasphaera*. In *Bergey's Manual of Systematics of Archaea and Bacteria* (eds Trujillo, M. E. *et al.*) 1–16 (Wiley, 2015).

Acknowledgements

This study is part of the Joint Programming Initiative on Agriculture, Food Security and Climate Change (FACCE-JPI)'s “GLOBAL NETWORK” project and the “Feeding and Nutrition Network” (<http://animalscience.psu.edu/fnn>) of the Livestock Research Group within the Global Research Alliance for Agricultural Greenhouse Gases (www.globalresearchalliance.org).

Author contributions

Z.Y. conceived and designed the study. P.J. provided the data. Z.Y., B.Z., and L.M. set up the database. B.Z. did the modeling. L.M. and Z.Y. supervise the study. B.Z. wrote the manuscript with input from all co-authors. All authors read and approved the final manuscript.

Funding

The authors gratefully acknowledge funding for this project from the USDA National Institute of Food and Agriculture (Award number: 2014-67003-21979). The animal and microbial data originated from a study funded by the Pastoral Greenhouse Gas Research Consortium (www.pggrc.co.nz).

Competing interests

The authors declare no competing interests.

Additional information

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1038/s41598-023-48449-y>.

Correspondence and requests for materials should be addressed to S.L. or Z.Y.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023