

*Anthrenus (Anthrenus) quernerii*  
(Coleoptera: Dermestidae: Megatominae),  
a new species from Austria

Article

Published Version

Creative Commons: Attribution-Noncommercial 3.0

Open Access

Holloway, G. J. ORCID: <https://orcid.org/0000-0003-0495-0313>  
(2024) *Anthrenus (Anthrenus) quernerii* (Coleoptera:  
Dermestidae: Megatominae), a new species from Austria.  
*Insecta Mundi*. 1060. ISSN 1942-1354 Available at  
<https://centaur.reading.ac.uk/116976/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Published version at: <https://journals.flvc.org/mundi/article/view/136792>

Publisher: Center for Systematic Entomology, Inc.

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

A journal of world insect systematics

# INSECTA MUNDI

---

---

1060

*Anthrenus (Anthrenus) quernerii* (Coleoptera:  
Dermestidae: Megatominae), a new species from Austria

Graham J. Holloway

Cole Museum of Zoology, School of Biological Sciences, Health and Life Sciences Building,  
The University of Reading, Whiteknights, Reading, Berkshire, RG6 6EX, UK

Date of issue: June 28, 2024

Center for Systematic Entomology, Inc., Gainesville, FL

**Holloway GJ. 2024.** *Anthrenus (Anthrenus) quernerii* (Coleoptera: Dermestidae: Megatominae), a new species from Austria. *Insecta Mundi* 1060: 1–6.

Published on June 28, 2024 by  
**Center for Systematic Entomology, Inc.**  
P.O. Box 141874  
Gainesville, FL 32614-1874 USA  
<http://centerforsystematicentomology.org/>

**INSECTA MUNDI** is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

*Insecta Mundi* is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

**Chief Editor:** David Plotkin, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Assistant Editor:** Paul E. Skelley, [insectamundi@gmail.com](mailto:insectamundi@gmail.com)

**Layout Editor:** Robert G. Forsyth

**Editorial Board:** Davide Dal Pos, M. J. Paulsen, Felipe Soto-Adames

**Founding Editors:** Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

**Review Editors:** Listed on the *Insecta Mundi* webpage

**Printed copies (ISSN 0749-6737) annually deposited in libraries**

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA

The Natural History Museum, London, UK

National Museum of Natural History, Smithsonian Institution, Washington, DC, USA

Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

**Electronic copies (online ISSN 1942-1354) in PDF format**

Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.  
<https://creativecommons.org/licenses/by-nc/3.0/>

# *Anthrenus (Anthrenus) quernerii* (Coleoptera: Dermestidae: Megatominae), a new species from Austria

Graham J. Holloway

Cole Museum of Zoology, School of Biological Sciences, Health and Life Sciences Building,  
The University of Reading, Whiteknights, Reading, Berkshire, RG6 6EX, UK  
g.j.holloway@reading.ac.uk

🌐 <https://orcid.org/0000-0003-0495-0313>

**Abstract.** A new species, *Anthrenus (Anthrenus) quernerii* Holloway (Coleoptera: Dermestidae: Megatominae) from Austria is described and compared with *A. pimpinellae* (Fabricius, 1775), the most similar species. Images of the habitus, ventrites, antenna, aedeagus, and sternite IX are presented. *Anthrenus quernerii* represents the 26<sup>th</sup> species belonging to the Palaearctic *Anthrenus pimpinellae* complex.

**Key words.** Carpet beetle, *Anthrenus amandae*, *Anthrenus chikatunovi*, aedeagus, median lobe, sternite, taxonomy, distribution.

**ZooBank registration.** urn:lsid:zoobank.org:pub:EF3828C0-DDD2-4619-9F86-BC8626522465

## Introduction

The Dermestidae contain over 1900 named species (Háva 2024). One of the larger genera within the family is *Anthrenus* Geoffroy, 1762 with nearly 300 species, including the Palaearctic *Anthrenus pimpinellae* (Fabricius, 1775) complex. All species within the complex have a fascia of white scales across the elytra, although the fascia is reduced in some species, e.g. *A. pfefferi* Kalík, 1954 and *A. mroczkowskii* Kalík, 1954. Some species are sufficiently alike to cause confusion (Háva and Herrmann 2019, 2020), which has likely constrained the rate of discovery of new species. Around the middle of the 20<sup>th</sup> century, only about nine species had been described belonging to the complex (see data in Háva 2024). Since the beginning of the 21<sup>st</sup> century, the number of known species has increased to 25 (Holloway and Herrmann 2024). This number will undoubtedly continue to increase, largely because species recognition within the complex is now predicated on the study of genitalia, principally male genitalia (Kadej et al. 2007; Holloway 2019, 2020, 2021; Holloway and Herrmann 2024; Holloway et al. 2024).

Many historic houses and museums in Austria are routinely surveyed for pest insect species (such as Dermestidae) as part of integrated pest management programs. During the spring survey period of 2022, a specimen was trapped and misidentified as *A. pimpinellae pimpinellae*. It is now recognized as a new species belonging to the *A. pimpinellae* complex and is here described as *Anthrenus* (s. str.) *quernerii*.

## Materials and Methods

In many Austrian museums and historic houses, sticky (blunder) traps are deployed to capture pest beetles and other species at ground level. About 5000 traps are regularly set across 65 buildings. Traps are renewed every six months, February or March and October or November. During the spring and summer, the traps are checked every 4 to 8 weeks and replaced if full of insects or dust. In the historic Schönbrunn Palace, all traps were set on 24<sup>th</sup> February 2022 and in the ‘Millions Room’ (sticky trap number 63) a single, slightly damaged *A. pimpinellae* complex individual was captured between 3<sup>rd</sup> May and 2<sup>nd</sup> June. The specimen was lifted from the trap glue using ethyl acetate. The specimen was clean and needed no further chemical treatment.

The Austrian specimen is held in the Natural History Museum, London (NHML). Twenty-nine *Anthrenus pimpinellae* specimens from across Europe were examined (NHML) along with 19 Greek *A. pimpinellae* held in the author’s collection. All specimens were macerated in a solution of 2% acetic acid for five days to soften prior to dissection. Dissection was carried out under a Brunel BMSL zoom stereo LED microscope and

involved detaching the abdomen from the rest of the insect using two entomological pins. The soft tergites were then peeled away from the harder ventrites to expose the genitalia. The aedeagus was detached from the ring sclerite, and sternite IX was detached from the ring sclerite and the aedeagus. Habitus and ventrite images were captured at  $\times 20$  magnification using a Canon EOS 2000D camera mounted on the BMSL microscope. Images of aedeagus and sternite IX were captured at  $\times 200$  magnification for measurement using a Canon EOS 1300D camera mounted on a Brunel monocular SP28 microscope. After dissection, all body parts were mounted on card. The antennae were imaged at  $\times 200$  magnification through the SP28 microscope. All images were fed through Helicon Focus Pro version 8.2.2 focus-stacking software. All measurements were made using DsCap.Ink software version 3.90. Measurements taken:

- Body length (BL): distance from anterior margin of pronotum to the apex of the elytra
- Body width (BW): maximum distance across the single elytron  $\times 2$
- Antennal club length (AL): length of the last three antennomeres
- Antennal club width (AW): maximum width across the terminal antennomere
- Paramere length (PL): distance from the anterior end of the paramere to the apex of the paramere
- Sternite IX length (SL): distance from the tip of one anterior horn to the tip of the posterior lobe.

Scale bars were added using ImageJ 1.53M (Schneider et al. 2012). The distribution map was generated using SimpleMappr (Shorthouse 2010).

## Results


### *Anthrenus (Anthrenus) quernerii* Holloway, new species

Figures 1–3

**Type specimens. Holotype male.** Schönbrunn Palace, Vienna, Austria (48.183, 16.307), 3<sup>rd</sup> May 2022 to 2<sup>nd</sup> June 2022, Pascal Querner leg. NHML.

**Paratypes.** None.

**Description, external characteristics.** Holotype *Anthrenus (Anthrenus) quernerii* (Fig 1A) (BL = 3.07 mm, BW = 2.26 mm) with a single pale brown ocellus on face well below the level of the top of the eyes. A large notch in the lower inner margin of each eye. Integument very dark brown to black. Head covered in black scales apart from small patch of white scales on vertex in center, and patches of brown scales scattered along the inner eye margins,



**Figure 1.** *Anthrenus quernerii*. **A)** Habitus (left elytron missing, scale bar = 1 mm). **B)** Abdominal ventrites (scale bar = 1 mm). **C)** Detached antenna (scale bar = 100  $\mu$ m).

especially in the notch. Pronotum covered in white and orange scales, especially on outer thirds, with patches of black scales mostly on inner third. Only the right elytron remains on the specimen. Prominent white fascia crosses the elytron extending down the elytral suture from the scutellum for about  $\frac{2}{5}$  of the length of the elytron, extending across the elytron as an expanding band to cover  $\frac{1}{2}$  of the length of the elytron at the margin. White fascia almost entirely an unbroken block of white scales apart from small patches of black scales sub-sutural and sub marginal,  $\frac{1}{3}$  down elytron in both cases, and a few black scales at the lower marginal edge of the white fascia.

Ventrites (Fig. 1B) covered in white tinted brown scales. Large patches of black scales at margins of each ventrite, largest on ventrite I which merges with the black marginal patch on ventrite II. Ventrite V mostly covered in black scales, pale scales only reaching the posterior margin at two points with just a few scales at the anterior edge of the marginal black spots on ventrite V. A small number of pale brown scales mixed in with the pale scales on ventrite V.

Eleven segmented antenna (Fig. 1C) (AL = 194  $\mu\text{m}$ , AW = 152  $\mu\text{m}$ ) has a broad, symmetrical rectangular club consisting of the terminal three antennomeres. The antenna and legs are dark red. Anterior faces of femora covered in white, with pale brown scales proximally and distally, and mostly black scales medially.

**Description, internal characteristics.** Aedeagus (Fig. 2A) has broad parameres (PL = 373  $\mu\text{m}$ ). Parameres have rounded lateral margins, curving into relatively sharp apices that angle in towards each other. The apical  $\frac{2}{5}$  of the inner margins are almost parallel where they narrow sharply and continue to the anterior end as relatively narrow curved rods. The surface of the broad, paddle shaped terminal  $\frac{2}{5}$  of each paramere is covered in long, narrow setae that point inwards towards the median lobe. Posterior tips of parameres pale. Median lobe very broad at base that narrows gradually to blunt, rounded, slightly expanded tip, and is slightly shorter than the parameres.

Sternite IX (Fig. 2B, SL = 473  $\mu\text{m}$ ) with broad posterior lobe. The apex is clear laterally, the rest of the sternite is pale brown. The medial apex of the posterior lobe is a smooth, shallow curve. The clear lateral components of the apex are covered in setae, longest towards the lateral corners and absent along the apical margin. The setae



**Figure 2.** *Anthrenus quernerii*. A) Aedeagus dorsal aspect. B) Sternite IX. Scale bar = 100  $\mu\text{m}$  in both cases.




**Figure 3.** Location of collection of *Anthrenus quernerii* holotype.

extend down the lateral margins becoming progressively shorter to the mid-point where the margins narrow to form a neck. From the neck, the margins diverge and continue as smooth curves to the anterior, ending in two narrow, curved horns.

**Distribution.** Only known from the holotype collected in the Schönbrunn Palace, Vienna, Austria. (Fig. 3).

**Etymology.** *Anthrenus* (*Anthrenus*) *quernerii* is named after the collector, Pascal Querner, who is the IPM manager for the Museum of Natural History, Vienna.

**Differential diagnosis.** Initially, *A. quernerii* was misidentified as *A. pimpinellae* (Fig. 4A). *Anthrenus pimpinellae*



**Figure 4.** *Anthrenus pimpinellae*. A) Habitus (scale bar = 1 mm). B) Abdominal ventrites (scale bar = 1 mm). C) Antenna (scale bar = 100  $\mu$ m).



white fascia is much narrower than *A. quernerii* and overall *A. pimpinellae* has a narrower profile than *A. quernerii*. Holloway and Bakaloudis (2020) provide a BW/BL value of  $0.688 \pm 0.029$  for *A. pimpinellae*, and BW/BL for *A. quernerii* holotype is 0.736, indicating that *A. quernerii* is broader than an average *A. pimpinellae*. It is not obvious that the ventrites (Fig. 4B) and the antennal structure (Fig. 4C) differ between *A. quernerii* and *A. pimpinellae*, although perhaps *A. quernerii* ventrites are covered in cleaner white scales than *A. pimpinellae*.

*Anthrenus pimpinellae* parameres (Fig. 5A) have more strongly curved lateral margins than *A. quernerii*. On the inner margin, the paramere paddles are longer than *A. quernerii*, extending more than half the length of the paramere, whereas the paddles of *A. quernerii* are considerably less than half the paramere length. *Anthrenus pimpinellae* median lobe is broad at the base but narrows more sharply than *A. quernerii* terminating in a long, slim apex, much slimmer than *A. quernerii*.

*Anthrenus pimpinellae* sternite IX (Fig. 5B) is almost entirely pale brown with only the very outer apical edges of the posterior lobe being clear. All setae are longer than *A. quernerii* and clustered along the margins rather than spreading into the posterior lobe disc as in *A. quernerii*.

## Discussion

The number of known species within the Palearctic *A. pimpinellae* complex continues to increase. Holloway and Herrmann (2024) list 25 species, this study increases the number to 26. This is the third *A. pimpinellae* lookalike species described from Europe, the others being *A. amandae* Holloway, 2019 from Mallorca and *A. chikatunovi* Holloway, 2020 from the Pyrennees. As well as similar color patterns, all also have quadrate antennal clubs. Insect



**Figure 5.** *Anthrenus pimpinellae*. A) Aedeagus dorsal aspect. B) Sternite IX. Scale bar = 100  $\mu$ m in both cases.

species in Europe are extensively studied so it does beg the question, how many more cryptic *A. pimpinellae* complex species remain to be discovered, particularly from less studied parts of the Palaearctic.

*Anthrenus quernerii* is compared here with a typically patterned *A. pimpinellae*. It should be borne in mind that a small proportion of *A. pimpinellae* specimens from Greece have broader white bands than shown in Fig. 4A, so the width of the white band might not always be a reliable identification cue. Only a single *A. quernerii* is currently known. More specimens need to be found and studied to discover more and better ways to distinguish between *A. quernerii* and *A. pimpinellae* using external features alone.

## Acknowledgments

The author is very grateful to Andreas Herrmann, Ivan Cañada Luna, and Dr. Paul Skelley for reviewing the manuscript so efficiently, and to Dr. David Plotkin for his effective editorial work. Finally, many thanks to Professor D.E. Bakaloudis for collecting examples of *A. pimpinellae* from Greece and making them available for study.

## Literature Cited

- Háva J. 2024.** Dermestidae World (Coleoptera). Available at <http://dermestidae.wz.cz/world-dermestidae> (Last accessed 22<sup>nd</sup> February 2024.)
- Háva J, Herrmann A. 2019.** New faunistic records and remarks on Dermestidae (Coleoptera) – Part 19. *Natura Somogyiensis* 33: 27–36.
- Háva J, Herrmann A. 2020.** New faunistic records and remarks on Dermestidae (Coleoptera) – Part 20. *Natura Somogyiensis* 34: 9–20.
- Holloway GJ. 2019.** *Anthrenus* (s. str.) *amandae* (Coleoptera: Dermestidae): a new species from Mallorca, Spain. *Zootaxa* 4543(4): 595–599. <http://doi.org/10.11646/zootaxa.4543.4.9>
- Holloway GJ. 2020.** *Anthrenus* (s. str.) *chikatunovi* (Coleoptera: Dermestidae): a new species from southern France. *Israel Journal of Entomology* 50: 69–75. <http://doi.org/10.5281/zenodo.4088743>
- Holloway GJ. 2021.** *Anthrenus* (s. str.) *corona* (Coleoptera, Dermestidae, Anthrenini): a new species in the *A. pimpinellae* (Fabricius, 1775) complex from Turkey. *Zootaxa* 4991(3): 555–560. <https://doi.org/10.11646/zootaxa.4991.3.7>
- Holloway GJ, Bakaloudis DE. 2020.** A comparative morphological study of *Anthrenus pimpinellae* (Fabricius, 1775) and *Anthrenus amandae* Holloway, 2019 (Coleoptera: Dermestidae). *The Coleopterists Bulletin* 74(2): 315–321. <https://doi.org/10.1649/0010-065X-74.2.315>
- Holloway GJ, Herrmann A. 2024 (in press).** A new species of the genus *Anthrenus* Geoffroy, 1762 (Coleoptera: Dermestidae) from Turkey and Lebanon. *Annales Zoologici*
- Holloway GJ, Thanasoulis G, Herrmann A. 2024.** A new species *Anthrenus bakaloudisi* sp. nov. (Coleoptera, Dermestidae, Megatominae) from Macedonia, Greece and comparison with *Anthrenus pfefferi* Kalik, 1954, *Anthrenus delicatus* Kiesenwetter, 1851, and *Anthrenus warchalowskii* Kadej, Háva & Kalík, 2007. *Baltic Journal of Coleopterology* 23(1): 41–50. [https://doi.org/10.59893/blc.23\(1\).003](https://doi.org/10.59893/blc.23(1).003)
- Kadej M, Háva J, Kalík V. 2007.** Review of the *Anthrenus pimpinellae* species group from Palaearctic region (Coleoptera: Dermestidae: Anthrenini). *Genus* 18(4): 721–750.
- Schneider CA, Rasband WS, Eliceiri KW. 2012.** NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671–675.
- Shorthouse DP. 2010.** SimpleMappr, an online tool to produce publication-quality point maps. Available at <https://www.simplemappr.net> (Last accessed 22<sup>nd</sup> February 2024.)

Received March 24, 2024; accepted June 3, 2024.

Review editor Paul E. Skelley.