

Minimisers of supremal functionals and mass-minimising 1-currents

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Katzourakis, N. and Moser, R. (2025) Minimisers of supremal functionals and mass-minimising 1-currents. *Calculus of Variations and Partial Differential Equations*, 64 (1). 26. ISSN 1432-0835 doi: <https://doi.org/10.1007/s00526-024-02892-5> Available at <https://centaur.reading.ac.uk/119518/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s00526-024-02892-5>

Publisher: Springer-Verlag

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Minimisers of supremal functionals and mass-minimising 1-currents

Nikos Katzourakis¹ · Roger Moser²

Received: 3 May 2024 / Accepted: 12 November 2024 / Published online: 6 December 2024
© The Author(s) 2024

Abstract

We study vector-valued functions that minimise the L^∞ -norm of their derivatives for prescribed boundary data. We construct a vector-valued, mass minimising 1-current (i.e., a generalised geodesic) in the domain such that all solutions of the problem coincide on its support. Furthermore, this current can be interpreted as a streamline of the solutions. The construction relies on a p -harmonic approximation. In the case of scalar-valued functions, it is closely related to a construction of Evans and Yu (Commun Partial Differ Equ 30:1401–1428, 2005). We therefore obtain an extension of their theory.

Mathematics Subject Classification 49K10 · 49Q15

1 Introduction

For $n \in \mathbb{N}$, let $\Omega \subseteq \mathbb{R}^n$ be a bounded domain with smooth boundary. Given $N \in \mathbb{N}$, we study functions $u: \Omega \rightarrow \mathbb{R}^N$ that minimise the functional

$$E_\infty(u) = \operatorname{ess\,sup}_\Omega |Du|$$

for prescribed boundary data, where $|\cdot|$ denotes the Frobenius norm of an $(N \times n)$ -matrix.

Variational problems of this sort go back to the pioneering work of Aronsson [2–5]. The scalar case $N = 1$ is now quite well understood. For the above functional E_∞ , it gives rise to the Aronsson equation, which has a well-developed theory in the framework of viscosity solutions. It has unique solutions for given boundary data, which correspond not just to minimisers of the functional, but to so-called absolute minimisers [21]. These are characterised by the condition that they minimise $\operatorname{ess\,sup}_{\Omega'} |Du|$ in suitable subsets $\Omega' \subseteq \Omega$

Communicated by A. Mondino.

✉ Roger Moser
r.moser@bath.ac.uk
Nikos Katzourakis
n.katzourakis@reading.ac.uk

¹ Department of Mathematics and Statistics, University of Reading, Whiteknights, Pepper Lane, Reading RG6 6AX, UK

² Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK

for their own boundary data. There is a regularity theory for solutions of the equation as well [14, 16, 17, 30, 35].

Much less is known for the vector-valued case $N \geq 2$, despite some work by the first author [22–27]. (This problem should not be confused with the problem of vector-valued optimal Lipschitz extensions [36], which amounts to replacing the Frobenius norm with the operator norm. For $n = 1$ or $N = 1$, the two problems are equivalent, but in general, they are not.) The vector-valued counterpart to the Aronsson equation is easy to write down, but much more difficult to make sense of. Above all, there is no meaningful interpretation of viscosity solutions. It is not known if absolute minimisers exist in general or if they are unique. We therefore take a somewhat different point of view in this paper. Rather than trying to determine a distinguished solution of the problem, we ask what all minimisers of E_∞ for given boundary data have in common.

The natural space for the functional E_∞ is the Sobolev space $W^{1,\infty}(\Omega; \mathbb{R}^N)$. We prescribe boundary data given in terms of a fixed Lipschitz continuous function $u_0: \mathbb{R}^n \rightarrow \mathbb{R}^N$, and then we are interested in the subset $u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$. We define

$$e_\infty = \inf_{u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)} E_\infty.$$

We are thus interested in the following problem.

Problem *Study all maps $u_\infty \in u_0 + W^{1,\infty}(\Omega; \mathbb{R}^N)$ such that $E_\infty(u_\infty) = e_\infty$.*

Since u_0 is Lipschitz continuous, it has a tangential derivative at almost every boundary point with respect to the $(n - 1)$ -dimensional Hausdorff measure. We write $D'u_0$ for the tangential derivative of u_0 , and we define

$$e'_\infty = \operatorname{ess\,sup}_{\partial\Omega} |D'u_0|$$

(where the essential supremum is taken with respect to the $(n - 1)$ -dimensional Hausdorff measure on $\partial\Omega$). The case where $e'_\infty < e_\infty$ is particularly interesting for the reasons explained below.

One of the key ingredients in our analysis is a measure derived as a limit from p -harmonic approximations. This tool essentially goes back to an idea of Evans [15] and has been studied by Evans and Yu [18] for the case $N = 1$, albeit in a different form. We show here that there is an interesting geometric structure behind this limit measure. Moreover, it is useful for the vector-valued problem, too, even though this requires an approach rather different from the one taken by Evans and Yu.

Roughly speaking, we will find some generalised length-minimising ‘curves’ in the domain, along which all solutions of the problem must coincide. In order to formulate a precise statement, however, we need some tools from geometric measure theory, above all the notion of currents.

Since we will work only with 1-currents (and their boundaries, given by 0-currents), readers not familiar with geometric measure theory may think of vector-valued distributions instead, or indeed of vector-valued measures in most cases. This will, however, obscure some of the geometric content of our results.

Definition 1 (Current, mass, boundary) For $j = 0, \dots, n$, let $\mathcal{D}^j(\mathbb{R}^n)$ denote the space of smooth, compactly supported j -forms in \mathbb{R}^n , endowed with the topology induced by locally uniform convergence of all derivatives. A j -current in \mathbb{R}^n is an element of its dual space $\mathcal{D}_j(\mathbb{R}^n) = (\mathcal{D}^j(\mathbb{R}^n))^*$. Given a j -current $T \in \mathcal{D}_j(\mathbb{R}^n)$, its mass is¹

$$\mathbf{M}(T) = \sup \left\{ T(\omega) : \omega \in \mathcal{D}^j(\mathbb{R}^n) \text{ with } \sup_{\mathbb{R}^n} |\omega| \leq 1 \right\}.$$

If $j \geq 1$, its boundary is the $(j - 1)$ -current ∂T such that $\partial T(\sigma) = T(d\sigma)$ for every $\sigma \in \mathcal{D}^{j-1}(\mathbb{R}^n)$.

We are interested above all in 1-currents supported on $\overline{\Omega}$ and their boundaries (which are 0-currents and can be regarded as distributions). These can be interpreted as generalised oriented curves in $\overline{\Omega}$. Indeed, given an oriented C^1 -curve $\Gamma \subset \overline{\Omega}$ of finite length, we can define a corresponding 1-current T by integration of 1-forms over Γ , i.e.,

$$T(\omega) = \int_{\Gamma} \omega. \tag{1}$$

In this situation, the mass $\mathbf{M}(T)$ will be the length of Γ . More generally, if μ is a Radon measure on $\overline{\Omega}$ and τ is a vector field that is integrable with respect to μ , then we can define a 1-current $T = [\mu, \tau]$ by the formula

$$T(\omega) = \int_{\overline{\Omega}} \omega(\tau) d\mu. \tag{2}$$

In this case,

$$\mathbf{M}(T) = \int_{\overline{\Omega}} |\tau| d\mu$$

and

$$\partial T(\sigma) = \int_{\overline{\Omega}} d\sigma(\tau) d\mu$$

for a 0-form σ on \mathbb{R}^n . For example, suppose that we have the above oriented curve Γ . Write \mathcal{H}^1 for the one-dimensional Hausdorff measure and define $\mu = \mathcal{H}^1 \llcorner \Gamma$. Furthermore, let τ denote the unit tangent vector field along Γ consistent with the orientation. Then the formulas (1) and (2) give rise to the same current.

We will also need to study N -tuples $T = (T_1, \dots, T_N)$ of 1-currents, which may also be regarded as vector-valued currents. The boundary ∂T is then taken component-wise. We use the following extension of the mass.

Definition 2 (Joint mass, normal) Let $T = (T_1, \dots, T_N) \in (\mathcal{D}_j(\mathbb{R}^n))^N$ be an N -tuple of j -currents in \mathbb{R}^n . Then the joint mass of T is

$$\mathbf{M}(T) = \sup \left\{ \sum_{k=1}^N T_k(\omega_k) : \omega_1, \dots, \omega_N \in \mathcal{D}^j(\mathbb{R}^n) \text{ with } \sup_{\mathbb{R}^n} \sum_{k=1}^N |\omega_k|^2 \leq 1 \right\}.$$

For $j \geq 1$, we say that T is normal if $\mathbf{M}(T) < \infty$ and $\mathbf{M}(\partial T) < \infty$.

¹ It is more common to define the mass in terms of the co-mass norm on the space of j -covectors, but this definition appears, e.g., in a book by Simon [37]. In the context of this paper, the distinction is inconsequential, because we will work only with 1-currents and their boundaries.

Given $T = (T_1, \dots, T_N) \in (D_1(\mathbb{R}^n))^N$ with $\mathbf{M}(T) < \infty$, we can always find a Radon measure $\|T\|$ on \mathbb{R}^n and $\|T\|$ -measurable vector fields $\vec{T}_1, \dots, \vec{T}_N$ such that $\sum_{k=1}^n |\vec{T}_k|^2 = 1$ almost everywhere and $T_k = [\|T\|, \vec{T}_k]$ for $k = 1, \dots, N$. Furthermore, this representation is unique (up to identification of vector fields that agree $\|T\|$ -almost everywhere). We then write $\vec{T} = (\vec{T}_1, \dots, \vec{T}_N)$.

Similarly, if $\mathbf{M}(\partial T) < \infty$, then ∂T is represented by an \mathbb{R}^N -valued Radon measure. For any continuous function $u: \mathbb{R}^n \rightarrow \mathbb{R}^N$ with compact support, we may then write

$$\partial T(u) = \sum_{k=1}^N \partial T_k(u_k).$$

If $\text{supp } \partial T \subseteq \overline{\Omega}$, then this also makes sense for any $u \in W^{1,\infty}(\Omega; \mathbb{R}^N)$, including the minimisers of E_∞ .

For a minimiser $u_\infty \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ of E_∞ , and for T as above, we want to be able to make statements about the ‘behaviour of Du_∞ along $\|T\|$ ’. Since the support of $\|T\|$ can be a null set with respect to the Lebesgue measure on Ω , and since Du_∞ is only well-defined up to null sets, this requires some explanation.

Definition 3 (Local L^2 -representative) Let μ be a (non-negative) Radon measure on Ω and let $f \in L^1_{\text{loc}}(\Omega)$. We say that $g \in L^2_{\text{loc}}(\mu)$ is the *local L^2 -representative* of f with respect to μ , and we write

$$g = \langle f \rangle_\mu,$$

if for any compact set $K \Subset \Omega$ and for any $\eta \in C_0^\infty(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} \eta(x) dx = 1$, the functions $\eta_\epsilon(x) = \epsilon^{-n} \eta(x/\epsilon)$ satisfy

$$\lim_{\epsilon \searrow 0} \int_K |g - \eta_\epsilon * f|^2 d\mu = 0.$$

The concept is defined similarly for vector-valued functions.

We require some more notation. For $x \in \mathbb{R}^n$ and $r > 0$, let $B_r(x)$ denote the open ball in \mathbb{R}^n with centre x and radius r . Given a measurable function $f: \Omega \rightarrow [0, \infty)$, we define $f^*: \overline{\Omega} \rightarrow [0, \infty]$ by

$$f^*(x) = \lim_{r \searrow 0} \text{ess sup}_{B_r(x) \cap \Omega} f$$

for $x \in \overline{\Omega}$.

We can now formulate our first main result.

Theorem 4 *There exists an N -tuple of 1-currents $T = (T_1, \dots, T_N)$ of finite joint mass with the following properties.*

- (i) $\text{supp } T \subseteq \overline{\Omega}$ and $\text{supp } \partial T \subseteq \partial \Omega$.
- (ii) For any minimiser $u = (u_1, \dots, u_N) \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ of E_∞ ,
 - (a) $|Du|^*(x) = e_\infty$ for $\|T\|$ -almost every $x \in \Omega$;
 - (b) $\langle Du \rangle_{\|T\|} = e_\infty \vec{T}$.
- (iii) Any two minimisers of E_∞ in $u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ coincide on $\text{supp } T$.
- (iv) If $e'_\infty < e_\infty$, then the following holds true.
 - (a) T is normal.

- (b) $\|T\|(\partial\Omega) = 0$ and $\|T\|(\Omega) > 0$.
- (c) $\partial T(u_0) = e_\infty \mathbf{M}(T)$.
- (d) Let S be a normal N -tuple of 1-currents such that $\text{supp } S \subseteq \overline{\Omega}$ and $\text{supp } \partial S \subseteq \partial\Omega$. If $\partial S = \partial T$, then $\mathbf{M}(S) \geq \mathbf{M}(T)$.

The theorem may appear technical, but there is a geometric interpretation. Statement (i) says that we have N generalised curves T_1, \dots, T_N in $\overline{\Omega}$ with no boundary in the interior of Ω . Along these generalised curves, according to (ii), any solution u of our variational problem will have a derivative of norm e_∞ (in a certain sense), with Du_k tangent to T_k . We therefore have a generalisation of what is called a *streamline* in the theory of ∞ -harmonic functions [4]. (For the vector-valued case, similar observations have been made by the first author for sufficiently smooth solutions [22].) Moreover, we have uniqueness of minimisers on the support of T by (iii).

If $e'_\infty < e_\infty$, then we can make additional statements on the structure of T according to (iv). It follows in particular that T is non-trivial in the interior of Ω by (b). Furthermore, statement (d) says that T minimises the joint mass (generalised length) among all competitors with the same boundary in this case. This means that we can think of *generalised length-minimising geodesics* rather than just generalised curves.

If $e'_\infty = e_\infty$, then the theorem may appear vacuous, because it does not rule out that T is supported completely on the boundary or even that $T = 0$. The proof, however, reveals some more information, which is omitted here for the sake of brevity. In our construction, the 1-current T will arise from a limit of p -harmonic functions $u_p: \Omega \rightarrow \mathbb{R}^N$, or more precisely, from renormalisations of the currents $[|Du_p|^{p-2} \mathcal{L}^n, Du_p]$, where \mathcal{L}^n is the Lebesgue measure on \mathbb{R}^n . It is possible that $\text{supp } T \subseteq \partial\Omega$, and then the theorem is indeed vacuous, but it can be interesting even if $e'_\infty = e_\infty$. In any case, $\|T\|$ coincides with the measure studied by Evans and Yu [18] for $N = 1$. They proved a condition that amounts to (i) and some additional properties related to (ii) and (iii). The analysis of this paper can therefore also be regarded as an extension of their results. Statement (iv) is completely new even if $N = 1$.

In the case $e'_\infty = e_\infty$, we still have a local variant of (iv).(d). Since its formulation is even more technical, however, we postpone it until Sect. 6 (see Theorem 19). We can further derive some information about the structure of T , which we will do in Sect. 7 (see Theorem 21). We employ standard results from geometric measure theory here, even though we consider non-standard objects. This result can be thought of as a consequence of the the mass minimising property in statement (iv).(d), but our construction allows a shortcut.

There is a simple observation that exemplifies some of the behaviour described in Theorem 4. Suppose that there are two boundary points $x, y \in \partial\Omega$ such that the line segment L between x and y is contained in $\overline{\Omega}$. Suppose further that $|u_0(x) - u_0(y)| \geq e_\infty |x - y|$. Then, if u is a minimiser of E_∞ in $u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$, we can see quite easily that the restriction of u to L is affine with a derivative of norm e_∞ . In general, such a pair of points need not exist, but the N -tuple of currents T from Theorem 4 has properties similar to the line segment L .

In the case $N = 1$, it was in fact proved by Aronsson [4, Theorem 2] that the set where all minimisers coincide is characterised by line segments as above. (Other related results also exist [9, 10].) This is not true for $N > 1$. Consider, e.g., the case $n = N$. If $u_0(x) = x$ for all $x \in \Omega$, then u_0 is p -harmonic for all $p < \infty$, and our analysis shows that it is also the unique minimiser of E_∞ in $u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^n)$. But for any $x, y \in \partial\Omega$, we clearly have the identity $|u_0(x) - u_0(y)| = |x - y|$, while $e_\infty = \sqrt{n}$ in this case.

We have interpreted T from Theorem 4 as a generalised curve, but in general it need not actually be 1-dimensional. For example, suppose that u_0 is an affine function. Then, as in the preceding example, we conclude that every p -harmonic function for these boundary data

coincides with u_0 , and an examination of our construction reveals that $\|T\|$ is a normalised version of the Lebesgue measure in Ω and \vec{T} is constant. We can then interpret T as the collective representation of many line segments in Ω .

If $e'_\infty = e_\infty$, then general statements on the behaviour of minimisers of E_∞ in the interior of Ω cannot be expected. In this case, the minimum value of E_∞ is in fact dictated by the local behaviour near a single boundary point.

Proposition 5 *If $e_\infty = e'_\infty$, then there exists $x \in \partial\Omega$ such that $|Du|^*(x) \geq e_\infty$ for any $u \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$.*

In this situation, we may have a minimiser $u \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ such that $\|Du\|_{L^\infty(K)} < e_\infty$ for any compact set $K \subset \Omega$. Then for any $\phi \in W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ with compact support, there exists $\delta > 0$ such that $u_\infty + t\phi$ is also a minimiser of E_∞ for all $t \in (-\delta, \delta)$. Thus the class of minimisers is simply too large to expect any general statements. (For example, let $n = 2$ and $N = 1$. Let $r = \sqrt{2}/(\sqrt{2} + 1)$ and consider the domain $\Omega = B_r(r, r) \subseteq \mathbb{R}^2$. Suppose that $u_0(x_1, x_2) = x_1^{4/3} - x_2^{4/3}$. This happens to be a minimiser of E_∞ for its boundary values [6]. We find that $e_\infty = e'_\infty = 4\sqrt{2}/3$, which is the value of $|Du_0|$ at $(1, 1)$ but nowhere else in $\overline{\Omega}$.) If $e'_\infty < e_\infty$, on the other hand, Theorem 4 rules out such behaviour.

Variational problems in L^∞ have long been studied predominantly with methods involving comparison arguments and viscosity solutions of the corresponding partial differential equations. Even so, measure theoretic arguments have also made an appearance in the literature in various contexts [11, 12, 18, 28]. This also includes a paper with a more geometric point of view by Daskalopoulos and Uhlenbeck [13], which is motivated by work of Thurston [38]. Their paper studies ‘ ∞ -harmonic maps’ from a hyperbolic manifold to the circle and shows (among other things) that the locus of maximum stretch is a geodesic lamination and is contained in the support of a certain $(n - 1)$ -current. (It should be noted here that $(n - 1)$ -currents can be identified with 1-currents via the Hodge star operator.) There is thus some overlap with the above theory, but the paper of Daskalopoulos and Uhlenbeck is restricted to one-dimensional targets and its scope is different. Further related results can be found in a work by Backus [7].

Notation *The following notation will be convenient. Given $r > 0$, we define $\Omega_r = \{x \in \Omega : \text{dist}(x, \partial\Omega) < r\}$. Given two matrices $A, B \in \mathbb{R}^{N \times n}$, we write $A : B$ for their Frobenius inner product.*

2 Currents and what they say about E_∞

In this section, we give some more information on the relationship between N -tuples of 1-currents and the functions minimising E_∞ in $u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$.

We first formulate a global version of Definition 3. We cannot simply use the convolution with a mollifying kernel any more, because this would cause problems at the boundary. Instead, we use the following.

Definition 6 (Regular mollifier) A family of linear operators $\mathcal{M}_\epsilon : L^\infty(\Omega) \rightarrow C^\infty(\overline{\Omega})$, for $\epsilon > 0$, is called a *regular mollifier* if

- (i) there exists $\theta : (0, \infty) \rightarrow [0, \infty)$ such that $\lim_{\epsilon \searrow 0} \theta(\epsilon) = 0$ and

$$|(\mathcal{M}_\epsilon f)(x)| \leq \|f\|_{L^\infty(\Omega \cap B_\epsilon(x))} + \theta(\epsilon)$$

- for all $x \in \overline{\Omega}$;
- (ii) if $f \in C^0(\overline{\Omega})$, then $\mathcal{M}_\epsilon f \rightarrow f$ uniformly as $\epsilon \searrow 0$; and
- (iii) if $f \in W^{1,\infty}(\Omega)$, then

$$\left| \mathcal{M}_\epsilon \frac{\partial f}{\partial x_j} - \frac{\partial x_j}{\partial \mathcal{M}_\epsilon} f \right| \rightarrow 0$$

in $C^0(\overline{\Omega})$ as $\epsilon \searrow 0$ for $j = 1, \dots, n$.

Regular mollifiers can be constructed by modifying the usual convolution with a mollifying kernel. For example, a suitable approach is used in a recent paper by the authors [29, Proof of Lemma 7]. Some tools for a different construction are discussed in a paper by the first author [28, Section 5].

Definition 7 (*L²-representative*) Let μ be a (non-negative) Radon measure on $\overline{\Omega}$ and let $f \in L^\infty(\Omega)$. We say that $g \in L^2(\mu)$ is the *L²-representative* of f with respect to μ , and we write

$$g = \llbracket f \rrbracket_\mu,$$

if for every regular mollifier \mathcal{M}_ϵ ,

$$\lim_{\epsilon \searrow 0} \int_{\overline{\Omega}} |g - \mathcal{M}_\epsilon f|^2 d\mu = 0.$$

The concept is defined similarly for vector-valued functions.

We have the following connection between currents and the functional E_∞ .

Proposition 8 *Suppose that $T = (T_1, \dots, T_N)$ is a normal N -tuple of 1-currents in \mathbb{R}^n with $\text{supp } T \subseteq \overline{\Omega}$. Let $u \in W^{1,\infty}(\Omega; \mathbb{R}^N)$. Then $\partial T(u) \leq E_\infty(u)\mathbf{M}(T)$, with equality if, and only if, $\llbracket Du \rrbracket_{\|T\|} = E_\infty(u)\vec{T}$.*

Proof If $E_\infty(u) = 0$, then the statement is trivial. We therefore assume that $E_\infty(u) > 0$.

Consider a regular mollifier $(\mathcal{M}_\epsilon)_{\epsilon>0}$ and set $u_\epsilon = \mathcal{M}_\epsilon u$. We define $e = E_\infty(u)$. Then we compute

$$\begin{aligned} \int_{\overline{\Omega}} |e\vec{T} - Du_\epsilon|^2 d\|T\| &= \int_{\overline{\Omega}} (e^2 + |Du_\epsilon|^2) d\|T\| - 2e \int_{\overline{\Omega}} \vec{T} : Du_\epsilon d\|T\| \\ &= \int_{\overline{\Omega}} (e^2 + |Du_\epsilon|^2) d\|T\| - 2e\partial T(u_\epsilon). \end{aligned} \tag{3}$$

By conditions (i) and (iii) in Definition 6, we know that

$$\limsup_{\epsilon \searrow 0} \|Du_\epsilon\|_{C^0(\overline{\Omega})} \leq \|Du\|_{L^\infty(\Omega)} \leq e.$$

It follows that

$$\limsup_{\epsilon \searrow 0} \int_{\overline{\Omega}} (e^2 + |Du_\epsilon|^2) d\|T\| \leq 2e^2\mathbf{M}(T).$$

On the other hand, we have the uniform convergence $u_\epsilon \rightarrow u$. As T is normal, it follows that $\partial T(u) = \lim_{\epsilon \searrow 0} \partial T(u_\epsilon)$. Taking the limsup in (3), we find that

$$\partial T(u) + \frac{1}{2e} \limsup_{\epsilon \searrow 0} \int_{\overline{\Omega}} |e\vec{T} - Du_\epsilon|^2 d\|T\| \leq e\mathbf{M}(T).$$

Hence $\partial T(u) \leq e\mathbf{M}(T)$. If we have equality, then it further follows that $\llbracket Du \rrbracket_{\|T\|} = e\vec{T}$.

Conversely, suppose that $\llbracket Du \rrbracket_{\|T\|} = e\vec{T}$. Then $\mathcal{M}_\epsilon Du \rightarrow e\vec{T}$ in $L^2(\|T\|)$. By property (iii) in Definition 6, this means that $Du_\epsilon \rightarrow e\vec{T}$ in $L^2(\|T\|)$ as well. Hence

$$\partial T(u) = \lim_{\epsilon \searrow 0} \partial T(u_\epsilon) = \lim_{\epsilon \searrow 0} \int_{\Omega} Du_\epsilon : \vec{T} d\|T\| = e \int_{\Omega} |\vec{T}|^2 d\|T\| = e\mathbf{M}(T),$$

as claimed. □

Remark These arguments also show that if $\mathcal{M}_\epsilon Du \rightarrow E_\infty(u)\vec{T}$ in $L^2(\|T\|)$ for some *specific* regular mollifier \mathcal{M}_ϵ , then $\llbracket Du \rrbracket_{\|T\|} = E_\infty(u)\vec{T}$.

Corollary 9 *Suppose that $T = (T_1, \dots, T_N)$ is a normal N -tuple of 1-currents in \mathbb{R}^n with $\text{supp } T \subseteq \bar{\Omega}$ and $\text{supp } \partial T \subseteq \partial\Omega$. Then $\partial T(u_0) \leq e_\infty\mathbf{M}(T)$. If equality holds and $e_\infty > 0$, then $\mathbf{M}(S) \geq \mathbf{M}(T)$ for any normal N -tuple of 1-currents S such that $\text{supp } S \subseteq \bar{\Omega}$ and $\partial S = \partial T$.*

Proof In order to prove the first statement, it suffices to choose a minimiser $u_\infty \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ of E_∞ and apply Proposition 8 to u_∞ . If we have equality and if $e_\infty > 0$, we apply the first statement to S . We conclude that $\mathbf{M}(T) = e_\infty^{-1}\partial T(u_0) = e_\infty^{-1}\partial S(u_0) \leq \mathbf{M}(S)$. □

The following are local versions of the above statements.

Corollary 10 *Suppose that $T = (T_1, \dots, T_N)$ is an N -tuple of 1-currents in \mathbb{R}^n with locally finite mass and with $\text{supp } \partial T \cap \Omega = \emptyset$. Let $u \in W^{1,\infty}(\Omega; \mathbb{R}^N)$. Then*

$$-T(ud\chi) \leq E_\infty(u) \int_{\Omega} \chi d\|T\|$$

for any $\chi \in C_0^\infty(\Omega)$ with $\chi \geq 0$. Equality holds for every such function χ if, and only if, $\llbracket Du \rrbracket_{\|T\|} = E_\infty(u)\vec{T}$.

Proof Fix $\chi \in C_0^\infty(\Omega)$ with $\chi \geq 0$. Consider $S \in (\mathcal{D}_1(\mathbb{R}^n))^N$ defined by $S_k(\omega) = T_k(\chi\omega)$ for $\omega \in (\mathcal{D}^1(\mathbb{R}^n))$ and for $k = 1, \dots, N$. Then S is normal with $\partial S(u) = -T(ud\chi)$ and

$$\mathbf{M}(S) = \int_{\Omega} \chi d\|T\|.$$

The first statement therefore follows immediately from Proposition 8.

Let $\eta \in C_0^\infty(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} \eta dx = 1$. Let $\eta_\epsilon(x) = \epsilon^{-n}\eta(x/\epsilon)$. Then we can construct a regular mollifier \mathcal{M}_ϵ such that $\mathcal{M}_\epsilon f = \eta_\epsilon * f$ in $\text{supp } \chi$ for all $f \in L^\infty(\Omega)$. Therefore, the second statement follows from Proposition 8 and the remark after its proof. □

Corollary 11 *Let $T = (T_1, \dots, T_N)$ be an N -tuple of 1-currents in \mathbb{R}^n with locally finite mass and with $\text{supp } \partial T \cap \Omega = \emptyset$. Let $u \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$. Suppose that*

$$-T(ud\chi) \geq e_\infty \int_{\Omega} \chi d\|T\|$$

for every $\chi \in C_0^\infty(\Omega)$ with $\chi \geq 0$. Then the following holds true.

(i) For any open set $\Omega' \subseteq \Omega$ with $\text{supp } T \cap \Omega' \neq \emptyset$,

$$\text{ess sup}_{\Omega'} |Du| \geq e_\infty.$$

(ii) Suppose that $e_\infty > 0$. Let $K \subset \Omega$ be a compact set. Let $S \in (\mathcal{D}_1(\mathbb{R}^n))^N$ with locally finite mass and with $\text{supp } \partial S \cap \Omega = \emptyset$. If $S(\omega) = T(\omega)$ for every $\omega \in (\mathcal{D}^1(\mathbb{R}^n))^N$ with $\omega = 0$ in K , then

$$\|T\|(K) \leq \|S\|(K).$$

Proof We may assume that $e_\infty > 0$, because (i) is trivial otherwise and (ii) excludes $e_\infty = 0$.

Given an open set $\Omega' \subseteq \Omega$ with $\text{supp } T \cap \Omega' \neq \emptyset$, choose $\chi \in C_0^\infty(\Omega')$ with $\chi \geq 0$ and $\int_{\Omega'} \chi d\|T\| > 0$. Choose an open set $\Omega'' \subseteq \Omega'$ with smooth boundary such that $\text{supp } \chi \subset \Omega''$. Applying Corollary 10 in Ω'' , we find that

$$-T(ud\chi) \leq \text{ess sup}_{\Omega''} |Du| \int_{\Omega''} \chi d\|T\|.$$

But by the assumptions, we also have

$$-T(ud\chi) \geq e_\infty \int_{\Omega''} \chi d\|T\|.$$

Hence $\text{ess sup}_{\Omega'} |Du_\infty| \geq e_\infty$.

If S and K are as in the second statement, then we choose $\chi \in C_0^\infty(\Omega)$ such that $\chi \equiv 1$ in K and $0 \leq \chi \leq 1$ everywhere. Then

$$\int_{\Omega} \chi d\|T\| \leq -\frac{1}{e_\infty} T(ud\chi) = -\frac{1}{e_\infty} S(ud\chi) \leq \int_{\Omega} \chi d\|S\|$$

by Corollary 10. Since $\|T\|(K)$ and $\|S\|(K)$ are the limits of such integrals when χ approaches the characteristic function of K , the claim follows. \square

If T has a mass minimising property as in Corollary 9 or Corollary 11, then we may think of it as a generalised length-minimising geodesic. There is an Euler-Lagrange equation for this variational problem, which amounts to the condition that

$$\sum_{k=1}^N \int_{\Omega} \vec{T}_k \cdot D_{\vec{T}_k} \psi d\|T\| = 0 \tag{4}$$

for all $\psi \in C_0^\infty(\Omega; \mathbb{R}^n)$, where $D_{\vec{T}_k}$ denotes the directional derivative in the direction of \vec{T}_k . This equation can be derived with the same arguments as for the more conventional (scalar-valued) mass minimising currents. (These tools are even more common in the theory of varifolds [1], but this is just another side of the same coin for this purpose.) We do not need to go into the details here, because we will obtain the equation in a different way. Instead, we formulate a consequence.

Proposition 12 Suppose that T is an N -tuple of 1-currents such that (4) holds true for all $\psi \in C_0^\infty(\Omega; \mathbb{R}^n)$. Let $w \in W_0^{1,\infty}(\Omega; \mathbb{R}^N)$. If $\langle Dw \rangle_{\|T\|} = 0$, then $w = 0$ on $\text{supp } T$.

Proof Let $r > 0$. Let $\chi \in C_0^\infty(\Omega)$ with $\chi \equiv 1$ in $\Omega \setminus \Omega_r$ and such that $|D\chi| \leq 2r$. Let $\eta \in C_0^\infty(\mathbb{R}^n)$ with $\int_{\mathbb{R}^n} \eta dx = 1$ and set $\eta_\epsilon(x) = \epsilon^{-n} \eta(x/\epsilon)$ for $\epsilon > 0$. Define $w_\epsilon = \eta_\epsilon * w$, assuming that w is extended (arbitrarily) to \mathbb{R}^n .

Set

$$\psi_\epsilon(x) = |w_\epsilon(x)|^2 \chi(x)$$

for $x \in \Omega$. Then we may test (4) with ψ_ϵ . This gives

$$0 = 2 \sum_{k=1}^N \int_{\Omega} x \cdot \vec{T}_k (w_\epsilon \otimes \vec{T}_k) : Dw_\epsilon \chi \, d\|T\| + \sum_{k=1}^N \int_{\Omega} x \cdot \vec{T}_k |w_\epsilon|^2 D\vec{T}_k \chi \, d\|T\| + \int_{\Omega} |w_\epsilon|^2 \chi \, d\|T\|.$$

Since $\langle Dw \rangle_{\|T\|} = 0$, we have the convergence

$$\int_{\text{supp } D\chi} |Dw_\epsilon|^2 \, d\|T\| \rightarrow 0$$

as $\epsilon \searrow 0$. Clearly $w_\epsilon \rightarrow w$ locally uniformly. Hence

$$0 = \sum_{k=1}^N \int_{\Omega} x \cdot \vec{T}_k |w|^2 D\vec{T}_k \chi \, d\|T\| + \int_{\Omega} |w|^2 \chi \, d\|T\|.$$

If we let $r \searrow 0$, then the first term vanishes in the limit, because

$$|w|^2 \leq N E_\infty(w) r^2$$

in Ω_r . Hence

$$0 = \int_{\Omega} |w|^2 \, d\|T\|.$$

It follows that $w = 0$ almost everywhere with respect to $\|T\|$. By the continuity of w , we conclude that $w = 0$ on $\text{supp } T$. □

3 Measure-function pairs and L^p -approximation

Here we introduce another tool from geometric measure theory, due to Hutchinson [20], that is convenient for our purpose. We only discuss the L^2 -version of Hutchinson’s theory here, because this is all we need.

In the second part of this section, we will apply it to minimisers of the functionals

$$E_p(u) = \left(\int_{\Omega} |Du|^p \, dx \right)^{1/p}. \tag{5}$$

The limit $p \rightarrow \infty$ will eventually produce not just a minimiser of E_∞ , but also the 1-currents from Theorem 4.

Definition 13 (Measure-function pair) *A measure-function pair* over $\overline{\Omega}$ with values in $\mathbb{R}^{N \times n}$ is a pair (μ, F) , where μ is a Radon measure on $\overline{\Omega}$ and $F \in L^2(\mu; \mathbb{R}^{N \times n})$.

Hutchinson further defines weak and strong convergence of measure-function pairs. His formulation of weak convergence is somewhat weaker than the following, but the strong convergence is the same.

Definition 14 (Weak and strong convergence) For $\ell \in \mathbb{N}$, let $M_\ell = (\mu_\ell, F_\ell)$ be measure-function pairs over $\overline{\Omega}$ with values in $\mathbb{R}^{N \times n}$. Let $M_\infty = (\mu_\infty, F_\infty)$ be another such measure-function pair.

(i) We have the *weak convergence* $M_\ell \rightarrow M_\infty$ as $\ell \rightarrow \infty$ if

$$\lim_{\ell \rightarrow \infty} \int_{\overline{\Omega}} (\eta + F_\ell : \phi) \, d\mu_\ell = \int_{\overline{\Omega}} (\eta + F_\infty : \phi) \, d\mu_\infty$$

for any $\eta \in C^0(\overline{\Omega})$ and any $\phi \in C^0(\overline{\Omega}; \mathbb{R}^{N \times n})$, and at the same time,

$$\limsup_{\ell \rightarrow \infty} \int_{\overline{\Omega}} (1 + |F_\ell|^2) \, d\mu_\ell < \infty.$$

(ii) We have the *strong convergence* $M_\ell \rightarrow M_\infty$ as $\ell \rightarrow \infty$ if

$$\lim_{a \rightarrow \infty} \int_{\{x \in \overline{\Omega} : |F_\ell(x)| \geq a\}} |F_\ell|^2 \, d\mu_\ell = 0$$

uniformly in ℓ and

$$\lim_{\ell \rightarrow \infty} \int_{\overline{\Omega}} \Phi(x, F_\ell(x)) \, d\mu_\ell(x) = \int_{\overline{\Omega}} \Phi(x, F_\infty(x)) \, d\mu_\infty$$

for all $\Phi \in C_0^0(\overline{\Omega} \times \mathbb{R}^{N \times n})$.

Some of the key statements in this theory are summarised in the following proposition, which amounts to a variant of [20, Theorem 4.4.2]. It generalises well-known results for weak and strong L^2 -convergence for a fixed measure μ .

Proposition 15 *For $\ell \in \mathbb{N}$, suppose that $M_\ell = (\mu_\ell, F_\ell)$ are measure-function pairs over $\overline{\Omega}$ with values in $\mathbb{R}^{N \times n}$. Let $M_\infty = (\mu_\infty, F_\infty)$ be another such measure function-pair. Let $\Phi : \overline{\Omega} \times \mathbb{R}^{N \times n} \rightarrow \mathbb{R}$ be a continuous function and suppose that there exists a constant $C > 0$ such that $|\Phi(x, z)| \leq C(|z|^2 + 1)$ for all $x \in \overline{\Omega}$ and $z \in \mathbb{R}^{N \times n}$.*

(i) *If*

$$\limsup_{\ell \rightarrow \infty} \int_{\overline{\Omega}} (|F_\ell|^2 + 1) \, d\mu_\ell < \infty,$$

then there exists a subsequence $(M_{\ell_m})_{m \in \mathbb{N}}$ that converges weakly.

(ii) *If $(M_\ell)_{\ell \in \mathbb{N}}$ converges weakly to M_∞ , then*

$$\|F_\infty\|_{L^2(\mu_\infty)} \leq \liminf_{\ell \rightarrow \infty} \|F_\ell\|_{L^2(\mu_\ell)}.$$

(iii) *If $(M_\ell)_{\ell \in \mathbb{N}}$ converges weakly to M_∞ and*

$$\|F_\infty\|_{L^2(\mu_\infty)} = \lim_{\ell \rightarrow \infty} \|F_\ell\|_{L^2(\mu_\ell)},$$

then the convergence is strong.

(iv) *If $(M_\ell)_{\ell \in \mathbb{N}}$ converges strongly to M_∞ , then*

$$\int_{\overline{\Omega}} \Phi(x, F_\infty(x)) \, d\mu_\infty(x) = \lim_{\ell \rightarrow \infty} \int_{\overline{\Omega}} \Phi(x, F_\ell(x)) \, d\mu_\ell(x).$$

Proof The first three statements are from [20, Theorem 4.4.2]. The final statement is a little stronger than what is stated in Hutchinson’s paper, but can be proved with the same arguments. (It is also quite easy to prove directly from the definition of strong convergence.) \square

We will apply these concepts to measures and functions generated by L^p -approximations of minimisers of E_∞ . For $2 \leq p < \infty$, we therefore consider the functionals E_p defined in (5). Since they are strictly convex, there exists a unique minimiser $u_p \in u_0 + W_0^{1,p}(\Omega; \mathbb{R}^n)$ of E_p for each $p < \infty$, which satisfies the Euler-Lagrange equation

$$\operatorname{div}(|Du_p|^{p-2} Du_p) = 0 \tag{6}$$

weakly in Ω . Furthermore, for any $\psi \in C_0^\infty(\Omega; \mathbb{R}^n)$, the condition

$$0 = \left. \frac{d}{dt} \right|_{t=0} E_p(u_p \circ (\operatorname{id}_\Omega + t\psi))$$

gives rise to

$$0 = \int_\Omega |Du_p|^{p-2} \left(\sum_{i,j=1}^n \frac{\partial u_p}{\partial x_i} \cdot \frac{\partial u_p}{\partial x_j} \frac{\partial \psi_i}{\partial x_j} - \frac{1}{p} |Du_p|^2 \operatorname{div} \psi \right) dx. \tag{7}$$

Fix $v \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$. If $2 \leq q < p < \infty$, then we observe that

$$E_q(u_q) \leq E_q(u_p) \leq E_p(u_p) \leq E_p(v) \leq E_\infty(v) \tag{8}$$

by Hölder’s inequality and the definition of u_p . Hence the family $(u_p)_{p \in [2, \infty)}$ is bounded in $W^{1,q}(\Omega; \mathbb{R}^N)$ for every $q < \infty$. Therefore, there exists a sequence $p_\ell \rightarrow \infty$ such that $u_{p_\ell} \rightarrow u_\infty$ weakly in all of these spaces for some limit $u_\infty \in u_0 + \bigcap_{q < \infty} W_0^{1,q}(\Omega; \mathbb{R}^N)$. We then estimate

$$E_\infty(u_\infty) = \lim_{q \rightarrow \infty} E_q(u_\infty) \leq \lim_{q \rightarrow \infty} \liminf_{\ell \rightarrow \infty} E_q(u_{p_\ell}) \leq \liminf_{\ell \rightarrow \infty} E_{p_\ell}(u_{p_\ell}) \leq E_\infty(v), \tag{9}$$

where we have used (8) in the last two steps. Hence $u_\infty \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$, and u is a minimiser of the functional E_∞ . In particular, it satisfies $E_\infty(u_\infty) = e_\infty$. We also define $e_p = E_p(u_p)$. Then as in (8), we see that

$$e_q \leq e_p \leq e_\infty$$

whenever $q \leq p \leq \infty$. Moreover, the estimates in (9) imply that $e_p \rightarrow e_\infty$ monotonically.

We now define

$$\mu_p = \frac{|Du_p|^{p-2}}{e_p^{p-2} \mathcal{L}^n(\Omega)} \mathcal{L}^n,$$

where \mathcal{L}^n denotes the Lebesgue measure on $\overline{\Omega}$. These measures have also been studied by Evans and Yu [18] in the case $N = 1$. They should be considered not on their own, but in conjunction with the function Du_p . Thus $M_p = (\mu_p, Du_p)$ naturally forms a measure-function pair over $\overline{\Omega}$.

We can make a few statements about M_p immediately. The Euler-Lagrange equation (6) now becomes

$$\int_\Omega Du_p : D\phi \, d\mu_p = 0 \tag{10}$$

for every $\phi \in C_0^\infty(\Omega; \mathbb{R}^N)$. We also have

$$\int_\Omega \left(\sum_{i,j=1}^n \frac{\partial u_p}{\partial x_i} \cdot \frac{\partial u_p}{\partial x_j} \frac{\partial \psi_i}{\partial x_j} - \frac{1}{p} |Du_p|^2 \operatorname{div} \psi \right) d\mu_p = 0 \tag{11}$$

for every $\psi \in C_0^\infty(\Omega; \mathbb{R}^n)$ because of (7). Moreover, we observe that

$$\int_{\Omega} |Du_p|^2 d\mu_p = e_p^{2-p} \int_{\Omega} |Du_p|^p dx = e_p^2 \leq e_\infty^2$$

and

$$\mu_p(\overline{\Omega}) = e_p^{2-p} \int_{\Omega} |Du_p|^{p-2} dx \leq e_p^{2-p} (E_p(u_p))^{p-2} = 1$$

by Hölder’s inequality. By Proposition 15, we may therefore replace $(p_\ell)_{\ell \in \mathbb{N}}$ with a subsequence such that $M_{p_\ell} \rightarrow M_\infty$ for some measure-function pair $M_\infty = (\mu_\infty, F_\infty)$ over $\overline{\Omega}$. Then

$$\int_{\overline{\Omega}} F_\infty : D\phi d\mu_\infty = \lim_{\ell \rightarrow \infty} \int_{\overline{\Omega}} Du_{p_\ell} : D\phi d\mu_{p_\ell} = 0 \tag{12}$$

for every $\phi \in C_0^\infty(\Omega; \mathbb{R}^N)$ by (10). We will see that (11) also gives a useful equation in the limit, but we need to establish strong convergence first.

4 Interesting boundary data

In this section, we examine the case $e'_\infty < e_\infty$ in more detail. Recall that

$$e_\infty = \inf_{u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)} E_\infty = E_\infty(u_\infty)$$

and

$$e'_\infty = \operatorname{ess\,sup}_{\partial\Omega} |D'u_0|.$$

If $e'_\infty < e_\infty$, then we have better control of the p -harmonic functions u_p near the boundary $\partial\Omega$. Therefore, we can prove additional properties of the limiting measure-function pair $M_\infty = (\mu_\infty, F_\infty)$. The following estimates rely on identity (11). We use measure-function pairs over $\partial\Omega$ with values in \mathbb{R}^N here, which are defined similarly to Definition 13.

Proposition 16 *Suppose that $e'_\infty < e_\infty$. Then there exist measure-function pairs (m_p, f_p) over $\partial\Omega$ with values in \mathbb{R}^N such that*

$$\limsup_{p \rightarrow \infty} \int_{\partial\Omega} (1 + |f_p|^2) dm_p < \infty \tag{13}$$

and

$$\int_{\Omega} Du_p : D\phi d\mu_p = \int_{\partial\Omega} f_p \cdot \phi dm_p \tag{14}$$

for all $\phi \in C^\infty(\mathbb{R}^n; \mathbb{R}^N)$.

Comparing the last identity with the usual integration by parts formula for p -harmonic functions, we may think of m_p as the restriction of $|Du_p|^{p-2}$ to the boundary (up to rescaling) and of f_p as a representation of $\nu \cdot Du_p$, where ν denotes the outer normal vector on $\partial\Omega$. In general, however, we do not know if we have enough regularity of u_p up to the boundary to write down the formulas in these terms.

Proof We first replace u_p by the solutions of a different boundary value problem, for which we have the better regularity theory. We fix $p \in [2, \infty)$ at first. For $\epsilon > 0$, define the function $H_{p,\epsilon} : \mathbb{R} \rightarrow \mathbb{R}$ by $H_{p,\epsilon}(t) = (t^2 + \epsilon^2)^{p/2}$ for $t \in \mathbb{R}$. Consider the functional

$$E_{p,\epsilon}(u) = \left(\int_{\Omega} H_{p,\epsilon}(|Du|) dx \right)^{1/p}.$$

Choose $c \in (e'_\infty, e_\infty)$. Let $u_{0,\epsilon} : \mathbb{R}^n \rightarrow \mathbb{R}^N$ be smooth functions such that

- $u_{0,\epsilon} \rightarrow u_0$ in $W^{1,p}(\Omega; \mathbb{R}^N)$ as $\epsilon \searrow 0$ and
- $|D'u_{0,\epsilon}|^2 + \epsilon^2 \leq c^2$ on $\partial\Omega$ when ϵ is sufficiently small.

Let $u_{p,\epsilon} \in u_{0,\epsilon} + W_0^{1,p}(\Omega; \mathbb{R}^N)$ be the unique minimiser of $E_{p,\epsilon}$ in this space. Then we have the Euler-Lagrange equation

$$\operatorname{div} \left((|Du_{p,\epsilon}|^2 + \epsilon^2)^{p/2-1} Du_{p,\epsilon} \right) = 0 \tag{15}$$

in Ω . There are theories for both interior and boundary regularity for this sort of problem. In particular, results of Uhlenbeck [39] show that $u_{p,\epsilon}$ is smooth in the interior of Ω . Results of Kristensen and Mingione [31, Theorem 1.1] show that $Du_{p,\epsilon}$ belongs to $W^{1/p+s,p}(\Omega; \mathbb{R}^{N \times n})$ and its trace on $\partial\Omega$ belongs to $W^{s,p}(\partial\Omega; \mathbb{R}^{N \times n})$ for some $s > 0$. This is enough to carry out the following computations.

Equation (15) implies that

$$\frac{\partial}{\partial x_j} H_{p,\epsilon}(|Du_{p,\epsilon}|) = p \operatorname{div} \left((|Du_{p,\epsilon}|^2 + \epsilon^2)^{p/2-1} \frac{\partial u_{p,\epsilon}}{\partial x_j} \cdot Du_{p,\epsilon} \right)$$

in Ω for $j = 1, \dots, n$. We choose a smooth vector field $\psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$ such that $\psi = \nu$ on $\partial\Omega$. Then

$$\begin{aligned} & \operatorname{div} \left(H_{p,\epsilon}(|Du_{p,\epsilon}|) \psi - p (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} D\psi u_{p,\epsilon} \cdot Du_{p,\epsilon} \right) \\ &= H_{p,\epsilon}(|Du_{p,\epsilon}|) \operatorname{div} \psi - p (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} \sum_{i,j=1}^n \frac{\partial u_{p,\epsilon}}{\partial x_i} \cdot \frac{\partial u_{p,\epsilon}}{\partial x_j} \frac{\partial \psi_i}{\partial x_j}. \end{aligned}$$

We write \mathcal{H}^{n-1} for the $(n - 1)$ -dimensional Hausdorff measure. Then

$$\begin{aligned} & \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} (|Du_{p,\epsilon}|^2 + \epsilon^2 - p|D_\nu u_{p,\epsilon}|^2) d\mathcal{H}^{n-1} \\ &= \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) \operatorname{div} \phi dx \\ & \quad - p \sum_{i,j=1}^n \int_{\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} \frac{\partial u_{p,\epsilon}}{\partial x_i} \cdot \frac{\partial u_{p,\epsilon}}{\partial x_j} \frac{\partial \psi_i}{\partial x_j} dx. \end{aligned}$$

Hence there exists a constant C , depending only on n and Ω , such that

$$\begin{aligned} & \left| \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} (|Du_{p,\epsilon}|^2 + \epsilon^2 - p|D_\nu u_{p,\epsilon}|^2) d\mathcal{H}^{n-1} \right| \\ & \leq Cp \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx. \end{aligned}$$

Write

$$f_{p,\epsilon} = D_\nu u_{p,\epsilon} \quad \text{and} \quad g_{p,\epsilon} = |D'u_{p,\epsilon}| = \sqrt{|Du_{p,\epsilon}|^2 - |f_{p,\epsilon}|^2}$$

on $\partial\Omega$. Then we may write the above inequality in the form

$$\begin{aligned} & \left| \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} \left(g_{p,\epsilon}^2 + \epsilon^2 - (p-1)|f_{p,\epsilon}|^2 \right) d\mathcal{H}^{n-1} \right| \\ & \leq Cp \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx. \end{aligned}$$

Hence

$$\begin{aligned} & \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} |f_{p,\epsilon}|^2 d\mathcal{H}^{n-1} \\ & \leq \frac{1}{p-1} \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} (g_{p,\epsilon}^2 + \epsilon^2) d\mathcal{H}^{n-1} \\ & \quad + \frac{Cp}{p-1} \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx, \end{aligned} \tag{16}$$

and

$$\begin{aligned} \int_{\partial\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} &= \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} (|f_{p,\epsilon}|^2 + g_{p,\epsilon}^2 + \epsilon^2) d\mathcal{H}^{n-1} \\ &\leq \frac{p}{p-1} \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} (g_{p,\epsilon}^2 + \epsilon^2) d\mathcal{H}^{n-1} \\ &\quad + \frac{Cp}{p-1} \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx \\ &\leq \frac{p-2}{p-1} \int_{\partial\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \\ &\quad + \frac{2}{p-1} \int_{\partial\Omega} (g_{p,\epsilon}^2 + \epsilon^2)^{p/2} d\mathcal{H}^{n-1} \\ &\quad + \frac{Cp}{p-1} \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx. \end{aligned}$$

Here we have used Young's inequality in the last step. It follows that

$$\int_{\partial\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \leq 2 \int_{\partial\Omega} (g_{p,\epsilon}^2 + \epsilon^2)^{p/2} d\mathcal{H}^{n-1} + Cp \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx. \tag{17}$$

Feeding this back into (16), we obtain

$$\begin{aligned} \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} |f_{p,\epsilon}|^2 d\mathcal{H}^{n-1} &\leq \frac{2}{p-1} \int_{\partial\Omega} (g_{p,\epsilon}^2 + \epsilon^2)^{p/2} d\mathcal{H}^{n-1} \\ &\quad + \frac{2Cp}{p-1} \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx. \end{aligned} \tag{18}$$

Define

$$\Gamma_{p,\epsilon}^1 = \left\{ x \in \partial\Omega : |f_{p,\epsilon}(x)|^2 \leq e_p^2 - c^2 \right\}$$

and $\Gamma_{p,\epsilon}^2 = \partial\Omega \setminus \Gamma_{p,\epsilon}^1$. By the choice of the boundary data, we have $g_{p,\epsilon}^2 + \epsilon^2 \leq c^2$ when ϵ is small, and thus $|Du_{p,\epsilon}|^2 + \epsilon^2 \leq e_p^2$ on $\Gamma_{p,\epsilon}^1$. Hence

$$\int_{\Gamma_{p,\epsilon}^1} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \leq e_p^p \mathcal{H}^{n-1}(\partial\Omega).$$

On $\Gamma_{p,\epsilon}^2$, we note that

$$|f_{p,\epsilon}|^2 \geq \frac{e_p^2 - c^2}{c^2} (g_{p,\epsilon}^2 + \epsilon^2)$$

for sufficiently small values of ϵ . Therefore,

$$|Du_{p,\epsilon}|^2 + \epsilon^2 = |f_{p,\epsilon}|^2 + g_{p,\epsilon}^2 + \epsilon^2 \leq \frac{e_p^2}{e_p^2 - c^2} |f_{p,\epsilon}|^2$$

on $\Gamma_{p,\epsilon}^2$. Hence

$$\int_{\Gamma_{p,\epsilon}^2} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \leq \frac{e_p^2}{e_p^2 - c^2} \int_{\partial\Omega} (|Du_{p,\epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1} |f_{p,\epsilon}|^2 d\mathcal{H}^{n-1}.$$

Using (18), we see that

$$\begin{aligned} & \int_{\Gamma_{p,\epsilon}^2} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \\ & \leq \frac{2e_p^2}{(p-1)(e_p^2 - c^2)} \left(\int_{\partial\Omega} (g_{p,\epsilon}^2 + \epsilon^2)^{p/2} d\mathcal{H}^{n-1} + Cp \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx \right) \\ & \leq \frac{2e_p^2}{(p-1)(e_p^2 - c^2)} \left(c^p \mathcal{H}^{n-1}(\partial\Omega) + Cp \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx \right). \end{aligned}$$

Since we know that $e_p \rightarrow e_\infty$ as $p \rightarrow \infty$, it follows that there exists a constant C' , depending on n, Ω, e_∞ , and c , such that

$$\int_{\partial\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) d\mathcal{H}^{n-1} \leq C' \left(e_p^p + \int_{\Omega} H_{p,\epsilon}(|Du_{p,\epsilon}|) dx \right), \tag{19}$$

provided that p is sufficiently large.

We now consider the limit $\epsilon \searrow 0$ (still for $p < \infty$ fixed). Since $E_{p,\epsilon}(u_{p,\epsilon}) \leq E_{p,\epsilon}(u_{0,\epsilon})$, it is clear that the family of functions $u_{p,\epsilon}$ is bounded in $W^{1,p}(\Omega; \mathbb{R}^N)$. We may therefore choose a sequence $\epsilon_k \searrow 0$ such that $u_{p,\epsilon_k} \rightharpoonup \tilde{u}_p$ weakly in this space. It is obvious that $\tilde{u}_p \in u_0 + W_0^{1,p}(\Omega; \mathbb{R}^N)$.

Let $v \in u_0 + W_0^{1,p}(\Omega; \mathbb{R}^N)$. Define $v_\epsilon = v + u_{0,\epsilon} - u_0$, in order to obtain a function in $u_{0,\epsilon} + W_0^{1,p}(\Omega; \mathbb{R}^N)$. Then

$$E_p(\tilde{u}_p) \leq \liminf_{k \rightarrow \infty} E_{p,\epsilon_k}(u_{p,\epsilon_k}) \leq \liminf_{k \rightarrow \infty} E_{p,\epsilon_k}(v_{\epsilon_k}) = E_p(v).$$

Hence \tilde{u}_p minimises E_p in $u_0 + W_0^{1,p}(\Omega; \mathbb{R}^N)$. The strict convexity of the functional implies that $\tilde{u}_p = u_p$. Inserting $v = u_p$ and using similar estimates, we also see that

$$\limsup_{k \rightarrow \infty} E_{p, \epsilon_k}(u_{p, \epsilon_k}) \leq E_p(u_p).$$

Hence the convergence $u_{p, \epsilon_k} \rightarrow u_p$ is strong in $W^{1,p}(\Omega; \mathbb{R}^N)$.

Consider the measures

$$m_{p, \epsilon} = \frac{(|Du_{p, \epsilon}|^2 + \epsilon^2)^{\frac{p}{2}-1}}{e_p^{p-2} \mathcal{L}^n(\Omega)} \mathcal{H}^{n-1} \llcorner \partial\Omega.$$

Using inequality (17), and recalling that $g_{p, \epsilon}^2 + \epsilon^2 \leq c$ when ϵ is small, we see that

$$\limsup_{\epsilon \searrow 0} \int_{\partial\Omega} (1 + |f_{p, \epsilon}|^2) dm_{p, \epsilon} < \infty$$

for any fixed $p < \infty$. Hence, by Proposition 15, we may assume that the measure-function pairs $(m_{p, \epsilon_k}, f_{p, \epsilon_k})$ over $\partial\Omega$ converge weakly to a measure-function pair (m_p, f_p) . Using (15) and passing to the limit, we obtain identity (14). Inequality (13) follows from (19) and Proposition 15. □

We also briefly consider the case $e'_\infty = e_\infty$. We conclude this section by proving Proposition 5, thus showing that in this situation, the minimum value of E_∞ is dictated locally near a single boundary point.

Proof of Proposition 5 Suppose that $e'_\infty = e_\infty$. Fix $u \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$. We extend u outside of Ω such that it is Lipschitz continuous globally in \mathbb{R}^n . Let L denote the Lipschitz constant of this extension.

Define

$$\alpha(x) = \lim_{r \searrow 0} \operatorname{ess\,sup}_{\partial\Omega \cap B_r(x)} |D'u_0|$$

for $x \in \partial\Omega$. This gives rise to an upper semicontinuous function with

$$e_\infty = \sup_{x \in \partial\Omega} \alpha(x).$$

Hence there exists $x \in \partial\Omega$ such that $e_\infty = \alpha(x)$. We fix this point now.

Let $\epsilon \in (0, \frac{1}{2}]$. Then there exists $r > 0$ such that

$$\operatorname{ess\,sup}_{\partial\Omega \cap B_r(x)} |D'u_0| > e_\infty - \epsilon.$$

Since the restriction of u_0 to $\partial\Omega$ is Lipschitz continuous, it is differentiable almost everywhere with respect to \mathcal{H}^{n-1} . Hence we can find a point $y \in \partial\Omega \cap B_r(x)$ such that $D'u_0(y)$ exists with

$$|D'u_0(y)| \geq e_\infty - \epsilon.$$

We may assume without loss of generality that $y = 0$ and $u_0(0) = 0$, and that the tangent space of $\partial\Omega$ at y is $\mathbb{R}^{n-1} \times \{0\}$. We write $A = D'u_0(y) = (a_{kj})_{1 \leq k \leq N, 1 \leq j \leq n-1}$. Then $|A| \leq \sqrt{N}L$.

If $s > 0$ is sufficiently small, then

$$\partial\Omega \cap [-s, s]^n \subseteq [-s, s]^{n-1} \times [-\epsilon s, \epsilon s] \quad \text{and} \quad [-s, s]^{n-1} \times [\epsilon s, s] \subseteq \Omega,$$

while at the same time,

$$|u_0(z') - Az'| \leq \epsilon s$$

for all $z' \in [-s, s]^n \cap \partial\Omega$. Set $\Sigma_\epsilon = [-s, s]^{n-1} \times [\epsilon s, 2\epsilon s]$. For any $z \in \Sigma_\epsilon$, there exists $z' \in [-s, s]^n \cap \partial\Omega$ such that $|z - z'| \leq 3\epsilon s$. Hence

$$|u(z) - Az| \leq |u(z) - u(z')| + |u_0(z') - Az'| + |A(z' - z)| \leq (3(\sqrt{N} + 1)L + 1)\epsilon s$$

for all $z \in \Sigma_\epsilon$.

Integrating $\frac{\partial u_k}{\partial x_j}$ along lines parallel to the x_j -axis, we see that there exists a constant C , depending only on n, N , and L , such that for $1 \leq k \leq N$ and for $1 \leq j \leq n - 1$, the inequality

$$\left| \int_{\Sigma_\epsilon} \frac{\partial u_k}{\partial x_j} dx - a_{kj} \right| \leq C\epsilon$$

holds true. Thus there exists a subset of Σ_ϵ of positive measure where $|Du| \geq |A| - C'\epsilon$ for another constant C' depending only on n, N , and L . (Otherwise we would conclude that

$$\left| \int_{\Sigma_\epsilon} Du dx - A \right| \geq |A| - \int_{\Sigma_\epsilon} |Du| dx \geq C'\epsilon,$$

using the reverse triangle inequality.) It follows that

$$\operatorname{ess\,sup}_{B_r(x)} |Du| \geq |A| - C'\epsilon \geq e_\infty - (C' + 1)\epsilon.$$

As ϵ was chosen arbitrarily, this implies the claim. □

5 Key estimates

Throughout this section and the rest of the paper, the function u_∞ and the measure-function pair $M_\infty = (\mu_\infty, F_\infty)$ are as constructed in Sect. 3 and are fixed. Recall that u_∞ is a minimiser of E_∞ . If $e'_\infty < e_\infty$, then we consider the measure-function pairs (m_p, f_p) from Proposition 16 as well. We may assume that (m_{p_k}, f_{p_k}) converges weakly to a limiting measure-function pair (m_∞, f_∞) over $\partial\Omega$, which will then satisfy

$$\int_\Omega F_\infty : D\phi d\mu_\infty = \int_{\partial\Omega} \phi \cdot f_\infty dm_\infty \tag{20}$$

for all $\phi \in C_0^\infty(\mathbb{R}^n; \mathbb{R}^N)$. If $e'_\infty = e_\infty$, then we choose an arbitrary measure-function pair (m_∞, f_∞) . Even then, we can still use equation (12) for $\phi \in C_0^\infty(\Omega; \mathbb{R}^N)$, and we conclude that (20) still holds true for these test functions.

We now analyse M_∞ in more detail. This will also reveal some information about other possible minimisers of E_∞ . Some of the key arguments in this section are similar to estimates due to Evans and Yu [18].

We prove the following statements.

Theorem 17 *Suppose that $v \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ is a minimiser of E_∞ .*

- (i) *Then $([Dv])_{\mu_\infty} = F_\infty$, and $|Dv|^*(x) = |F_\infty(x)| = e_\infty$ for μ_∞ -almost every $x \in \Omega$.*
- (ii) *If $e'_\infty < e_\infty$, then $([Dv])_{\mu_\infty} = F_\infty$, and $|Dv|^*(x) = |F_\infty(x)| = e_\infty$ for μ_∞ -almost every $x \in \bar{\Omega}$.*

- (iii) The convergence $M_{p_\ell} \rightarrow M_\infty$ is strong in every compact subset of Ω .
- (iv) If $e'_\infty < e_\infty$, then the convergence $M_{p_\ell} \rightarrow M_\infty$ is strong in $\overline{\Omega}$.

Before we prove these results, we note that statement (iii) has the following consequence. If we write $F_\infty = (F_{1\infty}, \dots, F_{N\infty})$, then (11) gives rise to the identity

$$\sum_{k=1}^N \int_{\Omega} F_{k\infty} \cdot D_{F_{k\infty}} \psi \, d\mu_\infty = 0 \tag{21}$$

for all $\psi \in C_0^\infty(\Omega; \mathbb{R}^n)$, owing to Proposition 15. This equation complements (12), and we will use it in Sect. 7 to say something about the structure of (M_∞, F_∞) , although we will formulate this in terms of a corresponding N -tuple of 1-currents.

In the framework of currents, equation (21) corresponds to (4). It is a weak formulation of the equation for geodesics and is one of key properties of the measure-function pair M_∞ .

For the proof of the theorem, we require the following lemma.

Lemma 18 For any $\xi \in C^0(\overline{\Omega})$ with $\xi \geq 0$, and for any $\alpha \in (0, 1)$,

$$\alpha^2 e_p^2 \int_{\Omega} \xi \, d\mu_p \leq \int_{\Omega} \xi |Du_p|^2 \, d\mu_p + \alpha^p e_p^2 \|\xi\|_{C^0(\Omega)}.$$

Proof Given $\alpha \in (0, 1)$, define $A_p = \{x \in \Omega : |Du_p| \leq \alpha e_p\}$. We first note that

$$\mu_p(A_p) = \frac{e_p^{2-p}}{\mathcal{L}^n(\Omega)} \int_{A_p} |Du_p|^{p-2} \, dx \leq \alpha^{p-2} \frac{\mathcal{L}^n(A_p)}{\mathcal{L}^n(\Omega)} \leq \alpha^{p-2}.$$

Hence

$$\begin{aligned} \int_{\Omega} \xi |Du_p|^2 \, d\mu_p &\geq \int_{\Omega \setminus A_p} \xi |Du_p|^2 \, d\mu_p \\ &\geq \alpha^2 e_p^2 \int_{\Omega \setminus A_p} \xi \, d\mu_p \\ &= \alpha^2 e_p^2 \left(\int_{\Omega} \xi \, d\mu_p - \int_{A_p} \xi \, d\mu_p \right) \\ &\geq \alpha^2 e_p^2 \left(\int_{\Omega} \xi \, d\mu_p - \|\xi\|_{C^0(\Omega)} \mu_p(A_p) \right) \\ &\geq \alpha^2 e_p^2 \int_{\Omega} \xi \, d\mu_p - \alpha^p e_p^2 \|\xi\|_{C^0(\Omega)}. \end{aligned}$$

This implies the desired inequality. □

Proof of Theorem 17 We first prove the local statements (i) and (iii), allowing for the possibility that $e'_\infty = e_\infty$. Let $\eta \in C_0^\infty(B_1(0))$ with $\int_{\mathbb{R}^n} \eta \, dx = 1$. Set $\eta_\epsilon(x) = \epsilon^{-n} \eta(x/\epsilon)$ and define $v_\epsilon = v * \eta_\epsilon$ (where for convenience v is extended arbitrarily outside of Ω , so that v_ϵ

is well-defined in Ω). Choose $\xi \in C_0^\infty(\Omega)$ with $0 \leq \xi \leq 1$. Also choose $\alpha \in (0, 1)$. Then

$$\begin{aligned} & \int_{\Omega} \xi |Dv_\epsilon - Du_p|^2 d\mu_p \\ &= \int_{\Omega} \xi (|Dv_\epsilon|^2 - |Du_p|^2) d\mu_p + 2 \int_{\Omega} \xi (Du_p - Dv_\epsilon) : Du_p d\mu_p \\ &= \int_{\Omega} \xi (|Dv_\epsilon|^2 - |Du_p|^2) d\mu_p - 2 \int_{\Omega} (u_p - v_\epsilon) \otimes D\xi : Du_p d\mu_p \\ &\leq \int_{\Omega} \xi |Dv_\epsilon|^2 d\mu_p - \alpha^2 e_p^2 \int_{\Omega} \xi d\mu_p + \alpha^p e_p^2 - 2 \int_{\Omega} (u_p - v_\epsilon) \otimes D\xi : Du_p d\mu_p. \end{aligned}$$

Here we have used Lemma 18 in the last step. Hence

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_{\Omega} \xi |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} &\leq \int_{\Omega} \xi |Dv_\epsilon|^2 d\mu_\infty - \alpha^2 e_\infty^2 \int_{\Omega} \xi d\mu_\infty \\ &\quad - 2 \int_{\Omega} (u_\infty - v_\epsilon) \otimes D\xi : F_\infty d\mu_\infty. \end{aligned}$$

Let $r > 0$. Choose ξ such that $\xi \equiv 1$ in $\Omega \setminus \Omega_r$ and $\xi \equiv 0$ in $\Omega_{r/2}$, and such that $|D\xi| \leq 4/r$ in all of Ω . Note that

$$\|u_\infty - v\|_{C^0(\overline{\Omega_r})} \leq 2\sqrt{N}re_\infty,$$

because v is a minimiser of E_∞ and thus $\|Dv\|_{L^\infty(\Omega)} = \|Du_\infty\|_{L^\infty(\Omega)} = e_\infty$. If $\epsilon < r$, then there exists a constant C , depending only on N, n , and η , such that

$$\|u_\infty - v_\epsilon\|_{C^0(\overline{\Omega_r})} \leq Cre_\infty.$$

It follows that

$$-2 \int_{\Omega} (u_\infty - v_\epsilon) \otimes D\xi : F_\infty d\mu_\infty \leq 8Ce_\infty \left(\mu_\infty(\Omega_r) \int_{\Omega} |F_\infty|^2 d\mu_\infty \right)^{1/2}.$$

Given a compact set $K \subset \Omega$ and given $\gamma > 0$, we may choose $r > 0$ such that $K \cap \Omega_r = \emptyset$ and

$$8Ce_\infty \left(\mu_\infty(\Omega_r) \int_{\Omega} |F_\infty|^2 d\mu_\infty \right)^{1/2} \leq \gamma.$$

For the above choice of ξ , we then conclude that

$$\limsup_{k \rightarrow \infty} \int_{\Omega} \xi |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \leq \int_{\Omega} \xi |Dv_\epsilon|^2 d\mu_\infty - \alpha^2 e_\infty^2 \int_{\Omega} \xi d\mu_\infty + \gamma \tag{22}$$

for all $\epsilon \leq r$. Clearly $|Dv_\epsilon| \leq \|Dv\|_{L^\infty(\Omega)} \leq e_\infty$ everywhere in Ω . Therefore,

$$\limsup_{k \rightarrow \infty} \int_K |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \leq (1 - \alpha^2)e_\infty^2 \int_{\Omega} \xi d\mu_\infty + \gamma \leq (1 - \alpha^2)e_\infty^2 + \gamma. \tag{23}$$

By Proposition 15,

$$\int_K |Dv_\epsilon - F_\infty|^2 d\mu_\infty \leq (1 - \alpha^2)e_\infty^2 + \gamma$$

as well. Since $\alpha \in (0, 1)$ and $\gamma > 0$ may be chosen arbitrarily here (and then the inequality holds for ϵ small enough, depending on γ), it follows that

$$\lim_{\epsilon \searrow 0} \int_K |Dv_\epsilon - F_\infty|^2 d\mu_\infty = 0.$$

According to Definition 3, this means that $F_\infty = \langle Dv \rangle_{\mu_\infty}$.

Because $|Dv_\epsilon| \leq e_\infty$ everywhere, this locally strong L^2 -convergence with respect to μ_∞ also implies that $|F_\infty| \leq e_\infty$ at μ_∞ -almost every point. Now we note that (22) further gives rise to the inequality

$$\int_\Omega \xi |F_\infty|^2 d\mu_\infty = \lim_{\epsilon \searrow 0} \int_\Omega \xi |Dv_\epsilon|^2 d\mu_\infty \geq e_\infty^2 \int_\Omega \xi d\mu_\infty. \tag{24}$$

Hence $|F_\infty| = e_\infty$ at μ_∞ -almost every point.

Inequality (24), together with the dominated convergence theorem, also has the consequence that

$$\begin{aligned} \int_\Omega \xi (|Dv|^*)^2 d\mu_\infty &= \int_\Omega \xi \lim_{\epsilon \searrow 0} \operatorname{ess\,sup}_{B_\epsilon(x)} |Dv|^2 d\mu_\infty(x) \\ &= \lim_{\epsilon \searrow 0} \int_\Omega \xi \operatorname{ess\,sup}_{B_\epsilon(x)} |Dv|^2 d\mu_\infty(x) \\ &\geq \lim_{\epsilon \searrow 0} \int_\Omega \xi |Dv_\epsilon|^2 d\mu_\infty \\ &\geq e_\infty^2 \int_\Omega \xi d\mu_\infty. \end{aligned} \tag{25}$$

Since we clearly have the pointwise bound $|Dv|^* \leq e_\infty$, we conclude that $|Dv|^* = e_\infty$ at μ_∞ -almost every point. All the claims of statement (i) are now proved.

The claim of statement (iii) relies on the same inequalities. We have shown, as a consequence of (23), that for any compact set $K \subset \Omega$, if $\delta > 0$ is given, then there exists $\epsilon_0 > 0$ such that for any $\epsilon \in (0, \epsilon_0]$,

$$\limsup_{k \rightarrow \infty} \int_K |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \leq \delta. \tag{26}$$

But clearly, for a fixed ϵ ,

$$\begin{aligned} &\limsup_{k \rightarrow \infty} \int_K |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \\ &= \limsup_{k \rightarrow \infty} \int_K (|Dv_\epsilon|^2 - 2Dv_\epsilon : Du_{p_k} + |Du_{p_k}|^2) d\mu_{p_k} \\ &= \int_K (|Dv_\epsilon|^2 - 2Dv_\epsilon : F_\infty) d\mu_\infty + \limsup_{k \rightarrow \infty} \int_K |Du_{p_k}|^2 d\mu_{p_k} \\ &= \int_K |Dv_\epsilon - F_\infty|^2 d\mu_\infty - \int_K |F_\infty|^2 d\mu_\infty + \limsup_{k \rightarrow \infty} \int_K |Du_{p_k}|^2 d\mu_{p_k}. \end{aligned}$$

Therefore, the above inequality (26) implies that

$$\limsup_{k \rightarrow \infty} \int_K |Du_{p_k}|^2 d\mu_{p_k} \leq \int_K |F_\infty|^2 d\mu_\infty. \tag{27}$$

By Proposition 15, this means that $M_{p_k} \rightarrow M_\infty$ strongly in K .

Finally, we assume that $e'_\infty < e_\infty$, and we prove the global statements (ii) and (iv) with global variants of the above arguments. To this end, consider a regular mollifier \mathcal{M}_ϵ . We now define $v_\epsilon = \mathcal{M}_\epsilon v$. We choose $\alpha \in (0, 1)$ again. Just as before, we compute

$$\begin{aligned} & \int_\Omega |Dv_\epsilon - Du_p|^2 d\mu_p \\ &= \int_\Omega (|Dv_\epsilon|^2 - |Du_p|^2) d\mu_p + 2 \int_{\partial\Omega} (u_p - v_\epsilon) \cdot f_p dm_p \\ &\leq \int_\Omega |Dv_\epsilon|^2 d\mu_p - \alpha^2 e_p^2 \mu_p(\Omega) + \alpha^p e_p^2 + 2 \int_{\partial\Omega} (u_p - v_\epsilon) \cdot f_p dm_p. \end{aligned}$$

Hence

$$\begin{aligned} \limsup_{k \rightarrow \infty} \int_\Omega |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} &\leq \int_\Omega |Dv_\epsilon|^2 d\mu_\infty - \alpha^2 e_\infty^2 \mu_\infty(\bar{\Omega}) \\ &\quad + 2 \int_{\partial\Omega} (u_\infty - v_\epsilon) \cdot f_\infty dm_\infty. \end{aligned}$$

Fix $\gamma > 0$. We know that $v_\epsilon \rightarrow v$ uniformly in $\bar{\Omega}$, and $v = u_\infty$ on $\partial\Omega$. Hence

$$2 \int_{\partial\Omega} (u_\infty - v_\epsilon) \cdot f_\infty dm_\infty \leq \gamma$$

whenever ϵ is sufficiently small. Therefore,

$$\limsup_{k \rightarrow \infty} \int_\Omega |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \leq \int_\Omega |Dv_\epsilon|^2 d\mu_\infty - \alpha^2 e_\infty^2 \mu_\infty(\bar{\Omega}) + \gamma. \tag{28}$$

By the properties of regular mollifiers, we know that

$$\limsup_{\epsilon \searrow 0} \|Dv_\epsilon\|_{C^0(\bar{\Omega})} \leq \|Dv\|_{L^\infty(\Omega)} \leq e_\infty. \tag{29}$$

Hence

$$\limsup_{k \rightarrow \infty} \int_\Omega |Dv_\epsilon - Du_{p_k}|^2 d\mu_{p_k} \leq (1 - \alpha^2) e_\infty^2 \mu_\infty(\bar{\Omega}) + 2\gamma$$

when ϵ is small enough. Proposition 15 then implies that

$$\lim_{\epsilon \searrow 0} \int_\Omega |Dv_\epsilon - F_\infty|^2 d\mu_\infty = 0.$$

It follows that $F_\infty = \llbracket Dv \rrbracket_{\mu_\infty}$.

Using (29) again, we conclude that the inequality $|F_\infty| \leq e_\infty$ holds μ_∞ -almost everywhere in $\bar{\Omega}$. By (28),

$$\int_\Omega |F_\infty|^2 d\mu_\infty = \lim_{\epsilon \searrow 0} \int_\Omega |Dv_\epsilon|^2 d\mu_\infty \geq e_\infty^2 \mu_\infty(\bar{\Omega}).$$

Hence $|F_\infty| = e_\infty$ at μ_∞ -almost every point of $\bar{\Omega}$.

As in (25), we see that

$$\begin{aligned} \int_{\overline{\Omega}} (|Dv|^*)^2 d\mu_\infty &= \lim_{\epsilon \searrow 0} \int_{\overline{\Omega}} \operatorname{ess\,sup}_{B_\epsilon(x) \cap \overline{\Omega}} |Dv|^2 d\mu_\infty(x) \\ &\geq \lim_{\epsilon \searrow 0} \int_{\overline{\Omega}} |Dv_\epsilon|^2 d\mu_\infty \\ &= \lim_{\epsilon \searrow 0} \int_{\overline{\Omega}} |F_\infty|^2 d\mu_\infty \\ &= e_\infty^2 \mu_\infty(\overline{\Omega}) \end{aligned}$$

by the properties of \mathcal{M}_ϵ . Hence $|Dv|^* = e_\infty$ at μ_∞ -almost every point of $\overline{\Omega}$. This completes the proof of (ii).

For the proof of (iv), we observe that the derivation of (27) now works for $K = \overline{\Omega}$ as well. Thus it suffices to use Proposition 15 again. □

6 Combining the tools

In this section, we prove Theorem 4. If $e_\infty = 0$, then we can choose any unit vector $\vec{T}_0 \in \mathbb{R}^{N \times n}$ and set $T = [\mathcal{L}^n \llcorner \overline{\Omega}, \vec{T}_0]$. The statements of the theorem will then be satisfied trivially. We therefore assume that $e_\infty > 0$ henceforth.

The current T in the theorem is then just another representation of the measure-function pair $M_\infty = (\mu_\infty, F_\infty)$ constructed above. Namely, we set $T = [\mu_\infty, F_\infty]$. Since we know that $|F_\infty(x)| = e_\infty$ for μ_∞ -almost every $x \in \Omega$ by Theorem 17, this means that $\|T\| \llcorner \overline{\Omega} = e_\infty \mu_\infty \llcorner \overline{\Omega}$ and $\vec{T} = e_\infty^{-1} F_\infty$ in Ω . If $e'_\infty < e_\infty$, then $\|T\| = e_\infty \mu_\infty$ and $\vec{T} = e_\infty^{-1} F_\infty$ on $\overline{\Omega}$ by the same theorem.

Theorem 4 now follows from the results in the previous sections. We will give the details below. First, however, we formulate and prove a local version of the mass minimising property in statement (iv).(d) in Theorem 4. This result also holds true if $e'_\infty = e_\infty$.

Theorem 19 *Suppose that S is an N -tuple of 1-currents with locally finite mass and with $\operatorname{supp} \partial S \cap \Omega = \emptyset$. If $K \subset \Omega$ is a compact set such that $S(\omega) = T(\omega)$ for all $\omega \in (\mathcal{D}^1(\Omega))^N$ with $\omega = 0$ in K , then $\|T\|(K) \leq \|S\|(K)$.*

Proof We first note that $\langle Du_\infty \rangle_{\|T\|} = e_\infty \vec{T}$ by Theorem 17. According to Corollary 10, this implies that

$$-T(u_\infty d\chi) = e_\infty \int_{\Omega} \chi d\|T\|$$

for any $\chi \in C_0^\infty(\Omega)$ with $\chi \geq 0$. Hence the claim follows from Corollary 11. □

For the proof of Theorem 4, we also require some additional tools, including the following estimate.

Lemma 20 *Suppose that $e'_\infty < e_\infty$. Then there exist $R > 0$ and $\beta \in (0, 1)$ such that $\|T\|(\Omega_{r/2} \cup \partial\Omega) \leq \beta \|T\|(\Omega_r \cup \partial\Omega)$ for all $r \in (0, R)$.*

Proof Choose $R > 0$ such that there exists a smooth nearest point projection $\varpi : \Omega_{2R} \rightarrow \partial\Omega$. We may replace u_0 (while still using the same notation) by a function such that $u_0(x) =$

$u_0(\varpi(x))$ for $x \in \Omega_R$. Then

$$c := \operatorname{ess\,sup}_{\Omega_R} |Du_0| < e_\infty,$$

provided that R is sufficiently small.

Now fix $r \in (0, R]$. Choose $\xi \in C_0^\infty(\mathbb{R}^n)$ with $0 \leq \xi \leq 1$ and such that $\xi \equiv 1$ in $\Omega_{r/2}$ and $\xi \equiv 0$ in a neighbourhood of $\Omega \setminus \Omega_r$, and such that $|D\xi| \leq 4/r$. Then

$$\begin{aligned} & \int_{\Omega} \xi |Du_p|^p dx \\ &= \int_{\Omega} \xi |Du_p|^{p-2} Du_p : Du_0 dx + \int_{\Omega} \xi |Du_p|^{p-2} Du_p : (Du_p - Du_0) dx \\ &= \int_{\Omega} \xi |Du_p|^{p-2} Du_p : Du_0 dx - \int_{\Omega} |Du_p|^{p-2} (u_p - u_0) \otimes D\xi : Du_p dx. \end{aligned}$$

Moreover, given $\alpha \in (0, 1)$, we can estimate

$$\int_{\Omega} \xi |Du_p|^{p-2} Du_p : Du_0 dx \leq \frac{p-1}{p} \alpha^{\frac{p}{p-1}} \int_{\Omega} \xi |Du_p|^p dx + \frac{1}{p\alpha^p} \int_{\Omega} \xi |Du_0|^p dx$$

by Young’s inequality. Hence

$$\begin{aligned} & \left(1 - \frac{p-1}{p} \alpha^{\frac{p}{p-1}}\right) \int_{\Omega} \xi |Du_p|^p dx \\ & \leq \frac{1}{p\alpha^p} \int_{\Omega} \xi |Du_0|^p dx - \int_{\Omega} |Du_p|^{p-2} (u_p - u_0) \otimes D\xi : Du_p dx \\ & \leq \frac{c^p}{p\alpha^p} \int_{\Omega} \xi dx - \int_{\Omega} |Du_p|^{p-2} (u_p - u_0) \otimes D\xi : Du_p dx. \end{aligned}$$

In terms of the measures μ_p , this means that

$$\begin{aligned} & \left(1 - \frac{p-1}{p} \alpha^{\frac{p}{p-1}}\right) \int_{\Omega} \xi |Du_p|^2 d\mu_p \\ & \leq \frac{c^p e_p^{2-p}}{p\alpha^p} \int_{\Omega} \xi dx - \int_{\Omega} (u_p - u_0) \otimes D\xi : Du_p d\mu_p. \end{aligned}$$

Choose $\alpha > \frac{c}{e_\infty}$. Restricting to p_k and letting $k \rightarrow \infty$, we then find that

$$(1 - \alpha) \int_{\Omega} \xi |F_\infty|^2 d\mu_\infty \leq \int_{\Omega} (u_\infty - u_0) \otimes D\xi : F_\infty d\mu_\infty$$

by Proposition 15. Since $u_\infty - u_0 \in W_0^{1,\infty}(\Omega; \mathbb{R}^N)$, there exists a constant C_1 such that $|u_\infty - u_0| \leq C_1 r$ in Ω_r . Hence

$$\int_{\Omega_{r/2} \cup \partial\Omega} |F_\infty|^2 d\mu_\infty \leq C_2 e_\infty \int_{\Omega_r \setminus \Omega_{r/2}} |F_\infty| d\mu_\infty,$$

where $C_2 = 4C_1 / ((1 - \alpha)e_\infty)$. By Theorem 17 and by the definition of T , this means that

$$\|T\|(\Omega_{r/2} \cup \partial\Omega) \leq C_2 \|T\|(\Omega_r \setminus \Omega_{r/2}).$$

Adding $C_2\|T\|(\Omega_{r/2} \cup \partial\Omega)$ on both sides of the inequality, we conclude that

$$\|T\|(\Omega_{r/2} \cup \partial\Omega) \leq \frac{C_2}{C_2 + 1} \|T\|(\Omega_r \cup \partial\Omega).$$

Thus we have the desired inequality for $\beta = C_2/(C_2 + 1)$. □

We now have everything in place for the proof of our main theorem.

Proof of Theorem 4 Recall that we use the assumption $e_\infty > 0$, as the theorem is trivial otherwise. We have a measure-function pair $M_\infty = (\mu_\infty, F_\infty)$, constructed in Sect. 3, that satisfies statements (i) and (iii) of Theorems 17, and (ii) and (iv) as well if $e'_\infty < e_\infty$. As described at the beginning of this section, we set $T = [\mu_\infty, F_\infty]$. Then Theorem 17 implies that $\|T\| \llcorner \Omega = e_\infty \mu_\infty \llcorner \Omega$ and $\vec{T} = e_\infty^{-1} F_\infty$ in Ω . If $e'_\infty < e_\infty$, then $\|T\| = e_\infty \mu_\infty$ and $\vec{T} = e_\infty^{-1} F_\infty$ on $\bar{\Omega}$. We conclude that $T \neq 0$ in the second case.

We now prove the individual statements of Theorem 4.

(i) It is clear that $\text{supp } T \subseteq \bar{\Omega}$. For any $\sigma \in (\mathcal{D}^0(\mathbb{R}^n))^N$ with compact support in Ω , we note that

$$T(d\sigma) = \int_\Omega F_\infty : D\sigma \, d\mu_\infty = 0$$

by (12). Hence $\text{supp } \partial T \subseteq \partial\Omega$.

(ii) Both statements follow immediately from Theorem 17.

(iii) We note that identity (21) for the measure-function pair M_∞ corresponds to (4) for T . Therefore, if we have two minimisers $u, v \in u_0 + W_0^{1,\infty}(\Omega; \mathbb{R}^N)$ of E_∞ , then we may apply Proposition 12 to the function $u - v$. Since we have already proved that $\langle Du \rangle_{\|T\|} = e_\infty \vec{T} = \langle Dv \rangle_{\|T\|}$, it follows that $u = v$ on $\text{supp } T$.

(iv) Here we assume that $e'_\infty < e_\infty$. Then statement (a) is a consequence of Proposition 16, as it follows that

$$\partial T(\sigma) = \lim_{\ell \rightarrow \infty} \int_\Omega Du_{p_\ell} : D\sigma \, d\mu_{p_\ell} = \int_{\partial\Omega} f_{p_\ell} \cdot \sigma \, dm_{p_\ell} = \int_{\partial\Omega} f_\infty \cdot \sigma \, dm_\infty$$

for all $\sigma \in (\mathcal{D}^0(\mathbb{R}^n))^N$. Here (m_∞, f_∞) is the measure-function pair over $\partial\Omega$ found at the beginning of Sect. 5.

Statement (b) is an obvious consequence of Lemma 20 and the fact that $T \neq 0$.

For the proof of statement (c), we invoke Theorem 17. We conclude that $\langle Du \rangle_{\|T\|} = e_\infty \vec{T}$. Hence Proposition 8 implies that $\partial T(u_0) = e_\infty \mathbf{M}(T)$.

Finally, we prove (d). Recall that we have already proved the identity $\partial T(u_0) = e_\infty \mathbf{M}(T)$ when $e'_\infty < e_\infty$. Thus Corollary 9 implies the desired statement. □

7 The structure of T

In this final section we give some more results about the structure of the N -tuple of 1-currents T constructed above. These are based on the condition (4) and are closely related to standard results on varifolds. Most of the results in the literature, however, do not apply to T , as it is somewhat unusual: it should be thought of as a 1-dimensional object, because it acts on 1-forms, but need not actually be 1-dimensional in any other sense. In contrast, most results on currents or varifolds in the literature assume rectifiability in the appropriate dimension.

We use some more concepts from geometric measure theory in the following, including the notion of a countably rectifiable measure. A definition can be found, e.g., in a book by Mattila [32, Definition 16.6].

Theorem 21 *The measure $\|T\| \llcorner \Omega$ is absolutely continuous with respect to the one-dimensional Hausdorff measure \mathcal{H}^1 . Moreover, for any $\|T\|$ -measurable set $A \subseteq \Omega$ such that $\mathcal{H}^1(A) < \infty$, the restriction $\|T\| \llcorner A$ is a countably 1-rectifiable measure. At $\|T\|$ -almost every point $x \in A$, the approximate tangent space of $\|T\| \llcorner A$ contains $\{\vec{T}_1(x), \dots, \vec{T}_N(x)\}$.*

This means that the dimension of T , say in the sense of Hausdorff dimension for the measure $\|T\|$ as defined by Mattila, Morán, and Rey [33], is at least 1 (unless $\|T\|(\Omega) = 0$). It can, however, be higher. (An example is discussed in the introduction.) If we restrict $\|T\|$ to a 1-dimensional set A , then we have the structure typically assumed in geometric measure theory. We then also find that the vectors $\vec{T}_1(x), \dots, \vec{T}_N(x)$ are tangential. Since we have a 1-dimensional approximate tangent space here, this means that they are parallel to one another (or vanish) at $\|T\|$ -almost every point $x \in A$.

For the proof, we use a monotonicity identity that is a standard tool in the theory of varifolds (see [37, §17]). It is difficult, however, to find a formulation in the literature for anything more general than a rectifiable varifold, even though the standard arguments apply more generally. We therefore provide a proof for the convenience of the reader. As mentioned earlier, varifolds and currents are closely related, and we can formulate the identity in terms of T .

Lemma 22 *For any $x_0 \in \Omega$ and for $0 < s < r \leq \text{dist}(x_0, \partial\Omega)$, the identity*

$$\frac{\|T\|(B_r(x_0))}{r} - \frac{\|T\|(B_s(x_0))}{s} = \int_{B_r(x_0) \setminus B_s(x_0)} \frac{1 - \left| \frac{x-x_0}{|x-x_0|} \cdot \vec{T} \right|^2}{|x-x_0|} d\|T\|$$

holds true.

Proof We may assume without loss of generality that $x_0 = 0$.

Choose a non-increasing function $\xi \in C^\infty(\mathbb{R})$ with $\xi \equiv 1$ in $(-\infty, 0]$ and $\xi \equiv 0$ in $[1, \infty)$. For $\rho > 0$, define

$$\eta_\rho(x) = \xi(|x|/\rho).$$

Test (4) with $\psi_\rho(x) = \eta_\rho(x)x$. This gives

$$\int_\Omega \left(\eta_\rho(x) + \xi'(|x|/\rho) \frac{|x|}{\rho} \left| \frac{x}{|x|} \cdot \vec{T} \right|^2 \right) d\|T\|(x) = 0.$$

Now we compute

$$\begin{aligned} \frac{d}{d\rho} \left(\frac{1}{\rho} \int_\Omega \eta_\rho d\|T\| \right) &= -\frac{1}{\rho^2} \int_\Omega \left(\eta_\rho(x) + \frac{|x|}{\rho} \xi'(|x|/\rho) \right) d\|T\|(x) \\ &= -\frac{1}{\rho^3} \int_\Omega \xi'(|x|/\rho) |x| \left(1 - \left| \frac{x}{|x|} \cdot \vec{T} \right|^2 \right) d\|T\|(x). \end{aligned}$$

Integrate with respect to ρ over the interval (s, r) :

$$\begin{aligned} & \frac{1}{r} \int_{\Omega} \eta_r d\|T\| - \frac{1}{s} \int_{\Omega} \eta_s d\|T\| \\ &= - \int_s^r \frac{1}{\rho^3} \int_{\Omega} \xi'(|x|/\rho)|x| \left(1 - \left|\frac{x}{|x|} \cdot \vec{T}\right|^2\right) d\|T\|(x) d\rho \\ &= - \int_{\Omega} \int_s^r \frac{|x|\xi'(|x|/\rho)}{\rho^3} d\rho \left(1 - \left|\frac{x}{|x|} \cdot \vec{T}\right|^2\right) d\|T\|(x). \end{aligned}$$

Set

$$\Phi(x) = - \int_s^r \frac{|x|\xi'(|x|/\rho)}{\rho^3} d\rho,$$

so that

$$\frac{1}{r} \int_{\Omega} \eta_r d\|T\| - \frac{1}{s} \int_{\Omega} \eta_s d\|T\| = \int_{\Omega} \Phi(x) \left(1 - \left|\frac{x}{|x|} \cdot \vec{T}\right|^2\right) d\|T\|(x).$$

Note that

$$\Phi(x) = \int_s^r \frac{d}{d\rho} \xi(|x|/\rho) \frac{d\rho}{\rho} = \frac{\xi(|x|/r)}{r} - \frac{\xi(|x|/s)}{s} + \int_s^r \xi(|x|/\rho) \frac{d\rho}{\rho^2}.$$

Now let ξ approximate the characteristic function of $(-\infty, 1)$. Then $\Phi(x)$ converges to

$$r^{-1} + \int_{|x|}^r \frac{d\rho}{\rho^2} = |x|^{-1}$$

for $s \leq |x| < r$ and to 0 else. Using the dominated convergence theorem, we obtain the formula in the statement. □

Proof of Theorem 21 A consequence of Lemma 22 is that for any $x \in \Omega$, the function $r \mapsto r^{-1} \|T\|(B_r(x))$ is monotone. Therefore, the limit

$$\Theta(x) = \lim_{r \searrow 0} \frac{\|T\|(B_r(x))}{r}$$

exists. Lemma 22 also implies that Θ is locally bounded. It follows from standard results on Hausdorff measures [19, Section 2.10.19] that $\|T\|$ is absolutely continuous with respect to \mathcal{H}^1 .

Let $\Sigma = \{x \in \Omega : \Theta(x) > 0\}$. Moreover, for $\ell \in \mathbb{N}$, let $\Sigma_{\ell} = \{x \in \Omega : \Theta(x) \geq 1/\ell\}$. Then it also follows from standard results [19, Section 2.10.19] that

$$\lim_{r \searrow 0} \frac{1}{r} \|T\|(B_r(x) \setminus \Sigma_{\ell}) = 0$$

for almost every $x \in \Sigma_{\ell}$ with respect to \mathcal{H}^1 (and therefore with respect to $\|T\|$ as well). Hence

$$\lim_{r \searrow 0} \frac{1}{r} \|T\|(B_r(x) \cap \Sigma_{\ell}) = \Theta(x)$$

for $\|T\|$ -almost every $x \in \Sigma_{\ell}$. The results of Preiss [34] now imply that the measure $\|T\| \llcorner \Sigma_{\ell}$ is countably 1-rectifiable. Hence Σ is a countably 1-rectifiable set.

Given $x_0 \in \Sigma$, we now consider tangent measures of $\|T\|$ at x_0 . To this end, define $\Omega_r = \frac{1}{r}(\Omega - x_0)$ for $r > 0$. Consider the measures λ_r on Ω_r defined by

$$\int_{\Omega_r} \eta \, d\lambda_r = \frac{1}{r} \int_{\Omega} \eta((x - x_0)/r) \, d\|T\|(x)$$

for $\eta \in C_0^0(\Omega_r)$, and consider the functions $\vec{T}_r : \Omega_r \rightarrow \mathbb{R}^{N \times n}$ with

$$\vec{T}_r(x) = \vec{T}(rx + x_0).$$

Then for any $R > 0$,

$$\lambda_r(B_R(0)) = \frac{1}{r} \|T\|(B_{Rr}(x_0)) \rightarrow R\Theta(x_0),$$

while $|\vec{T}_r| = 1$ almost everywhere with respect to λ_r . We may therefore pick a sequence $r_\ell \searrow 0$ such that the measure-function pairs $(\lambda_{r_\ell}, \vec{T}_{r_\ell})$ converge weakly to some measure-function pair (λ_0, \vec{T}_0) over \mathbb{R}^n . If x_0 is such that Σ has an approximate tangent line $L_{x_0} \subseteq \mathbb{R}^n$ at x_0 (which is $\|T\|$ -almost everywhere [37, Theorem 11.6]) and \vec{T} is approximately continuous at x_0 with respect to $\|T\|$ (also $\|T\|$ -almost everywhere [19, Theorem 2.9.13]), then the limit will be locally strong and of the form

$$\lambda_0 = \frac{1}{2} \Theta(x_0) \mathcal{H}^1 \llcorner L_{x_0} \tag{30}$$

and

$$\vec{T}_0(0) = \vec{T}(x_0). \tag{31}$$

Now suppose that $0 < S < R$. Let $\zeta \in C_0^0(B_R(0) \setminus B_S(0); \mathbb{R}^{N \times n})$ such that $x \cdot \zeta_k(x) = 0$ everywhere for $k = 1, \dots, N$. Then we conclude that

$$\begin{aligned} & \int_{\mathbb{R}^n} |x|^{-1} (\zeta : \vec{T}_0)^2 \, d\lambda_0 \\ &= \lim_{\ell \rightarrow \infty} \int_{\mathbb{R}^n} |x|^{-1} (\zeta : \vec{T}_{r_\ell})^2 \, d\lambda_{r_\ell}(x) \\ &= \lim_{\ell \rightarrow \infty} \int_{\Omega} |x - x_0|^{-1} \left(\zeta((x - x_0)/r_\ell) : \vec{T}(x) \right)^2 \, d\|T\|(x) \\ &\leq \|\zeta\|_{L^\infty(\mathbb{R}^n)}^2 \lim_{\ell \rightarrow \infty} \int_{B_{Rr_\ell}(x_0) \setminus B_{Sr_\ell}(x_0)} \frac{1 - \left| \frac{x-x_0}{|x-x_0|} \cdot \vec{T}(x) \right|^2}{|x - x_0|} \, d\|T\|(x) \\ &= \|\zeta\|_{L^\infty(\mathbb{R}^n)}^2 \lim_{\ell \rightarrow \infty} \left(\frac{1}{Rr_\ell} \|T\|(B_{Rr_\ell}(x_0)) - \frac{1}{Sr_\ell} \|T\|(B_{Sr_\ell}(x_0)) \right) \end{aligned}$$

by Lemma 22. As

$$\lim_{\ell \rightarrow \infty} \frac{\|T\|(B_{Rr_\ell}(x_0))}{Rr_\ell} = \Theta(x_0) = \frac{\|T\|(B_{Sr_\ell}(x_0))}{Sr_\ell},$$

we conclude that

$$\int_{\mathbb{R}^n} |x|^{-1} (\zeta : T_0)^2 \, d\lambda_0 = 0$$

for any ζ with the above properties. If x_0 is such that (30) and (31) hold true, then this means that $\vec{T}_k(x_0) \in L_{x_0}$ for $k = 1, \dots, N$. Recall that this is true for $\|T\|$ -almost every $x_0 \in \Sigma$.

Now the claims of the theorem follow almost immediately. We have already seen that $\|T\|$ is absolutely continuous with respect to \mathcal{H}^1 . For a set A as in the statement, we conclude that $\|T\|(A \setminus \Sigma) = 0$ [19, Section 2.10.19], so we may assume that $A \subseteq \Sigma$. The remaining statements then follow from what we know about Σ . \square

Acknowledgements We wish to thank A. Backus for his comments on a previous version of this paper.

Funding This work was partially supported by the Engineering and Physical Sciences Research Council (grant numbers EP/X017109/1 and EP/X017206/1).

Data availability Data sharing is not applicable to this article as no datasets were generated or analysed during the current study.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Ethics approval Ethics approval was not required for this study.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Allard, W.K.: On the first variation of a varifold. *Ann. Math.* **95**, 417–491 (1972)
- Aronsson, G.: Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$. *Ark. Mat.* **6**, 33–53 (1965)
- Aronsson, G.: Minimization problems for the functional $\sup_x F(x, f(x), f'(x))$. (II). *Ark. Mat.* **6**(4–5), 409–431 (1966). <https://doi.org/10.1007/BF02590964>
- Aronsson, G.: Extension of functions satisfying Lipschitz conditions. *Ark. Mat.* **6**, 551–561 (1967)
- Aronsson, G.: On the partial differential equation $u_x^2 u_{xx} + 2u_x u_y u_{xy} + u_y^2 u_{yy} = 0$. *Ark. Mat.* **7**(1968), 395–425 (1968)
- Aronsson, G.: On certain singular solutions of the partial differential equation $u_x^2 u_{xx} + 2u_x u_y u_{xy} + u_y^2 u_{yy} = 0$. *Manuscripta Math.* **47**, 133–151 (1984)
- Backus, A.: An ∞ -Laplacian for differential forms, and calibrated laminations, (2024). [arXiv:2404.02215](https://arxiv.org/abs/2404.02215) [math.AP]
- Bhattacharya, T., DiBenedetto, E., Manfredi, J.: Limits as $p \rightarrow \infty$ of $\Delta_p u_p = f$ and related extremal problems, 1989, Some topics in nonlinear PDEs (Turin, 1989), pp. 15–68 (1991)
- Brizzi, C.: On functions with given boundary data and convex constraints on the gradient, (2022). [arXiv:2209.01462](https://arxiv.org/abs/2209.01462) [math.AP]
- Brizzi, C., De Pascale, L.: A property of absolute minimizers in L^∞ calculus of variations and of solutions of the Aronsson-Euler equation. *Adv. Differ. Equ.* **28**, 287–310 (2023)
- Bungert, L., Korolev, Y.: Eigenvalue problems in L^∞ : optimality conditions, duality, and relations with optimal transport. *Comm. Amer. Math. Soc.* **2**, 345–373 (2022)
- Champion, T., De Pascale, L., Jimenez, C.: The ∞ -eigenvalue problem and a problem of optimal transportation. *Commun. Appl. Anal.* **13**, 547–565 (2009)
- Daskalopoulos, G., Uhlenbeck, K.: Transverse measures and best Lipschitz and least gradient maps, (2022). [arXiv:2010.06551](https://arxiv.org/abs/2010.06551) [math.DG]

14. Dong, H., Peng, F., Zhang, Y.R.-Y., Zhou, Y.: Jacobian determinants for nonlinear gradient of planar ∞ -harmonic functions and applications. *J. Reine Angew. Math.* **812**, 59–98 (2024)
15. Evans, L.C.: Three singular variational problems. In: *Viscosity Solutions of Differential Equations and Related Topics*, vol. 1323. Research Institute for the Mathematical Sciences, RIMS Kokyuroku (2003)
16. Evans, L.C., Savin, O.: $C^{1,\alpha}$ regularity for infinity harmonic functions in two dimensions. *Calc. Var. Partial Differ. Equ.* **32**, 325–347 (2008)
17. Evans, L.C., Smart, C.K.: Everywhere differentiability of infinity harmonic functions. *Calc. Var. Partial Differ. Equ.* **42**, 289–299 (2011)
18. Evans, L.C., Yu, Y.: Various properties of solutions of the infinity-Laplacian equation. *Comm. Partial Differ. Equ.* **30**, 1401–1428 (2005)
19. Federer, H.: *Geometric measure theory*. Springer, New York (1969)
20. Hutchinson, J.E.: Second fundamental form for varifolds and the existence of surfaces minimising curvature. *Indiana Univ. Math. J.* **35**, 45–71 (1986)
21. Jensen, R.: Uniqueness of Lipschitz extensions: minimizing the sup norm of the gradient. *Arch. Rational Mech. Anal.* **123**, 51–74 (1993)
22. Katzourakis, N.: L^∞ variational problems for maps and the Aronsson PDE system. *J. Differ. Equ.* **253**, 2123–2139 (2012)
23. Katzourakis, N.: Explicit 2D ∞ -harmonic maps whose interfaces have junctions and corners. *C. R. Math. Acad. Sci. Paris* **351**, 677–680 (2013)
24. Katzourakis, N.: ∞ -minimal submanifolds. *Proc. Amer. Math. Soc.* **142**, 2797–2811 (2014)
25. Katzourakis, N.: On the structure of ∞ -harmonic maps. *Comm. Partial Differ. Equ.* **39**, 2091–2124 (2014)
26. Katzourakis, N.: Nonuniqueness in vector-valued calculus of variations in L^∞ and some linear elliptic systems. *Commun. Pure Appl. Anal.* **14**, 313–327 (2015)
27. Katzourakis, N.: A characterisation of ∞ -harmonic and p -harmonic maps via affine variations in L^∞ . *Electron. J. Differ. Equ.* (2017)
28. Katzourakis, N.: Generalised vectorial ∞ -eigenvalue nonlinear problems for L^∞ functionals. *Nonlinear Anal.* **219**, 112806 (2022)
29. Katzourakis, N., Moser, R.: Variational problems in L^∞ involving semilinear second order differential operators, (2023). [arXiv:2303.15982](https://arxiv.org/abs/2303.15982) [math.AP]
30. Koch, H., Zhang, Y.R.-Y., Zhou, Y.: An asymptotic sharp Sobolev regularity for planar infinity harmonic functions. *J. Math. Pures Appl.* **132**, 457–482 (2019)
31. Kristensen, J., Mingione, G.: Boundary regularity in variational problems. *Arch. Ration. Mech. Anal.* **198**, 369–455 (2010)
32. Mattila, P.: *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*. Cambridge University Press (1995). <https://doi.org/10.1017/CBO9780511623813>
33. Mattila, P., Morán, M., Rey, J.-M.: Dimension of a measure. *Studia Math.* **142**, 219–233 (2000)
34. Preiss, D.: Geometry of measures in \mathbf{R}^n : distribution, rectifiability, and densities. *Ann. Math.* **125**(3), 537–643 (1987)
35. Savin, O.: C^1 regularity for infinity harmonic functions in two dimensions. *Arch. Ration. Mech. Anal.* **176**(3), 351–361 (2005)
36. Sheffield, S., Smart, C.K.: Vector-valued optimal Lipschitz extensions. *Comm. Pure Appl. Math.* **65**, 128–154 (2012)
37. Simon, L.: *Lectures on geometric measure theory*. Australian National University Centre for Mathematical Analysis, Canberra (1983)
38. Thurston, W. P.: Minimal stretch maps between hyperbolic surfaces, (1998). [arXiv:math/9801039](https://arxiv.org/abs/math/9801039) [math.GT]
39. Uhlenbeck, K.: Regularity for a class of non-linear elliptic systems. *Acta Math.* **138**, 219–240 (1977)