

Legacy of warming and cover crops on the response of soil microbial function to repeated drying and rewetting cycles

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Adekanmbi, A. A. ORCID: <https://orcid.org/0000-0003-3379-4161>, Zou, Y., Shu, X., Pietramellara, G., Pathan, S. I., Todman, L. ORCID: <https://orcid.org/0000-0003-1232-294X> and Sizmur, T. ORCID: <https://orcid.org/0000-0001-9835-7195> (2025) Legacy of warming and cover crops on the response of soil microbial function to repeated drying and rewetting cycles. *European Journal of Soil Science*, 76 (1). e70044. ISSN 1351-0754 doi: 10.1111/ejss.70044 Available at <https://centaur.reading.ac.uk/120204/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1111/ejss.70044>

Publisher: Wiley

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur



CentAUR

Central Archive at the University of Reading

Reading's research outputs online

RESEARCH ARTICLE OPEN ACCESS

Legacy of Warming and Cover Crops on the Response of Soil Microbial Function to Repeated Drying and Rewetting Cycles

Adetunji Alex Adekanmbi^{1,2}  | Yiran Zou¹ | Xin Shu¹ | Giacomo Pietramellara³ | Shamina Imran Pathan³ | Lindsay Todman⁴ | Tom Sizmur¹ 

¹Department of Geography and Environmental Science, University of Reading, Berkshire, UK | ²Department of Soil Science and Land Management, Federal University of Technology, Minna, Nigeria | ³Department of Agriculture, Food, Environment and Forestry, University of Florence, Firenze, Italy | ⁴Department of Sustainable Land Management, University of Reading, Berkshire, UK

Correspondence: Tom Sizmur (t.sizmur@reading.ac.uk)

Received: 7 December 2023 | **Revised:** 7 August 2024 | **Accepted:** 3 January 2025

Funding: This work was supported by the Biotechnology and Biological Sciences Research Council, BB/R006989/1.

Keywords: adaptation | climate change | cover crops | drought | perturbation | resistance | soil respiration | stress | warming

ABSTRACT

The response of soils to extreme weather events will become increasingly important in the future as more frequent and severe floods and droughts are expected to subject soils to drying and rewetting cycles as a result of climate change. These extreme events will be experienced against a backdrop of overall warming. Farmers are adopting cover cropping as a sustainable management practice to increase soil organic matter and benefit soil health. Cover crops may also increase the resilience of soils to help mitigate the impacts of climate change. We examined the legacy of warming and cover crops on the response of soil microbial function to repeated drying and rewetting cycles. We introduced open-top chambers to warm the soil surface of a field plot experiment in which cover crops (single-species monocultures and 4-species polycultures) were grown over the summer after harvest and before planting autumn sown cash crops in a cereal rotation. Soil samples were collected from warmed and ambient areas of the experimental plots in spring, before harvesting the cereal crop. Warming significantly increased, and cover crops significantly decreased, the abundance of genes encoding fungal β -glucosidase. We quantified respiration (a measure of soil microbial function) with high-frequency CO_2 flux measurements after 0, 1, 2, 4 or 8 wet/dry cycles imposed in the laboratory and the addition of barley grass powder substrate at a rate of 10 mg g^{-1} soil. We observed lower cumulative substrate-induced respiration in soils previously planted with cover crop mixtures than expected from the average of the same species grown in monoculture. Repeated drying and rewetting cycles increased the cumulative substrate-induced respiration rate observed, suggesting that repeated perturbations selected for a community adapted to processing the barley shoot powder more quickly. When we calculated the cumulative respiration after 8 wet/dry cycles, relative to cumulative respiration after 0 wet/dry cycles (which we infer represents the extent to which microbial communities adapted to repeated drying and rewetting cycles), our data revealed that the legacy of warming significantly reduced soil microbial community adaptation, but the legacy of cover crops significantly increased, soil microbial community adaptation. This adaptation of the soil microbial community was positively correlated with the concentration of water-extractable organic carbon in the soils before imposing the drying and rewetting cycles and/or adding the substrate. We conclude that cover crops may enhance the ability of the soil microbial community to adapt to drought events and mitigate the impact of warming, possibly due to the provision of labile organic carbon for the synthesis of osmolytes which then prime the decomposition of labile plant material upon rewetting.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2025 The Author(s). *European Journal of Soil Science* published by John Wiley & Sons Ltd on behalf of British Society of Soil Science.

Summary

- We observed the legacy of warming and cover crops on substrate-induced respiration after wet/dry cycles.
- Repeated drying and rewetting cycles increased the cumulative substrate-induced respiration rate observed.
- Warming reduced, but cover crops increased, microbial community adaptation to wet/dry cycles.
- Cover crops enhance soil microbial community adaptation to drought and mitigate warming.

1 | Introduction

Global warming of up to 6.4°C is expected during the 21st century unless mitigation measures are implemented (Carey et al. 2018; Li et al. 2022). Warming could increase soil respiration, and increase CO₂ flux from soils to the atmosphere, resulting in positive feedback that results in further warming (Rustad et al. 2001; Bardgett, Freeman, and Ostle 2008; Dutta and Dutta 2016). Field experiments that elevate soil temperatures by 0.3°C–6°C in various ecosystems have consistently shown increases in soil respiration (Rustad et al. 2001; Kuffner et al. 2012). Winter warming has been observed to outpace summer warming in the northern hemisphere, particularly in the United Kingdom (Vogelsang and Franses 2005). Kreyling et al. (2019) showed that warming of soil by up to 1.7°C from October to March affected several ecological processes, including plant performance, soil respiration, and soil biological and chemical properties. A rise in temperatures between 0.5°C and 2.0°C has also resulted in increases in soil pH and available phosphorus, but reduced phosphatase, catalase, and urease activities (Guoju et al. 2012). Low-level winter warming (a surface temperature increase of 0.69°C) was observed to improve the availability of soil carbon and nitrogen and increase the soil microbial biomass and greenhouse gas emissions (Liu et al. 2023). Since warming influences soil microbial communities and their emissions, further understanding of these processes may enhance our ability to predict the impact of climate change on soil respiration (Kreyling 2010; Kreyling et al. 2019) and ecological processes.

As a result of the overall increase in global temperatures, a greater frequency and severity of extreme weather events is being experienced globally (Rahmstorf and Coumou 2011). In the United Kingdom and much of Europe, this manifests as a greater incidence of drought (Lavalle et al. 2009), causing concern about the resilience of soils and crop yields to perturbations caused by extreme weather events (Harkness et al. 2020). Therefore, our agroecosystems must be resilient to extreme events as well as higher temperatures. The ability of soil microbial communities to withstand and recover from perturbations and disturbances (including drought) is vital to ensure their continued ability to deliver multiple ecosystem services (Philippot, Griffiths, and Langenheder 2021). Long-term strategies are required, in particular, to increase agricultural resilience to drought in temperate environments (Holman et al. 2021). When soil microorganisms are exposed to drought conditions, they increase their internal

solute potential by synthesising low molecular weight compounds and osmolytes, which help to maintain homeostasis and turgor (Schimel 2018). Upon re-wetting, a pulse of CO₂ is often observed (Zhang et al. 2020), the origins of which have been much debated in the literature (Barnard, Blazewicz, and Firestone 2020). However, it is proposed that this flush may, in part, be because of the metabolism of osmolytes which primes the decomposition of other substrates (Warren 2016). It is possible that repeated exposure of soil microbial communities to drought conditions may result in adaptation of the microbial community to better maintain soil function after drought (Allison 2023), in much the same way that we observe thermal adaptation of soil microbial communities to warming (Bradford 2013).

Cover crops are increasingly used in European cropping systems to enhance soil quality, soil biodiversity and reduce CO₂ emissions (Papp et al. 2018; Radicetti et al. 2019). Such cover crops are planted as a subsidiary crop to a cash crop to enhance the overall conditions of the soil and it has been suggested that cover crops can help mitigate the impacts of climate change (Kaye and Quemada 2017). Cover crops can enhance soil microbial biomass through carbon supplied by root exudates, shoots and roots (Gyssels et al. 2005; Paterson et al. 2007; Calderón et al. 2016; Papp et al. 2018). Growing mixtures of cover crops, rather than single species monocultures, can result in greater increases in soil microbial functional diversity (Drost et al. 2020) and biomass (Shu et al. 2022) because of the provision of more diverse substrates. However, it remains to be investigated whether cover crop mixtures (or monocultures) can increase the resistance and resilience of soil microbial communities to the effects of global warming, or extreme weather events.

In this study, we were broadly interested in whether cover crops mitigated the impact of winter warming on a soil's resilience to wet/dry cycles. Specifically, we investigated whether (i) repeated wet/dry cycles increase or decrease microbial substrate-induced respiration by adapting to these repeated perturbations, (ii) the legacy of winter warming influenced this adaptation of the microbial community, (iii) the legacy of cover crops influenced this adaptation of the microbial community, and (iv) the legacy of cover crops mitigates the legacy of winter warming in terms of the soil's capacity to adapt to wet/dry cycles. To answer these questions we introduced Open Top Chambers (OTCs) to warm the soil surface of an ongoing field plot experiment in which cover crops were grown as single-species monocultures or 4-species polycultures over the summer in between autumn sown cash crops in a cereal rotation. We compared warmed and ambient plots where cover crops had been grown in monoculture, polyculture, or absent. We measured soil respiration using a high-frequency respirometer after subjecting the soil to repeated (0, 1, 2, 4 or 8) drying and rewetting cycles and then applying a substrate of fresh barley shoot powder, following the assay described by Todman et al. (2018). We hypothesized that the legacy effect of warming would suppress substrate-induced respiration after eight repeated drying and rewetting cycles, relative to 0 cycles. We also hypothesized that cover crops would mitigate the impact of warming by supporting soil microbial activity and that this effect would be more prevalent when cover crops were grown in polyculture than monoculture.

2 | Materials and Methods

2.1 | Field Experiment Design

The field plot experiment was located on the University of Reading farm at Sonning in Berkshire, UK, and was established in August 2018. Each summer, after the harvest of a cereal crop, cover crops were planted and then terminated and incorporated before planting the next autumn sown cereal crop. The cereal rotation was Winter Wheat (2017/2018); Winter Barley (2018/2019); Winter Oats (2019/2020) and Winter Wheat (2020/2021). The cover crop treatments were control (no cover crops), buckwheat (*Fagopyrum esculentum*), berseem clover (*Trifolium alexandrinum*), oil radish (*Raphanus raphanistrum*), sunflower (*Helianthus annuus*) and a 4-species mixture of these four. The experiment was arranged in a randomised complete block design with four blocks. The field experiment included additional treatments not described here but this study focused on the 24 plots where cover crops were grown, and the residues incorporated into the soil. The selected plots are circled on the diagram shown in Figure S1.

2.2 | Passive Warming

Open Top Chambers (OTCs) were installed in identical locations after cereal crop establishment during the 2019/2020 and 2020/2021 growing seasons of the field experiment to act as passive warming devices by warming the soil under the OTCs. The OTCs were installed on 18 December 2019 and 24 November 2020, respectively, at the crop seedling stage when about 5 leaves were unfolded. Each chamber was placed on the southeast end of each of the 24 plots to allow yield measurements to be made on the rest of the plot. Each OTC was a hexagon made from clear extruded Perspex acrylic plates with the following dimensions: 5 mm thickness with a 100 cm base, 57.74 cm top, 62 cm side cut at an angle 71.16° and each side was 50 cm high (Figure S2). The diameter of the base of the chamber at its shortest orientation is 1.73 m. Our design for the OTC chamber conforms to the characteristics of the International Tundra Experiment (ITEX), with a similar shape and reinforcement as the hexagon chamber described by Marion et al. (1997). Clear Perspex acrylic sheets can transmit in excess of 92% of visible light and have higher light transmission capacity than glass. We monitored the effect of chambers on the temperature within the top 10 cm of soil by inserting temperature probes coupled to Plus 2 Tinytag data loggers (Gemini, UK) in the centre of a chamber (warmed treatment) and at the equivalent location at the other end of the same plot (ambient treatment) of a control plot. The loggers were set to log temperature every 15 min while the OTCs were in place. The OTCs warmed the soil by, on average, $0.75^{\circ}\text{C} \pm 0.92^{\circ}\text{C}$ in the control plot in 2020/2021. This is broadly similar to the warming of $0.5^{\circ}\text{C} - 2.0^{\circ}\text{C}$ observed by Guoju et al. (2012) and greater than the warming of 0.3°C observed by Whitehead et al. (1995). The difference in soil temperature under the OTCs, compared with the ambient measurements is shown in Figure S4.

2.3 | Soil Sampling

We sampled soils on 17 May 2021 from the top 10 cm underneath the OTSs (warmed treatment), and the equivalent location

at the other end (ambient treatment) of each plot circled in Figure S1 using a trowel. The soil samples were sieved, fresh, to 4 mm and divided into four subsamples. The first subsample was refrigerated at 4°C before KCl extraction to determine NH_4^+ and NO_3^- availability. The second subsample was used for DNA extraction, and quantification of gene abundance of microbial communities using real-time qPCR. The third subsample was air-dried for soil chemical analysis. The fourth subsample was used to assess the response of soil microbial respiration following repeated drying and rewetting cycles and the addition of barley shoot powder as a substrate.

2.4 | Soil Chemical Analysis

Soil pH was determined by shaking soil samples with deionised water (1:2.5 mass/volume ratio) for 10 min and leaving the mixture to stand for 2 min before pH was measured using a digital type DMP-2 mV/pH meter (Thermo Orion). Total N and C concentrations were determined using a C/N Elemental Analyser (Thermo Flash 2000 EA). The C/N ratio was then calculated from total C and N. NH_4^+ and NO_3^- were extracted in 1 M KCl and then analysed using a Continuous Flow Analyser (San++ Automated Wet Chemistry Analyser—SKALAR). Available N was calculated from the sum of extractable NH_4^+ and NO_3^- . Moisture content and loss on ignition were determined by weight loss at 105°C and 500°C, respectively. Hot water extractable carbon (HWEOC) and cold water extractable carbon (CWEOC) were analysed as described by Ghani, Dexter, and Perrott (2003). Approximately 3 g air-dried soil samples of known moisture content were accurately weighed into 50 mL polypropylene centrifuge tubes. Thirty millilitres of ultra-pure water was then added to each tube before mixing on a rotary shaker at 30 rpm for 30 min at 20°C. This was followed by centrifuging at 3500 rpm for 20 min at 20°C. The supernatants were then entirely removed using polypropylene syringes and passed through 0.2 µm cellulose nitrate membrane filters into polypropylene universal tubes, discarding the first 3 mL of the filtrate each time. A further 30 mL of ultra-pure water was then added to each centrifuge tube before vortexing for 10 s and leaving in an 80°C water bath overnight and the supernatants were removed, as described above. Both supernatants were analysed for CWEOC and HWEOC, respectively, using a Shimadzu TOC analyser.

2.5 | DNA Extraction and qPCR

Total soil DNA was extracted from 230 mg of soil using the DNeasy PowerSoil kit (Qiagen). Genomic DNA concentrations were estimated by a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific, USA) and the yield of soil-extracted DNA was checked by 1% agarose gel electrophoresis and quality control. The abundance of total bacterial 16S rRNA and total fungal ITS genes as well as bacterial and fungal beta-glucosidases genes were quantified using real-time PCR technique. Genes that encode beta-glucosidase were selected since beta-glucosidase is a major class of extracellular enzymes responsible for the hydrolysis of cellobiose to glucose, plays an important role in the decomposition of plant material in soils and is a key indicator of soil quality. Specific primers were used to amplify each targeted gene sequence. Primer sets and references reporting the cycling conditions are shown in Table S1. In addition, the melting curve conditions were

performed from 55°C to 95°C, with increments of 0.5°C every 5 s. Each 25 µL of PCR reaction contained 40 ng of the DNA, 400 nM of each primer and 12.5 µL 2×ITAQ SYBER Green Supermix (Bio-Rad, Munich, Germany). Standard curves were constructed using a purified PCR product of known concentration from pure culture, *Bacillus subtilis* and *Trichoderma reesei* for eubacteria (16S rRNA genes) and fungi (ITS genes) respectively. Standard curves for fungal and bacterial beta-glucosidases for GH1 (glycoside hydrolase family 1) and GH3 (glycoside hydrolase family 3) were constructed using purified PCR products of known concentration of common DNA mixture (equal amount of DNA from all samples collected in this study). The concentration of each purified product was measured by Picodrop Microliter UV/Vis Spectrophotometer. Ten-fold dilutions ranging from 10⁰ to 10⁻⁷ ng per 1 µL⁻¹ were applied for the standard curve construction. All standards and samples were run in triplicate and duplicate, respectively. Quantification was performed on a CFX Connect™ Real-Time PCR detection System using CFX Manager™ software 3.1 (Bio-Rad, Munich, Germany). Further, the quantified DNA concentration (ng g⁻¹ soil) of each sample was converted into units of copy number per g soil according to the SC method (Brankatschk et al. 2012).

2.6 | Imposition of Drying and Rewetting Cycles and Substrate Induced Respiration

The water-holding capacity of each soil sample was determined and soils were adjusted to 85% of their water-holding capacity (30.3% soil moisture) and pre-incubated at ~22°C for 1 week before the drying and rewetting assay to avoid artefacts caused by the soil preparation. The assay is based on the approach described by Fraser, Todman, et al. (2016) and Todman et al. (2018). To undertake the assay, the equivalent of 10 g (dry weight) of six fresh subsamples of each soil sample was weighed into tin trays (before pre-incubation). To one subsample (0 wet/dry cycles), 100 mg of barley shoot powder (C/N ratio 23.3) was added and mixed, following Kuan et al. (2007), to simulate the addition of fresh plant material to the soil. After mixing, CO₂ evolution was measured continuously for 96 h at hourly time intervals using an automated multichannel respirometer and an EGA60 multi-sample gas exchange system (ADC Bioscientific Ltd). Samples that had not received substrate were exposed to 1, 2, 4 or 8 drying and rewetting cycles before the addition of barley shoot powder and substrate-induced respiration being measured. Each drying and rewetting cycle consisted of 3 days of drying by enclosing samples in a sealed chamber with silica gel desiccant, followed by weighing and rewetting to 85% of the water holding capacity where they were maintained for a further 4 days, following Todman et al. (2018), as depicted in Figure S5. A preliminary trial indicated that these drying and rewetting cycles caused the soil moisture content to fluctuate between 10.9% and 30.3%. It is important to note that the barley shoot powder was only added once to each sub-sample, so the assay measured the CO₂ flux upon the addition of a novel substrate after being exposed to 0, 1, 2, 4 or 8 wet/dry cycles (Figure 1). CO₂ respired in (ppm) was converted to C—CO₂ (µg g⁻¹ soil h⁻¹) and the cumulative sum (µg g⁻¹ soil) of C—CO₂ respired per sample over 96 h was calculated. We then quantified cumulative respiration after 8 wet/dry cycles as a percentage of the cumulative respiration of a sub-sample from the same plot after 0 wet/dry cycles to

represent the extent to which the microbial community adapted to the drying and rewetting cycles.

2.7 | Statistical Analysis

The effect of experimental treatments (passive warming and cover crops) on cumulative respiration (and the percentage of cumulative respiration after 8 wet/dry cycles relative to 0 wet/dry cycles) was analysed using the following ANOVA model: Warming*Cover crop/(Mix/Type) in GenStat (2021 version). The *Warming* factor had two levels (warmed and ambient). The *Cover crop* factor also had two levels (control and cover crops). The *Mix* factor had two levels (monoculture and 4-species mixture) with a dummy level representing the control treatment. The *Type* factor had four levels (buckwheat, berseem clover, oil radish and sunflower) with dummy levels representing the control and mixture treatments. Treatments with *p* < 0.05 were considered statistically significant. The relationships between the percentage of cumulative respiration after 8 wet/dry cycles relative to 0 wet/dry cycles and soil chemical and biological properties were analysed using Pearson's correlation. The extent to which soil chemical and biological properties predict the cumulative respiration after 8 wet/dry cycles relative to 0 wet/dry cycles was explored using stepwise multivariate regression (stepwise selection of terms was based on *p* < 0.15).

3 | Results

3.1 | Differences in Soil Properties Due to Warming and Cover Crops

There were no significant effects of *Warming*, *Cover Crop*, *Mix* or *Type* on HWEOC, CWEOC, %LOI, extractable NO₃⁻, C/N ratio, or abundance of total bacteria (16S rRNA gene), total fungi (ITS region gene), or genes encoding a bacterial β-glucosidase enzyme (GH1) in soils (Table 1). However, the presence of cover crops significantly decreased soil pH (Figure S6), %C (Figure S7) and the abundance of genes encoding fungal β-glucosidase enzymes (GH3) (Figure S11). There were no significant differences in soil properties between plots with the cover crop mixture and the average of the four monoculture plots (*Mix*; Table 1). However, cover crop species had a significant effect on %N and the abundance of genes encoding both fungal and bacterial β-glucosidase enzyme (GH3). Soils from plots planted with clover had the greatest %N (Figure S8) while radish plots had the highest abundance of the bacterial GH3 gene (Figure S9) and the lowest abundance of the fungal GH3 gene (Figure S11). Warming significantly increased the abundance of genes encoding fungal β-glucosidase (GH1) (Table 1 and Figure S10).

3.2 | Substrate Induced Respiration

When data from each drying and rewetting cycle was statistically analysed on its own, there was no significant effect of *Warming* or *Cover crops* on cumulative substrate-induced respiration. However, cumulative substrate-induced respiration was




FIGURE 1 | Schematic of the experimental design and procedure and average CO₂ concentrations measured during the 96 h after barley shoot powder addition plotted as an average of all 48 experimental treatments.

TABLE 1 | Summary table showing *F* values (with *p* values in parenthesis) for nested ANOVA testing the effect of statistical factors (*Warming*, *Cover crops*, *Mix* and *Type*) on soil properties.

Parameters	Warming	Cover crops	Mix	Type
HWEOC ($\mu\text{g g}^{-1}$)	1.50 (0.230)	0.10 (0.760)	1.45 (0.237)	0.67 (0.577)
CWEOC ($\mu\text{g g}^{-1}$)	0.99 (0.327)	0.15 (0.699)	0.58 (0.451)	2.15 (0.114)
Soil pH	0.02 (0.879)	4.39 (0.044)	0.00 (0.997)	1.25 (0.307)
% LOI	0.06 (0.811)	0.87 (0.359)	0.01 (0.922)	0.71 (0.555)
Extractable NO ₃ ⁻ (mg kg^{-1})	0.45 (0.508)	0.30 (0.586)	0.23 (0.632)	0.39 (0.764)
Total extractable N (mg kg^{-1})	0.68 (0.414)	0.06 (0.814)	0.14 (0.710)	0.22 (0.882)
%N	0.13 (0.720)	0.07 (0.790)	0.02 (0.897)	3.37 (0.030)
%C	2.45 (0.128)	4.38 (0.044)	0.13 (0.724)	1.17 (0.336)
C/N	0.60 (0.446)	1.25 (0.271)	0.15 (0.698)	2.14 (0.115)
Bacteria abundance (16S rRNA gene copies g^{-1})	0.11 (0.746)	4.29 (0.046)	1.27 (0.268)	2.30 (0.097)
Fungal abundance (ITS gene copies g^{-1})	0.28 (0.603)	2.42 (0.130)	0.59 (0.448)	0.31 (0.818)
Bacterial β -glucosidase (GH1 gene copies g^{-1})	2.49 (0.124)	4.24 (0.048)	0.41 (0.528)	1.06 (0.378)
Bacterial β -glucosidase (GH3 gene copies g^{-1})	3.98 (0.055)	0.01 (0.909)	0.31 (0.582)	4.52 (0.009)
Fungal β -glucosidase (GH1 gene copies g^{-1})	5.03 (0.032)	0.14 (0.713)	0.78 (0.384)	0.25 (0.858)
Fungal β -glucosidase (GH3 gene copies g^{-1})	1.10 (0.303)	13.91 (0.0001)	1.21 (0.280)	3.68 (0.022)

significantly lower in the soils planted with a quaternary mixture of cover crop species than the average of the four monoculture plots after all five drying and rewetting cycle treatments (0, 1, 2, 4

and 8 cycles), as shown in Figure 2. The cover crop species planted also had a significant effect on the substrate-induced respiration, regardless of wet/dry cycle treatment (Figure 2). Generally, oil

radish and sunflower plots resulted in greater substrate-induced microbial respiration than buckwheat and clover plots. The effect of the *Mix* factor was always greater than the effect of the *Type* factor on cumulative substrate-induced respiration.

In soils taken from all plots, more dry/wet cycles resulted in an earlier peak of CO₂ flux, a higher maximum respiration rate and greater cumulative substrate-induced respiration, as demonstrated in Figure 1. Hence, a higher cumulative respiration was

observed after 8 wet/dry cycles, relative to 0 wet/dry cycles, in contrast with our hypothesis. Warming significantly decreased the extent to which the cumulative respiration increased after 8 wet/dry cycles relative to 0 wet/dry cycles (Figure 3). However, Cover crops significantly increased the extent to which the cumulative respiration increased after 8 repeated dry/wet cycles relative to 0 dry/wet cycles (Figure 3). While the 4-species mixture of cover crops resulted in a slightly greater increase in cumulative respiration after 8 cycles, relative to 0 cycles, this was




FIGURE 2 | Cumulative substrate-induced microbial respiration after 0, 1, 2, 4 or 8 wet-dry cycles from soils subjected to warmed or ambient treatments and planted with no cover crops (control), cover crop monocultures (buckwheat, clover, radish and sunflower) or a 4-species cover crop mixture. *Warming* and *Cover crops* had no significant effect on cumulative respiration, but cover crop diversity (*Mix*) and species (*Type*) had a significant effect on soils subjected to each cycle. Error bars are standard deviations; $n = 4$.




FIGURE 3 | Effects of statistical factors (*Warming*, *Cover crops*, *Mix* and *Type*) on cumulative substrate-induced microbial respiration after 8 wet/dry cycles expressed as a percentage of the cumulative substrate-induced microbial respiration observed after 0 wet/dry cycles. Error bars are standard deviations; $n = 4$.

not statistically significant and there was no statistically significant difference observed between the four cover crop monoculture treatments (Figure 3).

3.3 | Relationship Between Cumulative Substrate-Induced Respiration and Soil Properties

Correlation coefficients for the relationships between soil chemical and biological properties and the cumulative respiration after 8 wet/dry cycles relative to cumulative respiration after 0 wet/dry cycles are presented in Table 2. The relative increase in microbial respiration was significantly positively correlated with cold water extractable organic carbon, and significantly negatively correlated with the abundance of the bacterial-GH1 gene encoding beta-glucosidase and also negatively correlated, albeit not statistically

TABLE 2 | Pearson correlation coefficients and p values for the relationships between the cumulative respiration after 8 wet/dry cycles relative to cumulative respiration after 0 wet/dry cycles and soil chemical and biological parameters.

Soil properties	Coefficient (r)	p
HWEOC ^a ($\mu\text{g g}^{-1}$)	0.041	0.784
CWEOC ^b ($\mu\text{g g}^{-1}$)	0.493	<0.001
Soil pH	-0.040	0.789
% LOI ^c	-0.103	0.492
Extractable NO_3^- (mg kg^{-1})	-0.196	0.187
Total extractable N (mg kg^{-1})	-0.206	0.164
%N	-0.113	0.450
%C	-0.168	0.260
C/N	-0.048	0.747
Bacteria abundance (16S rRNA gene copies g^{-1})	-0.165	0.268
Fungal abundance (ITS gene copies g^{-1})	-0.260	0.077
Bacterial β -glucosidase (GH1 gene copies g^{-1})	-0.414	0.004
Bacterial β -glucosidase (GH3 gene copies g^{-1})	-0.280	0.057
Fungal β -glucosidase (GH1 gene copies g^{-1})	0.007	0.960
Fungal β -glucosidase (GH3 gene copies g^{-1})	-0.277	0.059

^aHot water extractable carbon.

^bCold water extractable carbon.

^cLoss on ignition.

significantly, with the abundance of the bacterial-GH3 genes encoding beta-glucosidase (Table 2). Stepwise multivariate regression revealed only one soil chemical or biological property (cold water extractable organic carbon) that significantly predicted the relative increase in microbial respiration after 8 wet/dry cycles relative to 0 wet-dry cycles, with no other property significantly improving the regression model (Figure 4).

4 | Discussion

4.1 | The Influence of Cover Crop Species Diversity on Substrate-Induced Respiration

Our finding that the average legacy effect of planting four monocultures of cover crops resulted in significantly greater cumulative substrate-induced respiration than the cover crop mixture in all five drying and rewetting cycle treatments (Figure 2) implies that increasing plant diversity decreases soil microbial activity, in contrast to previous observations (Shu et al. 2021). Where positive relationships between plant diversity on ecosystem function are observed in field experiments, they can often be explained by greater plant biomass in more diverse plant communities (Zak




FIGURE 4 | Regression between the cumulative respiration after 8 wet/dry cycles relative to cumulative respiration after 0 wet/dry cycles and cold water extractable organic carbon (CWEOC) for all samples and experimental treatments combined ($n = 48$).

et al. 2003). None of the soil chemical or biological parameters measured were significantly affected by cover crop diversity (apart from substrate-induced respiration) to aid our mechanistic explanation of this result.

4.2 | The Influence of Cover Crop Species Type on Substrate-Induced Respiration

Rhizodeposition during the growth phase of the cover crop and residue decomposition after cover crop termination and incorporation both regulate cover crop legacy effects on the soil microbiome (Spedding et al. 2004; Nannipieri et al. 2023). Cover crop-induced legacy effects on the soil microbiome can manifest as a species/cultivar-specific shift in soil microbial community structure or the magnitude of soil microbial activity that persists after cover crop termination, during the growth and after the harvest of the following cash crop (Cazzaniga et al. 2023). We also found species-specific effects on cover crop-induced soil microbial legacy effects. For instance, soils previously planted with oil radish or sunflower resulted in significantly greater cumulative microbial substrate-induced respiration, compared with soil planted with buckwheat or clover (Figure 2). Cazzaniga et al. (2023) assessed the legacy effects of cover crops on soil microbial footprints and found that, while the effects of some other species persist only until the onset of cash crop, oil radish impacted soil microbial footprints until the harvesting of the cash crop. Specifically, they found that oil radish boosted both the presence and activity of microbial groups known for suppressing soil-borne fungal diseases of plants. In our experiment, the cover crop species did not significantly affect abundance of total bacteria (16S rRNA gene) or total fungi (ITS region gene), but we did find that plots previously planted with radish had the highest abundance of the bacterial β -glucosidase gene (GH3) (Figure S9) and the lowest relative abundance of the fungal β -glucosidase gene (GH3) (Figure S11), implying evidence of negative impacts of radish on fungal communities involved in the hydrolysis of cellobiose, as previously observed when growing

brassicas (Vukicevich et al. 2016; Tagele, Kim, and Shin 2021). We also observed greater concentrations of total N in plots previously planted with clover, which could be evidence of the legacy of N fixation by leguminous cover crops (Crotty and Stoate 2019; Castellano-Hinojosa and Strauss 2020).

4.3 | The Influence of Drying and Rewetting Cycles on Substrate-Induced Respiration

The largest impacts of drying and rewetting on substrate-induced respiration were observed within the first 20 h after the substrate was added to the soil. This observation is similar to that of Fraser, Corstanje, et al. (2016) who observed that the majority of the rapidly released CO_2 from substrate-induced respiration following drying and rewetting occurred within the first 24 h. Most of the C mineralised after substrate addition has been attributed to free sugars, soluble organic-N and fructans and plant residues containing more of these compounds will lead to greater respiration compared with those with lower proportions (Gunnarsson et al. 2008; Fraser, Todman, et al. 2016).

We found that substrate-induced respiration increased after multiple drying and rewetting cycles (Figure 1). This finding is in contrast to our hypothesis that drying and rewetting cycles would suppress the substrate-induced respiration. Soil respiration after rewetting is often explained more by substrate supply than by the soil microbial biomass (Wang et al. 2003). When soils face drought conditions the soil microorganisms produce extracellular polymeric substances and osmolytes; low molecular weight organic compounds that help them maintain cell integrity (Wood et al. 2001; Kakumanu, Ma, and Williams 2019; Bogati and Walczak 2022). Upon rewetting, the osmolytes, particularly trehalose, can be rapidly metabolised, either by the organisms that synthesised them, or by other microorganisms emerging from dormancy (Warren 2014, 2016, 2020), and this may result in priming of soil organic matter decomposition (Allison 2023). The role of osmolytes in drought

tolerance of the soil microbial community or its function may help explain the results we observed. The greater the number of drying and rewetting cycles, the greater the substrate-induced respiration. We postulate that multiple drying and rewetting cycles result in microorganisms becoming more adapted to drought (Preece et al. 2019; Evans and Wallenstein 2012). This adaptation could be because of selection for microorganisms more capable of undergoing dormancy or synthesising osmolytes and extracellular polymeric substances as a form of ecological memory because of genetic (Canarini et al. 2021) or non-genetic (Riber and Hansen 2021) diversity that increases the ability for individuals to enter and emerge from dormancy over time in populations exposed to repeated rounds of drying and rewetting cycles. Microorganisms may also be able to re-acquire osmolytes synthesised in response to the previous drying event or acquire compatible compounds already present in the soil solution (Malik and Bouskill 2022). This greater labile organic resource would have resulted in more carbon available to respire upon wetting and a greater potential to prime the decomposition of the barley shoot powder.

4.4 | The Legacy of Cover Crops on the Adaptation of Soil Microbial Function to Repeated Drying and Rewetting Cycles

We observed significantly lower %C in plots that had previously been planted with cover crops than control plots where cover crops were not established (Table 1). This finding contrasts with the results of a global meta-analysis which found that including cover crops within agricultural rotations increased %C, depending on soil texture and the increased %C positively correlated with levels of mineralisable carbon and nitrogen in soil (Jian et al. 2020). However, we observed no significant difference in the concentration of C_{WEOC} in cover cropping plots (227.6 µg C g⁻¹ soil), compared with control plots (224.5 µg C g⁻¹ soil), despite differences in total %C. It is therefore likely that a much greater proportion of the %C in the soils where cover crops were grown is labile. It is expected that, if a drying and rewetting event is preceded by greater availability of labile C, then the microorganisms have more resources available to invest in the production of osmolytes and extracellular polymeric substances to help protect them from desiccation (Kakumanu, Ma, and Williams 2019). This means that there is more carbon available to respire upon wetting and prime the decomposition of the barley shoot powder.

4.5 | The Legacy of Warming on the Adaptation of Soil Microbial Function to Repeated Drying and Rewetting Cycles

Soil warming is known to result in faster metabolism of microbially available organic carbon (Bradford et al. 2008; Adekanmbi et al. 2022), but may have also been affected by faster plant growth under the OTCs. The observed decrease in substrate-induced respiration after 8 dry/wet cycles, relative to 0 dry-wet cycles, because of warming may be because of warming-induced depletion of dissolved organic carbon as a legacy effect. Our finding therefore implies that there will be lower soil microbial activity when soil previously exposed to warming is further exposed to extremes of drying and rewetting

events, whereas cover crops may help sustain soil microbial activity after drying and rewetting. In addition, the legacy effect of warming resulted in an increase in the abundance of genes encoding fungal β-glucosidase (GH1 gene copies g⁻¹). This result could mean that warming depleted substrate availability and stimulated the fungal community to produce extracellular enzymes, as observed by Hou et al. (2016).

5 | Conclusions

We found that cover crop mixtures created a negative legacy effect in the soil which resulted in lower cumulative substrate-induced respiration than expected from the average of the same species grown in monoculture. However, some monocultures (radish and sunflower) left a greater positive legacy on substrate-induced respiration than others (buckwheat and clover). Exposure of the soil to multiple drying and rewetting cycles resulted in greater substrate-induced respiration and this response was increased by the legacy of cover crops and decreased by the legacy of soil warming. It is likely that exposure of soils to drying and rewetting cycles induces the microbial synthesis of osmolytes and extracellular polymeric substances to help microorganisms avoid desiccation during periods of dormancy and select for a microbial community more capable of entering and emerging from dormancy. There was a positive correlation between the cumulative respiration after 8 wet/dry cycles relative to cumulative respiration after 0 wet/dry cycles and cold water extractable carbon (C_{WEOC}), highlighting the importance of substrate availability for microbial adaptation to drying and rewetting cycles. Labile carbon in soils previously planted with cover crops may provide the substrates to synthesise osmolytes and extracellular polymeric substances. The depletion of labile carbon by warming may decrease the substrates available to synthesise these compounds. Therefore, cover crops may be able to mitigate the impact of warming on the ability of soil microorganisms to adapt to drying and rewetting cycles.

Author Contributions

Adetunji Alex Adekanmbi: conceptualization, data curation, formal analysis, investigation, methodology, visualization, writing – original draft. **Yiran Zou:** investigation, methodology, writing – review and editing. **Xin Shu:** investigation, methodology, writing – review and editing. **Giacomo Pietramellara:** funding acquisition, resources, supervision, writing – review and editing. **Shamina Imran Pathan:** data curation, formal analysis, investigation, methodology, writing – review and editing. **Lindsay Todman:** conceptualization, methodology, formal analysis, writing – review and editing. **Tom Sizmur:** conceptualization, funding acquisition, project administration, supervision, visualization, writing – original draft.

Acknowledgements

The authors acknowledge the technical support provided by Richard Casebow, Caroline Handley, Mike Charij, Francesco Leva, Anne Dudley, Fengjuan Xiao, Marta O'Brian, Laurence Dale, Caroline Thomas and Luci Corbett.

Data Availability Statement

The data that support the findings of this study are openly available at DOI: [10.17632/m7nxy3747f.1](https://doi.org/10.17632/m7nxy3747f.1).

References

- Adekanmbi, A. A., X. Shu, Y. Zou, and T. Sizmur. 2022. "Legacy Effect of Constant and Diurnally Oscillating Temperatures on Soil Respiration and Microbial Community Structure." *European Journal of Soil Science* 73: 1–14.
- Allison, S. D. 2023. "Microbial Drought Resistance May Destabilize Soil Carbon." *Trends in Microbiology* 31: 1–8. <https://doi.org/10.1016/j.tim.2023.03.002>.
- Bardgett, R. D., C. Freeman, and N. J. Ostle. 2008. "Microbial Contributions to Climate Change Through Carbon Cycle Feedbacks." *ISME Journal* 2: 805–814.
- Barnard, R. L., S. J. Blazewicz, and M. K. Firestone. 2020. "Rewetting of Soil: Revisiting the Origin of Soil CO₂ Emissions." *Soil Biology and Biochemistry* 147: 107819. <https://doi.org/10.1016/j.soilbio.2020.107819>.
- Bogati, K., and M. Walczak. 2022. "The Impact of Drought Stress on Soil Microbial Community, Enzyme Activities and Plants." *Agronomy* 12: 1–26.
- Bradford, M. A. 2013. "Thermal Adaptation of Decomposer Communities in Warming Soils." *Frontiers in Microbiology* 4: 1–16.
- Bradford, M. A., C. A. Davies, S. D. Frey, et al. 2008. "Thermal Adaptation of Soil Microbial Respiration to Elevated Temperature." *Ecology Letters* 11: 1316–1327.
- Brankatschk, R., N. Bodenhausen, J. Zeyer, and H. Burgmann. 2012. "Simple Absolute Quantification Method Correcting for Quantitative PCR Efficiency Variations for Microbial Community Samples." *Applied and Environmental Microbiology* 78: 4481–4489.
- Calderón, F. J., D. Nielsen, V. Acosta-Martínez, M. F. Vigil, and D. Lyon. 2016. "Cover Crop and Irrigation Effects on Soil Microbial Communities and Enzymes in Semiarid Agroecosystems of the Central Great Plains of North America." *Pedosphere* 26: 192–205.
- Canarini, A., H. Schmidt, L. Fuchslueger, et al. 2021. "Ecological Memory of Recurrent Drought Modifies Soil Processes via Changes in Soil Microbial Community." *Nature Communications* 12: 1–14. <https://doi.org/10.1038/s41467-021-25675-4>.
- Carey, J. C., K. D. Kroeger, B. Zafari, and J. Tang. 2018. "Passive Experimental Warming Decouples Air and Sediment Temperatures in a Salt Marsh." *Limnology and Oceanography: Methods* 16: 640–648.
- Castellano-Hinojosa, A., and S. L. Strauss. 2020. "Impact of Cover Crops on the Soil Microbiome of Tree Crops." *Microorganisms* 8: 328.
- Cazzaniga, S. G., S. van den Elsen, C. Lombaers, et al. 2023. "On the Legacy of Cover Crop-Specific Microbial Footprints." *Soil Biology and Biochemistry* 184: 109080. <https://doi.org/10.1016/j.soilbio.2023.109080>.
- Crotty, F. V., and C. Stoate. 2019. "The Legacy of Cover Crops on the Soil Habitat and Ecosystem Services in a Heavy Clay, Minimum Tillage Rotation." *Food and Energy Security* 8: 1–16.
- Drost, S. M., M. Rutgers, M. Wouterse, W. de Boer, and P. L. E. Bodelier. 2020. "Decomposition of Mixtures of Cover Crop Residues Increases Microbial Functional Diversity." *Geoderma* 361: 114060. <https://doi.org/10.1016/j.geoderma.2019.114060>.
- Dutta, H., and A. Dutta. 2016. "The Microbial Aspect of Climate Change." *Energy, Ecology and Environment* 1: 209–232.
- Evans, S. E., and M. D. Wallenstein. 2012. "Soil Microbial Community Response to Drying and Rewetting Stress: Does Historical Precipitation Regime Matter?" *Biogeochemistry* 109: 101–116. <https://doi.org/10.1007/s10533-011-9638-3>.
- Fraser, F. C., R. Corstanje, L. K. Deeks, et al. 2016. "On the Origin of Carbon Dioxide Released From Rewetted Soils." *Soil Biology and Biochemistry* 101: 1–5. <https://doi.org/10.1016/j.soilbio.2016.06.032>.
- Fraser, F. C., L. C. Todman, R. Corstanje, et al. 2016. "Distinct Respiratory Responses of Soils to Complex Organic Substrate Are Governed Predominantly by Soil Architecture and Its Microbial Community." *Soil Biology and Biochemistry* 103: 493–501.
- Ghani, A., M. Dexter, and K. W. Perrott. 2003. "Hot-Water Extractable Carbon in Soils: A Sensitive Measurement for Determining Impacts of Fertilisation, Grazing and Cultivation." *Soil Biology and Biochemistry* 35: 1231–1243.
- Gunnarsson, S., H. Marstorp, A. S. Dahlin, and E. Witter. 2008. "Influence of Non-Cellulose Structural Carbohydrate Composition on Plant Material Decomposition in Soil." *Biology and Fertility of Soils* 45: 27–36.
- Guoju, X., Z. Qiang, B. Jiangtao, et al. 2012. "The Relationship Between Winter Temperature Rise and Soil Fertility Properties." *Air, Soil and Water Research* 5: 15–22.
- Gyssels, G., J. Poesen, E. Bochet, and Y. Li. 2005. "Impact of Plant Roots on the Resistance of Soils to Erosion by Water: A Review." *Progress in Physical Geography* 29: 189–217.
- Harkness, C., M. A. Semenov, F. Areal, et al. 2020. "Adverse Weather Conditions for UK Wheat Production Under Climate Change." *Agricultural and Forest Meteorology* 282–283, 107862: 107862. <https://doi.org/10.1016/j.agrformet.2019.107862>.
- Holman, I. P., T. M. Hess, D. Rey, and J. W. Knox. 2021. "A Multi-Level Framework for Adaptation to Drought Within Temperate Agriculture." *Frontiers in Environmental Science* 8: 1–14.
- Hou, R., Z. Ouyang, D. Maxim, G. Wilson, and Y. Kuzyakov. 2016. "Lasting Effect of Soil Warming on Organic Matter Decomposition Depends on Tillage Practices." *Soil Biology and Biochemistry* 95: 243–249. <https://doi.org/10.1016/j.soilbio.2015.12.008>.
- Jian, J., X. Du, M. S. Reiter, and R. D. Stewart. 2020. "A Meta-Analysis of Global Cropland Soil Carbon Changes due to Cover Cropping." *Soil Biology and Biochemistry* 143: 107735.
- Kakumanu, M. L., L. Ma, and M. A. Williams. 2019. "Drought-Induced Soil Microbial Amino Acid and Polysaccharide Change and Their Implications for C-N Cycles in a Climate Change World." *Scientific Reports* 9: 1–12.
- Kaye, J. P., and M. Quemada. 2017. "Using Cover Crops to Mitigate and Adapt to Climate Change. A Review." *Agronomy for Sustainable Development* 37: 1–7.
- Kreyling, J. 2010. "Winter Climate Change: A Critical Factor for Temperate Vegetation Performance." *Ecology* 91: 1939–1948.
- Kreyling, J., K. Grant, V. Hammerl, et al. 2019. "Winter Warming Is Ecologically More Relevant Than Summer Warming in a Cool-Temperate Grassland." *Scientific Reports* 9: 14632.
- Kuan, H. L., P. D. Hallett, B. S. Griffiths, A. S. Gregory, C. W. Watts, and A. P. Whitmore. 2007. "The Biological and Physical Stability and Resilience of a Selection of Scottish Soils to Stresses." *European Journal of Soil Science* 58: 811–821.
- Kuffner, M., B. Hai, T. Rattei, et al. 2012. "Effects of Season and Experimental Warming on the Bacterial Community in a Temperate Mountain Forest Soil Assessed by 16S rRNA Gene Pyrosequencing." *FEMS Microbiology Ecology* 82: 551–562.
- Lavalle, C., F. Micale, T. D. Houston, et al. 2009. "Climate Change in Europe. 3. Impact on Agriculture and Forestry. A Review." *Agronomy for Sustainable Development* 29: 433–446.
- Li, J., P. Areveso, X. Wang, S. Jian, and L. Gamage. 2022. "Simulating Temperature in a Soil Incubation Experiment." *Journal of Visualized Experiments* 188: 64081. <https://doi.org/10.3791/64081>.
- Liu, C., Y. Wang, H. Chen, Q. Sun, Q. Jiang, and Z. Wang. 2023. "High Level of Winter Warming Aggravates Soil Carbon, Nitrogen Loss and Changes Greenhouse Gas Emission Characteristics in Seasonal Freeze-Thaw Farmland Soil." *Science of the Total Environment* 905: 167180. <https://doi.org/10.1016/j.scitotenv.2023.167180>.

- Malik, A. A., and N. J. Bouskill. 2022. "Drought Impacts on Microbial Trait Distribution and Feedback to Soil Carbon Cycling." *Functional Ecology* 36: 1442–1456.
- Marion, G. M., G. H. R. Henry, D. W. Freckman, et al. 1997. "Open-Top Designs for Manipulating Field Temperature in High-Latitude Ecosystems." *Global Change Biology* 3: 20–32.
- Nannipieri, P., S. E. Hannula, G. Pietramellara, M. Schloter, T. Sizmur, and S. Imran. 2023. "Legacy Effects of Rhizodeposits on Soil Microbiomes: A Perspective." *Soil Biology and Biochemistry* 184: 109107. <https://doi.org/10.1016/j.soilbio.2023.109107>.
- Papp, R., S. Marinari, M. C. Moscatelli, et al. 2018. "Short-Term Changes in Soil Biochemical Properties as Affected by Subsidiary Crop Cultivation in Four European Pedo-Climatic Zones." *Soil and Tillage Research* 180: 126–136. <https://doi.org/10.1016/j.still.2018.02.019>.
- Paterson, E., T. Gebbing, C. Abel, A. Sim, and G. Telfer. 2007. "Rhizodeposition Shapes Rhizosphere Microbial Community Structure in Organic Soil." *New Phytologist* 173: 600–610.
- Philippot, L., B. S. Griffiths, and S. Langenheder. 2021. "Microbial Community Resilience Across Ecosystems and Multiple Disturbances." *Microbiology and Molecular Biology Reviews* 85: 1–24.
- Preece, C., E. Verbruggen, L. Liu, J. T. Weedon, and J. Peñuelas. 2019. "Effects of Past and Current Drought on the Composition and Diversity of Soil Microbial Communities." *Soil Biology and Biochemistry* 131: 28–39. <https://doi.org/10.1016/j.soilbio.2018.12.022>.
- Radicetti, E., O. Osipitan, A. Langeroodi, S. Marinari, and R. Mancinelli. 2019. "CO₂ Flux and C Balance due to the Replacement of Bare Soil With Agro-Ecological Service Crops in Mediterranean Environment." *Agriculture* 9: 71.
- Rahmstorf, S., and D. Coumou. 2011. "Increase of Extreme Events in a Warming World." *Proceedings of the National Academy of Sciences of the United States of America* 108: 17905–17909.
- Riber, L., and L. H. Hansen. 2021. "Epigenetic Memories: The Hidden Drivers of Bacterial Persistence?" *Trends in Microbiology* 29: 190–194. <https://doi.org/10.1016/j.tim.2020.12.005>.
- Rustad, L. E., J. L. Campbell, G. M. Marion, et al. 2001. "A Meta-Analysis of the Response of Soil Respiration, Net Nitrogen Mineralization, and Aboveground Plant Growth to Experimental Ecosystem Warming." *Oecologia* 126: 543–562.
- Schimel, J. P. 2018. "Life in Dry Soils: Effects of Drought on Soil Microbial Communities and Processes." *Annual Review of Ecology, Evolution, and Systematics* 49: 409–432.
- Shu, X., Y. Zou, L. J. Shaw, L. Todman, M. Tibbett, and T. Sizmur. 2021. "Cover Crop Residue Diversity Enhances Microbial Activity and Biomass With Additive Effects on Microbial Structure." *Soil Research* 60: 349–359.
- Shu, X., Y. Zou, L. J. Shaw, L. Todman, M. Tibbett, and T. Sizmur. 2022. "Applying Cover Crop Residues as Diverse Mixtures Increases Initial Microbial Assimilation of Crop Residue-Derived Carbon." *European Journal of Soil Science* 73: e13232.
- Spedding, T. A., C. Hamel, G. R. Mehuys, and C. A. Madramootoo. 2004. "Soil Microbial Dynamics in Maize-Growing Soil Under Different Tillage and Residue Management Systems." *Soil Biology and Biochemistry* 36: 499–512.
- Tagele, S. B., R. H. Kim, and J. H. Shin. 2021. "Interactions Between Brassica Biofumigants and Soil Microbiota: Causes and Impacts." *Journal of Agricultural and Food Chemistry* 69: 11538–11553.
- Todman, L. C., F. C. Fraser, R. Corstanje, et al. 2018. "Evidence for Functional State Transitions in Intensively-Managed Soil Ecosystems." *Scientific Reports* 8: 1–10. <https://doi.org/10.1038/s41598-018-29925-2>.
- Vogelsang, T. J., and P. H. Franses. 2005. "Are Winters Getting Warmer?" *Environmental Modelling and Software* 20: 1449–1455.
- Vukicevich, E., T. Lowery, P. Bowen, J. R. Úrbez-Torres, and M. Hart. 2016. "Cover Crops to Increase Soil Microbial Diversity and Mitigate Decline in Perennial Agriculture. A Review." *Agronomy for Sustainable Development* 36: 1–4. <https://doi.org/10.1007/s13593-016-0385-7>.
- Wang, W. J., R. C. Dalal, P. W. Moody, and C. J. Smith. 2003. "Relationships of Soil Respiration to Microbial Biomass, Substrate Availability and Clay Content." *Soil Biology and Biochemistry* 35: 273–284.
- Warren, C. R. 2014. "Response of Osmolytes in Soil to Drying and Rewetting." *Soil Biology and Biochemistry* 70: 22–32. <https://doi.org/10.1016/j.soilbio.2013.12.008>.
- Warren, C. R. 2016. "Do Microbial Osmolytes or Extracellular Depolymerisation Products Accumulate as Soil Dries?" *Soil Biology and Biochemistry* 98: 54–63. <https://doi.org/10.1016/j.soilbio.2016.03.021>.
- Warren, C. R. 2020. "Pools and Fluxes of Osmolytes in Moist Soil and Dry Soil That Has Been Re-Wet." *Soil Biology and Biochemistry* 150: 108012. <https://doi.org/10.1016/j.soilbio.2020.108012>.
- Whitehead, D., K. P. Hogan, G. N. D. Rogers, et al. 1995. "Performance of Large Open-Top Chambers for Long-Term Field Investigations of Tree Response to Elevated Carbon Dioxide Concentration." *Journal of Biogeography* 22: 307.
- Wood, J. M., E. Bremer, L. N. Csonka, et al. 2001. "Osmosensing and Osmoregulatory Compatible Solute Accumulation by Bacteria." *Comparative Biochemistry and Physiology. Part A, Molecular and Integrative Physiology* 130: 437–460.
- Zak, D. R., W. E. Holmes, D. C. White, A. D. Peacock, and D. Tilman. 2003. "Plant Diversity, Soil Microbial Communities, and Ecosystem Function: Are There Any Links?" *Ecology* 84: 2042–2050.
- Zhang, S., Z. Yu, J. Lin, and B. Zhu. 2020. "Responses of Soil Carbon Decomposition to Drying-Rewetting Cycles: A Meta-Analysis." *Geoderma* 361: 114069. <https://doi.org/10.1016/j.geoderma.2019.114069>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.