

Integrating multiple evidence streams to understand insect biodiversity change

Article

Accepted Version

Cooke, R., Outhwaite, C. L., Bladon, A. J. ORCID: <https://orcid.org/0000-0002-2677-1247>, Millard, J., Rodger, J. G., Dong, Z., Dyer, E. E., Edney, S., Murphy, J. F., Dicks, L. V., Hui, C., Jones, J. I., Newbold, T., Purvis, A., Roy, H. E., Woodcock, B. A. and Isaac, N. J. B. (2025) Integrating multiple evidence streams to understand insect biodiversity change. *Science*, 388 (6742). eadq2110. ISSN 1095-9203 doi: 10.1126/science.adq2110 Available at <https://centaur.reading.ac.uk/121700/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1126/science.adq2110>

Publisher: American Association for the Advancement of Science

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Title: Integrating multiple evidence streams to understand insect biodiversity change

Authors: Rob Cooke^{1†} & Charlotte L. Outhwaite^{2,3†}, Andrew J. Bladon^{4,5}, Joseph Millard^{4,6}, James G. Rodger⁷, Zhaoke Dong^{4,8}, Ellie E. Dyer^{1,2}, Siobhan Edney¹, John F. Murphy⁹, Lynn V. Dicks⁴, Cang Hui^{7,10}, J. Iwan Jones⁹, Tim Newbold², Andy Purvis^{6,11}, Helen E. Roy^{1,12}, Ben A. Woodcock¹ & Nick J. B. Isaac^{1*}

Affiliations:

¹UK Centre for Ecology & Hydrology, Maclean Building, Crowmarsh Gifford, Wallingford, OX10 8BB, UK

²Centre for Biodiversity and Environment Research, Department of Genetics, Evolution & Environment, University College London, London, WC1E 6BT, UK

³Institute of Zoology, Zoological Society of London, Regent's Park, London, NW1 4RY

⁴Department of Zoology, University of Cambridge, Downing Street, Cambridge, CB2 3EJ, UK

⁵Ecology and Evolutionary Biology Division, School of Biological Sciences, University of Reading, Reading, RG6 6EX, UK

⁶Biodiversity Futures Lab, Natural History Museum, Cromwell Road, London, SW7 5BD, UK

⁷Department of Mathematical Sciences, Stellenbosch University, Stellenbosch, 7602, South Africa

⁸College of Plant Health and Medicine, Qingdao Agricultural University, Qingdao, 266109, China

⁹School of Biological and Behavioural Sciences, Queen Mary University of London, London, E1 4NS, UK

¹⁰Centre for Invasion Biology, African Institute for Mathematical Sciences, National Institute for Theoretical and Computational Sciences, Cape Town 7945, South Africa

¹¹Georgina Mace Centre for the Living Planet, Imperial College London, Silwood Park, Ascot, SL5 7PY, UK

¹²Center for Ecology and Conservation, University of Exeter, Penryn Campus, Cornwall, TR10 9FE, UK

†Rob Cooke and Charlotte L. Outhwaite have contributed equally to this work and share first authorship (order of names determined alphabetically)

*Corresponding author: Nick Isaac, UK Centre for Ecology & Hydrology, Maclean Building, Crowmarsh Gifford, Wallingford, Oxfordshire, OX10 8BB, UK

Email: njbi@ceh.ac.uk

Abstract

Insects dominate animal species diversity yet face many threats from anthropogenic drivers of change. Many features of insect ecology make them a challenging group, and the fragmented state of knowledge compromises our ability to make general statements about their status. Here, we discuss the challenges of assessing insect biodiversity change. We describe how multiple lines of evidence – time series, spatial comparisons, experiments, and expert opinion – can be integrated to provide a synthesis overview of how insect biodiversity responds to drivers. Applying this approach will generate testable predictions of insect biodiversity across space, time, and changing drivers. Given the urgency of accelerating human impacts across the environment, this approach could yield a much-needed rapid assessment of insect biodiversity change.

Introduction

Insects are an extraordinarily diverse and abundant group of animals (1, 2), and are essential to terrestrial and freshwater ecosystem functioning (3, 4) that is critical to human wellbeing (5). Most of the world's flowering plant species depend on animal pollinators to reproduce, with an estimated 82% of species pollinated exclusively by insects (6, 7). Moreover, insects are a crucial trophic node linking primary production to higher trophic levels: insects comprise the bulk of food sources for many birds, bats, reptiles, freshwater fishes, and other vertebrates (8).

There is overwhelming evidence of changes in insect communities in recent decades. These changes include rapid local and regional declines in abundance, occupancy, biomass, and diversity, as well as the reorganization of communities (9–17). Concern over these changes has permeated across scientific, public and policy sectors (11, 15–18). While some narratives have likely exaggerated the extent of insect declines (22, 23), insects are clearly threatened by a combination of widespread anthropogenic drivers, including land-use change, climate change, agricultural intensification, pollution, and introduced species (14, 24).

Despite the growing recognition of the ecological and economic importance of insects (14, 25–28) research on insect biodiversity has been limited (29, 30) and under-funded (31). As a result, the available evidence describing insect trends is spatially, temporally, and taxonomically unrepresentative (32, 33), such that the overall magnitude of the problem remains unclear (11, 23). Current evidence is biased towards human-dominated landscapes in Europe and North America (13, 34, 35). Taxonomic bias favors easily observed or identified groups, such as butterflies, bees, and dragonflies, while more taxonomically intractable or geographically isolated groups (e.g., parasitoid wasps, earwigs, or icebugs) have been neglected (33). Available time series are generally short, leading to extreme and potentially spurious inferences (36). The high interannual volatility of insect populations means that long-term trends and historical drivers of insect biodiversity change are particularly poorly captured by short time series (37). New approaches are required to understand and quantify changes in insect biodiversity and its drivers, to better support global policy recommendations and to target resources effectively to mitigate threats to insects. However, these challenges are exacerbated by the fact that insects are hyper-diverse, have complex lifecycles, and experience substantial population fluctuations.

For these reasons, a detailed global perspective of insect biodiversity change and its drivers remains elusive (21), making it challenging to develop specific measurable targets and goals for insects in large-scale biodiversity discussions and strategies (38). For example, Goal A of the Kunming-Montreal Global Biodiversity Framework refers to metrics such as population abundance and extinction risk, for which there are excellent data for vertebrates (39). In contrast, a mere ~1.2% of insect species (12,100 species out of ~1 million described insect species (1)) have undergone the International Union for Conservation of Nature (IUCN) Red List assessments necessary to calculate extinction risk, of which approximately a quarter (3,107 species) were evaluated as Data Deficient (40). An even smaller fraction has been assessed more than once, undermining any ability to understand changes in extinction risk (40).

These shortfalls underline the need for new approaches to deliver a more comprehensive, globally representative picture of the state of insects to inform scientific research, public understanding, and biodiversity policies. Sparked by reports of unexpectedly large insect declines (9), there has been a surge in the compilation and analysis of insect time-series data. While this effort has led to valuable insights, disagreement remains as to whether, where and why insects are declining. The 'why' question is particularly important because any attempts to reverse declines will need to address the

drivers of change. Moreover, a synoptic overview of insect biodiversity change and its drivers is a prerequisite for making testable predictions about the effects of conservation or policy actions. For example, how much would insects benefit if we reduced global pesticide usage. Indeed, prediction is essential for demonstrating scientific understanding (41). A predictive understanding of insect biodiversity change would make it possible to project the potential future state of insect biodiversity and the consequences for ecosystems (e.g., the risk of reduced pollination services) and, thus, inform evidence-based policy recommendations and conservation action.

Here, we address the grand challenge of understanding insect biodiversity change from fragmentary data. Recent reviews on insect biodiversity have focused on the state of insects (2, 42), the drivers of insect declines (13, 14), and advocated priorities for data generation (18, 37, 43). In this paper we lay out the principles by which a more comprehensive understanding is possible via the integration of multiple imperfect lines of evidence that are already available.

We first outline the features of insect biology that make them challenging to study. We then explore the strengths and weaknesses of different evidence types for understanding how and why insect biodiversity is changing. Finally, we identify what is required to harness the breadth of currently available evidence to build a better understanding of the state of insect biodiversity. Our approach provides a basis for advances in understanding the drivers of insect biodiversity change that are not constrained by the limitations of only one evidence type. Our overall goal is to lay a roadmap toward a solid understanding of insect biodiversity change on which conservation policies can be built, without having to wait decades for comprehensive monitoring data.

The challenge of understanding insect biodiversity change

Insects are hyper-diverse, comprising up to 90% of all multicellular animal species, with the majority still undescribed (1, 2). Beyond the numbers lie a staggering diversity of ecological adaptations and functional roles, from eusocial pollinators, to specialized parasitoids, to scavenging detritivores (3, 4, 44), in addition to highly complex lifecycles (45). Thus, we should expect insect biodiversity change to be extremely heterogeneous. Moreover, insect population sizes are more stochastic than for other taxonomic groups, so more datapoints are required to capture fluctuations than for vertebrates (46, 47). Stochasticity is high both within and between years, with even small differences in the timing of annual monitoring resulting in dramatic differences in reported species abundance (20, 21).

The complexity of insect biodiversity change is further compounded by a system of interacting drivers (48, 49). At the local to landscape scale, there is strong evidence for land use, climate change, pesticides, disease, and invasive species as key drivers of insect biodiversity change (2, 14, 50, 51). What is lacking is a quantitative assessment of the relative importance of the various drivers across scales, as well as knowledge on how these drivers interact to cause change. It is only by understanding these complex and dynamic driver-response relationships that it will be possible to identify when and where declines in insect biodiversity may be reversible (52) and deliver a synoptic view of insect biodiversity change and its consequences.

Ecology and Traits

The impact of direct drivers of change is mediated by insect ecology and evolution. Species traits have long been used to provide mechanistic insights into how biodiversity responds to drivers (53). Traits are likely to be especially valuable for understanding differential responses to drivers in insects, given the diversity of insect life histories and dearth of high-quality data. Indeed, as many insects undergo metamorphosis, their ecological niche changes dramatically during their lifecycle,

such that environmental constraints on one life stage can be quite different to those on another (54). Thus, insect life histories are highly variable, encompassing inherent differences in life stages (e.g., larva vs adult), as well as in longevity, dormancy, synchronization (55), and reproductive strategies (sexual vs asexual; egg laying vs viviparity). Moreover, understanding how species with shared traits respond to drivers provides insights about the whole trait group, even when specific data are not available for all (which is often the case for insects). For example, whether forest-dwelling insects prefer open glades, closed understory, or forest canopy will influence how they respond to deforestation (56, 57), and species preferences combined with rates of deforestation could be used to infer insect trends.

Species' ecology can also interact with characteristics of the surrounding landscape to increase susceptibility to decline (Fig. 1), with poorly dispersing or wingless species (58, 59) or those unable to persist in certain habitat matrices (e.g. intensive agriculture) unable to survive (60). Better evidence on how insect traits and life stages mediate the impact of drivers is crucial to understanding their ongoing and future responses to global change (61).

Evolution and Plasticity

Global drivers subject insects to changing, and often strong, selection pressures, leading to evolutionary responses that also mediate the effects of drivers on insect diversity (62). Thus, understanding and predicting insect responses to environmental change is complicated by evolutionary dynamics and vice versa (63, 64). For instance, dispersal, driven by land-use change, can shape the spatial pattern of insect genetics (65). Similarly, crop domestication has selected for some insects to become agricultural 'pests' (66, 67).

Insects exhibit various examples of beneficial adaptive plasticity in response to environmental change, including diapause induction or suppression of reproductive output under harsh conditions (68). However, insects may also pursue maladaptive pathways (i.e., developmental traps). For example, climate change in the temperate zone is causing insects to complete their lifecycles more quickly, leading some to attempt additional generations (and often fail) before winter sets in (69). These short-term changes in turn impact insect population dynamics and can be translated into long-term adaptation. Temperature changes are a key driver of evolutionary adaptation in insects, and as suitable climates shift, insects are challenged with adapting to new conditions or tracking suitable ones (72). Conversely, pesticide resistance may actively select for certain species in agricultural settings, increasing their abundance while the majority are adversely affected (73). We must therefore consider adaptive capacity and plasticity in response to environmental drivers, since evolutionary rescue effects will be more apparent for insects than for longer-lived organisms (76, 77).

Interaction networks

Insect populations are influenced by a range of antagonistic (e.g., predation or herbivory) and facilitative (e.g., pollination) interactions (Fig. 1). The consequences of environmental change for insects depend both on how they interact with other species and the structure of the whole network. Insect species often have specialized interactions, including as parasitoids, herbivores, or pollinators (78, 79). For example, 27% of tropical herbivorous insects feed exclusively on a single host plant species, while 48% feed on plants within one genus and 60% within one family (80). Holometabolous insects (those with complete metamorphosis) experience shifting interactions throughout their lifecycles (e.g., many hoverflies change from predatory larvae to facultative pollinators as adults). This makes insects especially vulnerable to co-extinctions with their interaction partners (81, 82). Furthermore, the loss of generalist insect species may have wide

ranging consequences on network resilience and robustness through cascading extinctions (63), potentially resulting in coextinction of species with no direct interaction (84). These complexities mean that drivers that may superficially appear unimportant for certain species may still have consequences through indirect effects (82).

<FIGURE 1 HERE>

Emerging questions

These layers of complexity highlight the magnitude of the challenge in modeling insect biodiversity change from fragmentary data, but also suggest some emerging questions: which drivers are most important, for which insects, and what is the form and timing of the response? Can we extract sufficient signal to understand – broadly, if not precisely – the drivers of global change in insects, in models that are good enough to be useful for informing policy change or mitigation measures? Given the urgent need to address insect declines (85), we need to make better use of the data that is currently available.

Time series are not enough

Time series are comparable estimates of a biodiversity metric (e.g., population abundance, biomass) at a location at multiple points in time. Time series are therefore a direct form of evidence for biodiversity change, capturing the actual temporal dynamics of interest, and evidence from time series has formed a major part of current knowledge on insect change (11, 86). Individual time series vary in their temporal, spatial, and taxonomic coverage, while time series are undertaken for diverse reasons, including monitoring associated with recovery following restoration, post pollution or weather events, or in habitats known to be under threat. Collectively, the portfolio of existing insect time series is neither spatially nor taxonomically representative (35). These issues hinder our ability to use time series to quantify how insect biodiversity is changing at global and regional scales, and limit how well we can identify the causes of change (43).

The temporal coverage of individual time series varies in terms of the *frequency* (number of sampling occasions) and the *span* (the length of time between the first and last samples) of the period assessed (Fig. 2). Ideally, the frequency of data collection would be sufficient to capture change from one generation to another, whilst accounting for within-season variation (e.g., as adults emerge and then die). Instead, most insect time series contain gaps (35, 37), and many represent ‘snapshot’ resurveys of locations sampled once in the past (13% of studies analyzed in van Klink et al. (11); Fig. 2). Snapshot surveys are particularly problematic for insects, where year-to-year population variation can be much higher than for plants or vertebrates (20), creating substantial noise in the trend estimate and extreme sensitivity to the baseline conditions (21, 55). This noise can introduce bias if there is any non-randomness in the circumstances of either the original or repeat survey (87). As a result, intermittently sampled (i.e., less frequent than annual sampling) time series spanning fewer than ten years are unlikely to provide reliable estimates of change in insect populations (21, 46, 87). But with little support for long-term monitoring (20), data are more typically collected over much shorter periods. Even for longer time series, the frequency and span of available time series (Fig. 2) means they often lack the statistical power to detect trends (21, 36, 52, 88) or isolate associated drivers (43).

<FIGURE 2 HERE>

An essential quality for time-series data is that the sampling protocol should be consistent over time, so that comparable estimates of biodiversity are obtained (or that variation, e.g., due to surveyor identity, can be statistically accounted for). However, consistency is not sufficient for producing data that can inform the wider question of why insect biodiversity is changing. Collections of time-series data will only provide such information if they are representative of environmental gradients and the range of exposure to threats (90). Instead, site selection is often non-random, with many time series located at field stations or other areas likely to be buffered from ongoing threats (35, 87) (e.g., 34% of plots in the meta-analysis by van Klink et al. (11) are located in protected areas). By contrast, other time series may have been initiated in response to an environmental perturbation, either natural (such as a weather event), catastrophic (e.g., pollution) or artificial (e.g., experimental, restoration or changing management). In the absence of matching controls, these data have the potential to skew our perception of how and why insect biodiversity is changing (32). Resource limitation typical to long-term sampling strategies may also mean that sampling effort is limited, raising questions as to the power to identify trends in some populations.

A further issue is that most time series lack scale-relevant information on external drivers and how they changed over time. Where this information is available, these drivers are often correlated and difficult to disentangle, which limits the ability of time-series data to attribute the causes of biodiversity change (10, 91, 92). Without quantification of how drivers affect insect diversity (e.g., driver decomposition; (93, 94)) we cannot hope to predict how alternative policies and actions will affect future trends.

Sparked by the debate on insect biodiversity change, there have been calls to address the issues outlined above by greatly expanding the network of insect monitoring schemes (18, 37, 52). For instance, Didham et al. (21) suggested that intensive annual monitoring (e.g., (95)) could be complemented by extensive but infrequent occupancy surveys at large numbers of sites. A step-change in insect monitoring is indeed urgently required, and well-designed, long time series will ultimately provide the best quantification of biodiversity change (although attribution to drivers will remain difficult). However, the cost of ongoing inaction will be high, and we cannot wait decades to accumulate these data (17, 34, 86); there is enough evidence about insect declines in some regions to demand immediate remedial action (85). To target action where it is most needed, we require a rapid synthesis on the causes of insect biodiversity change using data from a broad range of evidence types, not from just time series.

Evidence beyond time series: developing threat-response models

Time series provide the best evidence that change has happened but struggle to tell us why that change occurred. Understanding why biodiversity has changed is critical if we are to reverse declines, prevent extinctions, and maintain ecosystem function into the future (41, 96). To build a predictive understanding of insect biodiversity change we need to first model the relationships between biodiversity metrics and direct drivers. We refer to these relationships as threat-response models (TRMs; *sensu* (97), also known as pressure-impact relationships (98)). By quantifying the relationships between drivers and biodiversity metrics, it becomes possible to project the biodiversity response across spatial domains where the driver intensity is known. In this way, TRMs can be used to summarize the total impact of a driver within a given domain, and to make testable predictions for regions where no biodiversity data exists. Projecting TRMs in time makes it possible to re-evaluate historical baseline conditions (99) and compare biodiversity trends under a range of plausible futures of climate and socioeconomic change (96).

Well-established TRMs, such as those developed as part of the GLOBIO (98) and PREDICTS (100) projects, are parameterized largely using data from vertebrates. However, the ability of TRMs to simplify multidimensional responses using trait-based approaches makes them particularly valuable for insects (and other hyper-diverse groups). The notion of using models to link insect biodiversity with threat gradients has been gaining traction in recent years, with insights from studies at local (101), regional (102) and global (48) scales. More broadly, insect conservation biologists have argued that experimentation (43) or spatial comparisons (52) can be employed to build models and reveal the most important drivers of insect biodiversity change. Where rigorous data is lacking, models linking insect populations with threats can also be derived from eliciting the opinion of taxon experts, using techniques from the social sciences (51, 103). Such approaches are particularly valuable for understudied taxa or geographic regions where structured expert opinion processes can provide key insights about population declines in the absence of quantitative data. These and other types of approaches all provide useful evidence, in the form of TRMs, to augment what can be learned from population time series.

The ideal dataset for building TRMs would measure change over a long period, have broad taxonomic and spatial coverage, and represent the full range of drivers. In reality, few biodiversity datasets come close to this ideal, although for well-studied groups such as birds and mammals they are sufficient for building useful TRMs. The evidence streams available for insects occupy different spaces along these three axes (Fig. 3): in other words, the available evidence types have complementary strengths and weaknesses.

<FIGURE 3 HERE>

To illustrate these complementarities, we evaluate each evidence type against six 'ideal properties' (Fig. 4) for understanding and predicting insect biodiversity change. These are 1) the ability to describe long-term trends, 2) the ability to capture transient dynamics, 3) whether the results of analyzes using the evidence type can be decomposed or aggregated across taxonomic and spatial scales, 4) whether the results of analyzes using the evidence type can be generalized and transferred, 5) whether corresponding driver information is available at the same scale as the evidence type, and 6) whether the evidence type can unpick mechanistic or causal threat-response relationships, including interacting (e.g., antagonistic, synergistic, additive) drivers.

<FIGURE 4 HERE>

Experiments

The strength of experiments lies in their ability to confirm causal links between drivers and biodiversity metrics (43). Experiments vary along a spectrum from lab-based, to plot or semi-field studies, up to more challenging landscape-scale manipulations (104). Whilst experiments deliver strong evidence, it is difficult to translate the outcomes to real world spatial and temporal scales (91). For instance, lab-based experiments are often targeted at individual species (minimal taxonomic breadth; Fig. 3) to characterize their response to a single factor e.g., thermal optima or agrochemical exposure (49, 105), and are often short-term (106) (low temporal breadth; Fig. 3).

Whilst experiments provide an opportunity to identify casual relationships, the scope of environmental drivers manipulated is typically limited (i.e., low environmental breadth; Fig 3) and so may only partially capture causes of declines. Moreover, experiments often fail to capture how

responses to drivers change when species are embedded in real interaction networks (Fig. 1) (although natural experiments are increasingly common (107)). Thus, simply scaling up the results of small-scale experiments may not be informative (108).

Ultimately, while experiments are common, there are few at a large enough spatial and temporal scale to fully capture real-world effects (91), reflecting practical and financial constraints (109). These limitations can be partially mitigated in distributed collaborative experiments (99) or transcended by synthesizing results across multiple experiments using meta-analysis. Combining multiple experiments provides a means to quantify generalized relationships between anthropogenic threats and insect diversity (90). Still, meta-analyses are subject to their own limitations, such as publication bias (110), inferential errors (111) and lack of data availability (112, 113). Nevertheless, meta-analytic approaches have already been effective in assessing the average response of insects to drivers including urbanization (114), dams (115), and nutrient enrichment (116, 117).

Spatial comparisons

Spatial comparisons – the comparison of biodiversity across sites – are often used to investigate insect biodiversity change (48, 118, 119). Their use assumes that patterns in space can shed light on patterns over time (120, 121). For example, Hallmann *et al.* (9) synthesized data from 63 sites within protected areas in Germany – most visited only once – to infer a 27-year time series of flying insect biomass.

Spatial comparisons have high statistical power to identify relationships between drivers and biodiversity declines and quantify their potential impact (52). Indeed, comparisons across whole landscapes allow sites to be compared across multiple driver gradients, e.g. (118). In addition, larger-scale changes can be modeled by bringing such comparisons together. For example, the PREDICTS project has compiled a global database of hundreds of studies comparing sites differing in land use types and intensity (122). TRMs built with these data have estimated average responses of biodiversity metrics across land-use gradients, including for insects (48, 123), and projected these responses in space and time (100).

Spatial comparisons have limitations for assessing changes in insect biodiversity (91, 124). They are correlative, limiting causal inference, since correlation between drivers makes their effects hard to separate (although see (125)). By taking snapshots in time they risk being confounded by transient dynamics and legacies of site history, and they overlook gains or losses in the regional species pool (an aspect of ‘shifting baseline syndrome’ (126)). The advantage, however, lies in the relative simplicity of assembling the evidence base: they often sample more sites, can be undertaken in more locations (91) and, given the statistical power with which they link spatial variation in biodiversity and drivers, have so far been undervalued as evidence of insect biodiversity change (52). Perhaps most importantly, their collection can be achieved in very short time scales and can be reactive in terms of their focus, including under-represented regions, habitats, or taxonomic groups.

Expert elicitation

Information from experts (including scientists, indigenous people, and other non-scientist specialists) reflects their accumulated knowledge and experience. Expert elicitation around biodiversity trends, and the drivers thereof, is particularly valuable for poorly studied insect groups, in the absence of more direct forms of evidence. Expert elicitation may also provide insights into the impact of multiple interacting drivers on better-studied groups, which may be difficult to disentangle quantitatively but can be teased apart conceptually (127, 128). A major limitation is the subjective nature of expert knowledge, which makes assessing the reliability and repeatability of results

challenging, especially when very few experts are available (14, 30). Similarly, the process of expert elicitation may introduce its own bias through survey design and choice of questions as well as interactions between participants in group meetings, although approaches such as the Delphi technique can be used to reduce some of these issues (129).

Given sufficient expertise, expert elicitation has the potential for high taxonomic and geographic coverage. It provides information that may be costly to gather in other ways (e.g. large-scale, long-term monitoring), although detailed species assessments (e.g., IUCN Red List assessments) can be time, data and cost intensive. At a broader scale, Miličić *et al.* (51) used a top-down expert elicitation process to gather information from 413 respondents on the most relevant threats to insect biodiversity in general, highlighting key differences between regions and taxa. This indirect evidence can be used to link the state of insect biodiversity with specific threats, especially where direct evidence is not available, or as a comparison to other types of evidence.

Harnessing the breadth of evidence

We have explored how TRMs based on experiments, spatial comparisons and expert elicitation provide alternative and complementary sources of evidence linking insect biodiversity change with drivers. Crucially, none of these evidence types alone can cover the full taxonomic, spatial, or temporal dimensions required for a synoptic and predictive understanding of insect biodiversity change (Figs. 3 and 4). Thus, there is an urgent need to harness this breadth of evidence, to synthesize the inferences from each evidence type in ways that maximize the complementarities and overcome the limitations of individual evidence types. To do this, we need to understand how to integrate diverse evidence streams and explore the limits to which we can combine TRMs and extend their inferences to different taxonomic, spatial, or temporal settings. Without attempts to generalize, the literature could become a descriptive ‘stamp collection’ of case studies (90).

One avenue for combining evidence is through model-based data integration, in which the relationship between data types is defined in terms of model parameters (130, 131). Integrated modeling is particularly well developed in the context of species distribution models. For instance, Kotta *et al.* (132) combined experimentally defined tolerance levels with spatial data to model climate change effects on a macroalgal species and its herbivore. Domisch *et al.* (133) combined freshwater species occurrence information with expert range data leading to improved spatial predictions. Whilst integrated modeling is conceptually attractive, there are analytical challenges, particularly when combining data that differ in both quality and quantity (134) and where there are mismatches in spatial scale among data types (135). As a result, formal data integration may currently only be possible for well-studied regions and taxa where data availability is highest. This starting point, however, is fertile ground for new modelling tools to be developed (136).

A tractable alternative is to compare the predictions from different types of TRM against one another, by projecting the models across space (e.g., using spatial data layers) and taxa (e.g., using species traits). Regions of parameter space where the model projections agree about the direction and/or magnitude of biodiversity change indicate higher confidence in the state of insect biodiversity. Regions where they disagree highlight priority research gaps or model inadequacies. Regions where direct evidence of biodiversity change is available (i.e., time series) provide an opportunity for external validation (90, 137) and a chance to demonstrate predictive understanding (41). This opens up a range of options for iterative model refinement to achieve the best fit to empirical data (known as a ‘digital twin’ (138)). For example, one could apply weights to each evidence type (139), then solve for the weights that optimize fit to the observed time-series trends, or test for lagged responses to increases in threat intensity (140). Similarly, when integrating TRMs

for different threat types, one could compare models that assume additivity among threats with those in which the threats act synergistically or antagonistically.

Synthetic TRMs integrating evidence types and drivers would provide a starting point to answer some of the biggest questions about insect biodiversity change (Fig. 5). Projecting the model in space would make it possible to assess the magnitude of insect biodiversity change across large scales and to identify probable hotspots of decline. Projecting the model in time would make it possible to explore plausible future scenarios for insect biodiversity (96). Aggregating insects by functional traits (e.g., body size, diet, fecundity, etc.) (141) that underpin ecosystem processes would make it possible to explore the consequences of insect biodiversity change for higher trophic levels (especially insectivorous bats and birds) and on benefits that people derive from insects (including pollination, pest control and nutrient cycling, amongst others (3, 4)). The modeling framework described here is also easily extensible to incorporate indirect drivers and the effects of mitigation strategies such as protected areas (96, 142). Based on these principles, a broad-brush quasi-global model of insect responses to land-use is within reach (sensu (48, 100)), as are regional models of multiple threats for well-studied Orders.

<FIGURE 5 HERE>

Whilst powerful, models have limitations. We see four major challenges for the development of large-scale and credible TRMs for insects. The first is around the technical aspects of model-fitting. There are well-known difficulties of modeling biodiversity using heterogeneous data (144), especially where the scale of inference is different from the scale of the data (135) or where there is context-dependency (111). A second challenge is around scale: TRMs built to address global questions (e.g. about the net magnitude of change) will not be suitable for addressing local questions (e.g. about the effect of a given intervention). Thus, TRMs at different scales will be appropriate for addressing different questions, but identifying the appropriate level of detail for a particular application may not always be obvious. The third challenge is around communication: it will be important for models to capture and convey the many forms of uncertainty and validity (90, 145). However, inferences around biodiversity change are often confounded by temporal, spatial and phylogenetic clustering within the data. Capturing all these uncertainties risks rendering the model uninformative for decision-making (145, 146). Similarly, it will be important to avoid making unwarranted claims about generality (90), particularly when presenting models as being of global relevance (147). Formal analysis of the degree to which models can be validly extended (i.e., transferability) is useful to place limits on the potential inference and should become standard (148, 149). Finally, there are ecological challenges that will be hard to overcome: certain aspects of insect ecology will be especially hard to model (as noted in Section 2). Early generations of insect biodiversity projection models will be unable to capture complex life histories or species interactions, but these and other processes could be added later (143). There are taxonomic groups and parts of the world where almost nothing is known about the insect fauna. Thus, one of the goals for the first generation of synthetic insect TRMs would be to highlight the remaining gaps (gaps not covered by data or by valid model inferences) where new data are urgently needed.

Outlook

Even with the caveats outlined above, there is sufficient evidence of changes in insect biodiversity to demand urgent action (85). This action needs to be informed and directed by a strong evidence base (i.e., sufficient extent, depth, and representativeness of evidence across time, space, and taxonomy, with levels of uncertainty that allow for clear decisions). Intergovernmental commitments to avert

biodiversity loss will always be limited, so resources must be used effectively. Even when commitments are made, the historic track record of meeting such goals has been less than impressive (150). The recent Global Biodiversity Framework provides new impetus for action, with an ambitious target for the reduction of pressures on natural systems by 2030, and a goal (amongst others) to achieve healthy and resilient populations of wild species by 2050. There is an urgent need for the biodiversity modeling community to evaluate whether the scale of ambition defined in the targets is sufficient to put us on a pathway toward the 2050 goal, i.e. to ‘bend the curve’ (96). Due to the sheer number of insects, their vast ecological diversity and the paucity of available data, the challenge of understanding global insect biodiversity change is an immense task. To date, limited resources have been directed toward this task. We are not the first to point out that time series of insect populations alone are insufficient to address this problem (43, 52). We have gone further in arguing that a broad suite of evidence types, brought into a common analytical framework, is required to evaluate the scale of the problem facing insect biodiversity, as well as providing a triage system to identify the highest priority taxa, places, and threats, and thus leverage points where mitigation can be most effective. Only through harnessing the full breadth of available evidence can we piece together the fragmentary data into a coherent picture. It is a small but important first step toward an insect-positive future.

References

1. N. E. Stork, How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? *Annu. Rev. Entomol.* **63**, 31–45 (2017).
2. P. Eggleton, The state of the world’s insects. *Annu. Rev. Environ. Resour.* **45**, 8.1-8.22 (2020).
3. C. M. Prather, S. L. Pelini, A. Laws, E. Rivest, M. Woltz, C. P. Bloch, I. Del Toro, C. K. Ho, J. Kominoski, T. A. S. Newbold, S. Parsons, A. Joern, Invertebrates, ecosystem services and climate change. *Biol. Rev.* **88**, 327–348 (2013).
4. A. Jankielsohn, The Importance of Insects in Agricultural Ecosystems. *Adv. Entomol.* **06**, 62–73 (2018).
5. S. Díaz, J. Settele, E. Brondízio, H. T. Ngo, M. Guèze, J. Agard Trinidad, A. Arneth, P. Balvanera, K. Brauman, S. Butchart, K. Chan, L. Garibaldi, K. Ichii, J. Liu, M. S. Subramanian, G. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. R. Chowdhury, Y. Shin, I. Visseren-Hamakers, K. Willis, C. Zayas, “Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES)” (2019); <https://zenodo.org/records/3553579>.
6. J. G. Rodger, J. M. Bennett, M. Razanajatovo, T. M. Knight, M. van Kleunen, T. L. Ashman, J. A. Steets, C. Hui, G. Arceo-Gómez, M. Burd, L. A. Burkle, J. H. Burns, W. Durka, L. Freitas, J. E. Kemp, J. Li, A. Pauw, J. C. Vamosi, M. Wolowski, J. Xia, A. G. Ellis, Widespread vulnerability of flowering plant seed production to pollinator declines. *Sci. Adv.* **7**, 3524–3537 (2021).
7. J. Ollerton, Winfree, R, Tarrant, S., How many flowering plants are pollinated by animals? *Oikos* **120**, 321–326 (2011).
8. D. W. Tallamy, W. G. Shriver, Are declines in insects and insectivorous birds related? *Ornithol. Appl.* **123**, duaa059 (2021).

- 473 9. C. A. Hallmann, M. Sorg, E. Jongejans, H. Siepel, N. Hofland, H. Schwan, W. Stenmans, A.
474 Müller, H. Sumser, T. Hörren, D. Goulson, H. De Kroon, More than 75 percent decline over 27
475 years in total flying insect biomass in protected areas. *PLoS ONE* **12**, e0185809 (2017).
- 476 10. S. Seibold, M. M. Gossner, N. K. Simons, N. Blüthgen, J. Müller, D. Ambarlı, C. Ammer, J.
477 Bauhus, M. Fischer, J. C. Habel, K. E. Linsenmair, T. Nauss, C. Penone, D. Prati, P. Schall, E. D.
478 Schulze, J. Vogt, S. Wöllauer, W. W. Weisser, Arthropod decline in grasslands and forests is
479 associated with landscape-level drivers. *Nature* **574**, 671–674 (2019).
- 480 11. R. van Klink, D. E. Bowler, K. B. Gongalsky, A. B. Swengel, A. Gentile, J. M. Chase, Meta-
481 analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*
482 **368**, 417–420 (2020).
- 483 12. P. Soroye, T. Newbold, J. Kerr, Climate change contributes to widespread declines among
484 bumble bees across continents. *Science* **367**, 685–688 (2020).
- 485 13. D. L. Wagner, Insect Declines in the Anthropocene. *Annu. Rev. Entomol.* **65**, 457–480 (2020).
- 486 14. D. L. Wagner, E. M. Grames, M. L. Forister, M. R. Berenbaum, D. Stopak, Insect decline in the
487 Anthropocene: Death by a thousand cuts. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2023989118
488 (2021).
- 489 15. G. D. Powney, C. Carvell, M. Edwards, R. K. A. Morris, H. E. Roy, B. A. Woodcock, N. J. B. Isaac,
490 Widespread losses of pollinating insects in Britain. *Nat. Commun.* **10**, 1018 (2019).
- 491 16. C. L. Outhwaite, R. D. Gregory, R. E. Chandler, B. Collen, N. J. B. Isaac, Complex long-term
492 biodiversity change among invertebrates, bryophytes and lichens. *Nat. Ecol. Evol.* **4**, 384–392
493 (2020).
- 494 17. M. S. Crossley, A. R. Meier, E. M. Baldwin, L. L. Berry, L. C. Crenshaw, G. L. Hartman, D. Lagos-
495 Kutz, D. H. Nichols, K. Patel, S. Varriano, W. E. Snyder, M. D. Moran, No net insect abundance
496 and diversity declines across US Long Term Ecological Research sites. *Nat. Ecol. Evol.* **4**, 1368–
497 1376 (2020).
- 498 18. J. A. Harvey, R. Heinen, I. Armbrrecht, Y. Basset, J. H. Baxter-Gilbert, T. M. Bezemer, M. Böhm,
499 R. Bommarco, P. A. V. Borges, P. Cardoso, V. Clausnitzer, T. Cornelisse, E. E. Crone, M. Dicke,
500 K.-D. B. Dijkstra, L. Dyer, J. Ellers, T. Fartmann, M. L. Forister, M. J. Furlong, A. Garcia-Aguayo,
501 J. Gerlach, R. Gols, D. Goulson, J.-C. Habel, N. M. Haddad, C. A. Hallmann, S. Henriques, M. E.
502 Herberstein, A. Hochkirch, A. C. Hughes, S. Jepsen, T. H. Jones, B. M. Kaydan, D. Kleijn, A.-M.
503 Klein, T. Latty, S. R. Leather, S. M. Lewis, B. C. Lister, J. E. Losey, E. C. Lowe, C. R. Macadam, J.
504 Montoya-Lerma, C. D. Nagano, S. Ogan, M. C. Orr, C. J. Painting, T.-H. Pham, S. G. Potts, A.
505 Rauf, T. L. Roslin, M. J. Samways, F. Sanchez-Bayo, S. A. Sar, C. B. Schultz, A. O. Soares, A.
506 Thancharoen, T. Tschardtke, J. M. Tylianakis, K. D. L. Umbers, L. E. M. Vet, M. E. Visser, A.
507 Vujic, D. L. Wagner, M. F. WallisDeVries, C. Westphal, T. E. White, V. L. Wilkins, P. H. Williams,
508 K. A. G. Wyckhuys, Z.-R. Zhu, H. de Kroon, International scientists formulate a roadmap for
509 insect conservation and recovery. *Nat. Ecol. Evol.* **4**, 174–176 (2020).
- 510 19. M. E. Saunders, J. K. Janes, J. C. O’Hanlon, Moving On from the Insect Apocalypse Narrative:
511 Engaging with Evidence-Based Insect Conservation. *BioScience* **70**, 80–89 (2020).
- 512 20. G. A. Montgomery, R. R. Dunn, R. Fox, E. Jongejans, S. R. Leather, M. E. Saunders, C. R.
513 Shortall, M. W. Tingley, D. L. Wagner, Is the insect apocalypse upon us? How to find out. *Biol.*
514 *Conserv.* **241**, 108327 (2020).

- 515 21. R. K. Didham, Y. Basset, C. M. Collins, S. R. Leather, N. A. Littlewood, M. H. M. Menz, J. Müller,
516 L. Packer, M. E. Saunders, K. Schönrogge, A. J. A. Stewart, S. P. Yanoviak, C. Hassall,
517 Interpreting insect declines: seven challenges and a way forward. *Insect Conserv. Divers.* **13**,
518 103–114 (2020).
- 519 22. C. D. Thomas, T. H. Jones, S. E. Hartley, “Insectageddon”: A call for more robust data and
520 rigorous analyses. *Glob. Change Biol.* **25**, 1891–1892 (2019).
- 521 23. R. K. Didham, F. Barbero, C. M. Collins, M. L. Forister, C. Hassall, S. R. Leather, L. Packer, M. E.
522 Saunders, A. J. A. Stewart, Spotlight on insects: trends, threats and conservation challenges.
523 *Insect Conserv. Divers.* **13**, 99–102 (2020).
- 524 24. S. G. Potts, J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, W. E. Kunin, Global
525 pollinator declines: trends, impacts and drivers. *Trends Ecol. Evol.* **25**, 345–353 (2010).
- 526 25. J. E. Losey, Ma. Vaughan, The Economic Value of Ecological Services Provided by Insects.
527 *BioScience* **56**, 311 (2006).
- 528 26. T. D. Schowalter, J. A. Noriega, T. Tscharnkte, Insect effects on ecosystem services—
529 Introduction. *Basic Appl. Ecol.* **26**, 1–7 (2018).
- 530 27. J. A. Noriega, J. Hortal, F. M. Azcárate, M. P. Berg, N. Bonada, M. J. I. Briones, I. Del Toro, D.
531 Goulson, S. Ibanez, D. A. Landis, M. Moretti, S. G. Potts, E. M. Slade, J. C. Stout, M. D. Ulyshen,
532 F. L. Wackers, B. A. Woodcock, A. M. C. Santos, Research trends in ecosystem services
533 provided by insects. *Basic Appl. Ecol.* **26**, 8–23 (2018).
- 534 28. S. H. Luke, H. E. Roy, C. D. Thomas, L. A. N. Tilley, S. Ward, A. Watt, M. Carnaghi, C. C.
535 Jaworski, M. P. T. G. Tercel, C. Woodrow, S. Aown, J. A. Banfield-Zanin, S. L. Barnsley, I.
536 Berger, M. J. F. Brown, J. C. Bull, H. Campbell, R. A. B. Carter, M. Charalambous, L. J. Cole, M.
537 J. Ebejer, R. A. Farrow, R. S. Fartyal, M. Grace, F. Highet, J. K. Hill, A. S. C. Hood, E. S. Kent, F.-T.
538 Krell, S. R. Leather, D. J. Leybourne, N. A. Littlewood, A. Lyons, G. Matthews, L. Mc Namara, R.
539 Menéndez, P. Merrett, S. Mohammed, A. K. Murchie, M. Noble, M.-R. Paiva, M. J. Pannell, C.-
540 K. Phon, G. Port, C. Powell, S. Rosell, F. Sconce, C. R. Shortall, E. M. Slade, J. P. Sutherland, J. C.
541 Weir, C. D. Williams, N. B. Zielonka, L. V. Dicks, Grand challenges in entomology: Priorities for
542 action in the coming decades. *Insect Conserv. Divers.* **16**, 173–189 (2023).
- 543 29. P. Cardoso, P. a.V. Borges, K. a. Triantis, M. a. Ferrández, J. L. Martín, Adapting the IUCN Red
544 List criteria for invertebrates. *Biol. Conserv.* **144**, 2432–2440 (2011).
- 545 30. J. A. Clark, R. M. May, Taxonomic Bias in Conservation Research. *Science* **297**, 191–192 (2002).
- 546 31. S. Mammola, N. Riccardi, V. Prié, R. Correia, P. Cardoso, M. Lopes-Lima, R. Sousa, Towards a
547 taxonomically unbiased European Union biodiversity strategy for 2030. *Proc. R. Soc. B* **287**,
548 20202166 (2020).
- 549 32. J. Troudet, P. Grandcolas, A. Blin, R. Vignes-Lebbe, F. Legendre, Taxonomic bias in biodiversity
550 data and societal preferences. *Sci. Rep.* **7**, 1–14 (2017).
- 551 33. E. Garcia-Rosello, J. Gonzalez-Dacosta, C. Guisande, J. M. Lobo, GBIF falls short of providing a
552 representative picture of the global distribution of insects. *Syst. Entomol.* **48**, 489–497 (2023).
- 553 34. A. Jeliaskov, D. Mijatovic, S. Chantepie, N. Andrew, R. Arlettaz, L. Barbaro, N. Barsoum, A.
554 Bartonova, E. Belskaya, N. Bonada, A. Brind’Amour, R. Carvalho, H. Castro, D. Chmura, P.

- Choler, K. Chong-Seng, D. Cleary, A. Cormont, W. Cornwell, R. de Campos, N. de Voogd, S. Doledec, J. Drew, F. Dziock, A. Eallonardo, M. J. Edgar, F. Farneda, D. F. Hernandez, C. Frenette-Dussault, G. Fried, B. Gallardo, H. Gibb, T. Gonçalves-Souza, J. Higuti, J. Y. Humbert, B. R. Krasnov, E. L. Saux, Z. Lindo, A. Lopez-Baucells, E. Lowe, B. Marteinsdottir, K. Martens, P. Meffert, A. Mellado-Díaz, M. H. M. Menz, C. F. J. Meyer, J. R. Miranda, D. Mouillot, A. Ossola, R. Pakeman, S. Pavoine, B. Pekin, J. Pino, A. Pocheville, F. Pomati, P. Poschlod, H. C. Prentice, O. Purschke, V. Raevel, T. Reitalu, W. Renema, I. Ribera, N. Robinson, B. Robroek, R. Rocha, S. H. Shieh, R. Spake, M. Staniaszek-Kik, M. Stanko, F. L. Tejerina-Garro, C. ter Braak, M. C. Urban, R. van Klink, S. Villéger, R. Wegman, M. J. Westgate, J. Wolff, J. Żarnowiec, M. Zolotarev, J. M. Chase, A global database for metacommunity ecology, integrating species, traits, environment and space. *Sci. Data* **7**, 1–15 (2020).
35. M. Desquilbet, L. Gaume, M. Grippa, R. Céréghino, J. F. Humbert, J. M. Bonmatin, P. A. Cornillon, D. Maes, H. V. Dyck, D. Goulson, Comment on "meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science* **370**, eabd8947 (2020).
36. G. N. Daskalova, A. B. Phillimore, I. H. Myers-Smith, Accounting for year effects and sampling error in temporal analyses of invertebrate population and biodiversity change: a comment on Seibold et al. 2019. *Insect Conserv. Divers.* **14**, 149–154 (2021).
37. E. M. Grames, G. A. Montgomery, D. H. Boyes, L. V. Dicks, M. L. Forister, T. A. Matson, S. Nakagawa, K. S. Prendergast, N. G. Taylor, M. W. Tingley, D. L. Wagner, T. E. White, P. Woodcock, C. S. Elphick, A framework and case study to systematically identify long-term insect abundance and diversity datasets. *Conserv. Sci. Pract.* **4**, e12687 (2022).
38. P. Donkersley, L. Ashton, G. P. A. Lamarre, S. Segar, Global insect decline is the result of wilful political failure: A battle plan for entomology. *Ecol. Evol.* **12**, e9417 (2022).
39. CBD Secretariat, "Post-2020 Global Biodiversity Framework" (Convention on Biological Diversity, Montreal, 2022); <https://www.cbd.int/doc/c/409e/19ae/369752b245f05e88f760aeb3/wg2020-05-l-02-en.pdf>.
40. IUCN, The IUCN Red List of Threatened Species (2022). <https://www.iucnredlist.org>.
41. J. E. Houlahan, S. T. McKinney, T. M. Anderson, B. J. McGill, The priority of prediction in ecological understanding. *Oikos* **126**, 1–7 (2017).
42. D. Goulson, The insect apocalypse, and why it matters. *Curr. Biol. CB* **29**, R967–R971 (2019).
43. W. Weisser, N. Blüthgen, M. Staab, R. Achury, J. Müller, Experiments are needed to quantify the main causes of insect decline. *Biol. Lett.* **19**, 20220500 (2023).
44. Saunders, Manu E., R. Rader, Ecosystem Services of Social Insects. *Encycl. Soc. Insects*, 352–358 (2021).
45. J. Kingsolver, H. Arthur Woods, L. B. Buckley, Potter, Kristen A., H. J. Maclean, J. K. Higgins, Complex Life Cycles and the Responses of Insects to Climate Change. *Integr. Comp. Biol.* **51**, 719–732 (2011).
46. E. R. White, Minimum time required to detect population trends: The need for long-term monitoring programs. *BioScience* **69**, 40–46 (2019).

- 594 47. R. Fox, C. A. Harrower, J. R. Bell, C. R. Shortall, I. Middlebrook, R. J. Wilson, Insect population
595 trends and the IUCN Red List process. *J. Insect Conserv.* **23**, 269–278 (2019).
- 596 48. C. L. Outhwaite, P. McCann, T. Newbold, Agriculture and climate change are reshaping insect
597 biodiversity worldwide. *Nature* **605**, 97–102 (2022).
- 598 49. R. J. Wilson, R. Fox, Insect responses to global change offer signposts for biodiversity and
599 conservation. *Ecol. Entomol.* **46**, 699–717 (2021).
- 600 50. L. V. Dicks, T. D. Breeze, H. T. Ngo, D. Senapathi, J. An, M. A. Aizen, P. Basu, D. Buchori, L.
601 Galetto, L. A. Garibaldi, B. Gemmill-Herren, B. G. Howlett, V. L. Imperatriz-Fonseca, S. D.
602 Johnson, A. Kovács-Hostyánszki, Y. J. Kwon, H. M. G. Lattorff, T. Lungharwo, C. L. Seymour, A.
603 J. Vanbergen, S. G. Potts, A global-scale expert assessment of drivers and risks associated with
604 pollinator decline. *Nat. Ecol. Evol.* **2021**, 1–9 (2021).
- 605 51. M. Miličić, S. Popov, V. V. Branco, P. Cardoso, Insect threats and conservation through the
606 lens of global experts. *Conserv. Lett.* **14**, e12814 (2021).
- 607 52. N. Blüthgen, M. Staab, R. Achury, W. W. Weisser, Unravelling insect declines: can space
608 replace time? *Biol. Lett.* **18**, 20210666 (2022).
- 609 53. D. O. Fisher, I. P. F. Owens, The comparative method in conservation biology. *Trends Ecol.*
610 *Evol.* **19**, 391–8 (2004).
- 611 54. E. Ashe-Jepson, M. P. Hayes, G. E. Hitchcock, K. Wingader, E. C. Turner, A. J. Bladon, Day-flying
612 lepidoptera larvae have a poorer ability to thermoregulate than adults. *Ecol. Evol.* **13**, e10623
613 (2023).
- 614 55. E. Jarzembowski, Insecta (Insects). *Encycl. Life Sci.*, doi: 10.1038/NPG.ELS.0001608 (2001).
- 615 56. S. R. P. J. Ross, F. H. Garcia, G. Fischer, M. K. Peters, Selective logging intensity in an East
616 African rain forest predicts reductions in ant diversity. *Biotropica* **50**, 768–778 (2018).
- 617 57. S. J. Willott, The effects of selective logging on the distribution of moths in a Bornean
618 rainforest. *Philos Trans R Soc Lond B Biol Sci* **354**, 59–66 (2000).
- 619 58. B. A. Woodcock, I. N. Vogiatzakis, D. B. Westbury, C. S. Lawson, A. R. Edwards, A. J. Brook, S. J.
620 Harris, K. A. Lock, N. Maczey, G. Masters, V. K. Brown, S. R. Mortimer, The role of
621 management and landscape context in the restoration of grassland phytophagous beetles. *J.*
622 *Appl. Ecol.* **47**, 366–376 (2010).
- 623 59. Z.-H. Zhao, C. Hui, D.-H. He, B.-L. Li, Effects of agricultural intensification on ability of natural
624 enemies to control aphids. *Sci. Rep.* **5**, 8024 (2015).
- 625 60. D. A. Driscoll, Is the matrix a sea? Habitat specificity in a naturally fragmented landscape. *Ecol.*
626 *Entomol.* **30**, 8–16 (2005).
- 627 61. T. Essens, F. van Langevelde, R. A. Vos, C. A. M. Van Swaay, M. F. WallisDeVries, Ecological
628 determinants of butterfly vulnerability across the European continent. *J. Insect Conserv.* **21**,
629 439–450 (2017).
- 630 62. S. P. Otto, Adaptation, speciation and extinction in the Anthropocene. *Proc. R. Soc. B Biol. Sci.*
631 **285**, 20182047 (2018).

- 632 63. P. Nosil, R. Villoutreix, C. F. De Carvalho, T. E. Farkas, V. Soria-Carrasco, J. L. Feder, B. J. Crespi,
633 Z. Gompert, Natural selection and the predictability of evolution in *Timema* stick insects.
634 *Science* **359**, 765–770 (2018).
- 635 64. A. Mas, Y. Lagadeuc, P. Vandenkoornhuyse, Reflections on the Predictability of Evolution:
636 Toward a Conceptual Framework. *iScience* **23**, 101736 (2020).
- 637 65. R. A. Sánchez-Guillén, A. Córdoba-Aguilar, B. Hansson, J. Ott, M. Wellenreuther, Evolutionary
638 consequences of climate-induced range shifts in insects. *Biol. Rev.* **91**, 1050–1064 (2016).
- 639 66. J. S. Bernal, A. M. Dávila-Flores, R. F. Medina, Y. H. Chen, K. E. Harrison, K. A. Berrier, Did
640 maize domestication and early spread mediate the population genetics of corn leafhopper?
641 *Insect Sci.* **26**, 569–586 (2019).
- 642 67. Y. H. Chen, Crop domestication, global human-mediated migration, and the unresolved role
643 of geography in pest control. *Elementa* **2016**, 106 (2016).
- 644 68. P. Gibert, V. Debat, C. K. Ghalambor, Phenotypic plasticity, global change, and the speed of
645 adaptive evolution. *Curr. Opin. Insect Sci.* **35**, 34–40 (2019).
- 646 69. H. Van Dyck, D. Bonte, R. Puls, K. Gotthard, D. Maes, The lost generation hypothesis: could
647 climate change drive ectotherms into a developmental trap? *Oikos* **124**, 54–61 (2015).
- 648 70. H. Weaving, J. S. Terblanche, P. Pottier, S. English, Meta-analysis reveals weak but pervasive
649 plasticity in insect thermal limits. *Nat. Commun.* **13**, 5292 (2022).
- 650 71. C. M. Sgrò, J. S. Terblanche, A. A. Hoffmann, What Can Plasticity Contribute to Insect
651 Responses to Climate Change? *Annu. Rev. Entomol.* **61**, 433–451 (2016).
- 652 72. S. C. Mason, G. Palmer, R. Fox, S. Gillings, J. K. Hill, C. D. Thomas, T. H. Oliver, Geographical
653 range margins of many taxonomic groups continue to shift polewards. *Biol. J. Linn. Soc.* **115**,
654 586–597 (2015).
- 655 73. S. Huang, L. Wang, L. Liu, Q. Fu, D. Zhu, Nonchemical pest control in China rice: a review.
656 *Agron. Sustain. Dev.* **34**, 275–291 (2014).
- 657 74. C. D. Thomas, E. J. Bodsworth, R. J. Wilson, A. D. Simmons, Z. G. Davies, M. Musche, L.
658 Conradt, Ecological and evolutionary processes at expanding range margins. *Nature* **411**, 577–
659 581 (2001).
- 660 75. J. K. Hill, H. M. Griffiths, C. D. Thomas, Climate Change and Evolutionary Adaptations at
661 Species' Range Margins. *Annu. Rev. Entomol.* **56**, 143–159 (2011).
- 662 76. A. Bush, K. Mokany, R. Catullo, A. Hoffmann, V. Kellermann, C. Sgrò, S. McEvey, S. Ferrier,
663 Incorporating evolutionary adaptation in species distribution modelling reduces projected
664 vulnerability to climate change. *Ecol. Lett.* **19**, 1468–1478 (2016).
- 665 77. H. A. Lindsey, J. Gallie, S. Taylor, B. Kerr, Evolutionary rescue from extinction is contingent on
666 a lower rate of environmental change. *Nature* **494**, 463–467 (2013).
- 667 78. Y. Aartsma, A. Cusumano, M. Fernández de Bobadilla, Q. Rusman, I. Vosteen, E. H. Poelman,
668 Understanding insect foraging in complex habitats by comparing trophic levels: insights from
669 specialist host-parasitoid-hyperparasitoid systems. *Curr. Opin. Insect Sci.* **32**, 54–60 (2019).

- 670 79. L. C. Jones, M. A. Rafter, G. H. Walter, Host interaction mechanisms in herbivorous insects –
671 life cycles, host specialization and speciation. *Biol. J. Linn. Soc.* **137**, 1–14 (2022).
- 672 80. V. Novotny, Y. Basset, Host specificity of insect herbivores in tropical forests. *Proc. R. Soc. B*
673 *Biol. Sci.* **272**, 1083–1090 (2005).
- 674 81. M. L. Moir, L. Hughes, P. A. Vesk, M. C. Leng, Which host-dependent insects are most prone
675 to coextinction under changed climates? *Ecol. Evol.* **4**, 1295–1312 (2014).
- 676 82. R. Kehoe, E. Frago, D. Sanders, Cascading extinctions as a hidden driver of insect decline. *Ecol.*
677 *Entomol.*, e212985 (2021).
- 678 83. J. Memmott, N. M. Waser, M. V. Price, Tolerance of pollination networks to species
679 extinctions. *Proc. R. Soc. Lond. B Biol. Sci.* **271**, 2605–2611 (2004).
- 680 84. L. P. Martins, D. Garcia-Callejas, H. R. Lai, K. L. Wootton, J. M. Tylianakis, The propagation of
681 disturbances in ecological networks. *Trends Ecol. Evol.*, doi: 10.1016/j.tree.2024.01.009
682 (2024).
- 683 85. M. L. Forister, E. M. Pelton, S. H. Black, Declines in insect abundance and diversity: We know
684 enough to act now. *Conserv. Sci. Pract.* **1**, e80 (2019).
- 685 86. M. S. Warren, D. Maes, C. A. M. van Swaay, P. Goffart, H. van Dyck, N. A. D. Bourn, I. Wynhoff,
686 D. Hoare, S. Ellis, The decline of butterflies in Europe: Problems, significance, and possible
687 solutions. *Proc. Natl. Acad. Sci. U. S. A.* **118**, e2002551117 (2021).
- 688 87. A. M. V. Fournier, E. R. White, S. B. Heard, Site-selection bias can drive apparent population
689 declines in long-term studies. *Conserv. Biol.* **33**, 1370–1379 (2019).
- 690 88. A. Gonzalez, B. J. Cardinale, G. R. H. Allington, J. Byrnes, K. Arthur Endsley, D. G. Brown, D. U.
691 Hooper, F. Isbell, M. I. O'Connor, M. Loreau, K. A. Endsley, D. G. Brown, D. U. Hooper, F. Isbell,
692 M. Loreau, A. Arbor, S. Paul, E. E. Station, Estimating local biodiversity change: a critique of
693 papers claiming no net loss of local diversity. *Ecology* **97**, 1949–1960 (2016).
- 694 89. R. van Klink, D. E. Bowler, O. Comay, M. M. Driessen, S. K. M. Ernest, A. Gentile, F. Gilbert, K.
695 B. Gongalsky, J. Owen, G. Pe'er, I. Pe'er, V. H. Resh, I. Rochlin, S. Schuch, A. B. Swengel, S. R.
696 Swengel, T. J. Valone, R. Vermeulen, T. Wepprich, J. L. Wiedmann, J. M. Chase, InsectChange:
697 a global database of temporal changes in insect and arachnid assemblages. *Ecology* **102**,
698 e03354 (2021).
- 699 90. R. Spake, R. E. O'Dea, S. Nakagawa, C. P. Doncaster, M. Ryo, C. T. Callaghan, J. M. Bullock,
700 Improving quantitative synthesis to achieve generality in ecology. *Nat. Ecol. Evol.*, 1–11
701 (2022).
- 702 91. A. De Palma, K. Sanchez-Ortiz, P. A. Martin, A. Chadwick, G. Gilbert, A. E. Bates, L. Börger, S.
703 Contu, S. L. L. Hill, A. Purvis, Challenges With Inferring How Land-Use Affects Terrestrial
704 Biodiversity: Study Design, Time, Space and Synthesis. *Adv. Ecol. Res.* **58**, 163–199 (2018).
- 705 92. W. E. Kunin, Robust evidence of declines in insect abundance and biodiversity. *Nature* **574**,
706 641–642 (2019).
- 707 93. C. S. Oedekoven, D. A. Elston, P. J. Harrison, M. J. Brewer, S. T. Buckland, A. Johnston, S.
708 Foster, J. W. Pearce-Higgins, Attributing changes in the distribution of species abundance to

709 weather variables using the example of British breeding birds. *Methods Ecol. Evol.*, doi:
710 10.1111/2041-210X.12811 (2017).

711 94. U. Bradter, A. Johnston, W. M. Hochachka, A. Saultan, J. E. Brommer, E. Gaget, J. A. Kålås, A.
712 Leikoinen, Å. Lindström, S. Piirainen, D. Pavón-Jordán, T. Pärt, I. J. Øien, B. K. Sandercock,
713 Decomposing the spatial and temporal effects of climate on bird populations in northern
714 European mountains. *Glob. Change Biol.* **28**, 6209–6227 (2022).

715 95. D. Karlsson, E. Hartop, M. Forshage, M. Jaschhof, F. Ronquist, The Swedish Malaise Trap
716 Project: A 15 Year Retrospective on a Countrywide Insect Inventory. *Biodivers. Data J.* **8**,
717 e47255 (2020).

718 96. D. Leclère, M. Obersteiner, M. Barrett, S. H. M. Butchart, A. Chaudhary, A. D. Palma, F. A. J.
719 DeClerck, M. D. Marco, J. C. Doelman, M. Dürauer, R. Freeman, M. Harfoot, T. Hasegawa, S.
720 Hellweg, J. P. Hilbers, S. L. L. Hill, F. Humpenöder, N. Jennings, T. Krisztin, G. M. Mace, H.
721 Ohashi, A. Popp, A. Purvis, A. M. Schipper, A. Tabeau, H. Valin, H. van Meijl, W.-J. van Zeist, P.
722 Visconti, R. Alkemade, R. Almond, G. Bunting, N. D. Burgess, S. E. Cornell, F. D. Fulvio, S.
723 Ferrier, S. Fritz, S. Fujimori, M. Grooten, T. Harwood, P. Havlík, M. Herrero, A. J. Hoskins, M.
724 Jung, T. Kram, H. Lotze-Campen, T. Matsui, C. Meyer, D. Nel, T. Newbold, G. Schmidt-Traub, E.
725 Stehfest, B. B. N. Strassburg, D. P. van Vuuren, C. Ware, J. E. M. Watson, W. Wu, L. Young,
726 Bending the curve of terrestrial biodiversity needs an integrated strategy. *Nature* **585**, 551–
727 556 (2020).

728 97. D. J. Ingram, G. B. Ferreira, K. E. Jones, G. M. Mace, Targeting Conservation Actions at Species
729 Threat Response Thresholds. *Trends Ecol. Evol.* **36**, 216–226 (2021).

730 98. A. M. Schipper, J. P. Hilbers, J. R. Meijer, L. H. Antão, A. Benítez-López, M. M. J. de Jonge, L. H.
731 Leemans, E. Scheper, R. Alkemade, J. C. Doelman, S. Mylius, E. Stehfest, D. P. van Vuuren, W.
732 J. van Zeist, M. A. J. Huijbregts, Projecting terrestrial biodiversity intactness with GLOBIO 4.
733 *Glob. Change Biol.* **26**, 760–771 (2020).

734 99. K. Klein Goldewijk, A. Beusen, G. van Drecht, M. de Vos, The HYDE 3.1 spatially explicit
735 database of human-induced global land-use change over the past 12,000 years. *Glob. Ecol.*
736 *Biogeogr.* **20**, 73–86 (2011).

737 100. T. Newbold, L. N. Hudson, S. L. L. Hill, S. Contu, I. Lysenko, R. A. Senior, L. Borger, D. J.
738 Bennett, A. Choimes, B. Collen, J. Day, A. D. Palma, S. Di, M. J. Edgar, A. Feldman, M. Garon,
739 M. L. K. Harrison, T. Alhusseini, S. Echeverria-london, D. J. Ingram, Y. Itescu, J. Kattge, V.
740 Kemp, L. Kirkpatrick, M. Kleyer, D. Laginha, P. Correia, C. D. Martin, S. Meiri, M. Novosolov, Y.
741 Pan, H. R. P. Phillips, D. W. Purves, A. Robinson, J. Simpson, S. L. Tuck, E. Weiher, H. J. White,
742 R. M. Ewers, G. M. Mace, J. P. W. Scharlemann, A. Purvis, Global effects of land use on local
743 terrestrial biodiversity. *Nature* **520**, 45–50 (2015).

744 101. S. R. P.-J. Ross, F. H. Garcia, G. Fischer, M. K. Peters, Selective logging intensity in an East
745 African rain forest predicts reductions in ant diversity. *Biotropica* **50**, 768–778 (2018).

746 102. Z.-H. Zhao, C. Hui, D.-H. He, B.-L. Li, Effects of agricultural intensification on ability of natural
747 enemies to control aphids. *Sci. Rep.* **5**, 8024 (2015).

748 103. N. Mukherjee, A. Zabala, J. Huge, T. O. Nyumba, B. Adem Esmail, W. J. Sutherland,
749 Comparison of techniques for eliciting views and judgements in decision-making. *Methods*
750 *Ecol. Evol.* **9**, 54–63 (2018).

- 751 104. Y. F. Wiersma, A review of landscape ecology experiments to understand ecological
752 processes. *Ecol. Process.* **11**, 57 (2022).
- 753 105. M. S. Heard, J. Baas, J. L. Dorne, E. Lahive, A. G. Robinson, A. Rortais, D. J. Spurgeon, C.
754 Svendsen, H. Hesketh, Comparative toxicity of pesticides and environmental contaminants in
755 bees: Are honey bees a useful proxy for wild bee species? *Sci. Total Environ.* **578**, 357–365
756 (2017).
- 757 106. A. K. Knapp, M. D. Smith, S. E. Hobbie, S. L. Collins, T. J. Fahey, G. J. A. Hansen, D. A. Landis, K.
758 J. La Pierre, J. M. Melillo, T. R. Seastedt, G. R. Shaver, J. R. Webster, Past, Present, and Future
759 Roles of Long-Term Experiments in the LTER Network. *BioScience* **62**, 377–389 (2012).
- 760 107. C. Herrick, K. Bell, The social life of natural experiments in epidemiology and public health.
761 *Sociol. Health Illn.* **46**, 276–294 (2024).
- 762 108. J. H. Lawton, Are There General Laws in Ecology? *Oikos* **84**, 177 (1999).
- 763 109. T. M. Fayle, E. C. Turner, Y. Basset, R. M. Ewers, G. Reynolds, V. Novotny, Whole-ecosystem
764 experimental manipulations of tropical forests. *Trends Ecol. Evol.* **30**, 334–346 (2015).
- 765 110. R. Rosenthal, The file drawer problem and tolerance for null results. *Psychol. Bull.* **86**, 638–
766 641 (1979).
- 767 111. R. Spake, D. E. Bowler, C. T. Callaghan, S. A. Blowes, C. P. Doncaster, L. H. Antão, S. Nakagawa,
768 R. McElreath, J. M. Chase, Understanding ‘it depends’ in ecology: a guide to hypothesising,
769 visualising and interpreting statistical interactions. *Biol. Rev.* **98**, 983–1002 (2023).
- 770 112. V. Stodden, J. Seiler, Z. Ma, An empirical analysis of journal policy effectiveness for
771 computational reproducibility. *Proc. Natl. Acad. Sci.* **115**, 2584–2589 (2018).
- 772 113. D. G. Roche, L. E. B. Kruuk, R. Lanfear, S. A. Binning, Public Data Archiving in Ecology and
773 Evolution: How Well Are We Doing? *PLoS Biol.* **13**, e1002295 (2015).
- 774 114. M. S. Fenoglio, M. R. Rossetti, M. Videla, Negative effects of urbanization on terrestrial
775 arthropod communities: A meta-analysis. *Glob. Ecol. Biogeogr.* **29**, 1412–1429 (2020).
- 776 115. J. Wang, C. Ding, J. Heino, X. Jiang, J. Tao, L. Ding, W. Su, M. Huang, D. He, What explains the
777 variation in dam impacts on riverine macroinvertebrates? A global quantitative synthesis.
778 *Environ. Res. Lett.* **15** (2020).
- 779 116. M. P. Nessel, T. Konnovitch, G. Q. Romero, A. L. González, Decline of insects and arachnids
780 driven by nutrient enrichment: A meta-analysis. *Ecology* **104**, e3897 (2023).
- 781 117. J. Gallego-Zamorano, M. M. J. de Jonge, K. Runge, S. H. Huls, J. Wang, M. A. J. Huijbregts, A.
782 M. Schipper, Context-dependent responses of terrestrial invertebrates to anthropogenic
783 nitrogen enrichment: A meta-analysis. *Glob. Change Biol.* **29**, 4161–4173 (2023).
- 784 118. C. Ganuza, S. Redlich, J. Uhler, C. Tobisch, S. Rojas-Botero, M. K. Peters, J. Zhang, C. S.
785 Benjamin, J. Englmeier, J. Ewald, U. Fricke, M. Haensel, J. Kollmann, R. Riebl, L. Uphus, J.
786 Müller, I. Steffan-Dewenter, Interactive effects of climate and land use on pollinator diversity
787 differ among taxa and scales. *Sci. Adv.* **8**, 9359 (2022).

- 788 119. J. Millard, C. L. Outhwaite, R. Kinnersley, R. Freeman, R. D. Gregory, O. Adedaja, S. Gavini, E.
789 Kioko, M. Kuhlmann, J. Ollerton, Z.-X. Ren, T. Newbold, Global effects of land-use intensity on
790 local pollinator biodiversity. *Nat. Commun.* 2021 121 **12**, 1–11 (2021).
- 791 120. B. A. Woodcock, J. M. Bullock, S. R. Mortimer, T. Brereton, J. W. Redhead, J. A. Thomas, R. F.
792 Pywell, Identifying time lags in the restoration of grassland butterfly communities: A multi-
793 site assessment. *Biol. Conserv.* **155**, 50–58 (2012).
- 794 121. Grimbacher, P.S., Catterall, C.P., How much do site age, habitat structure and spatial isolation
795 influence the restoration of rainforest beetle species assemblages? *Biol. Conserv.* **135**, 107–
796 118 (2007).
- 797 122. A. Purvis, T. Newbold, A. De Palma, S. Contu, S. L. L. Hill, K. Sanchez-Ortiz, H. R. P. Phillips, L. N.
798 Hudson, I. Lysenko, L. Börger, J. P. W. Scharlemann, Modelling and Projecting the Response of
799 Local Terrestrial Biodiversity Worldwide to Land Use and Related Pressures: The PREDICTS
800 Project. *Adv. Ecol. Res.* **58**, 201–241 (2018).
- 801 123. A. De Palma, M. Kuhlmann, R. Bugter, S. Ferrier, A. J. Hoskins, S. G. Potts, S. P. M. Roberts, O.
802 Schweiger, A. Purvis, Dimensions of biodiversity loss: Spatial mismatch in land-use impacts on
803 species, functional and phylogenetic diversity of European bees. *Divers. Distrib.* **23**, 1435–
804 1446 (2017).
- 805 124. C. Damgaard, A Critique of the Space-for-Time Substitution Practice in Community Ecology.
806 *Trends Ecol. Evol.* **34**, 416–421 (2019).
- 807 125. P. Batáry, R. Gallé, F. Riesch, C. Fischer, C. F. Dormann, O. Mußhoff, P. Császár, S. Fusaro, C.
808 Gayer, A.-K. Happe, K. Kurucz, D. Molnár, V. Rösch, A. Wietzke, T. Tschardtke, The former Iron
809 Curtain still drives biodiversity–profit trade-offs in German agriculture. *Nat. Ecol. Evol.* **1**,
810 1279–1284 (2017).
- 811 126. M. Soga, K. J. Gaston, Shifting baseline syndrome: causes, consequences, and implications.
812 *Front. Ecol. Environ.* **16**, 222–230 (2018).
- 813 127. H. Siviter, E. J. Bailes, C. D. Martin, T. R. Oliver, J. Koricheva, E. Leadbeater, M. J. F. Brown,
814 Agrochemicals interact synergistically to increase bee mortality. *Nature* **596**, 389–392 (2021).
- 815 128. F. Sánchez-Bayo, D. Goulson, F. Pennacchio, F. Nazzi, K. Goka, N. Desneux, Are bee diseases
816 linked to pesticides? — A brief review. *Environ. Int.* **89–90**, 7–11 (2016).
- 817 129. N. Mukherjee, J. Hugé, W. J. Sutherland, J. McNeill, M. Van Opstal, F. Dahdouh-Guebas, N.
818 Koedam, J. Hugé, W. J. Sutherland, J. McNeill, M. Van Opstal, F. Dahdouh-Guebas, N.
819 Koedam, The Delphi technique in ecology and biological conservation: applications and
820 guidelines. *Methods Ecol. Evol.* **6**, 1097–1109 (2015).
- 821 130. D. A. W. Miller, K. Pacifici, J. S. Sanderlin, B. J. Reich, The recent past and promising future for
822 data integration methods to estimate species' distributions. *Methods Ecol. Evol.* **10**, 22–37
823 (2019).
- 824 131. N. J. B. Isaac, M. A. Jarzyna, P. Keil, L. I. Dambly, P. H. Boersch-Supan, E. Browning, S. N.
825 Freeman, N. Golding, G. Guillera-Aroita, P. A. Henrys, S. Jarvis, J. Lahoz-Monfort, J. Pagel, O.
826 L. Pescott, R. Schmucki, E. G. Simmonds, R. B. O'Hara, Data Integration for Large-Scale Models
827 of Species Distributions. *Trends Ecol. Evol.* **35**, 56–67 (2020).

- 828 132. J. Kotta, J. Vanhatalo, H. Jänes, H. Orav-Kotta, L. Rugiu, V. Jormalainen, I. Bobsien, M.
829 Viitasalo, E. Virtanen, A. N. Sandman, M. Isaeus, S. Leidenberger, P. R. Jonsson, K.
830 Johannesson, Integrating experimental and distribution data to predict future species
831 patterns. *Sci. Rep.* **9**, 1821 (2019).
- 832 133. S. Domisch, A. M. Wilson, W. Jetz, Model-based integration of observed and expert-based
833 information for assessing the geographic and environmental distribution of freshwater
834 species. *Ecography* **39**, 1078–1088 (2016).
- 835 134. E. G. Simmonds, S. G. Jarvis, P. A. Henrys, N. J. B. Isaac, R. B. O’Hara, Is more data always
836 better? A simulation study of benefits and limitations of integrated distribution models.
837 *Ecography* **43**, 1413–1422 (2020).
- 838 135. K. Pacifici, B. J. Reich, D. A. W. Miller, B. S. Pease, Resolving misaligned spatial data with
839 integrated species distribution models. *Ecology*, e02709 (2019).
- 840 136. K. P. Adjei, C. Carvell, N. J. B. Isaac, F. Mancini, R. B. O’Hara, Integrating data from different
841 taxonomic resolutions to better estimate community alpha diversity. *Ecography* **n/a**, e07182
842 (2024).
- 843 137. S. Piirainen, A. Lehikoinen, M. Husby, J. A. Kålås, Å. Lindström, O. Ovaskainen, Species
844 distributions models may predict accurately future distributions but poorly how distributions
845 change: A critical perspective on model validation. *Divers. Distrib.* **29**, 654–665 (2023).
- 846 138. K. De Koning, J. Broekhuijsen, I. Kühn, O. Ovaskainen, F. Taubert, D. Endresen, D. Schigel, V.
847 Grimm, Digital twins: dynamic model-data fusion for ecology. *Trends Ecol. Evol.*,
848 S0169534723000903 (2023).
- 849 139. R. J. Fletcher, T. J. Hefley, E. P. Robertson, B. Zuckerberg, R. A. McCleery, R. M. Dorazio, A
850 practical guide for combining data to model species distributions. *Ecology*, e02710 (2019).
- 851 140. G. N. Daskalova, I. H. Myers-Smith, A. D. Bjorkman, S. A. Blowes, S. R. Supp, A. E. Magurran,
852 M. Dornelas, Landscape-scale forest loss as a catalyst of population and biodiversity change.
853 *Science* **368**, 1341–1347 (2020).
- 854 141. A. Greenop, B. A. Woodcock, C. L. Outhwaite, C. Carvell, R. F. Pywell, F. Mancini, F. K.
855 Edwards, A. C. Johnson, N. J. B. Isaac, Patterns of invertebrate functional diversity highlight
856 the vulnerability of ecosystem services over a 45-year period. *Curr. Biol.* **31**, 4627–4634.e3
857 (2021).
- 858 142. R. Cooke, F. Mancini, R. J. Boyd, K. L. Evans, A. Shaw, T. J. Webb, N. J. B. Isaac, Protected areas
859 support more species than unprotected areas in Great Britain, but lose them equally rapidly.
860 *Biol. Conserv.* **278**, 109884 (2023).
- 861 143. M. C. Urban, J. M. J. Travis, D. Zurell, P. L. Thompson, N. W. Synes, A. Scarpa, P. R. Peres-Neto,
862 A.-K. Malchow, P. M. A. James, D. Gravel, L. De Meester, C. Brown, G. Bocedi, C. H. Albert, A.
863 Gonzalez, A. P. Hendry, Coding for Life: Designing a Platform for Projecting and Protecting
864 Global Biodiversity. *BioScience* **72**, 91–104 (2022).
- 865 144. N. J. B. Isaac, A. J. van Strien, T. A. August, M. P. de Zeeuw, D. B. Roy, Statistics for citizen
866 science: extracting signals of change from noisy ecological data. *Methods Ecol. Evol.* **5**, 1052–
867 1060 (2014).

- 868 145. T. F. Johnson, A. P. Beckerman, D. Z. Childs, T. J. Webb, K. L. Evans, C. A. Griffiths, P. Capdevila,
869 C. F. Clements, M. Besson, R. D. Gregory, G. H. Thomas, E. Delmas, R. P. Freckleton, Revealing
870 uncertainty in the status of biodiversity change. *Nature*, 1–7 (2024).
- 871 146. B. Leung, A. Gonzalez, Global monitoring for biodiversity: Uncertainty, risk, and power
872 analyses to support trend change detection. *Sci. Adv.* **10**, eadj1448 (2024).
- 873 147. C. Wyborn, M. C. Evans, Conservation needs to break free from global priority mapping. *Nat.*
874 *Ecol. Evol.* **2021** 510 **5**, 1322–1324 (2021).
- 875 148. K. L. Yates, P. J. Bouchet, M. J. Caley, K. Mengersen, C. F. Randin, S. Parnell, A. H. Fielding, A. J.
876 Bamford, S. Ban, A. M. Barbosa, C. F. Dormann, J. Elith, C. B. Embling, G. N. Ervin, R. Fisher, S.
877 Gould, R. F. Graf, E. J. Grev, P. N. Halpin, R. K. Heikkinen, S. Heinänen, A. R. Jones, P. K.
878 Krishnakumar, V. Lauria, H. Lozano-Montes, L. Mannocci, C. Mellin, M. B. Mesgaran, E.
879 Moreno-Amat, S. Mormede, E. Novaczek, S. Oppel, G. Ortuño Crespo, A. T. Peterson, G.
880 Rapacciuolo, J. J. Roberts, R. E. Ross, K. L. Scales, D. Schoeman, P. Snelgrove, G. Sundblad, W.
881 Thuiller, L. G. Torres, H. Verbruggen, L. Wang, S. Wenger, M. J. Whittingham, Y. Zharikov, D.
882 Zurell, A. M. M. Sequeira, Outstanding Challenges in the Transferability of Ecological Models.
883 *Trends Ecol. Evol.* **33**, 790–802 (2018).
- 884 149. H. Meyer, E. Pebesma, Predicting into unknown space? Estimating the area of applicability of
885 spatial prediction models. *Methods Ecol. Evol.* **12**, 1620–1633 (2021).
- 886 150. S. H. M. Butchart, M. Walpole, B. Collen, A. van Strien, J. P. W. Scharlemann, R. E. A. Almond,
887 J. E. M. Baillie, B. Bomhard, C. Brown, J. Bruno, K. E. Carpenter, G. M. Carr, J. Chanson, A. M.
888 Chenery, J. Csirke, N. C. Davidson, F. Dentener, M. Foster, A. Galli, J. N. Galloway, P. Genovesi,
889 R. D. Gregory, M. Hockings, V. Kapos, J.-F. Lamarque, F. Leverington, J. Loh, M. a McGeoch, L.
890 McRae, A. Minasyan, M. Hernández Morcillo, T. E. E. Oldfield, D. Pauly, S. Quader, C. Revenga,
891 J. R. Sauer, B. Skolnik, D. Spear, D. Stanwell-Smith, S. N. Stuart, A. Symes, M. Tierney, T. D.
892 Tyrrell, J.-C. Vié, R. Watson, Global biodiversity: indicators of recent declines. *Science* **328**,
893 1164–8 (2010).

894 Acknowledgments

895 **Funding:** All authors were supported by award numbers NE/V007548/1, NE/V007173/1,
896 NE/V006533/1, NE/V006800/1 and NE/V006886/1 from the UK Natural Environment Research
897 Council. JM is now supported by an Early Career Fellowship from the Leverhulme Trust.

898

899 Author contributions:

900 Conceptualization: All authors

901 Formal analysis: RC

902 Writing - Original Draft: RC, CLO, AJB, JM, JGR, ZD, EED, TN, AP, BAW, NJBI

903 Writing - Review & Editing: All authors

904 Visualization: RC, CLO, NJBI

905 Project administration: RC, CLO, NJBI

906 Funding acquisition: CLO, LVD, CH, JIJ, TN, AP, HER, BAW, NJBI

907

908 **Competing interests:** Dr. Lynn Dicks is a Board Member of Natural England, the UK Government's
909 statutory advisor for the natural environment in England. Dr. Nick Isaac is a member of the
910 Biodiversity Expert Committee of the UK Government Department of Agriculture, Food and Rural
911 Affairs, and of the European Commission expert group on Pollinating Insects.

912

913 **Data and materials availability:** All data used in the manuscript are previously published (89).

Figure Legends

Fig. 1. The complexity of insects in a changing world and the role of interactions. The center ring shows an example insect lifecycle, with curved arrows between life stages (here, larva, pupa, adult). Connected to the lifecycle are a simplified number of interacting species including a variety of taxonomic groups. The middle ring represents the landscape in which this lifecycle and the local interactions occur, often incorporating multiple habitat types and land uses. The outer ring shows some of the major drivers that act across the landscape and local scales, affecting the species found in those areas. These can act alone but can also interact.

Fig. 2. Frequency vs span of time series in a large insect time series database. The points represent the 1,657 sites (i.e., time series) included in the InsectChange database (89), which underpins a large-scale meta-analysis on insect trends (11). Note that only time series covering ~10 or more years were included in this database. The color underneath the points reflects the density of time series across the plot (red = highest density, white = lowest density). The lines represent approximately annual sampling (i.e., 10 sample events for a 10-year time series), ~seasonal sampling (i.e., 40 sample events for a 10-year time series), ~monthly sampling (i.e., 120 sample events for a 10-year time series), ~weekly sampling (i.e., 520 sample events for a 10-year time series).

Fig. 3. The power of attribution and breadth of coverage of different evidence types for insect biodiversity change across temporal, spatial/environmental, and taxonomic dimensions. Power of attribution refers to the potential for evidence types to identify causal links between insect biodiversity change and any external driver based on existing data. Experiments have high power because they are designed to quantify specific effects, whereas time series are typically set up without regard to the drivers present. Coverage captures the extent, depth, and representativeness of the evidence type in the three dimensions of time, space, and taxonomy. For example, taxonomic coverage reflects the number of insect Orders included (extent), the number of species within Orders (depth), and the representation of species across Orders (representativeness). Although inherently subjective, positioning of the shapes illustrates our interpretation of the strengths and weaknesses of the four evidence types; the size of each shape indicates the approximate variation in power of attribution and coverage within each evidence type. N.B. Evidence types are not mutually exclusive, e.g., some studies can be both experiments and time series, depending on how data were collected and analyzed.

Fig. 4. Properties of available data for understanding insect biodiversity change and underlying drivers. Grades and shading (A – greatest, dark to D – least, light) reflect the fulfilment of the ideal property for each evidence type. Each box contains our rationale for the assignment of grades to each property and evidence type.

Fig. 5. Establishing a synoptic overview of insect biodiversity change to inform policy and conservation action. Integrating multiple evidence types on the state of and changes in insect biodiversity with information on the drivers of this change can provide a synoptic overview of current patterns. This overview would underpin projections of insect biodiversity change across space and through time, as well as of the potential consequences of insect biodiversity change.