

*Two new Globicornis Latreille species
from eastern Mediterranean (Coleoptera:
Dermestidae: Megatominae)*

Article

Published Version

Creative Commons: Attribution-Noncommercial 4.0

Open Access

Holloway, G. J. and Herrmann, A. (2025) Two new Globicornis Latreille species from eastern Mediterranean (Coleoptera: Dermestidae: Megatominae). Insecta Mundi, 1141. pp. 1-10. ISSN 1942-1354 doi: 10.64338/im.1141.4c4u7 Available at <https://centaur.reading.ac.uk/125545/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

Identification Number/DOI: 10.64338/im.1141.4c4u7
<<https://doi.org/10.64338/im.1141.4c4u7>>

Publisher: Center for Systematic Entomology, Inc.

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

A journal of world insect systematics

INSECTA MUNDI

1141

Two new *Globicornis* Latreille species from eastern
Mediterranean (Coleoptera: Dermestidae: Megatominae)

Graham J. Holloway

Cole Museum of Zoology, Biological Sciences, HLS Building, University of Reading, Whiteknights, Reading RG6 6EX, UK

Andreas Herrmann

Bremervörder Strasse 123, 21682 Stade, Germany

Date of issue: August 29, 2025

Center for Systematic Entomology, Inc., Gainesville, FL

Holloway GJ, Herrmann A. 2025. Two new *Globicornis* Latreille species from eastern Mediterranean (Coleoptera: Dermestidae: Megatominae). *Insecta Mundi* 1141: 1–10.

Published on August 29, 2025 by
Center for Systematic Entomology, Inc.
P.O. Box 141874
Gainesville, FL 32614-1874 USA
<http://centerforsystematicentomology.org/>

INSECTA MUNDI is a journal primarily devoted to insect systematics, but articles can be published on any non-marine arthropod. Topics considered for publication include systematics, taxonomy, nomenclature, checklists, faunal works, and natural history. *Insecta Mundi* will not consider works in the applied sciences (i.e. medical entomology, pest control research, etc.), and no longer publishes book reviews or editorials. *Insecta Mundi* publishes original research or discoveries in an inexpensive and timely manner, distributing them free via open access on the internet on the date of publication.

Insecta Mundi is referenced or abstracted by several sources, including the Zoological Record and CAB Abstracts. *Insecta Mundi* is published irregularly throughout the year, with completed manuscripts assigned an individual number. Manuscripts must be peer reviewed prior to submission, after which they are reviewed by the editorial board to ensure quality. One author of each submitted manuscript must be a current member of the Center for Systematic Entomology.

Guidelines and requirements for the preparation of manuscripts are available on the *Insecta Mundi* website at <http://centerforsystematicentomology.org/insectamundi/>

Chief Editor: David Plotkin, insectamundi@gmail.com

Assistant Editor: Paul E. Skelley, insectamundi@gmail.com

Layout Editor: Robert G. Forsyth

Editorial Board: Davide Dal Pos, José Martínez, Alex Orfinger, M. J. Paulsen, Felipe Soto-Adames

Founding Editors: Ross H. Arnett, Jr., J. H. Frank, Virendra Gupta, John B. Heppner, Lionel A. Stange, Michael C. Thomas, Robert E. Woodruff

Review Editors: Listed on the *Insecta Mundi* webpage

Printed copies (ISSN 0749-6737) annually deposited in libraries

Florida Department of Agriculture and Consumer Services, Gainesville, FL, USA
The Natural History Museum, London, UK
National Museum of Natural History, Smithsonian Institution, Washington, DC, USA
Zoological Institute of Russian Academy of Sciences, Saint-Petersburg, Russia

Electronic copies (online ISSN 1942-1354) in PDF format

Archived digitally by Portico.

Florida Virtual Campus: <http://purl.fcla.edu/fcla/insectamundi>

University of Nebraska-Lincoln, Digital Commons: <http://digitalcommons.unl.edu/insectamundi/>

Goethe-Universität, Frankfurt am Main: <http://nbn-resolving.de/urn/resolver.pl?urn:nbn:de:hebis:30:3-135240>

This is an open access article distributed under the terms of the Creative Commons, Attribution Non-Commercial License, which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.
<https://creativecommons.org/licenses/by-nc/3.0/>

Two new *Globicornis* Latreille species from eastern Mediterranean (Coleoptera: Dermestidae: Megatominae)

Graham J. Holloway

Cole Museum of Zoology, Biological Sciences, HLS Building, University of Reading, Whiteknights, Reading RG6 6EX, UK
g.holloway@reading.ac.uk

https://orcid.org/0000-0003-0495-0313

Andreas Herrmann

Bremervörder Strasse 123, 21682 Stade, Germany
herrmann@coleopterologie.de

https://orcid.org/0000-0001-5700-1125

Abstract. Two new species, *Globicornis* (*Globicornis*) *busei* Holloway and Herrmann and *Globicornis* (*Globicornis*) *lienemanni* Holloway and Herrmann (Coleoptera: Dermestidae: Megatominae) from eastern Mediterranean are described and compared with *G. picta* (Küster, 1851), the most likely confounding species because of geographical overlap and external similarity. Images of the habitus, ventrites, antenna, aedeagus, and sternite IX are presented. Ten *Globicornis* (*Globicornis*) Latreille, 1829 species are now known from the eastern Mediterranean.

Key words. Taxonomy, dissection, genitalia, *busei*, *lienemanni*, *picta*.

ZooBank registration. urn:lsid:zoobank.org:pub:A93526EA-9A01-4ADE-AFBB-25FA451B41E9

DOI. https://doi.org/10.64338/im.1141.4c4u7

Introduction

The family Dermestidae Latreille, 1804 contains 1930 species (Háva 2024). The number of species has grown by about 1000 since Mroczkowski (1968), an increase of about 55% in the space of just 60 years, with most species discovered since the year 2000 (Háva 2024). The largest subfamily within the Dermestidae is Megatominae Leach, 1815, and within Megatominae sits the genus *Globicornis* Latreille, 1829. *Globicornis* is split into four subgenera: *Elania* Mulsant and Rey, 1868 with two species, *Globicornis* with 20 species, *Hadrotoma* Erichson, 1846 with six species, and *Pseudomesalia* Ganglbauer, 1900 with five species, a total of 33 species within the genus *Globicornis*. *Globicornis* is entirely Palaearctic with species distributed from western Europe and into central Asia. Only one species, *G. yushuensis* Háva, 2016 occurs as far east as China (Háva 2024).

The earliest species described was *G. nigripes* (Fabricius, 1792) with a further two species described prior to 1850. An additional 14 species were added between 1851 and 1900, and another six species were added during the 20th century. Only 10 new *Globicornis* species have been described during the 21st century, well below the rate of species description across the Dermestidae as a whole, suggesting perhaps that *Globicornis* hasn't received as much attention as some other genera within the family. The most recently described species is *G. peckhamae* Holloway and Cañada Luna, 2023. Holloway and Cañada Luna (2023) described *G. peckhamae* from Mallorca, splitting the new species from *G. bifasciata* (Perris, 1866), and suggested a reason for the lack of recent species discovery. *Globicornis bifasciata* and *G. peckhamae* are very similar externally, in particular with respect to the possession of a large, characteristic antennal club, reinforcing the belief that the species was well-known and providing little incentive to look further than this. Indeed, Holloway and Cañada Luna (2023) only discovered *G. peckhamae* whilst carrying out work to better describe *G. bifasciata*; the discovery of a second species was unexpected. This raises the possibility that the ranges of other named *Globicornis* species encompass several look-alike species. This situation has been found in other Megatominae, including *Anthrenus pimpinellae* (Fabricius, 1775) from which similar western Palaearctic species have been split (Holloway 2019, 2020, 2024a, 2024b), and the *A. maculifer* group of species from Asia (Holloway 2025). Studies like these facilitate a better understanding of true distribution (Holloway et al. 2021, 2023, 2025; Holloway and Herrmann 2024).

Globicornis picta (Küster, 1851) is another species from the subgenus *Globicornis* to be described in the mid-19th century (along with *G. variegata* (Küster, 1851), although Küster initially assigned both species to *Trogoderma* (Dejean, 1821)). *Globicornis picta* is believed to have one of the widest ranges within the genus, ranging from Italy in the west, east into Turkey and south to Egypt (Háva 2024), making it a good species for further investigative study. The purpose of the current study was to provide a more extensive description of *G. picta* and to consider whether more than one species exists within the reported *G. picta* distribution (Háva 2024). In addition to *G. picta*, two new species were found and are described herein: *G. busei* and *G. lienemanni*.

Materials and Methods

Thirty-three specimens from Andreas Herrmann's private collection (AHEC) were examined although dissection and species recognition was based on the ten male specimens.

All specimens were macerated in a solution of 2% acetic acid for five days to allow removal from staging prior to dissection. Dissection was carried out under a Brunel BMSL zoom stereo LED microscope and involved detaching the abdomen from the rest of the insect using two entomological pins. The soft tergites were then peeled away from the harder ventrites to expose the genitalia. The aedeagus was detached from the ring sclerite, and then sternite IX was detached from the ring sclerite and the aedeagus. Prior to imaging, aedeagi were cleaned in 5% KOH for 12 hours. Images of habitus, both upper and lower sides, were captured at ×20 magnification using a Canon EOS 2000D camera mounted on the BMSL microscope. Images of aedeagi and sternite IX were captured at ×200 magnification for measurement using a Canon EOS 1300D camera mounted on a Brunel monocular SP28 microscope. After dissection, all body parts were mounted on card. The antennae were teased out and images were taken at ×200 magnification through the SP28 microscope. All images were fed through Helicon Focus Pro version 8.2.2 focus-stacking software. Habitus measurements were made using a calibrated eyepiece. All other measurements were made using DsCap.Ink software version 3.90. Measurements taken:

- Body length (BL): distance from anterior margin of pronotum to the apex of the elytra.
- Paramere length (PL): distance from the anterior end of the parameres to the apex of the parameres
- Median lobe length (ML): distance from posterior tip to tip of one anterior stirrup.
- Sternite IX length (SL): distance from the tip of one anterior horn to the tip of the posterior lobe

The data for the distribution maps (Shorthouse 2010) were derived from the data labels on the studied specimens and Háva (2024). Scale bars were added using ImageJ 1.53M (Schneider et al. 2014).

Results

Taxonomy

Dermestidae Latreille, 1803

Megatominae Dalla Torre, 1911

Megatomini Leach, 1815

Megatomina Leach, 1815

Globicornis Latreille in Cuvier, 1829

Globicornis (*Globicornis*) *busei* Holloway and Herrmann, new species

(Fig. 1–2)

Type specimens. Holotype. Male. Israel, Upper Galilee, Biriya Forest/Safed, 32.97N, 35.47E. Window trap on *Pinus brutia*. 14.v.2007, J. Buse leg. AHEC.

Paratypes. None.

Other material examined. Male, Cyprus, Nicosia district, Cedar Valley, 35.03, 32.66, 1200 m. *Quercus alnifolia* 7.v.2014, David Frank leg. Male, Southern Turkey, Umg. Kemer, 16.v.1993, A. Weigel leg. Male, Cyprus, Panagia/Paphos, 34.919, 32.631. 14.iv.2010, W. Ziegler leg. AHEC.




Figure 1. *Globicornis busei* sp. nov., holotype. **A)** Habitus (scale bar = 1 mm). **B)** Ventrites (scale bar = 1 mm). **C)** Antenna (scale bar = 100 μ m).

Description, external characteristics. Holotype *Globicornis* (*Globicornis*) *busei* (Fig. 1A), BL = 3.2 mm, BW = 1.65 mm). Single brown ocellus just above level of top of eyes. Eyes oblong, margins complete, bulbous. Integument of head and pronotum dark brown, sculptured, pustulate. Face from ocellus, laterally to eyes and down to labrum with yellowish hairs. Lateral pronotal margins evenly curved to anterior angles, anterior margin slightly sinuate, posterior margin sinuate with shallow projection above scutellar shield. Dark brown scutellar shield equilateral triangle. Pronotum with yellow hairs, slightly paler, slightly flattened (scale like), and denser at pronotal hind angles, yellower, not flattened hairs around anterior angles and across anterior margin, and above scutellar shield, sparser yellow hairs on disk of pronotum.

Elytral integument dark brown at base, becoming paler towards apices. Three reddish fasciae across elytra: pre-medial, medial, and post-medial. Pre-medial fascia from lateral margin complete, sweeping downwards and then up towards but not reaching scutellum. Medial fascia consists of a series of spots. Post-medial fascia consists of two larger spots on each elytron. Medial and post-medial fasciae linked by reddish integument running along elytral margin. Reddish margin continues to elytral apices from post-medial fascia to form large apical spots, Apical spots and post-medial fascia also connected by reddish integument along elytral suture. Reddish integument covered in scale-like whitish hairs, brown integument covered in brown hairs. Ventrites (Fig. 1B) shining, mid-brown, covered in evenly spaced, short, yellowish hairs.

Antenna with 10 antennomeres (Fig. 1C) with mid-brown, globular antennomeres 1 and 2. Antennomeres 3–7 yellowish, antennomeres 3 and 4 very elongate, antennomere 5 slightly elongate, antennomere 6 transverse, antennomere 7 disc-shaped with brown edging to disc, antennomeres 8–10 brown, antennomeres 8 and 9 disc-shaped, antennomere 10 triangular with 90° angle between posterior and inner lateral margin, connected by an evenly curved anterior margin. Femora brown, tibiae and tarsi red.

Description, internal characteristics. The overall shape of the aedeagus (Fig. 2A) is broad oval, especially evident in ventral aspect. Parameres (PL = 466 μ m) connected at anterior base by two narrow rods. From anterior, parameres diverge and $\frac{1}{3}$ along broaden and continue to curve inwards at posterior apices. Inner surfaces of parameres with long, straight, posteriad angled hairs that continue posterior becoming progressively shorter to paramere



Figure 2. *Globicornis busei* sp. nov., holotype. A) Aedeagus. B) Sternite IX. Scale bar = 100 μ m in both cases.

tips. Anterior end of parameres pale brown, becoming progressively paler to whitish at rounded posterior tips. Paramere tips point distinctly towards posterior without curving over the top of the median lobe. Medium lobe (ML = 351 μ m) with steadily converging, pale brown, rod-like margins, inner margins of rods connected by pale, thin membrane. Marginal rods widely spaced at base, curve dorsally, joining at posterior apex forming a hook and falling just short of the paramere tips. Median lobe attached to parameres about $\frac{1}{3}$ along from anterior and terminates in two straight, short stirrups. Sternite IX (Fig. 2B; SL = 512 μ m) with single anterior attachment point, that extends posteriad as two narrow marginal rods that diverge forming concave outer margins for first $\frac{1}{2}$ of sternite. From widest point and towards posterior, concave lateral margins become membranous and form rounded, slightly pointed posterior margin. Membranous margin carrying small number of stout, dark setae with about 10 such setae away from margin on sternite disk. Sternite membrane very pale brown in anterior half, paler, almost white in posterior half.

Etymology. The species is named after the collector of the holotype, the coleopterist Dr. Jörn Buse (Germany).

***Globicornis (Globicornis) lienemanni* Holloway and Herrmann, new species**

(Fig. 3–4)

Type specimens. Holotype. Male. Cyprus, Kyrenia, Magar, 600 m, 35.292, 33.521. *Pastinaca savita*, 2.vi.1992, K. Lienemann, leg. AHEC.

Paratype. Female, same data as holotype, AHEC.

Other material examined. Male, Turkey, Mersin, 36.798, 34.63. 6.v.2004, V. Brachat H. Meybohm, leg. AHEC.

Description, external characteristics. Holotype *Globicornis (Globicornis) lienemanni* (Fig. 3A), BL = 2.4 mm, BW = 1.25 mm). Single brown ocellus above level of top of eyes. Eyes oblong, margins complete, bulbous. Integument of head and pronotum dark brown, sculptured, pustulate. A few weak yellowish hairs of face. Lateral pronotal margins evenly curved to anterior angles, anterior margin slightly sinuate, posterior margin sinuate with shallow projection above scutellar shield. Dark brown scutellar shield equilateral triangle. Pronotum with yellow hairs, slightly paler, slightly flattened (scale like) towards pronotal hind angles, yellow, (not flattened) hairs around anterior angles and across posterior margin above scutellar shield. Very few yellow hairs on anterior part of disk of pronotum, apart from a few hairs on anterior pronotal corners.

Elytral integument dark brown at base, becoming paler towards apices. Three reddish fasciae across elytra: pre-medial, medial, and post-medial. No fascia reaching lateral margin. Pre-medial fascia complete, sweeping downwards from sub-margin, and then up towards but not reaching scutellum. Medial fascia like pre-medial fascia. Post-medial fascia consists of tightly connected spots. Fasciae not connected with strips of clear reddish integument along the lateral margin or elytral suture. Ventrites (Fig. 3B) shining, mid-brown, covered in evenly spaced, yellowish hairs.

Antenna with 10 antennomeres (Fig. 3C) with mid-brown, globular antennomeres 1 and 2. Antennomeres 3–7 red, antennomeres 3 and 4 elongate, antennomere 5 slightly elongate, antennomeres 6 and 7 transverse, antennomeres 8–10 brown, antennomeres 8 and 9 disc-shaped, antennomere 10 rounded (Antennomere 10 folded inwards which likely affects its appearance). Femora brown, tibiae and tarsi red.

Description, internal characteristics. Overall shape of aedeagus (Fig. 4A) elongated oblong. Parameres (PL = 415 μ m) connected at anterior base by two narrow rods. From anterior, parameres diverge and $\frac{1}{3}$ along broaden,




Figure 3. *Globicornis lienemanni* sp. nov., holotype. **A)** Habitus (scale bar = 1 mm). **B)** Ventrites (scale bar = 1 mm). **C)** Antenna (scale bar = 100 μ m).




Figure 4. *Globicornis lienemanni* sp. nov., holotype. **A)** Aedeagus. **B)** Sternite IX. Scale bar = 100 µm in both cases.

run parallel to each other, and curve inwards at posterior apices. Straight setae along inner margin of parameres angled posteriad, long setae on paramere tips especially on outer surface. Paramere tips narrow and curve inwards over median lobe. Medium lobe (ML = 317 µm) with brown, rod-like margins that run parallel for $\frac{2}{3}$ length before converging, curving dorsally and ending in hook at posterior end. Median lobe attached to parameres less than $\frac{1}{3}$ along from anterior and terminates in two straight, short stirrups. Sternite IX (Fig. 4B; SL = 391 µm) with short, single anterior attachment point. Lateral margins diverge from attachment point for less than $\frac{1}{2}$ of sternite. From widest point, lateral margins are straight and converge slightly to slightly extended lobe with flat (or notched) posterior margin. Short, black setae mostly restricted to posterior lobe extension, mostly at margins with about 10 setae scattered across disc of posterior lobe. Sternite membrane very pale brown, slightly paler towards posterior margin.

Etymology. The species is named after the collector of the holotype, Dr. Klaus Lienemann (Germany).

Differential diagnosis. Both *G. busei* and *G. lienemanni* occur within the presumed range of *G. picta* (Háva 2024) and were mixed in with a sample of specimens all tentatively identified as *G. picta*. Externally, from the sample of specimens examined there was little to differentiate the three species from each other, although *G. picta* tended to have fewer pale setae on the pronotum than *G. busei* and *G. lienemanni* (Fig. 5). Of course, there




Figure 5. *Globicornis picta*. **A)** Habitus (scale bar = 1 mm). **B)** Ventriles (scale bar = 1 mm). **C)** Antenna (scale bar = 100 µm).

is often colour pattern variation in Dermestidae (e.g., Holloway et al. 2022), so a larger sample of the three species will be required to confirm whether identification is possible without dissection. Dissection and inspection of the aedeagus remains the most reliable way to differentiate between *G. picta*, *G. busei*, and *G. lienemanni*. The overall aedeagus shape in *G. lienemanni* (Fig. 4) and *G. picta* (Fig. 6) is less oval than *G. busei* (Fig. 2). In *G. picta*, the median lobe is not as broad as *G. busei* and wider than *G. lienemanni*. The hairs on *G. picta* paramere inner margin are evenly spaced, but longer, denser and more brush-like in *G. busei* and *G. lienemanni*.

Distribution. Figure 7 illustrates the distribution of the specimens from the current study overlaying the distribution reported by Háva (2024).

Discussion

Globicornis (*Globicornis*) *picta* is considered to have the widest distribution of the subgenus *Globicornis*, a region of about 3000 × 4000 km, so there was a chance that more than one species fell under the name *G. picta* from that very large area. From the sample of specimens examined, three species were found: *G. picta* from the western section of the distribution, *G. busei* from the eastern section of the distribution, and *G. lienemanni* from Cyprus and into Turkey. The sample size of specimens examined was relatively small, so a more extensive survey might reveal greater species distribution overlap.

Externally, the three species described here are very similar and the main differences are found in the male genitalia. The current study illustrates a further example of cryptic species within Dermestidae and bears similarity to the *Globicornis* study from Mallorca where a cryptic species, *G. peckhamae*, was found within the *G. bifasciata* range (Holloway and Canada Luna 2023). The discovery of two new species brings the number of known *Globicornis* (*Globicornis*) species from the Mediterranean region to 15, those from eastern Mediterranean (east of Croatia: 10) are in bold:

Globicornis bifasciata (Perris, 1866)

***Globicornis breviclavus* Reitter in Schneider and Leder, 1878**

***Globicornis busei* Holloway and Herrmann, new species**




Figure 6. *Globicornis picta*. A) Aedeagus. B) Sternite IX. Scale bar = 100 μ m in both cases.

***Globicornis cypriensis* Háva and Herrmann, 2014**

Globicornis fasciata (Fairmaire, 1859)

***Globicornis kaffai* Háva, 2000**

***Globicornis lienemanni* Holloway and Herrmann, new species**

Globicornis luckowi Herrmann, Háva and Kadej, 2011

Globicornis maroccana Kadej, Háva, Herrmann, Benyahia and Brustal, 2016

***Globicornis nigripes* (Fabricius, 1792)**

***Globicornis picta* Küster, 1851**

***Globicornis rufoguttatus* Pic, 1899**

***Globicornis signatipennis* Pic, 1899**

***Globicornis tristis* Reitter, 1881**




Figure 7. Specimen localities of the three *Globicornis* species discussed in this study, plotted on the distribution of *Globicornis picta* according to Háva (2024).

Globicornis variegata (Küster, 1851)

All additional eastern Mediterranean species highlighted above differ from the new species described in the current study by having a very large, triangular terminal antennomere, and/or having fewer than three orange fasciae, or, in the case of *G. tristis*, having a bulbous median lobe (Herrmann 2025).

Acknowledgments

The authors are very grateful to everybody who donated specimens to AHEC which makes studies, such as the one reported here, possible. The authors are also grateful to Dr C.W. Foster, Ivan Cañada Luna and the editors for very constructive comments of how the manuscript could be improved.

Literature Cited

- Háva J. 2024.** Dermestidae World (Coleoptera). World Dermestidae | Dermestidae world (Coleoptera), Megatominae. Available at <https://dermestidae.wz.cz/wp-content/uploads/2023/04/Subfamily-Megatominae.pdf> (Last accessed February 2025.)
- Herrmann A. 2025.** Dermestidae (Coleoptera) of the World. Dermestidae (Coleoptera)—Homepage of Andreas Herrmann. Available at <http://www.dermestidae.com/> (Last accessed March 2025.)
- Holloway GJ. 2019.** *Anthrenus* (s. str.) *amandae* (Coleoptera: Dermestidae): a new species from Mallorca, Spain. Zootaxa 4543(4): 595–599. <https://doi.org/10.11646/zootaxa.4543.4.9>
- Holloway GJ. 2020.** *Anthrenus* (s. str.) *chikatunovi* (Coleoptera: Dermestidae): a new species from southern France. Israel Journal of Entomology 50: 69–75. <https://doi.org/10.5281/zenodo.4088743>
- Holloway GJ. 2024a.** *Anthrenus* (*Anthrenus*) *quernereri* (Coleoptera: Dermestidae: Megatominae), a new species from Austria. Insecta Mundi 1060: 1–6.
- Holloway GJ. 2024b.** *Anthrenus* (*Anthrenus*) *algeriensis* (Coleoptera, Dermestidae, Megatominae), a new species from Algeria. Baltic Journal of Coleopterology 24(1): 33–41. [https://doi.org/10.59893/bjc.24\(1\).004](https://doi.org/10.59893/bjc.24(1).004)
- Holloway GJ. 2025.** A review of the *Anthrenus maculifer* Reitter, 1881 species group (Coleoptera: Dermestidae: Megatominae). Insecta Mundi 1118: 1–13.

- Holloway GJ, Bakaloudis DE, Cocks L. 2022.** Colour pattern plasticity in *Anthrenus isabellinus* (Coleoptera, Dermestidae). *Bulletin of Insectology* 75(1): 131–136.
- Holloway GJ, Bakaloudis DE, Foster CW. 2021.** *Anthrenus dorsatus* new to the United States and a comparison with *Anthrenus pimpinellae* ssp. *pimpinellae* (Coleoptera: Dermestidae). *Journal of the Kansas Entomological Society* 93(2): 153–163. <https://doi.org/10.2317/0022-8567-93.2.153>
- Holloway GJ, Cañada Luna I. 2023.** *Globicornis peckhamae* (Coleoptera, Dermestidae, Megatominae), a new species from Mallorca, and a description of *Globicornis bifasciata*. *Zootaxa* 5306(2): 297–300. <https://doi.org/10.11646/zootaxa.5306.2.9>
- Holloway GJ, Herrmann A. 2024.** A new species of the genus *Anthrenus* Geoffroy, 1762 (Coleoptera: Dermestidae) from Turkey and Lebanon. *Annales Zoologici* 74(4): 641–649. <https://doi.org/10.3161/00034541ANZ2024.74.4.008>
- Holloway GJ, Maclure CJ, Foster CW. 2023.** Palaearctic distributions of *Anthrenus pimpinellae* (Fabricius) and *Anthrenus isabellinus* Küster (Coleoptera: Dermestidae). *Entomologist's Monthly Magazine* 159(4): 23–28. <https://doi.org/0.31184/M00138908.1594.4207>
- Mroczkowski M. 1968.** Distribution of the Dermestidae (Coleoptera) of the world with a catalogue of all known species. *Annales Zoologici* 26(3): 15–192.
- Schneider CA, Rasband WS, Eliceiri KW. 2012.** NIH Image to ImageJ: 25 years of image analysis. *Nature Methods* 9: 671–675. <https://doi.org/10.1038/nmeth.2089>
- Shorthouse DP. 2010.** SimpleMappr, an online tool to produce publication-quality point maps. Available at <https://www.simplemappr.net> (Last accessed March 2025.)

Received June 23, 2025; accepted August 1, 2025.

Review editor David Plotkin.