

Beachy Head Woman: clarifying her origins using a multiproxy anthropological and biomolecular approach

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Walton, A., Marsh, W., Strang, A., Seaman, J., Van Doorn, K., Eckardt, H. ORCID: <https://orcid.org/0000-0001-9288-5624>, Wilkinson, C., Barnes, I. and Brace, S. (2026) Beachy Head Woman: clarifying her origins using a multiproxy anthropological and biomolecular approach. *Journal of Archaeological Science*, 186. 106445. ISSN 1095-9238 doi: 10.1016/j.jas.2025.106445 Available at <https://centaur.reading.ac.uk/127376/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.jas.2025.106445>

Publisher: Elsevier



All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online



Beachy Head Woman: clarifying her origins using a multiproxy anthropological and biomolecular approach

Andy Walton ^{a,b,1} , William Marsh ^{a,c,1} , Alex Strang ^{a,b,c} , Jonathan Seaman ^d, Kelly Van Doorn ^e, Hella Eckardt ^f , Caroline Wilkinson ^g , Ian Barnes ^{a,c} , Selina Brace ^{a,c,*}

^a Scientific Research, Natural History Museum, London, UK

^b UCL Genetics Institute, Department of Genetics, Evolution and Environment, University College London, London, UK

^c Centre for Human Evolutionary Research, Natural History Museum, London, UK

^d Mint House, High Street, Pevensey, UK

^e Heritage Eastbourne, Eastbourne Borough Council, Town Hall, Grove Road, Eastbourne, UK

^f Department of Archaeology, University of Reading, Reading, UK

^g Forensic Research Institute and Face Lab, Liverpool John Moores University, Liverpool, UK

ARTICLE INFO

Keywords:

Ancient DNA
Roman Britain

ABSTRACT

The skeletal remains of an individual colloquially referred to as Beachy Head Woman (BHW) were re-discovered in the Eastbourne Town Hall collection in 2012, and have remained the subject of significant public interest since. Radiocarbon dating yielded a calibrated date of between 129 and 311 calCE indicating that she lived during the period of the Roman occupation of Britain and, over more than a decade, there have been several attempts to unravel her geographical origins and ancestry. Here, we present results of all bioanthropological and biomolecular analyses performed to date. Initial osteological analyses indicated possible sub-Saharan origin, with BHW thus presented as one of the earliest Africans in Britain. However, her story was complicated by subsequent (unpublished) biomolecular analyses, which suggested she likely grew up on the south coast of Britain and had recent European ancestry. Here we present high quality ancient DNA data indicating that Beachy Head Woman has a strong genetic affinity to individuals from rural Britain during the Roman occupation and modern day Britons. We find no signals of admixture that would suggest recent sub-Saharan ancestry. Phenotypic predictions suggest she had blue eyes, intermediate (between pale and dark) skin pigmentation and light hair. Combined, our multiproxy approach indicates that Beachy Head Woman was of local British ancestry.

1. Introduction

In the early 2010s, the 'Eastbourne Ancestors Project' (EAP) sought to reveal the stories of over 200 skeletons in the collection of Eastbourne Council in the UK (Seaman and Forsyth, 2014). As part of this project, in 2012, a skeleton was discovered in a box in the basement of Eastbourne Town Hall. The only contextual information available for the remains was an English Heritage label detailed 'Beachy Head (1959)'. Despite an extensive search of museum archives, no other contextual information or records of any associated excavation could be uncovered. As the skeleton had been morphologically assessed to be female, it was dubbed 'Beachy Head Woman' (BHW), with Beachy Head referring to the stretch of coastline just outside Eastbourne in the South of England (Fig. 1).

BHW's origins would soon garner public interest when results of early (2013) craniofacial analyses (see 1.3 Bioarchaeological Assessment and S.1.2) suggested she was of recent sub-Saharan ancestry. The discovery of the 'first black Briton known to us' (Olusoga, 2016, p. 33) gained traction across several media outlets (e.g. BBC News, 2014; Campbell, 2020; Steafel, 2021; The Times, 2014; Van der Pool, 2016), non-fiction books (e.g. Adi, 2022; Olusoga, 2016, pp. 31–33; Sethi, 2021, p. 285), educational resources (e.g. Hackney Learning Trust, n.d.; Khan and Torrington, 2023; School History, n.d.; The Royal Mint, n.d.) and academic publications (e.g. Goodwin, 2020; Osborne, 2020, p. 250; Swank, 2022, p. 533; Vrikki et al., 2020, p. 87). However, over the proceeding decade as further analyses of BHW's remains were conducted (but not previously published), this interpretation began to shift.

* Corresponding author. Centre for Human Evolutionary Research, Natural History Museum, London, UK.

E-mail address: s.brace@nhm.ac.uk (S. Brace).

¹ Andy Walton and William Marsh are co-first authors and contributed equally.

Here, we present the entire suite of biomolecular and morphometric work that has been performed on the individual over that period, alongside the analysis of newly generated ancient DNA data. We detail how the interpretation of BHW ancestry has changed over the last decade as a result of recent advances in scientific methods and technology.

Part 1 Early Analysis: 2012–2024

Authors of this paper involved in this phase of analyses include Caroline Wilkinson (craniofacial), Ian Barnes and Selina Brace (aDNA).

1.1. Site and samples

Beachy Head refers to a stretch of coastline two miles east of the town of Eastbourne on the south coast of the UK (Fig. 1). The surrounding area has an extensive archaeological history, notably a well-excavated rural settlement immediately north of Beachy Head at Bullock Down, which was occupied from the conquest to the end of the Roman period (Rudling, 1982). To the west are possible rural sites at Birling and East Dean attested by occupation scatters (East Sussex HER: MES7411, MES7408, MES7407 and MES3046). To the east of Beachy Head are the palatial villa at Eastbourne (Rudling, 1982, p. 343; Russell, 2006, pp. 143–144, 264; Sutton, 1952) and the Roman fort at Pevensey, constructed in the very late 3rd century (Fulford and Rippon, 2011).

The burial evidence for the immediate region is sparse (Rudling, 1979, p. 353 Fig. 1; 1982, Figure 68; 2008, p. 128) but includes seven apparently Romano-British inhumations found in the late 19th century (Seaman, 2018; Whitley, 1890, pp. 113–115; East Sussex HER: MES3045). While grave goods were donated to Eastbourne Museum (Sussex Archaeological Society, 1892; Whitley, 1890, p. 114) they are thought to have been destroyed during WWII; there is no mention of skeletons being deposited.

1.2. Radiocarbon dating

C^{14} radiocarbon dating from a sample of BHW was performed at Oxford University's Radiocarbon Accelerator Unit (ORAU) which produced an uncalibrated date of 1831 ± 24 BP (OxA-28343). Using the IntCal20 calibration curve (Reimer *et al.*, 2020) generated a calibrated date of between 129 and 311 calCE (95 % CI; SI Fig. 1), a period that corresponds with evidence for increased Roman activity in the region.

1.3. Bioarchaeological Assessment

The osteological assessment of BHW was undertaken according to

established guidelines from the Biological Association for British Anthropologists; Institute of Archaeologists; and English Heritage (Seaman and Forsyth, 2014). This included macroscopic analysis to determine the individual's biological sex, age, stature, and any pathology or trauma visually evident. Methods and results were detailed in a report for Eastbourne Borough Council, the relevant details (those pertaining to BHW) are included in the SI (S.1). BHW's biological sex was categorised as 'possible female'. Her age range was assigned as young adult, aged between 18 and 25 at death and she was estimated to be 1.52m tall. Cause of death could not be established from the skeleton, however, there was evidence of antemortem trauma including an ossified hematoma present on the midshaft of the right femur.

In 2013, macroscopic craniofacial analysis was carried out at the University of Dundee to reconstruct BHWs appearance, (for original report see S.1.2). The cranium was almost complete, except for the loss of both maxillary and mandibular incisors, and canines. Whilst some traits such as a wide palate, mild brow ridges and a marked nuchal line were suggestive of a male, cranial features were overall indicative that the skull was female. The skull was dolichocephalic (cephalic index of 73; 131.5/180), with a post-bregmatic depression, rectangular orbits, wide interorbital distance, low nasal root, absent nasal spine, nasal guttering, moderate prognathism, wide palate, rounded mandibular border, and shallow coronoid notch. These details were found to be consistent with an individual of sub-Saharan African ancestry. Though, it was emphasised that any such determination is difficult given the wide variation in cranial morphology seen both within and between populations (Cartmill, 1998; Sauer, 1992).

This would make BHW one of the earliest individuals with recent sub-African ancestry identified thus far in Britain. Population movement from Northern Africa into Britain during the Roman period has been well established, both on an individual and military unit level. This is evidenced by inscribed altars, tombstones and military diplomas (Eckardt, 2014, pp. 67–77, Fig. 3.3). During the Roman period, North Africa was a culturally complex region characterised by centuries of mixing of Phoenician, Italian, Berber and African populations (Shaw, 2014; also Brett and Fentress, 1996; Mattingly and Hitchner, 1995, pp. 171–174). Sub-Saharan Africans were known as 'Aethiopians' and are depicted quite frequently in Roman art, with the cultural significance of these figurines, vessels and other objects still much debated (e.g. Bell, 2022, with extensive references to earlier work). Given military recruitment and trading patterns, individuals of African descent buried in Roman Britain are perhaps more likely to come from the mixed populations of North Africa than the sub-Saharan region.




Fig. 1. Location of Beachy Head and notable Roman sites surrounding the area. Map produced by Sarah Lambert-Gates.

1.4. Facial reconstruction

Alongside the 2013 craniofacial analysis, a 3D facial depiction of BHW was produced at the University of Dundee (for original report see S.1.2). This followed the Manchester method (Wilkinson, 2004; Wilkinson and Rynn, 2012) with facial feature shape determined using published anatomical and anthropometrical standards (Fedosyutkin and Nainys, 1993; Gerasimov, 1955; Rynn *et al.*, 2010; Wilkinson and Mautner, 2003). Facial features were estimated based entirely on skeletal detail using standards that were not reliant on ancestry/population affinity (Rynn *et al.*, 2012).

When performing facial reconstruction from archaeological remains, the most likely hair, skin and eye pigments are determined from the bioanthropological determination, and relevant historical and contextual data. Given the uncertainty regarding any historical phenotype and the variation in pigmentation both across and within populations, researchers chose pigments that are represented in sub-Saharan populations, but also frequently occur across other global populations. Thus, dark eye and hair colours, wavy hair type and a mid-tone skin pigment were depicted. Plastic eyeballs with brown iris colour were mounted in the head model and the skin was painted using acrylic paints. A dark brown wig with long, loose curls was attached to the model (SI figure S1.2.8).

1.5. Bulk isotope analysis

In 2013, as part of the EAP, several samples from Anglo Saxon inhumations from the Eastbourne area, including BHW, were submitted to the NERC Isotope Geosciences Laboratories (NIGL) for carbon and nitrogen analysis ($\delta^{13}\text{C}$ and $\delta^{15}\text{N}$) of collagen (dietary reconstruction), and strontium and oxygen isotope ($^{87}\text{Sr}/^{86}\text{Sr}$ and $\delta^{18}\text{O}$) analysis of tooth enamel (geographical location during childhood).

The NIGL report (full methods and results in S.2) plots the strontium and oxygen isotope results for the EAP individuals including BHW alongside data from Ringlemere Anglo Saxon individuals (Brettell *et al.*, 2012) and a previous study of Anglo Saxons from Eastbourne (Millard *et al.*, 2005) (Fig. 2a). The strontium and oxygen isotopes for all EAP individuals, including BHW, are consistent with an origin on the south coast of Britain (BHW: $^{87}\text{Sr}/^{86}\text{Sr} = 0.709100$; $\delta^{18}\text{O}_p = 17.84$; SI Fig. 2), though the results cannot exclude areas of the continent with similar values (S.2: NIGL report; SI S.2). The carbon and nitrogen isotope data from BHW ($\delta^{13}\text{C} = -18.8$) and the EAP individuals were plotted with published British individuals from the Iron Age (Wetwang Slack) (Jay and Richards, 2006), the Late Medieval but with a high fish diet (Fishergate) (Müldner and Richards, 2007) and Anglo Saxon sites (Berinsfield) (Privat *et al.*, 2002) (Fig. 2b). The EAP individuals, including BHW, had higher nitrogen isotope ratios than both the Iron Age farmers and the (inland) Anglo Saxons, suggesting a greater marine component in her diet, which would be consistent with coastal living. Relative to the other individuals, BHW had an unusual $\delta^{13}\text{C}$ bulk value (-18.8‰) suggesting a comparatively increased contribution of C4 plants in her diet.

1.6. Ancient DNA analysis

In 2017, preliminary ancient DNA (aDNA) analysis on BHW was conducted by researchers from the Natural History Museum. The low-level sequencing carried out at this time generated an insufficient amount of data based on the number of reads/nuclear DNA to analyse the DNA to publication standard. The results, however, were suggestive that she did not have recent sub-Saharan African ancestry but was more likely to be of east Mediterranean origin. Along with the isotope data, the proposal at this time was that she may have been born in Cyprus and moved to Eastbourne as a young child. Whilst these preliminary results were not published in a scientific journal they were subsequently reported in the media.




Fig. 2. The isotopic profile of Beachy Head Woman. (a) Sr and O ratios plotted to indicate mobility patterns and (b) C and N plotted against one another to indicate dietary profile. Isotopic results from individuals incorporated into the wider Eastbourne Ancestors Project (EAP) from coastal South England, alongside individuals from Roman-era Britain are plotted alongside BHW for context.

Part 2 Recent Analysis: 2024–2025

2. Methods and Analyses (2024-2025)

2.1. Ancient DNA

2.1.1. Laboratory methods (2017 and 2024)

DNA extraction and initial library building were carried out in 2017. Forty six mg of bone powder was drilled from the petrous bone of BHW using a Dremel™ high powered drill in the ancient DNA lab of the Natural History Museum; a dedicated aDNA facility following standard protocols (reviewed in Orlando *et al.*, 2021). DNA extraction method followed Dabney *et al.* (2013), modified so that the Zymo-Spin V columns were replaced with Roche High Pure Viral Nucleic Acid Large Volume spin columns, and two final elution steps of 50 μl TET buffer (total 100 μl) were performed. Double stranded library build was implemented (20 PCR cycles) using dual indexes (Meyer and Kircher, 2010). The library was screened on a NovaSeq550 2x75BP; Mid-Output at the Natural History Museum.

In 2024, a human specific DNA capture method (TWIST Biosciences) that was specifically designed for degraded human material, and targeted variable sites on the human nuclear genome (Rohland *et al.*, 2022) was employed. Laboratory methods followed the default protocol and

used a bead-based DNA isolation method to replace the standard drying down protocol, prior to the hybridization reaction. The suggested number of PCR cycles for highly degraded DNA ($n = 23$) were performed post hybridization. The captured amplified library was quantified using Tapestation 4200, High Res. and sequenced on a NovaSeq X Plus system (10B; 150x2 PE; Novogene, Cambridge).

2.1.2. Computational analysis - alignment and authentication

Paired end reads were demultiplexed and transferred onto the NHM high performance computing (hpc) cluster and the CropDiversity hpc cluster (Percival-Alwyn *et al.*, 2024). We trimmed read adapters and collapsed paired end reads (adapterremoval 2.0; Schubert *et al.*, 2016). Dual index combinations were checked, with collapsed reads showing more than two nucleotide index mismatches discarded. Collapsed reads were aligned to the nuclear human reference genome hg38 (Schneider *et al.*, 2017) and the Revised Cambridge Reference Sequence (rCRS; NC_012920) (Andrews *et al.*, 1999) using BWA 0.7.17 (Li and Durbin, 2009), relaxed parameters (-q 10) and a random seed (-l 1024). Aligned reads were subsequently sorted, filtered for mapping and base quality ($Q \geq 30$), and duplicate reads removed using samtools (Danecek *et al.*, 2021). This produced the final filtered bam file used for all subsequent computational analysis.

Mapping statistics (SI Table 1; Supplementary Data 1) for the nuclear and mitochondrial reference genomes were calculated using samtools and qualimap (Okonechnikov *et al.*, 2016), and aligned reads visually inspected using Tablet (Milne *et al.*, 2013). mapDamage 2.0 (Jónsson *et al.*, 2013) was used to assess DNA damage patterns using default settings. PMDtools (Skoglund *et al.*, 2014) was used to further authenticate DNA, with reads filtered for a PMD score >3 . Initial analyses used both filtered and unfiltered datasets to check for modern human contamination. To allow for comparison between BHW and local populations of the same time period, we downloaded recently published rural British Roman-era genomes ($n = 96$) (Scheib *et al.*, 2024) from the ENA database, and processed them in the same manner as above. The final bam file was imputed with GLIMPSE2 (Rubinacci *et al.*, 2023), using the 1000 Genome Project Phase 3 reference panel (Auton *et al.*, 2015).

2.1.3. DNA preservation and authentication

DNA was found to be well preserved in BHW, with an endogenous content of $\sim 13\%$ from initial whole genome screening (WGS) data (SI Table 1). DNA capture and sequencing resulted in over 80 % of variable 1240k SNP sites being covered by 5 or more unique reads (Table 1). DNA damage patterns seen for both WGS and capture data are in keeping with the provenance of the material (SI Fig. 3). PMD filtering of the readset shows a high proportion of retained reads (45 %; Skoglund *et al.*, 2013), and similar results of ancestry analyses when using filtered and

Table 1
Phenotypic analysis of Beachy Head Woman. Predicted eye, hair and skin pigmentation as calculated using HiIrisPlex-S.

Phenotype	Likelihood	AUC loss
Blue eye	0.911	0
Intermediate eye	0.057	0
Brown eye	0.032	0
Blond hair	0	0.025
Brown hair	0	0.015
Red hair	0	0.102
Black hair	0	0.012
Light hair	0.958	0.007
Dark hair	0.042	0.007
Very pale skin	0.067	0.064
Pale skin	0.136	0.012
Intermediate skin	0.706	0.025
Dark skin	0.05	0.005
Dark to black skin	0.041	0.003

unfiltered readsets (SI Table 1). Schmutzi (Renaud *et al.*, 2015) was run to estimate mitochondrial contamination, giving a contamination estimate of between 1 and 3 % (SI Fig. 4). As this was an acceptable range, all subsequent analyses used the unfiltered readset.

2.1.4. Sexing and MT haplogroup assignment

Sex classification was performed using Skoglund *et al.* (2013) and implemented following Anastasiadou *et al.* (2024). We considered the asymmetrical targeting of X and Y reads seen when performing targeted hybridisation capture (-capture option). As expected, we found BHW to be genetically female (XX chromosomes; SI Table 2).

A consensus mitochondrial genome for BHW was assembled using all unique q30 aligned mitochondrial reads and ANGSD (-do-fasta 2; -min. depth [2 and 3]; -trim 2). Haplogrep v3 (Schönherr *et al.*, 2023) was used to determine her mitochondrial haplogroup, using the latest MT Phylotree (v17 - Forensic Update). BHW was shown to carry a mitochondrial haplotype deriving from the K1a haplogroup (K1a26; Q = 82 %). This haplogroup is associated with modern-day populations from Northern European and the British Isles, and identified in several individuals from rural Iron Age Britain (Scheib *et al.*, 2024; Schiffels *et al.*, 2016).

2.1.5. Population genetic analyses

Prior to population level analysis, all aligned read data for BHW and all processed rural British Roman-era individuals were trimmed of terminal ends (bamUtils; -n 5). Pseudo-haploid calling was performed at nucleotide positions on the human genome shown previously to be variable across populations (Mathieson *et al.*, 2015) using samtools mpileup (Danecek *et al.*, 2021) and pileupCaller (<https://github.com/sts/chiff/sequenceTools>). These data were merged with the Allen Ancient DNA Resource (S. Mallick and Reich, 2023; v62.0), creating a dataset that contained over 20,000 modern and ancient human individuals spanning tens of thousands of years.

Initial analysis to determine the population affinity of BHW was performed using ADMIXTOOLS 2 (Maier and Patterson, 2024). Principal Component Analysis (PCA) was performed using smartpca (Patterson *et al.*, 2006; Price *et al.*, 2006), where a set of modern genomes were used to calculate eigenvectors (allsnps: YES), and BHW subsequently projected. Following the results of this initial analysis, we subset the list of modern genomes to only include those from West Eurasian and repeated the PCA to offer higher resolution. For this analysis, we projected relevant ancient Eurasian individuals alongside BHW.

Outgroup-f3 statistics ($qp3Pop$) were calculated to clarify the genetic affinity of BHW to modern populations. We used global modern genomes as comparative populations and both Chimp and Mbuti as outgroups, before analysis was refined adhoc to include all modern West Eurasians and ancient Eurasian individuals dated to the Roman period and 2000 years prior (SI Table 3). Chimp was used as an outgroup as it is asymmetrically related to all human populations, with Mbuti used for subsequent analysis focused on non-African populations given this population is equally related to all non-Africans.

In order to test the hypotheses, as suggested by previous craniofacial results, of admixture into BHW and to compare her levels of drift with contemporary British individuals we used $qpDstat$'s f4 mode in ADMIXTOOLS 2 (Maier and Patterson, 2024) and ADMIXTURE (Alexander *et al.*, 2009). F4-statistics were used to explicitly test for evidence of recent African admixture in BHW, and to infer the similarity between the ancestry of contemporary Iron Age British populations and BHW when compared to contemporary individuals from Continental Europe and Africa. See SI 3.1 for details of ADMIXTURE and structured modelling in ADMIXTURE analyses including SI Figs. 5 and 6.

Initial f3-outgroup analysis using modern human individuals ($qp3Pop\{BHW, modern\ population, Mbuti/Chimp\}$) show BHW to have a strong affinity to modern day Europeans (Fig. 3a), and the highest genetic affinity to modern individuals from Great Britain (Fig. 3b).

PCA (Fig. 3b) shows BHW is most similar to modern populations




Fig. 3. The genetic affinity of Beachy Head Woman. (a) World map with global populations coloured by outgroup-f3 statistic value when: $f3\{\text{Beachy Head Woman, modern population, Chimp}\}$. Lighter colour denotes a greater genetic affinity between BHW and the population in question. Principal component analysis (PCA), with Beachy Head Woman (golden triangle) projected onto eigenvalues calculated using (b) modern global genomes, and (c) modern Eurasian genomes. In figure (b), populations representative of sub-Saharan African (i.e. Mbuti; Red), West Eurasian (i.e. French; Green), and East Asian (i.e. Han Chinese; Blue) ancestry are highlighted. Several relevant genomes dating to the Roman period (Scheib et al., 2024) are projected onto modern West Eurasian populations in figure (c). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

from Western Europe, with no attraction towards sub-Saharan Africans. The projection of additional Roman-era individuals (Fig. 3c) highlights the association between the ancestry of BHW and populations from Roman-era Britain and Northern Europe. This ancestry differs from individuals associated with the expansion of the Roman empire (i.e. those of Mediterranean origin) and instead resembles the local and/or rural communities in Germany, Poland, Belgium and the UK (Antonio et al., 2019; Scheib et al., 2024; Silva et al., 2024) during the Roman occupation.

D-statistics (f4-statistics) consistently show no evidence of recent gene-flow between BHW and sub-Saharan Africans, with highly similar results obtained when gene flow between local and/or rural Roman-era individuals from Britain and sub-Saharan Africans are also calculated (Fig. 4 & SI Fig. 7). Ad-hoc D-statistics found *Roman-era British* individuals share elevated levels of drift with BHW (and vice-versa) over non-British Iron Age individuals (SI Fig. 8; $f4\{\text{Mbuti/Chimp, BHW, Roman-era British, European/MENA/African Roman era individuals}\}$). Results of both unsupervised ($K = 6$) (SI Fig. 9) and supervised ($K = 6$) admixture (SI Fig. 11) modelling shows BHW to have the most similar pattern of ancestry with Iron Age northern Europeans.

2.1.6. Phenotypic traits

HiIrisPlex-S (Shirk and Cushman, 2011; Walsh et al., 2014, 2017) was used to predict the hair, eye and skin pigmentation of BHW. Variable sites targeted by TWIST and shown to be associated with phenotype ($n = 32$) were filtered and individually inspected. Of the 32 sites used by HiIrisPlex-S, 27 were covered by 5 or more bases, with any nucleotide positions with less than 5 unique reads covered set as empty (N/A) to avoid issues with low-coverage data. Genotype likelihoods were calculated (ANGSD); at all sites exceeding this threshold, allowing for the homo- ($0.2 \geq \text{maf}$) and hetero- ($\text{maf} > 0.2$) zygosity at each to be determined. These calls were input into HiIrisPlex-S and eye, skin and hair pigmentation predictions assessed, with skin pigmentation (intermediate), eyes (blue), and hair (light) could be predicted (Table 1). Further, manual investigation highlighted that SNP rs1426654 was covered by 10 unique reads and homozygous (A) for the allele state strongly associated with the *SLC24A5* gene and light skin pigmentation (Lamason et al., 2005; C. B. Mallick et al., 2013). These results were confirmed with the imputed data that produced diploid calls at all targeted phenotype positions ($n = 32$) used for assessment by HiIrisPlex-S (Supplementary Data 3).




Fig. 4. A comparison of shared genetic drift between Beachy Head Woman and sub-Saharan Africans using f_4 statistics, with the shared genetic drift observed between other Roman Era British individuals and sub-Saharan Africans. BHW and contemporary British individuals modelled with African populations from the Human Origins dataset and Iron Age England as a clade. Of note is the difference between BHW and the Middle Eastern individual found in Yorkshire included in Martiniano *et al.* (2016)

2.2. Updated facial depiction

In 2025, the original facial reconstruction of BHW (produced in 2013) was scanned using an Artec Space Spider™ handheld 3D scanner (see Fig. 5a). This scan then was imported into UnReal Engine 4.26, and MetaHuman Creator (<https://MetaHuman.unrealengine.com/>) used to create a digital model of BHW that matched the facial morphology of the original reconstruction. Intermediate skin pigmentation, blond hair and blue eye colour, as informed by ancient DNA results, were selected to

create the fully pigmented and textured digital 3D depiction (Fig. 5b).

3. Discussion

The decade-long investigation into Beachy Head Woman's origins has centred around how her story has shifted over time and has sparked important debates about diversity and how we portray individuals from our past. The results presented here will no doubt add to this, but the story of the investigation into BHW is not out of the ordinary. All science




Fig. 5. Second iteration facial reconstruction of BHW. (a) 3D scan of the iteration #1 facial reconstruction - physical model (b) Iteration #2 of the facial depiction of Beachy Head Woman. MetaHuman version with hair, skin and eye textures as suggested by the DNA analysis. Both images courtesy of Face Lab at Liverpool John Moores University.

is an iterative process and BHW represents a case study in how as methods and technologies evolve, so can the conclusions we draw from them. The case of BHW does, however, raise important questions as to how methods previously commonly applied in bioanthropology classify human diversity, and how these can be combined with multiproxy biomolecular analysis.

The field of bioanthropological classification has seen significant changes over the last 15 years, and has been moving away from the traditional morphological and anthropometrical methods of ancestry estimation due to their origins in 19th century biological classification of humans along racial hierarchies (Albanese and Saunders, 2006; Brace, 1982; Haller, 1971; Sauer *et al.*, 2016; Williams *et al.*, 2005). These traditional methods of ancestry estimation using cranial metrics can help reify outdated notions of the biological reality of race (Goodman, 1997; Johnson, 2016; Ross and Williams, 2021). Classifying individuals into discrete racial categories overlooks the continuous nature of most human variation, as seen in genetic data (e.g. Prugnolle *et al.*, 2005), blood groups (e.g. Howes *et al.*, 2011), and skin pigmentation (e.g. Jablonski, 2012). In addition, skeletal traits often have as much variation within as between ancestry groups, and do not cluster into discrete categories (Manica *et al.*, 2007). There has been much discussion of how to take this into account in anthropological ancestry estimation and the visualisation of ancient peoples (Armstrong-Fumero, 2014; Caspari, 2010; Edgar and Hunley, 2009; Ousley *et al.*, 2009; Warren *et al.*, 2018; Wilkinson, 2020). Although certain morphological features are still considered to identify populations, this is being increasingly carried out with an understanding of the clinal variation in these features and the invalidation of the biological concept of race (Adams & Pilloud 2021; Hefner *et al.*, 2024; Navega *et al.*, 2015; Spradley and Weisensee, 2013).

The case of BHW demonstrates the errors associated with traditional anthropology methods and explains why they are no longer used to establish ancestry estimation. Anthropologists have noted (Dunn *et al.*, 2020; Edgar and Pilloud, 2021; Parsons, 2021) that whilst they may detect discrete biological traits believed to correlate with geographic origin, there are inherent scientific, theoretical, and structural challenges associated with these methods, due to overlapping classifications, unrepresentative datasets and inflexible interpretation. It is important to stress that anthropology methods are not unique in evolving, genomic data generation and analyses have changed considerably over the last 20 years. Next generation sequencing (NGS) radically changed the field of ancient DNA and arguably most notably the field of ancient human DNA. This is due to the large quantity and low cost of data that can be produced with NGS, but also the ability to bioinformatically authenticate genuinely ancient DNA from modern DNA, a process that was not possible using the older methods of Sanger sequencing Polymerase Chain Reaction (PCR) products. Whilst the genomic results reported here were all generated from authenticated NGS data, the shift in ancestry from that of a possible Cypriot background when using low level sequencing data to English/northern continental Europe when deeper sequencing effort was achieved further highlights the importance of generating sufficient high resolution genomic data to more accurately determine an individual's ancestry. The results of this paper suggest that previous archaeological cases reliant on anthropological ancestry estimation, and/or non-authenticated genomic data from older studies, may need reassessment using multi-proxy biomolecular methods. For Roman archaeology, specifically, the results offer an opportunity to reassess bioarchaeological work carried out over recent decades, while still acknowledging the strong historical and epigraphic evidence for migration into Britain during this period.

4. Conclusions

The research presented here spans over a decade of technological advancements and highlights progressive analyses in osteological, morphometric and biomolecular techniques that have been applied to BHW. Although initial analysis using traditional anthropological

morphometric techniques deemed the remains of BHW to be likely of sub-Saharan origin, the results from recent genetic analysis indicate that she shows a close affinity to individuals from modern day England and contemporary Roman-era Iron Age individuals in England and Northern continental Europe. When combined with $^{87}\text{Sr}/^{86}\text{Sr}$ isotope results, which indicated BHW shared similar early-life mobility to individuals also excavated from the south coast and dated to the Roman-era Iron Age and early medieval periods, this suggests BHW likely came from the south coast of England. The decade-long investigation into BHW not only helps clarify her origin but also illustrates this iterative work in practice. By reporting each analysis conducted on BHW, even those whose conclusions we no longer support, we highlight how these efforts can lead to both more equitable and more accurate science.

CRediT authorship contribution statement

Andy Walton: Writing – review & editing, Writing – original draft, Methodology, Investigation, Formal analysis. **William Marsh:** Writing – review & editing, Methodology, Investigation, Formal analysis. **Alex Strang:** Writing – review & editing, Methodology, Formal analysis. **Jonathan Seaman:** Writing – review & editing, Resources, Conceptualization. **Kelly Van Doorn:** Writing – review & editing, Resources. **Hella Eckardt:** Writing – review & editing. **Caroline Wilkinson:** Writing – review & editing, Writing – original draft, Resources, Methodology, Formal analysis. **Ian Barnes:** Writing – review & editing, Supervision, Conceptualization. **Selina Brace:** Writing – review & editing, Writing – original draft, Supervision, Investigation, Conceptualization.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors acknowledge Research Computing at the James Hutton Institute for providing computational resources and technical support for the “UK’s Crop Diversity Bioinformatics HPC” (BBSRC grants BB/S019669/1 and BB/X019683/1), use of which has contributed to the results reported within this paper. We thank Heritage Eastbourne, of Eastbourne Council, and their research staff for their help and cooperation throughout this project. We would also like to thank The Calleva Foundation for supporting several authors of this paper. AW is grateful to the Adelphi Genetics Forum for their funding support.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jas.2025.106445>.

Data availability

DNA sequence data generated for this project has been deposited in the European Nucleotide Archive under project accession number PRJEB102740.

References

Adams, D.M., Pilloud, M.A., 2021. The (mis) appropriation of biological anthropology in race science and the implications for forensic anthropology. *Forensic Anthropology* 4 (4), 97.

Adi, H., 2022. *African and Caribbean People in Britain: a History*. Penguin Books.

Albanese, J., Saunders, S.R., 2006. Is it possible to escape racial typology in forensic identification? In: Schmitt, A., Cunha, E., Pinheiro, J. (Eds.), *Forensic Anthropology and Medicine: Complementary Sciences from Recovery to Cause of Death*. Humana Press, pp. 281–316. https://doi.org/10.1007/978-1-59745-099-7_12.

Alexander, D.H., Novembre, J., Lange, K., 2009. Fast model-based estimation of ancestry in unrelated individuals. *Genome Res.* 19 (9), 1655–1664.

Anastasiadou, K., Silva, M., Booth, T., Speidel, L., Audsley, T., Barrington, C., Buckberry, J., Fernandes, D., Ford, B., Gibson, M., Gilardet, A., Glocke, I., Keefe, K., Kelly, M., Masters, M., McCabe, J., McIntyre, L., Ponce, P., Rowland, S., et al., 2024. Detection of chromosomal aneuploidy in ancient genomes. *Commun. Biol.* 7 (1), 14. <https://doi.org/10.1038/s42003-023-05642-z>.

Andrews, R.M., Kubacka, I., Chinnery, P.F., Lightowler, R.N., Turnbull, D.M., Howell, N., 1999. Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nat. Genet.* 23 (2). <https://doi.org/10.1038/13779>, 147–147.

Antonio, M.L., Gao, Z., Moots, H.M., Lucci, M., Candilio, F., Sawyer, S., Oberreiterer, V., Calderon, D., Devitofranceschi, K., Aikens, R.C., Aneli, S., Bartoli, F., Bedini, A., Cheromet, O., Cotter, D.J., Fernandes, D.M., Gasperetti, G., Griffoni, R., Guidi, A., et al., 2019. Ancient Rome: a genetic crossroads of Europe and the Mediterranean. *Science* 366 (6466), 708–714. <https://doi.org/10.1126/science.aay6826>.

Armstrong-Fumero, F., 2014. 'Even the Most Careless Observer': race and visual discernment in physical anthropology from Samuel Morton to Kennewick man. *Am. Stud.* 53 (2), 5–29. JSTOR.

Auton, A., Abecasis, G.R., Alshuler, D.M., Durbin, R.M., Abecasis, G.R., Bentley, D.R., Chakravarti, A., Clark, A.G., Donnelly, P., Eichler, E.E., Flieck, P., Gabriel, S.B., Gibbs, R.A., Green, E.D., Hurles, M.E., Knoppers, B.M., Korbel, J.O., Lander, E.S., Lee, C., et al., 2015. A global reference for human genetic variation. *Nature* 526 (7571), 68–74. <https://doi.org/10.1038/nature15393>.

BBC News, 2014. Centuries old Beachy Head Lady's face revealed. BBC News. <https://www.bbc.com/news/uk-england-sussex-25962183>.

Bell, S.W., 2022. Images and interpretation of Africans in Roman art and social practice. In: Cline, L.K., Elkins, N.T. (Eds.), *The Oxford Handbook of Roman Imagery and Iconography*. Oxford University Press, pp. 425–463. <https://doi.org/10.1093/oxfordhb/9780190850326.013.25>.

Brace, C.L., 1982. The roots of the race concept in American physical anthropology. *A History of American Physical Anthropology, 1930–1980* 11–29.

Brett, Michael, Fentress, Elizabeth, 1996. *The Berbers*. Blackwell.

Brettell, R., Evans, J., Marzinik, S., Lamb, A., Montgomery, J., 2012. 'Impious easterners': can oxygen and strontium isotopes serve as indicators of provenance in early medieval European cemetery populations? *Eur. J. Archaeol.* 15 (1), 117–145. <https://doi.org/10.1179/146195712Y.0000000001>. Cambridge Core.

Campbell, L., 2020. Ten black history events that should be taught to every pupil. *Guardian* 17. ProQuest Central.

Cartmill, M., 1998. The status of the race concept in physical anthropology. *Am. Anthropol.* 100 (3), 651–660. <https://doi.org/10.1525/aa.1998.100.3.651>.

Caspari, R., 2010. Deconstructing race: racial thinking, geographic variation, and implications for biological anthropology. In: *A Companion to Biological Anthropology*, pp. 104–123. <https://doi.org/10.1002/9781444320039.ch6>.

Dabney, J., Knapp, M., Glocke, I., Gansauge, M.-T., Weihmann, A., Nickel, B., Valdiosera, C., García, N., Pääbo, S., Arsuaga, J.-L., Meyer, M., 2013. Complete mitochondrial genome sequence of Middle Pleistocene cave bear reconstructed from ultrashort DNA fragments. *Proc. Natl. Acad. Sci.* 110 (39), 15758–15763. <https://doi.org/10.1073/pnas.1314445110>.

Danecek, P., Bonfield, J.K., Liddle, J., Marshall, J., Ohan, V., Pollard, M.O., Whitwham, A., Keane, T., McCarthy, S.A., Davies, R.M., Li, H., 2021. Twelve years of SAMtools and BCFtools. *GigaScience* 10 (2), giab008. <https://doi.org/10.1093/gigascience/giab008>.

Dunn, R.R., Spiros, M.C., Kamnikar, K.R., Plemmons, A.M., Hefner, J.T., 2020. Ancestry estimation in forensic anthropology: a review. *Wiley Interdiscip. Rev.: Forensic Sci.* 2 (4), e1369. <https://doi.org/10.1002/wfs2.1369>.

Eckardt, H., 2014. Objects and Identities: Roman Britain and the North-Western Provinces. Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199693986.001.0001>.

Edgar, H.J.H., Hunley, K.L., 2009. Race reconciled?: how biological anthropologists view human variation. *Am. J. Phys. Anthropol.* 139 (1), 1–4. <https://doi.org/10.1002/ajpa.20995>.

Edgar, H., Pilloud, M., 2021. A reassessment of assessing Race: "Ancestry" estimation and its implications for forensic anthropology and beyond. *Forensic Anthropology* (University of Florida) 4 (4).

Fedorystukin, B.A., Nainys, J.V., 1993. The relationship of skull morphology to facial features. *Forensic Analysis of the Skull* 199–213.

Fulford, Michael, Rippon, S., 2011. Pevensey Castle, Sussex: excavations in the Roman fort and medieval keep, 1993–95. *Wessex Archaeology*. No. 26. 1–185. ISBN: 978-1-874350-55-2.

Gerasimov, M., 1955. The reconstruction of the face from the basic structure of the skull. *Trans. W. Tshernezky (1975) Russia*.

Goodman, A.H., 1997. Bred in the bone? *The Sciences* 37 (2), 20–25. <https://doi.org/10.1002/j.2326-1951.1997.tb03296.x>.

Goodwin, K.A., 2020. Modern Ethnicity and Ancient Culture: The Persistence of White Narratives in Displays of the Roman Past [Ph.D., University of Kent (United Kingdom)]. In: *PQDT - Global* (3073246540). ProQuest Dissertations & Theses Global. <https://doi.org/10.22244/UniKent/01.02.87602>.

Hackney Learning Trust. (n.d.). Home learning: history activity KS3. https://www.hackneyservicesforschools.co.uk/system/files?file=extranet/TheBeachyHeadLady_KS3_blackhistory%20task.pdf.

Haller Jr., J.S., 1971. Attitudes of racial inferiority in nineteenth-century anthropometry. In: *Outcasts from Evolution: Scientific Attitudes of Racial Inferiority*. University of Illinois Press, pp. 1859–1900.

Hefner, J.T., Ousley, S., Richardson, R., 2024. MaMD Analytical 1.0: a Computer Program for macromorphoscopic trait analysis. *Forensic Anthropology* 7 (2–3), 119.

Howes, R.E., Patil, A.P., Piel, F.B., Nyangiri, O.A., Kabaria, C.W., Gething, P.W., Zimmerman, P.A., Barnadas, C., Beall, C.M., Gebremedhin, A., Ménard, D., Williams, T.N., Weatherall, D.J., Hay, S.I., 2011. The global distribution of the Duffy blood group. *Nat. Commun.* 2 (1), 266. <https://doi.org/10.1038/ncomms1265>.

Jablonski, N.G., 2012. Living Color, first ed. University of California Press; JSTOR <http://www.jstor.org/stable/10.1525/j.ctt1pn64b>.

Jay, M., Richards, M.P., 2006. Diet in the Iron Age cemetery population at Wetwang Slack, East Yorkshire, UK: carbon and nitrogen stable isotope evidence. *J. Archaeol. Sci.* 33 (5), 653–662. <https://doi.org/10.1016/j.jas.2005.09.020>.

Johnson, H., 2016. Craniofacial reconstruction and its socio-ethical implications. *Mus. Soc. Issues* 11 (2), 97–113. <https://doi.org/10.1080/15596893.2016.1204598>.

Jónsson, H., Ginolhac, A., Schubert, M., Johnson, P.L.F., Orlando, L., 2013. mapDamage2.0: fast approximate Bayesian estimates of ancient DNA damage parameters. *Bioinformatics* 29 (13), 1682–1684. <https://doi.org/10.1093/bioinformatics/btt193>.

Khan, I., Torrington, A., 2023. Teachers Pack 2023. Commissioned by Lambeth Council. https://content.govdelivery.com/attachments/UKLAMBETH/2024/06/18/file_attachments/2910735/Windrush%202023_Teacher%27s%20Pack_June%202023.pdf.

Lamason, R.L., Mohideen, M.-A.P.K., Mest, J.R., Wong, A.C., Norton, H.L., Aros, M.C., Jurneyc, M.J., Mao, X., Humphreville, V.R., Humbert, J.E., Sinha, S., Moore, J.L., Jagadeeswaran, P., Zhao, W., Ning, G., Makalowska, I., McKeigue, P.M., O'Donnell, D., Kittles, R., et al., 2005. SLC24A5, a putative cation exchanger, affects pigmentation in zebrafish and humans. *Science* 310 (5755), 1782–1786. <https://doi.org/10.1126/science.1116238>.

Li, H., Durbin, R., 2009. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25 (14), 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>.

Maier, R., Patterson, N., 2024. Admixtools: inferring demographic history from genetic data. <https://github.com/uqrmaie1/admixtools>.

Mallick, C.B., Iliescu, F.M., Möls, M., Hill, S., Tamang, R., Chaubey, G., Goto, R., Ho, S.Y., W., Gallego Romero, I., Crivellaro, F., Hudjashov, G., Rai, N., Metspalu, M., Mascie-Taylor, C.G.N., Pitchappan, R., Singh, L., Mirazon-Lahr, M., Thangaraj, K., Villem, R., Kivisild, T., 2013. The light skin allele of SLC24A5 in South Asians and Europeans shares identity by descent. *PLoS Genet.* 9 (11), e1003912. <https://doi.org/10.1371/journal.pgen.1003912>.

Mallick, S., Reich, D., 2023. *The Allen Ancient DNA Resource (AADR): a curated compendium of ancient human genomes* (Version V8). Harvard DataVerse. <https://doi.org/10.7910/DVN/FFIDCW> [Data set].

Manica, A., Amos, W., Balloux, F., Hanilhara, T., 2007. The effect of ancient population bottlenecks on human phenotypic variation. *Nature* 448 (7151), 346–348. <https://doi.org/10.1038/nature05951>.

Martiniano, R., Caffell, A., Holst, M., Hunter-Mann, K., Montgomery, J., Müldner, G., McLaughlin, R.L., Teasdale, M.D., van Rheezen, W., Veldink, J.H., van den Berg, L.H., Hardiman, O., Carroll, M., Roskams, S., Oxley, J., Morgan, C., Thomas, M.G., Barnes, I., McDonnell, C., et al., 2016. Genomic signals of migration and continuity in Britain before the Anglo-Saxons. *Nat. Commun.* 7 (1), 10326. <https://doi.org/10.1038/ncomms10326>.

Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg, S.A., Harney, E., Stewardson, K., Fernandes, D., Novak, M., Sirak, K., Gamba, C., Jones, E.R., Llamas, B., Dryomov, S., Pickrell, J., Arsuaga, J.L., de Castro, J.M.B., Carbonell, E., et al., 2015. Genome-wide patterns of selection in 230 ancient Eurasians. *Nature* 528 (7583), 499–503. <https://doi.org/10.1038/nature16152>.

Mattingly, D.J., Hitchner, R.B., 1995. Roman Africa: an archaeological review. In: *Journal of Roman Studies*, vol. 85. Cambridge Core, pp. 165–213. <https://doi.org/10.2307/301062>.

Meyer, M., Kircher, M., 2010. Illumina sequencing library preparation for highly multiplexed target capture and sequencing. *Cold Spring Harb. Protoc.* 2010 (6), prot5448. <https://doi.org/10.1101/pdb.prot5448>.

Millard, A.R., Roberts, C.A., Hughes, S.S., 2005. Isotopic evidence for migration in medieval England: the potential for tracking the introduction of disease. *Soc. Biol. Hum. Aff.* 70 (1).

Milne, I., Stephen, G., Bayer, M., Cock, P.J.A., Pritchard, L., Cardle, L., Shaw, P.D., Marshall, D., 2013. Using Tablet for visual exploration of second-generation sequencing data. *Briefings Bioinf.* 14 (2), 193–202. <https://doi.org/10.1093/bib/bbs012>.

Müldner, G., Richards, M.P., 2007. Diet and diversity at later medieval fishergate: the isotopic evidence. *Am. J. Phys. Anthropol.* 134 (2), 162–174. <https://doi.org/10.1002/ajpa.20647>.

Navega, D., Coelho, C., Vicente, R., Ferreira, M.T., Wasterlain, S., Cunha, E., 2015. AncestryTrees: ancestry estimation with randomized decision trees. *Int. J. Leg. Med.* 129 (5), 1145–1153.

Okonechnikov, K., Conesa, A., García-Alcalde, F., 2016. Qualimap 2: advanced multi-sample quality control for high-throughput sequencing data. *Bioinformatics* 32 (2), 292–294. <https://doi.org/10.1093/bioinformatics/btv566>.

Olusoga, D., 2016. *Black and British: a Forgotten History*. Macmillan.

Orlando, L., Allaby, R., Skoglund, P., Der Sarkissian, C., Stockhammer, P.W., Ávila-Arcos, M.C., Fu, Q., Krause, J., Willerslev, E., Stone, A.C., Warinner, C., 2021. Ancient DNA analysis. *Nat. Rev. Methods Primers* 1 (1), 14. <https://doi.org/10.1038/s43586-020-00011-0>.

Osborne, D., 2020. "Civis Romana Sum" Bernardine Evaristo's the emperor's babe and the emancipatory poetics of (Multi-) cultural citizenship. In: Wyatt, J., George, S. (Eds.), *Reading Contemporary Black British and African American Women Writers*, first ed. Routledge. <https://doi.org/10.4324/9780429199271-16>.

Ousley, S., Jantz, R., Freid, D., 2009. Understanding race and human variation: why forensic anthropologists are good at identifying race. *Am. J. Phys. Anthropol.* 139 (1), 68–76. <https://doi.org/10.1002/ajpa.21006>.

Parsons, H.R., 2021. Ancestry estimation in practice: an evaluation of forensic anthropology reports in the United States. *Forensic Anthropology* 4 (4), 192.

Patterson, N., Price, A.L., Reich, D., 2006. Population structure and eigenanalysis. *PLoS Genet.* 2 (12), e190. <https://doi.org/10.1371/journal.pgen.0020190>.

Percival-Alwyn, L., Barnes, I., Clark, M.D., Cockram, J., Coffey, M.P., Jones, S., Kersey, P. J., Kidner, C.A., Kosiol, C., Li, B., Marsh, W.A., Zhou, J., Caccamo, M., Milne, I., 2024. UKCropDiversity-HPC: a collaborative high-performance computing resource approach for sustainable agriculture and biodiversity conservation. *PLANTS, PEOPLE, PLANET.* <https://doi.org/10.1002/ppp3.10607>. n/a(n/a).

Price, A.L., Patterson, N.J., Plenge, R.M., Weinblatt, M.E., Shadick, N.A., Reich, D., 2006. Principal components analysis corrects for stratification in genome-wide association studies. *Nat. Genet.* 38 (8), 904–909. <https://doi.org/10.1038/ng1847>.

Privat, K.L., O'connell, T.C., Richards, M.P., 2002. Stable isotope analysis of human and faunal remains from the Anglo-Saxon cemetery at Berinsfield, Oxfordshire: dietary and social implications. *J. Archaeol. Sci.* 29 (7), 779–790. <https://doi.org/10.1006/jasc.2001.0785>.

Prugnolle, F., Manica, A., Balloux, F., 2005. Geography predicts neutral genetic diversity of human populations. *Curr. Biol.* 15 (5), R159–R160. <https://doi.org/10.1016/j.cub.2005.02.038>.

Reimer, P.J., Austin, W.E.N., Bard, E., Bayliss, A., Blackwell, P.G., Bronk Ramsey, C., Butzin, M., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M., Guilderson, T.P., Hajdas, I., Heaton, T.J., Hogg, A.G., Hughen, K.A., Kromer, B., Manning, S.W., Muscheler, R., et al., 2020. The IntCal20 Northern hemisphere radiocarbon Age calibration curve (0–55 cal kBP). *Radiocarbon* 62 (4), 725–757. <https://doi.org/10.1017/RDC.2020.41>. Cambridge Core.

Renaud, G., Slon, V., Duggan, A.T., Kelso, J., 2015. Schmutzi: estimation of contamination and endogenous mitochondrial consensus calling for ancient DNA. *Genome Biol.* 16 (1), 224. <https://doi.org/10.1186/s13059-015-0776-0>.

Rohland, N., Mallick, S., Mah, M., Maier, R., Patterson, N., Reich, D., 2022. Three assays for in-solution enrichment of ancient human DNA at more than a million SNPs. *Genome Res.* 32 (11–12), 2068–2078. <https://doi.org/10.1101/gr.276728.122>.

Ross, A.H., Williams, S.E., 2021. Ancestry studies in forensic anthropology: back on the frontier of racism. *Biology* 10 (7). <https://doi.org/10.3390/biology10070602>.

Rubinacci, S., Hofmeister, R.J., Sousa da Mota, B., Delaneau, O., 2023. Imputation of low-coverage sequencing data from 150,119 UK Biobank genomes. *Nat. Genet.* 55 (7), 1088–1090. <https://doi.org/10.1038/s41588-023-01438-3>.

Rudling, D., 1979. Invasion and response: downland settlement in East Sussex. In: Burnham, B.C., Johnson, H.B. (Eds.), *Invasion and Response: the Case of Roman Britain*. BAR, pp. 339–356.

Rudling, D., 1982. The Romano-British farm on Bullock Down. In: Drewett, P. (Ed.), *The Archaeology of Bullock Down, Eastbourne, East Sussex: the Development of a Landscape*. Sussex Archaeological Society, pp. 97–142.

Russell, Miles, 2006. *Roman Sussex*. Tempus.

Rynn, C., Balueva, T., Veselovskaya, E., 2012. Relationships between the skull and face. In: Wilkinson, C., Rynn, C. (Eds.), *Craniofacial Identification*. Cambridge University Press, Cambridge Core, pp. 193–202. <https://doi.org/10.1017/CBO9781139049566.016>.

Rynn, C., Wilkinson, C.M., Peters, H.L., 2010. Prediction of nasal morphology from the skull. *Forensic Sci. Med. Pathol.* 6, 20–34.

Sauer, N.J., 1992. Forensic anthropology and the concept of race: if races don't exist, why are forensic anthropologists so good at identifying them? *Soc. Sci. Med.* 34 (2), 107–111. [https://doi.org/10.1016/0277-9536\(92\)90086-6](https://doi.org/10.1016/0277-9536(92)90086-6).

Sauer, N.J., Wankmiller, J.C., Hefner, J.T., 2016. The assessment of ancestry and the concept of race. <https://doi.org/10.4324/9781315528939-21>.

Scheib, C.L., Hui, R., Rose, A.K., D'Atanasio, E., Inskip, S.A., Dittmar, J., Cessford, C., Griffith, S.J., Solnik, A., Wiseman, R., Neil, B., Biers, T., Harknett, S.-J., Sasso, S., Biagini, S.A., Runfeldt, G., Duhig, C., Evans, C., Metspalu, M., et al., 2024. Low genetic impact of the roman occupation of Britain in rural communities. *Mol. Biol. Evol.* 41 (9), msae168. <https://doi.org/10.1093/molbev/msae168>.

Schiffels, S., Haak, W., Paajanen, P., Llamas, B., Popescu, E., Loe, L., Clarke, R., Lyons, A., Mortimer, R., Sayer, D., Tyler-Smith, C., Cooper, A., Durbin, R., 2016. Iron Age and Anglo-Saxon genomes from East England reveal British migration history. *Nat. Commun.* 7 (1), 10408. <https://doi.org/10.1038/ncomms10408>.

Schneider, V.A., Graves-Lindsay, T., Howe, K., Bouk, N., Chen, H.-C., Kitts, P.A., Murphy, T.D., Pruitt, K.D., Thibaud-Nissen, F., Albracht, D., Fulton, R.S., Kremitzki, M., Magrini, V., Markovic, C., McGrath, S., Steinberg, K.M., Auger, K., Chow, W., Collins, J., et al., 2017. Evaluation of GRCh38 and de novo haploid genome assemblies demonstrates the enduring quality of the reference assembly. *Genome Res.* 27 (5), 849–864.

Schönherr, S., Weissensteiner, H., Kronenberg, F., Forer, L., 2023. Haplogrep 3—An interactive haplogroup classification and analysis platform. *Nucleic Acids Res.* 51 (W1), W263–W268. <https://doi.org/10.1093/nar/gkad284>.

School History. Beachy head lady Facts & Worksheets. n.d. <https://schoolhistory.co.uk/ancient-world/beachy-head-lady/>. (Accessed 6 April 2025).

Schubert, M., Lindgreen, S., Orlando, L., 2016. AdapterRemoval v2: rapid adapter trimming, identification, and read merging. *BMC Res. Notes* 9 (1), 88. <https://doi.org/10.1186/s13104-016-1900-2>.

Seaman, J., 2018. The mystery of Beachy Head Lady. *Museum Crush*. <https://museumcrush.org/the-mystery-of-beachy-head-lady-a-roman-african-from-eastbourne>.

Seaman, J., Forsyth, F., 2014. Eastbourne Ancestors Project Report. Eastbourne Council. <https://www.visiteastbourne.com/dbimngs/EA%20Report%20HF%20JS%20edited%20K.pdf>.

Sethi, A., 2021. *I Belong Here: a Journey Along the Backbone of Britain*. Bloomsbury Publishing Plc. <http://ebookcentral.proquest.com/lib/ucl/detail.action?docID=6561627>.

Shaw, B.D., 2014. Who are you? Africa and Africans. In: McInerney, J. (Ed.), *A Companion to Ethnicity in the Ancient Mediterranean*. Wiley-Blackwell, pp. 527–540.

Shirk, A.J., Cushman, S.A., 2011. sGD: software for estimating spatially explicit indices of genetic diversity. *Molecular Ecology Resources* 11 (5), 922–934. <https://doi.org/10.1111/j.1755-0998.2011.03035.x>.

Silva, M., Booth, T., Moore, J., Anastasiadou, K., Walker, D., Gilardet, A., Barrington, C., Kelly, M., Williams, M., Henderson, M., 2024. An individual with Sarmatian-related ancestry in Roman Britain. *Curr. Biol.* 34 (1), 204–212.

Skoglund, P., Northoff, B.H., Shunkov, M.V., Derevianko, A.P., Pääbo, S., Krause, J., Jakobsson, M., 2014. Separating endogenous ancient DNA from modern day contamination in a Siberian Neandertal. *Proc. Natl. Acad. Sci.* 111 (6), 2229–2234. <https://doi.org/10.1073/pnas.1318934111>.

Skoglund, P., Storå, J., Götherström, A., Jakobsson, M., 2013. Accurate sex identification of ancient human remains using DNA shotgun sequencing. *J. Archaeol. Sci.* 40 (12), 4477–4482. <https://doi.org/10.1016/j.jas.2013.07.004>.

Spradley, M.K., Weisensee, K.A., 2013. Why do forensic anthropologists estimate ancestry, and why is it so controversial. *Forensic anthropology: An introduction* 231–243.

Steafel, E., 2021. The forgotten stories of the Black tudors. In: *The Daily Telegraph*, vol 51644. The Telegraph Historical Archive, p. 23.

Sussex Archaeological Society, 1892. *Sussex Archaeological Collections Relating to the History and Antiquities of the County*, vol. XXXVIII. Farncombe & Co.

Sutton, T., 1952. The Eastbourne Roman Villa, vol. 90, pp. 1–12. <https://doi.org/10.5284/1085426>.

Swank, K., 2022. "Moor" and "Saracen" in medieval and contemporary arthurian texts. In: Coldham-Fussel, V., Edlich-Muth, M., Ward, R. (Eds.), *The Arthurian World*. Routledge.

The Royal Mint. Diversity built Britain Black British history teachers' notes. n.d. <https://www.royalmint.com/globalassets/kids/activity-packs/diversity-built-britain—black-british-history.pdf>.

The Times, 2014. Mystery of African Woman who Died in Roman Britain. *the Times*, 7111, 14. The Times Digital Archive.

Van der Pool, J. (Director), 2016. Black and British: a Forgotten history: a Forgotten history, Episode 1 [video recording]. In: *Academic Video Online* (2158276380). BBC Worldwide; Academic Video Online; ProQuest One Black Studies. <https://www.proquest.com/audio-video-works/first-encounters-africa-britain-forgotten-history/doc/view/2158276380/se-2?accountid=14511>.

Vrikki, P., Malik, S., Jagannathan, A., 2020. Creative anti-racisms. In: Malik, S., Mahn, C., Pierse, M., Rogaly, B. (Eds.), *Creativity and Resistance in a Hostile World*. Manchester University Press; JSTOR, pp. 81–112. <http://www.jstor.org/stable/j.ctv1c3pd6v.10>.

Walsh, S., Chaitanya, L., Breslin, K., Muralidharan, C., Bronikowska, A., Pospiech, E., Koller, J., Kovatsi, L., Wollstein, A., Branicki, W., Liu, F., Kayser, M., 2017. Global skin colour prediction from DNA. *Hum. Genet.* 136 (7), 847–863. <https://doi.org/10.1007/s00439-017-1808-5>.

Walsh, S., Chaitanya, L., Clarisse, L., Wirken, L., Draus-Barini, J., Kovatsi, L., Maeda, H., Ishikawa, T., Sijen, T., de Knijff, P., Branicki, W., Liu, F., Kayser, M., 2014. Developmental validation of the HiRisPlex system: DNA-based eye and hair colour prediction for forensic and anthropological usage. *Forensic Sci. Int.: Genetics* 9, 150–161. <https://doi.org/10.1016/j.fsigene.2013.12.006>.

Warren, M.W., Friend, A.N., Stock, M.K., 2018. Navigating cognitive bias in forensic anthropology. In: Boyd, Clifford C.J., Boyd, D.C. (Eds.), *Forensic Anthropology: Theoretical Framework and Scientific Basis*. Wiley Press, pp. 39–52. <https://doi.org/10.1002/9781119226529.ch3>.

Whitley, H.M., 1890. Recent archaeological discoveries in the eastbourne district. *Sussex Archaeological Collections* 37, 111–115.

Wilkinson, C., 2004. *Forensic Facial Reconstruction*. Cambridge University Press.

Wilkinson, C., 2020. Cognitive bias and facial depiction from skeletal remains. *Bioarchaeology International* 4 (1), 1–14. <https://doi.org/10.5744/bi.2020.1001>. ProQuest Central; Social Science Premium Collection.

Wilkinson, C., Mautner, S.A., 2003. Measurement of eyeball protrusion and its application in facial reconstruction. *J. Forensic Sci.* 48 (1), JFS2002053.

Wilkinson, C., Rynn, C. (Eds.), 2012. *Craniofacial Identification*. Cambridge University Press. <https://doi.org/10.1017/cbo9781139049566>.

Williams, F.L., Belcher, R.L., Armelagos, G.J., 2005. Forensic misclassification of Ancient Nubian crania: implications for assumptions about human variation. *Curr. Anthropol.* 46 (2), 340–346. <https://doi.org/10.1086/428792>.