

Reduced future North Atlantic eddy-driven jet variability in high-resolution, fully coupled global climate models

Article

Published Version

Creative Commons: Attribution 4.0 (CC-BY)

Open Access

Baker, A. J. ORCID: <https://orcid.org/0000-0003-2697-1350>, Lockwood, J. F., Athanasiadis, P. J. and Vidale, P. L. ORCID: <https://orcid.org/0000-0002-1800-8460> (2026) Reduced future North Atlantic eddy-driven jet variability in high-resolution, fully coupled global climate models. *Journal of Climate*, 39 (11). pp. 2925-2936. ISSN 1520-0442 doi: 10.1175/jcli-d-25-0418.1 Available at <https://centaur.reading.ac.uk/129189/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1175/jcli-d-25-0418.1>

Publisher: American Meteorological Society

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

Reduced Future North Atlantic Eddy-Driven Jet Variability in High-Resolution, Fully Coupled Global Climate Models

ALEXANDER J. BAKER^a, JULIA F. LOCKWOOD^b, PANOS J. ATHANASIADIS^c, AND PIER LUIGI VIDALE^a

^a National Centre for Atmospheric Science and Department of Meteorology, University of Reading, Reading, Berkshire, United Kingdom

^b Met Office Hadley Centre, Exeter, Devon, United Kingdom

^c Centro Euro-Mediterraneo sui Cambiamenti Climatici, Bologna, Italy

(Manuscript received 24 July 2025, in final form 9 February 2026, accepted 24 March 2026)

ABSTRACT: The westerly jet streams are a key component of North Atlantic climate, particularly during winter. Projections of European surface climate change depend largely on models' ability to capture jet behavior, which requires sufficient resolution in both the atmosphere and ocean. We evaluated the impact of model resolution on the winter climatological zonal wind and on eddy-driven jet position and speed simulated under historical (1950–2014) and future (2015–50) climate conditions in an ensemble of fully coupled global climate models, with resolution spanning from ~100 to ~25 km in the atmosphere and from 1° to 1/12° in the ocean. We find that increasing resolution improves the North Atlantic climatological zonal wind field at mid-high latitudes, but biases remain around the low-latitude, equatorward flank of the upper-level subtropical jet. By 2050, low-resolution models simulate a small equatorward shift in the midlatitude jet, reducing the meridional separation of the subtropical and midlatitude jets, but high-resolution models project a strengthening of the jets and a small poleward shift of the midlatitude jet. The analysis of the large-scale meridional temperature gradient over the North Atlantic suggests that tropical amplification influences the future zonal wind response, and there is some sensitivity of lower-level temperature gradient trends to ocean resolution. At low resolution, the eddy-driven jet shows little mean meridional shift, but increasing resolution reduces the jet's latitudinal variance. These results help clarify the role of model resolution in near-term North Atlantic climate projections and suggest further increases in atmosphere and ocean resolutions may advance the understanding of future jet behavior and its downstream impacts on surface climate over Europe.

SIGNIFICANCE STATEMENT: The westerly jet streams are an important part of large-scale atmospheric circulation. Over the North Atlantic, the eddy-driven component of this westerly flow steers midlatitude storm systems toward Europe. Projections of European surface climate, therefore, hinge on future jet behavior. However, projected North Atlantic circulation changes are uncertain. Increased model resolution in both the atmosphere and ocean may help reduce uncertainty. We find that projections of jet variability in latitude differ between low- and high-resolution models, and we also detect an influence of ocean resolution on trends in the large-scale meridional temperature gradients that govern the midlatitude atmospheric circulation response. These results highlight the need for high-resolution, coupled models in exploring future North Atlantic climate.

KEYWORDS Atmospheric circulation; Jets; General circulation models; Climate change

1. Introduction

The westerly jet streams are a key part of the large-scale atmospheric circulation over the North Atlantic. In the midlatitudes, the lower-level, eddy-driven component of this zonal flow steers synoptic, low pressure weather systems toward Europe (Hoskins and Hodges 2002; Woollings et al. 2018b). On the seasonal time scale, model skill in predicting jet position and the North Atlantic Oscillation are strongly related (Parker et al.

2019), highlighting the importance of the jet in modulating downstream weather. Projections of high-impact weather, therefore, depend largely on future jet behavior (Harvey et al. 2023), particularly in winter, when the eddy-driven jet is most active. However, despite the importance of the jet for European surface climate, projected North Atlantic circulation changes, based on coupled climate models, remain uncertain (Blackport and Fyfe 2022; Woollings 2010). Key sources of uncertainty are an insufficient physical understanding (Shepherd 2014), significant intermodel spread (Oudar et al. 2020), and the divergence of historical model simulations from observational data (Blackport and Fyfe 2022; Smith et al. 2020).

Climate models project a strengthening and narrowing of the eddy-driven jet in response to amplified near-surface warming in the Arctic and upper-level warming in the tropics (Oudar et al. 2020; Peings et al. 2018)—two opposing influences on the jet—as well as a downstream extension in winter (Harvey et al. 2023). Based on data from the Coupled Model Intercomparison Project (CMIP) phase 3–6, studies have described the competing effects of Arctic and tropical amplification, which

Denotes content that is immediately available upon publication as open access.

Supplemental information related to this paper is available at the Journals Online website: <https://doi.org/10.1175/JCLI-D-25-0418.s1>.

Corresponding author: Alexander J. Baker, alexander.baker@reading.ac.uk

DOI: 10.1175/JCLI-D-25-0418.1

© 2026 Author(s). This published article is licensed under the terms of a Creative Commons Attribution 4.0 International (CC BY 4.0) License



act to weaken and strengthen, respectively, the westerly zonal flow. Therefore, reducing uncertainty around the magnitude of regional warming (Steiner et al. 2020) will, in principle, constrain the midlatitude circulation response (Harvey et al. 2014; Peings et al. 2018). However, although zonal wind biases have improved across CMIP phases 3–6 (Harvey et al. 2020; Oudar et al. 2020), uncertainties associated with the responses of individual driving mechanisms have not reduced significantly (Harvey et al. 2023). Substantial spread in simulated Arctic and tropical amplification remains evident across CMIP6 models (Oudar et al. 2020). Simulated jet responses to climate change, therefore, exhibit similar spatial patterns across CMIP phases (Harvey et al. 2020). Moreover, climate models fail to capture the observed jet strengthening trend (as well as the downstream precipitation trend over Europe), indicating that projections of winter atmospheric circulation (and surface climate over Europe), based on models which fail to capture the jet's forced response, may be unreliable (Blackport and Fyfe 2022).

Models project a poleward jet shift due to tropical warming (Barnes and Polvani 2013; Oudar et al. 2020; Simpson et al. 2014), and similar trends in the Northern Hemisphere are emerging in satellite-era reanalyses (Woollings et al. 2023). Although weak, these trends are consistent with the predominance of tropical forcing, as opposed to Arctic warming, on the jet streams. However, trends in reanalyses lie only just within the spread of CMIP6 model trends, and reanalyses show stronger trends than models, despite a somewhat weaker positive temperature gradient trend compared with that simulated by models (Woollings et al. 2023). Weaker simulated trends may be due to models underrepresenting poleward eddy heat flux in the atmosphere, confining upper-tropospheric warming to the tropics, and insufficiently enhancing the meridional temperature gradient, inducing a weaker jet shift.

Capturing jet responses to anthropogenic climate change relies on models' ability to represent both large-scale variability and weather-scale processes. Increased horizontal resolution in climate models is, therefore, expected to improve the simulation of North Atlantic climate and increase confidence in climate projections (Baker et al. 2019). In fully coupled CMIP6 High-Resolution Model Intercomparison Project (HighResMIP) simulations, increasing atmosphere resolution from ~ 100 to ~ 25 km reduces sea surface temperature (SST) and, in turn, large-scale circulation biases (Athanasiadis et al. 2022). Increased resolution also improves the representation of North Atlantic weather regime clustering, although has little impact on models' ability to reproduce observed regime frequency and persistence (Fabiano et al. 2020). Additionally, Blackport and Fyfe (2022) found little difference in simulated jet speed trends (or trend spread) between high-resolution (HR) and low-resolution (LR) coupled models and found only small trend differences among simulations forced by observed SST, which suggests SST biases are unlikely to explain this disparity between models and observations. The impact of increasing resolution—in both the atmosphere and ocean—on large-scale circulation and, in turn, surface climate is, therefore, insufficiently well understood.

In this study, we assessed the impact of increased resolution on simulating the North Atlantic zonal wind and eddy-driven

jet in historical and near-future climates. Our aims are to (i) identify differences in climate change responses that arise when horizontal resolution is increased and (ii) assess whether HighResMIP provides evidence of a response of the westerly zonal flow to ocean resolution increase. To this end, we analyzed fully coupled simulations from HighResMIP over the period 1950–2050 and for boreal winter (i.e., December–February), focusing first on climatological zonal wind fields and then on the eddy-driven jet. This paper is structured as follows: Model data, observational data, and methods are described in section 2; results are presented in section 3, with further discussion and our conclusions summarized in section 4.

2. Data and methods

a. Model ensemble

This study is based on CMIP6 HighResMIP (Haarsma et al. 2016) fully coupled (tier 2) historical and future simulations from five global climate models (Table 1): CMCC-CM2 (Cherchi et al. 2019), CNRM-CM6.1 (Voldoire et al. 2019), EC-Earth3P (Haarsma et al. 2020), HadGEM3-GC3.1 (Roberts et al. 2019; Williams et al. 2018), and MPI-ESM1.2 (Gutjahr et al. 2019). Each simulation was performed under the Process-based Climate Simulation: Advances in High-Resolution Modeling and European Climate Risk Assessment (PRIMAVERA) program. Fully coupled historical (1950–2014) and future (2015–50) experiments are termed *hist-1950* and *highres-future*, respectively (Haarsma et al. 2016). We analyzed a single ensemble member for each model at each resolution for both *hist-1950* and *highres-future*. Additionally, we performed a separate analysis of multiple ensemble members available for HadGEM3-GC3.1 (Table S1 in the online supplemental material) to help assess the extent to which multimodel results are affected by internal variability. Seddon et al. (2023) described the curation of HighResMIP data.

Under the HighResMIP experimental protocol, minimal changes in model-tuning parameters were made between low- and high-resolution integrations to ensure that the analysis of resolution sensitivity is not confounded by substantial differences in model configurations between resolutions (Haarsma et al. 2016). Between low- and high-resolution configurations, no model-physics changes were made to the atmospheric components of CMCC-CM2, CNRM-CM6.1, and EC-Earth3P, and only minor adjustments were made to a single parameter in HadGEM3-GC3.1 (related to quasi-biennial oscillation period) and MPI-ESM1.2 (related to numerical stability). Shorter dynamical time steps were used in the high-resolution integrations of all models to ensure numerical stability. The effective resolutions of the models at high resolution resolve synoptic-scale dynamics in the atmosphere (Klaver et al. 2020). For ocean models, a key difference is that the effects of mesoscale eddies are parameterized at low resolutions of $\sim 1^\circ$ but are partially resolved at high resolutions of $\sim 0.25^\circ$ or finer (Roberts et al. 2019). Since this study concerns midlatitude phenomena, atmosphere resolutions are given as a model's regular mesh spacing at a latitude of 50° (Table 1). For convenience, we

TABLE 2. Reanalyses. Atmospheric mesh spacing at 50°N in units of kilometer is given in parentheses. 3(4)D-Var = 3(4)D variational data assimilation, GSI = Gridpoint Statistical Interpolation analysis system, and IAU = incremental analysis update.

Reanalysis	ERA5	JRA-55	MERRA-2	NCEP
Data assimilation	4D-Var	4D-Var	3D-Var, GSI, IAU	3D-Var, GSI
Resolution (grid spacing at 50°)	T639L137 (33 km)	TL319L60 (55 km)	Cubed sphere (50 km)	T382L64 (38 km)
Analysis period	1979–2018	1959–2017	1980–2018	1979–2016

Barnes and Polvani (2013) showed that patterns of jet variability are partly a function of jet position: As the jet shifts poleward, variability becomes less north–south and more akin to pulsing (i.e., variation in jet speed). This latitude–speed relationship is, therefore, a key descriptor of North Atlantic jet behavior and included in our analysis. We examined the inverse relationship between ϕ variance and jet speed (i.e., the standard deviation of ϕ binned by jet speed). At each identified ϕ , we computed jet speed, u' , as the square root of the sum of the squares of the low-pass-filtered zonal and meridional winds. Using both wind fields accounts for instances when the magnitude of jet speed is dominated by the meridional component. All data were regridded to the N96 grid of HadGEM3-GC3.1-LL to isolate resolution sensitivity from any improved sampling at higher resolution.

e. Significance testing

Significance was computed at the 99% level using Welch's unequal variances t test with respect to each individual model's internal year-to-year variability, as the HighResMIP ensemble size is insufficient to compute significance with respect to the ensemble.

3. Results

a. North Atlantic westerly zonal flow

We first focus on the climatological North Atlantic westerly zonal wind, quantifying multimodel-mean biases in, and the climate change response of, the vertical profile of the zonal flow during winter. The simulated zonal wind field is sensitive to model resolution. In all models except CMCC-CM2, weaker zonal winds are seen at high resolution at $\sim 50^\circ\text{N}$, with increased winds on the equatorward and poleward flanks of the jet (Fig. S1). This indicates a larger latitudinal range of the mid-latitude jet (i.e., a broader jet) when resolution is increased, consistent with Athanasiadis et al. (2022). At low resolution, several multimodel-mean zonal wind biases (vs the multireanalysis-mean wind field) are evident (Fig. 1a). For the subtropical jet, winds are too strong on the equatorward flank. For the midlatitude jet, winds are too strong on the equatorward flank and too weak on the poleward flank. At high resolution, mid- and high-latitude biases are significantly reduced, particularly around the jet core latitude at low levels and the subtropical winds throughout the depth of the troposphere, but winds at high resolution remain too strong on the equatorward flank of the midlatitude jet (Fig. 1b). Additionally, subtropical jet biases worsen somewhat, as noted by Moreno-Chamarro et al. (2022) for the annually averaged Northern Hemisphere subtropical jet. Overall, however, the climatological westerly zonal wind field across the North

Atlantic is significantly better represented at high resolution. Additionally, a separate analysis of multiple ensemble members available for HadGEM3-GC3.1 (Table S1) yields similar results at high resolution (Fig. S2), supporting the multimodel results.

Under climate change (to 2050), low-resolution models simulate a zonal wind decrease on the equatorward flank of the




FIG. 1. Multimodel-mean climatological biases in the vertical profile of the North Atlantic westerly zonal wind averaged for December–February in (a) low-resolution (LR) and (b) high-resolution (HR) simulations. Biases (i.e., hist-1950 minus reanalyses; shaded contours) were computed against the multireanalysis-mean climatology (black contours). Unit is meters per second, and teal stippling indicates where the bias is significant at the 99% level and where all models agree on the sign of the bias. All monthly data were regridded to the lowest-resolution model grid (HadGEM3-GC3.1-LL) to facilitate an intermodel comparison. For clarity, only selected isobaric levels are labeled on each panel's ordinate. An equivalent analysis of multiple HadGEM3-GC3.1 ensemble members (Fig. S2) yielded a similar overall pattern of biases at HR and at LR (excepting a negative bias at the equatorward flank of the subtropical jet) compared with the multimodel-mean results shown here.




FIG. 2. Multimodel-mean climatological vertical profile of the North Atlantic westerly zonal wind averaged for December–February in (a) LR and (b) HR simulations. Shown are the multimodel-mean historical climatology (black contours) (i.e., hist-1950) and the climate change response (shaded contours) (i.e., highres-future minus hist-1950). Differences were computed using monthly data for the 30-yr periods 1950–79 and 2020–49. Unit is meters per second, and teal stippling indicates where the response is significant at the 99% level and where all models agree on the sign of the response. All data were re-gridded to the lowest-resolution model grid (HadGEM3-GC3.1-LL) to facilitate an intermodel comparison. For clarity, only selected isobaric levels are labeled on each panel’s ordinate. An equivalent analysis of multiple HadGEM3-GC3.1 ensemble members (Fig. S4) yielded a similar overall pattern of responses compared with the multimodel-mean results shown here.

subtropical jet and poleward flank of the midlatitude jet, with an increase in the winds between the jets ($\sim 30^\circ\text{N}$)—both at upper levels. This indicates a future reduction and shallowing of the interjet separation (Fig. 2). The most favorable region for baroclinic wave development lies along the subtropical jet only when the subtropical jet is sufficiently strong; a moderate-strength subtropical jet is typically associated with greater meridional separation between the subtropical and eddy-driven jets (Lee and Kim 2003). At high resolution, the weakening of mid- and upper-level westerlies at higher latitudes (poleward of $\sim 60^\circ\text{N}$) in future is more significant (Fig. 2b). This indicates a shift toward reduced jet separation in future when resolution is increased, although this response varies across models and is clearest in CNRM-CM6.1, EC-Earth3P, and HadGEM3-GC3.1 (Fig. S3). Additionally, a separate analysis of multiple

HadGEM3-GC3.1 ensemble members reveals a similar ensemble-mean response for this model (Fig. S4), supporting the multimodel results. Overall, the climate change responses at low and high resolution are similar, except for the enhanced decrease in mid- and upper-level subtropical winds at high resolution, the region where a negative wind bias is significantly reduced at high resolution (Fig. 1b). Low-level responses in the midlatitudes are uncertain. Stronger winds at high resolution are seen at $\sim 50^\circ\text{N}$, but this response is not significant in the multimodel mean, indicating that a forced response will not emerge by 2050.

b. Trends in North Atlantic meridional temperature gradients

The response of the North Atlantic westerlies is sensitive to the local equator-to-pole temperature difference, dT/dy (Harvey et al. 2014). Therefore, to explain the zonal wind responses to future forcing, we analyzed trends in bulk meridional temperature gradients, computed throughout the troposphere over the simulated period (1950–2050). Negative trends exist at low levels due to Arctic amplification of warming in the lower troposphere, and positive trends are seen aloft due to the amplification of warming in the tropical upper troposphere (Fig. 3).

At low resolution, the vertical profile of multimodel-mean trends in HighResMIP is generally consistent with those derived from reanalysis datasets (Lee et al. 2019), albeit computed over a longer time frame here (Fig. 3). Several differences between low- and high-resolution simulations are apparent. At low resolution, the transition from negative to positive trends occurs between 400 and 500 hPa, consistent with reanalyses. However, this transition occurs at a lower level (between 850 and 700 hPa) at high resolution (Fig. 3). Additionally, individual high-resolution models exhibit their strongest positive trends at different levels in the mid- and upper troposphere (Fig. 3). The magnitudes of negative and positive trends in the lower- and upper troposphere, respectively, are weaker in the high-resolution multimodel mean, and this is also generally true of individual models, particularly EC-Earth3P and HadGEM3-GC3.1. These results may to some extent be influenced by the lack of model retuning of the global energy balance at high resolution, an aspect of the HighResMIP protocol (Haarsma et al. 2016).

There is an indication that ocean resolution influences bulk meridional temperature gradients. CMCC-CM2 and MPI-ESM1.2 show little change in low-level negative trends between atmosphere resolutions (Fig. 3), and these are models where ocean resolution is unchanged. For HadGEM3-GC3.1, simulations performed with different ocean resolutions (ranging from 1° to $1/12^\circ$) allow insight into the impact of partially resolving ocean eddies. HadGEM3-GC3.1 shows a clear decrease in the magnitudes of low-level negative trends with increasing ocean resolution (Fig. 3). Comparing simulations with ocean resolutions of 0.25° [medium-resolution atmosphere and medium-resolution ocean (MM)] and $1/12^\circ$ [medium-resolution atmosphere and high-resolution ocean (MH)] for the same atmosphere resolution (60 km) shows less pronounced low-level negative trends with the higher-resolution ocean. Conversely,




FIG. 3. Tropospheric vertical profiles of trends (1950–2050) in the winter-mean (December–February) meridional temperature difference over the North Atlantic, denoted ΔT , in hist-1950 and highres-future simulations. (first column) Multimodel-mean trends are shown at LR and HR, with error bars denoting the standard deviation among model trends. Individual model trends are separated by panel columns, and (from top to bottom) available resolutions are ordered from low to high, with error bars denoting the 95% confidence intervals for the linear trends. Red and blue colors denote positive and negative trends, respectively.

simulations with atmosphere resolutions of 60 (MH) and 25 km [high-resolution atmosphere and high-resolution ocean (HH)] for the same ocean resolution ($1/12^\circ$) exhibit highly similar vertical profiles. These results highlight the possible influence of ocean resolution on the bulk meridional temperature gradients which modulate midlatitude circulation. Potentially, this reflects higher poleward eddy heat transport at high ocean resolution (Roberts et al. 2019) acting to reduce the low-level gradients via air–sea interactions (Gulev et al. 2013). However, an initial analysis of surface turbulent heat fluxes (not shown) did not reveal similar sensitivity to ocean resolution. An important caveat here is that a larger multimodel ensemble, including multiple ocean resolutions, is required to investigate this.

In these coupled simulations, future zonal wind changes at each isobaric level with respect to changes at the surface (i.e., the vertical wind shear) are in near-exact thermal wind balance with the changes in meridional temperature gradient (Fig. S5). Blackport and Fyfe (2022) and Woollings et al. (2023) reported that dT/dy in the upper troposphere is stronger in models than observational data, yet concurrent changes in the upper-tropospheric zonal wind are smaller than expected from this temperature gradient. This implies that simulated

zonal wind trends are not in accordance with thermal wind balance. However, we find that thermal wind balance holds when the integral of dT/dy (over 1000 hPa to a given isobaric level) is taken instead of a single level.

Lee et al. (2019) showed that the compensating influences of the lower- and upper-level meridional temperature gradient trends result in no discernible historical trend in upper-level midlatitude zonal wind speed. Here, given thermal wind balance holds, we attempt to relate differences between the low- and high-resolution tropospheric trend profiles with differences between the low- and high-resolution zonal wind responses (Fig. 2). These trends in meridional temperature gradients are generally consistent with the simulated future responses of the North Atlantic zonal wind field. Stronger winds aloft at both low and high resolution (Fig. 2) may be explained by the positive trends in meridional temperature gradients at upper levels. At low resolution, strengthened future westerly winds occur above ~ 500 hPa, but strengthened midlatitude winds are present at lower levels (above ~ 850 hPa) at high resolution—both consistent with the approximate levels at which the transitions between negative and positive temperature trends occur. High-ResMIP models, therefore, provide further evidence that




FIG. 4. (a) Frequency distribution of December–February eddy-driven jet latitude ϕ computed using low-pass-filtered daily data following Woollings et al. (2010) for reanalyses and hist-1950 simulations (highres-future distributions are shown in Fig. S6). (b) Response of ϕ in highres-future simulations. Unit in (a) and (b) is normalized frequency density. (c) Standard deviation of ϕ as a function of jet speed u' (given as percentiles, with a bin width of 10%), computed from low-pass-filtered daily zonal and meridional wind fields following Woollings et al. (2018b). (d) Response of ϕ variance in highres-future simulations. Unit in (c) and (d) is degrees. Gray shading in (a) and (c) indicates the standard deviation of reanalyses. Shown in all panels are multimodel-mean (bold lines) and individual model (thin lines) curves for LR (solid) and HR (dashed) simulations. An equivalent analysis of multiple HadGEM3-GC3.1 ensemble members is shown in Fig. S7.

tropical amplification is potentially a stronger actor in determining the future zonal wind response. A caveat to this interpretation, however, is the relatively small ensemble size of HighResMIP.

c. North Atlantic eddy-driven jet

In this section, we analyze the eddy-driven jet using a jet latitude index, which describes three regimes: northern ($\sim 53^\circ$ – 60° N), central ($\sim 45^\circ$ – 50° N), and southern ($\sim 35^\circ$ – 40° N). This apparent trimodality of the eddy-driven jet is captured by HighResMIP models (Fig. 4a), an aspect not adequately simulated by a majority of CMIP5 models (Iqbal et al. 2018) nor by several CMIP6 models over the central and eastern North Atlantic (Dorrington et al. 2022; Oudar et al. 2020). The most frequent central jet latitude regime is better simulated at high resolution (Fig. 4a), confirming a complementary analysis of HighResMIP simulations (Athanasiadis et al. 2022) and atmosphere-only, single-model results (Baker et al. 2019). The northern regime is less prominent in the HighResMIP multimodel mean compared with reanalyses (Fig. 4a), which is also seen in CMIP6 models (Oudar et al. 2020). Furthermore, the distribution of jet-latitude variance as a function of jet speed (percentile), described by a characteristic negative curve, is also better captured at high resolution (Fig. 4c). At low resolution, simulated variance is too low compared with

reanalyses, although greater interreanalysis and intermodel spread are evident at lower speed percentiles.

Under climate change, the multimodel-mean response of jet latitude at low resolution is a frequency decrease in the central and northern regimes and an increase in the southern regime—in other words, an equatorward shift (Fig. 4b). At high resolution, however, responses differ: a frequency increase in the central and northern regimes and a decrease in the southern regime—in other words, a poleward shift (Fig. 4b). Strictly, these are shifts in the frequencies of each jet regime, rather than in the absolute latitudes at which regimes occur (Fig. S6). Additionally, reduced meridional variability is projected at high resolution (Fig. 4c), indicating a narrowing of the jet that is consistent with the mean westerly flow (Fig. 2). A separate analysis of the available HadGEM3-GC3.1 ensemble members (Table S1) reveals a poleward shift across resolutions for this model, and this shift is stronger at high resolution (Fig. S7a), consistent with Baker et al. (2019) and broadly corroborating multimodel results. The climate change response of variance in jet latitude is also sensitive to resolution. At low resolution, the response of jet-latitude variance is close to zero at all speed percentiles (Fig. 4d). At high resolution, however, a decrease in variance is simulated at nearly all speed percentiles (Fig. 4d), indicating a more persistent jet. For this diagnostic, high-resolution simulations are potentially more trustworthy because the historical latitude–speed relationship is ostensibly




FIG. 5. Bivariate distributions of December–February eddy-driven jet latitude ϕ and jet speed u' (percentiles) in (a) reanalyses and multimodel-mean biases in these quantities in (b) LR and (c) HR hist-1950 simulations; (d) sensitivity to resolution (i.e., HR minus LR); and the multimodel-mean climate change response (i.e., highres-future minus hist-1950) at (e) LR and (f) HR. Zonal and meridional wind fields were low-pass filtered as in Fig. 4, and the unit is normalized frequency density ($\times 10^{-3}$). Overlain black contour (at 0.4×10^{-3}) indicates the multireanalysis-mean field in (b) and (c) and the historical multimodel-mean field in (e) and (f). Reanalyses are shown individually in Fig. S8.

closer to reanalyses at ~ 25 -km resolution. A separate analysis of the HadGEM3-GC3.1 ensemble also shows little change of jet-latitude variance at low resolution but reduced variance across speed percentiles at high resolution (Fig. S7b), substantiating the multimodel result. This is consistent with a meridional narrowing of the westerly flow (Peings et al. 2018) and reduced winter blocking (Gao et al. 2025; Woollings et al. 2018a), although differs from a recent analysis of lower-resolution CMIP6 models (Dorrington et al. 2022). These results are somewhat inconsistent with an analysis of large-scale waviness in High-ResMIP (Yamamoto and Martineau 2024), which showed a significant reduction in atmospheric waviness at low resolution but a less clear response at high resolution, although their model ensemble differs from that of this study (Table 1). Additionally, historical HighResMIP simulations show limited improvement with increased resolution in capturing persistent episodes of European blocking (Schiemann et al. 2020), indicating that some sensitivity to the metric used to characterize large-scale variability exists.

To examine eddy-driven jet behavior in more detail, we analyzed bivariate distributions of jet latitude and speed. The multireanalysis-mean bivariate distribution shows that the central jet regime is associated with higher jet speeds compared with the northern and southern regimes (Fig. 5a), and the southern regime is associated particularly with low speeds.

Reanalyses show little variation (Fig. S8, upper row). To evaluate simulated bivariate distributions, multimodel-mean biases (Figs. 5b,c) and individual model biases (Fig. S8, middle and lower rows) were computed. At low resolution, there is a clear positive bias for the central jet regime at $\sim 45^\circ\text{N}$, and negative biases are seen for both the southern and northern jet regimes (Fig. 5b). At high resolution, the central jet regime bias is reduced (Fig. 5c), which is important because this is both the most frequent regime and the regime associated with the strongest steering flow for low pressure weather systems impacting Europe (Baker et al. 2019; Harvey et al. 2023; Woollings et al. 2018b). However, at high resolution, the negative northern regime bias persists, and a positive bias emerges between 30° and 40°N , where high-resolution models simulate high-speed southern jet instances that are not present in reanalyses. This low-latitude bias in the multimodel mean is predominantly due to CNRM-CM6.1 and MPI-ESM1.2 (Fig. S8). Additionally, simulated historical distributions show an equatorward shift in jet latitude when resolution is increased (Fig. 5d), with less frequent central and northern jet occurrences, particularly at higher speed percentiles, and more frequent southern jet occurrences (Fig. 5d), consistent with both the resolution sensitivity of the mean zonal wind (Fig. 1) and with a recent complementary analysis of the North Atlantic midlatitude storm track (Lockwood et al. 2025). Although the multimodel-

mean climatological variance in jet latitude as a function of jet speed is better represented at high resolution (Fig. 4c), diverse individual model biases are found in HighResMIP (Fig. S8). The occurrence of each jet latitude regime at a particular range in jet speed, as apparent in reanalyses, is not better captured when resolution is increased from ~ 100 to ~ 25 km in the midlatitudes.

Under climate change, changes in latitude–speed distributions are small. At low resolution, a small equatorward shift is projected, but the overall pattern is equivocal (Fig. 5e). At high resolution, projected increases in frequency across northern and central jet regimes are clearest, with decreases across the southern regime and smaller decreases in the lowest and highest latitudes (Fig. 5f), again resulting in a narrowed jet (and a relative poleward shift), consistent with Fig. 4d. Additionally, at high resolution, the eddy-driven jet’s response to climate change may be influenced by the equatorward-shifted historical mean state (Fig. 5d). The two models for which ocean resolution is unchanged (CMCC-CM2 and MPI-ESM1.2) exhibit little difference between low- and high-resolution responses (Fig. S9), again indicating that dynamical coupling with a higher-resolution ocean may influence the circulation response.

4. Summary and discussion

a. Main results

In this study, we assessed the impact of model resolution on the winter climatological zonal wind and eddy-driven jet position and speed, simulated under historical (1950–2014) and future (2015–50) climate conditions in an ensemble of fully coupled global climate models at resolutions ranging from ~ 100 to ~ 25 km in the atmosphere and 1° – $1/12^\circ$ in the ocean. Increasing resolution significantly improves the representation of the midlatitude jet and its eddy-driven component, and responses to climate change differ between low- and high-resolution simulations. The main findings of our work are as follows:

- Increasing resolution significantly improves (vs reanalyses) the North Atlantic climatological zonal wind field at mid- and high latitudes, but biases remain around the low-latitude, equatorward flank of the upper-level subtropical jet (Fig. 1).
- By 2050, low-resolution models simulate an equatorward shift in the midlatitude jet, reducing the upper-level meridional separation of the subtropical and midlatitude jets, but high-resolution models project a strengthening of the jets and a small poleward shift of the midlatitude jet (Fig. 2).
- Trends in the large-scale meridional temperature gradient over the North Atlantic suggests that tropical amplification influences the future zonal wind response, and there is an indication of sensitivity of lower-level temperature gradient trends to ocean resolution (Fig. 3).
- At low resolution, the eddy-driven jet shows little mean meridional shift under climate change. At high resolution, however, climate change reduces the jet’s latitudinal variance (Figs. 4 and 5).

b. Discussion and outlook

In HighResMIP simulations, future zonal wind changes at a given isobaric level with respect to changes at the surface (i.e.,

the vertical wind shear) are in thermal wind balance with the changes in meridional temperature gradient. To explain models’ zonal wind responses to future forcing, we analyzed trends in meridional temperature gradients. These trends were found to be generally consistent with the simulated future responses of the North Atlantic zonal wind field. Strengthened future westerly winds aloft (Fig. 2) may be explained by positive trends in meridional temperature gradients at upper levels. Stronger westerlies occur above ~ 500 hPa at low resolution and above ~ 850 hPa at high resolution, consistent with the approximate levels at which the transitions between negative and positive temperature gradient trends occur (Fig. 3). In HighResMIP, tropical amplification appears a stronger actor in determining the future zonal wind response, consistent with recent analyses (Oudar et al. 2020; Woollings et al. 2023). CMIP6 models overestimate warming of the tropical midtroposphere (Mitchell et al. 2020) and upper (Blackport and Fyfe 2022) troposphere. These biases are potentially due to the representation of convection in low-resolution CMIP6 models, which tend to overestimate the amount of upper-tropospheric warming for a given level of lower-tropospheric warming (Keil et al. 2021; Mitchell et al. 2020). This is due to low entrainment rates warming the upper troposphere by allowing convection and latent heating further aloft. Further increases in atmosphere resolution to reduce reliance on parameterized convection may resolve this (Blackport and Fyfe 2022).

We also analyzed the eddy-driven component of the westerly flow, focusing on jet position and on variance in jet position as a function of jet speed. HighResMIP models capture the trimodality of the jet latitude index, which CMIP5 and several CMIP6 models do not, and both the overall distribution of jet latitude and the variance of jet latitude as a function of jet speed are better simulated at high resolution (Fig. 4). This indicates that resolution is a key component of capturing eddy-driven jet behavior in models. However, the occurrence of each jet latitude regime at a typical range in jet speed, which is seen in reanalyses, is not well captured when resolution is increased from ~ 100 to 25 km in the midlatitudes. Under climate change, the eddy-driven jet shifts equatorward at low resolution but narrows and shifts poleward at high resolution. Increased ocean resolution may influence the high-resolution near-future response, but a key caveat to these results is that the ensemble size considered here is relatively small.

To help increase confidence in atmospheric circulation projections, future work should focus on expanding the ensemble size, by conducting ensemble simulations for multiple models, and further increasing model resolution in both the atmosphere and ocean. In HighResMIP, the use of an eddy-permitting ocean model in a fully coupled framework increases blocking frequency (Athanasiadis et al. 2022), and SST biases are improved significantly in models where ocean resolution is increased (or physics improved) alongside increased atmosphere resolution (Moreno-Chamarro et al. 2022). Finer resolution may improve model biases further, as eddy forcing is too weak in HighResMIP models (Dorrington et al. 2022), leading to underpersistent regime behavior (Fabiano et al. 2020). This study provides evidence that jet latitude regime persistence increases at ~ 25 km but, together with existing studies of the same simulations

(Blackport and Fyfe 2022; Dorrington et al. 2022; Fabiano et al. 2020; Moreno-Chamarro et al. 2022; Schiemann et al. 2020), indicates that the contribution of eddy forcing to North Atlantic climate variability remains insufficiently captured. Here, we see the evidence of influence of ocean resolution on trends in low-level meridional temperature gradients, which do not change in coupled models where ocean resolution is constant. With an eddy-permitting ocean resolution, low-level temperature gradients are reduced, an effect that may be enhanced by the higher eddy activity in response to warming seen in recent eddy-rich ocean modeling (Beech et al. 2022; Li et al. 2024).

Models may also underestimate the magnitude of multidecadal and centennial internal variability (Marcheggiani et al. 2023; Smith et al. 2020). Centennial-length simulations extending beyond 2050 are needed to address two outstanding concerns. First, this study indicates that climate change signals in eddy-driven jet latitude and speed may not emerge by 2050. The projected atmospheric circulation response to Arctic sea ice loss is also weak (Smith et al. 2022), and much uncertainty remains around the impact of Arctic amplification on midlatitude weather (Cohen et al. 2014). Second, proxy evidence points to high internal variability, enveloping recent observed jet shifts, which makes extracting anthropogenically forced changes challenging (Osman et al. 2021). Longer, higher-resolution simulations with fully coupled models, such as those proposed by Roberts et al. (2025), may help address the uncertainty around changes in North Atlantic atmospheric circulation and surface climate over the twenty-first century.

Acknowledgments. The authors are grateful to the modeling teams that produced the PRIMAVERA simulations and to J. Seddon for data curation. All authors received financial support from the PRIMAVERA project (European Commission Horizon2020 Grant Agreement 641727) with data access supported by IS-ENES3 (Grant Agreement 824084) via the JASMIN facility (jasmin.ac.uk), which is operated by the Science and Technology Facilities Council on behalf of the Natural Environment Research Council. A.J.B. received additional support from a Natural Environment Research Council national capability programme, the North Atlantic Climate System: Integrated Study (Grant Agreements NE/N018001/1, NE/N018044/1, NE/N018028/1, and NE/N018052/1). A. J. B. and P. L. V. conceived the study. The authors are grateful to the editor, R. C. J. Wills, and to K. Strommen and three anonymous reviewers for their constructive feedback on the initial draft of this paper. A. J. B. performed data analysis and figure preparation and wrote the manuscript. J. F. L. performed the analysis presented in Fig. S5. All authors approved the final manuscript draft. The authors declare no competing interests.

Data availability statement. Reanalysis data are available from cds.climate.copernicus.eu, rda.ucar.edu or disc.gsfc.nasa.gov. Model data are available from the Earth System Grid Federation nodes (esgf.llnl.gov). Data analysis and visualization code are available from the corresponding author upon request.

REFERENCES

- Athanasiadis, P. J., and Coauthors, 2022: Mitigating climate biases in the midlatitude North Atlantic by increasing model resolution: SST gradients and their relation to blocking and the jet. *J. Climate*, **35**, 6985–7006, <https://doi.org/10.1175/JCLI-D-21-0515.1>.
- Baker, A. J., and Coauthors, 2019: Enhanced climate change response of wintertime North Atlantic circulation, cyclonic activity, and precipitation in a 25-km-resolution global atmospheric model. *J. Climate*, **32**, 7763–7781, <https://doi.org/10.1175/JCLI-D-19-0054.1>.
- Barnes, E. A., and L. Polvani, 2013: Response of the midlatitude jets, and of their variability, to increased greenhouse gases in the CMIP5 models. *J. Climate*, **26**, 7117–7135, <https://doi.org/10.1175/JCLI-D-12-00536.1>.
- Beech, N., T. Rackow, T. Semmler, S. Danilov, Q. Wang, and T. Jung, 2022: Long-term evolution of ocean eddy activity in a warming world. *Nat. Climate Change*, **12**, 910–917, <https://doi.org/10.1038/s41558-022-01478-3>.
- Blackport, R., and J. C. Fyfe, 2022: Climate models fail to capture strengthening wintertime North Atlantic jet and impacts on Europe. *Sci. Adv.*, **8**, eabn3112, <https://doi.org/10.1126/sciadv.abn3112>.
- Cherchi, A., and Coauthors, 2019: Global mean climate and main patterns of variability in the CMCC-CM2 coupled model. *J. Adv. Model. Earth Syst.*, **11**, 185–209, <https://doi.org/10.1029/2018MS001369>.
- Cohen, J., and Coauthors, 2014: Recent Arctic amplification and extreme mid-latitude weather. *Nat. Geosci.*, **7**, 627–637, <https://doi.org/10.1038/ngeo2234>.
- Dorrington, J., K. Strommen, and F. Fabiano, 2022: Quantifying climate model representation of the wintertime Euro-Atlantic circulation using geopotential-jet regimes. *Wea. Climate Dyn.*, **3**, 505–533, <https://doi.org/10.5194/wcd-3-505-2022>.
- Duchon, C. E., 1979: Lanczos filtering in one and two dimensions. *J. Appl. Meteor.*, **18**, 1016–1022, [https://doi.org/10.1175/1520-0450\(1979\)018<1016:LFIOAT>2.0.CO;2](https://doi.org/10.1175/1520-0450(1979)018<1016:LFIOAT>2.0.CO;2).
- Fabiano, F., H. M. Christensen, K. Strommen, P. Athanasiadis, A. Baker, R. Schiemann, and S. Corti, 2020: Euro-Atlantic Weather Regimes in the PRIMAVERA coupled climate simulations: Impact of resolution and mean state biases on model performance. *Climate Dyn.*, **54**, 5031–5048, <https://doi.org/10.1007/s00382-020-05271-w>.
- Gao, Y., and Coauthors, 2025: Enhanced simulation of atmospheric blocking in a high-resolution Earth system model: Projected changes and implications for extreme weather events. *J. Geophys. Res. Atmos.*, **130**, e2024JD042045, <https://doi.org/10.1029/2024JD042045>.
- Gulev, S. K., M. Latif, N. Keenlyside, W. Park, and K. P. Koltermann, 2013: North Atlantic Ocean control on surface heat flux on multidecadal timescales. *Nature*, **499**, 464–467, <https://doi.org/10.1038/nature12268>.
- Gutjahr, O., D. Putrasahan, K. Lohmann, J. H. Jungclaus, J. S. von Storch, N. Brüggemann, H. Haak, and A. Stössel, 2019: Max Planck Institute Earth System Model (MPI-ESM1.2) for the High-Resolution Model Intercomparison Project (High-ResMIP). *Geosci. Model Dev.*, **12**, 3241–3281, <https://doi.org/10.5194/gmd-12-3241-2019>.
- Haarsma, R., and Coauthors, 2016: High Resolution Model Intercomparison Project (HighResMIP v1.0) for CMIP6. *Geosci. Model Dev.*, **9**, 4185–4208, <https://doi.org/10.5194/gmd-9-4185-2016>.

- , and Coauthors, 2020: HighResMIP versions of EC-Earth: EC-Earth3P and EC-Earth3P-HR—Description, model computational performance and basic validation. *Geosci. Model Dev.*, **13**, 3507–3527, <https://doi.org/10.5194/gmd-13-3507-2020>.
- Harvey, B., L. C. Shaffrey, and T. J. Woollings, 2014: Equator-to-pole temperature differences and the extra-tropical storm track responses of the CMIP5 climate models. *Climate Dyn.*, **43**, 1171–1182, <https://doi.org/10.1007/s00382-013-1883-9>.
- , P. Cook, L. C. Shaffrey, and R. Schiemann, 2020: The response of the Northern Hemisphere storm tracks and jet streams to climate change in the CMIP3, CMIP5, and CMIP6 climate models. *J. Geophys. Res. Atmos.*, **125**, e2020JD032701, <https://doi.org/10.1029/2020JD032701>.
- , E. Hawkins, and R. Sutton, 2023: Storylines for future changes of the North Atlantic jet and associated impacts on the UK. *Int. J. Climatol.*, **43**, 4424–4441, <https://doi.org/10.1002/joc.8095>.
- Hersbach, H., and Coauthors, 2020: The ERA5 global reanalysis. *Quart. J. Roy. Meteor. Soc.*, **146**, 1999–2049, <https://doi.org/10.1002/qj.3803>.
- Hoskins, B. J., and K. I. Hodges, 2002: New perspectives on the Northern Hemisphere winter storm tracks. *J. Atmos. Sci.*, **59**, 1041–1061, [https://doi.org/10.1175/1520-0469\(2002\)059<1041:NPTNH>2.0.CO;2](https://doi.org/10.1175/1520-0469(2002)059<1041:NPTNH>2.0.CO;2).
- Iqbal, W., W.-N. Leung, and A. Hannachi, 2018: Analysis of the variability of the North Atlantic eddy-driven jet stream in CMIP5. *Climate Dyn.*, **51**, 235–247, <https://doi.org/10.1007/s00382-017-3917-1>.
- Keil, P., H. Schmidt, B. Stevens, and J. Bao, 2021: Variations of tropical lapse rates in climate models and their implications for upper tropospheric warming. *J. Climate*, **34**, 9747–9761, <https://doi.org/10.1175/JCLI-D-21-0196.1>.
- Klaver, R., R. Haarsma, P. L. Vidale, and W. Hazeleger, 2020: Effective resolution in high resolution global atmospheric models for climate studies. *Atmos. Sci. Lett.*, **21**, e952, <https://doi.org/10.1002/asl.952>.
- Kobayashi, S., and Coauthors, 2015: The JRA-55 reanalysis: General specifications and basic characteristics. *J. Meteor. Soc. Japan*, **93**, 5–48, <https://doi.org/10.2151/jmsj.2015-001>.
- Lee, S., and H.-K. Kim, 2003: The dynamical relationship between subtropical and eddy-driven jets. *J. Atmos. Sci.*, **60**, 1490–1503, [https://doi.org/10.1175/1520-0469\(2003\)060<1490:TDRBSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(2003)060<1490:TDRBSA>2.0.CO;2).
- Lee, S. H., P. D. Williams, and T. H. A. Frame, 2019: Increased shear in the North Atlantic upper-level jet stream over the past four decades. *Nature*, **572**, 639–642, <https://doi.org/10.1038/s41586-019-1465-z>.
- Li, X., Q. Wang, S. Danilov, N. Koldunov, C. Liu, V. Müller, D. Sidorenko, and T. Jung, 2024: Eddy activity in the Arctic Ocean projected to surge in a warming world. *Nat. Climate Change*, **14**, 156–162, <https://doi.org/10.1038/s41558-023-01908-w>.
- Lockwood, J. F., and Coauthors, 2025: The effect of increasing model resolution on the Northern Hemisphere winter midlatitude storm track: An equatorward shift due to contraction of the Hadley cell. *J. Climate*, **38**, 4539–4551, <https://doi.org/10.1175/JCLI-D-24-0414.1>.
- Marcheggiani, A., J. Robson, P.-A. Monerie, T. J. Bracegirdle, and D. Smith, 2023: Decadal predictability of the North Atlantic eddy-driven jet in winter. *Geophys. Res. Lett.*, **50**, e2022GL102071, <https://doi.org/10.1029/2022GL102071>.
- Mitchell, D. M., Y. T. Eunice Lo, W. J. M. Seviour, L. Haimberger, and L. M. Polvani, 2020: The vertical profile of recent tropical temperature trends: Persistent model biases in the context of internal variability. *Environ. Res. Lett.*, **15**, 1040b4, <https://doi.org/10.1088/1748-9326/ab9af7>.
- Molod, A., L. Takacs, M. Suarez, and J. Bacmeister, 2015: Development of the GEOS-5 atmospheric general circulation model: Evolution from MERRA to MERRA2. *Geosci. Model Dev.*, **8**, 1339–1356, <https://doi.org/10.5194/gmd-8-1339-2015>.
- Moreno-Chamarro, E., and Coauthors, 2022: Impact of increased resolution on long-standing biases in HighResMIP-PRIMAVERA climate models. *Geosci. Model Dev.*, **15**, 269–289, <https://doi.org/10.5194/gmd-15-269-2022>.
- Osman, M. B., S. Coats, S. B. Das, J. R. McConnell, and N. Chellman, 2021: North Atlantic jet stream projections in the context of the past 1,250 years. *Proc. Natl. Acad. Sci. USA*, **118**, e2104105118, <https://doi.org/10.1073/pnas.2104105118>.
- Oudar, T., J. Cattiaux, and H. Douville, 2020: Drivers of the northern extratropical eddy-driven jet change in CMIP5 and CMIP6 models. *Geophys. Res. Lett.*, **47**, e2019GL086695, <https://doi.org/10.1029/2019GL086695>.
- Parker, T., T. Woollings, A. Weisheimer, C. O'Reilly, L. Baker, and L. Shaffrey, 2019: Seasonal predictability of the winter North Atlantic Oscillation from a jet stream perspective. *Geophys. Res. Lett.*, **46**, 10159–10167, <https://doi.org/10.1029/2019GL084402>.
- Peings, Y., J. Cattiaux, S. J. Vavrus, and G. Magnusdottir, 2018: Projected squeezing of the wintertime North-Atlantic jet. *Environ. Res. Lett.*, **13**, 074016, <https://doi.org/10.1088/1748-9326/aacc79>.
- Perez, J., A. C. Maycock, S. D. Griffiths, S. C. Hardiman, and C. M. McKenna, 2024: A new characterisation of the North Atlantic eddy-driven jet using two-dimensional moment analysis. *Wea. Climate Dyn.*, **5**, 1061–1078, <https://doi.org/10.5194/wcd-5-1061-2024>.
- Roberts, M. J., and Coauthors, 2019: Description of the resolution hierarchy of the global coupled HadGEM3-GC3.1 model as used in CMIP6 HighResMIP experiments. *Geosci. Model Dev.*, **12**, 4999–5028, <https://doi.org/10.5194/gmd-12-4999-2019>.
- , and Coauthors, 2025: High-Resolution Model Intercomparison Project phase 2 (HighResMIP2) towards CMIP7. *Geosci. Model Dev.*, **18**, 1307–1332, <https://doi.org/10.5194/gmd-18-1307-2025>.
- Saha, S., and Coauthors, 2014: The NCEP Climate Forecast System version 2. *J. Climate*, **27**, 2185–2208, <https://doi.org/10.1175/JCLI-D-12-00823.1>.
- Schiemann, R., and Coauthors, 2020: Northern Hemisphere blocking simulation in current climate models: Evaluating progress from the Climate Model Intercomparison Project Phase 5 to 6 and sensitivity to resolution. *Wea. Climate Dyn.*, **1**, 277–292, <https://doi.org/10.5194/wcd-1-277-2020>.
- Seddon, J., A. Stephens, M. S. Mizielinski, P. L. Vidale, and M. J. Roberts, 2023: Technology to aid the analysis of large-volume multi-institute climate model output at a central analysis facility (PRIMAVERA Data Management Tool V2.10). *Geosci. Model Dev.*, **16**, 6689–6700, <https://doi.org/10.5194/gmd-16-6689-2023>.
- Shepherd, T. G., 2014: Atmospheric circulation as a source of uncertainty in climate change projections. *Nat. Geosci.*, **7**, 703–708, <https://doi.org/10.1038/ngeo2253>.
- Simpson, I. R., T. A. Shaw, and R. Seager, 2014: A diagnosis of the seasonally and longitudinally varying midlatitude circulation response to global warming. *J. Atmos. Sci.*, **71**, 2489–2515, <https://doi.org/10.1175/JAS-D-13-0325.1>.

- Smith, D. M., and Coauthors, 2020: North Atlantic climate far more predictable than models imply. *Nature*, **583**, 796–800, <https://doi.org/10.1038/s41586-020-2525-0>.
- , and Coauthors, 2022: Robust but weak winter atmospheric circulation response to future Arctic sea ice loss. *Nat. Commun.*, **13**, 727, <https://doi.org/10.1038/s41467-022-28283-y>.
- Steiner, A. K., and Coauthors, 2020: Observed temperature changes in the troposphere and stratosphere from 1979 to 2018. *J. Climate*, **33**, 8165–8194, <https://doi.org/10.1175/JCLI-D-19-0998.1>.
- Volodire, A., and Coauthors, 2019: Evaluation of CMIP6 DECK experiments with CNRM-CM6-1. *J. Adv. Model. Earth Syst.*, **11**, 2177–2213, <https://doi.org/10.1029/2019MS001683>.
- Williams, K. D., and Coauthors, 2018: The Met Office global coupled model 3.0 and 3.1 (GC3.0 and GC3.1) configurations. *J. Adv. Model. Earth Syst.*, **10**, 357–380, <https://doi.org/10.1002/2017MS001115>.
- Woollings, T., 2010: Dynamical influences on European climate: An uncertain future. *Philos. Trans. Roy. Soc.*, **A368**, 3733–3756, <https://doi.org/10.1098/rsta.2010.0040>.
- , A. Hannachi, and B. Hoskins, 2010: Variability of the North Atlantic eddy-driven jet stream. *Quart. J. Roy. Meteor. Soc.*, **136**, 856–868, <https://doi.org/10.1002/qj.625>.
- , and Coauthors, 2018a: Blocking and its response to climate change. *Curr. Climate Change Rep.*, **4**, 287–300, <https://doi.org/10.1007/s40641-018-0108-z>.
- , and Coauthors, 2018b: Daily to decadal modulation of jet variability. *J. Climate*, **31**, 1297–1314, <https://doi.org/10.1175/JCLI-D-17-0286.1>.
- , M. Drouard, C. H. O'Reilly, D. M. H. Sexton, and C. McSweeney, 2023: Trends in the atmospheric jet streams are emerging in observations and could be linked to tropical warming. *Commun. Earth Environ.*, **4**, 125, <https://doi.org/10.1038/s43247-023-00792-8>.
- Yamamoto, A., and P. Martineau, 2024: On the driving factors of the future changes in the wintertime Northern-Hemisphere atmospheric waviness. *Geophys. Res. Lett.*, **51**, e2024GL108793, <https://doi.org/10.1029/2024GL108793>.