

Effects of caffeic acid and bovine serum albumin in reducing the rate of development of rancidity in oil-in-water and water-in-oil emulsions

Article

Accepted Version

Conde, E., Gordon, M. H., Moure, A. and Dominguez, H. (2011) Effects of caffeic acid and bovine serum albumin in reducing the rate of development of rancidity in oil-in-water and water-in-oil emulsions. *Food Chemistry*, 129 (4). pp. 1652-1659. ISSN 0308-8146 doi: <https://doi.org/10.1016/j.foodchem.2011.06.027> Available at <https://centaur.reading.ac.uk/22824/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.foodchem.2011.06.027>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

www.reading.ac.uk/centaur

CentAUR

Central Archive at the University of Reading

Reading's research outputs online

1 **Effects of Caffeic Acid and Bovine Serum Albumin in Reducing the Rate of**
2 **Development of Rancidity in Oil-in-water and Water-in-oil Emulsions**

3 ENMA CONDE¹, MICHAEL H.GORDON², ANDRÉS MOURE¹ and HERMINIA DOMINGUEZ¹

4
5 ¹ Departamento de Enxeñería Química, Universidade de Vigo (Campus Ourense), Edificio Politécnico

6 As Lagoas s/n, 32004, Ourense, Spain

7 ² Hugh Sinclair Unit of Human Nutrition, Department of Food and Nutritional Sciences, The
8 University of Reading, Whiteknights P.O. Box 226, Reading RG6 6AP, United Kingdom

9
10
11 AUTHOR EMAIL ADDRESS m.h.gordon@reading.ac.uk

12
13 RECEIVED DATE

14 TITLE RUNNING HEAD Effects of caffeic acid and bovine serum albumin on rancidity
15 development in emulsions

16 CORRESPONDING AUTHOR FOOTNOTE Phone +44 118 3786723 Fax: +44 118 9310 080

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

ABSTRACT

The antioxidant properties of caffeic acid and bovine serum albumin in oil-in-water and water-in-oil emulsions were studied. Caffeic acid (5 mmol/kg emulsion) showed good antioxidant properties in both 30% sunflower oil-in-water (OW) and 20% water-in-sunflower oil emulsions (WO), pH 5.4, during storage at 50 °C. Although bovine serum albumin (BSA) (0.2%) had a slight antioxidant effect, the combination of caffeic acid and BSA showed a synergistic reduction in the rate of development of rancidity, with significant reductions in concentration of total volatiles, peroxide value (PV) and p-anisidine value (PA) for both emulsion types. The synergistic increase in stability of the OW and WO emulsions containing BSA and caffeic acid was 102.9 and 50.4 % respectively based on TOTOX values, which are calculated as $2PV + PA$, with greater synergy calculated if based on formation of headspace volatiles, The OW emulsion was more susceptible to the development of headspace volatiles by oxidation than the WO emulsion, even though the degree of oxidation assessed by the TOTOX value was similar.

Keywords: Antioxidant activity; Bovine serum albumin; Caffeic acid; Emulsions; Oxidative stability

47

48

49 **1. INTRODUCTION**

50 Lipid oxidation is of great concern to the consumer because it causes not only changes in the quality
51 attributes of foods, including taste, texture, shelf life, appearance and flavor, but also due to the strong
52 decrease in the nutritional value and safety caused by the loss of antioxidants and the formation of
53 harmful components including free radicals and reactive aldehydes (Halliwell, Murcia, Chirico &
54 Aruoma, 1995).

55 Oxidation in food emulsions is of particular interest since oil is widely consumed in foods which are
56 either water-in-oil, for example, butter and margarine, or oil-in-water emulsions, for example,
57 mayonnaise, milk and cream. Lipid oxidation in emulsions is generally recognized as being more
58 complex than lipid oxidation in bulk oil systems, as the emulsification process will lead to the
59 formation of a large interfacial area, and lipid oxidation is initiated at the interface between oil and
60 water, where different non-polar and polar compounds in the system can interact (Frankel, 1998;
61 McClements & Decker, 2000).

62 One of the most effective methods of retarding lipid oxidation in fatty foods is to incorporate
63 antioxidants. The behavior of antioxidants is also more complex in emulsions than in bulk oil since
64 more variables are involved, including the emulsifier and pH (Mancuso, McClements & Decker, 1999;
65 Sørensen et al., 2008). Antioxidant efficacy in food systems depends on various factors including the
66 structure (i.e., number and position of hydroxyl groups bound to the aromatic ring and presence of
67 other substituents), chemical reactivity of the phenolics, interactions with food components, and
68 environmental conditions. The partitioning of the antioxidant into the different phases is an important
69 factor for emulsions (Decker, Warner, Richards & Shahidi, 2005). The efficacy of antioxidants in bulk
70 oil and in dispersed systems is affected by their polarity as described by the “polar paradox” (Frankel,
71 Huang, Kanner & German, 1994). According to the polar paradox, polar antioxidants like ascorbic

72 acid and Trolox are more active in non-polar media like bulk oils than less polar components including
73 ascorbyl palmitate and tocopherol. These apparent paradoxical effects of the antioxidants have been
74 reported as being due to the polar antioxidants being located at the air-oil interface in bulk oils, i.e.
75 situated where oxidation is suggested to take place or at least to be initiated. In addition, they may act
76 in non-polar media by chelation of metal ions. On the other hand, lipophilic antioxidants are more
77 active in polar systems, because they are located at the oil-water interface where oxidation is
78 propagated (Frankel et al, 1994; Huang, Hopia, Schwarz, Frankel & German, 1996; Porter, 1993).

79 Phenolic compounds, such as caffeic acid, have received increasing interest due to their antioxidant
80 activity, which improves the stability of lipid-containing foods and their possible beneficial effects on
81 human health. Caffeic acid may exert its antioxidant effect by donating a hydrogen atom to free
82 radicals, thereby acting as chain-breaking antioxidants, or may act as a metal chelator, which reduces
83 the activity of prooxidants (Gülçin, 2006; Leonardis & Macciola, 2003).

84 Although antioxidants have been frequently studied in oils, emulsions, and other foods, there have
85 been few reports of how proteins, which are commonly present in foods, may affect the activity of
86 antioxidants. Most antioxidants of interest for foods have one or more phenolic hydroxyl groups, and
87 several studies have demonstrated that molecules with this structure may bind to proteins. Polyphenols
88 may associate with proteins through hydrophobic interactions and hydrogen bonding (Oda, Kinoshita,
89 Nakayama & Kakehi, 1998), and a range of phenolic antioxidants has also been shown to bind to
90 bovine skin proteins (Wang & Goodman, 1999).

91 Proteins have been shown to have weak antioxidant activity including both metal chelating and radical
92 scavenging activity (Arcan & Yemenicioğlu. 2007). Bovine serum albumin (BSA), a minor whey
93 protein with M.W. 66 kDa has surface-active properties and has been used to stabilize model food
94 emulsions (Rampon, Lethuaut, Mouhous-Riou & Genot, 2001).

95 The aim of this study was to determine the influence of BSA on the total antioxidant activity of caffeic
96 acid in model food emulsions. The pH of the emulsions was 5.4, which is a typical pH for margarine
97 samples.

98

99 **2. MATERIALS AND METHODS**

100 ***2.1. Chemicals***

101 All solvents used were analytical or HPLC grade (Merck & Co. Inc., Darmstadt, Germany). Isooctane,
102 glacial acetic acid, ferrous sulfate, barium chloride, ammonium thiocyanate, polyoxyethylene sorbitan
103 monolaurate (Tween-20), p-anisidine, cumene hydroperoxide, bovine serum albumin (BSA), hexanal,
104 and bromobenzene were purchased from Sigma-Aldrich (Gillingham, Dorset, UK). Refined sunflower
105 oil was purchased from a local retail outlet. Fatty acids were purchased from Sigma Chemical Co.
106 (Saint Louis, MO, USA).

107 The fatty acid composition of the sunflower oil is shown in **Table 1**.

108 ***2.2. Removal of tocopherols from sunflower oil***

109 Tocopherols were removed from sunflower oil by column chromatography using activated alumina as
110 described by Yoshida, 1993.

111 ***2.3. Emulsion preparation***

112 Oil-in-water emulsions (30% oil) were prepared by dissolving Tween-20 (1%) in acetate buffer (0.1
113 M, pH 5.4), either with or without BSA (0.2%) and caffeic acid (5 mmol/kg emulsion). Water-in-oil
114 emulsions (20% water) were prepared by dissolving Dimodan® (1%) in acetate buffer (0.1 M, pH
115 5.4), either with or without BSA (0.2%) and caffeic acid (5 mmol/kg emulsion). Emulsions were
116 prepared by the dropwise addition of oil to the water phase, with sonication by a Vibracell sonicator
117 (Sonic & Materials, Newton, CT, USA) whilst cooling in an ice bath for 5 min.

118 ***2.4. Emulsion oxidation***

119 All emulsions were stored in triplicate in 25 mL glass bottles in the dark (inside an oven) at 50 °C, and
120 each replicate was analysed once.

121 Aliquots of each emulsion were removed periodically for peroxide value (PV), *p*-anisidine value (PA)
122 determinations and for analysis of volatile oxidation products by gas chromatography (GC).

123 ***2.5. Determination of fatty acid composition***

124 The fatty acid composition of sunflower oil was determined by GC on an HP Agilent 6890N
125 chromatograph equipped with a HP 7683 injector and a flame ionization detector (FID). Fatty acid
126 methyl esters (FAMES) were prepared by transesterification according to Shehata, de Man &
127 Alexander, 1970. Chromatographic separation was carried out using a SP-2560 column (100 m length,
128 0.25 mm ID and 0.20 µm film thickness; Supelco). The oven temperature was 120 °C for 5 min,
129 followed by temperature programming to 170 °C at 2 °C min⁻¹, 200 °C at 5 °C min⁻¹, and then
130 increased to 235 °C at 2 °C min⁻¹. Helium was used as carrier gas, the FID temperature was 280 °C and
131 the injection port was held at 260 °C.

132 ***2.6. Spectrophotometric determination of peroxide value (PV)***

133 Emulsion (0.3 mL) was added to isooctane/2-propanol (3:2 v/v, 1.5 mL) and the mixture was mixed
134 on a vortex mixer three times for 10 s each time. After centrifugation for 2 min at 1000 x g, the clear
135 upper layer (0.2 mL) was collected and peroxides were quantified using a method based on that of
136 Díaz, Dunn, McClements & Decker, 2003. Lipid peroxide concentration was determined using a
137 cumene hydroperoxide standard curve ($r^2=0.9977$).

138 ***2.7. Spectrophotometric determination of p-anisidine Value (PA)***

139 The PA value was determined by AOCS Official Method no. cd 18-90, 1989.

140 ***2.8. Solid phase microextraction (SPME) sampling conditions***

141 An aliquot of emulsion (0.2 g) was weighed in a 2mL vial. A magnetic follower was added and the
142 vial was capped with a Teflon-faced rubber septum and plastic cap. The vial was placed in a water
143 bath on a magnetic stirrer and the sample was equilibrated for 2 min at 60°C. The septum was

144 manually pierced with the SPME needle and the fiber was exposed to the emulsion headspace for 60
145 min and transferred to the gas chromatograph where the volatiles were desorbed in the injection port.
146 The desorption time in the injection port was 15 min.

147 **2.9. SPME/GC analysis**

148 Volatile oxidation compounds were monitored by headspace analysis with solid phase microextraction
149 (HS-SPME). A manual SPME fiber holder unit and 30 μm DVB-CAR-PDMS fiber (Sigma-Aldrich
150 Company Ltd., Dorset, UK) were used to adsorb volatiles from the emulsion in a closed vial at 60°C
151 with a sampling time of 60 min. GC analyses were performed with a HP 5890 series II gas
152 chromatograph (Agilent UK, South Queensferry, UK) equipped with FID detector and split/splitless
153 injector. Chromatographic separation was carried out using a HP-5-column (15 m length, 0.25 mm ID
154 and 0.25 μm film thickness; Agilent UK). The oven temperature was 40 °C for 10 min, followed by
155 temperature programming to 140 °C at 2.5 °C min⁻¹, and then increased to 300 °C at 20 °C min⁻¹.
156 Helium was used as carrier gas in the splitless mode. The FID temperature was 280 °C and the
157 injection port was held at 260 °C.

158 The identification of all compounds was based on the mass spectra determined by GC-MS using a HP
159 5890 series II gas chromatograph with MS detector and by comparison of their retention time with
160 those of authentic standards. Retention times for a series of n-alkanes (C₅-C₂₅) were determined and
161 used to calculate the Linear Retention Indices (LRI values) of detected compounds.

162 Bromobenzene (1 $\mu\text{L/g}$ emulsion) was added as an internal standard prior to analysis. The relative
163 concentrations of the investigated compounds were calculated by relating the area of the internal
164 standard to the area of the compound of interest, defined as:

$$165 \text{ Relative conc.} = \frac{\text{Peak area of particular compound}}{\text{Peak area of IS}} \times \text{IS conc.}$$

166 **2.10. Confocal microscopy**

167 Confocal microscopy was performed with a Leica TCS SP2 AOBS confocal laser scanning
168 microscope mounted on a Leica inverted DM IRE2 microscope for the oil-in-water emulsions and
169 with a Leica TCS SP5 confocal laser microscope for the water-in-oil emulsions.. A Nile Red/Nile Blue
170 mixture (0.01%) was added to an emulsion sample and examined using x10, x20 and x 40 objectives.
171 The image was obtained by exciting the sample with a 633nm Helium-Neon laser with an emission
172 band of 630-750nm.

173

174 **2.11. Calculation of synergy**

175 Synergy was calculated by comparing the times to a given level of deterioration, defined as induction
176 period for the calculation, as described in Almajano & Gordon, 2004.

$$177 \quad \% \text{synergism} = 100 \frac{[IP(a+p) - IP(c)] + [IPa - IPc] + (IPp - IPc)}{[IPa - IPc] + (IPp - IPc)}$$

178 where IP= induction period, a = antioxidant, p = protein and c = control.

179 **2.12. Statistical analysis**

180 The triplicate determinations were used to calculate the mean (m), standard deviation (sd), and
181 standard error of the mean (SEM).

182 Times to selected levels of deterioration were compared, and significant differences at the 5% level
183 were assessed by one-way ANOVA using commercial software (Statistica 6.0). To verify the
184 association among experimental data, Pearson correlation analysis was performed using the same
185 statistical package; p-values < 0.05 were considered significant.

186

187

188 **3. RESULTS AND DISCUSSION**

189 The sunflower oil used contained linoleic acid (64.17%), and oleic acid (25.01%) as the main
190 unsaturated fatty acids, with linolenic acid being present at 0.16% (**Table 1**).

191 The emulsions were stored at 50 °C to accelerate oxidative changes that would occur more slowly, at
192 lower temperatures. Reproducibility of the data was good with the SEM being <25% for 80% of the
193 samples analysed by PV, and the SEM being <25% for 90% of the samples analysed by PA.

194 The initial PV of the emulsion samples was similar with values of 0.05 mM cumene hydroperoxide for
195 OW emulsions and 0.07-0.17 mM for WO emulsions. After 14 days of oxidation the OW emulsions
196 reached significantly different PV values in the order: OW > OW-BSA > OW-CAF > OW-CAF-BSA.

197 The PV values reached in WO emulsions were in the same order up to 35 days but the PV of WO-
198 CAF was higher than WO-BSA at 42 days, since the PV of the control and BSA samples declined
199 sharply after 35 days with corresponding sharp increases in PA values indicating degradation of the
200 hydroperoxides.

201 As shown in **Figure 1**, the maximum PV reached was 10.43 mM hydroperoxide at 42 days for the OW
202 sample, followed by 7.72 mM hydroperoxide for OW-BSA. In WO emulsions the maximum value
203 was reached at 35 days for the WO sample followed by the WO-BSA sample (6.28 and 5.44 mM
204 hydroperoxide, respectively); then there was a decrease in PV for all WO emulsions as the rate of
205 hydroperoxide formation became less than hydroperoxide decomposition.

206 At 35 days of storage, the extent of formation of hydroperoxides in the OW-CAF and WO-CAF
207 samples was lower than in the appropriate control sample with inhibition of 66.7 and 34.7 %,
208 respectively, indicating the antioxidant effect of caffeic acid. The greater effect in the OW emulsion
209 may be due to type of emulsion, the higher caffeic acid:oil ratio (16.7 mmol/kg oil) in the OW
210 emulsion, which was 2.7 times higher than in the WO emulsion (6.25 mmol/kg oil) and the difference
211 between oil concentration in the emulsions. BSA increased the activity of caffeic acid in both
212 emulsions.

213

214 The PA value determination, a measure of secondary oxidation products (List, Evans, Kwolek,
215 Warner, Boundy & Cowan, 1974), confirmed the findings of the PV analysis. The PA test was used to
216 determine the level of aldehydes, principally 2-alkenals and 2,4-alkadienals, present in the emulsions.
217 The PA values were relatively stable for the samples with caffeic acid, and with both caffeic acid and
218 BSA throughout the study.

219 As shown in **Table 2**, in the presence of BSA, the stability of the OW emulsions containing caffeic
220 acid increased by more than 200%, when assessed by the time to PA = 0.15, but BSA did not
221 contribute any significant increase in stability to WO emulsions containing caffeic acid when assessed
222 by the PA values. The TOTOX value is a measure of both primary and secondary oxidation products,
223 where $TOTOX = 2PV + PA$. Synergy between the caffeic acid and bovine serum albumin, in retarding
224 oxidation of emulsions, was investigated by consideration of the TOTOX values. As shown in **Table**
225 **2**, the OW sample took 8.19 days to reach a TOTOX value of 3.7, followed by OW-BSA with 10.72
226 days, then OW-CAF with 23.7 days and finally OW-CAF-BSA, the most stable, with 42 days. The
227 WO emulsions showed similar times to reach a TOTOX value of 8.9.

228

229 Headspace solid phase microextraction (HS-SPME) was used to isolate the headspace volatile
230 products formed during secondary oxidation of OW and WO emulsions. Volatile aldehydes have great
231 importance as oxidation products because of their contribution to the aroma of oxidised oils and
232 emulsions. Hydrophobic flavor components can be perceived at lower concentrations in water than in
233 oil, since many of the lipid oxidation products have higher solubility in the oil phase. This means that,
234 for a fixed concentration of volatile components, their concentration in the headspace of an emulsion
235 decreases as the oil concentration increases. As a consequence, a low fat emulsion may be perceived
236 as more oxidized than a high fat emulsion, even though both emulsions have the same concentration of
237 volatile components.

238 **Figure 2** shows the change in concentration of hexanal and total volatile products with time expressed
239 as relative peak area. Hexanal has been widely used as a marker of volatile oxidation products of n-6
240 fatty acids in previous studies (Rouseff & Cadwallader, 2001). In both systems, hexanal was the main
241 volatile formed during storage. The OW emulsion was more susceptible to headspace volatile
242 formation than the WO emulsion, as shown by the considerably higher concentrations of total volatiles
243 that were formed, even though the degree of oxidation assessed by the TOTOX value was not
244 significantly different in the two emulsion types (**Table 3**). At day 42, the TOTOX value for the OW
245 emulsion containing caffeic acid and BSA was even lower than that of the analogous WO emulsion.
246 The difference in headspace volatile concentration is partly due to the reduction of vapour pressure of
247 each volatile component in the greater mass of oil present in the WO emulsions, since, according to
248 Raoult's law, the vapour pressure of a solute is proportional to its molar concentration. However,
249 differences in the relative importance of hydroperoxide decomposition pathways into volatile and non-
250 volatile products may also occur in the different emulsion types, since the headspace volatile: PA ratio
251 after 42 days storage was 6.7-7.9 times higher in the OW emulsion than in the corresponding WO
252 emulsion, for samples containing no additive, BSA or caffeic acid and 49.3 times higher for the
253 samples containing caffeic acid and BSA (Table 3). The higher oxidative susceptibility of OW
254 emulsions has been reported extensively, and it is attributed to the greater extent of interfacial
255 interactions that are possible between the lipid substrate and prooxidants including metal ions in the
256 aqueous phase as a consequence of the greater surface area presented by the emulsion droplets
257 (McClements & Decker, 2000; Jacobsen, Let, Nielsen & Meyer, 2008). It has been suggested that lipid
258 oxidation in WO emulsions will occur at a rate similar to that in bulk oils because the surface of the
259 lipid phase is exposed directly to air (McClements & Decker, 2000), but effects of antioxidants, metal
260 ions and chelating agents will differ in these media. In WO emulsions, caffeic acid inhibited the
261 formation of hexanal and total volatile compounds at the end of the storage period by 75.48 and 53.51
262 %, respectively. Caffeic acid showed good antioxidant properties in OW emulsions inhibiting the
263 formation of hexanal and total volatile compounds by 95.08 and 87.27 %, respectively. BSA increased

264 the antioxidant activity of caffeic acid in inhibiting the formation of hexanal and volatile compounds
265 in the later stages of storage in WO emulsions (by 18.49 and 26.49 %, respectively), but was less
266 effective in OW emulsions (2.51 and 3.32 %, respectively).

267 The synergistic increase in stability of the OW and WO emulsions prepared with BSA containing
268 caffeic acid was 277.34 and 468.53 %, respectively, based on the time for the total volatile
269 concentration to reach 585.37 and 364.56 μg bromobenzene/ g oil, respectively (**Table 2**).

270 Positive correlations between PV and total volatiles ($r=0.892$) and also between PA and both hexanal
271 and total volatiles ($r=0.992$, $r=0.981$, respectively) in OW emulsions were found. Weaker correlations
272 among parameters analyzed were found in WO emulsions (**Table 4**). Faster hydroperoxide
273 decomposition compared to hydroperoxide formation in the WO emulsions, which leads to a reduction
274 in PV but an increase in total volatiles, contributes to the weaker correlation between PV and total
275 volatiles in this emulsion system.

276 In the secondary oxidation stage, volatile compounds (e.g. alcohols and aldehydes) are formed by the
277 decomposition of lipid hydroperoxides. In particular, volatile aldehydes have a great importance as an
278 indicator of oxidation due to their considerable contribution to the aroma and flavor deterioration of
279 the final products. The concentration of volatile compounds identified in OW and WO emulsions at 21
280 and 42 days of oxidation is summarized in **Table 5**. The profile of volatiles was dominated by
281 oxidation products derived from linoleic acid. As confirmed by GC-MS analysis, hexanal, (*E*)-2-
282 heptenal, (*E*)-2-octenal and (*E,E*)-2,4-decadienal were the major volatile products in the autoxidation,
283 while pentanal, 1-octen-3-ol, nonanal and (*E*)-dec-2-enal were also present in significant amounts.
284 Hexanal, (*E*)-2-octenal, 2-heptenal, pentanal, and (*E,E*)-2,4-decadienal are major oxidation products of
285 linoleic acid (Selke, Rohwedder & Dutton, 1980; Frankel, 1998; Aidos, Jacobsen, Jensen, Luten, Van
286 der Padt & Boom, 2002; Jiménez-Álvarez, Giuffrida, Golay, Cotting, Destailats, Dionisi et al, 2008),
287 whereas nonanal, octanal, 1-heptanol, 1-octanol, and (*E*)-dec-2-enal are formed from triacylglycerols
288 containing oleic acid (Selke et al, 1980).

289 The combination of caffeic acid and BSA was most effective at reducing the formation of both
290 hexanal and total volatiles in both emulsion systems. Caffeic acid was very effective at reducing
291 hexanal and total volatile formation in the OW emulsion, but its effect was weaker in the WO
292 emulsion where it was no better than BSA in reducing total volatile formation up to 35 days. This is
293 consistent with the increased hydroperoxide decomposition observed in the WO emulsion containing
294 caffeic acid (Figure 1b). The relative concentrations of the volatiles changed to some extent during the
295 oxidation with E-2-octenal becoming a major volatile product at later stages of oxidation in the OW
296 emulsion.

297 The antioxidant power and other important biological properties of caffeic acid are well substantiated
298 (Gülçin, 2006; Yanishlieva & Marinova, 1995; Chen & Ho, 1997). Caffeic acid retarded the formation
299 of hydroperoxides in menhaden oil-in-water emulsions (Maqsood & Benjakul, 2010), and De
300 Leonardis & Macciola (De Leonardis & Macciola, 2003) reported that the antioxidant effectiveness of
301 caffeic acid was better than that of BHA in hydrophobic phases such as cod liver oil.

302 The interactions between polyphenols and albumin have attracted much attention due to their
303 importance in food (He, Liang, Wang & Luo, , 2010; Prigent, Voragen, Visser, Van Koningsveld &
304 Gruppen, 2007). Caffeic acid is known to bind to BSA during storage in air with antioxidant activity
305 being incorporated in the protein fraction (Bartolomé, Estrella & Hernández, 2000; Almajano &
306 Gordon, 2004; Almajano, Carbo, Delgado & Gordon, 2007) but this does not occur in samples stored
307 under nitrogen (unpublished data). Confocal microscopy showed that BSA was present at the interface
308 between the oil and water phases in both an oil-in-water emulsion (Figure 3a, b), and a water-in-oil
309 emulsion (c, d). BSA can be seen clearly since it enhances the Nile blue fluorescence at 670 nm (Lee,
310 Suh & Li, 2003). Hence, it can be deduced that caffeic acid is oxidized to a quinone, and binds to BSA
311 by reaction with amine groups in the side chains of amino acids such as lysine, as described by Rawel,
312 Rohn, Kruse & Kroll (2002). This transfers the caffeic acid from homogeneous solution in the aqueous
313 phase to the interface where it is more effective as an antioxidant. The caffeic acid retains a free
314 phenolic group, which is activated by a positively charged nitrogen atom as shown in scheme 1.

315

316 CONCLUSIONS

317 Development of headspace volatiles by lipid oxidation was significantly affected by emulsion type.
318 The oil-in-water emulsions had a higher headspace volatile concentration, despite the degree of total
319 oxidation assessed by the TOTOX value being similar to that of the water-in-oil emulsions. Although
320 BSA had a slight antioxidant effect, it was found that BSA exerted a synergistic effect with caffeic
321 acid in both emulsion types, and this is consistent with previous reports of synergy accompanied by
322 formation of a protein-antioxidant adduct during storage (Almajano & Gordon, 2004). BSA caused a
323 greater synergistic increase in the antioxidant activity of caffeic acid in a water-in-oil emulsion than in
324 an oil-in-water emulsion.

325

326 ACKNOWLEDGMENTS

327 EC thanks the Spanish Ministry of Education (AP2007-02164) for funding. We thank Andrew Dodson
328 and Stephen Pountney for technical assistance.

329 REFERENCES

- 330 Aidos, I.; Jacobsen, C., Jensen, B., Luten, J. B., Van der Padt, A., & Boom, R. M. (2002). Volatile
331 oxidation products formed in crude herring oil under accelerated oxidative conditions. *Eur. Food Res.*
332 *Technol.*, *104*, 808–818.
- 333 Almajano, M. P., Carbo, R., Delgado, M. E., & Gordon, M. H. (2007). Effect of pH on the
334 antimicrobial activity and oxidative stability of oil-in-water emulsions containing caffeic acid. *J. Food*
335 *Sci.*, *72*, 258-263.
- 336 Almajano, M. P., & Gordon, M. H. (2004). Synergistic effect of BSA on antioxidant activities in model
337 food emulsions. *J. Am. Oil Chem. Soc.*, *81*, 275-280.
- 338 AOCS Official Method cd 18-90. (1989). In *Official methods and recommended practices of the*
339 *American Oil Chemists' Society*, 4th Ed., Firestone, D. Ed., American Oil Chemists' Society:
340 Champaign, IL.
- 341 Arcan, I. & Yemenicioğlu, A. (2007). Antioxidant activity of protein extracts from heat-treated or
342 thermally processed chickpeas and white beans. *Food Chem.*, *103*, 301–312.
- 343 Bartolomé, B., Estrella, I., & Hernández, M. T. (2000). Interaction of low molecular weight phenolics
344 with proteins (BSA). *J. Food Sci.*, *65*, 617-621.

- 345 Chen, J. H., & Ho, C. T. (1997). Antioxidant activities of caffeic acid and its related hydroxycinnamic
346 acid compounds. *J. Agric. Food Chem.*, *45*, 2374–2378.
- 347 Decker, E. A., Warner, K., Richards, M. P., & Shahidi, F. (2005). Measuring antioxidant effectiveness
348 in food. *J. Agric. Food Chem.* *53*, 4303–4310.
- 349 De Leonardis A., & Macciola V. (2003). Effectiveness of caffeic acid as an anti-oxidant for cod liver
350 oil. *Int. J. Food Sci. Technol.*, *38*, 475–480.
- 351 Díaz, M., Dunn, C., McClements, D. J., & Decker, E. A. (2003). Use of caseinophosphopeptides as
352 natural antioxidants in oil-in-water emulsions. *J. Agric. Food Chem.*, *51*, 2365-2370.
- 353 Frankel, E. N. (1998). Lipid Oxidation. The Oily Press Ltd.: West Ferry, Dundee, Scotland.
- 354 Frankel, E. N., Huang, S.-W., Kanner, J., & German, J. B. (1994). Interfacial phenomena in the
355 evaluation of antioxidants: bulk oils vs emulsions. *J. Agric. Food Chem.*, *42*, 1054-1059.
- 356 Gülçin, I. (2006). Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicol.*, *217*,
357 213–220.
- 358 Halliwell, B., Murcia, M. A., Chirico, S., & Aruoma, O. I. (1995). Free radicals and antioxidants in
359 food and in vivo: what they do and how they work. *Crit. Rev. Food Sci. Nutr.* *35*, 7-20.
- 360 He, T., Liang, Q., Wang, Y., & Luo, G. (2010). Characterization of the interactions between natural
361 flavonoid compounds and bovine serum albumin by capillary electrophoresis and fluorescence
362 method. *J. Liq. Chromatogr. Rel. Technol.*, *33*, 548-562.
- 363 Huang, S.-W., Hopia A., Schwarz K., Frankel E. N., & German J. B. (1996). Antioxidant activity of a
364 tocopherol and trolox in different lipid substrates: bulk oils vs oil-in-water emulsions. *J. Agric. Food
365 Chem.*, *44*, 444-452.
- 366 Jacobsen, C., Let, M. B., Nielsen, N. S., & Meyer, A. S. (2008). Antioxidant strategies for preventing
367 oxidative flavour deterioration of foods enriched with n-3 polyunsaturated lipids: a comparative
368 evaluation. *Trends in Food Sci. & Technol.*, *19*, 76-93
- 369 Jiménez-Álvarez, D., Giuffrida, F., Golay, P. A., Cotting, C., Destailats, F., Dionisi, F., et al.; (2008).
370 Profiles of volatile compounds in milk containing fish oil analyzed by HS-SPMEGC/ MS. *Eur. J.
371 Lipid Sci. Technol.*, *110*, 277–283.
- 372 Lee, S.H., Suh, J.K., & Li, M. (2003). Determination of bovine serum albumin by its enhancement
373 effect of Nile blue fluorescence. *B. Kor. Chem.Soc.*, *24* (1), 45-48.
- 374 Leonardis, A. D., & Macciola, V. (2003). Effectiveness of caffeic acid as an antioxidant for cod liver
375 oil. *Int. J. Food Sci. Technol.*, *38*, 475–480.
- 376 List, G. R., Evans, C. D., Kwolek, W. F., Warner, K., Boundy, B. K., & Cowan, J. C. (1974).
377 Oxidation and quality of soybean oil: A preliminary study of the anisidine test. *J. Am. Oil Chem. Soc.*,
378 *51*, 17–21.
- 379 Mancuso, J. R., McClements, D. J., & Decker, E. A. (1999). The effects of surfactant type, pH, and
380 chelators on the oxidation of salmon oil-in- water emulsions. *J. Agric. Food Chem.* *47*, 4112–4116.
- 381 Maqsood, S., & Benjakul, S. (2010). Comparative studies of four different phenolic compounds on in
382 vitro antioxidative activity and the preventive effect on lipid oxidation of fish oil emulsion and fish
383 mince. *Food Chem.*, *119*, 123-132.
- 384 McClements, D. J., & Decker, E. A. (2000). Lipid oxidation in oil-in-water emulsions: impact of
385 molecular environment on chemical reactions in heterogeneous food systems. *J. Food Sci.* *65*, 1270-
386 1282.
- 387 Oda Y. M., Kinoshita, K., Nakayama, K., & Kakehi, K. (1998). Evaluation of fluorescence polarization
388 method for binding study in carbohydrate–lectin interaction. *Bio. Pharm. Bulletin.*, *21*, 1215–1217.

- 389 Porter, W. L. (1993). Paradoxical behavior of antioxidants in food and biological systems. *Toxicol. Ind.*
390 *Health.*, 9, 93-122.
- 391 Prigent, S. V. E., Voragen, A. G. J., Visser, A. J. W. G., Van Koningsveld, G. A., & Gruppen, H.
392 (2007). Covalent interactions between proteins and oxidation products of caffeoylquinic acid
393 (chlorogenic acid). *J. Sci. Food Agric.*, 87, 2502-2510.
- 394 Rampon, V., Lethuaut, L., Mouhous-Riou, N., & Genot, C. (2001). Interface characterization and
395 aging of bovine serum albumin stabilized oil-in-water emulsions as revealed by front-surface
396 fluorescence. *J. Agric. Food Chem.*, 49, 4046–4051.
- 397 Rawel, H.M., Rohn, S., Kruse, H.P., & Kroll, J. (2002). Structural changes induced in bovine serum
398 albumin by covalent attachment of chlorogenic acid. *Food Chem.*, 78 (4), 443-445.
- 399 Rouseff, R. L., & Cadwallader, K. R. (2001). Headspace volatile aldehydes as indicators of lipid
400 oxidation in food. In *Headspace techniques in foods and flavours*; Kluwer Academic/Plenum
401 Publishers: New York.
- 402 Selke, E., Rohwedder, W. K., & Dutton, H. J. (1980). Volatile components from trilinolein heated in
403 air, *J. Am. Oil Chem. Soc.*, 57, 25-30.
- 404 Shehata, A. J., de Man, J. M., & Alexander, J. C. (1970). A simple and rapid method for the
405 preparation of methyl esters of fats in milligram amounts for gas chromatography. *Can. Inst. Food Sci.*
406 *Technol. J.*, 3, 85-89.
- 407 Sørensen, A. M., Haahr, A., Becker, E. M., Skibsted, L. H., Bergenståhl, B., Nilsson, L. et al. (2008).
408 Interactions between iron, phenolic compounds, emulsifiers, and pH in omega-3-enriched oil-in-water
409 emulsions. *J. Agric. Food Chem.* 56, 1740-1750.
- 410 Wang, W.Q., & Goodman, M. T. (1999). Antioxidant property of dietary phenolic agents in a human
411 LDL oxidation ex vivo model: Interaction of protein binding activity, *Nutr. Res.*, 19, 191–202.
- 412 Yanishlieva, N., & Marinova, E. M. (1995). Effects of antioxidants on the stability of triacylglycerols
413 and methyl esters of fatty acids of sunflower oil. *Food Chem.*, 54, 377–382.
- 414 Yoshida, H. (1993). Influence of fatty acids of different unsaturation in the oxidation of purified
415 vegetable-oils during microwave irradiation. *J. Sci. Food Agric.*, 62, 41–47.

416 **Table 1.** Fatty acid composition of sunflower oil

Fatty acid name		Amount (%)
<i>Saturated (S)</i>		10.42
Myristic acid	C14:0	0.11 ± 0.02
Palmitic acid	C16:0	5.72 ± 0.04
Margaric acid	C17:0	0.05 ± 0.00
Stearic acid	C18:0	3.49 ± 0.05
Arachidic acid	C20:0	0.24 ± 0.00
Behenic acid	C22:0	0.57 ± 0.01
Tricosanoic acid	C23:0	0.05 ± 0.00
Lignoceric acid	C24:0	0.19 ± 0.01
<i>Unsaturated (U)</i>		89.58
<i>Monounsaturated</i>		25.25
Palmitoleic acid	C16:1	0.10 ± 0.00
Oleic acid	C18:1(n-9)	25.01 ± 0.63
Eicosenoic acid	C20:1(n-9)	0.14 ± 0.00
<i>Polyunsaturated</i>		64.33
Linoleic acid	C18:2(n-6)	64.17 ± 0.73
Linolenic acid	C18:3(n-3)	0.16 ± 0.00
<i>S : U</i>		0.12
<i>Omega 6</i>		64.17
<i>Omega 3</i>		0.16
<i>Omega 6 : Omega 3</i>		401.06

417

418

419

420

421

422

423

424

425

426 **Table 2.** Times in days for oil-in-water and water-in-oil emulsions to reach indicated values at 50 °C

427

<i>Oil-in-water emulsion (value at assessment time)</i>	PV (1.79 mM hydroperoxide)	PA (0.15)	TOTOX (3.7)	Hexanal (14.63 µg bromobenzene / g oil)	Total volatiles (585.37 µg bromobenzene / g oil)
OW	9.95	2.65	8.19	2.36	7.54
OW-BSA	11.22	8.84	10.72	6.12	10.99
OW-CAF	24.91	9.41	23.7	22.49	13.22
OW-CAF-BSA	42.00	42.00	42.00	42.00	42.00
% Synergy	97.47	202.60	102.9	65.95	277.34
<i>Water-in-oil emulsion (value at assessment time)</i>	PV (2.15 mM hydroperoxide)	PA (4.59)	TOTOX (9.5)	Hexanal (18.71 µg bromobenzene / g oil)	Total volatiles (364.56 µg bromobenzene / g oil)
WO	12.87	17.34	16.2	3.95	13.42
WO-BSA	15.82	17.26	17.7	13.77	15.84
WO-CAF	24.47	39.47	27.2	17.70	16.02
WO-CAF-BSA	42.00	42.0	35	42.00	42.00
% Synergy	100.20	12.10	50.4	61.49	468.53

428

429

430

431

432

433

434

435

436

437

438

439

440

441 **Table 3.** PV, PA, TOTOX and relative concentration of total volatiles for emulsions at 42 days

	PV (mM hydroperoxide)	PA	TOTOX	Total volatiles (μg bromobenzene / g oil)	Total volatiles: PA ratio
OW	10.43 \pm 1.97	17.29 \pm 1.68	38.15	6219.58 \pm 999.53	359.72
OW-BSA	7.72 \pm 1.76	11.77 \pm 1.94	27.21	5477.40 \pm 675.16	465.37
OW-CAF	4.82 \pm 0.03	0.71 \pm 0.12	10.35	791.53 \pm 48.42	1114.83
OW-CAF-BSA	1.79 \pm 0.32	0.15 \pm 0.04	3.73	585.40 \pm 48.15	3902.67
WO	3.97 \pm 0.33	34.03 \pm 0.65	41.97	1822.68 \pm 279.83	53.56
WO-BSA	2.87 \pm 0.60	23.49 \pm 2.11	29.23	1379.19 \pm 8.32	58.71
WO-CAF	3.42 \pm 0.34	5.11 \pm 0.08	11.95	847.29 \pm 95.04	165.81
WO-CAF-BSA	2.15 \pm 0.24	4.59 \pm 0.91	8.89	364.56 \pm 14.21	79.42

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457 **Table 4.** Significant Pearson's correlations among parameters analyzed (PV, PA, hexanal and total
458 volatiles)

Pearson's correlation:	<i>Oil-in-water emulsion</i>				<i>Water-in-oil emulsion</i>			
	PV	PA	Hexanal	Total volatiles	PV	PA	Hexanal	Total volatiles
PV	-	0.856	0.860	0.892	-	0.504	0.712	0.796
PA	0.856	-	0.992	0.981	0.504	-	0.904	0.885
Hexanal	0.860	0.992	-	0.987	0.712	0.904	-	0.922
Total volatiles	0.892	0.981	0.987	-	0.796	0.885	0.922	-

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

476 **Table 5.** Concentration (expressed as equivalent to bromobenzene) of volatile compounds identified in
 477 oil-in-water and water-in-oil emulsions during storage at 50 °C

Compound	Concentration (μg bromobenzene / g oil)			
	OW (21 days)	OW (42 days)	WO (21 days)	WO (42 days)
<i>Butanal</i>	20.97	46.74	6.36	24.05
<i>Pentanal</i>	15.19	71.97	14.69	45.71
<i>(E)-2-Pentenal</i>	18.39	141.37	nd	nd
<i>1-Pentanol</i>	5.91	88.85	3.37	28.61
<i>Hexanal</i>	70.66	606.9	61.7	310.26
<i>2-Hexenal</i>	nd	20.44	0.83	5.43
<i>2-Heptanone</i>	nd	10.95	1.25	7.38
<i>Heptanal</i>	nd	60.42	1.75	14.36
<i>(E)-2-Heptenal</i>	17.54	259.35	12.96	53.07
<i>1-Heptanol</i>	2.12	11.12	1.72	3.61
<i>1-Octen-3-ol</i>	5.93	117.9	3.64	15.92
<i>2-Pentyl-furan</i>	nd	nd	9.13	34.28
<i>Octanal</i>	8.36	65.37	2.18	16.81
<i>3-Octen-2-one</i>	2.35	28.01	1.53	14.08
<i>5-Ethylidihydro-2(3H)-furanone</i>	nd	nd	13.39	116.33
<i>(E)-2-Octenal</i>	15.08	550.12	2.89	28.28
<i>1-Octanol</i>	1.54	14.92	1.37	4.04
<i>Heptanoic acid</i>	2.99	10.37	0.87	7.23
<i>2-Nonanone</i>	5.18	43.57	4.07	20.49
<i>Nonanal</i>	10.84	26.5	3.93	13.61
<i>(E,E)-2,4-Octadienal</i>	0.07	5.81	nd	nd
<i>3-Nonen-2-one</i>	0.55	15.27	0.47	2.27
<i>(E)-2-Nonenal</i>	1.69	18.45	2.48	13.36
<i>2-Decanone</i>	0.91	13.77	1.07	2.67
<i>(E,E)-2,4-Nonadienal</i>	3.02	67.91	3.1	15.89
<i>3-Methylbut-2-enal</i>	11.63	85.71	3.02	9.98
<i>Methyl-cyclodecane</i>	1.28	7.55	0.55	2.07
<i>(E)-2-Decenal</i>	7.11	70.53	3.71	14.04
<i>Nonanoic acid</i>	1.35	32.75	1.11	7.45
<i>(E,E)-2,4-Decadienal</i>	70.46	443.56	19.32	60.09
<i>5-pentyl-5(H)-furan-2-one</i>	6.25	229.94	3.79	40.06
<i>(E)-2-Tridecenal</i>	7.18	75.28	3.48	11.95
<i>6-Dodecanone</i>	1.16	34.52	1.06	3.1
<i>2-Butyloct-2-enal</i>	8.14	100.88	2.51	7.71
<i>Total volatiles</i>	323.85	3376.8	193.3	954.19

480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502

FIGURE LEGENDS





Figure 1. Changes in peroxide value and *p*-anisidine value of a) OW emulsions and b) WO emulsions containing antioxidants during storage at 50 °C.

Figure 2. Changes in hexanal and total volatiles concentration of a) OW emulsions and b) WO emulsions containing antioxidants during storage at 50 °C, with concentration expressed as area equivalent to bromobenzene.

Figure 3. Oil-in-water emulsions stained with oil red- Nile blue mixture (a) with BSA; (b) no protein

A) Oil-in-water emulsion

B) Water-in-oil emulsion



—□— OW —■— OW-BSA ···△··· OW-CAF ···▲··· OW-CAF-BSA

—□— WO —■— WO-BSA ···△··· WO-CAF ···▲··· WO-CAF-BSA

503

504

505

506

507


508 **Figure 1.** Changes in peroxide value and *p*-anisidine value of a) OW emulsions and b) WO emulsions

509 containing antioxidants during storage at 50 °C

510

511

512
513
514
515
516
517
518




519 **Figure 2.** Changes in hexanal and total volatiles concentration of a) OW emulsions and b) WO
520 emulsions containing antioxidants during storage at 50 °C, with concentration expressed as area
521 equivalent to bromobenzene.


522

523


524




(a)



(b)



(c)



(d)


525

526 **Figure 3.** Emulsions stained with oil red- Nile blue mixture (a) Oil-in-water emulsion with BSA; (b)
527 Oil-in-water emulsion with no protein; (c) water-in-oil emulsion with BSA; (d) water-in-oil emulsion
528 with no protein

529

530

531



532

533

534 **Scheme 1.** Reaction of caffeic acid with lysine residue of BSA (P = protein backbone)