

# *The domestic garden: its contribution to urban green infrastructure*

Article

Accepted Version

Cameron, R., Blanusa, T., Taylor, J., Salisbury, A., Halstead, A., Henricot, B. and Thompson, K. (2012) The domestic garden: its contribution to urban green infrastructure. *Urban Forestry and Urban Greening*, 11 (2). pp. 129-137. ISSN 1618-8667 doi: 10.1016/j.ufug.2012.01.002 Available at <https://centaur.reading.ac.uk/26212/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1016/j.ufug.2012.01.002>

Publisher: Elsevier

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

# **The Domestic Garden – Its Contribution to Urban Green Infrastructure**

Ross W.F. CAMERON<sup>1</sup>, Tijana BLANUŠA<sup>2, 3, \*</sup>, Jane E. TAYLOR<sup>3</sup>, Andrew SALISBURY<sup>2</sup>,  
Andrew J. HALSTEAD<sup>2</sup>, Béatrice HENRICOT<sup>2</sup> and Ken THOMPSON<sup>4</sup>

<sup>1</sup>Department of Landscape, University of Sheffield, Sheffield S10 2TN, UK

<sup>2</sup>Plant Sciences Department, Royal Horticultural Society, Garden Wisley, Woking GU23  
6QB, UK

<sup>3</sup>School of Biological Sciences, University of Reading, Whiteknights, Reading RG6 6AS, UK

<sup>4</sup>Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

\*Author for correspondence: [tijanablanusa@rhs.org.uk](mailto:tijanablanusa@rhs.org.uk), Telephone: +44-(0)118-378-6467

## **Abstract**

Domestic gardens provide a significant component of urban green infrastructure but their relative contribution to eco-system service provision remains largely un-quantified. ‘Green infrastructure’ itself is often ill-defined, posing problems for planners to ascertain what types of green infrastructure provide greatest benefit and under what circumstances. Within this context the relative merits of gardens are unclear; however, at a time of greater urbanization where private gardens are increasingly seen as a ‘luxury’, it is important to define their role precisely. Hence, the nature of this review is to interpret existing information pertaining to gardens /gardening *per se*, identify where they may have a unique role to play and to highlight where further research is warranted. The review suggests that there are significant differences in both form and management of domestic gardens which radically influence the benefits. Nevertheless, gardens can play a strong role in improving the environmental impact of the domestic curtilage, e.g. by insulating houses against temperature extremes they can reduce domestic energy use. Gardens also improve localized air cooling, help mitigate flooding and provide a haven for wildlife. Less favourable aspects include contributions of gardens and gardening to greenhouse gas emissions, misuse of fertilizers and pesticides, and introduction of alien plant species. Due to the close proximity to the home and hence accessibility for many, possibly the greatest benefit of the domestic garden is on human health and well-being, but further work is required to define this clearly within the wider context of green infrastructure.

**Keywords:**

carbon footprint; ecosystem services; green space; private garden; well-being.

## **1. Introduction**

Recent research has highlighted the positive role urban green infrastructure can provide in terms of ecosystem services (de Groot et al., 2002; Tratalos et al., 2007) and health agendas (Tzoulas et al., 2007). Yet, urban green infrastructure (parks, public green space, allotments, green corridors, street trees, urban forests, roof and vertical greening, private gardens) covers a range of landscape types of varying complexity and morphology. Do all hold equal value? Domestic gardens are quoted as an important component of green infrastructure (e.g. Loram et al., 2007), but their specific contribution is rarely assessed, and hence their relative value within the wider urban green space is difficult to quantify. For example, in terms of green space provision for the future should urban planners be prioritising private gardens around building stock, or rather, invest in more open, communal green spaces? Therefore, the aims of this review are to examine the evidence relating private domestic gardens to ecosystem service provision, to understand how form, function and management of gardens may influence the extent of that service, and to identify where further research is required. The relationship between the domestic garden and the four broad strands of ecosystem service function (provisioning, regulating, cultural and supporting; MA, 2003) are incorporated, although much of the literature focuses mainly focuses on the first three of these. We anticipate that the existing data on gardens highlighted here, in addition to new research initiatives, will help to provide a greater precision to existing models relating green infrastructure to human and ecosystem 'health' (Tzoulas et al., 2007).

## **2. What is an urban domestic garden?**

For the purpose of this review we define the private domestic garden as the area adjacent to a domestic dwelling, which itself is either privately owned or rented. A key

element is that the resident/s have autonomy over the garden, albeit they may wish to delegate responsibility to others (professional designer, hired gardener etc.). We attempt to differentiate and exclude, 'open' green space and communal gardens, for example where management may be by committee, or at the discretion of a housing association or local authority. Conversely, locations that have been 'guerrilla gardened', i.e. public areas that have been illegally planted by individuals are also omitted, despite these often reflecting individual autonomy, self-expression etc. We also exclude public parks and gardens, although their design may in many ways reflect those of private domestic gardens. Similarly, allotment sites where local bylaws may influence what may / may not be grown also fall outside our definition. Lastly, the emphasis has been on urban gardens and their role, rather than encompass large rural gardens where the dividing line between garden and extensive landscaped estate can be difficult to distinguish.

In highly urbanised societies up to 90% of the population reside in urban areas, and in cultures such as the UK 87% of households have access to a domestic garden (Gibbons et al., 2011). Depending on age and location of cities, domestic gardens contribute between 22 and 36% of the total urban area (Gaston et al., 2005; Mathieu et al., 2007), and perhaps as much as 3-4% of total land mass (Alloway, 2004; Gibbons et al., 2011). Similarly, they constitute a significant proportion of the urban green space, more than 50% in Dunedin, New Zealand, (Mathieu et al., 2007) and 35- 47% in the UK (Loram et al., 2007). Urbanization, however, is decreasing the proportion of area dedicated to gardens (Mathieu et al., 2007; Smith, 2010), through infill development, i.e. existing gardens being sold for development, or newer housing stock having smaller gardens.

Gardens are highly heterogeneous in form and function. At one extreme, they may comprise a few square meters of multi-layered diverse vegetation to, at the other, large areas of single dimension paving with no vegetation at all. Loram et al. (2007; 2008) reported that

domestic garden size varied between 3.6 m<sup>2</sup> and 2290 m<sup>2</sup> within UK cities, with median garden size between 96 m<sup>2</sup> and 213 m<sup>2</sup>. Housing type and density influences the proportion of green space available (Whitford et al., 2001). Although these authors did not differentiate between private gardens and other local green space, namely communal lawns, street trees / roadside verges or vegetated pedestrian pathways the data are illustrative of how housing age and type can influence presence and access to local green space. The highest proportion of surrounding green space was associated with 1930's era semi-detached houses (57%), intermediate with the modern low rise flats (30%) and lowest with the modern terraced housing (13%) (Whitford et al., 2001). Grass was the dominant green cover but the percentages varied (32.6%, 14.4% and 6.1%, respectively); trees covered 11%, 0.3%, and 0.9% respectively. Within domestic gardens lawns constitute 60% (UK, Gaston et al., 2005) or 55% (New Zealand, Mathieu et al., 2007) of the area. In the USA, lawns (private, public and sports turf) cover between 8 and 16 million ha, far surpassing land coverage of major crops, (e.g. cotton, Robbins et al., 2001).

Greater housing density is linked to smaller garden sizes, but not necessarily less urban green space (Smith et al., 2009). Socio-economic variables (e.g. population and housing density, education, home ownership) were apparently better predictors of the extent and the type of vegetation cover in private gardens than solely biophysical variables (e.g. rainfall, soil fertility, solar radiation etc., Luck et al., 2009). Larger domestic gardens (Smith et al., 2005), those associated with older properties (Hope et al., 2003), or with higher income or tertiary-educated residents tend to have proportionally more vegetation, greater diversity of plants, and more complex garden styles (Daniels and Kirkpatrick, 2006).

Front and back gardens often differ in character, with visual impact more important in front gardens, with the back used for functional purposes; this is, however, probably an oversimplification and an outdated concept (Daniels and Kirkpatrick, 2006). For example,



UK trends over the last decade suggest that vegetation and aesthetic character have been lost from the front garden, due to off-road parking (Perry and Nawaz, 2008). Irrespective of location, vegetation is being replaced in favour of 'low maintenance' hard standings such as patios or decked areas (Smith, 2010).

### **3. Garden vegetation, temperature regulation and energy conservation**

The relationship between green infrastructure and the mitigation of the urban heat island effect is currently under close scrutiny (Takebayashi and Moriyama, 2007; Alexandri and Jones, 2008; Huang et al., 2008), as climate change models highlight the increased likelihood of heat waves (Meehl and Tebaldi, 2004). Modelling approaches suggest that increasing the proportion of green space within the urban matrix, can reduce surface, and hence air temperatures. Gill et al. (2007) suggest a 10% increase in urban green infrastructure would negate the 4°C increase predicted for Manchester, UK over the next 80 years. Type of urban green infrastructure and even plant species (Blanusa et al., unpublished), however, are likely to affect this cooling potential.

Urban trees are considered a major factor in providing cooling, *via* shade and evapo-transpiration, and are estimated to offer over 950 MJ (almost 270 kWh) cooling per day, per tree, due to evapo-transpiration effects alone (Huang et al., 1990). The extent to which gardens contribute to cooling is unclear. Domestic gardens account for 25% of the non-woodland trees in the urban matrix (Davies et al., 2009). Akbari et al. (1997) predicted that up to a quarter of the cooling effect by urban trees in USA cities may relate to garden/street trees contributing to direct cooling of the adjacent buildings. This will vary depending on the tree size, species, maturity, and architecture (Simpson, 2002), but more detailed research is

needed to understand which type of buildings benefit from which trees, and to determine specific requirements for a range of scenarios.

Reducing the energy load on buildings by shading and evaporative cooling can reduce the reliance on / improve efficiency of, mechanical air conditioning units. US data suggest that strategic positioning of plants could reduce domestic buildings' energy consumption by 20-40% (Akbari et al., 1997; 2001; Huang et al., 1990). However, it was also suggested that greater savings may be accomplished if highly reflective building surfaces were used instead (Akbari et al., 2001); it is of interest therefore, to evaluate plants with a particularly high albedo within this context.

Despite the potential for urban vegetation to provide cooling, effectiveness relates to soil water availability. As climate change events reduce summer precipitation (for Europe) and heat waves exacerbate soil moisture deficits (IPCC, 2007), cooling through evapotranspiration is likely to be compromised. For example, the cooling potential of turf is strongly linked to watering (McPherson et al., 1989). Hence, greater scrutiny is required in future research to elucidate the relative cooling associated with shade, albedo and evapotranspiration.

Beyond trees, a wider range of vegetation has potential to improve cooling, e.g. through planting next to, or on, a building. Sufficiently robust data are still lacking for green facades, especially as different mechanisms are used to irrigate them. Recent work with turf applied as vertical greening, however, showed that this reduced interior surface temperatures by  $> 2^{\circ}\text{C}$  (Cheng et al., 2010). Green roofs also reduce interior temperatures through improved insulation via the substrate as well as the potential for evaporative cooling (Santamouris et al., 2007; Sailor, 2008). There is a suggestion that the strategies to conserve energy in temperate climates cannot be generalized, due to the "complex interactive effects of shade and wind reduction" (McPherson et al., 1988). Indeed, Bowler et al. (2010) point that

there are insufficient data to provide specific recommendations. At the garden level, vegetation can influence the energy loads on individual buildings, but how this impacts on air temperatures across the wider urban environment is unclear. There also remain unresolved issues with respect to the scale on which urban climate parameters and vegetation are measured and how they are compared (Stewart, 2011).

In terms of winter temperature regulation and energy savings there is much less information available, although preliminary studies are encouraging (e.g. 17% saving in energy, Liu and Harris, 2008). Vegetation around houses enhances winter energy saving by reducing the velocity of air over buildings ('wind break'), removing draughts and minimizing temperature differences between existing and incoming air (McPherson et al., 1988). Careful planting design is paramount to ensure that wind tunnels are not directed towards the house, and that maximum solar gain is retained in winter (Knowles, 2003).

#### **4. Carbon balance**

Little information specifically identifies the role of the domestic garden in carbon dynamics, but the effectiveness of garden plants and soils for carbon storage will strongly depend on design (types of vegetation and coverage, see Figure 1), management, and the longevity of plantings. An average of  $2.5 \times 10^3 \text{ g m}^{-2}$  of carbon is stored in domestic gardens with 83% in soil (to 600mm depth), 16% in trees and shrubs and only 0.6% on average in grass and herbaceous plants (Jo and McPherson, 1995). Indeed, there may be as much carbon stored within the top 100mm of soil profile alone, as in the entire vegetation biomass above ground (e.g. compare data for forests and forest soils, Figure 1). Soil carbon storage, however, is reduced by soil disturbance during replanting as organic matter is oxidized (Bolinder et al., 2007). Gardens that mimic forest or low maintenance meadow landscapes, or

follow practices aligned with permaculture principles e.g. minimum soil cultivation, low chemical use, heterogeneous vegetation, use of recycled water and organic matter are likely to have the least impact in terms of CO<sub>2</sub> release (Pouyat et al., 2002; Favoino and Hogg, 2008). These suggestions, however, require testing in the domestic garden context. Even practices designed to aid carbon sequestration, such as tree planting, may have surprisingly long pay-back times. Due to the embedded energy associated with tree production and the volume of soil CO<sub>2</sub> released during planting, newly planted trees have been estimated to become 'carbon neutral' only after 3-10 years (Nowak and Crane, 2002; Schlosser et al., 2003).

Gardening is often promoted as an environmentally-friendly pastime, but this notion may be questioned when the intensity of garden activities are considered. Today's gardeners rely greatly on mechanized tools, synthetic fertilizers and pesticides (with high embodied energy, see section 5). In addition, potable water use has an energy load associated with its cleaning and distribution, and use of 'hard' landscaping materials bring their own embedded carbon, as well as the energy in growing and retail of nursery plants.

Perhaps the lawn provides the greatest illustration of the diverging attitudes to gardening in this respect with, at one extreme, naturalized grassland (meadows) having a low environmental footprint, in contrast to a heavily-maintained sward (Qian and Follett, 2002; Trudgill et al., 2010). The former requires a single annual cut, in contrast to high frequency mowing associated with formal lawns (e.g. weekly during optimum growth). Although accurate estimations of carbon use through mowing are difficult to attain (Reid et al., 2010), Jo and McPherson (1995) suggest petrol-powered mowing releases 1.5-fold the carbon sequestered by the lawn itself. Two stroke engines particularly are considered detrimental with a 7-fold higher by-product emission compared to four stroke engines (Volckens et al., 2008). Intensively-managed lawns are also associated with high fertilizer and herbicide inputs

(and in warmer, affluent countries, high water use – see section 7). Despite this, where urbanisation results in housing development on land previously degraded or heavily cultivated, the creation of a lawn can help replace soil organic carbon originally lost through oxidation (Zirkle et al., 2011).

Longer-lived woody plants can provide greater carbon sequestration than annual plants due to reduced tillage and storage in biomass. Comparison between five shrub species indicated carbon storage in biomass ranges from  $4.7\text{--}7.2 \times 10^6$  g carbon ha<sup>-1</sup>, and atmospheric carbon uptake ranges from  $0.56\text{--}0.80 \times 10^6$  g carbon ha<sup>-1</sup> year<sup>-1</sup> (Jo, 2002). Total carbon sequestered in urban trees in the USA is estimated to be  $700 \times 10^{12}$  g (Nowak and Crane, 2002). Assuming that the ratio of garden to street / parkland trees in the USA is similar to that of the UK (i.e. 25%), this may equate to  $175 \times 10^{12}$  g of carbon sequestered in urban garden trees. This could be an over-estimation as garden trees are often smaller than parkland specimens. Caution is required, however, as these data do not include carbon emissions associated with tree maintenance which can lead to a negative life-time carbon balance (Nowak et al., 2002); carbon sequestration is usually modelled on a small unit basis (ha) (Brainard et al., 2006) and involves numerous assumptions and uncertainties when scaled up.

While there is a significant volume of carbon in garden trees, it is minor compared to the carbon stored in the world's forests ( $359 \times 10^{15}$  g carbon, Dixon et al., 1994). This also needs to be viewed in light of total amount of excess carbon that needs to be sequestered annually to mitigate climate change,  $1.8 \pm 1.4 \times 10^{15}$  g CO<sub>2</sub>e (Dixon et al., 1994) [CO<sub>2</sub>e being the amount of all greenhouse gases, when converted into equivalent units of carbon dioxide]. Therefore, although increasing the number of urban trees provides a range of environmental benefits, justifying new plantings solely to offset global deforestation, or fossil fuel emissions, seems irresponsible.

## 5. Garden chemicals

Chemicals, primarily fertilizers and pesticides, have widespread use (e.g. 50-74% of all homeowners in the UK and USA respectively, Robbins et al., 2001; Grey et al., 2006). Their application, particularly on lawns, is apparently linked to home property values, with more affluent residents using them more frequently (Robbins et al., 2001). In the USA, local by-laws encourage the use of garden chemicals and promote intensive management (e.g. lawns regularly mown and weed free, Robbins et al., 2001; Clayton, 2007).

The production/use of artificial *fertilizers* contribute significantly to greenhouse gas emissions (Howarth et al., 2002), with impact being particularly high from nitrogen fertilizer production based on the Haber-Bosch process. In contrast, manufacture of phosphate and potassium fertilisers is 20-fold less energy intensive. Composted organic matter offers a lower carbon cost alternative for supplementing nitrogen than using an inorganic nitrogen fertiliser  $\text{NH}_4\text{NO}_3$  (Lillywhite and Rahn, 2008). Lawn fertilizer use in particular affects greenhouse gas emissions (Livesley et al., 2010) with lawns emitting up to 10 times more  $\text{N}_2\text{O}$  than neighbouring agricultural grassland. The higher emissions are thought to be due to the higher irrigation rates and soil temperatures found in urban lawns (Bijoor et al., 2008).

Garden *pesticides* are significant contributors to USA's non-point source water pollution (Robbins et al., 2001). Surface run-off, although not specifically quantified for domestic gardens, can include chemicals as well as particulate matter that can contribute to the pollution of watercourses (Overmyer et al., 2005). Parts of the gardening community are, therefore, encouraging lower chemical use and organic approaches, largely due to concerns over environmental issues. For example, the use of organic composts (green waste and spent mushroom) to support pest (e.g. aphid) control by encouraging predatory species (e.g. spiders

and beetles) has shown some success (Bell et al., 2008). Furthermore, green spaces themselves can reduce pollution through capture of particulates and dissolved pollutants (Davis et al., 2001), although effectiveness will depend on the properties of the pollutant and the magnitude of benefits is yet to be assessed in the domestic context.

## **6. Run-off and flooding**

Gardens provide storm attenuation ‘services’ to the urban matrix. Vegetation, trees especially, intercept intense precipitation, hold water temporarily within their canopy thus reducing peak flow and easing demand on urban drains (Xiao and McPherson, 2002). In addition, vegetation mitigates flood risk by increasing infiltration into the soil reducing surface flow (Dunne et al., 1991). Despite these advantages, hard paving in domestic gardens is increasing. Perry and Nawaz (2008) found a 13% increase in impervious surfaces over 30 years in the city of Leeds UK, 75% of which was due to paving of residential front gardens; this was linked to more frequent and severe flooding in the area. Pauleit and Duhme (2000) found that lower density housing with gardens had 3-fold less storm water run-off than higher density stock. The benefits of vegetated front gardens have been recognised by the UK planning system; from 2008 permission is required for impermeable paving (Anon, 2009).

## **7. Water use**

Gardens are frequently associated with heavy water use, particularly during periods of dry weather. The proportion of potable water used for the domestic garden escalates with increased aridity. In western Australia, 56% of domestic water use is associated with domestic gardens (Syme et al., 2004). Similar values were found for Barcelona, Spain; on average, 30% of household water use is consumed within the garden, and this rises to 50%

during the summer (Domene and Sauri, 2006). Climate change will increase demand for garden water in temperate regions, although surprisingly few studies have investigated details of this. Mulching flower beds minimises moisture evaporation and has the added benefit of limiting gaseous emissions (Livesley et al., 2010). In lawns, organic vs intensive style of turf management can reduce water consumption up to 10-fold (Morris and Bagby, 2008).

Water use in the domestic garden needs to be systematically assessed. Despite moves to increase efficiency of garden water use (e.g. seep hose irrigation), greater attention to water storage (e.g. water butts) and grey water use, it is likely, that potable water use in the garden will become increasingly restricted. This is partly due to increased water demand, e.g. greater housing density, climate change, and the carbon associated with mains water supply (Strutt et al., 2008).

## **8. Peat**

The use of peat in gardens, and horticulture in general, is controversial due to habitat destruction and carbon emissions linked with peat extraction (Alexander et al., 2008), despite the fact that the majority of peat harvested is consumed by power stations (Alanne and Saari, 2006). Peat used as a soil conditioner and as potting medium for the English amateur garden market was  $2.03 \times 10^6 \text{ m}^3$  in 2009 (Anon, 2010a), a consumption rate that has led the UK's government to propose stopping garden peat use by 2020 (Anon, 2010a).

## **9. Air quality**

Although urban trees are often cited as removing aerial pollutants (e.g. Yang et al., 2005; Nowak et al., 2006; Jim and Chen, 2008) and garden trees (and other vegetative forms) no doubt contribute to capturing overall pollutant loads, recent research suggests that



relatively wide belts of woodlands are required to elicit measurable benefits (Pataki et al., 2011). Domestic gardens therefore may have only a minor role in helping mitigate point pollutant sources such as roads, or industrial outlets. Some plants are also considered to be strong emitters of biogenic volatile organic compounds, implying their role in contributing to the formation of photochemical smog in the urban environment (Niinemets and Peñuelas, 2008; Peñuelas and Staudt, 2010). On balance, there is a high level of uncertainty about the role of the urban vegetation in air pollutant removal (Pataki et al., 2011); this is the case even more so for domestic gardens where there is no direct evidence available.

## **10. Biodiversity**

Fifty years ago there was a widespread belief that gardens, highly managed and dominated by alien plants, provided few resources for native animals (Elton, 1966). However, as published reports of wildlife in domestic gardens began to accumulate, it became clear that the value of urban gardens for biodiversity may be substantial (Davis, 1978; Vickery, 1995). Indeed, some declining species, once common in low-intensity farmland, are now more abundant in urban areas, and particularly in domestic gardens (Gregory and Baillie, 1998). For some bird species, urban gardens support a significant fraction of the total UK population providing valuable habitat and supporting some nationally important sub-sets of species (Chamberlain et al., 2009). In a comprehensive study, Owen (2010) revealed that her rather ordinary garden in Leicester (UK) was visited over a 30-yr period by around a quarter of all the insect species recorded from the entire country. By focusing for just three years on one family of parasitoid wasps (*Ichneumonidae*), Owen discovered 15 species new to the UK and four previously undescribed species. Gardens are such a good habitat for hoverflies (*Syrphidae*) that here they greatly outnumber their supposed wasp mimics (Azmeah et al.,

1998). The BUGS project which surveyed 61 domestic gardens in Sheffield, UK (Smith et al., 2006a; Thompson, 2007), confirmed Owen's conclusion that private gardens, taken together, might reasonably be described as the UK's most important nature reserve. The BUGS project also demonstrated that neither small size nor isolation from countryside seem to be a problem; small, city-centre gardens support much the same invertebrate wildlife as large, suburban ones. Other recent research has tended to endorse this view; in Manhattan, New York, gardens with sunny, flower-rich patches supported diverse pollinator communities (Matteson and Langellotto, 2010), and in Toronto, small 'microcosms' (soil-filled pots, with or without vegetation) introduced into gardens recruited plants, seeds and invertebrates in much the same way as those placed in grassland or forest (Sperling and Lortie, 2010).

The widespread perception of 50 years ago that gardens are of little wildlife value was clearly mistaken, but what *sort* of wildlife is supported by gardens? Gardens seem to be used by only a small proportion of mammals, although this minority may be very frequent garden visitors (Baker and Harris, 2007). Nor are the invertebrates that frequent gardens in large numbers a random selection. Among butterflies, whose ecology can be predicted with some certainty from the biology of their larval host plants (Dennis et al., 2004), it is clear that garden butterflies are largely generalist, mobile species, and that specialist, sedentary species are only rarely able to make use of urban gardens (Bergerot et al., 2010; Owen, 2010). Similarly, gardens sustain very large populations of a small group of ubiquitous, generalist bumblebees, while rarer specialists are confined to less-disturbed semi-natural habitats (Goulson et al., 2006).

Pessimists point out that the rather generalist, mobile and adaptable nature of garden wildlife means it is the fraction most likely to survive anyway, with or without the assistance of gardens and gardeners. Conversely, even these adaptable species are not always able to survive in the most intensively farmed landscapes: see, for example, the extraordinarily 'poor

bee', fauna of Dutch arable land (Kleijn et al., 2001), and the evidence that pollinators in general are increasingly beleaguered in agricultural landscapes outside gardens (Cussans et al., 2010). While gardens will never replace species-rich semi-natural habitats, they are nevertheless a useful complement to such habitats and, moreover, increasingly provide urban residents with their only close encounters with the natural world.

## **11. Plant composition and pest species**

Gardeners have always been drawn to exotic and unusual plants, and even today plant hunters search for unusual genotypes that can be introduced into the garden trade. This results in a large proportion of non-native plant species, selected genotypes (varieties) or accidental or intentionally bred hybrids (Smith et al., 2006b) in the garden; increasingly many of these 'exotic' garden plants are traded internationally (Smith et al., 2007). Historically, the garden trade has introduced a wide range of alien plant species (e.g. Reichard and White, 2001), a number are now invasive weeds, e.g. *Acacia spp.* in South Africa (Kull et al., 2007), *Clematis vitalba* in Australia and New Zealand (Williams and West, 2000), resulting in huge costs associated with eradication, e.g. over 34 bn US \$ in the USA (Olson, 2006). Increasing international trade in plants also introduces non-native pest and pathogen species (Henricot and Gorton, 2005; Tubby and Webber, 2010). In the UK, the ornamental trade can account for 90% of human-assisted introductions of plant pests (Smith et al., 2007) and 53% of new plant pathogens have been recorded on ornamental plants, compared to only 15% on native species (Jones and Baker, 2007).

The high diversity of plant species in gardens may also encourage invertebrate or pathogen species to jump host. This is supported by evidence that plants uncommon in the wild, but frequently cultivated, may be more susceptible to invertebrate herbivores in

gardens; for example, at least nine species of moth (*Lepidoptera*) larvae have been recorded feeding on *Potentilla fruticosa* in a UK urban garden (Owen, 2010) but in the wild no moth species had been recorded exploiting this plant (Thompson et al., 2003).

Climate change and associated impacts on the urban environment, e.g. heat islands, seem to be encouraging species survival and migration. Some pests that normally require the protected conditions of a glasshouse can now survive outdoors, while other pests may become more prevalent. *Icerya purchasi* (cottony cushion scale) has been long known as an occasional pest in UK glasshouses (Green, 1931), but it is now a problem on outdoor plants (Watson and Malumphy, 2004). Heat island and allied abiotic stresses have also been linked to a higher incidence of a pest species, such as clearwing moth, *Podesia syringae*, on *Fraxinus pennsylvatica*; providing evidence that elevated urban temperatures can alter species composition and pest abundance (Cregg and Dix, 2001).

## **12. Gardens, health and well-being**

Gardening as a pastime is often seen as a form of ‘retreat’, but also relates to ownership, identity and the ability to interact with nature (Clayton, 2007; Gross and Lane, 2007). Therefore, the gardens influence on an individual’s well-being may relate in part to that individual’s attitude to gardening. The health (and allied social) benefits associated with engagement with the green space are now well documented (e.g. Hartig et al., 2003; St Leger, 2003; van den Berg et al., 2010) and largely centring on the attention restoration theory (ART) (Kaplan, 1995), or the psychophysiological stress recovery theory (PSRT) (Ulrich et al., 1991). Cited benefits are wide ranging and include improvements in pain relief, blood pressure, heart rate, less frequent illness, improved cognitive function (Hartig et al., 2003; Tzoulas et al., 2007). In contrast, other studies suggest that generic links between health and

green infrastructure are weak (Lee and Maheswaran, 2011), or the response is dependent on factors such as sample population studied (Ottoosson and Grahn, 2005), the form, extent or quality of the green infrastructure (Milligan and Bingley, 2007; Mitchell and Popham, 2007), the nature of the 'green' activity undertaken (Barton and Pretty, 2010) or ancillary factors such as ease of access, degree of motivation or perceptions of safety (Lee and Maheswaran, 2011).

Some studies on green infrastructure include domestic gardens / gardeners within the experimental methodology employed, but many do not. Indeed, some that attempt to relate health and proximity to green space, actually exclude the private garden, as the resolution of the GIS systems employed fail to adequately account for them (Maas et al., 2008). So where do gardens and gardening fit in terms of the generic relationship between green space and health? Those studies that primarily focus on gardening (or activities that are directly analogous) highlight the benefits such as reduced mortality, higher bone density, lower blood pressure and cholesterol levels (Walsh et al., 2001; Milligan et al., 2004). Gardening can help reduce the risk of the onset of dementia (Simons et al., 2006) and aid those suffering from it (Lee and Kim, 2008). To maximize health benefits, Magnus et al. (1979) claim that consistent, moderate exercise throughout the year is advantageous; hence regular 'all season' gardening should be encouraged. Regular physical activity can reduce risk of coronary heart disease, ischemic stroke, Type 2 diabetes, hypertension, osteoporosis, anxiety, depression and certain types of cancer (Park et al., 2009). Gardening is one of a number of pastimes that are linked with encouraging greater physical activity, including sustaining long-term engagement through opportunities for creativity, communication and self-expression (Blair et al., 1991). The intensity of physical activity, however, will vary with type of gardening activity, age and ability of participant (Dallosso et al., 1988).

Despite the body of evidence supporting gardening as a healthy and restorative pastime, it is not risk free. Over a 30 day period, Powell et al. (1998) suggested that 2.1 million USA citizens (1.1% of the population) could suffer injury in the domestic garden. Gardening activities have been blamed for increased incidence of injury through misuse of tools (especially lawn mowers, van Duijne et al., 2008), dermatitis (McMullen and Gawkrödger, 2006), and certain pathogens (O'Connor et al., 2007). Although these negative factors can be substantial, they appear to be out-weighed by the advantages cited in relation to increased physical activity and perceived restorative effects.

### **13. Domestic gardens in a cultural context**

Although community gardens, particularly city orchards or allotments have been linked to improving social capital (Glover et al., 2005) and large 'landscape' or 'heritage' gardens are imbued with historical and cultural significance (e.g. Turner, 2011), the role of the urban private garden is more ambiguous. The domestic garden can be seen as an extension to the household (Alexander, 2002), and therefore a forum for family (and friends) to interact and spend leisure time. Positive memories of childhood are often linked with the family garden (Gross and Lane, 2007) and provide strong place attachment (Brook, 2003). One of the possible causes for an increase in gardening within western societies in recent years is that it may be seen as an anti-dote to anxieties and perceived risks associated with changes in lifestyle, including the development of technology, globalization and wider environmental degradation (Bhatti and Church, 2004). This relates back to the garden being a place where an individual can exert control, where perhaps this ability has been lost in other societal institutions and the political system. Domestic gardens at the interface between the home and public space are also considered to be places of real ambiguity, with a range of

conflicting paradoxes, e.g. between consumerism and environmentalism (Bhatti and Church, 2001; 2004; Longhurst, 2006). In contrast to community gardens, the design of some private gardens may act intentionally as a barrier to the wider environment and society (Alexander, 2002). The tall, stone boundary walls and deep hedges placed at the perimeter of 18<sup>th</sup> Century Victorian villa gardens (UK) are perhaps epitomising a highly class-conscious society.

The practical uses that a domestic garden is put to, may also strongly reflect culture or historical context. During World War Two, increased numbers of domestic gardens in the UK were used for food production, a trend that has repeated itself during subsequent periods of economic austerity, when food prices have risen (Lang, 1999, Carney, 2011). Also recent migrants from rural to urban environments may see the domestic garden primarily as a land resource for providing food and not as an aesthetic feature *per se* (Head et al., 2004). Indeed, Risbeth (2005) argues that attitudes to domestic gardens can be strongly cultural, with white, middle class residents in the UK appreciating the privacy and security offered by a private garden, whereas “Bengali and Afro-Caribbean participants play out their scenes in cafés, kitchens, streets and parks – never their back gardens”. Although attitudes to gardens *per se* may vary, an interesting juxtaposition is that one of the key elements within them, i.e. flowers, have universal appeal and can be a unifying factor across cultures (Risbeth, 2005).

#### **14. Gardens as therapeutic landscapes**

Increasingly, the broad principles relating health and green space are being acknowledged by policy makers e.g. World Health Organisation (Edwards and Tsouros, 2006) Greenspace Scotland (Croucher et al., 2007) and the Faculty of Public Health, UK (Anon 2010b) but practitioners are this still left with the dilemma as to what sort of landscapes and facilities should be provided? Do all green landscapes and activities hold

similar health benefits? Part of the problem is that green space has often been ill-defined in terms of specific components, scale or extent. Perhaps the infinite variability in natural landscapes may explain some of the inconsistencies in the data linking green space to well-being. Essentially, there is an argument that not all forms of green space act in the same way (Lee and Maheswaran, 2011), with corresponding implications for the role of domestic gardens and their design (Ottosson and Grahn, 2005; Ivarsson and Hagerhall, 2008). Despite conceptual models being developed to relate green infrastructure to health (e.g. Tzoulas et al., 2007), these too are dependent on the details associated with each type of green space being present to fully test the models, and help define relative benefits.

In terms of ‘attention restoration’ (stress relief) gardens are thought to facilitate some, or all, of the components identified with the ART (Ivarsson and Hagerhall, 2008). They may be particularly important in the prevention and recovery from illness, due to their general accessibility within an urban matrix (providing the opportunity for exercise and engagement with nature, quite literally, at many people’s ‘back door’). Surprisingly few studies though have focused on the private domestic garden and gardeners (Gross and Lane, 2007). In contrast, most reported studies relate to community gardens, or the value of gardens / gardening for those with defined health-related problems (Milligan et al., 2004; Park et al., 2009). In reality, the majority of those who garden may do it in relative isolation or in small groups, and may not be consciously doing so from a health benefit perspective. Therefore more research is required to categorize the benefits and drawbacks for the ‘average gardener’ and ideally, within the specific context of the domestic garden.

In their study on the restorative value of different landscapes, Ivarsson and Hagerhall (2008) indicated that landscapes with a high level of natural features, rather than e.g. urban elements, provided greatest restoration. Even within natural features hierarchy was evident (Herzog et al., 2003). Flowers and water had a higher correlation with quality of life traits



than animals, trees, hills, natural aromas or sounds (Ogunseitan, 2005). Interestingly, Barton and Pretty (2010) too found enhanced health benefits when 'green exercise' was performed within sight of water. If certain natural features are more beneficial than others from a health perspective, then those gardens that incorporate them may enhance the positive psychological responses (Stigsdotter and Grahn, 2002). Even for individuals not suffering health problems, design of garden may be important in offsetting stress associated with work, commuting, family life etc., but this warrants further investigation. It is also true though, that health *per se* may not be at the forefront of most gardeners objectives; aesthetic desires and functional criteria will also determine garden design (Chen et al., 2009). Furthermore, domestic gardeners usually have a large element of control over the design and management of gardens, and this can be linked with other positive psychological experiences including self-esteem, feeling of achievement and fulfilment of talent and skill.

## **15. Conclusions**

Domestic gardens are undoubtedly an important component in many people's lives, but attitudes towards them are not uniform. To some, they are an essential element of life providing opportunity for engagement with nature, self-actualisation, creativity or well-being; to others, they are at best a parking lot, or worse, represent an additional chore to an already busy lifestyle.

The environmental benefits of gardens appear mixed. Domestic gardens help provide improved thermal comfort to their residents, have potential to reduce domestic energy consumption and minimize storm run-off. They also clearly provide much needed habitat for wildlife within the urban environment, but management issues may result in gardens contributing to greenhouse gas emissions rather than offsetting them. Their role in sustainable

use of resources may veer to the negative side of the equation (use of water, pesticides, fertilizers, etc.); in some countries garden activities have been directly responsible for the introduction of invasive species, with huge consequences for native biodiversity and the economics associated with eradication measures. This report highlights a number of areas where future research should be targeted, but such future studies need to embrace the fact that domestic gardens, as with the wider green space, are variable, and the effects / influences of contrasting garden styles need to be incorporated into any methodology.

Perhaps the gardens' greatest benefit (to humans at least) is the potential to provide a forum for exercise, and to facilitate respite from stress. As outlined, green spaces appear to provide a range of benefits linked with health and well-being, but more specific information is required about whether and where the domestic garden fits in with some of these broad conclusions and to challenge some of the generic assumptions that all green spaces act in the same way. This is important, as western society faces the challenges of a more sedentary lifestyle, greater incidences of obesity and increasing occurrence of mental health problems. If domestic gardens do indeed provide a fundamental role in promoting physical exercise and well-being, then they are ideally placed to address these issues directly due to their relative accessibility for much of the population. This has implications for the planning process, as currently many new housing developments omit private gardens and prefer to incorporate communal green areas, or only provide minimal private garden space. Conversely, there may be little point in every house having a garden if their potential is not realised. Educational processes may be required to encourage engagement, especially as society becomes increasingly displaced from the natural world and food cultivation, but evidence is required of the clear benefits before such activity could be undertaken.

## **Acknowledgements**

We are grateful to the Plant for Life Initiative (sponsored by the European Union and the Horticultural Trades Association, UK) and the Royal Horticultural Society UK for supporting this study.

## **References**

- Akbari H, Kurn DM, Bretz SE, Hanford JW. 1997. Peak power and cooling energy savings of shade trees. *Energy and Buildings* 25, 139-148.
- Akbari H, Pomerantz M, Taha H. 2001. Cool surfaces and shade trees to reduce energy use and improve air quality in urban areas. *Solar Energy* 70, 295-310.
- Alanne K, Saari A. 2006. Distributed energy generation and sustainable development. *Renewable and Sustainable Energy Reviews* 10, 539-558.
- Alexander C. 2002. The garden as occasional domestic space. *Signs* 27, 857-871.
- Alexander PD, Bragg NC, Meade R, Padelopoulos G, Watts O. 2008. Peat in horticulture and conservation: the UK response to a changing world. *Mires and Peat* 3, 1-10.
- Alexandri E, Jones P. 2008. Temperature decreases in an urban canyon due to green walls and green roofs in diverse climates. *Building and Environment* 43, 480-493.
- Alloway BJ. 2004. Contamination of soils in domestic gardens and allotments: a brief overview. *Land Contamination & Reclamation* 12, 179-187.
- Anon. 2009. Guidance on the permeable surfacing of front gardens. Department for Communities and Local Government, London, UK. p 1-28.

- Anon. 2010a. Monitoring of the horticultural use of peat and progress towards the UK biodiversity Action Plan target. In: Department for Environment Farming and Rural Affairs, London, UK. p 1-19.
- Anon. 2010b. Great Outdoors: How our natural health service uses green space to improve wellbeing - An action report. Faculty of Health, London, UK. p 1-18.
- Azmeh S, Owen J, Sorensen K, Grewcock D, Gilbert F. 1998. Mimicry profiles are affected by human-induced habitat changes. *Proceedings of the Royal Society of London Series B-Biological Sciences* 265, 2285-2290.
- Baker PJ, Harris S. 2007. Urban mammals: What does the future hold? An analysis of the factors affecting patterns of use of residential gardens in Great Britain. *Mammal Review* 37, 297-315.
- Barton J, Pretty J. 2010. What is the best dose of nature and green exercise for improving mental health? A multi-study analysis. *Environmental Science and Technology* 44, 3947-3955.
- Bell JR, Traugott M, Sunderland KD, Skirvin DJ, Mead A, Kravar-Garde L, Reynolds K, Fenlon JS, Symondson WOC. 2008. Beneficial links for the control of aphids: the effects of compost applications on predators and prey. *Journal of Applied Ecology* 45, 1266-1273.
- Bergerot B, Fontaine B, Renard M, Cadi A, Julliard R. 2010. Preferences for exotic flowers do not promote urban life in butterflies. *Landscape and Urban Planning* 96, 98-107.
- Bhatti M, Church A. 2001. Cultivating natures: Homes and gardens in late modernity. *Sociology-the Journal of the British Sociological Association* 35, 365-383.
- Bhatti M, Church A. 2004. Home, the culture of nature and meanings of gardens in late modernity. *Housing Studies* 19, 37-51.

- Bijoor NS, Czimeczik CI, Pataki DE, Billings SA. 2008. Effects of temperature and fertilization on nitrogen cycling and community composition of an urban lawn. *Global Change Biology* 14, 2119-2131.
- Blair D, Giesecke CC, Sherman S. 1991. A dietary, social and economic evaluation of the Philadelphia Urban Gardening Project. *Journal of Nutrition Education* 23, 161-167.
- Bolinder MA, Andren O, Katterer T, de Jong R, VandenBygaart AJ, Angers DA, Parent LE, Gregorich EG. 2007. Soil carbon dynamics in Canadian Agricultural Ecoregions: Quantifying climatic influence on soil biological activity. *Agriculture Ecosystems & Environment* 122, 461-470.
- Bowler DE, Buyung-Ali L, Knight TM, Pullin AS. 2010. Urban greening to cool towns and cities: A systematic review of the empirical evidence. *Landscape and Urban Planning* 97, 147-155.
- Brainard J, Lovett A, Bateman I. 2006. Sensitivity analysis in calculating the social value of carbon sequestered in British grown Sitka spruce. *Journal of Forest Economics* 12, 201-228.
- Brook I. 2003. Making Here Like There: Place Attachment, Displacement and the Urge to Garden. *Ethics, Place & Environment* 6, 227-234.
- Carney M. 2011. Compounding crises of economic recession and food insecurity: a comparative study of three low-income communities in Santa Barbara County. *Agriculture and Human Values* DOI 10.1007/s10460-011-9333-y, 1-17.
- Chamberlain DE, Cannon AR, Toms MP, Leech DI, Hatchwell BJ, Gaston KJ. 2009. Avian productivity in urban landscapes: a review and meta-analysis. *Ibis* 151, 1-18.
- Chen B, Adimo OA, Bao Z. 2009. Assessment of aesthetic quality and multiple functions of urban green space from the users' perspective. The case of Hangzhoo Flower Garden, China. *Landscape and Urban Planning* 93, 76-82.

- Cheng CY, Cheung KKS, Chu LM. 2010. Thermal performance of a vegetated cladding system on facade walls. *Building and Environment* 45, 1779-1787.
- Clayton S. 2007. Domesticated nature: Motivations for gardening and perceptions of environmental impact. *Journal of Environmental Psychology* 27, 215-224.
- Cregg BM, Dix ME. 2001. Tree moisture stress and insect damage in urban areas in relation to heat island effects. *Journal of Arboriculture* 27, 8-17.
- Croucher K, Myers L, Bretherton J. 2007. Transforming urban spaces. The links between greenspace and health: A critical literature review. *Greenspace Scotland Research Report*. p. 1-61.
- Cussans J, Goulson D, Sanderson R, Goffe L, Darvill B, Osborne JL. 2010. Two bee-pollinated plant species show higher seed production when grown in gardens compared to arable farmland. *Plos One* 5, e11753.
- Dallosso HM, Morgan K, Bassey EJ, Ebrahim SB, Fentem PH, Arie TH. 1988. Levels of customary physical activity among the old and the very old living at home. *Journal of Epidemiology and Community Health* 42, 121-127.
- Daniels GD, Kirkpatrick JB. 2006. Comparing the characteristics of front and back domestic gardens in Hobart, Tasmania, Australia. *Landscape and Urban Planning* 78, 344-352.
- Davies ZG, Fuller RA, Loram A, Irvine KN, Sims V, Gaston KJ. 2009. A national scale inventory of resource provision for biodiversity within domestic gardens. *Biological Conservation* 142, 761-771.
- Davis AP, Shokouhian M, Sharma H, Minami C. 2001. Laboratory Study of Biological Retention for Urban Stormwater Management. *Water Environment Research* 73, 5-14.
- Davis BNK. 1978. Urbanisation and the diversity of insects. In: Mound LA, Waloff N, editors. *Diversity of Insect Faunas*. Oxford: Blackwell. p 126-138.

- de Groot RS, Wilson MA, Boumans RMJ. 2002. A typology for the classification, description and valuation of ecosystem functions, goods and services. *Ecological Economics* 41, 393-408.
- Dennis RLH, Hodgson JG, Grenyer R, Shreeve TG, Roy DB. 2004. Host plants and butterfly biology. Do host-plant strategies drive butterfly status? *Ecological Entomology* 29, 12-26.
- Dixon RK, Brown S, Houghton RA, Solomon AM, Trexler MC, Wisniewski J. 1994. Carbon pools and flux of global forest ecosystems. *Science* 263, 185-191.
- Domene E, Sauri D. 2006. Urbanisation and Water Consumption: Influencing Factors in the Metropolitan Region of Barcelona. *Urban Studies* 43, 1605-1623.
- Dunne T, Zhang W, Aubry BF. 1991. Effects of rainfall, vegetation, and microtopography on infiltration and runoff. *Water Resources Research* 27, 2271-2285.
- Edwards P, Tsouros A. 2006. The solid facts. Promoting physical activity and active living in urban environments. Copenhagen, Denmark: World Health Organisation. p. 1-55.
- Elton CS. 1966. *The Pattern of Animal Communities*. London: Methuen.
- Euliss Jr NH, Gleason RA, Olness A, McDougal RL, Murkin HR, Robarts RD, Bourbonniere RA, Warner BG. 2006. North American prairie wetlands are important nonforested land-based carbon storage sites. *Science of The Total Environment* 361, 179-188.
- Favoino E, Hogg D. 2008. The potential role of compost in reducing greenhouse gases. *Waste Management Research* 26, 61-39.
- Gaston KJ, Warren PH, Thompson K, Smith RM. 2005. Urban Domestic Gardens (IV): The Extent of the Resource and its Associated Features. *Biodiversity and Conservation* 14, 3327-3349.

- Gibbons S, Mourato S, Resende G. 2011. The amenity value of English nature: a hedonic price approach. London, UK: Spatial Economics Research Centre (SERC), London School of Economics and Political Sciences.
- Gill SE, Handley JF, Ennos AR, Pauleit S. 2007. Adapting cities for climate change: the role of green infrastructure. *Built Environment* 33, 115-133.
- Glover TD, Parry DC, Shinew KJ. 2005. Building relationships, accessing resources: mobilizing social capital in community garden contexts. *Journal of Leisure Research* 37, 450-474.
- Gough CM, Fritz EA. 2009. Management and social indicators of soil carbon storage in a residential ecosystem, Midlothian, VA. *Virginia Journal of Science* 60, 181-194.
- Goulson D, Hanley M, Darvill B, Ellis J. 2006. Biotope associations and the decline of bumblebees (*Bombus spp.*). *Journal of Insect Conservation* 10, 95-103.
- Green EE. 1931. Observations on British *Coccidae*. XIII. *Entomologist's Monthly Magazine* 67, 99-106.
- Gregory RD, Baillie SR. 1998. Large-scale habitat use of some declining British birds. *Journal of Applied Ecology* 35, 785-799.
- Grey CNB, Nieuwenhuijsen MJ, Golding J, Team A. 2006. Use and storage of domestic pesticides in the UK. *Science of the Total Environment* 368, 465-470.
- Gross H, Lane N. 2007. Landscapes of the lifespan: Exploring accounts of own gardens and gardening. *Journal of Environmental Psychology* 27, 225-241.
- Hartig T, Evans GW, Jamner LD, David DS, Garling T. 2003. Tracking restoration in natural and urban field settings. *Journal of Environmental Psychology* 23, 109-123.
- Head L, Muir P, Hampel E. 2004. Australian backyard gardens and the journey of migration. *Geographical Review* 94, 326-347.
- Henricot B, Gorton C. 2005. Plant Pathogens on the move. *Microbiology Today* 32, 100.



- Herzog TR, Maguire CP, Nebel MB. 2003. Assessing the restorative components of environments. *Journal of Environmental Psychology* 23, 159-170.
- Hope D, Gries C, Zhu W, Fagan WF, Redman CL, Grimm NB, Nelson AL, Martin C, Kinzig A. 2003. Socioeconomics drive urban plant diversity. *Proceedings of the National Academy of Sciences USA* 100, 8788-8792.
- Howarth RW, Boyer EW, Pabich WJ, Galloway JN. 2002. Nitrogen use in the United States from 1961–2000 and potential future trends. *AMBIO* 31, 88-96.
- Huang L, Li J, Zhao D, Zhu J. 2008. A fieldwork study on the diurnal changes of urban microclimate in four types of ground cover and urban heat island of Nanjing, China. *Building and Environment* 43, 7-17.
- Huang YJ, Akbari H, Taha AA. 1990. The wind shielding and shading effects of trees on residential heating and cooling requirements; *Proceedings of the Winter meeting of the American Society of Heating, Refrigerating and Air-Conditioning Engineers, Inc: Atlanta, GA (USA)*. p. 22.
- IPCC, 2007: *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Core Writing Team, Pachauri, R.K and Reisinger, A. (eds.)]. IPCC, Geneva, Switzerland, 104 pp.
- Ivarsson CT, Hagerhall CM. 2008. The perceived restorativeness of gardens - Assessing the restorativeness of a mixed built and natural scene type. *Urban Forestry & Urban Greening* 7, 107-118.
- Jim CY, Chen WY. 2008. Assessing the ecosystem service of air pollutant removal by urban trees in Guangzhou (China). *Journal of Environmental Management* 88, 665-676.
- Jo H-K. 2002. Impacts of urban greenspace on offsetting carbon emissions for middle Korea. *Journal of Environmental Management* 64, 115-126.

- Jo H-K, McPherson EG. 1995. Carbon storage and flux in urban residential greenspace. *Journal of Environmental Management* 45, 109-133.
- Jones DR, Baker RHA. 2007. Introductions of non-native plant pathogens into Great Britain, 1970-2004. *Plant Pathology* 56, 891-910.
- Kaplan S. 1995. The restorative benefits of nature - toward an integrative framework. *Journal of Environmental Psychology* 15, 169-182.
- Kleijn D, Berendse F, Smit R, Gilissen N. 2001. Agri-environment schemes do not effectively protect biodiversity in Dutch agricultural landscapes. *Nature* 413, 723-725.
- Knowles RL. 2003. The solar envelope: its meaning for energy and buildings. *Energy and Buildings* 35, 15-25.
- Kull CA, Tassin J, Rambeloarisoa G, Sarrailh J-M. 2007. Invasive Australian acacias on western Indian Ocean islands: a historical and ecological perspective. *African Journal of Ecology* 46, 684-689.
- Lang T. 1999. The complexities of globalization: The UK as a case study of tensions within the food system and the challenge to food policy. *Agriculture and Human Values* 16, 169-185.
- Lee ACK, Maheswaran R. 2011. The health benefits of urban green spaces: a review of the evidence. *Journal of Public Health* 33, 212-222.
- Lee Y, Kim S. 2008. Effects of indoor gardening on sleep, agitation, and cognition in dementia patients—a pilot study. *International Journal of Geriatric Psychiatry* 23, 485-489.
- Lillywhite RD, Rahn CR. 2008. True cost of using fertilizers on the land. *Horticulture week*(17 April), 37-38.
- Liu Y, Harris DJ. 2008. Effects of shelterbelt trees on reducing heating-energy consumption of office buildings in Scotland. *Applied Energy* 85, 115-127.

- Livesley S, Dougherty B, Smith A, Navaud D, Wylie L, Arndt S. 2010. Soil-atmosphere exchange of carbon dioxide, methane and nitrous oxide in urban garden systems: impact of irrigation, fertiliser and mulch. *Urban Ecosystems* 13, 273-293.
- Longhurst R. 2006. Plots, plants and paradoxes: contemporary domestic gardens in Aotearoa New-Zealand. *Social & Cultural Geography* 7, 581-593.
- Loram A, Tratalos J, Warren PH, Gaston KJ. 2007. Urban domestic gardens (X): the extent & structure of the resource in five major cities. *Landscape Ecology* 22, 601-615.
- Loram A, Warren PH, Gaston KJ. 2008. Urban Domestic Gardens (XIV): The Characteristics of Gardens in Five Cities *Environmental Management* 42, 361-376.
- Luck G, Smallbone L, O'Brien R. 2009. Socio-Economics and Vegetation Change in Urban Ecosystems: Patterns in Space and Time. *Ecosystems* 12, 604-620.
- MA (Millennium Ecosystem Assessment). 2003. Ecosystems and human well-being: a framework for assessment. Washington, DC: Island Press.
- Maas J, Verheij RA, Spreeuwenberg P, Groenewegen PP. 2008. Physical activity as a possible mechanism behind the relationship between green space and health: A multilevel analysis. *BMC Public Health* 8, 206.
- Magnus K, Matroos A, Strackee J. 1979. Walking, cycling, or gardening, with or without seasonal interruption, in relation to coronary events. *American Journal of Epidemiology* 110, 724-733.
- Mathieu R, Freeman C, Aryal J. 2007. Mapping private gardens in urban areas using object-oriented techniques and very high-resolution satellite imagery. *Landscape and Urban Planning* 81, 179-192
- Matteson KC, Langellotto GA. 2010. Determinates of inner city butterfly and bee species richness. *Urban Ecosystems* 13, 333-347.

- McMullen E, Gawkrödger DJ. 2006. Physical friction is under-recognized as an irritant that can cause or contribute to contact dermatitis. *British Journal of Dermatology* 154, 154-156.
- McPherson EG, Herrington LP, Heisler GM. 1988. Impacts of vegetation on residential heating and cooling. *Energy and Buildings* 12, 41-51.
- McPherson EG, Simpson JR, Livingston M. 1989. Effects of 3 landscape treatments on residential energy and water-use in Tucson, Arizona. *Energy and Buildings* 13, 127-138.
- Meehl GA, Tebaldi C. 2004. More intense, more frequent, and longer lasting heat waves in the 21st century. *Science* 305, 994-997.
- Milligan C, Bingley A. 2007. Restorative places or scary spaces? The impact of woodland on the mental well-being of young adults. *Health & Place* 13, 799-811.
- Milligan C, Gatrell A, Bingley A. 2004. 'Cultivating health': therapeutic landscapes and older people in northern England. *Social Science & Medicine* 58, 1781-1793.
- Mitchell R, Popham F. 2007. Greenspace, urbanity and health: relationships in England. *Journal of Epidemiology and Community Health* 61, 681-683.
- Morris J, Bagby J. 2008. Measuring environmental value for natural lawn and garden care practices. *International Journal of Life Cycle Assessment* 13, 226-234.
- Niinemets U, Peñuelas J. 2008. Gardening and urban landscaping: Significant players in global change. *Trends in Plant Science* 13, 60-65.
- Nowak DJ, Crane DE. 2002. Carbon storage and sequestration by urban trees in the USA. *Environmental Pollution* 116, 381-389.
- Nowak DJ, Crane DE, Stevens JC. 2006. Air pollution removal by urban trees and shrubs in the United States. *Urban Forestry and Urban Greening* 4, 115-123.

- Nowak DJ, Stevens JC, Sisinni SM, Luley CJ. 2002. Effects of urban tree management and species selection on atmospheric carbon dioxide. *Journal of Arboriculture* 28, 113-122.
- O'Connor BA, Carman J, Eckert K, Tucker G, Givney R, Cameron S. 2007. Does using potting mix make you sick? Results from a *Legionella longbeachae* case-control study in South Australia. *Epidemiology and Infection* 135, 34-39.
- Ogunseitan OA. 2005. Topophilia and the quality of life. *Environmental Health Perspectives* 113, 143-148.
- Olson LJ. 2006. The Economics of Terrestrial Invasive Species: A Review of the Literature. *Agricultural and Resource Economics Review* 35, 178-194.
- Ottosson J, Grahn P. 2005. A comparison of leisure time spent in a garden with leisure time spent indoors: On measures of restoration in residents in geriatric care. *Landscape Research* 30, 23-55.
- Overmyer JP, Noblet R, Armbrust KL. 2005. Impacts of lawn-care pesticides on aquatic ecosystems in relation to property value. *Environmental Pollution* 137, 263-272.
- Owen J. 2010. *Wildlife of a Garden: A Thirty-Year Study*. London: Royal Horticultural Society.
- Park S-A, Shoemaker CA, Haub MD. 2009. Physical and Psychological Health Conditions of Older Adults Classified as Gardeners or Nongardeners. *HortScience* 44, 206–210.
- Pataki DE, Carreiro MM, Cherrier J, Grulke NE, Jennings V, Pincetl S, Pouyat RV, Whitlow TH, Zipperer WC. 2011. Coupling biogeochemical cycles in urban environments: ecosystem services, green solutions, and misconceptions. *Frontiers in Ecology and the Environment* 9, 27-36.
- Pauleit S, Duhme F. 2000. Assessing the environmental performance of land cover types for urban planning. *Landscape and Urban Planning* 52, 1-20.

- Peñuelas J, Staudt M. 2010. BVOCs and global change. *Trends in Plant Science* 15, 133-144.
- Perry T, Nawaz R. 2008. An investigation into the extent and impacts of hard surfacing of domestic gardens in an area of Leeds, United Kingdom. *Landscape and Urban Planning* 86, 1-13.
- Pouyat R, Groffman P, Yesilonis I, Hernandez L. 2002. Soil carbon pools and fluxes in urban ecosystems. *Environmental Pollution* 116, S107-S118.
- Powell KE, Heath GW, Kresnow M-J, Sacks JJ, Branche CM. 1998. Injury rates from walking, gardening, weightlifting, outdoor bicycling, and aerobics. *Medicine & Science in Sports & Exercise* 30, 1246-1249.
- Qian YL, Follett RF. 2002. Assessing soil carbon sequestration in turfgrass systems using long-term soil testing data. *Agronomy Journal* 94, 930-935.
- Reichard SH, White P. 2001. Horticulture as a pathway of invasive plant introductions in the United States. *BioScience* 51, 103-113.
- Reid SB, Pollard E, Sullivan DC, Shaw SL. 2010. Improvements to lawn and garden equipment emissions estimates for Baltimore, Maryland. *Journal of the Air & Waste Management Association* 60, 1452-1462.
- Risbeth C. 2005. Gardens of ethnicity. In: Kingsbury N, Richardson T, editors. *Vista: The culture and politics of gardens*. London: Frances Lincoln. p 191.
- Robbins P, Polderman A, Birkenholtz T. 2001. Lawns and toxins - An ecology of the city. *Cities* 18, 369-380.
- Sailor DJ. 2008. A green roof model for building energy simulation programs. *Energy and Buildings* 40, 1466-1478.
- Santamouris M, Pavlou C, Doukas P, Mihalakakou G, Synnefa A, Hatzibiros A, Patargias P. 2007. Investigating and analysing the energy and environmental performance of an

- experimental green roof system installed in a nursery school building in Athens, Greece. *Energy* 32, 1781-1788.
- Schlosser WE, Bassman JH, Wandschneider PR, Everett RL. 2003. A carbon balance assessment for containerized *Larix gmelinii* seedlings in the Russian Far East. *Forest Ecology and Management* 173, 335-351.
- Schuman GE, Janzen HH, Herrick JE. 2002. Soil carbon dynamics and potential carbon sequestration by rangelands. *Environmental Pollution* 116(3), 391-396.
- Simons LA, Simons J, McCallum J, Friedlander Y. 2006. Lifestyle factors and risk of dementia: Dubbo Study of the elderly. *MJA* 184, 3.
- Simpson JR. 2002. Improved estimates of tree-shade effects on residential energy use. *Energy and Buildings* 34, 1067-1076.
- Smith C. 2010. London: Garden city? London: London Wildlife Trust, Greenspace Information for Greater London, Greater London Authority. 12 p.
- Smith C, Clayden A, Dunnett N. 2009. An exploration of the effect of housing unit density on aspects of residential landscape sustainability in England. *Journal of Urban Design* 14, 163–187.
- Smith RM, Baker RHA, Malumphy CP, Hockland S, Hammon RP, Ostojá-Starzewski JC, Collins DW. 2007. Recent non-native invertebrate plant pest establishments in Great Britain: origins, pathways, and trends. *Agricultural and Forest Entomology* 9, 307-326.
- Smith RM, Gaston KJ, Warren PH, Thompson K. 2005. Urban domestic gardens (V): relationships between landcover composition, housing and landscape. *Landscape Ecology* 20, 235-253.


- Smith RM, Gaston KJ, Warren PH, Thompson K. 2006a. Urban domestic gardens (VIII): environmental correlates of invertebrate abundance. *Biodiversity and Conservation* 15, 2515-2545.
- Smith RM, Thompson K, Hodgson JG, Warren PH, Gaston KJ. 2006b. Urban domestic gardens (IX): Composition and richness of the vascular plant flora, and implications for native biodiversity. *Biological Conservation* 129, 312-322.
- Sperling CD, Lortie CJ. 2010. The importance of urban backgardens on plant and invertebrate recruitment: a field microcosm experiment. *Urban Ecosystems* 13, 223-235.
- St Leger L. 2003. Health and nature – new challenges for health promotion. *Health Promotion International* 18, 173-175.
- Stewart ID. 2011. A systematic review and scientific critique of methodology in modern urban heat island literature. *International Journal of Climatology* 31, 200-217.
- Stigsdotter UA, Grahn P. 2002. What Makes a Garden a Healing Garden? *Journal of Therapeutic Horticulture* 13, 60-69.
- Strutt J, Wilson S, Shorney-Darby H, Shaw A, Byers A. 2008. Assessing the carbon footprint of water production. *American Water Works Association Journal* 100, 80-91.
- Syme GJ, Shao QX, Po M, Campbell E. 2004. Predicting and understanding home garden water use. *Landscape and Urban Planning* 68, 121-128.
- Takebayashi H, Moriyama M. 2007. Surface heat budget on green roof and high reflection roof for mitigation of urban heat island. *Building and Environment* 42, 2971-2979.
- Thompson K. 2007. No nettles required. The reassuring truth about wildlife gardening. London: Transworld.



- Thompson K, Austin KC, Smith RM, Warren PH, Angold PG, Gaston KJ. 2003. Urban domestic gardens (I): Putting small-scale plant diversity in context. *Journal of Vegetation Science* 14, 71-78.
- Thuille A, Buchmann N, Schulze E-D. 2000. Carbon stocks and soil respiration rates during deforestation, grassland use and subsequent Norway spruce afforestation in the Southern Alps, Italy. *Tree Physiology* 20, 849-857.
- Tratalos J, Fuller RA, Warren PH, Davies RG, Gaston KJ. 2007. Urban form, biodiversity potential and ecosystem services. *Landscape and Urban Planning* 83, 308-317.
- Trudgill S, Jeffery A, Parker J. 2010. Climate change and the resilience of the domestic lawn. *Applied Geography* 30, 177-190.
- Tubby K, Webber J. 2010. Pest and diseases threatening urban trees under a changing climate. *Forestry* 83, 451-459.
- Turner T. 2011. *European Gardens: History, Philosophy and Design*. Abingdon, UK: Routledge. 416 p.
- Tzoulas K, Korpela K, Venn S, Yli-Pelkonen V, Kazmierczak A, Niemela J, James P. 2007. Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and Urban Planning* 81, 167-178.
- Ulrich RS, Simons RF, Losito BD, Fiorito E, Miles MA, Zelson M. 1991. Stress recovery during exposure to natural and urban environments. *Journal of Environmental Psychology* 11, 201-230.
- van den Berg AE, Maas J, Verheij RA, Groenewegen PP. 2010. Green space as a buffer between stressful life events and health. *Social Science and Medicine* 70, 1203-1210.
- van Duijne FH, Kanis H, Hale AR, Green BS. 2008. Risk perception in the usage of electrically powered gardening tools. *Safety Science* 46, 104-118.

- Vickery ML. 1995. Gardens: the neglected habitat. In: Pullin AS, editor. Ecology and Conservation of Butterflies London: Chapman and Hall. p 123-134.
- Volckens J, Olson DA, Hays MD. 2008. Carbonaceous species emitted from handheld two-stroke engines. Atmospheric Environment 42, 1239-1248.
- Walsh JME, Rogo-Pressman A, Cauley JA, Browner WS. 2001. Predictors of physical activity in community-dwelling elderly white women. Journal of General Internal Medicine 16, 721-727.
- Watson GW, Malumphy CP. 2004. *Icerya purchasi* Maskell, cottony cushion scale (Hemiptera: *Margarodidae*), causing damage to ornamental plants growing outdoors in London. British Journal of Entomology and Natural History 17, 105-109.
- Whitford V, Ennos AR, Handley JF. 2001. “City form and natural process”—indicators for the ecological performance of urban areas and their application to Merseyside, UK Landscape and Urban Planning 57, 91-103.
- Williams JA, West CJ. 2000. Environmental weeds in Australia and New Zealand: issues and approaches to management. Australian Ecology 25, 425–444
- Xiao Q, McPherson EG. 2002. Rainfall interception by Santa Monica's municipal urban forest. Urban Ecosystems 6, 291-302.
- Yang J, McBride J, Zhou J, Sun Z. 2005. The urban forest in Beijing and its role in air pollution reduction. Urban Forestry and Urban Greening 3, 65-78.
- Zirkle G, Lal R, Augustin B. 2011. Modelling carbon sequestration in home lawns. HortScience 46, 808-814.

**Figure 1.** Typical carbon levels ( $\text{g} \times 10^6 \text{ ha}^{-1}$ ) stored in soils (upper 100mm only) and vegetation. Garden vegetation types compared to forest vegetation, including less densely planted urban forests.



Data derived from Jo and McPherson, 1995; Thuille et al., 2000; Schuman et al., 2002; Nowak and Crane, 2002; Euliss et al., 2006; Gough and Fritz, 2009.

**NOTE:**

1. Soil data shown is normalized for upper 10 cm of soil profile only, to facilitate comparisons. Total soil carbon storage, however, will be considerably greater in soils that have deeper organic profiles. For example, deeper root zones associated with prairie or meadow vegetation may increase soil organic matter by 3-4 fold over domestic lawns.
2. Values depicted are examples to provide relative scales only. *In situ* values vary considerably based on soil type, temperature, precipitation and density of vegetation.
3. 'Urban Forest proportional' relates to a scenario where the discrete, separate sections of urban forestry join and provide continuous cover across the urban matrix (Nowak and Crane, 2002). Carbon sequestration would be greater than conventional non-thinned forests because:
  - a. Planting densities are less thus allowing individual trees to access more light / atmospheric  $\text{CO}_2$ .
  - b. Average tree size is larger resulting in greater carbon storage within individual trees.
  - c. There are greater localized concentrations of  $\text{CO}_2$  in urban environments due to anthropogenic factors.