

# *Cannabinol and cannabidiol exert opposing effects on rat feeding patterns*

Article

Accepted Version

Farrimond, J. A., Whalley, B. J. and Williams, C. M. ORCID: <https://orcid.org/0000-0003-4452-671X> (2012) Cannabinol and cannabidiol exert opposing effects on rat feeding patterns. *Psychopharmacology*, 223 (1). pp. 117-129. ISSN 0033-3158 doi: 10.1007/s00213-012-2697-x Available at <https://centaur.reading.ac.uk/28010/>

It is advisable to refer to the publisher's version if you intend to cite from the work. See [Guidance on citing](#).

To link to this article DOI: <http://dx.doi.org/10.1007/s00213-012-2697-x>

Publisher: Springer Verlag

All outputs in CentAUR are protected by Intellectual Property Rights law, including copyright law. Copyright and IPR is retained by the creators or other copyright holders. Terms and conditions for use of this material are defined in the [End User Agreement](#).

[www.reading.ac.uk/centaur](http://www.reading.ac.uk/centaur)

**CentAUR**

Central Archive at the University of Reading

Reading's research outputs online

**Cannabinol and cannabidiol exert opposing effects on rat feeding patterns.**

**Jonathan A. Farrimond<sup>1,2</sup>, Benjamin J. Whalley<sup>1</sup> and Claire M. Williams<sup>2</sup>.**

<sup>1</sup>School of Pharmacy,  
University of Reading,  
Whiteknights,  
Reading,  
Berkshire,  
RG6 6AJ,  
U.K.  
0118 378 7088

<sup>2</sup>School of Psychology and Clinical Language Sciences  
University of Reading,  
Earley Gate,  
Whiteknights,  
Reading,  
Berkshire,  
RG6 6AL,  
U.K.  
0118 378 7540

Corresponding author: Claire Williams (claire.williams@reading.ac.uk)

This research was supported in part by the University of Reading Research Endowment Trust Fund (to JAF).

## Abstract

### *Rationale*

Increased food consumption following  $\Delta^9$ tetrahydrocannabinol-induced cannabinoid type 1 receptor agonism is well documented. However, possible non- $\Delta^9$ tetrahydrocannabinol phytocannabinoid-induced feeding effects have yet to be fully investigated. Therefore, we have assessed the effects of the individual phytocannabinoids, cannabigerol, cannabidiol and cannabinol upon feeding behaviors.

### *Methods*

Adult male rats were treated (p.o.) with cannabigerol, cannabidiol, cannabinol or cannabinol plus the CB<sub>1</sub>R antagonist, SR141716A. Prior to treatment, rats were satiated and food intake recorded following drug administration. Data were analyzed for hourly intake and meal microstructure.

### *Results*

Cannabinol induced a CB<sub>1</sub>R-mediated increase in *appetitive* behaviors via significant reductions in the latency to feed, and increases in *consummatory* behaviors via increases in meal one size and duration. Cannabinol also significantly increased the intake during hour 1 and total chow consumed during the test. Conversely, cannabidiol significantly reduced total chow consumption over the test period. Cannabigerol administration induced no changes to feeding behavior.

### *Conclusion*

This is the first time cannabinol has been shown to increase feeding. Therefore, cannabinol could, in the future, provide an alternative to currently used and psychotropic  $\Delta^9$ tetrahydrocannabinol-based medicines since cannabinol is currently considered to be non-psychotropic. Furthermore, cannabidiol reduced food intake in line with some existing reports, supporting the need for further mechanistic and behavioral work examining possible anti-obesity effects of cannabidiol.

**Keywords:** *cannabis, cannabigerol, cannabidiol, cannabinol, phytocannabinoids, feeding, appetite, behavio(u)r*

## 1 Abbreviations

2-AG	2-arachidonoylglycerol
$\Delta^9$ THC	$\Delta^9$ -tetrahydrocannabinol
$\Delta^9$ THCV	$\Delta^9$ -tetrahydrocannabivarin
AEA	Anandamide
ANOVA	Analysis of variance
BDS	Botanical drug substance
<i>C. sativa</i>	<i>Cannabis sativa</i>
CB <sub>1</sub> R	Cannabinoid type 1 receptor
CB <sub>2</sub> R	Cannabinoid type 2 receptor
CBD	Cannabidiol
CBG	Cannabigerol
CBN	Cannabinol
CNS	Central nervous system
eCB	Endocannabinoid
i.p.	Intraperitoneal
pCB	Phytocannabinoid
p.o.	Per ora
s.c.	Subcutaneous

## 1 Introduction

2 While *Cannabis sativa* (*C. sativa*) has been used on the Indian subcontinent and in China for thousands of years  
3 as a medicine, its use has been a source of controversy in Western medicine since its introduction in the 19<sup>th</sup>  
4 century due to widespread recreational use and abuse (O'Shaughnessey 1843; Wang et al. 2008). *C. sativa*'s  
5 pharmacological actions and psychotropic properties include sedation, analgesia, hypothermia, catalepsy and  
6 euphoria (Martin et al. 1981) alongside ravenous eating (Abel 1975).

7 Since the original identification of the cannabinoid type 1 and 2 receptors (CB<sub>1</sub> and CB<sub>2</sub>R; Devane et al. 1988;  
8 Matsuda et al. 1990; Munro et al. 1993) and confirmation of an endogenous cannabinoid system following the  
9 discovery of the endogenous cannabinoids (eCBs; anandamide (AEA; Devane et al. 1992) and 2-arachidonyl  
10 glycerol (2-AG; Mechoulam et al. 1995; Sugiura et al. 1995)) research has largely focused on the effects of  
11  $\Delta^9$ tetrahydrocannabinol ( $\Delta^9$ THC; Gaoni and Mechoulam 1964). Indeed, only limited research has considered the  
12 effects of the numerous other phytocannabinoids (pCBs) also present (Izzo et al. 2009). More recently, research  
13 has begun to examine the effects of these individual pCBs (for a review of cannabinoid pharmacology see Izzo  
14 et al. 2009). Currently, a range of possible cannabinoid-based therapies are being considered for a number of  
15 disorders (e.g. neurological and neurodegenerative, multiple sclerosis and anti-obesity (Glass 2001; Pryce et al.  
16 2003; Van Gaal et al. 2005), for review see Amar 2006). Interestingly, this new research has made it apparent  
17 that these pCBs are likely to act at sites other than CB<sub>1</sub> and CB<sub>2</sub>R due to their low binding affinities at these  
18 receptor sub-types (with the exceptions of  $\Delta^9$ -tetrahydrocannabivarin ( $\Delta^9$ THCV) and cannabinal (CBN);  
19 Petrosino et al. 2009). Importantly, the currently available literature gives no indication of non- $\Delta^9$ THC pCB  
20 psychoactivity (for reviews see Amar, 2006; Izzo et al., 2009).

21 Specifically in terms of feeding, and unlike the other pCBs,  $\Delta^9$ THC has been relatively well studied. Indeed,  
22 some time ago it became apparent that CB<sub>1</sub>R sites in the central nervous system (CNS; Herkenham et al. 1991)  
23 were responsible for  $\Delta^9$ THC-mediated increases in feeding (Williams and Kirkham 2002a; Williams and  
24 Kirkham 2002b; Williams et al. 1998).  $\Delta^9$ THC-induced CB<sub>1</sub>R-mediated hyperphagia following a prefeed  
25 process is classically described by increases in consumption during the first hour of testing due to significant  
26 decreases in the latency to feed without concomitant increases in meal size and duration (Williams and Kirkham  
27 2002b; Williams et al. 1998).  $\Delta^9$ THC-induced hyperphagia has been shown to be CB<sub>1</sub>R-mediated in

1 experiments which co-administered  $\Delta^9$ THC alongside the CB<sub>1</sub>R antagonist SR141716A (Rinaldi-Carmona *et*  
2 *al.*, 1994) even though in the same paradigm SR141716A alone was unable to alter feeding patterns (Williams  
3 and Kirkham 2002b). Similar alterations to feeding patterns have also been observed following exogenous AEA  
4 administration where AEA reduced the latency to feed but also increased meal size and duration (Hao *et al.*  
5 2000; Jamshidi and Taylor 2001; Williams and Kirkham 2002a). As such, it has been suggested that CB<sub>1</sub>R-  
6 mediated alterations to feeding patterns can be divided into *consummatory* (those which control intake quantity)  
7 and *appetitive* (those which control feeding pattern) behaviors (Farrimond *et al.* 2011a).

8 Recently in our lab, we have described alterations to feeding behaviors induced by a variety of non- $\Delta^9$ THC  
9 pCBs when administered as standardized cannabis extracts, i.e. botanical drug substances (BDS). Following  
10 administration of a  $\Delta^9$ THC-rich standardized extract (high- $\Delta^9$ THC BDS; 67%  $\Delta^9$ THC, 6.5% other pCBs), we  
11 observed a reduction in  $\Delta^9$ THC-induced hyperphagia when our extract was compared to purified  $\Delta^9$ THC alone  
12 (Farrimond *et al.* 2010a). Interestingly, in subsequent trials we demonstrated that a  $\Delta^9$ THC-free extract analogue  
13 (non- $\Delta^9$ THC pCB content matched to the high- $\Delta^9$ THC BDS; Farrimond *et al.* 2011b) and a second standardized  
14 extract which contained little  $\Delta^9$ THC (low- $\Delta^9$ THC BDS; 6.9%  $\Delta^9$ THC, 14.2% other pCBs; Farrimond *et al.*  
15 2010b) administered at  $\Delta^9$ THC doses below those previously observed to alter feeding patterns could both  
16 increase feeding in male rats. Importantly, the  $\Delta^9$ THC-free extract analogue altered feeding behaviors by  
17 reducing the latency to feed and increasing the quantity of food consumed during both the first hour of testing  
18 and the first meal in the same manner as the high- $\Delta^9$ THC BDS and purified  $\Delta^9$ THC did, but without increases in  
19 first meal duration. However, the low- $\Delta^9$ THC BDS significantly increased *appetitive* behaviors, only inducing  
20 hyperphagia as a result of a highly significant decrease in the latency to feed but without concomitant increases  
21 in meal size and duration. These data have led us to suggest that non- $\Delta^9$ THC pCBs can not only modulate the  
22 feeding effects of  $\Delta^9$ THC but also induce alterations to feeding behaviors by themselves. However, our previous  
23 data shed no light on the specific contributions made by the individual pCBs found in our BDS' to changes in  
24 feeding behaviors.

25 To date, there has only been limited research of the effects of the non- $\Delta^9$ THC pCBs on feeding behaviors. In  
26 1976, Sofia and Knobloch reported that CBN (50.0 mg/kg; intraperitoneal injection (i.p.)) reduced food intake in  
27 rats, an effect that has yet to be recapitulated (Sofia and Knobloch 1976). However, one might expect that CBN  
28 could elicit hyperphagia because, like  $\Delta^9$ THC, it exhibits CB<sub>1</sub>R agonist properties (Felder *et al.* 1995). In

contrast, cannabidiol (CBD) exerts a superfluity of intracellular effects *in vitro* (e.g. modulation of Ca<sup>2+</sup> homeostasis; Ryan et al. 2009 and AEA reuptake and FAAH inhibition; De Filippis 2008; Izzo et al. 2009) and has been employed in a small number of feeding studies. Wiley *et al.*, (2005) reported that CBD (3, 10, 30 and 100 mg/kg; i.p.) did not affect food intake in mice, a result confirmed by Scopinho et al. (2011) who demonstrated that CBD (1, 10 or 20 mg/kg; i.p.) did not affect feeding in rats. Similar data, in mice, were also recently described by Riedel (2009, 10.0 mg/kg; i.p.). Conversely however, Sofia and Knobloch (1976) reported a CBD-induced (50 mg/kg; i.p.) reduction in feeding in rats. Very recently, these data have been supported by the observation that CBD (2.5 and 5 mg/kg; i.p.) can reduce body weight gain in relatively young (260 ± 20 g at the start of testing) rats over a period of two weeks, a finding which suggests either reduced food consumption or increased activity over the test period (Ignatowska-Jankowska et al. 2010). As such, data describing the effects of CBD on feeding remains inconclusive and the mechanisms by which it could increase or decrease intake and/or body weight remain to be elucidated.

To our knowledge the possible effects of cannabigerol (CBG) on feeding have yet to be examined although such investigation is warranted since CBG shows partial agonism at CB<sub>1</sub>R and/or CB<sub>2</sub>R sites (Pertwee 2008), possible antagonism at CB<sub>1</sub>R sites (Cascio et al. 2010), phospholipase A2 activation (Evans et al. 1987) and/or AEA reuptake inhibition (Ligresti et al. 2006). Therefore, it is conceivable that CBG administration could induce either hyper- or hypo-phagic effects.

Considering the poor side-effect profile of current Δ<sup>9</sup>THC-based anti-anorectic agents (e.g. hallucinations; BNF 2006), and given the drive to produce new anti-obesity agents which do not cause unwanted side effects (*viz.* SR141716A; EMA 2009 or MK-0364; Clark 2009) it is clear that further research examining the possible feeding effects of pCB could prove therapeutically useful. Furthermore, considering the myriad of protocols thus far used to test possible feeding effects of pCBs, direct comparisons of these data are limited. To address this, we have administered CBD, CBG and CBN individually using the same prefeed protocol that we have successfully used to highlight hyperphagic actions of Δ<sup>9</sup>THC. Furthermore, in order to assess possible CB<sub>1</sub>R-mediation of any observed CBN effects we have also performed a CBN and SR141716A co-administration trial. We present an analysis of hourly intakes and critical meal parameters following drug administration.



## General Methods

### *Animals*

Thirty adult, male Lister-hooded rats ( $P > 40$ , 200 – 250 g at the start of testing, Harlan UK Ltd, England) were maintained in a temperature controlled environment (21 - 22 °C) under a 12:12 hour light:dark cycle (red light on at 10:30 hrs). Given the distinct pharmacological profiles of CBD, CBG and CBN (reviewed in Farrimond et al, 2011a), direct comparisons between the drugs on feeding behaviour would yield little pertinent data, thus rats were split into three groups of ten animals; with each group acting as its own control and receiving a different test substance (see table 1). Normal laboratory chow (PCD Mod C, Special Diet Services, Witham, England) was available *ad libitum* but on test days was removed for a three hour period and replaced with a prefeed mash for a two hour period (see ‘*Prefeed Procedure*’) which was followed by one hour of food deprivation immediately after drug administration. Fresh tap water was available *ad libitum*. All procedures were performed in compliance with the requirements of the United Kingdom Animals (Scientific Procedures) Act 1986.

### *Test Environment*

All tests were performed during the dark light phase under low intensity red light (~4 lx). Testing took place in standard plastic cages, each fitted with a modified food hopper connected, via a strain gauge weighing device, to a computer running data acquisition and analysis software (The Feeding and Drinking Monitor v 2.16, TSE Systems GmbH, Bad Homburg, Germany) which permitted continuous monitoring of food intake. In addition, each cage was fitted with a CCTV camera positioned above each cage to allow an unimpeded view of rat behavior (distance from cage to camera approximately 10 cm). Food intake data were analyzed to provide information on hourly food intakes as well as critical meal parameters such as latency to onset of meals, individual meal size and duration. For the purposes of this study, a meal was defined as any feeding episode causing a change in food weight of  $\geq 0.1$  g, lasting at least 1 minute and separated by at least 15 minutes from any subsequent episode. These criteria have been previously used to facilitate the visualization and interpretation of drug effects on feeding behavior and to distinguish prolonged eating episodes from more transient, exploratory contacts with food (Williams and Kirkham 2002a). Consecutive feeding events separated by intervals of  $< 15$  minutes were considered to be part of a single meal. Due to these criteria, in some

instances, animals have or have not chosen to consume meals during different test hours. Therefore, some ANOVA results have different degrees of freedom and F values than might be expected.

### *Drugs*

Fresh solutions of CBG, CBD and CBN (GW Pharmaceuticals, Salisbury, England) were prepared 15 minutes before administration on each test day. All pCBs were dissolved in a sesame seed oil vehicle (Sainsbury's Supermarkets Ltd, London, England) and the doses specified in Table 1 were administered. The presented pCB doses were based on multiples of 1x (low), 10x (medium) and 100x (high) times the concentrations present in a low- $\Delta^9$ THC cannabis extract that we have previously shown to induce increases in appetitive behaviors (Farrimond et al. 2010b). Phytocannabinoids were delivered orally via a syringe placed into the rat's cheek pouch (*per ora*; p.o.).

In a second experiment, because of the likelihood of CB<sub>1</sub>R involvement in any observed CBN effects, we co-administered 26.0 mg/kg CBN with SR141716A (1.0 mg/kg) and compared these data to that collected previously following CBN alone administration. SR141716A was administered in a 1:1:18 vehicle made as 1 part ethanol (Fisher Scientific UK Ltd, Loughborough, UK), 1 part cremophor (Sigma-Aldrich, St Louis, USA) and 18 parts 0.9% sodium chloride (Fisher Scientific UK Ltd, Loughborough, UK) saline via subcutaneous injection (s.c.).

Both administration methods (p.o. and s.c.) were calculated to have an injection volume of 1.0 ml/kg. Each group of animals received their drug treatments according to a Latin Square design, counterbalanced for phytocannabinoid dose, with at least 48 hours between successive treatments. All drug groups used vehicle controls as part of the Latin square design. Drug administration began only after animals had been habituated to housing conditions, oral dosing, s.c. injections and all subsequent test procedures.

Throughout these tests, no non-specific behavioral effects of any drug at any dose were evident

### *Prefeed Procedure*

1 In all experiments, rats were transferred from home cages to individual test cages immediately after dark onset  
2 (10:30 hrs) and presented with 30 g of a wet mash diet for 120 minutes as a prefeed. Any remaining wet mash  
3 and spillage was recovered after 120 minutes and weighed. Animals were fully habituated to the prefeed  
4 procedure before testing began and drug administration did not begin until prefeed intakes were stable as  
5 assessed by a non-significant one-way analysis of variance (ANOVA).

## 6 7 *Procedure*

8 Following removal of the prefeed at 12:30 hrs, all drugs were administered to the rats according to a Latin  
9 square design. Rats were then deprived of food until 13:30 hrs to allow for drug assimilation. At 13:30 hrs, 30 g  
10 of normal laboratory chow were placed into the food hoppers. Subsequent hourly food intake (calculated from:  
11 starting food mass – (remaining food mass + spillage)) was measured for four consecutive hours.

## 12 13 *Statistical Analysis*

14 Hourly food intake was analyzed by two-way ANOVA with four dose levels (vehicle, low, medium and high)  
15 and four time points (hours 1, 2, 3 and 4), where appropriate this analysis was followed by individual one-way  
16 ANOVA tests for each time point and bonferroni post-hoc tests. The data collected for each meal parameter  
17 following test substance administration were separately analyzed using one-way ANOVA with four dose levels  
18 (vehicle, low, medium and high), with bonferroni post-hoc tests performed where appropriate. Following  
19 SR141716A plus CBN co-administration, the same hourly intake and meal parameter data were analyzed by  
20 further one-way ANOVA with three drug levels (vehicle, CBN alone and CBN plus SR141716A), Bonferroni  
21 post-hoc tests were carried out when appropriate. All tests were performed using IBM SPSS Statistics 19  
22 (International Business Machines Corp., Armonk, USA).

## Results

### *Cannabinol alone and cannabinol co-administered with SR141716A*

Before testing began, prefeed intakes were stabilized for both CBN-only administration ( $F(12,122)=1.277$ ,  $p=0.242$ ) and CBN-SR141716A co-administration ( $F(4,49)=1.538$ ,  $p=0.207$ ). During testing animals consumed 19.40 ( $\pm 0.57$ ) and 19.50 ( $\pm 0.46$ ) g of prefeed per day respectively. Furthermore, upon rearrangement of prefeed intakes by dose, no significant differences were apparent between any prefeed intakes for any individual dose of either CBN alone, CBN plus SR141716A or their respective vehicle-treatments ( $F(5,57)=0.113$ ,  $p=0.989$ ).

### *Hourly intake*

Two-way analysis of variance failed to show significant effects of either dose ( $F(3,60)=0.973$ ,  $p=0.411$ ) or time ( $F(3,20)=0.807$ ,  $p=0.505$ ), however, there was a significant time by dose interaction ( $F(9, 60)= 2.704$ ,  $p=0.010$ ). Subsequent analysis of effects for each individual hour showed that CBN significantly increased chow consumption during the first hour (Figure 1 (panel A, white bars);  $F(3,34)=7.663$ ,  $p=0.001$ ) from a vehicle-treated intake of  $0.86 \pm 0.51$  g to  $2.87 \pm 0.45$  g at the 26.0 mg/kg dose. Post-hoc analysis revealed that intake following 26.0 mg/kg CBN was significantly greater than after vehicle treatment ( $p=0.010$ ), no other doses induced significant hyperphagic effects. During the second hour of testing, a marginal effect was apparent (Figure 1 (panel A, light grey bars);  $F(3,34)=2.391$ ,  $p=0.088$ ) which is most likely due to the small increases in feeding seen following the 2.60 mg/kg dose compared to vehicle treatment; intakes increased from  $0.86 \pm 0.51$  g following vehicle treatment to  $1.44 \pm 0.44$  g at the 2.6 mg/kg dose. However, post-hoc tests show no significant differences between chow intakes following any CBN treatment when compared to vehicle treatments ( $p \geq 0.938$  in all cases) during hour two. Significant increases were observed in all cumulative combinations of hourly intake (Figure 1 (panel B; light grey, grey and black bars), hours one and two;  $F(3,34)=3.590$ ,  $p=0.025$ , hours one, two and three;  $F(3,34)=4.635$ ,  $p=0.009$  and all four hours;  $F(3,34)=3.509$ ,  $p=0.027$ ).

Co-administration of SR141716A with CBN blocked the previously observed CBN-mediated increases in hour one intake (Figure 2 (panel A, white bars);  $p=0.696$ ). Furthermore, SR141716A co-administered with CBN also blocked the marginally significant increase in chow consumption observed during the second hour of testing

(Figure 2 (panel A, light grey bars);  $F(2,27)=2.099$ ,  $p=0.114$ ) and during each consecutive cumulative hourly arrangement of animals' intakes (Figure 2 (panel B; light grey, grey and black bars), hours one and two;  $F(2,27)=1.351$ ,  $p=0.227$ , hours one, two and three;  $F(2,27)=0.974$ ,  $p=0.392$  and all four hours;  $F(2,27)=2.112$ ,  $p=0.142$ ). However, during the third hour of testing intakes recorded for the two vehicle-treated conditions (those animals which received both the sesame oil and 1:1:18 ethanol:cremophor:saline vehicles) varied such that SR141716A plus CBN co-administration vehicle-treated animal intakes were ~2 g higher than their CBN vehicle-treated counterparts (0.1 g). As such, during the third hour of CBN plus SR141716A treatment, a significant reduction in chow consumption compared to control was apparent ( $F(2,27)=3.940$ ,  $p=0.033$ ) such that SR141716A plus CBN treated animals displayed significantly reduced intakes versus vehicle-treated animals ( $p=0.038$ ).

#### *Alterations to meal pattern*

Following CBN administration, the observed increase in hour one food consumption (Figure 1, panel A, white bars) was due to a significant dose-dependent increase in the size of the first meal (Figure 1 (panel C);  $F(3,34)=4.377$ ,  $p=0.011$ ) and a reduction in the latency to the first meal (Figure 1, (panel D, light grey bars);  $F(3,34)=5.217$ ,  $p=0.005$ ) which shifted feeding into the first hour of the test. However, post-hoc Bonferroni tests revealed no significant differences in meal one size following any individual dose of CBN versus vehicle treatments, whilst the latency to meal one was significantly reduced from  $96.7 \pm 26.7$  to  $10.8 \pm 4.6$  min (at a dose of 26.0 mg/kg CBN;  $p=0.038$ ). In conjunction with the dose-dependent increase in the size of the first meal, its duration was also significantly increased in a dose-dependent manner (Figure 1 (panel D, light grey bars);  $F(3,34)=2.963$ ,  $p=0.047$ ). Consistent with the previously reported abolition of the CBN effect upon hourly intake, SR141716A blocked the CBN induced increases in meal one size (Figure 2 (panel C):  $p=0.374$ ), meal one duration (Figure 2 (panel D):  $p=1.000$ ) and the latency to feed (Figure 2 (panel D):  $p=1.000$ ), supporting a  $CB_1R$ -mediated mechanism for CBN. Analysis of second, third and inter-meal interval parameters is not included as less than four rats consumed second or third meals in any given dose group during this test.

## *Cannabidiol administration*

After habituation to test procedures prefeed intakes were stabilized ( $F(6,69)=1.282$ ,  $p=0.279$ ) and CBD administration commenced. During the test period animals consumed  $16.57 \pm 0.46$  g of prefeed per test day.

## *Hourly intake*

Here, 2-way ANOVA failed to show a significant effect of CBD treatment ( $F(3,108)=1.380$ ,  $p=0.253$ ) or any dose by time interaction ( $F(9,108)=1.412$ ,  $p=0.192$ ). However, a significant effect of time was seen ( $F(3,36)=7.338$ ,  $p=0.001$ ) indicating that chow intake did alter over the course of the experiment. However, 1-way ANOVA for each individual hour showed that chow consumption following CBD administration did not vary significantly from those observed for vehicle treatments during any individual hour (Figure 3, panel A: hour 1 (white bars);  $F(3,37)=0.394$ ,  $p=0.758$ , hour 2 (light grey bars);  $F(3,37)=2.088$ ,  $p=0.120$ , hour 3 (grey bars);  $F(3,37)=0.868$ ,  $p=0.467$  or hour 4 (black bars);  $F(3,37)=0.481$ ,  $p=0.698$ ). Cumulative food intakes in hours one and two ( $0.77 \pm 0.19$  g) and one, two and three ( $2.31 \pm 0.26$  g) also showed no significant variation from vehicle treatments induced by CBD administration (Figure 3, panel B, light grey and grey bars;  $F(3,39)=1.837$ ,  $p=0.158$  and  $F(3,39)=1.033$ ,  $p=0.390$  respectively). However importantly, CBD induced significant dose-dependent reductions in total food intake over the total four hour test period (Figure 3 (panel B; black bars);  $F(3,39)=3.343$ ,  $p=0.030$ ). Vehicle treated animals consumed  $4.06 \pm 0.44$  g of chow which was reduced following administration of the highest CBD dose (4.40 mg/kg) to  $2.59 \pm 0.36$  g during four hours.

## *Alterations to meal pattern*

Whilst CBD administration significantly reduced the total amount of food consumed in all meals combined, it had no effect on all other meal parameters. Specifically, no significant effects of CBD administration were observed for the latency to meal one ( $121.8 \pm 10.8$  min, Figure 3 D;  $F(3,37)=1.635$ ,  $p=0.196$ ), the intake during ( $1.88 \pm 0.21$  g) or duration ( $7.8 \pm 0.9$  min) of meal one (Figure 3 C;  $F(3,37)=0.570$ ,  $p=0.638$  and Figure 3 D;  $F(3,37)=0.523$ ,  $p=0.670$  respectively), the cumulative intakes or durations of meals one and two combined ( $3.04 \pm 0.25$  g;  $F(3,37)=0.957$ ,  $p=0.424$  and  $13.4 \pm 1.5$  min;  $F(3,37)=1.250$ ,  $p=0.307$  respectively) or total duration of

1 all consumed meals ( $17.1 \pm 2.1$  min;  $F(3,37)=1.523$ ,  $p=0.226$ ). Please note that quoted values are averages  $\pm$   
2 S.E.M. collapsed by dose. Analysis of second, third and inter-meal interval parameters is not included as less  
3 than four rats consumed second or third meals in any given dose group during this test.

4

## *CBG administration*

Prefeed intakes were stabilized before testing began ( $F(9,99)=1.395$ ,  $p=0.202$ ). On each test day animals receiving CBG consumed  $18.94 \pm 0.44$  g.

## *Hourly intake*

Two-way ANOVA failed to show any significant effect of dose ( $F(3,72)=0.872$ ,  $p=0.460$ ), time ( $F(3,24)=2.135$ ,  $p=0.122$ ) or time by dose interaction ( $F(3,72)=0.990$ ,  $p=0.456$ ). CBG administration induced no significant changes from vehicle-treated animal intakes during any hour of the test (Figure 4 panel A: hour 1 (white bars);  $F(3,33)=0.739$ ,  $p=0.537$ , hour 2 (light grey bars);  $F(3,33)=2.105$ ,  $p=0.121$ , hour 3 (grey bars);  $F(3,33)=1.278$ ,  $p=0.300$  and hour 4 (black bars);  $F(3,33)=1.473$ ,  $p=0.242$ ) or in any cumulative hourly arrangement of chow intakes (Figure 4 panel B: hour 1 (white bars);  $F(3,33)=0.739$ ,  $p=0.537$ , hours 1 and 2 (light grey bars);  $F(3,33)=0.810$ ,  $p=0.498$ , hours 1, 2 and 3 (grey bars);  $F(3,33)=0.834$ ,  $p=0.486$  and total intake (black bars);  $F(3,39)=1.563$ ,  $p=0.215$ ).

## *Alterations to meal pattern*

In conjunction with hourly intake quantities, CBG administration had no effect on meal patterns. Indeed, meal one intake remained constant at  $2.20 \pm 0.23$  g (Figure 4, panel C:  $F(3,29)=0.488$ ,  $p=0.694$ ), the latency to the first meal at  $110.9 \pm 14.4$  min (Figure 4, panel D:  $F(3,29)=0.597$ ,  $p=0.622$ ) and the duration of the first meal at  $12.4 \pm 1.9$  min (Figure 4, panel D:  $F(3,26)=0.123$ ,  $p=0.945$ ). Analysis of second, third and inter-meal interval parameters is not included as less than six rats consumed second or third meals in any given dose group during this test.



## Discussion

In this study, the effects of CBD, CBG and CBN on feeding patterns in adult male rats were investigated. Here, the results obtained demonstrate that CBN can stimulate feeding and alter meal patterns in rats whilst, CBD significantly reduces intake. CBG had no effect upon feeding patterns using the experimental paradigm employed here. It should be noted, however, that non-significant differences in intake under vehicle control conditions exist between our three experiments and as a consequence these differences may limit the extent of interpretation of the drug effects.

CBN induced a dose-dependent increase in chow consumption during the first hour, as illustrated by a significant increase in intake versus vehicle-treated intakes at its highest dose. This significant increase in first hour intake can be attributed to significant decreases in the latency to feed which altered the temporal arrangement of feeding such that the first meal occurred in the first, rather than the second, hour of testing (Figure 1). Furthermore, CBN increased the size and duration of the first meal but, importantly and unlike  $\Delta^9$ THC (Farrimond et al. 2010a), also increased the total amount of food consumed during the test period. Indeed, in the present case, total chow intake following CBN administration was significantly increased by ~60% compared to vehicle-treatments during the test period, whereas previously, we observed a non-significant change of ~2% following  $\Delta^9$ THC administration compared to vehicle-treatments over the same four hour period (Farrimond et al. 2010a). CBN's effects upon appetitive aspects of feeding (i.e. decreased latency to feed resulting in increased intake during the first hour of testing) and increases in the total amount of chow consumed mirror the behavioral effects of administration of the eCB, AEA, which have been shown to be CB<sub>1</sub>R-mediated (Hao et al. 2000; Jamshidi and Taylor 2001; Koch and Matthews 2001; Williams and Kirkham 1999; 2002a). Indeed, radioligand binding has demonstrated that CBN is a CB<sub>1</sub>R agonist (Rhee et al. 1997) which justified our co-administration of the CB<sub>1</sub>R antagonist, SR141716A, with CBN. This co-administration duly blocked first hour and first meal intake increases and the reduction in the latency to feed. These results conclusively demonstrate that the changes to feeding patterns seen following CBN administration alone were CB<sub>1</sub>R-mediated. Indeed, we have demonstrated that SR141716A blocked CBN-induced changes to all meal parameters and hour one intake and observed no significant effect of CBN alone administration in any subsequent hour. Therefore, even though during the third hour of SR141716A CBN co-administration testing there were

differences in third hour vehicle-treated intakes between the CBN alone and SR141716A CBN co-administration trials, such differences do not hinder the analysis of our data

It is interesting that CBN and  $\Delta^9$ THC administration induced different changes to feeding patterns, even though both have been shown to affect feeding via a solely CB<sub>1</sub>R-mediated mechanism (see Williams and Kirkham 2002b for CB<sub>1</sub>R involvement in  $\Delta^9$ THC-mediated hyperphagia). In this test we have seen that the hourly effects of CBN administration do not exhibit the reduction in second hour chow consumption which is characteristic of  $\Delta^9$ THC-mediated modulation of feeding patterns. This lack of compensatory effect has led to increased total chow consumption in this study. Indeed,  $\Delta^9$ THC has previously been found to significantly increase intakes during the first hour of testing, but be followed by a significant reduction in feeding in the second hour; at the highest administered  $\Delta^9$ THC dose (2.68 mg/kg; p.o.), intakes during hour two were ~17% of vehicle-treated intakes (Farrimond et al. 2010a). The reason for this lack of compensatory mechanism may suggest that CBN remained at higher concentrations in the brain for an increased period of time compared to  $\Delta^9$ THC (due to the comparatively higher administered doses), or could be due to differences in the psychotropic properties of the two pCBs. Due to CBN's lack of observed psychotropic side effects it is possible to administer CBN at higher doses than  $\Delta^9$ THC without disruptions to feeding patterns caused by non-specific behaviors (i.e. motor incoordination). These comparatively higher doses of CBN may have led to increased feeding behaviors with a longer duration of action.

It is also intriguing that CBN's significant hyperphagic effects only manifested at a dose of 26 mg/kg versus vehicle-treatments, and not at any lower dose. While CBN is an agonist at CB<sub>1</sub>R, its disassociation constant ( $K_i$ ) is approximately five times greater than that of  $\Delta^9$ THC (CBN: 211.2 nM; Rhee et al. 1997 versus  $\Delta^9$ THC: 39.5 nM; Bayewitch et al. 1996). Therefore, CBN's lower affinity at CB<sub>1</sub>R could explain the observed difference in effective doses when compared with  $\Delta^9$ THC where a maximal effect in this paradigm is seen at 2.68 mg/kg; Farrimond et al. 2010a). Furthermore, CBN's lower affinity for CB<sub>1</sub>R could also explain why no evidence supports psychotropic effects of CBN which are commonly associated with CB<sub>1</sub>R activation. However, we must accept that the gross visual analysis of behaviors we used here does not preclude the possibility of non-specific behavioral side effects. Therefore, while we believe it is highly unlikely that the administered CBN had any non-specific behavioral side effects for the previously mentioned reasons, we suggest that further experiments

1 be performed using a battery of behavioral tests (e.g. balance bars) which would fully determine any  
2 psychoactive properties of CBN.

3 Previously, we administered CBN (0.26 mg/kg) with  $\Delta^9$ THC (0.27 mg/kg) and various other pCBs and observed  
4 significant hyperphagia (Farrimond et al. 2010b). When CBN was administered alone at 0.26 mg/kg in this  
5 study, and purified  $\Delta^9$ THC alone at 0.34 mg/kg previously (Farrimond et al. 2010a), we observed no significant  
6 alterations to feeding patterns. This comparison clearly suggests that  $\Delta^9$ THC and CBN interact synergistically in  
7 some way to induce changes to feeding patterns at doses which have previously been shown to be ineffective  
8 when administered alone. Further studies are required to fully characterize the behavioral adjustments induced  
9 by CBN  $\Delta^9$ THC co-administration and any mechanisms via which CBN and  $\Delta^9$ THC may interact to alter  
10 feeding patterns. Co-administration of  $\Delta^9$ THC and CBN could have the therapeutic advantage that similar  
11 increases in feeding could be induced by doses of  $\Delta^9$ THC below those currently used which induce unwanted  
12 side effects.

13 The data presented here contradicts, to an extent, that previously published by Sofia and Knobloch (Sofia and  
14 Knobloch 1976) where a significant reduction in feeding at a CBN dose twice as high as the highest presented  
15 here (50.0 mg/kg; i.p.) was reported. However, not only were Sofia's experiments conducted over a  
16 considerably different time scale (daily food intake measurements over a six day period, rather than a four hour  
17 test) but the route of administration (i.p.) would have caused a more rapid increase in plasma/brain  
18 concentrations of CBN which would have reached a higher maximum concentration than via p.o. administration  
19 employed here.

20 It must also be noted that repeated  $\Delta^9$ THC administration has previously been linked to sensitization effects in  
21 rat models. Both Cadoni *et al.*, (2001) and Runbino *et al.*, (2001) have observed that if rats are pretreated twice a  
22 day for three or five days respectively with  $\Delta^9$ THC then, after a washout period, they react more strongly to  
23 further  $\Delta^9$ THC administration compared to untreated controls, an effect that can be removed by administration  
24 of SR141716A. As such sensitization has not yet been demonstrated following CBN, CBD or CBG  
25 administration and given the lower affinity of CBN for CB<sub>1</sub>R  $\Delta^9$ THC compared to  $\Delta^9$ THC, the distinct  
26 pharmacologies of both CBD and CBG and since both Cadoni *et al.*, (2001) and Runbino *et al.*, (2001)  
27 demonstrated sensitization following i.p. not p.o. administration in non-feeding behavioral tests it is unlikely  
28 that non- $\Delta^9$ THC pCB-induced behavioral sensitization is affecting the presented results.

1 Here we also present results which demonstrate that CBD administration can induce significant reductions in  
2 chow consumption over a four hour period. Specifically, CBD administration induced only subtle, non-  
3 significant reductions in animal intake during any individual hour of the test; however, together this led to a  
4 significant reduction in total chow intake over the test period due to significant reductions in intake during all  
5 meals. It is worthy of mention that these apparent late-onset of suppressive effects may reflect the relatively  
6 slow pharmacokinetic profile of CBD. Indeed, Deiana et al (2011) recently showed that brain levels of CBD  
7 continued to rise progressively for 4 hours following a 120 mg/kg oral dose. Despite these effects on hourly  
8 intakes, CBD administration had no significant effect on any other critical meal parameter. Such behaviors have  
9 been intimately linked to CB<sub>1</sub>R activation, and since it is currently thought that CBD is unlikely to interact with  
10 CB<sub>1</sub>R (Hill *et al.*, 2011), these data may suggest that CBD can affect a feeding pathway which is unrelated to  
11 CB<sub>1</sub>R. Such data fit well with the reductions in chow consumption previously reported by Sofia and Knobloch  
12 in 1976 who also demonstrated a CBD-mediated reduction in chow consumption. Very recently, Ignatowska-  
13 Jankowska has shown that CBD (2.5 and 5.0 mg/kg; i.p.) can reduce body weight gain in young rats, suggesting  
14 either reduced food intake or increased activity, and therefore indirectly supporting the reductions observed by  
15 Sofia and Knobloch and the results presented here (Ignatowska-Jankowska et al. 2010). However, data which  
16 describe the effects of CBD administration on feeding patterns are not yet conclusive. Recently, Wiley and  
17 colleagues (Wiley et al. 2005) and Scopinho and colleagues (Scopinho et al. 2011) observed no effect of CBD  
18 on feeding patterns in food restricted mice and normally fed and fasted rats respectively. Wiley used 3.0 – 100.0  
19 mg/kg CBD, whilst Scopinho used 1.0 – 20.0 mg/kg CBD; both administered i.p.. Therefore, it seems likely that  
20 the differences between the experimental protocols used by Wiley and Scopinho and those presented here could  
21 feasibly explain the differences in reported effects. Indeed, due to the route of administration (p.o.) used in our  
22 study, it is likely that peak cerebrospinal fluid concentrations were considerably lower than those achieved with  
23 the i.p. route used by Wiley and Scopinho. Furthermore, since neither Wiley nor Scopinho used pre-fed animals  
24 it is likely that differences in endocannabinergic tone between the models will have altered feeding behaviors,  
25 and consequently, the animals' responses to CBD administration.

26 It seems apparent from the available literature that the functional effect of CBD to induce significant decreases  
27 in chow intake could arise from the numerous intra- and extra-cellular mechanisms with which it is known to  
28 interact, but are likely to be unrelated to traditional CB<sub>1</sub>R-mediated feeding control. Such a theory is supported  
29 by CBD administration's failure to affect any meal parameter or individual hourly intake in this test.

1 Unfortunately, the relatively wide spectrum of cellular and molecular mechanisms that have been proposed but  
2 not definitively established *in vivo* make such suggestions highly speculative and further investigations that  
3 probe the discrete mechanisms potentially involved are required to confirm the mechanisms underlying the  
4 observed functional effects. Clearly, it would be most interesting to establish a non-CB<sub>1</sub>R-mediated feeding  
5 pathway that is modulated by a pCB, such as CBD, although the lack of pharmacological tools with which to  
6 block CBD's putative AEA reuptake, FAAH inhibition and antagonism of  $\Delta^9$ THC at CB<sub>1</sub>R separately renders  
7 such an experiment challenging to undertake.

8 We believe this to be the first time that possible CBG effects on feeding have been examined, although no  
9 significant CBG-mediated effects were observed. It is unlikely that CBG administration can exert any effects via  
10 direct CB<sub>1</sub>R binding since it has a very low affinity for CB<sub>1</sub>R (disassociation constant: 440 nM; Gausson et al.  
11 2007, c.f.  $\Delta^9$ THC: 39.5 nM; Bayewitch et al. 1996). However, CBG is a known AEA reuptake inhibitor  
12 (Ligresti et al. 2006) such that CBG could induce increased brain concentrations of AEA which could  
13 conceivably produce similar effects to that seen following administration of exogenous AEA. However, in the  
14 presented experimental paradigm, we had reduced eCB tone using a prefeed. Therefore, even if CBG was  
15 inducing functionally effective AEA reuptake blockade, little endogenous AEA would be present in the CNS  
16 and so reuptake blockade would be unable to potentiate CBG-mediated behavioral effects. *Id est*, were CBG to  
17 be tested using a food restricted paradigm which elevated eCB production, then its putative effects on AEA  
18 reuptake inhibition may begin to induce significant effects on feeding patterns. As such, while the data we now  
19 present suggests that CBG administration has no effect on feeding patterns, different results may be found using  
20 a different experimental paradigm.

21 In summary, following the administration of CBN alone we observed significant increases in appetitive,  
22 consummatory and total intake behaviors. Thus we suggest that a balance exists between endocannabinergic  
23 tone and pCB-mediated CB<sub>1</sub>R activation. This balance manifests as increasing feeding behaviors (appetitive,  
24 consummatory and total chow intake increase) with increasing CB<sub>1</sub>R activation, and decreasing feeding  
25 behaviors with decreasing CB<sub>1</sub>R activation. The data we have presented suggests that as CB<sub>1</sub>R activation is  
26 reduced, feeding behaviors decay and the weakest behaviors are lost first (increased total chow consumption <  
27 increased meal one duration < increased meal one chow intake < increased hourly intake & reduced latency to  
28 the first meal). Such a theory is supported by currently available literature since only recently have significant

1 effects on  $\Delta^9$ THC-mediated meal pattern changes in rats been observed, but AEA-induced increases in total  
2 chow intake, appetitive and consummatory behaviors have been demonstrated (see Farrimond et al. 2011a for  
3 review). Furthermore, we have also demonstrated significant, short-term CBD-mediated reductions in feeding  
4 which, we suggest, are due to reduced consummatory behaviors following CBD administration. However, given  
5 CBDs pharmacological profile, such effects are unlikely to be CB<sub>1</sub>R-mediated. Finally, we have observed that  
6 the administration of CBG induces no significant alterations to feeding patterns in the presented paradigm. A  
7 direct comparison between these three drug treatments is necessarily limited by the large variability in response  
8 seen under vehicle conditions, and as such, the robustness of the effects we describe here should be confirmed  
9 by a fully randomized replication of our study.

10

## Conclusion

Using a prefeed paradigm, CBN induced significant CB<sub>1</sub>R-mediated hyperphagia in male rats via significant reductions in the latency to feed and significant increases in the food consumed during the first hour and meal, alongside significant increases in the total amount of food consumed when compared to vehicle-treatments. Conversely, CBD administration reduced total feeding over a four hour period. Neither  $\Delta^9$ THCV nor CBG administration exerted effects on feeding behaviors in this paradigm.

As CBN has not so far been shown to have psychoactive properties it could be a useful anti-anorexic agent, since in this study CBN administration significantly increased intake over the total test period. Clearly, further experiments are required to fully characterize the effects of both chronic and acute CBN administration on food consumption and body weight. Moreover, the data we have presented here when compared to some of our previous data (Farrimond et al. 2010b) suggests that CBN and  $\Delta^9$ THC, when co-administered, may synergistically induce powerful hyperphagic effects. Therefore, co-administration of CBN and  $\Delta^9$ THC may also exhibit anti-anorexic properties.

Given CBD's well documented non-psychotropic nature and its high tolerability in humans, further characterization of its effects on feeding reduction and the mechanisms via which CBD induces such effects are also clearly warranted. Such tests may provide an interesting insight into the subtle feeding effects of CBD we have observed here and it would be particularly interesting to identify a non-eCB system-mediated mechanism of action of CBD in relation to feeding behaviors.

1    **Acknowledgements**

2    This research was supported in part by the University of Reading Research Endowment Trust Fund (to JAF).  
3    The authors thank Ms. Pam Rummings and her team for technical assistance and GW Pharmaceuticals for the  
4    kind gift of purified phytocannabinoids.

5

6    **Ethical compliance**

7    All procedures were performed in compliance with the requirements of the United Kingdom Animals (Scientific  
8    Procedures) Act 1986 and all other applicable laws and standards in the U.K.

9



## References

- Abel E (1975) Cannabis: effects on hunger and thirst. *Behavioral biology* 15: 255.
- Amar B (2006) Cannabinoids in medicine: A review of their therapeutic potential. *J Ethnopharmacol* 105: 1-25.
- Bayewitch M, Rhee MH, Avidor-Reiss T, Breuer A, Mechoulam R, Vogel Z (1996) (-)-Delta9-tetrahydrocannabinol antagonizes the peripheral cannabinoid receptor-mediated inhibition of adenylyl cyclase. *Journal of Biological Chemistry* 271: 9902-5.
- BNF (2006) British National Formulary. BMJ Publishing Group, London
- Cadoni C, Pisanu A, Solinas M, Acquas E, Di Chiara G (2001) Behavioural sensitization after repeated exposure to Delta 9-tetrahydrocannabinol and cross-sensitization with morphine. *Psychopharmacology* 158(3):259-66.
- Cascio MG, Gauson LA, Stevenson LA, Ross RA, Pertwee RG (2010) Evidence that the plant cannabinoid cannabigerol is a highly potent  $\alpha$ 2-adrenoceptor agonist and moderately potent 5HT1A receptor antagonist. *British Journal of Pharmacology* 159: 129-141.
- Clark RT (2009) Annual Report Pursuant to Section 13 or 15(d) of the Securities Exchange Act of 1934 For the Fiscal Year Ended December 31, 2008 United States Securities and Exchange Commission. Merck & Co., Inc., Washington
- De Filippis Dea (2008) Effect of cannabidiol on sepsis-induced motility disturbances in mice: involvement of CB1 receptors and fatty acid amide hydrolase. *Neurogastroenterology and Motility* 20: 919-927.
- Deiana S, Watanabe A, Yamasaki Y, Amada N, Arthur M, Fleming S, Woodcock H, Dorward P, Pigliacampo B, Close S, Platt B, Riedel G (2011) Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarin (CBDV),  $\Delta^9$ -tetrahydrocannabinol (THCV) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive-compulsive behaviour. *Psychopharmacology*, DOI 10.1007/s00213-011-2415-0
- Devane W, Dysarz F, Johnson MR, Melvin LS, Howlett A (1988) Determination and characterisation of a cannabinoid receptor in the rat brain. *Molecular Pharmacology* 34: 605-613.
- Devane WA, Hanus L, Breuer A, Pertwee RG, Stevenson LA, Griffin G, Gibson D, Mandelbaum A, Etinger A, Mechoulam R (1992) Isolation and structure of a brain constituent that binds to the cannabinoid receptor. *Science* 258: 1946-1949.
- EMA (2009) Procedural steps taken and scientific information after the authorisation Acomplia. In: Agency EM (ed) EMEA/H/C/000666/A20/0012. European Union, Brussels
- Engeli S, Böhnke J, Feldpausch M, Gorzelnik K, Janke J, Bátkai S, Pacher P, Harvey-White J, Luft F, Sharma A (2005) Activation of the peripheral endocannabinoid system in human obesity. *Diabetes* 54: 2838.
- Evans AT, Formukong E, Evans FJ (1987) Activation of phospholipase A2 by cannabinoids: Lack of correlation with CNS effects. *FEBS Letters* 211: 119-122.
- Farrimond J, Hill A, Whalley B, Williams C (2010a) Cannabis constituents modulate 9-tetrahydrocannabinol-induced hyperphagia in rats. *Psychopharmacology* 210: 97-106.
- Farrimond JA, Mercier MS, Whalley BJ, Williams CM (2011a) Cannabis sativa and the endogenous cannabinoid system: therapeutic potential for appetite regulation. *Phytotherapy Research* 25: 18.
- Farrimond JA, Whalley BJ, Williams CM (2010b) A low  $\Delta^9$ tetrahydrocannabinol cannabis extract induces hyperphagia in rats. *Behavioural pharmacology* 21: 769-773.
- Farrimond JA, Whalley BJ, Williams CM (2011b) Non- $\Delta^9$ tetrahydrocannabinol phytocannabinoids are effective modulators of rat feeding patterns in vivo. *Behavioural pharmacology* in press.
- Felder CC, Joyce KE, Briley EM, Mansouri J, Mackie K, Blond O, Lai Y, Ma AL, Mitchell RL (1995) Comparison of the pharmacology and signal transduction of the human cannabinoid CB1 and CB2 receptors. *Molecular Pharmacology* 48: 443-50.
- Gaoni Y, Mechoulam R (1964) Isolation, Structure, and Partial Synthesis of an Active Constituent of Hashish. *Journal of the American Chemical Society* 86: 1646-1647.
- Gauson L, Stevenson L, Thomas A, Baillie G, Ross R, Pertwee R (2007) Cannabigerol behaves as a partial agonist at both CB1 and CB2 receptors. *Symposium on the Cannabinoids. International Cannabinoid Research Society, Burlington, Vermont, USA*, pp 206
- Glass M (2001) The role of Cannabinoids in neurodegenerative diseases. *Progress In Neuro-psychopharmacology and Biological Psychiatry* 25: 743-765.
- Hao S, Avraham Y, Mechoulam R, Berry EM (2000) Low dose anandamide affects food intake, cognitive function, neurotransmitter and corticosterone levels in diet-restricted mice. *European Journal of Pharmacology* 392: 147-156.


- 1 Herkenham M, Lynn AB, Johnson MR, Melvin LS, de Costa BR, Rice KC (1991) Characterization and  
2 localization of cannabinoid receptors in rat brain: a quantitative in vitro autoradiographic study. *Journal*  
3 *of Neuroscience* 11: 563-83.
- 4 Hill A, Williams C, Whalley B, Stephens G (2011) Phytocannabinoids as novel therapeutic agents in CNS  
5 disorders. *Pharmacology & Therapeutics* in press.
- 6 Ignatowska-Jankowska B, Jankowski M, Swiergiel A (2010) Cannabidiol decreases body weight gain in rats:  
7 involvement of CB2 receptors. *Neuroscience Letters* 490: 82-84.
- 8 Izzo AA, Borrelli F, Capasso R, Di Marzo V, Mechoulam R (2009) Non-psychotropic plant cannabinoids: new  
9 therapeutic opportunities from an ancient herb. *Trends in Pharmacological Sciences* 30: 515-527.
- 10 Jamshidi N, Taylor DA (2001) Anandamide administration into the ventromedial hypothalamus stimulates  
11 appetite in rats. *British Journal of Pharmacology* 134: 1151-1154.
- 12 Koch JE, Matthews SM (2001) 9-Tetrahydrocannabinol Stimulates Palatable Food Intake in Lewis Rats: Effects  
13 of Peripheral and Central Administration. *Nutritional Neuroscience* 4: 179-187.
- 14 Ligresti A, Moriello A, Starowicz K, Matias I, Pisanti S, De Petrocellis L, Laezza C, Portella G, Bifulco M, Di  
15 Marzo V (2006) Antitumor activity of plant cannabinoids with emphasis on the effect of cannabidiol on  
16 human breast carcinoma. *Journal of Pharmacology and Experimental Therapeutics* 318: 1375.
- 17 Martin BR, Balster RL, Razdan RK, Harris LS, Dewey WL (1981) Behavioral comparisons of the stereoisomers  
18 of tetrahydrocannabinols. *Life Science* 29: 565-74.
- 19 Matsuda LA, Lolait SJ, Brownstein MJ, Young AC, Bonner TI (1990) Structure of a cannabinoid receptor and  
20 functional expression of the cloned cDNA. *Nature* 346: 561-4.
- 21 Mechoulam R, Ben-Shabat S, Hanus L, Ligumsky M, Kaminski NE, Schatz AR, Gopher A, Almog S, Martin  
22 BR, Compton DR (1995) Identification of an endogenous 2-monoglyceride, present in canine gut, that  
23 binds to cannabinoid receptors. *Biochemical Pharmacology* 50: 83-90.
- 24 Munro S, Thomas KL, Abu-Shaar M (1993) Molecular characterization of a peripheral receptor for  
25 cannabinoids. *Nature* 365: 61-65.
- 26 Nahas G, Harvey DJ, Sutin K, Turndorf H, Cancro R (2002) A molecular basis of the therapeutic and  
27 psychoactive properties of cannabis ( $\Delta^9$ -tetrahydrocannabinol), *Progress in Neuro-*  
28 *Psychopharmacology and Biological Psychiatry*, 26:721-730.
- 29 O'Shaughnessy WB (1843) On the Preparations of the Indian Hemp, or gunjah: Cannabis Indica Their Effects  
30 on the Animal System in Health, and their Utility in the Treatment of Tetanus and other Convulsive  
31 Diseases. *Provincial Medical Journal* 5.
- 32 Pertwee RG (2008) The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids:  $\Delta^9$ -  
33 tetrahydrocannabinol, cannabidiol and  $\Delta^9$ -tetrahydrocannabivarin. *British Journal of Pharmacology*  
34 153: 199-215.
- 35 Petrosino S, Ligresti A, Di Marzo V (2009) Endocannabinoid chemical biology: a tool for the development of  
36 novel therapies. *Current Opinion in Chemical Biology* 13: 309-320.
- 37 Pryce G, Ahmed Z, Hankey D, Jackson S, Croxford J, Pocock J, Ledent C, Petzold A, Thompson A,  
38 Giovannoni G, Cuzner M, Baker D (2003) Cannabinoids inhibit neurodegeneration in models of  
39 multiple sclerosis. *Brain* 126: 2191-2202.
- 40 Rhee MH, Vogel Z, Barg J, Bayewitch M, Levy R, Hanus L, Breuer A, Mechoulam R (1997) Cannabinol  
41 derivatives: binding to cannabinoid receptors and inhibition of adenylylcyclase. *Journal of Medicinal*  
42 *Chemistry* 40: 3228-33.
- 43 Riedel G, Fadda P, McKillop-Smith S, Pertwee R, Platt B, Robinson L (2009) Synthetic and plant-derived  
44 cannabinoid receptor antagonists show hypophagic properties in fasted and non-fasted mice. *British*  
45 *Journal of Pharmacology* 156: 1154-1166.
- 46 Rinaldi-Carmona M, Barth F, Héaulme M, Shire D, Calandra B, Congy C, et al. (1999) SR141716A, a potent  
47 and selective antagonist of the brain cannabinoid receptor. *FEBS letters* 350(2-3):240-44.
- 48 Rubino T, Viganò D, Massi P, Parolaro D (2001) The psychoactive ingredient of marijuana induces behavioural  
49 sensitization. *European Journal of Neuroscience* 14(5):884-86.
- 50 Ryan D, Drysdale AJ, Lafourcade C, Pertwee RG, Platt B (2009) Cannabidiol Targets Mitochondria to Regulate  
51 Intracellular Ca<sup>2+</sup> Levels. *Journal of Neuroscience* 29: 2053-2063.
- 52 Scopinho AA, Guimaraes FS, Correa F, Resstel L (2011) Cannabidiol inhibits the hyperphagia induced by  
53 cannabinoid-1 or serotonin-1A receptor agonists. *Pharmacology Biochemistry and Behavior*.
- 54 Sofia RD, Knobloch LC (1976) Comparative effects of various naturally occurring cannabinoids on food,  
55 sucrose and water consumption by rats. *Pharmacology Biochemistry and Behavior* 4: 591-599.
- 56 Sugiura T, Kondo S, Sukagawa A, Nakane S, Shinoda A, Itoh K, Yamashita A, Waku K (1995) 2-  
57 Arachidonoylglycerol: a possible endogenous cannabinoid receptor ligand in brain. *Biochemical and*  
58 *Biophysical Research Communications* 215: 89-97.

- 1 Thomas A, Stevenson LA, Wease KN, Price MR, Baillie G, Ross RA, Pertwee RG (2005) Evidence that the  
2 plant cannabinoid  $\Delta^9$ -tetrahydrocannabivarin is a cannabinoid CB<sub>1</sub> and CB<sub>2</sub> receptor antagonist. *British*  
3 *Journal of Pharmacology* 7: 917-26.
- 4 Van Gaal LF, Rissanen AM, Scheen AJ, Ziegler O, Rössner S (2005) Effects of the cannabinoid-1 receptor  
5 blocker rimonabant on weight reduction and cardiovascular risk factors in overweight patients: 1-year  
6 experience from the RIO-Europe study. *Lancet* 365: 1389-1397.
- 7 Wang T, Collet J, Shapiro S, Ware M (2008) Adverse effects of medical cannabinoids: a systematic review.  
8 *Canadian Medical Association Journal* 178: 1669.
- 9 Wiley J, Burston J, Leggett D, Alekseeva O, Razdan R, Mahadevan A, Martin B (2005) CB1 cannabinoid  
10 receptor-mediated modulation of food intake in mice. *British Journal of Pharmacology* 145: 293-300.
- 11 Williams CM, Kirkham TC (1999) Anandamide induces overeating: mediation by central cannabinoid CB<sub>1</sub>  
12 receptors. *Psychopharmacology* 143: 315-317.
- 13 Williams CM, Kirkham TC (2002a) Observational analysis of feeding induced by Delta9-THC and anandamide.  
14 *Physiology and Behaviour* 76: 241-50.
- 15 Williams CM, Kirkham TC (2002b) Reversal of D<sup>9</sup>-THC hyperphagia by SR141716 and naloxone but not  
16 dexfenfluramine. *Pharmacology Biochemistry and Behavior* 71: 333-340.
- 17 Williams CM, Rogers PJ, Kirkham TC (1998) Hyperphagia in pre-fed rats following oral D<sup>9</sup>-THC. *Physiology*  
18 *and Behavior* 65: 343-346.

19


#	Phytocannabinoid	Doses (mg/kg)
1	Cannabidiol	0.00, 0.04, 0.44 & 4.40
2	Cannabigerol	0.00, 0.176, 1.76 & 17.60
3	Cannabinol	0.00, 0.26, 2.60 & 26.00 & 26.00 + 1.00 SR141716A

**Table 1: Phytocannabinoid doses employed in this study. All phytocannabinoids were administered p.o. while SR141716A was administered s.c.. All drugs were administered at an injection volume of 1.0 ml/kg, one hour before testing began.**




**Figure 1: Mean hourly chow (A), cumulative hourly chow (B) and meal one chow consumption (C) and meal pattern (D) following administration of CBN (0, 0.26, 2.60 and 26.00 mg/kg; p.o.).**

CBN administration significantly increased hour one intake (A: white bars) and chow consumption over all cumulative hourly arrangements (B). Furthermore, following CBN administration significant increases in chow consumption during the first meal (C) and highly significant decreases in the latency to feed (D) were observed. No statistical analyses have been performed on second meal data as animals consumed too few second meals. Meal 2 bars are included for reference only. In panel D meal duration is represented by the length of bar and is provided numerically above each bar (min ± SEM). Chow intake (A, B and C) is represented as mean intake ± SEM. \* denotes  $p \leq 0.05$  in Bonferroni post-hoc test following a significant one-way ANOVA result versus vehicle-treatment. – denotes a significant Bonferroni post-hoc test following a significant one-way ANOVA on meal 1 latency.




**Figure 2: Mean hourly chow (A), cumulative hourly chow (B) and meal one chow consumption (C) and meal pattern (D) following administration of CBN and SR14171A (0 and 26.00 CBN mg/kg; p.o. and 1.0 mg/kg SR141716A; s.c.).**

The response recorded previously (see figure 1) following the highest dose of CBN (26.00 mg/kg; p.o.) is compared to those following CBN (26.00 mg/kg; p.o.) and SR141716A (1.0 mg/kg; s.c.) co-administration. Co-administration of SR141716A with CBN blocked CBN-mediated increases in hour one intake (A: white bars), meal one size (C) and duration (D) and the latency to feed (D). No statistical analyses have been performed on second meal data as animals consumed too few second meals. Meal 2 bars are included for reference only. In panel D meal duration is represented by the length of bar and is provided numerically above each bar (min ± SEM). Chow intake (A, B and C) is represented as mean intake ± SEM. \* denotes  $p \leq 0.05$  in Bonferroni post-hoc test following a significant one-way ANOVA result versus vehicle-treatment. – denotes a significant Bonferroni post-hoc test following a significant one-way ANOVA on meal 1 latency.



**Figure 3: Mean hourly chow (A), cumulative hourly chow (B) and meal one chow consumption (C) and meal pattern (D) following administration of CBD (0.00, 0.044, 0.44 and 4.40 mg/kg; p.o.).**

CBD administration significantly reduced chow intake over the period of the test (B). No statistical analyses have been performed on second meal data as animals consumed two few second meals. Meal 2 bars are included for reference only. In panel D meal duration is represented by the length of bar and is provided numerically above each bar (min ± SEM). \* denotes  $p \leq 0.05$  in Bonferroni post-hoc test following a significant one-way ANOVA result versus vehicle-treatment. – denotes a significant Bonferroni post-hoc test following a significant one-way ANOVA on meal 1 latency.



**Figure 4: Mean hourly chow (A), cumulative hourly chow (B) and meal one chow consumption (C) and meal pattern (D) following administration of CBG (0.00, 0.176, 1.76 and 17.60 mg/kg; p.o.).**

Administration of CBG induced no significant deviations from vehicle-treatments for any measure. No statistical analyses have been performed on second meal data as animals consumed too few second meals. Meal 2 bars are included for reference only. In panel D meal duration is represented by the length of bar and is provided numerically above each bar (min ± SEM). Chow intake (A, B and C) is represented as mean intake ± SEM. \* denotes  $p \leq 0.05$  in Bonferroni post-hoc test following a significant one-way ANOVA result versus vehicle-treatment. – denotes a significant Bonferroni post-hoc test following a significant one-way ANOVA on meal 1 latency.